

A CALCULUS OF FOURIER INTEGRAL OPERATORS AND THE GLOBAL FUNDAMENTAL SOLUTION FOR A SCHRÖDINGER EQUATION

HITOSHI KITADA

(Received January 12, 1981)

Introduction

In Kitada and Kumano-go [6] we studied a theory of Fourier integral operators and constructed the fundamental solution $U_h(t, s_0)$ for a pseudo-differential equation of Schrödinger's type:

$$(1) \quad \begin{cases} L_h u \equiv \left(\frac{1}{i} \frac{\partial}{\partial t} + H_h(t, X, D_x) \right) u = 0, \\ u|_{t=s_0} = f \in \mathcal{D} \quad (s_0 \in R^1), \end{cases}$$

in the form of a Fourier integral operator for t near s_0 . Here

$$(2) \quad H_h(t, x, \xi) = h^{\delta-\rho} H(t, h^{-\delta} x, h^\rho \xi) \quad (0 < h < 1, 0 \leq \delta \leq \rho \leq 1)$$

covers a rather general class of smooth time-dependent potentials $V(t, x)$ if $H(t, x, \xi)$ is of the form $H(t, x, \xi) = \frac{1}{2} |\xi|^2 + V(t, x)$. However, contrary to the generality of $H(t, x, \xi)$ that we can deal with, the time range in which we can represent $U_h(t, s_0)$ as a single Fourier integral operator was very small. The similar situations are also the case in Fujiwara's construction ([2], [3]) of the fundamental solution, except the results in [3, §4].

In this paper we shall make a rather strong restriction on the potential $V(t, x)$ (see Assumption (A) in section 3), and construct the fundamental solution $U_h(t, s_0)$ for (1) with $H(t, x, \xi) = \frac{1}{2} |\xi|^2 + V(t, x)$ in the form of a single (conjugate) Fourier integral operator for all $t \geq s_0$, when s_0 is sufficiently large.

To do so, in sections 1 and 2 we shall introduce a class of (conjugate) Fourier integral operators and investigate their calculus, which is also our purpose in the present paper. The symbol class for our Fourier integral operators is the same as in [6], while the class of phase functions is different from [6] (see Definition 1.1). The characteristic feature of our phase functions $\phi_h(x, \xi)$ is, roughly speaking, that the function $J_h(x, \xi) \equiv \phi_h(x, \xi) - x \cdot \xi$ is "small" in the

sense that only the derivatives of $\nabla_x J_h(x, \xi)$ are small, while in [6] we assumed that the derivatives of both $\nabla_x J_h(x, \xi)$ and $\nabla_\xi J_h(x, \xi)$ are small. This relaxation is possible, because, in the present paper, we restrict ourselves to considering only the conjugate Fourier integral operators of the form

$$(3) \quad P_h(\phi_h^*)f(x) = 0_s - \iint e^{i(x \cdot \xi - \phi_h(x', \xi))} p_h(\xi, x') f(x') dx' d\xi,$$

while in [6] we considered both Fourier, and conjugate Fourier, integral operators.

Section 2 is devoted to proving a theorem concerning the calculus of conjugate Fourier integral operators, which is different from [6] in the point that we shall treat $\nu+1$ conjugate Fourier integral operators directly, while in [6] the product of two Fourier integral operators and that of Fourier and conjugate Fourier integral operators were basic. This result will allow us in section 4 to make a global calculus in time of the local fundamental solutions represented as conjugate Fourier integral operators.

In section 3, we shall in turn consider the Schrödinger equation (1) with $H(t, x, \xi) = \frac{1}{2} |\xi|^2 + V(t, x)$, where $V(t, x)$ is assumed to satisfy

$$(4) \quad \sup_{x \in \mathbb{R}^n} |\partial_x^\alpha V(t, x)| \leq C_\alpha (1 + |t|)^{-|\alpha| - \varepsilon}$$

for $|\alpha| \neq 0$ with $\varepsilon > 0$. We shall first give several estimates concerning the classical orbit $(q, p)(t, s; x, \xi)$ defined as the solution of the Hamilton equation

$$(5) \quad \begin{cases} \frac{dq}{dt}(t, s) = p(t, s), \\ \frac{dp}{dt}(t, s) = -\nabla_x V(t, q(t, s)) \end{cases}$$

with the initial condition $(q, p)(s, s) = (x, \xi)$. From this $(q, p)(t, s; x, \xi)$, we shall construct the phase function $\phi(s, t; x, \xi)$ as the solution of the eikonal equation

$$(6) \quad \begin{cases} \partial_s \phi(s, t; x, \xi) + H(s, x, \nabla_x \phi(s, t; x, \xi)) = 0, \\ \phi(t, t; x, \xi) = x \cdot \xi, \end{cases}$$

which can be solved globally for $t \geq s$ when s is sufficiently large, as well as locally for $|t-s| \leq \delta_0 (\ll 1)$. Then we shall define the global and local approximate fundamental solutions of order m ($m=0$ or ∞) in the sense of [6] in the form

$$(7) \quad \begin{aligned} & E_h^m(\phi_h(s, t)^*)f(x) \\ &= 0_s - \iint e^{i(x \cdot \xi - \phi_h(s, t; x', \xi))} e_h^m(t, s; \xi, x') f(x') dx' d\xi \end{aligned}$$

for $t \geq s$ or $|t-s| \leq \delta_0$, where $\phi_h(s, t; x', \xi) \equiv h^{\delta-\rho} \phi(s, t; h^{-\delta} x', h^\rho \xi)$, and $m=0$ in

case $0 \leq \delta \leq \rho \leq 1$ and $m = \infty$ in case $0 \leq \delta < \rho \leq 1$. We shall then summarize the important estimates concerning these approximate fundamental solutions as Theorem 3.11 at the end of section 3.

Using these estimates, in section 4 we shall first construct the local fundamental solution $U_h(t, s)$ for $|t-s| \leq \delta_0$ as a conjugate Fourier integral operator in quite a similar way to [6]. Then using the global solution $\phi(s, t; x, \xi)$ of (6) and the results of section 2 on the calculus, we shall represent the global fundamental solution $U_h(t, s_0) = U_h(t, t_\nu)U_h(t_\nu, t_{\nu-1}) \cdots U_h(t_1, s_0)$ ($0 < t_j - t_{j-1} \leq (t-s_0)/(\nu+1) \leq \delta_0, s_0 < t_1 < \cdots < t_\nu \leq t$) as a single conjugate Fourier integral operator for sufficiently large s_0 . For general s_0 , we can therefore represent the global fundamental solution $U_h(t, s_0)$ as a product of a finite number of conjugate Fourier integral operators, the number being independent of t but dependent on s_0 . At the same time, we shall also give some estimates for the differences between the fundamental solution $U_h(t, s)$ and the global approximate fundamental solutions $E_h^m(\phi_h(s, t)^*)$ when $t \geq s$ for sufficiently large s . One of these estimates played a crucial role in the proof of the completeness of modified wave operators in [5].

We note that our assumption on the potential $V(t, x)$, hence on the Hamiltonian $H(t, x, \xi)$, is not symmetric in x and ξ , while the assumption adopted in [6] was symmetric. Moreover, under our present assumption (4), the classical orbit $y = q(s, t; x, \xi)$ in the configuration space is uniquely determined by its initial and final positions x and y for $t \geq s$ when s is sufficiently large (by (3.11) below), which makes it possible to construct the global phase function (compare this with the situations in §4 of Fujiwara [3]).

Recently, Nishiwada [9] gave an explicit expression, which is written by means of one or two integral transformations, of the global fundamental solution for a Schrödinger equation with a quadratic Hamiltonian. However his assumption and method are different from ours.

1. Fourier integral operators

In this and the next sections, we introduce a class of Fourier integral operators and investigate their properties, especially their calculus. We first explain some basic notations we shall use in the following. For any point $x = (x_1, \dots, x_n)$ in the n -dimensional Euclidian space R^n , we define its norm $|x|$ by $|x| = (\sum_{j=1}^n x_j^2)^{1/2}$, and for any $n \times n$ real matrix $A = (a_{ij})$ we define $|A| = \sup_{|x|=1, x \in R^n} |Ax|/|x|$. Let $\alpha = (\alpha_1, \dots, \alpha_n)$ be a multi-index whose components α_j are non-negative integers and let $x, y, z \in R^n$. Then we use the following notations:

$$|\alpha| = \alpha_1 + \cdots + \alpha_n, \quad \alpha! = \alpha_1! \cdots \alpha_n!, \quad x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n},$$

$$\partial_x^\alpha = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_n}^{\alpha_n}, \quad D_x^\alpha = D_{x_1}^{\alpha_1} \cdots D_{x_n}^{\alpha_n}, \quad \partial_{x_j} = \frac{\partial}{\partial x_j}, \quad D_{x_j} = \frac{1}{i} \frac{\partial}{\partial x_j},$$

$$\begin{aligned} \nabla_x &= {}^t(\partial_{x_1}, \dots, \partial_{x_n}), \quad \vec{\nabla}_x = {}^t\nabla_x, \\ \langle x \rangle &= \sqrt{1+|x|^2}, \quad \langle x; y \rangle = \sqrt{1+|x|^2+|y|^2}, \\ \langle x; y; z \rangle &= \sqrt{1+|x|^2+|y|^2+|z|^2}. \end{aligned}$$

By \mathcal{S} we denote the Schwartz space of rapidly decreasing functions on R^n . For $f \in \mathcal{S}$ we define its Fourier transform $\hat{f}(\xi) = \mathcal{F}f(\xi)$ by

$$\mathcal{F}f(\xi) = \int e^{-ix \cdot \xi} f(x) dx, \quad x \cdot \xi = \sum_{j=1}^n x_j \xi_j.$$

The inverse Fourier transform $\mathcal{F}^{-1}f$ of $f \in \mathcal{S}$ is given by

$$(\mathcal{F}^{-1}f)(x) = \int e^{ix \cdot \xi} f(\xi) d\xi, \quad d\xi = (2\pi)^{-n} d\xi.$$

DEFINITION 1.1. 1° Let $0 \leq \tau < 1, 0 \leq \sigma < \infty$ and $0 \leq \delta \leq \rho \leq 1$. A family $\{\phi_h(x, \xi)\}_{0 < h < 1}$ of C^∞ -functions $\phi_h(x, \xi)$ in $R^n \times R^n$ is said to belong to the class $\{P_{\rho, \delta}^{[\tau]}(\tau, \sigma; h)\}_{0 < h < 1}$, if the function $\tilde{J}_h(x, \xi)$ defined by

$$(1.1) \quad \begin{cases} \tilde{J}_h(x, \xi) = \tilde{\phi}_h(x, \xi) - x \cdot \xi = h^{\rho-\delta} J_h(h^\delta x, h^{-\rho} \xi), \\ \tilde{\phi}_h(x, \xi) = h^{\rho-\delta} \phi_h(h^\delta x, h^{-\rho} \xi), \\ J_h(x, \xi) = \phi_h(x, \xi) - x \cdot \xi \end{cases}$$

satisfies

$$(1.2) \quad \begin{cases} \text{i)} & \sup_{h, x, \xi} \{|\nabla_\xi \tilde{J}_h(x, \xi)| / \langle \xi \rangle\} + \sup_{h, x, \xi} |\nabla_x \tilde{J}_h(x, \xi)| < \infty, \\ \text{ii)} & \sup_{h, x, \xi} |\vec{\nabla}_\xi \nabla_\xi \tilde{J}_h(x, \xi)| \leq \sigma, \\ \text{iii)} & \max_{|\alpha+\beta|=1} \sup_{h, x, \xi} |\partial_\xi^\alpha D_x^\beta \nabla_x \tilde{J}_h(x, \xi)| \leq \tau \end{cases}$$

and

$$(1.3) \quad \sup_{h, x, \xi} |\tilde{J}_{h(\beta)}^{(\alpha)}(x, \xi)| < \infty \quad \text{for } |\alpha+\beta| \geq 3,$$

where $\tilde{J}_{h(\beta)}^{(\alpha)}(x, \xi) = \partial_\xi^\alpha D_x^\beta \tilde{J}_h(x, \xi)$. For simplicity we also write this as $\phi_h(x, \xi) \in P_{\rho, \delta}^{[\tau]}(\tau, \sigma; h)$.

2° For $\phi_h(x, \xi) \in P_{\rho, \delta}^{[\tau]}(\tau, \sigma; h)$ we define a semi-norm $|J_h|_{l, m}$ for integers $l, m \geq 0$ by

$$(1.4) \quad |J_h|_{l, m} = \max_{l \leq |\alpha+\beta| \leq l+m} \sup_{h, x, \xi} |\tilde{J}_{h(\beta)}^{(\alpha)}(x, \xi)|.$$

REMARK. In section 4 we shall also use the class $\{P_{\rho, \delta}(\tau, l; h)\}_{0 < h < 1}$ ($0 \leq \tau < 1, l=0, 1, 2, \dots$) defined in Kitada and Kumano-go [6]. Here for the sake of the later convenience, we state its definition. $\{\phi_h(x, \xi)\}_{0 < h < 1} \in \{P_{\rho, \delta}(\tau, l; h)\}_{0 < h < 1}$ or $\phi_h(x, \xi) \in P_{\rho, \delta}(\tau, l; h)$ means that J_h and \tilde{J}_h defined by (1.1) satisfy

$$|J_h|_l \equiv \sum_{|\alpha+\beta| \leq l} \sup_{h, x, \xi} \{|\tilde{J}_{h(\beta)}^{(\alpha)}(x, \xi)| / \langle x; \xi \rangle^{2-|\alpha+\beta|}\}$$

$$+ \sum_{2 \leq |\alpha + \beta| \leq 2+l} \sup_{h, x, \xi} |\tilde{J}_{h(\beta)}^{(\alpha)}(x, \xi)| \leq \tau$$

and

$$\sup_{h, x, \xi} |\tilde{J}_{h(\beta)}^{(\alpha)}(x, \xi)| < \infty \quad \text{for } |\alpha + \beta| \geq 3.$$

We next define the symbol classes which are the same as those introduced in [6].

DEFINITION 1.2. 1° Let $m \in R^1$ and $0 \leq \delta \leq \rho \leq 1$. A family $\{p_h(x, \xi, x')\}_{0 < h < 1}$ of C^∞ -functions $p_h(x, \xi, x')$ is said to belong to the class $\{B_{\rho, \delta}^m(h)\}_{0 < h < 1}$ if $\{p_h\}_{0 < h < 1}$ satisfies

$$(1.5) \quad \begin{aligned} |p_h|_l^{(m)} &= |\{p_h\}_{0 < h < 1}|_l^{(m)} \\ &\equiv \max_{|\beta + \alpha + \beta'| \geq l} \sup_{h, x, \xi} h^{-m - \rho|\alpha| + \delta|\beta + \beta'|} |p_{h(\beta, \beta')}^{(\alpha)}(x, \xi, x')| < \infty \end{aligned}$$

for any integer $l \geq 0$, where $p_{h(\beta, \beta')}^{(\alpha)} = D_x^\beta \partial_\xi^\alpha D_{x'}^{\beta'} p_h(x, \xi, x')$. We write this also as $p_h(x, \xi, x') \in B_{\rho, \delta}^m(h)$.

2° For $m \in R^1, r \geq 0$ and $0 \leq \delta \leq \rho \leq 1$, we say that a family $\{p_h(x, \xi, x')\}_{0 < h < 1}$ of C^∞ -functions belongs to the class $\{B_{\rho, \delta}^{m, r}(h)\}_{0 < h < 1}$ if $\langle h^{-\delta} x; h^\rho \xi; h^{-\delta} x' \rangle^{-r} p_h(x, \xi, x')$ belongs to $B_{\rho, \delta}^m(h)$.

REMARK. 1° $B_{\rho, \delta}^{m, 0}(h) = B_{\rho, \delta}^m(h)$.

2° When $p_h(x, \xi, x') = p_h(x, \xi)$ (independent of x') [resp. $p_h(x, \xi, x') = p_h(\xi, x')$ (independent of x)], $p_h(x, \xi, x') \in B_{\rho, \delta}^{m, r}(h)$ is equivalent to $\langle h^{-\delta} x; h^\rho \xi \rangle^{-r} p_h(x, \xi) \in B_{\rho, \delta}^m(h)$ [resp. $\langle h^\rho \xi; h^{-\delta} x' \rangle^{-r} p_h(\xi, x') \in B_{\rho, \delta}^m(h)$]. Such symbols are called single symbols.

Proposition 1.3. Let $p_{j, h}(x, \xi, x') \in B_{\rho, \delta}^{m_j}(h)$ ($j=0, 1, 2, \dots$) such that $m_0 \leq m_1 \leq \dots \leq m_j \leq \dots \rightarrow \infty$ and let χ be a C^∞ -function on $[0, \infty]$ such that $0 \leq \chi(\theta) \leq 1$ on $[0, \infty)$ and $\chi(\theta) = 1$ (for $0 \leq \theta \leq 1/2$), $= 0$ (for $\theta \geq 1$). Then there exists a decreasing sequence $\{\varepsilon_j\}_{j=0}^\infty$ tending to zero as $j \rightarrow \infty$ such that

$$(1.6) \quad p_h(x, \xi, x') = \sum_{j=0}^\infty \chi(\varepsilon_j^{-1} h) p_{j, h}(x, \xi, x')$$

converges in $B_{\rho, \delta}^{m_0}(h)$ and

$$p_h(x, \xi, x') - \sum_{j=0}^{N-1} \chi(\varepsilon_j^{-1} h) p_{j, h}(x, \xi, x') \in B_{\rho, \delta}^{m_N}(h)$$

for any $N \geq 1$. Furthermore such $p_h \in B_{\rho, \delta}^{m_0}(h)$ is unique modulo $B^\infty(h) \equiv \bigcap_{m \in R^1} B_{\rho, \delta}^m(h)$ (independent of ρ, δ).

For the proof see Theorem 1.3 of [6].

Proposition 1.4. Let $\phi_h(x, \xi) \in P_{\rho, \delta}^{[\tau]}(\tau, \sigma; h)$ and $p_h(x, \xi, x') \in B_{\rho, \delta}^{m, r}(h)$ for $0 \leq \tau < 1, 0 \leq \sigma < \infty, 0 \leq \delta \leq \rho \leq 1, m \in R^1$ and $r \geq 0$. Then for any $f \in \mathcal{S}$ and $\chi \in$

\mathcal{S} with $\chi(0)=1$, the integrals

$$(1.7) \quad \begin{cases} P_{h,\varepsilon}[f](x) = \iint e^{i(\phi_h(x,\xi)-x'\cdot\xi)} p_h(x, \xi, x') f(x') \chi(\varepsilon\xi) dx' d\xi, \\ \bar{P}_{h,\varepsilon}[f](x) = \iint e^{i(x\cdot\xi-\phi_h(x',\xi))} p_h(x, \xi, x') f(x') \chi(\varepsilon\xi) dx' d\xi \end{cases}$$

have the limits $P_h[f](x)$ and $\bar{P}_h[f](x)$ for $\varepsilon \downarrow 0$, which are independent of χ . Moreover P_h and \bar{P}_h define continuous linear mappings from \mathcal{S} into \mathcal{S} for each $h \in (0, 1)$. We write those limits as

$$(1.8) \quad \begin{cases} P_h[f](x) = 0_s - \iint e^{i(\phi_h(x,\xi)-x'\cdot\xi)} p_h(x, \xi, x') f(x') dx' d\xi, \\ \bar{P}_h[f](x) = 0_s - \iint e^{i(x\cdot\xi-\phi_h(x',\xi))} p_h(x, \xi, x') f(x') dx' d\xi. \end{cases}$$

Proof. Putting $\psi_h(x, \xi, x') = x \cdot \xi - \phi_h(x', \xi) = (x - x') \cdot \xi - J_h(x', \xi)$, we see from (1.1)–(1.2) that $\langle \nabla_{x'} \psi_h \rangle \geq C_h \langle \xi \rangle$ for some constant $C_h > 0$. Thus $L \equiv \langle \nabla_x, \psi_h \rangle^{-1} (1 - i \nabla_x \cdot \psi_h \cdot \nabla_{x'})$ is well-defined and we have for any $l \geq 0$

$$(1.9) \quad \bar{P}_{h,\varepsilon}[f](x) = \iint e^{i\psi_h({}^tL)} [p_h(x, \xi, x') f(x') \chi(\varepsilon\xi)] dx' d\xi,$$

where tL is the transposed operator of L . Then taking $l > n + r$, noting $f \in \mathcal{S}$ and letting $\varepsilon \downarrow 0$, we have

$$(1.10) \quad \bar{P}_h[f](x) = \iint e^{i\psi_h({}^tL)} [p_h(x, \xi, x') f(x')] dx' d\xi,$$

which is independent of χ . Therefore we get

$$\begin{aligned} & x^\alpha D_x^\beta (\bar{P}_h[f])(x) \\ &= \sum_{\beta^1 + \beta^2 = \beta} \iint e^{ix\cdot\xi} D_\xi^\alpha \{ e^{-i\phi_h(x',\xi)} \xi^{\beta^1} ({}^tL)^l [D_x^{\beta^2} p_h(x, \xi, x') f(x')] \} dx' d\xi. \end{aligned}$$

We see from (1.2)–(1.3) that this is uniformly bounded in $x \in R^n$ for each fixed $h \in (0, 1)$, if l is taken sufficiently large.

For $P_h[f]$, putting $\varphi_h(x, \xi, x') = \phi_h(x, \xi) - x' \cdot \xi = (x - x') \cdot \xi + J_h(x, \xi)$, we have $\nabla_{x'} \varphi_h = -\xi$. So letting $L \equiv \langle \nabla_x \varphi_h \rangle^{-1} (1 - i \nabla_x \varphi_h \cdot \nabla_{x'})$ we have (1.10) for $P_h[f]$ and

$$\begin{aligned} & |x^\alpha D_x^\beta (P_h[f])(x)| \\ &\leq C_h \iint \langle x'; \xi \rangle^{|\alpha|} \langle \xi \rangle^{|\beta^1|} \sum_{\alpha' \leq \alpha, \beta' \leq \beta} |D_\xi^{\alpha'} ({}^tL)^l [D_x^{\beta'} p_h(x, \xi, x') f(x')]| dx' d\xi, \end{aligned}$$

which shows $P_h[f] \in \mathcal{S}$ if l is taken sufficiently large. The continuity of P_h and \bar{P}_h in \mathcal{S} is clear by the above discussions. \square

DEFINITION 1.5. 1° For $p_h(x, \xi, x') \in B_{\rho, \delta}^{m, r}(h)$ ($0 \leq \delta \leq \rho \leq 1, m \in R^1, r \geq 0$) we

define a family of pseudo-differential operators $P_h = p_h(X, D_x, X')$ by

$$(1.11) \quad P_h f(x) = 0_s - \iint e^{i(x-x') \cdot \xi} p_h(x, \xi, x') f(x') dx' d\xi$$

for $f \in \mathcal{S}$, and write this as $\{P_h\}_{0 < h < 1} \in \{B_{\rho, \delta}^{m, r}(h)\}_{0 < h < 1}$ or simply as $P_h \in B_{\rho, \delta}^{m, r}(h)$.

2° For $\phi_h(x, \xi) \in P_{\rho, \delta}^{[\tau, \sigma]}(\tau, \sigma; h)$ ($0 \leq \tau < 1, 0 \leq \sigma < \infty$) and $p_h(x, \xi, x') \in B_{\rho, \delta}^{m, r}(h)$, we define a family of Fourier, and conjugate Fourier, integral operators $P_h(\phi_h) = p_h(\phi_h; X, D_x, X')$ and $P_h(\phi_h^*) = p_h(\phi_h^*; X, D_x, X')$ by

$$(1.12) \quad \begin{cases} P_h(\phi_h)f(x) = 0_s - \iint e^{i(\phi_h(x, \xi) - x' \cdot \xi)} p_h(x, \xi, x') f(x') dx' d\xi, \\ P_h(\phi_h^*)f(x) = 0_s - \iint e^{i(x \cdot \xi - \phi_h(x, \xi))} p_h(x, \xi, x') f(x') dx' d\xi \end{cases}$$

for $f \in \mathcal{S}$. We write this as $\{P_h(\phi_h)\}_{0 < h < 1} \in \{B_{\rho, \delta}^{m, r}(\phi_h)\}_{0 < h < 1}$ and $\{P_h(\phi_h^*)\}_{0 < h < 1} \in \{B_{\rho, \delta}^{m, r}(\phi_h^*)\}_{0 < h < 1}$, or simply as $P_h(\phi_h) \in B_{\rho, \delta}^{m, r}(\phi_h)$ and $P_h(\phi_h^*) \in B_{\rho, \delta}^{m, r}(\phi_h^*)$.

REMARK. 1° If we define $q_h(x, \xi, x') = \overline{p_h(x', \xi, x)}$ for $p_h(x, \xi, x') \in B_{\rho, \delta}^{m, r}(h)$, then we have $q_h(x, \xi, x') \in B_{\rho, \delta}^{m, r}(h)$ and $(P_h f, g)_{L^2} = (f, Q_h g)_{L^2}$ for $f, g \in \mathcal{S}$.

2° For single symbols $p_h(x, \xi)$ and $q_h(\xi, x') \in B_{\rho, \delta}^{m, \alpha}(h)$, we have from Proposition 1.4 that

$$(1.13) \quad \begin{cases} P_h(\phi_h)f(x) = \int e^{i\phi_h(x, \xi)} p_h(x, \xi) \hat{f}(\xi) d\xi, \\ \widehat{Q_h(\phi_h^*)f}(\xi) = \int e^{-i\phi_h(x', \xi)} q_h(\xi, x') f(x') dx' \end{cases}$$

for $f \in \mathcal{S}$ and $\phi_h \in P_{\rho, \delta}^{[\tau, \sigma]}(\tau, \sigma; h)$.

Theorem 1.6. Let $r \geq 0$, and denote by \bar{r} the minimum integer not less than r . Let $p_h(\xi, x') \in B_{\rho, \delta}^{m, r}(h)$ and $\phi_h(x, \xi) \in P_{\rho, \delta}^{[\tau, \sigma]}(\tau, \sigma; h)$ ($0 \leq \tau < 1, 0 \leq \sigma < \infty, 0 \leq \delta \leq \rho \leq 1$) and assume that

$$(1.14) \quad p_{h(\beta)}^{(\alpha)}(\xi, x') \in B_{\rho, \delta}^{m + |\alpha| - \delta|\beta|, r - |\alpha| + \beta|}(h)$$

for $|\alpha + \beta| \leq \bar{r}$. Let $Q_h(\phi_h^*) = q_h(\phi_h^*; D_x, X') \in B_{\rho, \delta}^{m'}(\phi_h^*)$. Set

$$\begin{cases} s_h(\xi, x', \xi', x'') = p_h(\xi, x' + \nabla_{\xi} J_h(\xi, x'', \xi')) q_h(\xi', x''), \\ \nabla_{\xi} J_h(\xi, x'', \xi') = \int_0^1 \nabla_{\xi} J_h(x'', \xi' + \theta(\xi - \xi')) d\theta, \end{cases}$$

and define $r_h(\xi, x'')$ by

$$(1.15) \quad r_h(\xi, x'') = 0_s - \iint e^{-iy \cdot \eta} s_h(\xi, x'' + y, \xi - \eta, x'') dy d\eta.$$

Then we have $r_h(\xi, x'') \in B_{\rho, \delta}^{m + m', r}(h)$ and $R_h(\phi_h^*) \equiv r_h(\phi_h^*; D_x, X') = P_h Q_h(\phi_h^*)$. More precisely we have for $N \geq \bar{r}$

$$\begin{aligned}
 (1.16) \quad r_h(\xi, x'') &= \sum_{|\alpha| < \mathcal{N}} \frac{(-1)^{|\alpha|}}{\alpha!} \partial_{\xi'}^{\alpha} \{p_{h(\omega)}(\xi, \tilde{\nabla}_{\xi} \phi_h(\xi, x'', \xi')) q_h(\xi', x'')\}_{|\xi'=\xi} \\
 &= N \sum_{|\gamma| = \mathcal{N}} \frac{(-1)^{|\gamma|}}{\gamma!} \int_0^1 (1-\theta)^{N-1} t_{\gamma, h}(\xi, x''; \theta) d\theta \in B_{\rho, \delta}^{m+m'+(\rho-\delta)N}(h),
 \end{aligned}$$

where

$$\begin{aligned}
 (1.17) \quad t_{\gamma, h}(\xi, x''; \theta) &= 0_s - \iint e^{-iy \cdot \eta} \partial_{\xi'}^{\gamma} \{p_{h(\gamma)}(\xi, \theta y + \tilde{\nabla}_{\xi} \phi_h(\xi, x'', \xi')) q_h(\xi', x'')\}_{|\xi'=\xi-\eta} dy d\eta, \\
 \tilde{\nabla}_{\xi} \phi_h(\xi, x'', \xi') &= \int_0^1 \nabla_{\xi} \phi_h(x'', \xi' + \theta(\xi - \xi')) d\theta.
 \end{aligned}$$

If, in particular $q_h(\xi, x') = 1$, we have for $\tilde{N} \geq \mathcal{N}/2$

$$\begin{aligned}
 (1.16)' \quad r_h(\xi, x'') &= \sum_{|\alpha| < \tilde{N}} \frac{(-1)^{|\alpha|}}{\alpha!} \partial_{\xi'}^{\alpha} \{p_{h(\omega)}(\xi, \tilde{\nabla}_{\xi} \phi_h(\xi, x'', \xi'))\}_{|\xi'=\xi} \\
 &= \tilde{N} \sum_{|\gamma| = \tilde{N}} \frac{(-1)^{|\gamma|}}{\gamma!} \int_0^1 (1-\theta)^{\tilde{N}-1} t_{\gamma, h}(\xi, x''; \theta) d\theta \in B_{\rho, \delta}^{m+m'+2(\rho-\delta)N}(h),
 \end{aligned}$$

where

$$\begin{aligned}
 (1.17)' \quad t_{\gamma, h}(\xi, x''; \theta) &= 0_s - \iint e^{-iy \cdot \eta} \partial_{\xi'}^{\gamma} \{p_{h(\gamma)}(\xi, \theta y + \tilde{\nabla}_{\xi} \phi_h(\xi, x'', \xi'))\}_{|\xi'=\xi-\eta} dy d\eta.
 \end{aligned}$$

Proof is similar to that of Theorem 3.7 and Proposition 5.6 of [6].

Theorem 1.7. Let $\phi_h(x, \xi) \in P_{\rho, \delta}^{[x]}(\tau, \sigma; h)$ and $p_h(\xi, x') \in B_{\rho, \delta}^m(h)$ with $0 \leq \tau < 1$, $0 \leq \sigma < \infty$, $0 \leq \delta \leq \rho \leq 1$ and $m \in \mathbb{R}^1$. Then for $P_h(\phi_h^*) \equiv p_h(\phi_h^*; D_x, X') \in B_{\rho, \delta}^m(\phi_h^*)$ we have

$$\begin{aligned}
 (1.18) \quad &\|P_h(\phi_h^*)\|_{L^2 \rightarrow L^2} \\
 &\leq Ch^m |p_h|_M^{(m)} (1 + \max_{1 \leq |\alpha| + \beta| \leq M+1} \sup_{h, x, \xi} |\partial_{\xi}^{\alpha} D_x^{\beta} \nabla_x \tilde{J}_h(x, \xi)|)^{(M+1)/2},
 \end{aligned}$$

where $M = 2([n/2] + [5n/4] + 2)$; \tilde{J}_h is defined by (1.1); and C is a positive constant independent of $h \in (0, 1)$, $\{\phi_h\}_{0 < h < 1}$ and $\{p_h\}_{0 < h < 1}$.

Proof. For $f \in \mathcal{S}$ we have from (1.13)

$$P_h(\widehat{\phi_h^*})f(\xi) = \int e^{-i\phi_h(x', \xi)} p_h(\xi, x') f(x') dx'.$$

Thus we have

$$\|P_h(\phi_h^*)f\|_{L^2}^2 = (K_h f, f)_{L^2},$$

where

$$K_h f(x) = 0_s - \iint e^{i(\phi_h(x,\xi) - \phi_h(y,\xi))} p_h(\xi, y) \overline{p_h(\xi, x)} f(y) dy d\xi.$$

Noting that $\phi_h(x, \xi) - \phi_h(y, \xi) = (x - y) \cdot \tilde{\nabla}_x \phi_h(x, \xi, y) \equiv (x - y) \cdot \int_0^1 \nabla_x \phi_h(y + \theta(x - y), \xi) d\theta$ and that the mapping $\xi \mapsto \eta = \tilde{\nabla}_x \phi_h(x, \xi, y)$ has the inverse C^∞ -diffeomorphism $\eta \mapsto \tilde{\nabla}_x \phi_h^{-1}(x, \eta, y)$, since $|\tilde{\nabla}_\xi \tilde{\nabla}_x \phi_h(x, \xi, y) - I| = |\tilde{\nabla}_\xi \tilde{\nabla}_x J_h(h^{-\delta} x, h^\rho \xi, h^{-\delta} y)| \leq \sigma < 1$ by (1.2)-iii), we make a change of variable: $\eta = \tilde{\nabla}_x \phi_h(x, \xi, y)$. Then we obtain

$$K_h f(x) = 0_s - \iint e^{i\eta \cdot (x - y)} p_h(\tilde{\nabla}_x \phi_h^{-1}(x, \eta, y), y) \times \\ \times \overline{p_h(\tilde{\nabla}_x \phi_h^{-1}(x, \eta, y), x)} \left| \frac{D(\tilde{\nabla}_x \phi_h^{-1})(x, \eta, y)}{D(\eta)} \right| f(y) dy d\eta.$$

Putting

$$r_h(\tilde{x}, \tilde{\eta}, \tilde{y}) = h^{-2m} \tilde{p}_h(\tilde{\nabla}_x \tilde{\phi}_h^{-1}(\tilde{x}, h^{\rho - \delta} \tilde{\eta}, \tilde{y}), \tilde{y}) \times \\ \times \overline{\tilde{p}_h(\tilde{\nabla}_x \tilde{\phi}_h^{-1}(\tilde{x}, h^{\rho - \delta} \tilde{\eta}, \tilde{y}), \tilde{x})} \left| \frac{D(\tilde{\nabla}_x \tilde{\phi}_h^{-1})(\tilde{x}, h^{\rho - \delta} \tilde{\eta}, \tilde{y})}{D(\tilde{\eta})} \right|,$$

where $\tilde{p}_h(\xi, x) = p_h(h^{-\rho} \xi, h^\delta x)$, and making again a change of variables $x = h^\delta \tilde{x}$, $\eta = h^{-\delta} \tilde{\eta}$, $y = h^\delta \tilde{y}$, we obtain

$$K_h f(h^\delta \tilde{x}) = h^{2m} 0_s - \iint e^{i\tilde{\eta} \cdot (\tilde{x} - \tilde{y})} r_h(\tilde{x}, \tilde{\eta}, \tilde{y}) f(h^\delta \tilde{y}) d\tilde{y} d\tilde{\eta}.$$

Thus by the Calderón-Vaillancourt theorem ([1]) we have

$$\|K_h f(h^\delta \tilde{x})\|_{L^2(R_\tilde{x}^n)} \\ \leq Ch^{2m} \max_{|\beta + \alpha + \beta'| \leq M} \sup_{\tilde{x}, \tilde{\eta}, \tilde{y}} |\partial_{\tilde{x}}^\beta \partial_{\tilde{\eta}}^\alpha \partial_{\tilde{y}}^{\beta'} r_h(\tilde{x}, \tilde{\eta}, \tilde{y})| \|f(h^\delta \tilde{y})\|_{L^2(R_\tilde{y}^n)}$$

for some constant $C > 0$ independent of $h \in (0, 1)$. From $\rho - \delta \geq 0$ and the definition of r_h we get

$$|\partial_{\tilde{x}}^\beta \partial_{\tilde{\eta}}^\alpha \partial_{\tilde{y}}^{\beta'} r_h(\tilde{x}, \tilde{\eta}, \tilde{y})| \\ \leq C(1 + \max_{2 \leq |\alpha + \beta| \leq M+1} \sup_{h, x, \xi} |\partial_\xi^\alpha D_x^\beta \nabla_x J_h(x, \xi)|)^{M+1} (|p_h|_M^{(n)})^2$$

for $|\beta + \alpha + \beta'| \leq M$, where C is independent of $h \in (0, 1)$. Thus we have (1.18). \square

2. Multi-products of conjugate Fourier integral operators

Now we turn to the study of the multi-products of conjugate Fourier integral operators. We first introduce the following condition (#) for $(\nu + 1)$ -tuple $(\phi_{1,h}, \dots, \phi_{\nu+1,h})$ ($\nu \geq 1$, integer) of phase functions $\phi_{j,h} \in P_{\rho,\delta}^{[x]}(\tau_j, \sigma_j; h)$ ($j = 1, \dots, \nu + 1$):

(#) For each fixed $h \in (0, 1)$, there exists a unique C^∞ solution $\{X_{j,h}^i,$

$\Xi_{v,h}^j\}_{j=1}^v(x, \xi)$ of the equation

$$(2.1) \quad \begin{cases} X_{v,h}^j = \nabla_{\xi} \phi_{j,h}(X_{v,h}^{j-1}, \Xi_{v,h}^j), \\ \Xi_{v,h}^j = \nabla_x \phi_{j+1,h}(X_{v,h}^j, \Xi_{v,h}^{j+1}) \end{cases} \quad (j = 1, \dots, v)$$

where $X_{v,h}^0 = x$ and $\Xi_{v,h}^{v+1} = \xi$.

DEFINITION 2.1. For $(v+1)$ -tuple $(\phi_{1,h}, \dots, \phi_{v+1,h})$ of phase functions $\phi_{j,h} \in P_{\rho,\delta}^{[x]}(\tau_j, \sigma_j; h)$ satisfying $(\#)$, we define its $\#$ - $(v+1)$ product $\Phi_{v+1,h} = \phi_{1,h} \# \dots \# \phi_{v+1,h}$ by

$$(2.2) \quad \begin{aligned} & \Phi_{v+1,h}(x, \xi) \\ &= \sum_{j=1}^v (\phi_{j,h}(X_{v,h}^{j-1}, \Xi_{v,h}^j) - X_{v,h}^j \cdot \Xi_{v,h}^j) + \phi_{v+1,h}(X_{v,h}^v, \xi), \end{aligned}$$

where $X_{v,h}^0 = x$ and $\{X_{v,h}^j, \Xi_{v,h}^j\}_{j=1}^v(x, \xi)$ is the assumed solution of (2.1).

REMARK. 1° The condition $(\#)$ is satisfied by the phase functions defined as the solution of some Hamilton-Jacobi equations (see Proposition 4.3 of section 4).

2° Let $\{\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^j\}_{j=1}^v(x, \xi) \equiv \{h^{-\delta} X_{v,h}^j, h^{\rho} \Xi_{v,h}^j\}_{j=1}^v(h^{\delta} x, h^{-\rho} \xi)$. Then $\{\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^j\}_{j=1}^v(x, \xi)$ is the solution of (2.1) with $\phi_{j,h}(x, \xi)$ replaced by $\tilde{\phi}_{j,h}(x, \xi) \equiv h^{\rho-\delta} \phi_{j,h}(h^{\delta} x, h^{-\rho} \xi)$. Thus $\tilde{\phi}_{j,h} \in P_{\rho,\delta}^{[x]}(\tau_j, \sigma_j; h)$ ($j=1, 2, \dots$) satisfy the condition $(\#)$ and we can define $\#$ - $(v+1)$ product $\tilde{\Phi}_{v+1,h} = \tilde{\phi}_{1,h} \# \dots \# \tilde{\phi}_{v+1,h}$ by (2.2) with $\phi_{j,h}$ and $\{X_{v,h}^j, \Xi_{v,h}^j\}_{j=1}^v$ replaced by $\tilde{\phi}_{j,h}$ and $\{\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^j\}_{j=1}^v$. In this case we have the relation $\tilde{\Phi}_{v+1,h}(x, \xi) = h^{\rho-\delta} \Phi_{v+1,h}(h^{\delta} x, h^{-\rho} \xi)$.

We next prepare a technical key lemma.

Lemma 2.2. Let $x^0, x^j, \xi^j, u^j, v^j \in R^n$ and let r_j, s_j, \tilde{s}_j and t_j be $n \times n$ real matrices for $j=1, 2, \dots$ such that

$$(2.3) \quad |r_j| \leq \sigma, \quad |t_j|, |s_j|, |\tilde{s}_j| \leq \tau_j$$

for some $0 \leq \sigma < \infty$ and $0 \leq \tau_j < \infty$ ($j=1, 2, \dots$). Then we have for any integer $v \geq 1$

$$(2.4) \quad \begin{aligned} & |x^1 - (I + \tilde{s}_1)x^0 - r_1 \xi^1 - u^1| \\ &+ \sum_{j=2}^v \{ |x^j - (I + \tilde{s}_j)x^{j-1} - r_j \xi^j - u^j| + |\xi^{j-1} - (I + s_j)\xi^j - t_j x^{j-1} - v^j| \} \\ &+ |\xi^v - (I + s_{v+1})\xi^{v+1} - t_{v+1}x^v - v^{v+1}| \\ &\geq (1 - 2\sigma \bar{\tau}_{v+1} - \bar{\tau}_{v+1}) \sum_{j=1}^v |\xi^j - \xi^{j+1}| - (2\sigma \bar{\tau}_{v+1} + \bar{\tau}_{v+1}) |\xi^{v+1}| \\ &+ (1 - 2\bar{\tau}_{v+1}) \sum_{j=1}^v |x^j - x^{j-1} - r_j \xi^j| - 2\bar{\tau}_{v+1} |x^0| \\ &- \sum_{j=1}^v (|u^j| + |v^{j+1}|), \end{aligned}$$

provided that $\bar{\tau}_{\nu+1} = \sum_{k=1}^{\nu+1} k\tau_k$ and $\bar{\tau}_{\nu+1} = \sum_{k=1}^{\nu+1} \tau_k$ satisfy $0 \leq 2(\sigma \bar{\tau}_{\nu+1} + \bar{\tau}_{\nu+1}) < 1$. The inequality (2.4) also holds for $n \times n$ real matrices x^0, x^j, ξ^j, u^j , and v^j .

Proof. We first observe that the left hand side of (2.4) is bounded from below by $\Xi + X + U$, where

$$(2.5) \quad \begin{cases} \Xi = \sum_{j=2}^{\nu+1} |\xi^{j-1} - \xi^j| - \sum_{j=2}^{\nu+1} \tau_j |\xi^j|, \\ X = \sum_{j=1}^{\nu} |x^j - x^{j-1} - r_j \xi^j| - 2 \sum_{j=1}^{\nu} \tau_j |x^{j-1}| - \tau_{\nu+1} |x^\nu|, \\ U = - \sum_{j=1}^{\nu} (|u^j| + |v^{j+1}|). \end{cases}$$

Put $\tau_{\nu,j} = \tau_\nu + \dots + \tau_j$. Then X is estimated as

$$\begin{aligned} X &= \sum_{j=1}^{\nu} (1 - (\tau_{\nu+1} + 2\tau_{\nu,j+1})) |x^j - x^{j-1} - r_j \xi^j| \\ &\quad + \sum_{j=1}^{\nu} (\tau_{\nu+1} + 2\tau_{\nu,j+1}) |x^j - x^{j-1} - r_j \xi^j| - 2 \sum_{j=1}^{\nu} \tau_j |x^{j-1}| - \tau_{\nu+1} |x^\nu| \\ &\geq (1 - 2\bar{\tau}_{\nu+1}) \sum_{j=1}^{\nu} |x^j - x^{j-1} - r_j \xi^j| + \sum_{j=1}^{\nu} (\tau_{\nu+1} + 2\tau_{\nu,j+1}) |x^j| \\ &\quad - \sum_{j=1}^{\nu} (\tau_{\nu+1} + 2\tau_{\nu,j+1}) (|x^{j-1}| + \sigma |\xi^j|) \\ &\quad - 2 \sum_{j=1}^{\nu} \tau_j |x^{j-1}| - \tau_{\nu+1} |x^\nu| \\ &\geq (1 - 2\bar{\tau}_{\nu+1}) \sum_{j=1}^{\nu} |x^j - x^{j-1} - r_j \xi^j| - 2\sigma \sum_{j=1}^{\nu} \tau_{\nu+1,j+1} |\xi^j| - 2\bar{\tau}_{\nu+1} |x^0|. \end{aligned}$$

Thus we have

$$(2.6) \quad X + \Xi \geq A + B,$$

where

$$(2.7) \quad \begin{cases} A = (1 - 2\bar{\tau}_{\nu+1}) \sum_{j=1}^{\nu} |x^j - x^{j-1} - r_j \xi^j| - 2\bar{\tau}_{\nu+1} |x^0|, \\ B = \sum_{j=2}^{\nu+1} |\xi^{j-1} - \xi^j| - \sum_{j=2}^{\nu+1} \tau_j |\xi^j| - 2\sigma \sum_{j=1}^{\nu} \tau_{\nu+1,j+1} |\xi^j|. \end{cases}$$

Here noting $\sum_{l=1}^{j+1} \tau_{\nu+1,l} \leq \bar{\tau}_{\nu+1}$ for $1 \leq j \leq \nu$, we have

$$\begin{aligned} B &\geq \sum_{j=1}^{\nu} |\xi^j - \xi^{j+1}| - \sum_{j=1}^{\nu} (2\sigma \tau_{\nu+1,j+1} + \tau_j) |\xi^j| - \tau_{\nu+1} |\xi^{\nu+1}| \\ &\geq \sum_{j=1}^{\nu} (1 - (2\sigma \sum_{l=1}^{j+1} \tau_{\nu+1,l} + \bar{\tau}_j)) |\xi^j - \xi^{j+1}| \\ &\quad + \sum_{j=1}^{\nu} (2\sigma \sum_{l=1}^{j+1} \tau_{\nu+1,l} + \bar{\tau}_j) (|\xi^j| - |\xi^{j+1}|) \end{aligned}$$

$$\begin{aligned}
 & - \sum_{j=1}^{\nu} (2\sigma\tau_{\nu+1,j+1} + \tau_j) |\xi^j| - \tau_{\nu+1} |\xi^{\nu+1}| \\
 & \geq \sum_{j=1}^{\nu} (1 - (2\sigma\bar{\tau}_{\nu+1} + \bar{\tau}_{\nu+1})) |\xi^j - \xi^{j+1}| - (2\sigma\bar{\tau}_{\nu+1} + \bar{\tau}_{\nu+1}) |\xi^{\nu+1}|.
 \end{aligned}$$

Combining this with (2.5)–(2.7) proves the lemma. \square

Proposition 2.3. *Let $\phi_{j,h} \in P_{\rho,\delta}^{[k]}(\tau_j, \sigma_j; h)$ ($j=1, 2, \dots$) satisfy the condition (#) for any integer $\nu \geq 1$. Let $\{\tau_j\}_{j=1}^{\infty}$ and $\{\sigma_j\}_{j=1}^{\infty}$ satisfy*

$$(2.8) \quad \begin{cases} 0 \leq \sigma_j \leq \sigma_0 \\ 0 \leq 2(\sigma_0\bar{\tau}_{\infty} + \bar{\tau}_{\infty}) \leq \tau_0 \end{cases}$$

for some $0 \leq \sigma_0 < \infty$ and $0 \leq \tau_0 < 1$, where $\bar{\tau}_{\infty} = \sum_{k=1}^{\infty} \tau_k$ and $\bar{\tau}_{\infty} = \sum_{k=1}^{\infty} k\tau_k$. Let $\{\tilde{X}_{\nu,h}^j\}$

$\{\tilde{\Xi}_{\nu,h}^j\}_{j=1}^{\nu}(x, \xi) = \{h^{-\delta} X_{\nu,h}^j, h^{\sigma} \Xi_{\nu,h}^j\}_{j=1}^{\nu}(h^{\delta} x, h^{-\rho} \xi)$, where $\{X_{\nu,h}^j, \Xi_{\nu,h}^j\}_{j=1}^{\nu}$ is the assumed solution of (2.1). Then the following estimates hold.

i) For any $\nu \geq 1$ there exists a constant $C_{\nu} > 0$ such that for $j=1, \dots, \nu$, $h \in (0, 1)$ and $(x, \xi) \in R^{2n}$

$$(2.9) \quad \begin{cases} |\tilde{X}_{\nu,h}^j - \tilde{X}_{\nu,h}^{j-1}| \leq C_{\nu} \langle \xi \rangle, \\ |\tilde{\Xi}_{\nu,h}^j - \tilde{\Xi}_{\nu,h}^{j+1}| \leq C_{\nu}. \end{cases}$$

ii) For any $\nu \geq 1$, $1 \leq k \leq \nu$, $h \in (0, 1)$ and $(x, \xi) \in R^{2n}$, one has

$$(2.10) \quad \begin{cases} \text{i) } \sum_{j=1}^k |\nabla_x(\tilde{X}_{\nu,h}^j - \tilde{X}_{\nu,h}^{j-1})| \leq \frac{\tau_0}{1-\tau_0} (1 + \bar{\sigma}_k), \quad |\nabla_x \tilde{X}_{\nu,h}^k - I| \leq \frac{\tau_0}{1-\tau_0} (1 + \bar{\sigma}_k), \\ \text{ii) } \sum_{j=1}^k |\nabla_x(\tilde{\Xi}_{\nu,h}^j - \tilde{\Xi}_{\nu,h}^{j+1})| \leq \frac{\tau_0}{1-\tau_0}, \quad |\nabla_x \tilde{\Xi}_{\nu,h}^k| \leq \frac{\tau_0}{1-\tau_0}, \end{cases}$$

and

$$(2.11) \quad \begin{cases} \text{i) } \sum_{j=1}^k |\nabla_{\xi}(\tilde{X}_{\nu,h}^j - \tilde{X}_{\nu,h}^{j-1})| \leq \frac{1 + \bar{\sigma}_k}{1 - \tau_0}, \quad |\nabla_{\xi} \tilde{X}_{\nu,h}^k| \leq \frac{1 + \bar{\sigma}_k}{1 - \tau_0}, \\ \text{ii) } \sum_{j=1}^k |\nabla_{\xi}(\tilde{\Xi}_{\nu,h}^j - \tilde{\Xi}_{\nu,h}^{j+1})| \leq \frac{\tau_0}{1 - \tau_0}, \quad |\nabla_{\xi} \tilde{\Xi}_{\nu,h}^k - I| \leq \frac{\tau_0}{1 - \tau_0}, \end{cases}$$

where $\bar{\sigma}_k = \sigma_1 + \dots + \sigma_k$.

iii) For any α, β satisfying $|\alpha + \beta| \geq 1$, one has

$$(2.12) \quad \begin{aligned} & |\partial_x^{\alpha} \partial_{\xi}^{\beta}(\tilde{X}_{\nu,h}^j - x, \tilde{\Xi}_{\nu,h}^j - \xi)| \\ & \leq C_{\alpha,\beta} \left(\frac{1 + \bar{\sigma}_{\nu}}{1 - \tau_0}\right)^{2|\alpha + \beta| - 1} \left(\sum_{j=1}^{\nu+1} |J_{j,h}|_{3,|\alpha + \beta| - 1}\right)^{|\alpha + \beta| - 1}, \end{aligned}$$

where the constant $C_{\alpha,\beta}$ is independent of $\nu \geq j \geq 1$, $h \in (0, 1)$, $(x, \xi) \in R^{2n}$, $\{\sigma_j\}_{j=1}^{\infty}$, $\{\tau_j\}_{j=1}^{\infty}$, $0 \leq \sigma_0 \leq 1$ and $0 \leq \tau_0 \leq 1$.

Proof. i) Since $\{\tilde{X}_{\nu,h}^j, \tilde{\Xi}_{\nu,h}^j\}_{j=1}^{\nu}$ is the solution of (2.1) with $\phi_{j,h}$ replaced by

$\tilde{\phi}_{j,h}$, we have

$$(2.13) \quad \begin{cases} \tilde{X}_{v,h}^j - \tilde{X}_{v,h}^{j-1} = \nabla_{\xi} \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j), \\ \tilde{\Xi}_{v,h}^j - \tilde{\Xi}_{v,h}^{j+1} = \nabla_x \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}), \end{cases} \quad (j = 1, \dots, \nu)$$

where $\tilde{X}_{v,h}^0 = x$ and $\tilde{\Xi}_{v,h}^{\nu+1} = \xi$. From this and (1.2) we have (2.9).

ii) Differentiating (2.13) we have for $j=1, \dots, \nu$

$$(2.14) \quad \begin{cases} \nabla_x \tilde{X}_{v,h}^j = (I + \tilde{\nabla}_x \nabla_{\xi} \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j)) \cdot \nabla_x \tilde{X}_{v,h}^{j-1} \\ \quad + \tilde{\nabla}_{\xi} \nabla_{\xi} \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j) \cdot \nabla_x \tilde{\Xi}_{v,h}^j, \\ \nabla_x \tilde{\Xi}_{v,h}^j = (I + (\tilde{\nabla}_{\xi} \nabla_x \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}))) \cdot \nabla_x \tilde{\Xi}_{v,h}^{j+1} \\ \quad + \tilde{\nabla}_x \nabla_x \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}) \cdot \nabla_x \tilde{X}_{v,h}^j \end{cases}$$

and

$$(2.15) \quad \begin{cases} \nabla_{\xi} \tilde{X}_{v,h}^j = (I + \tilde{\nabla}_x \nabla_{\xi} \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j)) \cdot \nabla_{\xi} \tilde{X}_{v,h}^{j-1} \\ \quad + \tilde{\nabla}_{\xi} \nabla_{\xi} \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j) \cdot \nabla_{\xi} \tilde{\Xi}_{v,h}^j, \\ \nabla_{\xi} \tilde{\Xi}_{v,h}^j = (I + \tilde{\nabla}_{\xi} \nabla_x \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1})) \cdot \nabla_{\xi} \tilde{\Xi}_{v,h}^{j+1} \\ \quad + \tilde{\nabla}_x \nabla_x \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}) \cdot \nabla_{\xi} \tilde{X}_{v,h}^j. \end{cases}$$

Writing $y^j = \nabla_x \tilde{X}_{v,h}^j$, $\eta^j = \nabla_x \tilde{\Xi}_{v,h}^j$ and putting $\tilde{s}_j = \tilde{\nabla}_x \nabla_{\xi} \tilde{J}_{j,h}$, $s_{j+1} = \tilde{\nabla}_{\xi} \nabla_x \tilde{J}_{j+1,h}$, $t_{j+1} = \tilde{\nabla}_x \nabla_x \tilde{J}_{j+1,h}$ and $r_j = \tilde{\nabla}_{\xi} \nabla_{\xi} \tilde{J}_{j,h}$, we can rewrite (2.14) as

$$(2.16) \quad \begin{cases} y^j = (I + \tilde{s}_j) y^{j-1} + r_j \eta^j, \\ \eta^j = (I + s_{j+1}) \eta^{j+1} + t_{j+1} y^j, \end{cases} \quad (j = 1, \dots, \nu)$$

where $y^0 = I$ and $\eta^{\nu+1} = 0$. Since \tilde{s}_j , s_j , t_j and r_j satisfy $|\tilde{s}_j|$, $|s_j|$, $|t_j| \leq \tau_j$ and $|r_j| \leq \sigma_j \leq \sigma_0$, we can apply Lemma 2.2 and we have

$$(2.17) \quad 0 \geq (1 - \tau_0) \sum_{j=1}^{\nu} (|\eta^j - \eta^{j+1}| + |y^j - y^{j-1} - r_j \eta^j|) - \tau_0 |\eta^{\nu+1}| - \tau_0 |y^0|,$$

hence

$$(2.18) \quad \sum_{j=1}^{\nu} (|\eta^j - \eta^{j+1}| + |y^j - y^{j-1} - r_j \eta^j|) \leq \frac{\tau_0}{1 - \tau_0}.$$

Thus we obtain (2.10)-ii). So using (2.18) again we obtain for $k=1, \dots, \nu$,

$$(2.19) \quad \begin{aligned} \sum_{j=1}^k |y^j - y^{j-1}| &\leq \sum_{j=1}^k (|y^j - y^{j-1} - r_j \eta^j| + \sigma_j |\eta^j|) \\ &\leq \frac{\tau_0}{1 - \tau_0} + \sum_{j=1}^k \sigma_j \sum_{l=j}^{\nu} |\eta^l - \eta^{l+1}| \\ &\leq \frac{\tau_0}{1 - \tau_0} (1 + \sigma_k), \end{aligned}$$

which proves (2.10).

We next prove (2.11). Under obvious notations, (2.15) is rewritten as (2.16) with $y^0=0$ and $\eta^{v+1}=I$. Applying Lemma 2.2 we obtain (2.17) hence (2.18), from which we get (2.11)-ii). So using (2.18) again we have for $k=1, \dots, \nu$

$$(2.20) \quad \begin{aligned} \sum_{j=1}^k |y^j - y^{j-1}| &\leq \frac{\tau_0}{1-\tau_0} + \sum_{j=1}^k \sigma_j \left(\sum_{l=j}^{\nu} |\eta^l - \eta^{l+1}| + |\eta^{v+1}| \right) \\ &\leq \frac{\tau_0 + \bar{\sigma}_k}{1-\tau_0} \leq \frac{1 + \bar{\sigma}_k}{1-\tau_0}, \end{aligned}$$

which proves (2.11).

iii) For any multi-indices α, β with $|\alpha + \beta| \geq 1$, differentiating (2.15) we have for $j=1, \dots, \nu$

$$(2.21) \quad \begin{cases} \partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{X}_{v,h}^j = (I + \vec{\nabla}_x \nabla_{\xi} \tilde{J}_{j,h}) \cdot \partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{X}_{v,h}^{j-1} \\ \quad + \vec{\nabla}_{\xi} \nabla_{\xi} \tilde{J}_{j,h} \cdot \partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{\Xi}_{v,h}^j + U_j, \\ \partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{\Xi}_{v,h}^j = (I + \vec{\nabla}_{\xi} \nabla_x \tilde{J}_{j+1,h}) \cdot \partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{\Xi}_{v,h}^{j+1} \\ \quad + \vec{\nabla}_x \nabla_x \tilde{J}_{j+1,h} \cdot \partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{X}_{v,h}^j + V_{j+1}, \end{cases}$$

where U_j and V_j are the polynomial of $\partial_{\xi}^{\gamma} \partial_x^{\delta} \tilde{J}_{j,h} (3 \leq |\gamma + \delta| \leq |\alpha + \beta| + 2)$ and $\partial_{\xi}^{\gamma'} \partial_x^{\delta'} \tilde{X}_{v,h}^{j-1}, \partial_{\xi}^{\gamma'} \partial_x^{\delta'} \tilde{\Xi}_{v,h}^j (1 \leq |\gamma' + \delta'| \leq |\alpha + \beta| + 2)$ of order $|\alpha + \beta| + 2$; especially the orders of $\partial_x \tilde{X}_{v,h}^{j-1}, \partial_{\xi} \tilde{X}_{v,h}^{j-1}, \partial_x \tilde{\Xi}_{v,h}^j$ and $\partial_{\xi} \tilde{\Xi}_{v,h}^j$ are at most $|\alpha + \beta| + 1$ and the order of $\partial_{\xi}^{\gamma} \partial_x^{\delta} \tilde{J}_{j,h}$ is at most 1. Moreover the sum of $|\gamma + \delta|$ of $\partial_{\xi}^{\gamma} \partial_x^{\delta} \tilde{X}_{v,h}^{j-1}$ and $\partial_{\xi}^{\gamma} \partial_x^{\delta} \tilde{\Xi}_{v,h}^j (1 \leq |\gamma + \delta| \leq |\alpha + \beta|)$ in every term of U_j and V_j does not exceed $|\alpha + \beta| + 1$. Similar results hold for the differentials of (2.14). Thus using Lemma 2.2 and ii) we obtain by induction

$$(2.22) \quad \begin{aligned} &|\partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{X}_{v,h}^j|, |\partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_{\xi} \tilde{\Xi}_{v,h}^j|, |\partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_x \tilde{X}_{v,h}^j|, |\partial_{\xi}^{\alpha} \partial_x^{\beta} \nabla_x \tilde{\Xi}_{v,h}^j| \\ &\leq C_{\alpha,\beta} \left(\sum_{j=1}^{\nu+1} |J_{j,h}|_{3,|\alpha+\beta|} \right)^{|\alpha+\beta|} \left(\frac{1 + \bar{\sigma}_{\nu}}{1 - \tau_0} \right)^{2|\alpha+\beta|+1}. \quad \square \end{aligned}$$

Proposition 2.4. Let $\phi_{j,h} \in P_{\rho,\delta}^{[x]}(\tau_j, \sigma_j; h) (j=1, 2, \dots)$ satisfy the condition (#) for any integer $\nu \geq 1$. Let $\{\tau_j\}_{j=1}^{\infty}$ and $\{\sigma_j\}_{j=1}^{\infty}$ satisfy

$$(2.23) \quad \begin{cases} 0 \leq \sigma_j \leq \sigma_0, \\ 0 \leq 2(\sigma_0 \bar{\tau}_{\infty} + \bar{\tau}_{\infty}) \leq \tau_0 \end{cases}$$

for some $0 \leq \sigma_0 < \infty$ and $0 \leq \tau_0 \leq 1/4$. Let $\Phi_{\nu+1,h} = \phi_{1,h} \# \dots \# \phi_{\nu+1,h}$ be defined by (2.2). Then $\Phi_{\nu+1,h}$ satisfies the following properties:

i) We have

$$(2.24) \quad \Phi_{\nu+1,h} \in P_{\rho,\delta}^{[x]}(3\tau_0, \frac{5}{3}(1 + \bar{\sigma}_{\nu+1}); h)$$

for any $\nu \geq 1$.

ii) We have for any $\nu \geq 1$

$$(2.25) \quad \begin{cases} \nabla_x \Phi_{v+1,h}(x, \xi) = \nabla_x \phi_{1,h}(x, \Xi_{v,h}^1), \\ \nabla_h \Phi_{v+1,h}(x, \xi) = \nabla_\xi \phi_{v+1,h}(X_{v,h}^v, \xi) \end{cases}$$

and

$$(2.26) \quad \begin{cases} \Delta_x J_{v+1,h}(x, \xi) = \nabla_x J_{1,h}(x, \Xi_{v,h}^1) + \Xi_{v,h}^1 - \xi, \\ \nabla_\xi J_{v+1,h}(x, \xi) = \nabla_\xi J_{v+1,h}(X_{v,h}^v, \xi) + X_{v,h}^v - x, \end{cases}$$

where $J_{v+1,h}(x, \xi) \equiv \Phi_{v+1,h}(x, \xi) - x \cdot \xi$.

Proof. Since the properties except (2.24) can be proved without using (2.24) in quite the same way as in the proof of Theorem 2.7 of [6], we only prove (2.24). Since

$$(2.27) \quad \begin{cases} \nabla_x \tilde{J}_{v+1,h}(x, \xi) = \nabla_x \tilde{J}_{1,h}(x, \tilde{\Xi}_{v,h}^1) + \tilde{\Xi}_{v,h}^1 - \xi, \\ \nabla_\xi \tilde{J}_{v+1,h}(x, \xi) = \nabla_\xi \tilde{J}_{v+1,h}(X_{v,h}^v, \xi) + X_{v,h}^v - x, \end{cases}$$

we easily see from (2.9) that

$$(2.28) \quad \sup_{h,x,\xi} \{ |\nabla_\xi \tilde{J}_{v+1,h}(x, \xi)| \langle \xi \rangle + |\nabla_x \tilde{J}_{v+1,h}(x, \xi)| \} < \infty.$$

From (2.27) we have

$$(2.29) \quad \begin{cases} \nabla_x \nabla_x \tilde{J}_{v+1,h}(x, \xi) = \nabla_x \nabla_x \tilde{J}_{1,h}(x, \tilde{\Xi}_{v,h}^1) \\ \qquad \qquad \qquad \qquad \qquad + (\nabla_\xi \nabla_x \tilde{J}_{1,h}(x, \tilde{\Xi}_{v,h}^1) + I) \cdot \nabla_x \tilde{\Xi}_{v,h}^1, \\ \nabla_\xi \nabla_x \tilde{J}_{v+1,h}(x, \xi) = (\nabla_\xi \nabla_x \tilde{J}_{1,h}(x, \tilde{\Xi}_{v,h}^1) + I) \cdot \nabla_\xi \tilde{\Xi}_{v,h}^1 - I, \\ \nabla_\xi \nabla_\xi \tilde{J}_{v+1,h}(x, \xi) = (\nabla_x \nabla_\xi \tilde{J}_{v+1,h}(X_{v,h}^v, \xi) + I) \cdot \nabla_\xi X_{v,h}^v \\ \qquad \qquad \qquad \qquad \qquad + \nabla_\xi \nabla_\xi \tilde{J}_{v+1,h}(X_{v,h}^v, \xi). \end{cases}$$

Thus from (2.10) and (2.11) we get

$$(2.30) \quad \left\{ \begin{aligned} |\nabla_x \nabla_x \tilde{J}_{v+1,h}(x, \xi)| &\leq \tau_1 + (\tau_1 + 1) \frac{\tau_0}{1 - \tau_0} \leq 3\tau_0, \\ |\nabla_\xi \nabla_x \tilde{J}_{v+1,h}(x, \xi)| &\leq (\tau_1 + 1) \frac{\tau_0}{1 - \tau_0} + \tau_1 \leq 3\tau_0, \\ |\nabla_\xi \nabla_\xi \tilde{J}_{v+1,h}(x, \xi)| &\leq (\tau_{v+1} + 1) \frac{1 + \bar{\sigma}_v}{1 - \tau_0} + \sigma_{v+1} \\ &\leq \frac{1 + \tau_0}{1 - \tau_0} (1 + \bar{\sigma}_{v+1}) \leq \frac{5}{3} (1 + \bar{\sigma}_{v+1}). \end{aligned} \right.$$

Differentiating (2.29) further and using (2.10)–(2.12) we can easily see that (1.3) holds for our $J_{v+1,h}$. Thus we have proved (2.24). \square

Before stating a theorem concerning the calculus of conjugate Fourier integral operators we prepare a lemma.

Lemma 2.5. Let $\phi_{j,h} \in P_{\rho,\xi}^{[x]}(\tau_j, \sigma_j; h)$ ($j=1, 2, \dots$) satisfy the condition (#) for any $\nu \geq 1$, and assume that $0 \leq \sigma_j \leq \sigma_0$ ($j=1, 2, \dots$) and $0 \leq 2(\sigma_0 \bar{\sigma}_\infty + \bar{\sigma}_\infty) \leq \tau_0$ for some $0 \leq \sigma_0 < \infty$ and $0 \leq \tau_0 \leq 1/4$. Let $\nu \geq 1$, and let $\{\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^j\}_{j=1}^\nu(x, \xi)$ and $\tilde{\Phi}_{\nu+1,h}$ be defined as in 2° of the remark after Definition 2.1. Define $\tilde{\varphi}_h$ by

$$(2.31) \quad \begin{aligned} & \tilde{\varphi}_h(y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu; x, \xi) \\ &= \sum_{j=1}^\nu (\tilde{\Phi}_{j,h}(\tilde{X}_{v,h}^{j-1}(x, \xi) + y^{j-1}, \tilde{\Xi}_{v,h}^j + \eta^j) - \tilde{X}_{j,h}^j(x, \xi) \cdot \tilde{\Xi}_{v,h}^j(x, \xi)) \\ & \quad + \tilde{\Phi}_{\nu+1,h}(\tilde{X}_{v,h}^\nu(x, \xi) + y^\nu, \xi) - \tilde{\Phi}_{\nu+1,h}(x, \xi) \end{aligned}$$

with $y^0=0$ and $\tilde{X}_{v,h}^0=x$. Then the following estimates hold:

i) For any $y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu, x, \xi \in R^n, \sigma \in R^1, h \in (0, 1)$, and $\nu \geq 1$

$$(2.32) \quad \begin{aligned} & 2\nu + h^{-2\sigma} \sum_{j=1}^\nu (|\nabla_{y^j} \tilde{\varphi}_h|^2 + |\nabla_{\eta^j} \tilde{\varphi}_h|^2) \\ & \geq \frac{(1-\tau_0)^2}{4\nu} \left\{ \sum_{j=1}^\nu (2 + h^{-\sigma} |\eta^j - \eta^{j+1}| + h^{-\sigma} |y^j - y^{j-1} - r_j \eta^j|) \right\}^2 \\ & \geq \nu(1-\tau_0)^2 \prod_{j=1}^\nu \langle h^{-\sigma} (\eta^j - \eta^{j+1}) \rangle^{1/\nu} \langle h^{-\sigma} (y^j - y^{j-1} - r_j \eta^j) \rangle^{1/\nu}, \end{aligned}$$

where $r_j = \vec{\nabla}_\xi \nabla_\xi \tilde{J}_{j,h}(\tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1}, \tilde{\Xi}_{v,h}^j + \eta^j)$.

ii) For any $y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu, x, \xi \in R^n, h \in (0, 1)$, and for any multi-indices $\alpha, \beta, \alpha^1, \beta^1, \dots, \alpha^\nu, \beta^\nu$ and any integer j with $1 \leq j \leq \nu$, we have

$$(2.33) \quad \begin{aligned} & |\partial_\xi^\alpha \partial_x^\beta \partial_\eta^{\alpha^1} \partial_{y^1}^{\beta^1} \dots \partial_{\eta^\nu}^{\alpha^\nu} \partial_{y^\nu}^{\beta^\nu} \nabla_{y^j} \tilde{\varphi}_h| \\ & \leq C_{\alpha, \beta, \alpha^j+1, \beta^j} \left(\frac{1+\bar{\sigma}_y}{1-\tau_0} \right)^{(2|\alpha+\beta|-1)\vee 0} \times \\ & \quad \times \left(1 + \sum_{k=1}^{j+1} |J_{k,h}|_{3, (|\alpha+\beta|-1)\vee 0} \right)^{|\alpha+\beta|} \times \\ & \quad \times \left(1 + |J_{j+1,h}|_{2, |\alpha+\beta+\alpha^j+1+\beta^j|} \right) \langle y^j; \eta^j; \eta^{j+1} \rangle \end{aligned}$$

and

$$(2.34) \quad \begin{aligned} & |\partial_\xi^\alpha \partial_x^\beta \partial_\eta^{\alpha^1} \partial_{y^1}^{\beta^1} \dots \partial_{\eta^\nu}^{\alpha^\nu} \partial_{y^\nu}^{\beta^\nu} \nabla_{\eta^j} \tilde{\varphi}_h| \\ & \leq C_{\alpha, \beta, \alpha^j, \beta^j-1} \left(\frac{1+\bar{\sigma}_y}{1-\tau_0} \right)^{(|\alpha+\beta|-1)\vee 0} \times \\ & \quad \times \left(1 + \sum_{k=1}^{j+1} |J_{k,h}|_{3, (|\alpha+\beta|-1)\vee 0} \right)^{|\alpha+\beta|} \times \\ & \quad \times \left(1 + |J_{j,h}|_{2, |\alpha+\beta+\alpha^j+\beta^j-1|} \right) \langle y^{j-1}; y^j; \eta^j \rangle, \end{aligned}$$

where $a \vee b = \max(a, b)$ for $a, b \in R^1; y^0 = \eta^{\nu+1} = 0$; and $\alpha^{\nu+1} = \beta^0 = 0$.

iii) For any $y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu, x, \xi \in R^n, h \in (0, 1)$ and any multi-indices α, β with $|\alpha+\beta| \geq 1$, we have

$$(2.35) \quad \begin{aligned} & |\partial_\xi^\alpha \partial_x^\beta \tilde{\varphi}_h| \\ & \leq C_{\alpha, \beta} \left(\frac{1+\bar{\sigma}_y}{1-\tau_0} \right)^{2|\alpha+\beta|-1} \left(1 + \sum_{k=1}^{j+1} |J_{k,h}|_{3, |\alpha+\beta|-1} \right)^{|\alpha+\beta|} \times \end{aligned}$$

$$\times \left\{ \sum_{j=1}^{\nu} (|y^j| + |\eta^j|) \right\}^2.$$

Proof. i) We can rewrite (2.31) using (2.2) as

$$\begin{aligned} & \Phi_h(y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu; x, \xi) \\ (2.36) \quad &= \sum_{j=1}^{\nu} \{ (\tilde{\phi}_{j,h}(\tilde{X}_{v,h}^{j-1} + y^{j-1}, \tilde{\Xi}_{v,h}^j + \eta^j) - \tilde{\phi}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j)) \\ & \quad - ((\tilde{X}_{v,h}^j + y^j) \cdot (\tilde{\Xi}_{v,h}^j + \eta^j) - \tilde{X}_{v,h}^j \cdot \tilde{\Xi}_{v,h}^j) \} \\ & \quad + (\tilde{\phi}_{\nu+1,h}(\tilde{X}_{v,h}^\nu + y^\nu, \xi) - \tilde{\phi}_{\nu+1,h}(\tilde{X}_{v,h}^\nu, \xi)). \end{aligned}$$

From this we have for $j=1, \dots, \nu$

$$(2.37) \quad \begin{cases} \nabla_y \tilde{\phi}_h = -\eta^j + (I + \vec{\nabla}_\xi \nabla_x \tilde{J}_{j+1,h}(\tilde{\Xi}_{v,h}^{j+1}, \tilde{X}_{v,h}^j + y^j, \tilde{\Xi}_{v,h}^{j+1} + \eta^{j+1})) \cdot \eta^{j+1} \\ \quad + \vec{\nabla}_x \nabla_x \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}, \tilde{X}_{v,h}^j + y^j) \cdot y^j, \\ \nabla_{\eta^j} \tilde{\phi}_h = -y^j + (I + \vec{\nabla}_x \nabla_\xi \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1})) \cdot y^{j-1} \\ \quad + \vec{\nabla}_\xi \nabla_\xi \tilde{J}_{j,h}(\tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1}, \tilde{\Xi}_{v,h}^j + \eta^j) \cdot \eta^j, \end{cases}$$

where $y^0 = \eta^{\nu+1} = 0$, $\tilde{X}_{v,h}^0 = x$ and $\tilde{\Xi}_{v,h}^{\nu+1} = \xi$. Putting $s_j = \vec{\nabla}_\xi \nabla_x \tilde{J}_{j,h}$, $\tilde{s}_j = \vec{\nabla}_x \nabla_\xi \tilde{J}_{j,h}$, $t_j = \vec{\nabla}_x \nabla_x \tilde{J}_{j,h}$, and $r_j = \vec{\nabla}_\xi \nabla_\xi \tilde{J}_{j,h}$, we have

$$(2.38) \quad \begin{cases} \nabla_{y^j} \tilde{\phi}_h = -\eta^j + (I + s_{j+1})\eta^{j+1} + t_{j+1}y^j, \\ \nabla_{\eta^j} \tilde{\phi}_h = -y^j + (I + \tilde{s}_j)y^{j-1} + r_j\eta^j, \end{cases} \quad (j = 1, \dots, \nu, y^0 = \eta^{\nu+1} = 0)$$

and

$$(2.39) \quad |s_j|, |\tilde{s}_j|, |t_j| \leq \tau_j, \quad |r_j| \leq \sigma_j.$$

Then applying Lemma 2.2 we have

$$\begin{aligned} & 4\nu[2\nu + h^{-2\sigma} \sum_{j=1}^{\nu} (|\nabla_{y^j} \tilde{\phi}_h|^2 + |\nabla_{\eta^j} \tilde{\phi}_h|^2)] \\ (2.40) \quad & \geq 2[(2\nu)^2 + \left\{ \sum_{j=1}^{\nu} (|h^{-\sigma} \nabla_{y^j} \tilde{\phi}_h| + |h^{-\sigma} \nabla_{\eta^j} \tilde{\phi}_h|) \right\}^2] \\ & \geq \{2\nu + \sum_{j=1}^{\nu} (|h^{-\sigma} \nabla_{y^j} \tilde{\phi}_h| + |h^{-\sigma} \nabla_{\eta^j} \tilde{\phi}_h|)\}^2 \\ & \geq \{(1 - \tau_0) \sum_{j=1}^{\nu} (2 + h^{-\sigma} |\eta^j - \eta^{j+1}| + h^{-\sigma} |y^j - y^{j-1} - r_j \eta^j|)\}^2 \\ & \geq \{2\nu(1 - \tau_0) \prod_{j=1}^{\nu} \langle h^{-\sigma} (\eta^j - \eta^{j+1}) \rangle^{1/(2\nu)} \langle h^{-\sigma} (y^j - y^{j-1} - r_j \eta^j) \rangle^{1/(2\nu)}\}^2, \end{aligned}$$

which proves (2.32).

ii) is easily seen from (2.37) and Proposition 2.3-iii).

iii) Using Taylor's expansion formula of order two we see from (2.36) and (2.1) that

$$\Phi_h(y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu; x, \xi)$$

$$\begin{aligned}
 (2.41) \quad &= \sum_{j=1}^{\nu} \{ \vec{\nabla}_x^2 \tilde{\phi}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}, \tilde{X}_{v,h}^j + y^j) y^j \cdot y^j \\
 &\quad + \vec{\nabla}_\xi^2 \tilde{\phi}_{j,h}(\tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1}, \tilde{\Xi}_{v,h}^j + \eta^j) \eta^j \cdot \eta^j \\
 &\quad + \vec{\nabla}_x \nabla_\xi \tilde{\phi}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1}) y^{j-1} \cdot \eta^j - y^j \cdot \eta^j \} ,
 \end{aligned}$$

where $\tilde{\Xi}_{v,h}^{\nu+1} = \xi$, $\tilde{X}_{v,h}^0 = x$, $y^0 = 0$, and

$$\begin{aligned}
 (2.42) \quad &\left\{ \begin{aligned} \vec{\nabla}_x^2 f(x, \xi, y) &= \int_0^1 (1-\theta) \vec{\nabla}_x \nabla_x f(x + \theta(y-x), \xi) d\theta , \\ \vec{\nabla}_\xi^2 f(\xi, x, \eta) &= \int_0^1 (1-\theta) \vec{\nabla}_\xi \nabla_\xi f(x, \xi + \theta(\eta-\xi)) d\theta \end{aligned} \right.
 \end{aligned}$$

for any C^2 -function $f(x, \xi)$. If we use $\tilde{J}_{j,h}$ we get another expression of $\tilde{\phi}_h$:

$$\begin{aligned}
 (2.43) \quad &\tilde{\phi}_h(y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu; x, \xi) \\
 &= \sum_{j=1}^{\nu} \{ \vec{\nabla}_x^2 \tilde{J}_{j+1,h}(\tilde{X}_{v,h}^j, \tilde{\Xi}_{v,h}^{j+1}, \tilde{X}_{v,h}^j + y^j) y^j \cdot y^j \\
 &\quad + \vec{\nabla}_\xi^2 \tilde{J}_{j,h}(\tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1}, \tilde{\Xi}_{v,h}^j + \eta^j) \eta^j \cdot \eta^j \\
 &\quad + \vec{\nabla}_x \nabla_\xi \tilde{J}_{j,h}(\tilde{X}_{v,h}^{j-1}, \tilde{\Xi}_{v,h}^j, \tilde{X}_{v,h}^{j-1} + y^{j-1}) y^{j-1} \cdot \eta^j \\
 &\quad + (y^{j-1} - y^j) \cdot \eta^j \} .
 \end{aligned}$$

From this and Proposition 2.3-iii) we obtain for $|\alpha + \beta| \geq 1$

$$\begin{aligned}
 (2.44) \quad &|\partial_\xi^\alpha \partial_x^\beta \tilde{\phi}_h| \\
 &\leq C_{\alpha,\beta} \left(\sum_{j=1}^{\nu+1} |\tilde{J}_{j,h}|_{3,|\alpha+\beta|-1} \right) \left(\frac{1 + \bar{\sigma}_\nu}{1 - \tau_0} \right)^{2|\alpha+\beta|-1} \times \\
 &\quad \times \left(1 + \sum_{k=1}^{\nu+1} |J_{k,h}|_{3,|\alpha+\beta|-1} \right)^{|\alpha+\beta|-1} (|y^j|^2 + |\eta^j|^2 + |y^{j-1}| |\eta^j|) \\
 &\leq C_{\alpha,\beta} \left(\frac{1 + \bar{\sigma}_\nu}{1 - \tau_0} \right)^{2|\alpha+\beta|-1} \left(1 + \sum_{k=1}^{\nu+1} |J_{k,h}|_{3,|\alpha+\beta|-1} \right)^{|\alpha+\beta|} \times \\
 &\quad \times \sum_{j=1}^{\nu} (|y^j|^2 + |\eta^j|^2) ,
 \end{aligned}$$

which proves iii). \square

Now we can state and prove the main result of this section.

Theorem 2.6. *Let $\phi_{j,h} \in P_{\rho,\delta}^{[x]}(\tau_j, \sigma_j; h)$ ($j=1, 2, \dots$) satisfy the condition (#) for any $\nu \geq 1$, and assume that $0 \leq \sigma_j \leq \sigma_0$ ($j=1, 2, \dots$) and $0 \leq 2(\sigma_0 \bar{\tau}_\infty + \bar{\tau}_\infty) \leq \tau_0$ for some $0 \leq \sigma_0 < \infty$ and $0 \leq \tau_0 \leq 1/4$. Let $\nu \geq 1$ and put $\Phi_{\nu+1,h} = \phi_{1,h} \# \dots \# \phi_{\nu+1,h}$. Let $p_{j,h}(\xi, x') \in B_{\rho,\delta}^{m_j}(h)$ for $j=1, \dots, \nu+1$. Then there exists a symbol $r_{\nu+1,h}(\xi, x') \in B_{\rho,\delta}^{\bar{m}_{\nu+1}}(h)$ ($\bar{m}_{\nu+1} = m_1 + \dots + m_{\nu+1}$) such that*

$$(2.45) \quad P_{\nu+1,h}(\phi_{\nu+1,h}^*) \cdots P_{1,h}(\phi_{1,h}^*) = R_{\nu+1,h}(\Phi_{\nu+1,h}^*)$$

and

$$\begin{aligned}
 & |r_{\nu+1,h}|^{(\bar{m}_{\nu+1})} \\
 (2.46) \quad & \leq C_l c_0^{6\nu n+8l+3} \prod_{m=1}^l (4\nu n+4l+1+m) \cdot \nu^{14\nu n+2\nu+21l+7} \times \\
 & \times (1+\sigma_\nu)^{6\nu n+16l+3} (1-\tau_0)^{-8\nu n-16l-4} (1+\sum_{s=1}^{\nu+1} |J_{s,h}|_{3,l})^{3l} \times \\
 & \times (1+\max_{1 \leq j \leq \nu+1} |J_{j,h}|_{2,2\nu n+3l+1})^{6\nu n+8l+3} \prod_{j=1}^{\nu+1} |p_{j,h}|_{2\nu n+3l+1}^{(m_j)}
 \end{aligned}$$

for any integer $l \geq 0$, where $P_{j,h}(\phi_{j,h}^*) = p_{j,h}(\phi_{j,h}^*; D_x, X')$; $R_{\nu-1,h}(\Phi_{\nu+1,h}^*) = r_{\nu+1,h}(\Phi_{\nu+1,h}^*; D_x, X')$; and $c_0 > 1$ is a constant.

Proof. We can write formally for $f \in \mathcal{S}$

$$\begin{aligned}
 & P_{\nu+1,h}(\phi_{\nu+1,h}^*) \cdots P_{1,h}(\phi_{1,h}^*) f(x^{\nu+1}) \\
 & = 0_s - \iint e^{i(x^{\nu+1}, \xi^{\nu+1} - \Phi_{\nu+1,h}(x^0, \xi^{\nu+1}))} r_{\nu+1,h}(\xi^{\nu+1}, x^0) \times \\
 & \quad \times f(x^0) dx^0 d\xi^{\nu+1},
 \end{aligned}$$

where

$$\begin{aligned}
 & r_{\nu+1,h}(\xi^{\nu+1}, x^0) \\
 & = 0_s - \int \cdots \int \exp \frac{1}{i} \left[\sum_{j=1}^{\nu} (\phi_{j,h}(x^{j-1}, \xi^j) - x^j \cdot \xi^j) \right. \\
 & \quad \left. + \phi_{\nu+1,h}(x^\nu, \xi^{\nu+1}) - \Phi_{\nu+1,h}(x^0, \xi^{\nu+1}) \right] \times \\
 & \quad \times p_{\nu+1,h}(\xi^{\nu+1}, x) \cdots p_{1,h}(\xi^1, x^0) dx^1 \cdots dx^\nu d\xi^1 \cdots d\xi^\nu.
 \end{aligned}$$

So we get (2.45) by limit process as in the proof of Proposition 1.4 if we show that $r_{\nu+1,h}(\xi^{\nu+1}, x^0)$ is well-defined as an oscillatory integral and satisfies (2.46). Set

$$\begin{aligned}
 (2.47) \quad & \tilde{r}_{\nu+1,h}(\xi^{\nu+1}, x^0) \equiv r_{\nu+1,h}(h^{-\rho} \xi^{\nu+1}, h^\delta x^0) \\
 & = h^{-2\nu n} 0_s - \int \cdots \int \exp \frac{h^{-2\sigma}}{i} \left[\sum_{j=1}^{\nu} (\tilde{\phi}_{j,h}(x^{j-1}, \xi^j) - x^j \cdot \xi^j) \right. \\
 & \quad \left. + \tilde{\phi}_{\nu+1,h}(x^\nu, \xi^{\nu+1}) - \tilde{\Phi}_{\nu+1,h}(x^0, \xi^{\nu+1}) \right] \times \\
 & \quad \times \tilde{p}_{\nu+1,h}(\xi^{\nu+1}, x^\nu) \cdots \tilde{p}_{1,h}(\xi^1, x^0) dx^1 \cdots dx^\nu d\xi^1 \cdots d\xi^\nu,
 \end{aligned}$$

where $\tilde{\phi}_{j,h}(x, \xi) = h^{\rho-\delta} \phi_{j,h}(h^\delta x, h^{-\rho} \xi)$, $\tilde{\Phi}_{\nu+1,h}(x, \xi) = h^{\rho-\delta} \Phi_{\nu+1,h}(h^\delta x, h^{-\rho} \xi)$, $\tilde{p}_{j,h}(\xi, x) = p_{j,h}(h^{-\rho} \xi, h^\delta x) \in B_{0,0}^{m_j}(h)$, and $\sigma = (\rho - \delta)/2$. So since $|\tilde{r}_{\nu+1,h}|^{(\bar{m}_{\nu+1})}$ (in $B_{0,0}^{\bar{m}_{\nu+1}}(h)$) $= |r_{\nu+1,h}|^{(\bar{m}_{\nu+1})}$ (in $B_{\rho,\delta}^{\bar{m}_{\nu+1}}(h)$), we have to prove $\tilde{r}_{\nu+1,h}(\xi, x) \in B_{0,0}^{\bar{m}_{\nu+1}}(h)$ and to estimate $|\tilde{r}_{\nu+1,h}|^{(\bar{m}_{\nu+1})}$ in $B_{0,0}^{\bar{m}_{\nu+1}}(h)$.

In (2.47) making a change of variables:

$$\begin{cases} x^j = \tilde{X}_{\nu,h}^j(x^0, \xi^{\nu+1}) + y^j \equiv h^{-\delta} X_{\nu,h}^j(h^\delta x^0, h^{-\rho} \xi^{\nu+1}) + y^j, \\ \xi^j = \tilde{\Xi}_{\nu,h}^j(x^0, \xi^{\nu+1}) + \eta^j \equiv h^\rho \Xi_{\nu,h}^j(h^\delta x^0, h^{-\rho} \xi^{\nu+1}) + \eta^j \end{cases}$$

for $j=1, \dots, \nu$, we get

$$\begin{aligned}
 & \tilde{r}_{\nu+1,h}(\xi, x) \\
 (2.48) \quad &= h^{-2\nu n\sigma} 0_s - \int \dots \int e^{-ih^{-2\sigma}\tilde{\varphi}_h(y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu; x, \xi)} \times \\
 & \quad \times \prod_{j=1}^{\nu+1} q_{j,h}(\xi, x; \eta^j, y^{j-1}) dy^1 \dots dy^\nu d\eta^1 \dots d\eta^\nu,
 \end{aligned}$$

where $\tilde{\varphi}_h$ is defined by (2.31) in Lemma 2.5 and

$$\begin{aligned}
 (2.49) \quad & q_{j,h}(\xi, x; \eta^j, y^{j-1}) \\
 &= \tilde{p}_{j,h}(\tilde{\Xi}_{j,h}^j(x, \xi) + \eta^j, \tilde{X}_{j,h}^{j-1}(x, \xi) + y^{j-1})
 \end{aligned}$$

for $j=1, \dots, \nu+1$ with $y^0 = \eta^{\nu+1} = 0$, $\tilde{\Xi}_{\nu,h}^{\nu+1} = \xi$ and $\tilde{X}_{\nu,h}^0 = x$. Now setting

$$\begin{cases}
 \Gamma_h = 2\nu + h^{-2\sigma} \sum_{j=1}^{\nu} (|\nabla_{y^j} \tilde{\varphi}_h|^2 + |\nabla_{\eta^j} \tilde{\varphi}_h|^2), \\
 L_h = \Gamma_h^{-1} \{2\nu - i \sum_{j=1}^{\nu} (\nabla_{y^j} \tilde{\varphi}_h \cdot \nabla_{y^j} + \nabla_{\eta^j} \tilde{\varphi}_h \cdot \nabla_{\eta^j})\},
 \end{cases}$$

we can write

$$\begin{aligned}
 & \tilde{r}_{\nu+1,h}(\xi, x) \\
 (2.51) \quad &= h^{-2\nu n\sigma} \int \dots \int e^{-ih^{-2\sigma}\tilde{\varphi}_h({}^tL_h)^{(\nu+1)} q_{j,h}(\xi, x; \eta^j, y^{j-1})} \times \\
 & \quad \times dy^1 \dots dy^\nu d\eta^1 \dots d\eta^\nu
 \end{aligned}$$

for $l \geq 0$ at least formally, where tL_h is the transposed operator of L_h . We shall show that the right hand side of (2.51) converges absolutely for $l \geq 2\nu n$.

Noting

$$\begin{aligned}
 (2.52) \quad {}^tL_h &= -\frac{1}{\Gamma_h} \sum_{j=1}^{\nu} (\nabla_{y^j} \tilde{\varphi}_h \cdot \nabla_{y^j} + \nabla_{\eta^j} \tilde{\varphi}_h \cdot \nabla_{\eta^j}) \\
 & \quad + \frac{1}{\Gamma_h^2} \{2\nu \Gamma_h + i \sum_{j=1}^{\nu} (\nabla_{y^j} \tilde{\varphi}_h \cdot \nabla_{y^j} \Gamma_h + \nabla_{\eta^j} \tilde{\varphi}_h \cdot \nabla_{\eta^j} \Gamma_h)\}
 \end{aligned}$$

and

$$\begin{cases}
 \nabla_{y^j} \Gamma_h = 2h^{-2\sigma} \{ \nabla_{y^j} \tilde{\varphi}_h \cdot \tilde{\nabla}_x \nabla_x \tilde{J}_{j+1,h} + \nabla_{\eta^{j+1}} \tilde{\varphi}_h \cdot (I + \tilde{\nabla}_x \nabla_\xi \tilde{J}_{j+1,h}) \\
 \quad - \nabla_{\eta^j} \tilde{\varphi}_h + [\nabla_{y^j} \tilde{\varphi}_h \cdot \nabla_{y^j} \tilde{\nabla}_x \nabla_x \tilde{J}_{j+1,h} + \nabla_{\eta^{j+1}} \tilde{\varphi}_h \cdot \nabla_{y^j} \tilde{\nabla}_x \nabla_\xi \tilde{J}_{j+1,h}] y^j \}, \\
 \nabla_{\eta^j} \Gamma_h = 2h^{-2\sigma} \{ \nabla_{y^{j-1}} \tilde{\varphi}_h \cdot (I + \tilde{\nabla}_\xi \nabla_x \tilde{J}_{j+1,h}) + \nabla_{\eta^j} \tilde{\varphi}_h \cdot \tilde{\nabla}_\xi \nabla_\xi \tilde{J}_{j,h} \\
 \quad - \nabla_{y^j} \tilde{\varphi}_h + [\nabla_{y^{j-1}} \tilde{\varphi}_h \cdot \nabla_{\eta^j} \tilde{\nabla}_\xi \nabla_x \tilde{J}_{j,h} + \nabla_{\eta^j} \tilde{\varphi}_h \cdot \nabla_{\eta^j} \tilde{\nabla}_\xi \nabla_\xi \tilde{J}_{j,h}] \eta^j \},
 \end{cases}$$

and using (2.37) we see by induction that $({}^tL_h)^l$ has the form

$$(2.54) \quad ({}^tL_h)^l = \frac{1}{\Gamma_h^{2l}} \sum_{\substack{|\mu| \leq 2l \\ |\rho| \leq l}} a_{\mu,\rho,h}^{(l)} (h^{-\sigma}(\mathbf{y}, \boldsymbol{\eta}))^\mu \partial_{(\mathbf{y}, \boldsymbol{\eta})}^\rho,$$

where $(\mathbf{y}, \boldsymbol{\eta}) = (y^1, \dots, y^\nu, \eta^1, \dots, \eta^\nu)$ and $a_{\mu,\rho,h}^{(l)}$ is a polynomial of $\partial_{(\mathbf{y}, \boldsymbol{\eta})}^\alpha \nabla_{(x, \xi)}^2 \tilde{J}_{j,h}$

($|\alpha| \leq l - |\rho|, 1 \leq j \leq \nu + 1$) of order $3l, \nabla_{(x,\xi)}^2 \tilde{J}_{j,h}$ denoting one of the terms $\tilde{\nabla}_x \nabla_\xi \tilde{J}_{j,h}(\tilde{X}_{\nu,h}^{j-1}, \tilde{\Xi}_{\nu,h}^j, \tilde{X}_{\nu,h}^{j-1} + y^{j-1})$, etc. in (2.37), and satisfies

$$(2.55) \quad \begin{aligned} & |\partial_\xi^\alpha \partial_x^\beta a_{\mu,\rho,h}^{(l)}| \\ & \leq C_{\alpha,\beta} c^l \left(\frac{1 + \bar{\sigma}_\nu}{1 - \tau_0} \right)^{2|\alpha+\beta|} (1 + \max_{1 \leq j \leq \nu+1} |J_{j,h}|_{2,l+|\alpha+\beta|})^{3l} \times \\ & \quad \times (1 + \sum_{s=1}^{\nu+1} |J_{s,h}|_{3, (|\alpha+\beta|-1) \vee 0})^{(|\alpha+\beta|-1) \vee 0} \end{aligned}$$

for some constant $c < 1$, where we have used Proposition 2.3-iii). Thus taking $l > 2\nu n$, and using (2.32) and the inequalities

$$(2.56) \quad \begin{cases} |\eta^j| \leq \sum_{k=j}^\nu |\eta^k - \eta^{k+1}| \leq \sum_{k=1}^\nu |\eta^k - \eta^{k+1}|, \\ |y^j| \leq \sum_{k=1}^j |y^k - y^{k-1}| \leq \sum_{k=1}^j |y^k - y^{k-1} - r_k \eta^k| + \sum_{k=1}^j \sigma_j |\eta^k| \\ \leq (1 + \bar{\sigma}_j) \sum_{k=1}^j (|y^k - y^{k-1} - r_k \eta^k| + |\eta^k - \eta^{k+1}|), \end{cases}$$

where $y^0 = \eta^{\nu+1} = 0$ and $r_k = \tilde{\nabla}_\xi \nabla_\xi \tilde{J}_{k,h}(\tilde{\Xi}_{\nu,h}^k, \tilde{X}_{\nu,h}^{k-1} + y^{k-1}, \tilde{\Xi}_{\nu,h}^k + \eta^k)$, we see that the integral (2.51) is well-defined.

Thus we have again formally for any α, β

$$(2.57) \quad \begin{aligned} & \partial_\xi^\alpha \partial_x^\beta \tilde{r}_{\nu+1,h}(\xi, x) \\ & = h^{-2\nu n \sigma} \sum_{\substack{\alpha^1 + \alpha^2 = \alpha \\ \beta^1 + \beta^2 = \beta}} \int \dots \int_{\sim 2\nu n} \partial_\xi^{\alpha^1} \partial_x^{\beta^1} (e^{-ih^{-2\sigma} \tilde{\varphi}_h}) \times \\ & \quad \times \partial_\xi^{\alpha^2} \partial_x^{\beta^2} [({}^t L_h)^{(\nu+1)} q_{j,h}(\xi, x; \eta^j, y^{j-1})] dy^1 \dots dy^\nu d\eta^1 \dots d\eta^\nu. \end{aligned}$$

From Lemma 2.5-iii) we get for $|\alpha^1 + \beta^1| \geq 1$

$$(2.58) \quad \begin{aligned} & |\partial_\xi^{\alpha^1} \partial_x^{\beta^1} (e^{-ih^{-2\sigma} \tilde{\varphi}_h})| \\ & \leq \sum_{l=1}^{|\alpha^1 + \beta^1|} \sum_{k_1 + \dots + k_l = |\alpha^1 + \beta^1|} \prod_{j=1}^l \{C_{k_j} \left(\frac{1 + \bar{\sigma}_\nu}{1 - \tau_0} \right)^{2k_j - 1} \times \\ & \quad \times (1 + \sum_{s=1}^{\nu+1} |J_{s,h}|_{3, k_j - 1})^{k_j} (h^{-\sigma} \sum_{s=1}^\nu (|y^s| + |\eta^s|))\} \\ & \leq C_{\alpha^1, \beta^1} \left(\frac{1 + \bar{\sigma}_\nu}{1 - \tau_0} \right)^{2|\alpha^1 + \beta^1| - 1} (1 + \sum_{s=1}^{\nu+1} |J_{s,h}|_{3, |\alpha^1 + \beta^1| - 1})^{|\alpha^1 + \beta^1|} \times \\ & \quad \times (1 + h^{-\sigma} \sum_{s=1}^\nu (|y^s| + |\eta^s|))^{2|\alpha^1 + \beta^1|}. \end{aligned}$$

Using (2.56) and i) and ii) of Lemma 2.5 we have for $|\alpha + \beta| \geq 1$

$$|\partial_\xi^\alpha \partial_x^\beta \left(\frac{1}{\Gamma_h^{2l}} \right)|$$

$$\begin{aligned}
 &\leq \sum_{m=1}^{|\alpha+\beta|} 2l(2l+1)\cdots(2l+m-1)\Gamma_h^{-2l-m} \sum_{\substack{k_1+\cdots+k_m=|\alpha+\beta| \\ k_i \geq 1}} \prod_{i=1}^m \sum_{j=1}^{\nu} \times \\
 (2.59) \quad &\times \sum_{a^1+a^2=k_i} h^{-2\sigma} \{ |\partial_{(\xi,x)}^{a^1} \nabla_{y^j} \tilde{\Phi}_h| |\partial_{(\xi,x)}^{a^2} \nabla_{y^j} \tilde{\Phi}_h| \\
 &\quad + |\partial_{(\xi,x)}^{a^1} \nabla_{\eta^j} \tilde{\Phi}_h| |\partial_{(\xi,x)}^{a^2} \nabla_{\eta^j} \tilde{\Phi}_h| \} \\
 &\leq C_{\alpha,\beta} 2l(2l+1)\cdots(2l+|\alpha+\beta|-1) \nu^{3|\alpha+\beta|} \left(\frac{1+\bar{\sigma}_\nu}{1-\tau_0}\right)^{4|\alpha+\beta|} \times | \\
 &\quad \times (1+\sum_{s=1}^{\nu+1} |J_{s,h}|_{3,|\alpha+\beta|})^{|\alpha+\beta|} (1+\max_{1 \leq j \leq \nu+1} |J_{j,h}|_{2,|\alpha+\beta|})^{2|\alpha+\beta|} \frac{1}{\Gamma_h^{2l}}.
 \end{aligned}$$

On the other hand, from Proposition 2.3-iii) we get

$$\begin{aligned}
 &|\partial_\xi^\gamma \partial_x^\delta \partial_{\eta^j}^a \partial_{y^{j-1}}^b q_{j,h}(\xi, x; \eta^j, y^{j-1})| \\
 (2.60) \quad &\leq C_{\gamma,\delta} \left(\frac{1+\bar{\sigma}_\nu}{1-\tau_0}\right)^{2|\gamma+\delta|} (1+\sum_{s=1}^{\nu+1} |J_{s,h}|_{3,|\gamma+\delta|})^{|\gamma+\delta|} h^{m_j} |p_{j,h}|_{|\gamma+\delta+a+b|}^{(m_j)}.
 \end{aligned}$$

Then using (2.54), (2.55), (2.59) and (2.60) we obtain

$$\begin{aligned}
 &|\partial_\xi^{\alpha^2} \partial_x^{\beta^2} [({}^t L_h)^\nu (\prod_{j=1}^{\nu+1} q_{j,h}(\xi, x; \eta^j, y^{j-1}))]| \\
 &\leq C_{\alpha^2,\beta^2} c^l \nu^{3l} 2l(2l+1)\cdots(2l+|\alpha^2+\beta^2|-1) \nu^{3|\alpha^2+\beta^2|} \times \\
 (2.61) \quad &\times \left(\frac{1+\bar{\sigma}_\nu}{1-\tau_0}\right)^{8|\alpha^2+\beta^2|} (1+\max_{1 \leq j \leq \nu+1} |J_{j,h}|_{2,l+|\alpha^2+\beta^2|})^{3l+2|\alpha^2+\beta^2|} \times \\
 &\times (1+\sum_{s=1}^{\nu+1} |J_{s,h}|_{3,|\alpha^2+\beta^2|})^{3|\alpha^2+\beta^2|} \frac{|h^{-\sigma}(\mathbf{y}, \boldsymbol{\eta})|^{3l}}{\Gamma_h^{2l}} h^{\bar{m}_{\nu+1}} \times \\
 &\times \prod_{j=1}^{\nu+1} |p_{j,h}|_{|\alpha^2+\beta^2+l|}^{(m_j)}.
 \end{aligned}$$

Thus from (2.58) and (2.61) we have

$$\begin{aligned}
 &|\partial_\xi^{\alpha^2} \partial_x^{\beta^2} \tilde{\mathcal{F}}_{\nu+1,h}(\xi, x)| \\
 &\leq C_{\alpha,\beta} c^l 2l(2l+1)\cdots(2l+|\alpha+\beta|-1) \nu^{3l+3|\alpha+\beta|} \left(\frac{1+\bar{\sigma}_\nu}{1-\tau_0}\right)^{8|\alpha+\beta|} \times \\
 (2.62) \quad &\times (1+\sum_{s=1}^{\nu+1} |J_{s,h}|_{3,|\alpha+\beta|})^{3|\alpha+\beta|} (1+\max_{1 \leq j \leq \nu+1} |J_{j,h}|_{2,l+|\alpha+\beta|})^{3l+2|\alpha+\beta|} \times \\
 &\times h^{\bar{m}_{\nu+1}} \prod_{j=1}^{\nu+1} |p_{j,h}|_{|\alpha+\beta+l|}^{(m_j)} \int \cdots \int h^{-2\nu n \sigma} \Gamma_h^{-2l} \times \\
 &\times \{1+h^{-\sigma} \sum_{s=1}^{\nu} (|y^s|+|\eta^s|)\}^{2|\alpha+\beta|} |h^{-\sigma}(\mathbf{y}, \boldsymbol{\eta})|^{3l} d\mathbf{y} d\boldsymbol{\eta}.
 \end{aligned}$$

Since (2.56) shows

$$\begin{aligned}
 &\{1+h^{-\sigma} \sum_{s=1}^{\nu} (|y^s|+|\eta^s|)\}^{2|\alpha+\beta|} |h^{-\sigma}(\mathbf{y}, \boldsymbol{\eta})|^{3l} \\
 &\leq \{1+\sum_{s=1}^{\nu} h^{-\sigma} (|y^s|+|\eta^s|)\}^{2|\alpha+\beta|+3l}
 \end{aligned}$$

$$\leq \{2\nu(1+\bar{\sigma}_\nu) [1+h^{-\sigma} \sum_{k=1}^{\nu} (|\eta^k-\eta^{k+1}|+|y^k-y^{k-1}-r_k\eta^k|)]\}^{2|\alpha+\beta|+3l},$$

we see from (2.32) that the integrand of the right hand side of (2.62) is bounded by

$$(2.63) \quad 2^{6l+4|\alpha+\beta|} \nu^{4l+4|\alpha+\beta|} (1+\bar{\sigma}_\nu)^{3l+2|\alpha+\beta|} (1-\tau_0)^{-4l} \times \\ \times h^{-2\nu n\sigma} \prod_{j=1}^{\nu} (\langle h^{-\sigma}(\eta^j-\eta^{j+1}) \rangle \langle h^{-\sigma}(y^j-y^{j-1}-r_j\eta^j) \rangle)^{-(l-2|\alpha+\beta|)/2\nu}.$$

This is integrable in $(\mathbf{y}, \boldsymbol{\eta})$ uniformly in $h \in (0, 1)$ whenever $l > 2\nu n + 2|\alpha + \beta|$, which shows (2.57) is actually valid. Thus from (2.62) and (2.63) we have for any $l > 2\nu n + 2|\alpha + \beta|$ and some constant $c_0 > 1$

$$(2.64) \quad |\partial_{\xi}^{\alpha} \partial_x^{\beta} \tilde{r}_{\nu+1,h}(\xi, x)| \\ \leq C_{\alpha,\beta} c_0^{3l+2|\alpha+\beta|} 2l(2l+1) \dots (2l+|\alpha+\beta|-1) \nu^{7l+7|\alpha+\beta|} \times \\ \times (1+\bar{\sigma}_\nu)^{3l+10|\alpha+\beta|} (1-\tau_0)^{-4l-8|\alpha+\beta|} (1+\sum_{s=1}^{\nu+1} |J_{s,h}|_{3,|\alpha+\beta|})^{3|\alpha+\beta|} \times \\ \times (1+\max_{1 \leq j \leq \nu+1} |J_{j,h}|_{2,l+|\alpha+\beta|})^{3l+2|\alpha+\beta|} h^{\bar{m}_{\nu+1}} \prod_{j=1}^{\nu+1} |p_{j,h}|_{l+|\alpha+\beta|}^{(m_j)} \times \\ \times \int \dots \int \prod_{j=1}^{\nu} (\langle \eta^j-\eta^{j+1} \rangle \langle y^j-y^{j-1}-r_j\eta^j \rangle)^{-(l-2|\alpha+\beta|)/(2\nu)} d\mathbf{y} d\boldsymbol{\eta}.$$

Taking $l=2\nu n+2|\alpha+\beta|+1$ and noting that the integral is bounded by $c_1^{\nu} \nu^{2\nu}$ for some constant $c_1 > 1$ prove the theorem. \square

3. Approximate fundamental solutions

We consider the Hamiltonian $H(t, x, \xi)$ of the form

$$(3.1) \quad H(t, x, \xi) = \frac{1}{2} |\xi|^2 + V(t, x),$$

where the time-dependent potential $V(t, x)$ satisfies the following assumption (A):

ASSUMPTION (A)

- i) For each $t \in R^1$, $V(t, x)$ is a real-valued C^∞ -function of $x \in R^n$.
- ii) For any multi-index α , $\partial_x^\alpha V(t, x)$ is continuous in $(t, x) \in R^1 \times R^n$.
- iii) There exists a constant $\varepsilon > 0$ such that for any multi-index α with $|\alpha| \neq 0$

$$|\partial_x^\alpha V(t, x)| \leq C_\alpha \langle t \rangle^{-|\alpha|-\varepsilon},$$

where the constant $C_\alpha > 0$ is independent of t, x .

This assumption is the same as in [5], where we have studied the scattering

problem by $V(t, x)$. For the examples covered by this assumption, see [5, §1].

Let $0 \leq \delta \leq \rho \leq 1$ and set

$$(3.2) \quad H_h(t, x, \xi) = h^{\delta-\rho} H(t, h^{-\delta}x, h^\rho \xi).$$

We consider the Schrödinger equation

$$(3.3) \quad \begin{cases} L_h u \equiv (D_t + H_h(t, X, D_x))u = 0 & \text{on } R^1, \\ u|_{t=s_0} = f(\in \mathcal{D}) & (s_0 \in R^1). \end{cases}$$

In this and the next sections we shall construct the fundamental solution of (3.3) globally in time in the form of a product of a certain finite number of (conjugate) Fourier integral operators, the number depending on s_0 but not on t . Here by the fundamental solution of (3.3) we mean an operator $U_h(t, s_0)$ such that

$$(3.4) \quad \begin{cases} U_h(s_0, s_0) = I, \\ L_h U_h(t, s_0) = 0 & (t, s_0 \in R^1). \end{cases}$$

It is easily seen from (3.1) that $H(t, X, D_x)$ is symmetric in $L^2(R^n)$. Thus we have by 1° of the remark after Definition 1.5

$$(3.5) \quad \begin{aligned} H_h(t, X, D_x)f(x) &= H_h(t, X', D_x)f(x) \\ &= 0_s - \iint e^{i(x-x') \cdot \xi} H_h(t, x', \xi) f(x') dx' d\xi \end{aligned}$$

for $f \in \mathcal{D}$. So we consider the Cauchy problem

$$(3.3)' \quad \begin{cases} L'_h u \equiv (D_t + H_h(t, X', D_x))u = 0 & \text{on } R^1, \\ u|_{t=s_0} = f(\in \mathcal{D}) & (s_0 \in R^1) \end{cases}$$

instead of (3.3). In the following, for the sake of simplicity we restrict ourselves to considering only the case $t \geq s_0$, since the other case can be dealt with similarly.

Let $(q(t, s; x, \xi), p(t, s; x, \xi))$ be the solution of the Hamilton equation

$$(3.6) \quad \begin{cases} \frac{dq}{dt}(t, s) = \nabla_\xi H(t, q(t, s), p(t, s)), \\ \frac{dp}{dt}(t, s) = -\nabla_x H(t, q(t, s), p(t, s)) \end{cases}$$

on R^1 with the initial condition

$$(3.7) \quad q(s, s) = x, \quad p(s, s) = \xi \quad (s \in R^1).$$

The equation (3.6)–(3.7) is equivalent to the integral equation

$$(3.8) \quad \begin{cases} q(t, s) = x + \int_s^t p(\tau, s) d\tau, \\ p(t, s) = \xi - \int_s^t \nabla_x V(\tau, q(\tau, s)) d\tau. \end{cases}$$

Then we can easily prove the following proposition by successive approximation. Let $\mathcal{B}^{k, \infty}(R^m)$ denote the Fréchet space of C^∞ -functions $f(y)$ on R^m such that $\partial_y^\alpha f(y)$ ($|\alpha| \geq k$) are all bounded on R^m with semi-norms $|f|_{k, l}$ ($l=0, 1, 2, \dots$) defined by

$$|f|_{k, l} = \sum_{|\alpha| \leq k-1} \sup_y \{ |\partial_y^\alpha f(y)| \langle y \rangle^{k-|\alpha|} \} + \sum_{k \leq |\alpha| \leq k+l} \sup_y |\partial_y^\alpha f(y)|.$$

We often write $\mathcal{B}^\infty(R^m) = \mathcal{B}^{0, \infty}(R^m)$. We also use the class $C^l(\Omega | \mathcal{B}^{k, \infty}(R_y^m))$ for a domain $\Omega \subset R^p$ which consists of the functions $f(\omega, y)$ on $\Omega \times R^m$ such that for each $\omega \in \Omega$ $f(\omega, y)$ is in $\mathcal{B}^{k, \infty}(R_y^m)$ and any derivative $\partial_y^\alpha f(\omega, y)$ is in $C^l(\Omega \times R^m)$. Then:

Proposition 3.1. *There exists a unique solution of (3.8). The solution (q, p) $(t, s; x, \xi)$ belongs to $C^1(R_t^1 \times R_s^1 | \mathcal{B}^{1, \infty}(R_x^n \times R_\xi^n))$. Furthermore there exist positive constants T_0 and C_0 such that the following estimates hold:*

i) For any $t \geq s \geq T_0$ and $x, \xi \in R^n$

$$(3.9) \quad \begin{cases} |q(s, t; x, \xi) - x| + |q(t, s; x, \xi) - x| \leq C_0(t-s) (\langle s \rangle^{-\varepsilon} + |\xi|), \\ |p(s, t; x, \xi) - \xi| + |p(t, s; x, \xi) - \xi| \leq C_0 \langle s \rangle^{-\varepsilon}; \end{cases}$$

$$(3.10) \quad \begin{cases} |\nabla_x q(s, t; x, \xi) - I| \leq C_0 \langle s \rangle^{-\varepsilon}, \quad |\nabla_x q(t, s; x, \xi) - I| \leq C_0(t-s) \langle s \rangle^{-1-\varepsilon}, \\ |\nabla_x p(s, t; x, \xi)| + |\nabla_x p(t, s; x, \xi)| \leq C_0 \langle s \rangle^{-1-\varepsilon}; \end{cases}$$

$$(3.11) \quad \begin{cases} |\nabla_\xi q(s, t; x, \xi) - (s-t)I| \leq C_0(t-s) \langle s \rangle^{-\varepsilon}, \\ |\nabla_\xi p(s, t; x, \xi) - I| \leq C_0(t-s) \langle s \rangle^{-1-\varepsilon}; \end{cases}$$

and

$$(3.12) \quad \begin{cases} |\nabla_\xi q(t, s; x, \xi) - (t-s)I| \leq C_0(t-s) \langle s \rangle^{-\varepsilon}, \\ |\nabla_\xi p(t, s; x, \xi) - I| \leq C_0 \langle s \rangle^{-\varepsilon}. \end{cases}$$

In particular, when $0 \leq t-s \leq 1$ and $s \geq T_0$, we have

$$(3.13) \quad \begin{cases} |\nabla_x q(s, t; x, \xi) - I| \leq C_0(t-s)^2 \langle s \rangle^{-2-\varepsilon}, \\ |\nabla_x p(t, s; x, \xi)| \leq C_0(t-s) \langle s \rangle^{-2-\varepsilon}, \\ |\nabla_\xi p(t, s; x, \xi) - I| \leq C_0(t-s)^2 \langle s \rangle^{-2-\varepsilon}. \end{cases}$$

ii) For any α, β with $|\alpha + \beta| \geq 2$, there is a constant $C_{\alpha, \beta}$ independent of $t \geq s (\geq T_0)$ and x, ξ such that

$$(3.14) \quad \begin{cases} |\partial_{\xi}^{\alpha} \partial_x^{\beta} q(s, t; x, \xi)| \leq C_{\alpha, \beta} (t-s)^{|\alpha|} \langle s \rangle^{-1-\varepsilon}, \\ |\partial_{\xi}^{\alpha} \partial_x^{\beta} p(s, t; x, \xi)| \leq C_{\alpha, \beta} (t-s)^{|\alpha|} \langle s \rangle^{-2-\varepsilon}, \end{cases}$$

and

$$(3.15) \quad \begin{cases} |\partial_{\xi}^{\alpha} \partial_x^{\beta} q(t, s; x, \xi)| \leq C_{\alpha, \beta} (t-s) \langle s \rangle^{-\varepsilon}, \\ |\partial_{\xi}^{\alpha} \partial_x^{\beta} p(t, s; x, \xi)| \leq C_{\alpha, \beta} \langle s \rangle^{-\varepsilon}. \end{cases}$$

Proof. (3.13) follows from (3.10)–(3.12) and the equalities obtained by differentiating (3.8) with respect to x or ξ . For the proof of the other results, see the proof of Proposition 2.1 of [5]. \square

From this proposition, we can easily get the following important proposition.

Proposition 3.2. *Take $T > T_0$ so large that $C_0 \langle T \rangle^{-\varepsilon} < 1/2$ for the constant C_0 in Proposition 3.1. Then for $t \geq s \geq T$ there exist the inverse C^∞ diffeomorphisms $x \mapsto y(s, t; x, \xi)$ and $\xi \mapsto \eta(t, s; x, \xi)$ of the mappings $y \mapsto x = q(s, t; y, \xi)$ and $\eta \mapsto \xi = p(t, s; x, \eta)$, respectively. These mappings y and η belong to $C^1(A_T | \mathcal{D}^{1, \infty}(R_x^n \times R_\xi^n))$, where $A_T \equiv \{(t, s) | t \geq s \geq T\}$, and they satisfy the following properties:*

$$i) \quad q(s, t; y(s, t; x, \xi), \xi) = x, \quad p(t, s; x, \eta(t, s; x, \xi)) = \xi.$$

$$ii) \quad \begin{cases} q(t, s; x, \eta(t, s; x, \xi)) = y(s, t; x, \xi), \\ p(s, t; y(s, t; x, \xi), \xi) = \eta(t, s; x, \xi). \end{cases}$$

$$iii) \quad \text{There is a constant } C_1 > 1 \text{ such that for any } (t, s) \in A_T \text{ and } x, \xi \in R^n$$

$$(3.16) \quad \begin{cases} |\eta(t, s; x, \xi) - \xi| \leq C_1 \langle s \rangle^{-\varepsilon}, \\ |y(s, t; x, \xi) - x| \leq C_1 (t-s) (\langle s \rangle^{-\varepsilon} + |\xi|); \end{cases}$$

$$(3.17) \quad \begin{cases} |\nabla_x y(s, t; x, \xi) - I| \leq C_1 \langle s \rangle^{-\varepsilon}, \\ |\nabla_\xi y(s, t; x, \xi) - (t-s)I| \leq (C_1 - 1) (t-s) \langle s \rangle^{-\varepsilon}; \end{cases}$$

and

$$(3.18) \quad \begin{cases} |\nabla_x \eta(t, s; x, \xi)| \leq C_1 \langle s \rangle^{-1-\varepsilon}, \\ |\nabla_\xi \eta(t, s; x, \xi) - I| \leq C_1 \langle s \rangle^{-\varepsilon}. \end{cases}$$

If, in particular $0 \leq t-s \leq 1$, we have

$$(3.19) \quad \begin{cases} |\nabla_x y(s, t; x, \xi) - I| \leq C_1 (t-s)^2 \langle s \rangle^{-2-\varepsilon}, \\ |\nabla_x \eta(t, s; x, \xi)| \leq C_1 (t-s) \langle s \rangle^{-2-\varepsilon}, \\ |\nabla_\xi \eta(t, s; x, \xi) - I| \leq C_1 (t-s)^2 \langle s \rangle^{-2-\varepsilon}. \end{cases}$$

iv) For any α, β with $|\alpha + \beta| \geq 2$, there is a constant $C_{\alpha, \beta}$ such that for any $(t, s) \in A_T$ and $x, \xi \in R^n$

$$(3.20) \quad \begin{cases} |\partial_{\xi}^{\alpha} \partial_x^{\beta} \eta(t, s; x, \xi)| \leq C_{\alpha, \beta} \langle s \rangle^{-\varepsilon}, \\ |\partial_{\xi}^{\alpha} \partial_x^{\beta} y(s, t; x, \xi)| \leq C_{\alpha, \beta} (t-s+1) \langle s \rangle^{-\varepsilon}. \end{cases}$$

Proof. The proof except for (3.19) is similar to that of Proposition 2.2 of [5]. (3.19) follows from (3.13), (3.10) and (3.12) of Proposition 3.1 by virtue of the relations in i). \square

When $|t-s|$ is small, we have the following estimates for (q, p) and (y, η) (see [4, §3]).

Proposition 3.3. *There exists a small constant $0 < \tilde{\delta} < 1$ such that the following assertions hold:*

i) *We have*

$$(3.21) \quad (q, p)(t, s; x, \xi) \in C^1(B_{\tilde{\delta}} | \mathcal{B}^{1,\infty}(R^{2n})),$$

where $B_{\tilde{\delta}} \equiv \{(t, s) | t, s \in R^1, |t-s| \leq \tilde{\delta}\}$, and

$$(3.22) \quad \text{“}\{[(q, p)(t, s; x, \xi) - (x, \xi)] / (t-s)\}_{(t,s) \in B_{\tilde{\delta}}}\text{ is bounded in } \mathcal{B}^{1,\infty}(R^{2n}).\text{”}$$

ii) *For $(t, s) \in B_{\tilde{\delta}}$, there exist the inverse C^∞ diffeomorphisms $x \mapsto y(t, s; x, \xi)$ and $\xi \mapsto \eta(t, s; x, \xi)$ of the mappings $y \mapsto x = q(t, s; y, \xi)$ and $\eta \mapsto \xi = p(t, s; x, \eta)$, respectively, and they satisfy*

$$(3.23) \quad (y, \eta)(t, s; x, \xi) \in C^1(B_{\tilde{\delta}} | \mathcal{B}^{1,\infty}(R^{2n})),$$

and

$$(3.24) \quad \text{“}\{[(y, \eta)(t, s; x, \xi) - (x, \xi)] / (t-s)\}_{(t,s) \in B_{\tilde{\delta}}}\text{ is bounded in } \mathcal{B}^{1,\infty}(R^{2n}).\text{”}$$

DEFINITION 3.4. For $(t, s) \in A_T \cup B_{\tilde{\delta}}$, define

$$(3.25) \quad \phi(s, t; x, \xi) = u(s, t; y(s, t; x, \xi), \xi),$$

where

$$(3.26) \quad u(s, t; y, \eta) = y \cdot \eta + \int_t^s (\xi \cdot \nabla_\xi H - H)(\tau, q(\tau, t; y, \eta), p(\tau, t; y, \eta)) d\tau.$$

Proposition 3.5. *Let $(t, s) \in A_T$ (or $B_{\tilde{\delta}}$). Then $\phi(s, t; x, \xi)$ defined above satisfies*

$$(3.27) \quad \begin{cases} \partial_s \phi(s, t; x, \xi) + H(s, x, \nabla_x \phi(s, t; x, \xi)) = 0, \\ \phi(t, t; x, \xi) = x \cdot \xi; \end{cases}$$

$$(3.28) \quad \partial_t \phi(s, t; x, \xi) - H(t, \nabla_\xi \phi(s, t; x, \xi), \xi) = 0;$$

and

$$(3.29) \quad \begin{cases} \nabla_x \phi(s, t; x, \xi) = \eta(t, s; x, \xi), \\ \nabla_\xi \phi(s, t; x, \xi) = y(s, t; x, \xi). \end{cases}$$

Furthermore we have

$$(3.30) \quad \phi(s, t; x, \xi) \in C^1(A_T | \mathcal{B}^{2,\infty}(R^{2n})) \text{ (or } \in C^1(B_{\tilde{\delta}} | \mathcal{B}^{2,\infty}(R^{2n}))) .$$

Proof. (3.27)–(3.29) can be shown by direct calculations (or see Kumano-go [7] and Kumano-go, Taniguchi, Tozaki [8]). (3.30) follows from Proposition 3.2 (or Proposition 3.3). \square

DEFINITION 3.6. Let $\phi_h(s, t; x, \xi)$ be defined by

$$(3.31) \quad \phi_h(s, t; x, \xi) = h^{\delta-\rho} \phi(s, t; h^{-\delta}x, h^\rho \xi) \quad (0 \leq \delta \leq \rho \leq 1)$$

for $(t, s) \in A_T \cup B_{\tilde{\delta}}$.

In the following, we switch to another large $T > T_0$ such that $C_1 \langle T \rangle^{-\epsilon} < 1$, if necessary.

Proposition 3.7. i) For $(t, s) \in A_T$, we have

$$(3.32) \quad \phi_h(s, t; x, \xi) \in P_{\rho, \delta}^{[x]}(C_1 \langle s \rangle^{-\epsilon}, C_1(t-s); h) .$$

When $(t, s) \in A_T$ and $|t-s| \leq 1$, we have

$$(3.33) \quad \phi_h(s, t; x, \xi) \in P_{\rho, \delta}^{[x]}(C_1(t-s) \langle s \rangle^{-2-\epsilon}, C_1(t-s); h) .$$

ii) For any $l \geq 0$, there exist constants $0 < \tilde{\delta}_l \leq \tilde{\delta}$ and $c_l \geq 1$ such that $c_l \tilde{\delta}_l < 1$ and

$$(3.34) \quad \phi_h(s, t; x, \xi) \in P_{\rho, \delta}(c_l |t-s|, l; h)$$

for any $(t, s) \in B_{\tilde{\delta}_l}$.

Proof. i) is clear from Propositions 3.5 and 3.2. ii) is also clear from Propositions 3.5 and 3.3. \square

In the sequel we switch to another small $\tilde{\delta} > 0$ such that $\tilde{\delta} \leq \tilde{\delta}_0$, if necessary. We next solve the transport equations.

DEFINITION 3.8. For $(t, s) \in A_T \cup B_{\tilde{\delta}}$, we define a sequence of functions $a_j(t, s; \xi, x')$ ($j=0, 1, \dots$) inductively as follows:

$$(3.35) \quad \begin{aligned} & a_0(t, s; \xi, x') \\ &= \exp \left\{ -\frac{1}{2} \sum_{l,k=1}^n \int_t^s (\partial_{x_k} \partial_{x_l} H)(\tau, y(s, \tau; x', \Xi(\tau)), \Xi(\tau)) \times \right. \\ & \quad \left. \times (\partial_{\xi_k} \partial_{\xi_l} \phi)(s, \tau; x', \Xi(\tau)) d\tau \right\} \end{aligned}$$

and for $j \geq 1$

$$(3.36) \quad a_j(t, s; \xi, x') = -a_0(t, s; \xi, x') \int_t^s \frac{B_j(\tau, s; \Xi(\tau), x')}{a_0(\tau, s; \Xi(\tau), x')} d\tau ,$$

where

$$(3.37) \quad \Xi(\tau) = p(\tau, s; x', \eta(t, s; x', \xi))$$

and

$$(3.38) \quad \begin{aligned} & B_j(t, s; \xi, x') \\ &= \sum_{2 \leq |\alpha| \leq j+1} \frac{1}{\alpha!} \partial_{\xi'}^\alpha \{ (\partial_x^\alpha H)(t, \tilde{\nabla}_\xi \phi(s, t; \xi, x', \xi'), \xi) \times \\ & \quad \times a_{j+1-|\alpha|}(t, s; \xi', x') \}_{|\xi'=\xi}. \end{aligned}$$

Moreover, in case $0 \leq \delta < \rho \leq 1$, we define $e_h(t, s; \xi, x')$ by

$$(3.39) \quad e_h(t, s; \xi, x') = \sum_{j=0}^{\infty} \chi(\varepsilon_j^{-1}h) (ih^{\rho-\delta})^j a_j(t, s; h^\rho \xi, h^{-\delta} x'),$$

where χ and $\{\varepsilon_j\}_{j=0}^{\infty}$ are taken as in Proposition 1.3.

Let $C^l(\Omega | B_{\rho,\delta}^m(h))$ ($\Omega \subset R^p$ domain) denote the set of families $\{f_h(\omega, y)\}_{0 < h < 1}$ of functions $f_h(\omega, y)$ such that for each $\omega \in \Omega$ $\{f_h(\omega, y)\}_{0 < h < 1} \in \{B_{\rho,\delta}^m(h)\}_{0 < h < 1}$ and the derivative $\partial_y^\alpha f_h(\omega, y)$ belongs to $C^l(\Omega \times R_j^k)$ for each $h \in (0, 1)$.

Proposition 3.9. i) *The function $a_j(t, s; \xi, x')$ ($j=0, 1, 2, \dots$) belongs to $C^1(A_T \cup B_{\tilde{\delta}} | \mathcal{B}^\infty(R^{2n}))$ and is the solution of the transport equation*

$$(3.40) \quad \begin{aligned} & -\partial_t a_j(t, s; \xi, x') + \sum_{k=1}^n (\partial_{x_k} H)(t, y(s, t; x', \xi), \xi) (\partial_{\xi_k} a_j)(rt, s; \xi, x') \\ & + \frac{1}{2} \sum_{i,k=1}^n (\partial_{x_i} \partial_{x_j} H)(t, y(s, t; x', \xi), \xi) (\partial_{\xi_k} \partial_{\xi_l} \phi)(s, t; x', \xi) \times \\ & \times a_j(t, s; \xi, x') + B_j(t, s; \xi, x') = 0 \end{aligned}$$

for $(t, s) \in A_T \cup B_{\tilde{\delta}}$ with the initial condition

$$(3.41) \quad a_0(s, s; \xi, x') = 1, \quad a_j(s, s; \xi, x') = 0 \quad (j = 1, 2, \dots),$$

where we put $B_0(t, s; \xi, x') = 0$. More precisely $a_j(t, s; \xi, x')$ satisfies the estimates

$$(3.42) \quad \begin{cases} |\partial_{\xi'}^\alpha \partial_{x'}^\beta (a_0(t, s; \xi, x') - 1)| \leq C_{\alpha,\beta} \langle s \rangle^{-\varepsilon} \text{ (or } \leq C_{\alpha,\beta} |t-s|^2), \\ |\partial_{\xi'}^\alpha \partial_{x'}^\beta a_j(t, s; \xi, x')| \leq C_{\alpha,\beta} \langle s \rangle^{-1-\varepsilon} \text{ (or } \leq C_{\alpha,\beta} |t-s|^2) \quad (j \geq 1) \end{cases}$$

for $(t, s) \in A_T$ (or $\in B_{\tilde{\delta}}$), where the constant $C_{\alpha,\beta} > 0$ is independent of ξ, x' and $(t, s) \in A_T$ (or $\in B_{\tilde{\delta}}$).

ii) *When $0 \leq \delta < \rho \leq 1$, $e_h(t, s; \xi, x')$ of (3.39) is well-defined and belongs to $C^1(A_T \cup B_{\tilde{\delta}} | B_{\rho,\delta}^0(h))$. Moreover the following estimate holds: For any $(t, s) \in A_T$ (or $\in B_{\tilde{\delta}}$)*

$$(3.43) \quad |e_h(t, s) - 1|^{(0)} \leq C_l \langle s \rangle^{-\varepsilon} \text{ (or } \leq C_l |t-s|^2)$$

in $B_{\rho,\delta}^0(h)$, where the constant $C_l > 0$ is independent of t and s .

Proof. i) We have only to prove (3.42), since the others are obvious from

the theory of the first order differential equations. We first note by (3.37), (3.10), (3.12), (3.15), and (3.20) that

$$(3.44) \quad |\partial_{\xi}^{\alpha} \partial_x^{\beta} \Xi(\tau)| \leq C_{\alpha, \beta}$$

for some constant $C_{\alpha, \beta} > 0$ independent of t, τ, s, ξ and x' . Thus from (3.35) we have the first estimate of (3.42). Then by induction we get the second estimate of (3.42) and

$$(3.45) \quad |\partial_{\xi}^{\alpha} \partial_x^{\beta} B_j(t, s; \xi, x')| \leq C_{\alpha, \beta} \langle t \rangle^{-2-\epsilon} \text{ (or } \leq C_{\alpha, \beta} |t-s| \text{)}$$

for $(t, s) \in A_T$ (or $\in B_{\delta}$), using (3.36), (3.38), (3.29), (3.17) and (3.20) (or (3.24)).

ii) is then easily seen from (3.39) and (3.42). \square

DEFINITION 3.10. For $(t, s) \in A_T \cup B_{\delta}$ and $f \in \mathcal{S}$, we define

$$(3.46) \quad E_h^0(\phi_h(s, t)^*)f(x) = 0_s - \iint e^{i(x \cdot \xi - \phi_h(s, t; x', \xi))} f(x') dx' d\xi$$

in case $0 \leq \delta \leq \rho \leq 1$, and

$$(3.47) \quad \begin{aligned} & E_h^{\infty}(\phi_h(s, t)^*)f(x) \\ &= 0_s - \iint e^{i(x \cdot \xi - \phi_h(s, t; x', \xi))} e_h(t, s; \xi, x') f(x') dx' d\xi \end{aligned}$$

in case $0 \leq \delta < \rho \leq 1$.

Theorem 3.11. *Let Assumption (A) be satisfied. Let $(t, s) \in A_T$ (or $\in B_{\delta}$) and $f \in \mathcal{S}$, and define*

$$(3.48) \quad G_h^m(\phi_h(s, t)^*)f(x) = \frac{1}{i} ((D_t + H_h(t, X', D_x)) E_h^m(\phi_h(s, t)^*)f(x),$$

where $m=0$ in case $0 \leq \delta \leq \rho \leq 1$ and $m=\infty$ in case $0 \leq \delta < \rho \leq 1$. Then:

i) We have

$$(3.49) \quad E_h^m(\phi_h(s, s)^*) = I$$

and

$$(3.50) \quad g_h^m(t, s; \xi, x') \in C^0(A_T | B_{\rho, \delta}^m(h)) \text{ (or } \in C^0(B_{\delta} | B_{\rho, \delta}^m(h)))$$

for $m=0$ or ∞ . Here $g_h^m(t, s; \xi, x') = \sigma(G_h^m(\phi_h(s, t)^*))(\xi, x')$ is the symbol function of $G_h^m(\phi_h(s, t)^*)$:

$$(3.51) \quad \begin{aligned} & G_h^m(\phi_h(s, t)^*)f(x) \\ &= 0_s - \iint e^{i(x \cdot \xi - \phi_h(s, t; x', \xi))} g_h^m(t, s; \xi, x') f(x') dx' d\xi. \end{aligned}$$

ii) More precisely, we have for $(t, s) \in A_T$ (or $\in B_{\delta}$) and any $N \geq 1$

$$(3.52) \quad |g_h^0(t, s)|_i^{(0)} \leq C_i \langle t \rangle^{-1-\epsilon} \text{ (or } \leq C_i |t-s| \text{)}$$

and

$$(3.53) \quad |g_h^\infty(t, s)|^{(0)} \leq C_{l,N} h^N \langle t \rangle^{-2-\varepsilon} \text{ (or } \leq C_{l,N} h^N |t-s| \text{)}.$$

Here $|\cdot|^{(0)}$ denotes the semi-norm of $B_{p,\delta}^0(h)$, and the constants C_l and $C_{l,N}$ are independent of t, s and h . Hence we have for $(t, s) \in A_T$ (or $\in B_{\bar{\delta}}$) and any $N \geq 1$

$$(3.54) \quad \|G_h^0(\phi_h(s, t)^*)\|_{L^2 \rightarrow L^2} \leq C \langle t \rangle^{-1-\varepsilon} \text{ (or } \leq C |t-s| \text{)}$$

and

$$(3.55) \quad \|G_h^\infty(\phi_h(s, t)^*)\|_{L^2 \rightarrow L^2} \leq C_N h^N \langle t \rangle^{-2-\varepsilon} \text{ (or } \leq C_N h^N |t-s| \text{)},$$

where C and C_N are independent of t, s and h .

REMARK. Theorem 3.11-i) says that $E_h^m(\phi_h(s, t)^*)$ is the approximate fundamental solution of order $m(m=0$ or $\infty)$ in the sense of Kitada and Kumano-go (see [6, §5]), though the condition (3.50) is weaker than (5.29)-i) and (5.30)-i) of [6].

Proof. (3.49) is obvious by definition. So we have only to prove (3.52) and (3.53), since (3.50), (3.54) and (3.55) follows from them by Theorem 1.7. Using Theorem 1.6 and (3.28) we have for $(t, s) \in A_T \cup B_{\bar{\delta}}$

$$(3.56) \quad \begin{aligned} &g_h^0(t, s; \xi, x') \\ &= \sum_{i,k=1}^n \int_0^1 \int_0^1 e^{-iy^\eta} (\partial_{x_k} \partial_{x_i} H_h)(t, \theta y + \nabla_\xi \phi_h(s, t; \xi, x', \xi - \eta), \xi) \times \\ &\quad \times \left(\int_0^1 r(\partial_{\xi_k} \partial_{\xi_i} \phi_h)(s, t; x', \xi - r\eta) dr \right) dy d\eta d\theta, \end{aligned}$$

from which follows (3.52) by virtue of the estimates in Propositions 3.2 and 3.3 and our Assumption (A)-iii).

Similarly using Theorem 1.6, (3.28) and (3.40) we get for $N \geq 2$

$$(3.57) \quad \begin{aligned} &g_h^\infty(t, s; \xi, x') \\ &= -i \sum_{j=0}^\infty \chi(\varepsilon_j^{-1} h) (ih^{\rho-\delta})^j B_j(t, s; h^\rho \xi, h^{-\delta} x') \\ &\quad + i \sum_{m=1}^\infty (ih^{\rho-\delta})^m \sum_{2 \leq |\alpha| \leq \min(N-1, m+1)} \frac{1}{\alpha!} \chi(\varepsilon_{m+1-|\alpha|}^{-1} h) \times \\ &\quad \times \partial_{\xi'}^\alpha \{(\partial_x^\alpha H)(t, \nabla_\xi \phi(s, t; h^\rho \xi, h^{-\delta} x', \xi'), h^\rho \xi) \times \\ &\quad \quad \times a_{m+1-|\alpha|}(t, s; \xi', h^{-\delta} x')\}_{|\xi' = h^\rho \xi} \\ &\quad + h^{(N-1)(\rho-\delta)} N \sum_{|\gamma|=N} \frac{(-1)^{|\gamma|}}{\gamma!} \int_0^1 (1-\theta)^{N-1} t_{\gamma,h}(\xi, x'; \theta) d\theta, \end{aligned}$$

where

$$(3.58) \quad \begin{aligned} &t_{\gamma,h}(\xi, x'; \theta) \\ &= 0_s - \int \int e^{-iy^\eta} \partial_{\xi'}^\gamma \{ (D_x^\gamma H)(t, h^{-\delta} \theta y + \nabla_\xi \phi(s, t; h^\rho \xi, h^{-\delta} x', \xi'), \xi) \times \end{aligned}$$

$$\times e_h(t, s; h^{-\rho}\xi', x') \Big|_{|\xi' = h^{\rho}(\xi - \eta)} dy d\eta.$$

Thus by (3.42), (3.45), (3.43) and Proposition 3.2 (or 3.3), we obtain (3.53). \square

4. Fundamental solution global in time

We first construct the fundamental solution locally in time. For this purpose we record a theorem concerning the multi-products of conjugate Fourier integral operators which is a version of Theorem 4.3 of [6].

Theorem 4.1. *Let $n_0 > n$ be an even integer and put $\bar{l} = 21n_0 + 1$. Let $\bar{\tau} > 0$ be sufficiently small as in Theorem 3.8 of [6]. Let $\phi_{j,h}(x, \xi) \in P_{\rho, \delta}(\tau_j, \bar{l}; h)$ for $j = 1, 2, \dots$, and let $\bar{\tau}_\infty \equiv \sum_{j=1}^\infty \tau_j \leq \bar{\tau}$. Let $\nu \geq 1$ be an integer and put $\Phi_{\nu+1,h} = \phi_{1,h} \# \dots \# \phi_{\nu+1,h}$. Let $p_{j,h}(\xi, x') \in B_{\rho, \delta}^{m_j}(h)$ for $j = 1, \dots, \nu + 1$. Then there exists a symbol $r_{\nu+1,h}(\xi, x') \in B_{\rho, \delta}^{\bar{m}_{\nu+1}}(h)$ ($\bar{m}_{\nu+1} = m_1 + \dots + m_{\nu+1}$) such that*

$$(4.1) \quad P_{\nu+1,h}(\phi_{\nu+1,h}^*) \cdots P_{1,h}(\phi_{1,h}^*) = R_{\nu+1,h}(\Phi_{\nu+1,h}^*)$$

and

$$(4.2) \quad \begin{aligned} & |r_{\nu+1,h}|^{(\bar{m}_{\nu+1})} \\ & \leq \bar{C}_l^{l+2} \exp(\bar{c}_l(1 + \sum_{s=1}^{\nu+1} |J_{s,h}|_{2,k_1})^{k_1+2}) \times \\ & \quad \times \sum_{l_1 + \dots + l_{\nu+1} \leq l + 2n_0} \prod_{j=1}^{\nu+1} |p_{j,h}|_{\vartheta_{n_0+l_j}}^{(m_j)}, \end{aligned}$$

where $P_{j,h}(\phi_{j,h}^*) = p_{j,h}(\phi_{j,h}^*; D_x, X')$; $R_{\nu+1,h}(\Phi_{\nu+1,h}^*) = r_{\nu+1,h}(\Phi_{\nu+1,h}^*; D_x, X')$; $J_{j,h} = \phi_{j,h} - x \cdot \xi$; $k_1 = 2l + 25n_0 + 1$; and \bar{C}_l are \bar{c}_l positive constants.

Proof. This theorem follows from Theorem 4.3 of [6], if we note the following fact: For $P_h(\phi_h^*) = p_h(\phi_h^*; D_x, X') \in B_{\rho, \delta}^m(\phi_h^*)$, $\phi_h \in P_{\rho, \delta}(\tau, 0; h)$, we have for $f \in \mathcal{S}$

$$(4.3) \quad P_h(\phi_h^*)^* f(x) = q_h(\phi_h; X, D_x) f(x),$$

where $q_h(x, \xi) = \overline{p_h(\xi, x)}$. \square

Now we construct the local fundamental solution.

Theorem 4.2. *Let Assumption (A) be satisfied. Then for a sufficiently small $0 < \delta_0 \leq \delta (< 1)$, there exists uniquely the fundamental solution $U_h(t, s)$ of the equation*

$$(4.4) \quad \begin{cases} U_h(s, s) = I & (s \in \mathbb{R}^1), \\ L_h U_h(t, s) = 0 & (|t - s| \leq \delta_0). \end{cases}$$

Moreover the fundamental solution $U_h(t, s)$ is uniquely represented as a conjugate Fourier integral operator with phase function $\phi_h(s, t)$ and a symbol of class

$C^1(B_{\delta_0} | B_{\rho, \delta}^0(h))$. More precisely there exist symbols $d_h^m(t, s; \xi, x') \in C^1(B_{\delta_0} | B_{\rho, \delta}^m(h))$ ($m=0$ in case $0 \leq \delta \leq \rho \leq 1$ and $m=\infty$ in case $0 \leq \delta < \rho \leq 1$) such that for $D_h^m(\phi_h(s, t)^*)$ defined by

$$(4.5) \quad \begin{aligned} & D_h^m(\phi_h(s, t)^*)f(x) \\ &= 0_s - \iint e^{i(x \cdot \xi - \phi_h(s, t) \cdot x' \cdot \xi)} \bar{d}_h^m(t, s; \xi, x') f(x') dx' d\xi \quad (f \in \mathcal{S}), \end{aligned}$$

we can write

$$(4.6) \quad U_h(t, s) = E_h^m(\phi_h(s, t)^*) + D_h^m(\phi_h(s, t)^*)$$

for $(t, s) \in B_{\delta_0}$. The operator $U_h(t, s)$ is extended to a unitary operator in $L^2(\mathbb{R}^n)$, and the following relations hold:

$$(4.7) \quad U_h(t, \theta)U_h(\theta, r) = U_h(t, r), \quad t, \theta, r \in [s - \delta_0/2, s + \delta_0/2],$$

$$(4.8) \quad D_s U_h(t, s) - U_h(t, s)H_h(s, X, D_x) = 0, \quad |t - s| \leq \delta_0.$$

Proof. We proceed quite similarly as in the proof of Theorem 6.1 of [6]. Let $n_0 > n$ be even and let $\bar{l} = 21n_0 + 1$. Let $\bar{\tau} > 0$ be sufficiently small so that Theorem 3.8 of [6] holds for our case. For $c_{\bar{\tau}}$ in Proposition 3.7-ii), we take $\delta_0 > 0$ as $c_{\bar{\tau}}\delta_0 \leq \bar{\tau}$. Then for any subdivision $\Delta: t > t_\nu > t_{\nu-1} > \dots > t_1 > s$ of $[s, t]$, we can easily see that

$$(4.9) \quad \phi_h(s, t_1) \# \phi_h(t_1, t_2) \# \dots \# \phi_h(t_\nu, t) = \phi_h(s, t)$$

holds (cf. Kumano-go, Taniguchi and Tozaki [8]). Now using Theorem 4.1, we define $W_{\nu, h}^m(\phi_h(s, t)^*)$ by

$$(4.10) \quad W_{1, h}^m(\phi_h(s, t)^*) = G_h^m(\phi_h(s, t)^*) = \frac{1}{i} L_h E_h^m(\phi_h(s, t)^*)$$

and

$$(4.11) \quad \begin{aligned} W_{\nu+1, h}^m(\phi_h(s, t)^*) &= \int_s^t W_{1, h}^m(\phi_h(\theta, t)^*) W_{\nu, h}^m(\phi_h(s, \theta)^*) d\theta \\ &= \int_s^{t'} \int_s^{t_\nu} \dots \int_s^{t_2} W_{1, h}^m(\phi_h(t_\nu, t)^*) W_{1, h}^m(\phi_h(t_{\nu-1}, t_\nu)^*) \dots \\ &\quad \dots W_{1, h}^m(\phi_h(s, t_1)^*) dt_1 \dots dt_\nu, \end{aligned}$$

where $m=0$ in case $0 \leq \delta \leq \rho \leq 1$ and $m=\infty$ in case $0 \leq \delta < \rho \leq 1$. Then, in quite the same way as in [6], we see from Theorem 3.11-i) that the following series converges in $C^0(B_{\delta_0} | B_{\rho, \delta}^m(h))$:

$$(4.12) \quad \bar{d}_h^m(t, s; \xi, x') = \sum_{\nu=1}^{\infty} w_{\nu, h}^m(t, s; \xi, x'),$$

where $w_{\nu, h}^m(t, s; \xi, x') \equiv \sigma(W_{\nu, h}^m(\phi_h(s, t)^*))(\xi, x') \in C^0(B_{\delta_0} | B_{\rho, \delta}^m(h))$. Hence setting $\bar{D}_h^m(\phi_h(s, t)^*) = \bar{d}_h^m(\phi_h(s, t)^*; t, s; D_x, X')$, we define

$$(4.13) \quad D_h^m(\phi_h(s, t)^*) = \int_s^t E_h^m(\phi_h(\theta, t)^*) \tilde{D}_h^m(\phi_h(s, \theta)^*) d\theta,$$

where we again use Theorem 4.1. Then $U_h(t, s)$ defined by (4.6) satisfies (4.4). The uniqueness, the unitarity, and the relations (4.7) and (4.8) are proved in a way quite similar to [6]. \square

REMARK. As can be easily seen from the proof, this theorem also holds under the same assumption on the Hamiltonian as in [6].

From this theorem we can construct uniquely the global fundamental solution $U_h(t, s_0)$ of the equation (3.4) for $t \in R^1$ by $U_h(t, s_0) = U_h(t, s_N) U_h(s_N, s_{N-1}) \cdots U_h(s_1, s_0)$, where $s_k = s_0 + k(t - s_0)/(N + 1)$ ($k = 1, \dots, N$) with N being an integer such that $N \geq |t - s_0|/\delta_0 - 1$. The operator $U_h(t, s_0)$ thus constructed is unitary in $L^2(R^n)$ and obviously satisfies the relations (4.7) and (4.8) for any $t, \theta, r, s \in R^1$. Especially we have $U_h(t, s)^{-1} = U_h(s, t)$ for $t, s \in R^1$.

We now study a simple expression of $U_h(t, s_0)$, restricting ourselves to considering only the case $t \geq s_0$. (The other case can be dealt with similarly.) Then we have

$$(4.14) \quad U_h(t, s_0) = U_h(t, t_\nu) U_h(t_\nu, t_{\nu-1}) \cdots U_h(t_1, \tilde{T}) \cdot U_h(\tilde{T}, s_L) U_h(s_L, s_{L-1}) \cdots U_h(s_1, s_0),$$

where $t_{j+1} = t_j + \delta_0$ ($j = 0, 1, \dots$), $t_0 = \tilde{T}$, $t_{\nu+1} \geq t > t_\nu$, $s_{l-1} = s - \delta_0$ ($l = 1, \dots, L + 1$), $s_{L+1} = \tilde{T}$, and $s_1 > s_0 \geq s_1 - \delta_0$, \tilde{T} being a large number. Then L is determined only by s_0 and \tilde{T} . So if we can represent $U_h(t, t_\nu) \cdots U_h(t_1, \tilde{T})$ as a single conjugate Fourier integral operator, then the fundamental solution $U_h(t, s_0)$ is represented as a product of a finite number of conjugate Fourier integral operators independently of t . Before proving this we prepare a proposition.

Proposition 4.3. *Let $T (> T_0)$ be as in section 3 and let $0 < \delta_0 < 1$ be as in Theorem 4.2. Then:*

i) *For any (t, s) satisfying $T \leq s \leq t \leq s + \delta_0$ we have $\phi_h(s, t) \in P_{\delta_0}^{[s]}(C_1 \delta_0 \langle s \rangle^{-2-\epsilon}, C_1 \delta_0; h)$, where C_1 is the constant in Proposition 3.2.*

ii) *For $s \geq T$ and $t > s + \delta_0$, let $\nu \geq 1$ be an integer such that $s + (\nu + 1)\delta_0 \geq t > s + \nu\delta_0$, and put $t_j = s + j\delta_0$ for $j = 0, 1, \dots, \nu$ and $t_{\nu+1} = t$. Then the $(\nu + 1)$ -tuple $(\phi_h(s, t_1), \phi_h(t_1, t_2), \dots, \phi_h(t_\nu, t))$ of phase functions satisfies the condition (#) of section 2. Moreover we have $\phi_h(s, t_1) \# \phi_h(t_1, t_2) \# \cdots \# \phi_h(t_\nu, t) = \phi_h(s, t)$.*

Proof. i) is clear from Proposition 3.7-i).

ii) Put $\phi_{j,h} = \phi_h(t_{j-1}, t_j)$ for $j = 1, \dots, \nu + 1$. Then by Definition 3.6 and Proposition 3.5-(3.29) the equation (2.1) is equivalent to

$$(4.15) \quad \begin{cases} \text{i)} & X_v^j = y(t_{j-1}, t_j; X_v^{j-1}, \Xi_v^j) \\ \text{ii)} & \Xi_v^j = \eta(t_{j+1}, t_j; X_v^j, \Xi_v^{j+1}) \end{cases} \quad (j = 1, \dots, \nu)$$

with $X_v^0 = x; \Xi_v^{v+1} = \xi$; and

$$(4.16) \quad \begin{cases} X_{v,h}^j(x, \xi) = h^\delta X_v^j(h^{-\delta}x, h^\rho \xi) \\ \Xi_{v,h}^j(x, \xi) = h^{-\rho} \Xi_v^j(h^{-\delta}x, h^\rho \xi). \end{cases} \quad (j = 1, \dots, \nu)$$

Assume that $\{X_v^j, \Xi_v^j\}_{j=1}^\nu(x, \xi)$ is the solution of (4.15). Then from (4.15) and Proposition 3.2 we have

$$(4.17) \quad \begin{cases} \text{i)} & X_v^{j-1} = q(t_{j-1}, t_j; X_v^j, \Xi_v^j) \\ \text{ii)} & \Xi_v^{j+1} = p(t_{j+1}, t_j; X_v^j, \Xi_v^j). \end{cases} \quad (j = 1, \dots, \nu)$$

On the other hand using Proposition 3.2-ii) we have from (4.15)

$$(4.18) \quad \begin{cases} \text{i)} & X_v^j = q(t_j, t_{j-1}; X_v^{j-1}, \Xi_v^{j-1}) \\ \text{ii)} & \Xi_v^j = p(t_j, t_{j+1}; X_v^{j+1}, \Xi_v^{j+1}), \end{cases} \quad (j = 1, \dots, \nu)$$

where we put $\Xi_v^0 = \eta(t_1, t_0; x, \Xi_v^1)$ and $X_v^{v+1} = \gamma(t_\nu, t_{\nu+1}; X_v^\nu, \xi)$. Thus from (4.17)-i) and (4.18)-ii) we get

$$(4.19) \quad \begin{aligned} (X_v^j, \Xi_v^j) &= (q, p)(t_j, t_{j+1}; X_v^{j+1}, \Xi_v^{j+1}) \\ &= (q, p)(t_j, t_{v+1}; X_v^{v+1}, \xi) \quad (j = 0, 1, \dots, \nu). \end{aligned}$$

On the other hand from (4.17)-ii) and (4.18)-i) we have

$$(4.20) \quad \begin{aligned} (X_v^j, \Xi_v^j) &= (q, p)(t_j, t_{j-1}; X_v^{j-1}, \Xi_v^{j-1}) \\ &= (q, p)(t_j, t_0; x, \Xi_v^0) \quad (j = 1, \dots, \nu+1). \end{aligned}$$

Hence we have from (4.19) and (4.20)

$$(4.21) \quad \begin{cases} x = X_v^0 = q(t_0, t_{v+1}; X_v^{v+1}, \xi), \\ \xi = \Xi_v^{v+1} = p(t_{v+1}, t_0; x, \Xi_v^0), \end{cases}$$

from which we get

$$(4.22) \quad \begin{cases} X_v^{v+1} = \gamma(t_0, t_{v+1}; x, \xi), \\ \Xi_v^0 = \eta(t_{v+1}, t_0; x, \xi). \end{cases}$$

Combining this with (4.18) and (4.17) gives

$$(4.23) \quad \begin{cases} X_v^j = q(t_j, t_0; x, \eta(t_{v+1}, t_0; x, \xi)) = q(t_j, t_{v+1}; \gamma(t_0, t_{v+1}; x, \xi), \xi), \\ \Xi_v^j = p(t_j, t_0; x, \eta(t_{v+1}, t_0; x, \xi)) = p(t_j, t_{v+1}; \gamma(t_0, t_{v+1}; x, \xi), \xi). \end{cases}$$

Obviously this is C^∞ in (x, ξ) and satisfies (4.15). Thus we have proved ii). \square

Now we can prove the main result of this paper.

Theorem 4.4. *Let Assumption (A) be satisfied. Then the following as-*

sertions hold.

i) There exists uniquely the fundamental solution $U_h(t, s_0)$ ($t, s_0 \in R^1$) of the equation (3.4). This operator $U_h(t, s_0)$ satisfies the relation

$$(4.24) \quad \begin{cases} U_h(t, \theta)U_h(\theta, s) = U_h(t, s), & t, \theta, s \in R^1, \\ D_s U_h(t, s) - U_h(t, s)H_h(s, X, D_x) = 0, & t, s \in R^1, \end{cases}$$

and is extended to a unitary operator in $L^2(R^n)$.

ii) Let $\tilde{T} (> T)$ be sufficiently large. Then the fundamental solution $U_h(t, s)$ for $t \geq s \geq \tilde{T}$ is uniquely represented as a single conjugate Fourier integral operator with phase function $\phi_h(s, t)$ ($\in P_{\rho, \delta}^{[x]}(C_1 \langle s \rangle^{-\varepsilon}, C_1(t-s); h)$) and a symbol of class $C^1(A_{\tilde{T}} | B_{\rho, \delta}^0(h))$. More precisely, let $0 < \delta_0 < 1$ be sufficiently small as in Theorem 4.2. For $t \geq s \geq \tilde{T}$ and $t > s + \delta_0$, let $\nu \geq 1$ be an integer such that $s + (\nu + 1)\delta_0 \geq t > s + \nu\delta_0$, and put $t_j = s + j\delta_0$ for $j = 0, 1, \dots, \nu$ and $t_{\nu+1} = t$. Then we have

$$(4.25) \quad \begin{aligned} U_h(t, s)f(x) &= U_h(t, t_\nu)U_h(t_\nu, t_{\nu-1}) \cdots U_h(t_1, s)f(x) \\ &= 0_s - \iint e^{i(x \cdot \xi - \phi_h(s, t; x', \xi))} u_h(t, s; \xi, x') f(x') dx' d\xi \end{aligned}$$

for $f \in \mathcal{S}$, where $u_h(t, s; \xi, x') \in C^1(A_{\tilde{T}} | B_{\rho, \delta}^0(h))$ is uniquely determined. Thus the fundamental solution $U_h(t, s_0)$ of (3.4) for $t \geq s_0$ is represented as a product of a finite number of conjugate Fourier integral operators, the number depending on s_0 but not on t .

iii) For $t \geq s \geq \tilde{T}$ define

$$(4.26) \quad F_h^m(\phi_h(s, t)^*) = U_h(t, s) - E_h^m(\phi_h(s, t)^*),$$

where $m=0$ in case $0 \leq \delta \leq \rho \leq 1$ and $m=\infty$ in case $0 \leq \delta < \rho \leq 1$. Then we have for $t \geq s \geq \tilde{T}$

$$(4.27) \quad f_h^m(t, s) \equiv \sigma(F_h^m(\phi_h(s, t)^*)) \in C^1(A_{\tilde{T}} | B_{\rho, \delta}^m(h))$$

and

$$(4.28) \quad \begin{cases} \|F_h^0(\phi_h(s, t)^*)\|_{L^2 \rightarrow L^2} \leq C \langle s \rangle^{-\varepsilon}, \\ \|F_h^\infty(\phi_h(s, t)^*)\|_{L^2 \rightarrow L^2} \leq C_N h^N \langle s \rangle^{-1-\varepsilon} \end{cases}$$

for any $N \geq 1$, where the constants C and C_N are independent of t, s and h .

Proof. i) is already proved.

ii) We have only to prove the second equality of (4.25). Since $0 \leq t_j - t_{j-1} \leq \delta_0$, from Proposition 4.3-i) we have $\phi_{j,h} \equiv \phi_h(t_{j-1}, t_j) \in P_{\rho, \delta}^{[x]}(\tau_j, \sigma_0; h)$ for $\sigma_0 = C_1 \delta_0$ and $\tau_j = C_1 \delta_0 \langle t_{j-1} \rangle^{-2-\varepsilon}$ ($j = 1, \dots, \nu + 1$). So putting $\phi_{k,h} = x \cdot \xi$ and $\tau_k = 0$ for $k \geq \nu + 2$, we have for the sequence $\{\phi_{j,h}\}_{j=1}^\infty$ of phase functions $\phi_{j,h} \in P_{\rho, \delta}^{[x]}(\tau_j, \sigma_0; h)$

$$\begin{aligned}
 \bar{\tau}_\infty &\equiv \sum_{j=1}^\infty \tau_j \leq \sum_{j=1}^\infty C_1 \delta_0 \langle t_{j-1} \rangle^{-2-\varepsilon} \\
 &\leq 2C_1 \sum_{j=1}^\infty \delta_0 \langle \tilde{T} + (j-1)\delta_0 \rangle^{-2-\varepsilon} \\
 (4.29) \quad &\leq 2C_1 \int_0^\infty \langle \tilde{T} + \tau \rangle^{-2-\varepsilon} d\tau \leq \frac{2C_1}{1+\varepsilon} \langle T \rangle^{-1-\varepsilon}
 \end{aligned}$$

and

$$\begin{aligned}
 \sigma_0 \bar{\tau}_\infty &\equiv \sigma_0 \sum_{j=1}^\infty j \tau_j \leq 2C_1^2 \sum_{j=1}^\infty j \delta_0^2 \langle \tilde{T} + (j-1)\delta_0 \rangle^{-2-\varepsilon} \\
 (4.30) \quad &\leq 2C_1^2 \sum_{j=1}^\infty (\delta_0 \langle \tilde{T} + (j-1)\delta_0 \rangle^{-1-\varepsilon} + \delta_0^2 \langle \tilde{T} + (j-1)\delta_0 \rangle^{-2-\varepsilon}) \\
 &\leq 2C_1^2 \int_0^\infty (\langle \tilde{T} + \tau \rangle^{-1-\varepsilon} + \delta_0 \langle \tilde{T} + \tau \rangle^{-2-\varepsilon}) d\tau \\
 &\leq 2C_1^2 \varepsilon^{-1} \langle \tilde{T} \rangle^{-\varepsilon} + 2C_1^2 (1+\varepsilon)^{-1} \langle \tilde{T} \rangle^{-1-\varepsilon}.
 \end{aligned}$$

Thus if we take $\tilde{T} (\geq T)$ sufficiently large, then we have $2(\sigma_0 \bar{\tau}_\infty + \bar{\tau}_\infty) \leq \tau_0$ for some $0 \leq \tau_0 \leq 1/4$ independently of ν . This, together with Proposition 4.3-ii), shows that Theorem 2.6 is applicable to the product $U_h(t, t_\nu) \cdots U_h(t_1, T)$. Thus by Theorem 2.6 we have (4.25). The smoothness of $u_h(t, s)$ in (t, s) at $t=t_j$ follows from the uniqueness by taking another small $\delta_0 > 0$.

iii) From (4.26) and (3.48) we have for $t \geq s \geq T$ and $f \in \mathcal{S}$

$$\begin{aligned}
 F_h^m(\phi_h(s, t)^*)f &= U_h(t, s) (I - U_h(t, s)^{-1} E_h^m(\phi_h(s, t)^*))f \\
 (4.31) \quad &= U_h(t, s) \int_t^s \frac{d}{d\theta} [U_h(\theta, s)^{-1} E_h^m(\phi_h(s, \theta)^*)f] d\theta \\
 &= U_h(t, s) \int_s^t U_h(\theta, s)^{-1} G_h^m(\phi_h(s, \theta)^*)f d\theta \\
 &= \int_s^t U_h(t, \theta) G_h^m(\phi_h(s, \theta)^*)f d\theta.
 \end{aligned}$$

From this and Theorem 3.11–(3.50) follows (4.27), if we use the expression (4.25) and Theorem 2.6. The estimate (4.28) also follows from (4.31) and (3.54)–(3.55) of Theorem 3.11. \square

References

- [1] A.P. Calderón and R. Vaillancourt: *A class of bounded pseudo-differential operators*, Proc. Nat. Acad. Sci. U.S.A. **69** (1972), 1185–1187.
- [2] D. Fujiwara: *A construction of the fundamental solution for the Schrödinger equation*, J. Analyse Math. **35** (1979), 41–96.
- [3] D. Fujiwara: *Remarks on convergence of the Feynman path integrals*, Duke Math. J. **47** (1980), 559–600.
- [4] H. Kitada: *On a construction of the fundamental solution for Schrödinger equations*, J. Fac. Sci. Univ. Tokyo Sec. IA **27** (1980), 193–226.

- [5] H. Kitada: *Scattering theory for Schrödinger equations with time-dependent potentials of long-range type*, J. Fac. Sci. Univ. Tokyo, Sec. IA, **29** (1982), 353–369.
- [6] H. Kitada and H. Kumano-go: *A family of Fourier integral operators and the fundamental solution for a Schrödinger equation*, Osaka J. Math. **18** (1981), 291–360.
- [7] H. Kumano-go: *A calculus of Fourier integral operators on R^n and the fundamental solution for an operator of hyperbolic type*, Comm. Partial Differential Equations **1** (1976), 1–44.
- [8] H. Kumano-go, K. Taniguchi and Y. Tozaki: *Multi-products of phase functions for Fourier integral operators with an application*, Comm. Partial Differential Equations **3** (1978), 349–380.
- [9] K. Nishiwada: *Explicit formulae for solutions of Schrödinger equations with quadratic Hamiltonians*, Kyoto Univ. (preprint, 1980).

Department of Pure and Applied Sciences
College of General Education
University of Tokyo
Komaba, Meguro-ku, Tokyo
153 Japan