

LINEARLY COMPACT MODULES OVER HNP RINGS

Dedicated to Professor Hiroshi Nagao for his 60th birthday

HIDETOSHI MARUBAYASHI

(Received May 4, 1984)

Let R be a hereditary noetherian prime ring (an HNP ring for short) and let F be a non-trivial right Gabriel topology on R , i.e., F consists of essential right ideals of R (see §1 of [9]). Then R is a topological ring with elements of F as a fundamental system of neighborhoods of 0 . Let M be a topological right R -module with a fundamental system of neighborhoods of 0 consisting of submodules. Then M is called *F-linearly compact* (*F-l.c.* for short) if

- (i) it is Hausdorff,
- (ii) if every finite subset of the set of congruences $x \equiv m_\alpha \pmod{N_\alpha}$, where N_α are closed submodules of M , has a solution in M , then the entire set of the congruences has a solution in M .

This paper is concerned with *F-l.c.* modules over HNP rings in the case F is special. Let A be a maximal invertible ideal of R and let F_A be the right Gabriel topology consisting of all right ideals containing some power of A . Then we give, in §2, a complete algebraic structure of F_A -l.c. modules by using Kaplansky's duality theorem and basic submodules. From this result we get: " F_A -l.c. modules" \Rightarrow " F_A^ω -pure injective modules". This implication is not necessary to hold for any right Gabriel topology as it is shown in §3. It is established that there is a duality between F_A -l.c. modules and left \hat{R}_A -modules, where \hat{R}_A is the completion of R with respect to A (see Theorem 2.6). Main results in this paper were announced without proofs in [11].

Concerning our terminologies and notations we refer to [8] and [9].

1. Throughout this paper, R denotes an HNP ring with quotient ring Q and $K=Q/R \neq 0$. Let F be any non-trivial right Gabriel topology on R ; "*trivial*" means that either all modules are F -torsion-free or all modules are F -torsion. Then F consists of essential right ideals of R (see [9, p. 96]). Let I be any essential right ideal of R . Define $(R: I)_l = \{q \in Q \mid qI \subseteq R\}$. Similarly $(R: J)_r = \{q \in Q \mid Jq \subseteq R\}$ for any essential left ideal J of R . An ideal X of R is called *invertible* if $(R: X)_l X = R = X(R: X)_r$. In this case we have $(R: X)_l = (R: X)_r$, denoted by X^{-1} . For any right Gabriel topology F , put $Q_F = \bigcup (R: I)_l$ ($I \in F$), the ring of quotients of R with respect to F . The family F_l of

left ideals J of R such that $Q_F J = Q_F$ is a left Gabriel topology on R , which is called the left Gabriel topology corresponding to F . It is clear that $Q_F = Q_{F_i} = \cup (R: J)_r$, ($J \in F_i$). Define $\hat{R}_F = \varprojlim R/I$ ($I \in F$), the inverse limit of the modules R/I , and $\hat{R}_{F_i} = \varprojlim R/J$ ($J \in F_i$). Then both \hat{R}_F and \hat{R}_{F_i} are rings (see [16, §4]). Let M be an \bar{F} -torsion module. Then it is an \hat{R}_F -module as follows; for any $m \in M$ and $\hat{r} = ([r_I + I]) \in \hat{R}_F$, we define $m\hat{r} = mr_L$, where L is any element in F contained in $O(m) = \{r \in R \mid mr = 0\}$. Similarly, an F_i -torsion left module is an \hat{R}_{F_i} -module. In [7], we studied F -l.c. modules over a Dedekind prime ring. All results in [7, §2] are carried over F -l.c. modules over any HNP rings without any changes of the proofs. Here we pick up some of them which are frequently used in §2. Let $\eta: R \rightarrow \hat{R}_F$ be the canonical map and $\hat{F} = \{\hat{L}: \text{right ideals of } \hat{R}_F \mid \hat{L} \supset \eta(I)\hat{R}_F \text{ for some } I \in F\}$. Then \hat{R}_F is a topological ring with elements of \hat{F} as a fundamental system of neighborhoods of 0. For any \hat{R}_F -module, we can define the concept of \hat{F} -l.c. modules.

(1.1) *A module is an F-l.c. module if and only if it is an \hat{R}_F -module and is an \hat{F} -l.c. module (see Proposition 2.10 of [7]).*

Let M be an F -l.c. module. Then M^* means the left module of all continuous homomorphisms from M into K_F ($=Q_F/R$), where K_F is equipped with the discrete topology. It is evident that an element $f \in \text{Hom}_R(M, K_F)$ is continuous if and only if $\text{Ker } f$ is open. Let G be a left \hat{R}_{F_i} -module. Then we denote by G^* the right module $\text{Hom}_{\hat{R}_{F_i}}(G, K_F)$ and define its finite topology by taking the submodules $\text{Ann}(N) = \{f \in G^* \mid (N)f = 0\}$ as a fundamental system of neighborhoods of zero, where N runs over all finitely generated \hat{R}_{F_i} -submodules of G .

(1.2) (Kaplansky's duality theorem) *Let M be an F-l.c. module. Then M^* is a left \hat{R}_{F_i} -module and M is isomorphic to M^{**} as topological modules, where M^{**} is equipped with the finite topology induced by M^* as the above (see Lemma 2.11 and Theorem 2.12 of [7]).*

2. Let A be a maximal invertible ideal of R and let $F_A = \{I: \text{right ideal of } R \mid I \supseteq A^n \text{ for some } n > 0\}$, a right Gabriel topology. Then $F_{A_i} = \{J: \text{left ideal of } R \mid J \supseteq A^m \text{ for some } m > 0\}$. We denote the inverse limit of the modules R/A^n ($n=1, 2, \dots$) by \hat{R} . Then $\hat{R}_{F_A} = \hat{R} = \hat{R}_{F_{A_i}}$, and it is an HNP ring with the Jacobson radical $\hat{A} = A\hat{R} = \hat{R}A$ and with quotient ring $\hat{Q} = Q \otimes_R \hat{R}$ (see Lemma 1.2 and Theorem 1.1 of [8]). F_A -l.c. modules and F_A -torsion modules are said to be A -l.c. modules and A -primary modules, respectively. We note that $K_{F_A} = \hat{Q}/\hat{R}$, because $K_{F_A} = \cup A^{-n}/R = (\cup A^{-n}/R) \otimes_R \hat{R} \cong \cup \hat{A}^{-n}/\hat{R} = \hat{Q}/\hat{R}$.

In this section, we shall give a complete algebraic structure of A -l.c. modules. We can see from (1.1) that a module is A -l.c. if and only if it is an \hat{R} -module and an \hat{A} -l.c. module. If A is a maximal ideal of R , then \hat{R} is a Dede-

kind prime ring with unique maximal ideal \hat{A} . Thus, in this case, the algebraic structure of A -l.c. modules has been characterized in Theorem 3.4 of [7]. If A is not maximal ideal, then $A=M_1 \cap \dots \cap M_p$, where M_1, \dots, M_p are all maximal idempotent ideals of R and is a cycle, i.e., $O_r(M_1)=O_l(M_2), \dots, O_r(M_p)=O_l(M_1)$, where $O_r(M_1)=\{q \in Q \mid M_1q \subseteq M_1\}$ and $O_l(M_2)=\{q \in Q \mid qM_2 \subseteq M_2\}$. Furthermore, we have the following (see Theorem 1.1 of [8] and Lemma 4 of [10]):

- (a) $\hat{R} = \overbrace{(e_1\hat{R} \oplus \dots \oplus e_1\hat{R})}^{k_1} \oplus \dots \oplus \overbrace{(e_p\hat{R} \oplus \dots \oplus e_p\hat{R})}^{k_p}$, where each $e_i\hat{R}$ is a uniform right ideal of \hat{R} , e_i is idempotent in \hat{R} , $e_i\hat{R}/e_i\hat{A}$ is a simple module annihilated by M_i and k_i is the Goldie dimension of R/M_i .
- (b) $\hat{A} = \hat{M}_1 \cap \dots \cap \hat{M}_p$, where $\hat{M}_1, \dots, \hat{M}_p$ are all maximal idempotent ideals of \hat{R} and is a cycle, and $\hat{M}_i = M_i\hat{R} = \hat{R}M_i$ for each i ($1 \leq i \leq p$).

Lemma 2.1. Under the same notations as in (a) and (b), we have the following

- (1) $(e_i\hat{A}^{-1} + \dots + e_i\hat{A}^{-1}) + \hat{R} = O_l(\hat{M}_{i+1}) = O_r(\hat{M}_i)$ ($1 \leq i \leq p$ and $p+1=1$).
- (2) $\hat{R}e_i/\hat{A}e_i$ is left M_i -primary, i.e., each element of $\hat{R}e_i/\hat{A}e_i$ is annihilated by M_i .

Proof. Firstly we note that $\hat{A}^{-1} = (e_1\hat{A}^{-1} \oplus \dots \oplus e_1\hat{A}^{-1}) \oplus \dots \oplus (e_p\hat{A}^{-1} \oplus \dots \oplus e_p\hat{A}^{-1})$ and $\hat{A}^{-1} = O_l(\hat{M}_1) + \dots + O_l(\hat{M}_p)$, because $O_l(\hat{M}_i) = (\hat{R} : \hat{M}_i)_l$. Thus we have

- (c) $\hat{A}^{-1}/\hat{R} = (e_i\hat{A}^{-1} + \dots + e_i\hat{A}^{-1} + \hat{R})/\hat{R} \oplus \dots \oplus (e_p\hat{A}^{-1} + \dots + e_p\hat{A}^{-1} + \hat{R})/\hat{R}$, and
- (d) $\hat{A}^{-1}/\hat{R} = O_l(\hat{M}_1)/\hat{R} \oplus \dots \oplus O_l(\hat{M}_p)/\hat{R}$.

It is clear that $O_l(\hat{M}_i)/\hat{R}$ is M_i -primary. Since $e_i\hat{Q}/e_i\hat{A}$ is a uniform and injective \hat{R} -module, it is a uniform and injective R -module by Lemma 2.4 of [8]. Thus we have $e_i\hat{A}^{-1}/e_i\hat{R}$ is M_{i+1} -primary by periodicity theorem and (a) (see Theorem 22 of [4]). It follows that $(e_i\hat{A}^{-1})\hat{M}_{i+1} \subseteq e_i\hat{R} \subseteq \hat{R}$ and $e_i\hat{A}^{-1} \subseteq O_l(\hat{M}_{i+1})$. Thus (1) follows from (c) and (d).

(2) Since $O_r(\hat{M}_i) = O_l(\hat{M}_{i+1})$, we have $\hat{M}_i(e_i\hat{A}^{-1}) \subseteq \hat{R}$ by (1) and hence $\hat{M}_i e_i \subseteq \hat{A}e_i$. This implies that $\hat{R}e_i/\hat{A}e_i$ is M_i -primary as left modules.

Let M be an \hat{R} -module. Then write $M^* = \text{Hom}_{\hat{R}}(M, K_{F_A})$.

Lemma 2.2. Under the same notations as in (a) and (b), we have

- (1) for any positive integer n and any i ($1 \leq i \leq p$), $(e_i\hat{R}/e_i\hat{A}^n)^* = \hat{R}e_j/\hat{A}^n e_j$ for some j ($1 \leq j \leq p$).
- (2) $(e_i\hat{R})^* = \hat{Q}e_i/\hat{R}e_i = E(\hat{R}e_{i-1}/\hat{A}e_{i-1})$, the injective hull of $\hat{R}e_{i-1}/\hat{A}e_{i-1}$, where $1 \leq i \leq p$ and $i-1=p$ if $i=1$.
- (3) $(e_i\hat{Q})^* = \hat{Q}e_i$ for each i ($1 \leq i \leq p$).
- (4) $(e_i\hat{Q}/e_i\hat{R})^* = \hat{R}e_i$ for each i ($1 \leq i \leq p$).

These modules are all A -l.c. modules.

Proof. (1) Clearly $(e_i\hat{R}/e_i\hat{A}^n)^* = \hat{A}^{-n}e_i + \hat{R}/\hat{R} = \hat{A}^{-n}e_i/\hat{R}e_i$ by left multiplications of elements in $\hat{A}^{-n}e_i$. $\hat{A}^{-n}e_i/\hat{R}e_i$ is a uniserial module of length n with composition factor modules $\hat{A}^{-k}e_i/\hat{A}^{-(k-1)}e_i$ ($1 \leq k \leq n$ and $\hat{A}^{-0} = \hat{R}$). There is j ($1 \leq j \leq p$) such that $\hat{A}^{-n}e_i/\hat{A}^{-(n-1)}e_i \cong \hat{R}e_j/\hat{A}e_j$ and then $\hat{R}e_j/\hat{A}e_j \cong \hat{A}^{-n}e_i/\hat{R}e_i$ by the periodicity theorem.

(2) The first isomorphism is also obtained by left multiplication of elements in $\hat{Q}e_i$. The second isomorphism follows from the periodicity theorem.

(3) Let $x = xe_i$ be any element of $\hat{Q}e_i$. Then a mapping $\lambda_x: e_i\hat{Q} \rightarrow K_{F_A}$ given by $\lambda_x(y) = [xy + \hat{R}]$ ($y \in e_i\hat{Q}$) is a homomorphism. Assume that $\lambda_x = 0$ and $x \neq 0$. Then $x\hat{Q} = xe_i\hat{Q} \subseteq \hat{R}$, that is, $x \in \hat{R}$. Hence $\hat{R}x\hat{R}\hat{Q} \subseteq \hat{R}$. But $\hat{R}x\hat{R}$ contains a regular element in \hat{R} and so $\hat{R}x\hat{R}\hat{Q} = \hat{Q}$, a contradiction. Hence we may assume that $\hat{Q}e_i \subseteq (e_i\hat{Q})^*$. Conversely, let f be any non zero element in $(e_i\hat{Q})^*$ and let $f(e_i) = [q + \hat{R}]$, where $q = qe_i \in \hat{Q}$. Since $(f - \lambda_q)(e_i\hat{R}) = 0$, $f - \lambda_q$ induces an element $\overline{f - \lambda_q}$ in $(e_i\hat{Q}/e_i\hat{R})^*$. Since $\hat{Q}/\hat{R} = e_i\hat{Q}/e_i\hat{R} \oplus (1 - e_i)\hat{Q}/(1 - e_i)\hat{R}$, we may consider that $\overline{f - \lambda_q} \in (\hat{Q}/\hat{R})^*$. By Proposition A.3 of [8], $\hat{R} \cong (\hat{Q}/\hat{R})^*$. Hence $\overline{f - \lambda_q} = \lambda_r$ for some $r \in \hat{R}$ and $f - \lambda_q = \lambda_r$. So we get that $(e_i\hat{Q})^* \subset \hat{Q}e_i$ and therefore $(e_i\hat{Q})^* = \hat{Q}e_i$.

(4) The exact sequence $0 \rightarrow e_i\hat{R} \rightarrow e_i\hat{Q} \rightarrow e_i\hat{Q}/e_i\hat{R} \rightarrow 0$ induces the exact sequence $0 \rightarrow (e_i\hat{Q}/e_i\hat{R})^* \rightarrow (e_i\hat{Q})^* \rightarrow (e_i\hat{R})^* \rightarrow 0$, because K_{F_A} is injective. The assertion follows from (2) and (3). The left modules in (1) and (2) are artinian and A -primary. So they are A -l.c. modules in the discrete topology by Lemma 2.1 of [7] (as it has been pointed out in §1, all results in [7, §2] hold in F -l.c. modules over any HNP rings). \hat{R} is an A -l.c. modules by Lemma 2.4 of [7]. Thus it follows that $\hat{R}e_i$ is also an A -l.c. module. Finally consider the exact sequence $0 \rightarrow \hat{R}e_i \rightarrow \hat{Q}e_i \rightarrow \hat{Q}e_i/\hat{R}e_i \rightarrow 0$. $\hat{Q}e_i$ is a topological module by taking as a fundamental system of 0 the submodules $\{\hat{A}^n e_i \mid n = 0, \pm 1, \pm 2, \dots\}$. Hence $\hat{Q}e_i$ is an A -l.c. module by Proposition 9 of [20].

Following [9], a submodule L of a module M is called F^ω -pure if $MJ \cap L = LJ$ for any $J \in F_i$. Let F_ω be the right Gabriel topology of all essential right ideals of R . Then "an F_ω -pure submodule" is merely called a *pure submodule*.

Consider the following condition:

(e) *all finitely generated F and F_i -torsion modules are a direct sum of cyclic modules.*

This condition is satisfied by any topologies F and F_i on R if R has enough invertible ideals and so, especially, if R has a non zero Jacobson radical (see Corollary 3.4 and Theorems 4.12, 4.13 of [3]). If all F and F_i -torsion modules are of bounded orders, i.e., unfaithful modules, then this condition is satisfied, because every factor ring of an HNP ring is serial (Corollary 3.2 of [1]). Note that [9, Lemma 1.2] is still valid for topologies F and F_i on any HNP ring R satisfying the condition (e). Furthermore, if R has a nonzero Jacobson radical, then a submodule L of a module M is pure if and only if $Mc \cap L = Lc$ for any

regular element c in R by Proposition 3 of [19] and the remark to Theorem 3.6 of [15].

Lemma 2.3. *Let R be an HNP ring with the Jacobson radical A and A be a maximal invertible ideal of R . If as hort exact sequence $0 \rightarrow L \xrightarrow{\alpha} M \xrightarrow{\beta} N \rightarrow 0$ is pure, then $0 \rightarrow \text{Hom}_R(N, K) \xrightarrow{\beta^*} \text{Hom}_R(M, K) \xrightarrow{\alpha^*} \text{Hom}_R(L, K) \rightarrow 0$ is pure as left R -modules.*

Proof. Let c be any regular element of R and let $cf = g\beta$ be any element in $c\text{Hom}_R(M, K) \cap (\text{Hom}_R(N, K))\beta^*$, where $f \in \text{Hom}_R(M, K)$ and $g \in \text{Hom}_R(N, K)$. Since $g\beta\alpha(L) = 0$, we have $\alpha(L) \subset \text{Ker } g\beta = \text{Ker } cf$. There is a nature number n such that $Rc \supset A^n$. It follows that $0 = Rc f \alpha(L) \cong A^n f \alpha(L)$. Put $f \alpha(L) = X/R$, where X is a submodule of Q containing R . Then $A^n X \subseteq R$ and so $X \subseteq (R : A^n)_r = A^{-n} = (R : A^n)_l$. Thus we have $XA^n \subseteq R$. This implies that $f \alpha(L) A^n = 0$. Put $\bar{M} = M / \alpha(L) A^n$. Then $\bar{L} = \alpha(L) / \alpha(L) A^n$ is pure in \bar{M} , because L is pure in M . It follows from Theorem 3 of [13] and Theorem 1.3 of [14] that \bar{L} is a direct summand of \bar{M} , because \bar{L} is of bounded order. Thus we have the following sequence;

$$M \xrightarrow{\eta} \bar{M} = L \oplus \bar{M}_1 \xrightarrow{\pi} \bar{M}_1 \xrightarrow{f_1} K,$$

where η is a natural homomorphism, π is a projection map from \bar{M} to \bar{M}_1 (M_1 is a submodule of M) and f_1 is the map induced by f (note that $f \alpha(L) A^n = 0$). Put $h = f_1 \pi \eta$ and let x be any element of M . Write $x = x_1 + x_2$ ($x_1 \in \alpha(L)$ and $x_2 \in M_1$). Then $ch(x) = cf_1 \pi \eta(x) = cf_1(x_2) = cf(x_2)$. Since $x - x_2 \in \alpha(L) + \alpha(L) A^n \subseteq \alpha(L)$ and $cf \alpha(L) = 0$, we have $ch(x) = cf(x_2) = cf(x)$. Therefore $ch = cf$. By the construction of h , $h(\alpha(L)) = 0$. This entails that h induces a map $k: N \rightarrow K$ such that $k\beta = h$. Hence we have $cf = ch = ck\beta \in c(\text{Hom}_R(N, K))\beta^*$, as desired.

Theorem 2.4. *Under the same notations as in (a) and (b), a module is an A -l.c. module if and only if it is isomorphic to a direct product of modules of the following types:*

$e_i \hat{R} / e_i \hat{A}^n$ ($n=1, 2, \dots$), $E(e_i \hat{R} / e_i \hat{A})$, the injective hull of $e_i \hat{R} / e_i \hat{A}$, $e_i \hat{R}$ and $e_i (Q \otimes_R \hat{R})$ ($1 \leq i \leq p$).

Proof. The sufficiency follows from Proposition 1 of [20] and Lemma 2.2. Conversely let M be an A -l.c. module. Then M^* is a left \hat{R} -module by (1.2). So M^* has a basic submodule B by Theorem 2.1 of [8]. Then B is a direct sum of modules of types; $\hat{R}e_i / \hat{A}^n e_i$ and $\hat{R}e_i$ ($1 \leq i \leq p$) and $n=1, 2, \dots$), and M^*/B is a direct sum of modules of types; $E(\hat{R}e_i / \hat{A}e_i)$ and $(Q \otimes_R \hat{R})e_i$ (see Theorem 2.2 of [8]). Then from pure exact sequence $0 \rightarrow B \rightarrow M^* \rightarrow M^*/B \rightarrow 0$, we derive the pure exact sequence $0 \rightarrow (M^*/B)^* \rightarrow M^{**} \rightarrow B^* \rightarrow 0$ (as right \hat{R} -modules) by Lemma 2.3. By Lemma 2.2, $(M^*/B)^*$ is a direct product of modules of types; $e_i(Q \otimes_R \hat{R})$ and $e_i \hat{R}$. Here $e_i(Q \otimes_R \hat{R})$ is an injective \hat{R} -module.

Since $\hat{R} \cong \text{Hom}_R(K_{F_A}, K_{F_A}) \cong \text{Hom}_{\hat{R}}(\hat{Q}/\hat{R}, \hat{Q}/\hat{R})$ (see Lemma 1.5 and Proposition A.3 of [8]), \hat{R} is a pure injective \hat{R} -module by Propositions A.5, A.6 of [8] and Theorem 3.5, the remark to Proposition A.5 of [9], i.e., \hat{R} has the injective property relative to the class of pure exact sequences and so is $\epsilon_i \hat{R}$. Hence $(M^*/B)^{\#}$ is also pure injective. This entails that $M^{**} \cong (M^*/B)^{\#} \oplus B^{\#}$, and the assertion follows from (1.2) and Lemma 2.2.

Lemma 2.5. *Let M be a left \hat{R} -module and let m be any non zero element of M . Then there is an element f in $M^{\#}$ such that $(m)f \neq 0$.*

Proof. $\hat{R}m$ is a finite direct sum of modules of types; $\hat{R}e_i/\hat{A}^n e_i$ and $\hat{R}e_i$ by Theorems 2.1 and 2.2 of [8]. Thus the assertion follows from Lemma 2.2, because K_{F_A} is an injective \hat{R} -module.

Theorem 2.6. *Let R be an HNP ring and let A be a maximal invertible ideal of R . Then*


- (1) *Let M be any A -l.c. module. Then M^* is a left \hat{R} -module and $M \cong M^{**}$.*
- (2) *Let M be any left \hat{R} -module. Then $M^{\#}$ is an A -l.c. module in a certain topology and $M \cong M^{**}$ ($M^{\#}$ is equipped with the finite topology).*

Proof. (1) is clear from (1.2).

(2) Let M be any left \hat{R} -module. Then $M^{\#}$ is a direct product of modules of types in Theorem 2.4 (this is proved in the same way as in Theorem 2.4 by using basic submodules). Thus $M^{\#}$ is an A -l.c. module. Now $M^{\#}$ is equipped with the finite topology (it is not requested that $M^{\#}$ is an A -l.c. module in the finite topology). Let $\beta: M \rightarrow M^{**}$ be the natural map given by $((m)\beta)(f) = (m)f$, where $m \in M$ and $f \in M^{\#}$. Note that $(m)\beta \in M^{**}$, because $\text{Ker}(m)\beta = \{g \in M^{\#} | (m)g = 0\}$. By Lemma 2.5, β is a monomorphism. To prove that β is an epimorphism, let g be any element in M^{**} . Since $\text{Ker } g$ is open in $M^{\#}$, there is a finitely generated left module N of M such that $\text{Ker } g \supseteq \text{Ann}(N)$. Write

$$(*) \quad N \cong \sum_{i=1}^p \sum_j \hat{R}e_i/\hat{A}^{n_{ij}} e_i \oplus \sum \hat{R}e_k \quad (1 \leq k \leq p),$$

where $n_{ij} \geq 0$. Thus N is a left A -l.c. module by Lemma 2.2. Consider the following commutative diagram;



where η is a natural map, \bar{g} is a map induced by g and $\delta([f + \text{Ann}(N)]) = f|N$, the restriction map of f to N ($f \in M^{\#}$). Let h be any element of $N^{\#}$. Then

there is a natural number n such that $\hat{A}^n(N)h=0$, because N is finitely generated. This entails that $\text{Ker } h \cong \sum_{i=1}^{\ell} \sum_j \hat{A}^n e_i / \hat{A}^{n+ij} e_i \oplus \sum \hat{A}^n e_k$, open in N (in the topology given in Lemma 2.2). Thus we have $h \in N^*$ and hence $N^* = N^\#$. It follows from (1.2) that $\alpha: N \cong N^{\#\#}$. So, for the element $g\delta^{-1} \in N^{\#\#}$ there is an element $n \in N$ such that $(n)\alpha = g\delta^{-1}$, i.e., $((n)\alpha)\delta = g$. Now let x be any element in $M^\#$. Then we have $g(x) = g\eta(x) = ((n)\alpha)\delta\eta(x) = (n)\{\delta\eta(x)\} = (n)\{\delta[x + \text{Ann}(N)]\} = (n)x = ((n)\beta)(x)$. Hence $g = (n)\beta$, as desired.

3. In this section, we study relationships between F -l.c. modules and F^ω -pure injective modules in case F is special. A module G is F^ω -pure injective if it has the injective property relative to the class of F^ω -pure exact sequences. Let A be a maximal invertible ideal of R . Then F_A^ω -pure injective modules are just called A -pure injective modules. The cancellation set of A , $C(A)$, is defined to be $\{c \in R \mid cx \in A \Rightarrow x \in A\} = \{c \in R \mid cx \in A \Rightarrow x \in A\}$. By [6], R satisfies the Ore condition with respect to $C(A)$ and the local ring R_A of R at A is an HNP ring with Jacobson radical $AR_A = R_A A$. Note that a module T is A -primary if and only if it is an R_A -module and torsion as R_A -modules (see the proof of Lemma 2.4 of [8]). Since R_A is R -flat and the inclusion map: $R \rightarrow R_A$ is an epimorphism, we have the following

Lemma 3.1. (1) *An exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is A -pure, then the induced sequence $0 \rightarrow L \otimes_R R_A \rightarrow M \otimes_R R_A \rightarrow N \otimes_R R_A \rightarrow 0$ is exact and is pure as R_A -modules.*

(2) *if an exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ of R_A -modules is pure as R_A -modules, then it is A -pure.*

Proof. Use (3) in Lemma 1.2 of [9].

Lemma 3.2. *Let F be a right Gabriel topology on R satisfying the condition (e) and let G be any F^ω -pure injective module. Then $G = D \oplus H$, where D is an injective module, and H is F -reduced, F^ω -pure injective and F^ω -complete. In particular, H is an \hat{R}_F -module.*

Proof. The proof of Theorem 3.2 of [9] may be used unaltered to yield this lemma.

Proposition 3.3. *Let G be a reduced module, i.e., G has no non zero injective submodules. Then*

(1) *G is A -pure injective if and only if G is an R_A -module and is pure injective as R_A -modules.*

(2) *G is A -pure injective if and only if $G \cong \hat{G} = \varprojlim G/GA^n$.*

Proof. (1) It is clear from Lemmas 3.1 and 3.2. (2) follows from Theorems 3.2.4 and 3.3.3 of [18] and (1), because R_A is a bounded HNP ring.

From Theorem 2.4 and Proposition 3.3, we have

Corollary 3.4. *A-l.c. modules are A-pure injective modules.*

In general, it is not necessary to hold that (*F-l.c. modules*) \Rightarrow (*F^ω-pure injective modules*). We will end up this paper with giving a counter example. To do this, let *B* be an idempotent ideal of *R*. Then write

$$(f) \quad F_1 = \{I \mid IO_r(B) = O_r(B), I: \text{right ideal of } R\} . \\ F_2 = \{I \mid IO_l(B) = O_l(B), I: \text{right ideal of } R\} .$$

Then $F_{1l} = \{J \mid O_r(B)J = O_r(B), J: \text{left ideal of } R\}$, and $F_{2l} = \{J \mid O_l(B)J = O_l(B), J: \text{left ideal of } R\}$. Since $BO_r(B) = B$, $O_r(B)B = O_r(B)$, $BO_l(B) = O_l(B)$ and $O_l(B)B = B$, we have $F_{1l} = \{J \mid J \supseteq B\}$, $F_{2l} = \{I \mid I \supseteq B\}$, $F_1 \ni B$ and $F_{2l} \ni B$.

Proposition 3.5. *Under the same notations as in (f), let *G* be any module. Then*

(1) *G is an F₁-l.c. module if and only if it is a direct product of modules of types (R: J)_r/R, where J is a left ideal of R containing B.*

(2) *G is an F₂-l.c. module, then it is a direct sum of modules of types R/I (I \supseteq B).*

Proof. (1) The sufficiency is evident from Proposition 1 of [20], Proposition A.1 of [8] and Lemma 2.1 of [7]. Let *G* be an *F₁-l.c.* module. Then G^* is a left \hat{R}_{F_1} (= *R/B*)-module by (1.2). Since *R/B* is a serial ring, G^* is a direct sum of cyclic modules (see Theorem 1.2 and Corollary 3.2 of [1]). Write $G^* = \sum \oplus R/J_i$ ($J_i \supseteq B$) and then $G \cong G^{**} = \Pi(R: J_i)_r/R$ by (1.2).

(2) is clear, because any *F₂-l.c.* module is an \hat{R}_{F_2} (= *R/B*)-module.

Let *C* be an idempotent ideal of *R* such that

$$(g) \quad O_r(B) = O_l(C) .$$

Then $F_1 = \{I \mid I \supseteq C, I: \text{right ideal of } R\}$. Note that there exists an idempotent ideal *C* of *R* satisfying the condition (g) for any idempotent ideal *B* of *R* if *R* has enough invertible ideals. In the absense of the condition of having enough invertible ideals, we can easily find a pair of idempotent ideals *B* and *C* satisfying the condition (g) (see [4]).

Lemma 3.6. *Under the same notations as in (f) and (g), let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be an exact sequence. Then it is F₁^ω-pure if and only if the induced sequence*

$$(i) \quad 0 \rightarrow L/LB \rightarrow M/MB$$

is splitting exact.

Proof. The sufficiency is clear. To prove the necessity, assume that

the sequence is F_1^ω -pure. Then (ι) is exact and is F_1^ω -pure. Since a module is a finitely presented \bar{R} ($=R/B$)-module if and only if it is a finitely generated and F_1 -torsion module, (ι) is pure as R -modules in the sense of [19] by Proposition 3 of [19] and Lemma 1.2 of [9]. Furthermore, since $\bar{M}=M/MB$ satisfies Singh's conditions (I), (II) and (III), \bar{L} is h -pure in the sense of Singh (see Theorem 1.3 of [14]). Hence \bar{L} is a direct summand of \bar{M} by Theorem 3 of [13], because \bar{L} is of bounded order.

Proposition 3.7. *Under the same notations as in (f) and (g), a reduced module is F_1^ω -pure injective if and only if it is an R/B -module.*

Proof. This is clear from Lemmas 3.2 and 3.6.

EXAMPLE 3.8. Under the same notations as (f) and (g), R/C is an F_1 -l.c. module in the discrete topology. But it is not an F_1^ω -pure injective module.

Proof. R/C is an artinian and F_1 -torsion module. So it is an F_1 -l.c. module in the discrete topology by Lemma 2.1 of [7]. Assume that it is F_1^ω -pure injective. Then it is an R/B -module by Proposition 3.7. This implies that $B \subseteq C$ and so $O_i(C) = O_r(B) = (R: B)_r \supset (R: C)_r = O_r(C)$. Thus we have $C = O_i(C)C \supset O_r(C)C = O_r(C) \supset R$, a contradiction.

References

- [1] D. Eisenbud and P. Griffith: *Serial rings*, J. Algebra **17** (1971), 389–400.
- [2] D. Eisenbud and J.C. Robson: *Modules over Dedekind prime rings*, J. Algebra **16** (1970), 67–85.
- [3] D. Eisenbud and J.C. Robson: *Hereditary noetherian prime rings*, J. Algebra **16** (1970), 86–104.
- [4] K.R. Goodearl and R.B. Warfield, Jr: *Simple modules over Hereditary noetherian prime rings*, J. Algebra **57** (1979), 91–117.
- [5] I. Kaplansky: *Dual modules over a valuation ring I*, Proc. Amer. Math. Soc. **4** (1953), 212–219.
- [6] J. Kuzmanovich: *Localizations of HNP rings*, Trans. Amer. Math. Soc. **173** (1972), 137–157.
- [7] H. Marubayashi: *Modules over Dedekind prime rings III*, Osaka J. Math. **11** (1974), 547–558.
- [8] H. Marubayashi: *Completions of hereditary noetherian prime rings*, Osaka J. Math. **17** (1980), 391–406.
- [9] H. Marubayashi: *Pure injective modules over hereditary noetherian prime rings with enough invertible ideals*, Osaka J. Math. **18** (1981), 95–107.
- [10] H. Marubayashi: *Completion of overrings of HNP rings and its application* (unpublished).

- [11] H. Marubayashi: *Linearly compact modules over HNP rings*, Proceedings of the 4th Symposium on Semigroup held at Yamaguchi University, Japan, 1980.
- [12] S. Singh: *Modules over hereditary noetherian prime rings*, *Canad. J. Math.* **XXVII** (1975), 867–883.
- [13] S. Singh: *Some decomposition theorems in abelian groups and their generalizations*, Ring Theory, Proceedings of Ohio Univ. Conference, Lecture Notes in Math., Marcel Dekker, 25 (1976), 183–189.
- [14] S. Singh: *Some decomposition theorems in abelian groups and their generalizations II*, *Osaka J. Math.* **16** (1979), 45–55.
- [15] S. Singh and S. Talwar: *Pure submodules of modules over bounded (hnp)-rings*, *Arch. Math.* **30** (1978), 570–577.
- [16] B.O. Stenström: *On the completion of modules in an additive topology*, *J. Algebra* **16** (1970), 523–540.
- [17] B.O. Stenström: *Rings of quotients*, Berlin-Heidelberg-New York, 1975.
- [18] S. Talwar: *Modules over hereditary noetherian prime rings*, Ph. D. Thesis, Guru Nanak Dev Univ., India, 1979.
- [19] R.B. Warfield: *Purity and algebraic compactness for modules*, *Pacific J. Math.* **28** (1969), 699–719.
- [20] D. Zelinsky: *Linearly compact modules and rings*, *Amer. J. Math.* **75** (1953), 79–90.

Department of Mathematics
College of General Education
Osaka University
Toyonaka, Osaka 560, Japan