

ON REGULARITY OF SOLUTIONS TO N-ORDER DIFFERENTIAL EQUATIONS OF PARABOLIC TYPE IN BANACH SPACES

ANGELO FAVINI and HIROKI TANABE

(Received June 16, 1992)

1. Introduction

The present paper is concerned with the following initial value problem for linear parabolic differential equations of higher order in the time variable in some Banach space X :

$$(P) \begin{cases} u^{(n)}(t) + A_{n-1}u^{(n-1)}(t) + \cdots + A_1u'(t) + A_0u(t) = f(t), & 0 \leq t \leq T < \infty, \\ u^{(k)}(0) = u_k, & k = 0, \dots, n-1. \end{cases}$$

There is an extensive literature on this subject and many authors have defined and studied some types of parabolicity for the problem (P). We want to quote M.K. Balaev [3], A Favini & E. Obrecht [9], A. Favini & H. Tanabe [10], S.G. Krein [12], E. Obrecht [15],[16],[17], H. Tanabe [22], S. Ya. Yakubov [24],[25] and the references in [9].

There are also at least two recent monographs to be mentioned on polynomial operator pencils and their applications, that is A.S. Markus [14] and L. Rodman [18, pp. 83–88].

In [24] Yakubov devised a method of transforming a second order equation to a first order system so that the new system is of parabolic type and known results on parabolic evolution equations are applicable to it (see also Krein [12], Chapter III, Theorem 3.3). In [3] Balaev, using a method analogous to the one introduced by Yakubov [24], has given some conditions on the operators $A_j, j=0,1,\dots,n-1$, so that after an introduction of suitable new unknown functions the problem (P) produces a first order system of parabolic type.

In this paper using a device similar to the one described in [3], to be compared with the methods discussed in [18] for n -order differential equations with coefficients of bounded linear operators, we transform (P) to a first order system of parabolic type in the product space $X \times X \times \cdots \times X$. To this new system we apply the existence, uniqueness and maximal regularity theorems established in many papers. In this way

we extend the results of [3] in several directions, namely we allow A_j to have a non dense domain, make weaker assumptions on the links between the operators A_j (no fractional power are used), and discuss regularity properties for the solution. As references we indicate [1],[2],[19] since they contain a rather complete and definitive description of these results. Their affirmations are translated in our situation and involve the introduction of the real interpolation spaces.

We note here that Balaev [3] treats also nonlinear equations. He relies on a perturbation argument, and the essential role is played by the linear part.

On the other hand we show that also another approach works, i.e. we adapt the technique in [9] to deal with the problem (P) directly without reducing it to a first order system. In fact we prove that under our basic assumptions the operator pencil $P(z) = \sum_{j=0}^n z^j A_j$, z a complex variable, $A_n = I =$ the identity operator in X , is parabolic in the sense of [9],[15],[16]. In this manner we extend some results in [9] on second order equations to equations of arbitrary order using the method of [15],[16] relative to classical or strict solutions of (P). In the particular case $n=2$, we show that even a weaker assumption on the domains of A_j , $j=0,1$, implies the parabolicity of the pencil, provided that A_1 is multiplied by a sufficiently large number ρ .

We show in section 2 the existence, uniqueness and regularity of solutions of the problem (P) by transforming it to a first order system. In section 3 we show the parabolicity of the associated operator pencil $P(z)$ so that we may solve (P) directly.

Section 4 contains a number of examples and applications to partial differential equations, both in L^p spaces and in spaces of continuous functions. In this second case, we use the well known results by H.B. Stewart [20], G. Da Prato and P. Grisvard [6] and others.

We are also able to treat in the space L^p , $p \neq 2$, some problems for second order equations of the type considered by S. Chen and R. Triaggiani [4],[5]. Further, we indicate by means of a simple example how to solve some initial-boundary-value problems, more general than those treated by [4],[5] in the setup of continuous functions.

To finish with, we mention that in the paper [10] the case $n=3$ is studied even with operators suitably depending on t too, and some different applications are given.

Notation. We denote by $\|\cdot\|_X$ the norm in the Banach space X . If Z is a linear subspace of X , Z^a denotes its closure in X . If Y is another Banach space $\mathcal{L}(X, Y)$ is the space of all bounded linear operators from

X into Y and we use $\mathcal{L}(X)$ for $\mathcal{L}(X, X)$.

For a compact interval I of R , $\mathcal{C}(I; X)$, $\mathcal{C}^\sigma(I; X)$, $0 < \sigma < 1$, $\mathcal{C}^m(I; X)$, $m \in N$, denote respectively the space of all X -valued functions on I which are continuous, which are σ -Hölder continuous, and which are m -times continuously differentiable.

$\mathcal{C}^{1,\sigma}(I; X)$, $0 < \sigma < 1$, is the space of all continuously differentiable X -valued functions on I whose derivative belongs to $\mathcal{C}^\sigma(I; X)$. Further, for $T > 0$, $\mathcal{C}^\sigma((0, T]; X) = \bigcap_{0 < \epsilon \leq T} \mathcal{C}^\sigma([\epsilon, T]; X)$ and $\mathcal{C}^{1,\sigma}((0, T]; X)$ is analogously defined. For $T > 0$, $B(0, T; X)$ denotes the space of all bounded and strongly measurable functions from $(0, T]$ into X .

For the definition and the main properties of the real interpolation spaces $(X, Y)_{\theta, \infty}$, $0 < \theta < 1$, we refer to [23]. In particular, we shall be interested in the space $(X, \mathcal{D}(A))_{\theta, \infty} = D_{-A}(\theta, \infty)$, when $-A$ generates an analytic semigroup in X with its domain $\mathcal{D}(A)$ not necessarily everywhere dense in X .

2. Reduction to a first order equation.

In the sequel we shall always suppose that $A_j, j = 0, \dots, n-1$, are closed linear operators in the complex Banach space X , with domain $\mathcal{D}(A_j)$, $u_j, j = 0, \dots, n-1$, are given elements of X and f is a continuous function from $[0, T]$ into X .

We then specify what we mean by a solution to (P) and we give the definitions as follows.

DEFINITION 1. We say that $u \in \mathcal{C}([0, T]; X)$ is a *classical* solution to problem (P) if $u \in \mathcal{C}^n((0, T]; X)$, $u^{(j)}(t) \in \mathcal{D}(A_j)$, $j = 0, \dots, n-1$, $0 < t \leq T$, $u^{(j)} \in \mathcal{C}((0, T]; \mathcal{D}(A_j))$, $j = 0, \dots, n-1$, and (P) is satisfied.

DEFINITION 2. A function $u \in \mathcal{C}([0, T]; X)$ is a *strict* solution to (P) if $u \in \mathcal{C}^n([0, T]; X)$, $u^{(j)}(t) \in \mathcal{D}(A_j)$, $j = 0, \dots, n-1$, $0 \leq t \leq T$, $u^{(j)} \in \mathcal{C}([0, T]; \mathcal{D}(A_j))$, $j = 0, \dots, n-1$ and (P) is true in the sense

$$\begin{cases} u^{(n)}(t) + \sum_{j=0}^{n-1} A_j u^{(j)}(t) = f(t), & 0 \leq t \leq T, \\ u^{(j)}(0) = u_j, & j = 0, 1, \dots, n-1. \end{cases}$$

We transform (P) into the first order Cauchy problem

$$(1) \quad \begin{cases} V'(t) = \mathcal{A}V(t) + F(t), & 0 \leq t \leq T, \\ V(0) = V_0 \end{cases}$$

in the Banach space $E = X \times \dots \times X$, n times, $V = {}^t(v_0, v_1, \dots, v_{n-1})$, $V_0 = {}^t(u_0, u_1, \dots, u_{n-1})$, $F(t) = {}^t(0, 0, \dots, 0, f(t))$,

$$\mathcal{A} = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \\ -A_0 - A_1 - A_2 \dots - A_{n-1} \end{bmatrix}$$

As it is well known, the operator \mathcal{A} not necessarily generates an analytic semigroup in E or even a \mathcal{C}_0 semigroup.

We are interested in the first case and we shall use a trick similar to the one in [3]; see [12,24] for $n=2$. It consists in observing that if we formally apply the operator K , defined by

$$K = \begin{bmatrix} A_1 A_2 A_3 \dots A_{n-1} I \\ 0 A_2 A_3 \dots A_{n-1} I \\ 0 0 A_3 \dots A_{n-1} I \\ \dots \\ 0 0 0 \dots A_{n-1} I \\ 0 0 0 \dots 0 I \end{bmatrix},$$

to the equation (1), we obtain the new problem

$$(2) \quad \begin{cases} KV'(t) = K\mathcal{A}V(t) + KF(t), & 0 < t \leq T, \\ KV(0) = KV_0. \end{cases}$$

Call $KV(t) = Z(t)$, and assume that K have a bounded inverse. This hypothesis is not restrictive and is ensured by the existence of $A_j^{-1} \in \mathcal{L}(X)$, $j=0,1,\dots,n-1$, as we shall always suppose. Then (2) becomes

$$(3) \quad \begin{cases} Z'(t) = K\mathcal{A}K^{-1}Z(t) + KF(t), & 0 < t \leq T, \\ Z(0) = KV_0. \end{cases}$$

Our goal is to have some conditions on the operators A_j so that $K\mathcal{A}K^{-1}$ generates an analytic semigroup in E and thus we can apply to (3) known existence and maximal regularity results to be translated to problem (P) by means of $V=K^{-1}Z$.

The conditions on A_j read as follows.

HYPOTHESIS (H): There are n closed linear operators $B_j, j=1, \dots, n$, in the space X such that

$$A_k = B_{n-k} \cdots B_2 B_1, \quad k=0, 1, \dots, n-1.$$

HYPOTHESIS (K): $-B_k$ for $k=1, \dots, n$, generates an analytic semigroup in $X, \mathcal{D}(B_k) \subseteq \mathcal{D}(B_{k+1}), k=1, \dots, n-1$, and there are $\alpha, C > 0$ such that

$$\|B_{k+1}(B_k + z)^{-1}; \mathcal{L}(X)\| \leq C|z|^{-\alpha}, \quad k=1, \dots, n-1,$$

for all $z \in \Sigma = \{z \in \mathbb{C}; \operatorname{Re} z \geq -a_0 |\operatorname{Im} z|\}$, where a_0 is another positive constant.

It is easily verified that $K\mathcal{A}K^{-1}$ coincides with $\mathcal{A}_1 + \mathcal{A}_2$, where $\mathcal{D}(\mathcal{A}_1) = \mathcal{D}(B_n) \times \mathcal{D}(B_{n-1}) \times \dots \times \mathcal{D}(B_1), \mathcal{D}(\mathcal{A}_2) = X \times \mathcal{D}(B_n) \times \dots \times \mathcal{D}(B_2)$, and

$$\mathcal{A}_1 = \begin{bmatrix} -B_n & 0 & 0 & 0 & \cdots & 0 & 0 \\ -B_n - B_{n-1} & 0 & 0 & \cdots & 0 & 0 \\ -B_n - B_{n-1} - B_{n-2} & 0 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -B_n - B_{n-1} - B_{n-2} - B_{n-3} \cdots \cdots - B_2 - B_1 \end{bmatrix},$$

$$\mathcal{A}_2 = \begin{bmatrix} 0 & B_n & 0 & 0 & \cdots & 0 & 0 \\ 0 & B_n & B_{n-1} & 0 & \cdots & 0 & 0 \\ 0 & B_n & B_{n-1} & B_{n-2} & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & B_n & B_{n-1} & B_{n-2} & \cdots & B_3 & 0 \\ 0 & B_n & B_{n-1} & B_{n-2} & \cdots & B_3 & B_2 \\ 0 & B_n & B_{n-1} & B_{n-2} & \cdots & B_3 & B_2 \end{bmatrix}.$$

Assume (H),(K) and let $z \in \mathbb{C}, \operatorname{Re} z > 0, |z|$ large. Then the equation

$$(-\mathcal{A}_1 + z)x = F,$$

where $x = {}^t(x_1, \dots, x_n)$, $F = {}^t(f_1, \dots, f_n) \in E$, has the unique solution x for any $f_j \in X$, with x_k given by

$$\begin{aligned} x_k = & - \sum_{j=1}^{k-1} z^{k-1-j} (B_{n-k+1} + z)^{-1} (B_{n-k+2} + z)^{-1} \dots (B_{n-k+j} + z)^{-1} \\ & B_{n-j+1} (B_{n-j+1} + z)^{-1} f_j + \\ & + (B_{n-k+1} + z)^{-1} f_k; \end{aligned}$$

furthermore,

$$\|x; E\| \leq C|z|^{-1} \|f; E\|.$$

On the other hand, $K\mathcal{A}K^{-1} - z = \mathcal{A}_1 - z + \mathcal{A}_2 = (I - \mathcal{A}_2(z - \mathcal{A}_1)^{-1})(\mathcal{A}_1 - z)$ and then all is reduced to show that \mathcal{A}_2 is \mathcal{A}_1 -bounded, with \mathcal{A}_1 -bound equal to zero, so that perturbation theory can be invoked [11]. Now this follows using some lengthy calculations from the fact that all the entries in the matrix $\mathcal{A}_2(z - \mathcal{A}_1)^{-1}$ are sums of

$$z^j B_{k+1} (B_k + z)^{-1} (B_{k+1} + z)^{-1} \dots (B_{k+j} + z)^{-1} B_{k+j+1} (B_{k+j+1} + z)^{-1}.$$

We then apply our hypotheses (H) and (K) to check that the norm of this term in the space $\mathcal{L}(X)$ is bounded by $C|z|^{-\alpha}$, $|z|$ sufficiently large.

We have proved

Theorem 1. *Under (H), (K), the operator $K\mathcal{A}K^{-1}$ generates an analytic semigroup in the space E , with (possibly non dense) domain $\mathcal{D}(B_n) \times \mathcal{D}(B_{n-1}) \times \dots \times \mathcal{D}(B_1)$.*

REMARK 1. In view of [7], the assumptions (H), (K) are fulfilled if $A_{n-1} = B^{k_1}$, where $k_1 \in \mathbf{N}$, $-B$ generates a bounded analytic semigroup of angle $\pi/2$ in the Banach space X , and $A_{n-j} = B^{k_1 + \dots + k_j}$ with $k_j \in \mathbf{N}$, $0 < k_n < k_{n-1} < \dots < k_1$.

REMARK 2. Assumption (K) can be weakened to suppose B_{k+1} to be B_k -bounded and to have zero as its B_k -bound.

We refer to [8].

We now describe some consequences of Theorem 1 relative to the solvability of problem (P).

Theorem 2. Assume (H), (K) and let $0 < \sigma < 1$. If $f \in C^\sigma((0, T]; X) \cap C([0, T]; X)$ and $u_k \in \mathcal{D}(A_{k+1}) = \mathcal{D}(B_{n-k-1} \cdots B_2 B_1)$, with $B_{n-k-1} \cdots B_1 u_k (= A_{k+1} u_k) \in \mathcal{D}(B_{n-k})^a$, then (P) has a unique classical solution u such that $u^{(n)} \in C^\sigma((0, T]; X)$.

Proof. We apply [1, Theorem 4.1, p. 40] to problem [3]. We then know that if $KV_0 \in \mathcal{D}(K \mathcal{A} K^{-1})^a$ and $KF \in C^\sigma((0, T], E) \cap C([0, T]; E)$, then (3) has a unique classical solution $Z \in C^{1,\sigma}((0, T]; E)$. Since $K^{-1} \in \mathcal{L}(E)$ and $V = K^{-1}Z$, V is differentiable and $KV' = (KV)' = Z$. Recall that $V = {}^t(u, u', \dots, u^{(n-1)})$, $V' = {}^t(u', u^{(2)}, \dots, u^{(n)})$.

One also observes that $\mathcal{D}(K \mathcal{A} K^{-1})^a = \mathcal{D}(B_n)^a \times \cdots \times \mathcal{D}(B_1)^a$,

$$KV_0 = {}^t \left(\sum_{j=1}^{n-1} A_j u_{j-1} + u_{n-1}, \sum_{j=2}^{n-1} A_j u_{j-1} + u_{n-1}, \dots, A_{n-1} u_{n-2} + u_{n-1}, u_{n-1} \right),$$

$$KF(t) = {}^t(f(t), \dots, f(t)),$$

and the conditions above give $u_{n-1} \in \mathcal{D}(B_1)^a = \mathcal{D}(A_{n-1})^a$, $B_1 u_{n-2} + u_{n-1} \in \mathcal{D}(B_2)^a, \dots, A_2 u_1 + A_3 u_2 + \cdots + A_{n-1} u_{n-2} + u_{n-1} \in \mathcal{D}(B_{n-1})^a$, $A_1 u_0 + A_2 u_1 + \cdots + A_{n-1} u_{n-2} + u_{n-1} \in \mathcal{D}(B_n)^a$, that is, $u_{n-1} \in \mathcal{D}(B_1)^a$, $B_1 u_{n-2} \in \mathcal{D}(B_2)^a$, $B_2 B_1 u_{n-3} \in \mathcal{D}(B_3)^a, \dots, B_{n-2} \cdots B_2 B_1 u_1 \in \mathcal{D}(B_{n-1})^a$, $B_{n-1} B_{n-2} \cdots B_2 B_1 u_0 \in \mathcal{D}(B_n)^a$.#

Theorem 3. Assume Hypotheses (H), (K) and let $0 < \sigma < 1$. If $f \in C^\sigma([0, T]; X)$ and $u_j \in \mathcal{D}(A_j)$, $j = 0, 1, \dots, n-1$, with

$$f(0) - \sum_{j=0}^k A_j u_j \in \mathcal{D}(B_{n-k})^a, \quad k = 0, 1, \dots, n-1,$$

then problem (P) has a unique strict solution u such that $u^{(n)} \in C^\sigma((0, T]; X)$.

Proof. We consider problem [3] and apply [1, Theorem 5.1, p.42]. In this case their condition

$$KF(0) + K \mathcal{A} K^{-1} KV_0 \in \mathcal{D}(K \mathcal{A} K^{-1})^a$$

translates into $u_j \in \mathcal{D}(A_j)$ and $f(0) - A_0 u_0 \in \mathcal{D}(B_n)^a$, $f(0) - A_0 u_0 - A_1 u_1 \in \mathcal{D}(B_{n-1})^a, \dots, f(0) - \sum_{k=0}^{n-1} A_k u_k \in \mathcal{D}(B_1)^a$; and this is precisely our assumption.#

Corollary 1. Under (H), (K), if $\mathcal{D}(B_j)$ is everywhere dense in X , then problem (P) has one strict solution u with $u^{(n)} \in C^\sigma((0, T]; X)$, provided

that $f^{(n)} \in C^\sigma([0, T]; X)$ and $u_j \in \mathcal{D}(A_j) = \mathcal{D}(B_{n-j} \cdots B_2 B_1)$.

Theorem 4. *If the assumptions (H), (K) hold, $f \in C^\sigma([0, T]; X)$, $0 < \sigma < 1$, and $u_j \in \mathcal{D}(A_j) = \mathcal{D}(B_{n-j} \cdots B_2 B_1)$, with*

$$f(0) - \sum_{j=0}^k A_j u_j \in D_{-B_{n-k}}(\sigma, \infty) = (X, \mathcal{D}(B_{n-k}))_{\sigma, \infty},$$

then the corresponding solution u to problem (P) has the maximal regularity property $u^{(n)} \in C^\sigma([0, T]; X)$.

Proof. In view of [19, Proposition 1.15], since $\mathcal{D}(K\mathcal{A}K^{-1}) = \mathcal{D}(B_n) \times \cdots \times \mathcal{D}(B_1) = \mathcal{D}(\mathcal{A}_0)$, where \mathcal{A}_0 is defined by

$$\mathcal{A}_0({}^t(x_1, \dots, x_n)) = {}^t(-B_n x_1, -B_{n-1} x_2, \dots, -B_1 x_n), \quad x = {}^t(x_1, \dots, x_n) \in \mathcal{D}(\mathcal{A}_0),$$

the space $D_{K\mathcal{A}K^{-1}}(\sigma, \infty)$ and $D_{\mathcal{A}_0}(\sigma, \infty)$ coincide.

By Theorem 5.3 in [1, p.45], the derivative Z' of the solution Z to problem (3) belongs to $C^\sigma([0, T]; E)$ if and only if

$$KF(0) + K\mathcal{A}K^{-1}KV_0 \in D_{\mathcal{A}_0}(\sigma, \infty) = D_{-B_n}(\sigma, \infty) \times D_{-B_{n-1}}(\sigma, \infty) \times \cdots \times D_{-B_1}(\sigma, \infty).$$

Now $V = K^{-1}Z$ and hence $V' = {}^t(u', u^{(2)}, \dots, u^{(n)}) \in C^\sigma([0, T]; E)$ if the above condition is verified. On the other hand, this is equivalent to what we have established, as one can easily see. #

Next theorems shall study what happens when the data are more regular with respect to the “space”.

Theorem 5. *Under assumptions (H), (K), if $f \in \mathcal{C}([0, T]; X) \cap B(0, T, D_{-B_1}(\sigma, \infty))$, $0 < \sigma < 1$, and $A_{j+1}u_j = B_{n-j-1} \cdots B_2 B_1 u_j \in \mathcal{D}(B_{n-j})^a$, then there is one classical solution u to (P).*

Proof. In view of Sinestrari’s paper [19, Theorem 5.4, p. 59], if $KF \in \mathcal{C}([0, T]; E) \cap B(0, T; D_{K\mathcal{A}K^{-1}}(\sigma, \infty))$ and $KV_0 \in \mathcal{D}(B_n)^a \times \cdots \times \mathcal{D}(B_1)^a$, problem (3) has a unique classical solution Z . Hence $V = K^{-1}Z$ satisfies (2) and (1), too. The condition on KV_0 gives $u_{n-1} \in \mathcal{D}(B_1)^a$, $u_{n-1} + A_{n-1}u_{n-2} \in \mathcal{D}(B_2)^a, \dots, u_{n-1} + A_{n-1}u_{n-2} + \cdots + A_1u_0 \in \mathcal{D}(B_n)^a$, that are precisely the hypotheses that we have done on the initial values. #

Theorem 6. *Under hypotheses (H), (K), if $f \in C([0, T]; X) \cap$*

$B(0, T, D_{-B_1}(\sigma, \infty)), 0 < \sigma < 1$, and $u_j \in \mathcal{D}(A_j) = \mathcal{D}(B_{n-j} \cdots B_2 B_1)$, with $\sum_{j=0}^k B_{n-j} \cdots B_2 B_1 u_j \in \mathcal{D}(B_{n-k})^a$, then problem (P) has one strict solution.

Proof. In view of [19, Theorem 5.5, p. 60], we need only repeat the same argument as in Theorem 5, because $K \mathcal{A} V_0 \in \mathcal{D}(B_n)^a \times \cdots \times \mathcal{D}(B_1)^a$ if

$$A_0 u_0 \in \mathcal{D}(B_n)^a, A_0 u_0 + A_1 u_1 \in \mathcal{D}(B_{n-1})^a, \dots, \sum_{k=0}^{n-1} A_k u_k \in \mathcal{D}(B_1)^a. \#$$

Theorem 7. Suppose (H), (K) and $0 < \sigma < 1$, if $f \in C([0, T]; X) \cap B(0, T; D_{-B_1}(\sigma, \infty))$ and $\sum_{j=0}^k B_{n-j} \cdots B_2 B_1 u_j \in D_{-B_{n-k}}(\sigma, \infty)$, $k = 0, 1, \dots, n-1$, then the strict solution u to (P) has the regularity

$$A_j u^{(j)} \in C^\sigma([0, T]; X), j = 0, 1, \dots, n-1,$$

$$A_j u^{(j)} \in B(0, T; D_{-B_{n-j+1}}(\sigma, \infty)), j = 1, \dots, n,$$

(where $A_n = I$), and $\sum_{j=0}^k A_j u^{(j)} \in B(0, T; D_{-B_{n-k}}(\sigma, \infty))$, $k = 0, 1, \dots, n-1$.

Proof. According to [19, Theorem 5.5], our assumptions imply that

$$(KV)' \in B(0, T; D_{K \mathcal{A} K^{-1}}(\sigma, \infty)) \text{ and } K \mathcal{A} V \in C^\sigma([0, T]; E) \cap B(0, T; D_{K \mathcal{A} K^{-1}}(\sigma, \infty)).$$

Theorem 7 then translates this statement in the present situation. #

3. A direct approach to problem (P).

In recent years some authors have studied higher order abstract differential equations of parabolic type by a direct approach, without any reduction to a system. We quote here mainly the papers [9, 15, 16, 17].

We recall that according to these authors, the operator pencil P defined by

$$P: \mathcal{D}(P) (= \bigcap_{j=0}^{n-1} \mathcal{D}(A_j)) \rightarrow X, P(z)x = \sum_{j=0}^n z^j A_j x, x \in \mathcal{D}(P), A_n = I,$$

is parabolic if

a) there exist $K \in \mathbb{R}^+$, $\theta \in (\pi/2, \pi)$, such that

$$\Sigma = \{z \in \mathbb{C}; |z| \geq K, |\arg z| \leq \theta_0\} \subset \rho(P),$$

where $\rho(P) = \{w \in \mathbb{C}; \exists P(w)^{-1} \in \mathcal{L}(X)\}$.

b) $\|z^j A_j P(z)^{-1}; \mathcal{L}(X)\| \leq M$, for all $z \in \Sigma, j=0, 1, \dots, n$,

where M is a suitable positive constant.

In what follows we shall show that if $A_j = B_{n-j} \cdots B_2 B_1, j=0, \dots, n-1$, and Hypotheses (H), (K) hold, then the operator pencil $\sum_{j=0}^n z^j A_j, A_n = I$, is in fact parabolic, that is,

Theorem 8. *If Hypotheses (H), (K) are satisfied, then the pencil*

$$\sum_{j=0}^n z^j A_j, A_j = B_{n-j} \cdots B_2 B_1, j=0, 1, \dots, n-1, A_n = I,$$

is parabolic.

Proof. We observe that for all $z \in \mathbb{C}$,

$$\begin{aligned} P(z) &= z^n + z^{n-1} B_1 + z^{n-2} B_2 B_1 + \cdots + B_n B_{n-1} \cdots B_2 B_1 = \\ &= z^n + (z^{n-1} + z^{n-2} B_2 + \cdots + B_n B_{n-1} \cdots B_2) B_1 = \\ &= (z^{n-1} + z^{n-2} B_2 + \cdots + B_n B_{n-1} \cdots B_2)(z + B_1) - z(z^{n-2} + z^{n-3} B_3 + \cdots \\ &\quad + B_n B_{n-1} \cdots B_3) B_2. \end{aligned}$$

Let us define

$$P_j(z) = z^j + z^{j-1} B_{n-j+1} + z^{j-2} B_{n-(j-1)+1} B_{n-(j-1)} + z^{j-3} B_{n-j+3} B_{n-j+2} B_{n-j+1} + \cdots + B_n B_{n-1} \cdots B_{n-j+1}, j=2, \dots, n,$$

so that

$$P(z) = P_n(z) = P_{n-1}(z)(I - zP_{n-1}(z)^{-1}P_{n-2}(z)B_2(z+B_1)^{-1})(B_1+z),$$

and, in this same way, provided that the inverses exist,

$$P_{n-1}(z)^{-1} = (B_2+z)^{-1}(I - zP_{n-2}(z)^{-1}P_{n-3}(z)B_3(B_2+z)^{-1})^{-1} P_{n-2}(z)^{-1}$$

implies that

$$P_n(z) = P_{n-1}(z)(I - z(B_2+z)^{-1}(I - zP_{n-2}(z)^{-1}P_{n-3}(z)B_3(B_2+z)^{-1})^{-1} B_2(B_1+z)^{-1})(B_1+z).$$

Consider

$$\begin{aligned} I - zP_{n-2}(z)^{-1}P_{n-3}(z)B_3(B_2+z)^{-1} &= \\ = I - z(B_3+z)^{-1}(I - zP_{n-3}(z)^{-1}P_{n-4}(z)B_4(B_3+z)^{-1})^{-1} B_3(B_2+z)^{-1}. \end{aligned}$$

If we continue this process, we verify that all is reduced to check the invertibility of

$$I - zP_3(z)^{-1}P_2(z)B_{n-2}(B_{n-3} + z)^{-1}.$$

Now,

$$P_3(z) = P_2(z)(I - zP_2(z)^{-1}(z + B_n)B_{n-1}(B_{n-2} + z)^{-1})(B_{n-2} + z)$$

and

$$P_2(z) = (z + B_n)(I - z(z + B_n)^{-1}B_n(B_{n-1} + z)^{-1})(z + B_{n-1}).$$

In view of (H) and (K), $P_2(z)$ is in fact invertible for all $z \in \Sigma$, $|z|$ sufficiently large. Furthermore, there exists $P_3(z)^{-1} \in \mathcal{L}(X)$ and

$$P_3(z)^{-1} = (B_{n-2} + z)^{-1}(I - z(z + B_{n-1})^{-1}(I - z(z + B^n)^{-1}B_n(z + B_{n-1})^{-1})^{-1}B_{n-1}(B_{n-2} + z)^{-1})^{-1}P_2(z)^{-1}.$$

By induction, we obtain

$$P_n(z)^{-1} = (B_1 + z)^{-1}R(z)P_{n-1}(z)^{-1},$$

where

$$R(z) = (I - zP_{n-1}(z)^{-1}P_{n-2}(z)B_2(B_1 + z)^{-1})^{-1},$$

and also

$$\begin{aligned} \|P_n(z)^{-1}; \mathcal{L}(X)\| &\leq C|z|^{-n}, \quad z \in \Sigma, \quad |z| \text{ large,} \\ \|B_1P_n(z)^{-1}; \mathcal{L}(X)\| &\leq C|z|^{-(n-1)}, \quad z \in \Sigma, \quad |z| \text{ large.} \end{aligned}$$

On the other hand, if

$$Q_j(z) = z^j + z^{j-1}B_1 + z^{j-2}B_2B_1 + \dots + B_jB_{j-1}\dots B_1, \quad j = 1, \dots, n,$$

then

$$\begin{aligned} P_n(z) &= zQ_{n-1}(z) + B_nB_{n-1}\dots B_1 = \\ &= (z + B_n - zB_n(z^{n-2} + z^{n-3}B_1 + \dots + B_{n-2}\dots B_1)Q_{n-1}(z)^{-1})Q_{n-1}(z) \\ &= (z + B_n)(I - z(z + B_n)^{-1}B_nQ_{n-2}(z)Q_{n-1}(z)^{-1})Q_{n-1}(z). \end{aligned}$$

We deduce that

$$Q_{n-1}(z) = (z + B_{n-1})(I - z(z + B_{n-1})^{-1}B_{n-1}Q_{n-3}(z)Q_{n-2}(z)^{-1})Q_{n-2}(z)$$

and at last

$$Q_3(z) = (z + B_3)(I - z(z + B_3)^{-1}B_3Q_1(z)Q_2(z)^{-1})Q_2(z).$$

Now,

$$\begin{aligned} Q_2(z) &= (z + B_2)(z + B_1) - zB_2 \\ &= (I - zB_2(z + B_1)^{-1})(z + B_2)(z + B_1) \end{aligned}$$

and hence

$$\begin{aligned} Q_1(z)Q_2(z)^{-1} &= (z + B_2)^{-1}(I - zB_2(z + B_1)^{-1}(z + B_2)^{-1})^{-1}, \\ Q_3(z) &= (z + B_3)(I - z(z + B_3)^{-1}B_3(z + B_2)^{-1}(I - zB_2(z + B_1)^{-1}(z + B_2)^{-1})^{-1})Q_2(z). \end{aligned}$$

Assumptions (H), (K) ensure that $Q_2(z)$ and $Q_1(z)$ have bounded inverses for any $z \in \Sigma$, $|z|$ large, and

$$\begin{aligned} \|B_2 B_1 Q_2(z)^{-1}; \mathcal{L}(X)\| &\leq \text{Const}, \quad \|z^3 Q_3(z)^{-1}; \mathcal{L}(X)\| \leq \text{Const}, \\ \|B_3 B_2 B_1 Q_3(z)^{-1}; \mathcal{L}(X)\| &\leq \text{Const}, \quad \|z B_2 B_1 Q_3(z)^{-1}; \mathcal{L}(X)\| \leq \text{Const}, \\ z \in \Sigma, |z| &\geq K_0. \end{aligned}$$

By induction on $n \in \mathbf{N}$, we see that

$$P(z)^{-1} = Q_{n-1}(z)^{-1} H(z) (z + B_n)^{-1},$$

where

$$\begin{aligned} H(z) &= (I - z(z + B_n)^{-1} B_n Q_{n-2}(z) Q_{n-1}(z)^{-1})^{-1} \\ \text{and } \|H(z); \mathcal{L}(X)\| &\leq \text{Const}. \end{aligned}$$

Hence, since $Q_j(z)$ has the same structure as $P(z)$, we conclude that

$$\begin{aligned} \|z B_{n-1} \cdots B_1 P(z)^{-1}; \mathcal{L}(X)\| &\leq \text{Const}, \quad \|z^2 B_{n-2} \cdots B_1 P(z)^{-1}; \mathcal{L}(X)\| \\ &\leq \text{Const}, \cdots, \\ \|z^{n-2} B_2 B_1 P(z)^{-1}; \mathcal{L}(X)\| &= \|z \cdot z^{n-3} B_2 B_1 Q_{n-1}(z)^{-1} H(z) (z + B_n)^{-1}; \\ \mathcal{L}(X)\| &\leq \text{Const}, \quad z \in \Sigma, |z| \text{ sufficiently large.} \end{aligned}$$

The estimate

$$\|B_n B_{n-1} \cdots B_1 P(z)^{-1}; \mathcal{L}(X)\| \leq \text{Const}$$

follows by the preceding ones and $A_0 P(z)^{-1} = I - \sum_{j=1}^n z^j A_j P(z)^{-1}$.

Alternatively, we could prove by induction that

$$\|B_{j+2} Q_j(z) Q_{j+1}(z)^{-1}; \mathcal{L}(X)\| \leq C |z|^{-\alpha}, \quad j = 1, \dots, n-2, \text{ if } z \in \Sigma, |z| \text{ large,}$$

noting that

$$\begin{aligned} Q_j(z) Q_{j+1}(z)^{-1} &= (I - z(z + B_{j+1})^{-1} B_{j+1} Q_{j-1}(z) Q_j(z)^{-1})^{-1} (z + B_{j+1})^{-1} \\ &= (z + B_{j+1})^{-1} + z(z + B_{j+1})^{-1} B_{j+1} Q_{j-1}(z) Q_j(z)^{-1} Q_j(z) Q_{j+1}(z)^{-1}, \end{aligned}$$

where we used the equality $(I - S)^{-1} = I + S(I - S)^{-1}$. Hence, we have for $z \in \Sigma$, $|z|$ large,

$$Q_{j+1}(z)^{-1} = Q_j(z)^{-1} (I - z(z + B_{j+1})^{-1} B_{j+1} Q_{j-1}(z) Q_j(z)^{-1})^{-1} (z + B_{j+1})^{-1}.$$

With the aid of this we can show by induction

$$\|z^{j-k} B_k \cdots B_1 Q_j(z)^{-1}; \mathcal{L}(X)\| \leq C, \quad k = 0, \dots, j, \text{ if } z \in \Sigma, |z| \text{ large, } j = 1, \dots, n.$$

In particular,

$$\|z^{n-k}B_k \cdots B_1 P(z)^{-1}; \mathcal{L}(X)\| \leq C, \quad k=0, \dots, n, \text{ if } z \in \Sigma, |z| \text{ large. } \#$$

If we carefully watch out the demonstration given above to Theorem 8, we realize that, at least in the case $n=2$, the strong connection between the domains of B_1 and B_2 contained in the assumptions (H), (K) can be weakened to $\mathcal{D}(B_1) = \mathcal{D}(B_2)$, on condition that B_1 is multiplied by a sufficiently large constant ρ .

Notice that S. Chen and R. Triggiani [4,5] have studied, via reduction to a first order system in a suitable product space, parabolicity of (P), with $n=2$ and in a Hilbert space, when A_0 is a positive selfadjoint operator A and A_1 is comparable to $A^{1/2}$.

We have

Theorem 9. *Let $-B_i, i=1,2$ be the generator of an analytic semigroup in X , with $\mathcal{D}(B_1) = \mathcal{D}(B_2)$. If $\rho > 0$ is sufficiently large, then the operator pencil $P_\rho(z) = z^2 + z\rho B_1 + B_2 B_1$ is parabolic.*

Proof. We write

$$P_\rho(z) = (z + \rho^{-1} B_2)(z + \rho B_1) - z\rho^{-1} B_2 = (z + \rho^{-1} B_2)Q(z)(z + \rho B_1),$$

where

$$Q(z) = I - z\rho^{-1}(z + \rho^{-1} B_2)^{-1} B_2 (z + \rho B_1)^{-1}, \quad z \in \Sigma.$$

On the other hand, if $z \in \Sigma$ and $|z| \geq \rho_0 > 0$ suitably chosen,

$$\begin{aligned} \|(z + \rho B_1)^{-1}; \mathcal{L}(X)\| &\leq M|z|^{-1}, \quad \|(z + \rho^{-1} B_2)^{-1}; \mathcal{L}(X)\| \leq M|z|^{-1}, \\ \|B_2(z + \rho B_1)^{-1}; \mathcal{L}(X)\| &\leq M\rho^{-1}. \end{aligned}$$

The conclusion is then obvious. #

4. Applications and examples

APPLICATION 1. We want to recall some basic results for which we refer to [23, pp. 333–334]. Let Ω be a bounded C^∞ domain in $\mathbf{R}^q, q \in \mathbf{N}$, and let

$$A(D)u = \sum_{|\alpha|=2m} c_\alpha D^\alpha u, \quad c_\alpha \in \mathbf{C},$$

$m \in \mathbf{N}$, be a differential operator. We are also given a normal system $\{B_j, j=1, \dots, m\}$ of boundary operators complemented with respect to $A(D)$

so that Theorem 4.9.1 in [23, p. 334] is satisfied. Notice that then

$$\mathbf{B}_j u(x) = \sum_{|\alpha| \leq k_j} b_{j\alpha}(x) D^\alpha u(x), \quad b_{j\alpha} \in C^\infty(\partial\Omega),$$

$j = 1, \dots, m$, and $0 \leq k_1 < k_2 < \dots < k_m < 2m$. (In particular, these assumptions hold when $\mathbf{B}_j u = \frac{\partial^{k+j-1} u}{\partial \nu^{k+j-1}}$, $j = 1, \dots, m$, where $k \in \{0, 1, \dots, m\}$ is fixed).

A remarkable result by R. Seeley characterizes the complex interpolation spaces between $L^p(\Omega)$, $1 < p < \infty$, and

$$H_{p, \{\mathbf{B}_j\}}^{2m} = \{u \in H_p^{2m}(\Omega); \mathbf{B}_j u|_{\partial\Omega} = 0, j = 1, \dots, m\}.$$

Precisely, [23, p. 321], if there does not exist a number k_j , $j = 1, \dots, m$, such that $2m\theta - 1/p = k_j$, then

$$\begin{aligned} [L^p(\Omega), H_{p, \{\mathbf{B}_j\}}^{2m}(\Omega)]_\theta &= H_{p, \{\mathbf{B}_j\}}^{2m\theta}(\Omega) \\ &= \{u \in H_p^{2m\theta}(\Omega); \mathbf{B}_j u|_{\partial\Omega} = 0 \text{ for } k_j < 2m\theta - 1/p\}. \end{aligned}$$

Furthermore, the operator $-A$, where

$$\mathcal{D}(A) = H_{p, \{\mathbf{B}_j\}}^{2m}(\Omega), \quad (Au)(x) = A(D)u(x),$$

generates an analytic semigroup in $L^p(\Omega)$ and the imaginary powers A^{it} of A , t real, are bounded linear operators in $L^p(\Omega)$ such that $\|A^{it}\|_{\mathcal{L}(L^p(\Omega))} \leq C e^{\gamma|t|}$, for certain $C > 0$ and $\gamma \geq 0$, [23, p. 334]. Therefore, if $2m\theta = h$ is a positive integer $< 2m$, in view of [23, p. 103],

$$[L^p(\Omega), H_{p, \{\mathbf{B}_j\}}^{2m}(\Omega)]_{h/2m} = \{u \in H_p^h(\Omega) = W^{h,p}(\Omega); \mathbf{B}_j u|_{\partial\Omega} = 0 \text{ for } k_j < h - 1/p\} = \mathcal{D}(A^{h/2m}).$$

(It is not too restrictive to suppose A to be a positive operator in $L^p(\Omega)$).

These results indicate a possible choice of the operators B_1, \dots, B_n appearing in assumptions (H), (K) above. We start with $B_1 = A$ and take as B_2 another operator of the same type as A , of lower order and such that $\mathcal{D}(B_2) \supseteq \mathcal{D}(A^{h/2m})$. Then we repeat this procedure to B_3, \dots, B_n .

To clarify the idea, let $n = 2$,

$$\mathcal{D}(B_1) = \{u \in H_p^4(\Omega); u|_{\partial\Omega} = \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} = 0\}, \quad B_1 u = \Delta^2 u, \quad u \in \mathcal{D}(B_1),$$

$$\mathcal{D}(B_2) = \{u \in H_p^2(\Omega); \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} = 0\}, \quad B_2 u = -\Delta u, \quad u \in \mathcal{D}(B_2).$$

Since $-B_1, -B_2$ generate analytic semigroups in $L^p(\Omega)$, $1 < p < \infty$ and

$$[L^p(\Omega), \mathcal{D}(B_1)]_{1/2} = \{u \in H_p^2(\Omega); u|_{\partial\Omega} = \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} = 0\} \subset \mathcal{D}(B_2),$$

we deduce that (H) and (K) hold and thus we can treat the problem

$$\left\{ \begin{array}{l} \frac{\partial^2 u}{\partial t^2} + \Delta^2 \frac{\partial u}{\partial t} - \Delta^3 u = f(t, x), \quad 0 < t \leq T, \quad x \in \Omega, \\ u(0, x) = u_0(x), \quad \frac{\partial u}{\partial t}(0, x) = u_1(x), \quad x \in \Omega, \\ \frac{\partial u}{\partial t}(t, \cdot) \in \mathcal{D}(B_1), \quad u(t, \cdot) \in \mathcal{D}(B_1) \cap H_p^6(\Omega), \\ \frac{\partial \Delta^2 u(t, \cdot)}{\partial \nu} \Big|_{\partial\Omega} = 0, \quad t > 0. \end{array} \right.$$

Of course, one could perturb the differential operators by means of derivatives of lower order and to this purpose we refer to [9]. Henceforth we describe other examples that could be generalized in various directions, too.

EXAMPLE 1. Referring to Remark 1, let B be the operator in $L^p(\Omega) = X$, $1 < p < \infty$, defined by

$$\mathcal{D}(B) = W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega), \quad Bu = -\Delta u, \quad u \in \mathcal{D}(B).$$

Then $-B$ generates an analytic semigroup in X satisfying the spectral properties in [7], so that for all $q \in \mathbf{N}$ the operator $-B_q$ given by

$$\mathcal{D}(B_q) = \{u \in W^{2q,p}(\Omega); u|_{\partial\Omega} = \Delta u|_{\partial\Omega} = \dots = \Delta^{q-1} u|_{\partial\Omega} = 0\}, \quad B^q = (-\Delta)^q$$

generates a holomorphic semigroup in X , too and one can apply Remark 1. This allows to treat the corresponding initial-boundary-value problem relative to the equation

$$\frac{\partial^n u}{\partial t^n} + c_1 (-\Delta)^{q_1} \frac{\partial^{n-1} u}{\partial t^{n-1}} + \dots + c_n (-\Delta)^{q_1 + \dots + q_n} u = f(t, x),$$

where $c_j > 0$ and $q_1 > q_2 > \dots > q_n > 0$.

In our results the initial conditions u_j shall belong to some interpolation

spaces that have been characterized in several papers by A. Lunardi [13], P. Acquistapace and B. Terreni [1,2] and others.

On the other hand, also the spaces $(X, \mathcal{D}(B_j))_{\theta, q}$, $1 < q < \infty$, have a complete description ([23]) and $(X, \mathcal{D}(B_j))_{\theta, q} \subseteq (X, \mathcal{D}(B_j))_{\theta, \infty}$. In this way we have a wide class of initial values to which our results apply directly.

EXAMPLE 2. Let Ω be a bounded domain in R^q , $q \in N$, with a smooth boundary $\partial\Omega$, $T > 0$, $\Sigma = (0, T] \times \partial\Omega$. Consider the problem

$$(4) \quad \begin{cases} u_{tt} + \Delta^2 u_t - \Delta^3 u = f(x, t), & 0 \leq t \leq T, \quad x \in \Omega, \\ u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x), & x \in \Omega, \\ u|_{\Sigma} = \frac{\partial u}{\partial \nu}|_{\Sigma} = \Delta^2 u|_{\Sigma} = 0 \quad \text{or} \quad u|_{\Sigma} = \frac{\partial u}{\partial \nu}|_{\Sigma} = \frac{\partial \Delta^2 u}{\partial \nu}|_{\Sigma} = 0, \end{cases}$$

This time we take $X = L^2(\Omega)$,

$$\mathcal{D}(B_1) = \{u \in H^4(\Omega); u|_{\partial\Omega} = \frac{\partial u}{\partial \nu}|_{\partial\Omega} = 0\}, \quad B_1 u = \Delta^2 u, \quad u \in \mathcal{D}(B_1),$$

$$\mathcal{D}(B_2) = \{u \in H^2(\Omega); u|_{\partial\Omega} = 0\} \quad \text{or} \quad \{u \in H^2(\Omega); \frac{\partial u}{\partial \nu}|_{\partial\Omega} = 0\},$$

$$B_2 u = -\Delta u, \quad u \in \mathcal{D}(B_2).$$

Then $\mathcal{D}(B_2) \supset \mathcal{D}(B_1^{1/2}) = [X, \mathcal{D}(B_1)]_{1/2} = H_0^2(\Omega)$, and therefore all our general results can be applied.

Of course, different exponents in the powers of the laplacian occuring in (4) are possible, provided that the assumptions (H) and (K) remain satisfied.

EXAMPLE 3. We take the same equation as in (4) but with other boundary conditions. Precisely, we assume $X = L^p(\Omega)$, $1 < p < \infty$,

$$\mathcal{D}(B_1) = \{u \in W^{4,p}(\Omega); u|_{\partial\Omega} = \frac{\partial u}{\partial \nu}|_{\partial\Omega} = 0\}, \quad B_1 u = \Delta^2 u, \quad u \in \mathcal{D}(B_1),$$

$$\mathcal{D}(B_2) = W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega), \quad B_2 u = -\Delta u.$$

Since $\mathcal{D}(B_1^{1/2}) = W_0^{2,p}(\Omega) \subset \mathcal{D}(B_2)$, the moment inequality (or interpolation theory) guarantees that

$$\|B_2(z + B_1)^{-1}; \mathcal{L}(X)\| \leq M |z|^{-1/2}, \quad z \in \rho(-B_1).$$

EXAMPLE 4. Let $a_0, a_1, a_2 > 0$. Consider in the space $X = L^p(\Omega)$, $1 < p < \infty$, the equation

$$\frac{\partial^3 u}{\partial t^3} - a_2 \Delta^3 \frac{\partial^2 u}{\partial t^2} - a_1 \Delta^5 \frac{\partial u}{\partial t} + a_0 \Delta^6 u = f(t, x), \quad 0 \leq t \leq T, \quad x \in \Omega.$$

Let us define the operator B_1 by

$$\mathcal{D}(B_1) = W^{6,p}(\Omega) \cap W_0^{3,p}(\Omega), \quad B_1 u = -a_2 \Delta^3 u, \quad u \in \mathcal{D}(B_1), \quad 1 < p < \infty,$$

so that

$$\mathcal{D}(B_1^{2/3}) = [X, \mathcal{D}(B_1)]_{2/3} = \{u \in W^{4,p}(\Omega); u|_{\partial\Omega} = \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} = \frac{\partial^2 u}{\partial \nu^2} \Big|_{\partial\Omega} = 0\}.$$

Let us choose B_2 with

$$\mathcal{D}(B_2) \supseteq \mathcal{D}(B_1^{2/3}), \quad B_2 u = (a_1/a_2) \Delta^2 u, \quad u \in \mathcal{D}(B_2),$$

such that $-B_2$ generates an analytic semigroup in X (e.g., $\mathcal{D}(B_2) = W^{4,p}(\Omega) \cap W_0^{2,p}(\Omega)$, or else). Finally define B_3 so that we have

$$\mathcal{D}(B_2^{1/2}) \subset \mathcal{D}(B_3) \subset W^{2,p}(\Omega), \quad B_3 u = -(a_0/a_1) \Delta u, \quad u \in \mathcal{D}(B_3),$$

and $-B_3$ generates an analytic semigroup in X .

Of course, if $\mathcal{D}(B_2) = W^{4,p}(\Omega) \cap W_0^{2,p}(\Omega)$, then $\mathcal{D}(B_2^{1/2}) = \{u \in W^{2,p}(\Omega); u|_{\partial\Omega} = \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} = 0\} = W_0^{2,p}(\Omega)$, and hence the elements of $\mathcal{D}(B_3)$ could satisfy either Dirichlet or Neumann boundary conditions.

EXAMPLE 5. Consider the initial-boundary-value problem

$$(5) \quad \begin{cases} u_{tt} - \rho \Delta u_t + \Delta^2 u = f = f(t, x), & 0 \leq t \leq T, \quad x \in \Omega, \\ u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x), & x \in \Omega, \\ u(t, \cdot)|_{\partial\Omega} = u_t(t, \cdot)|_{\partial\Omega} = 0, & 0 < t \leq T, \\ \Delta u(t, \cdot)|_{\partial\Omega} = 0, & 0 < t \leq T, \end{cases}$$

where Ω is a bounded domain in R^q , $q \in N$, with a smooth boundary $\partial\Omega$. Now we take $X = L^p(\Omega)$, $1 < p < \infty$, and define B_1, B_2 by

$$\mathcal{D}(B_1) = W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega), \quad B_1 u = -\Delta u, \quad u \in \mathcal{D}(B_1), \quad 1 < p < \infty, \quad B_2 = B_1.$$

Then Theorem 9 permits to handle (5) in the space X for any $\rho > 0$

sufficiently large. In this way we extend to $p \in (1, \infty)$ the treatment in $L^2(\Omega)$ given in the paper [5]. Notice that other boundary conditions can be associated to the laplacian to generate well posed problems.

APPLICATION 2. Let Ω be a bounded open subset of R^q , $q \in \mathbf{N}$, whose boundary $\partial\Omega$ is sufficiently smooth (see [20]), even if one could also treat the case Ω unbounded without too many changes. Let

$$A(x, D_x) = \sum_{|\alpha|=2m} a_\alpha(x) D^\alpha, \quad D^\alpha = \frac{\partial^{\alpha_1} \dots \partial^{\alpha_q}}{\partial x_1^{\alpha_1} \dots \partial x_q^{\alpha_q}}, \quad \alpha = (\alpha_1, \dots, \alpha_q),$$

be a uniformly strongly elliptic operator satisfying the roots condition, a_α being a real-valued continuous function on $\bar{\Omega}$. $\mathcal{C}(\bar{\Omega})$ shall denote the space of all complex-valued functions continuous on $\bar{\Omega}$. If one defines the operator A by means of

$$\mathcal{D}(A) = \{u \in \mathcal{C}(\bar{\Omega}) \cap W_{loc}^{2m,s}(\Omega); A(\cdot, D)u \in \mathcal{C}(\bar{\Omega}), D^\beta u = 0 \text{ on } \partial\Omega, 0 \leq |\beta| < m\},$$

where $A(\cdot, D)u$ is intended in the sense of distributions, $Au = A(\cdot, D)u$, $u \in \mathcal{D}(A)$, with $s > q$, then ([20, p. 152]) $-A$ generates an analytic semigroup in $\mathcal{C}(\bar{\Omega})$, with non dense domain.

Further, it is known [20] that there exist $M > 0$, $\lambda_0 \geq 0$, $\varepsilon > 0$ such that

$$\|u; \mathcal{C}(\bar{\Omega})\| + \sum_{0 < |\beta| < 2m} |z|^{-|\beta|/2m} \|D^\beta u; \mathcal{C}(\bar{\Omega})\| \leq M|z|^{-1} \|(A+z)u; \mathcal{C}(\bar{\Omega})\|,$$

for all complex numbers z , $|z| \geq \lambda_0$, $|\arg z| \leq \pi/2 + \varepsilon$. For sake of brevity, let $\mathcal{C}(\bar{\Omega}) = X$. Then we infer that

$$\|D^\beta u; X\| \leq M|z|^{-(1-|\beta|/2m)} \|Au; X\| + M|z|^{|\beta|/2m} \|u; X\|$$

and hence, taking $M|z|^{-(1-|\beta|/2m)} = \eta$, there is $C(\eta)$ such that

$$\|D^\beta u; X\| \leq \eta \|Au; X\| + C(\eta) \|u; X\|, \quad 0 \leq \beta < 2m, \text{ for any } u \in \mathcal{D}(A).$$

It follows that a differential operator of order $< 2m$ and with its coefficients continuous on $\bar{\Omega}$ defines an operator A -bounded and with A -bound equal to 0.

We continue observing that if $k \in \mathbf{N}$, $0 < k < m$, and $b_p(\cdot)$ is continuous on $\bar{\Omega}$, $|\beta| \leq 2k$, then the operator B given by

$$\mathcal{D}(B) = \{u \in X; u \in W_{loc}^{2k,s}(\bar{\Omega}); B(\cdot, D)u \in X, D^\beta u = 0 \text{ on } \partial\Omega, 0 \leq |\beta| < k\},$$

$$Bu = \sum_{|\beta| \leq k} b(\cdot) D^\beta u, \quad u \in \mathcal{D}(B),$$

has properties analogous to the ones of A , and it is A -bounded; moreover its A -bound is zero. These remarks indicate how it is possible to treat by our methods the problem

$$\begin{cases} u_{tt} + A(x,D)u_t + B(x,D)u = f(t,x), & 0 \leq t \leq T, \quad x \in \Omega, \\ u(0, \cdot) = u_0, \quad u_t(0, \cdot) = u_1, \\ + \text{ boundary conditions} \end{cases}$$

in the space of continuous functions, as well as its generalization to equations of order n in time.

EXAMPLE 6. If k is a positive integer, let us denote by $\mathcal{C}^k([0,1]) = \{u \in \mathcal{C}([0,1]); u \text{ is } k\text{-times continuously differentiable on } [0,1]\}$. We also recall the following results by [2].

$$\text{Let } \alpha_i, \beta_i \geq 0, \alpha_i + \beta_i > 0, i=0,1, a, b, c \in \mathcal{C}([0,1]; \mathbf{R}), \inf_{x \in [0,1]} a(x) > 0.$$

Define an operator K by

$$\begin{aligned} \mathcal{D}(K) &= \{u \in \mathcal{C}^2([0,1]); \alpha_0 u(0) - \beta_0 u'(0) = \alpha_1 u(1) + \beta_1 u'(1) = 0\}, \\ Ku &= a(\cdot)u'' + b(\cdot)u' + c(\cdot)u - \omega_0 u, \quad \omega_0 \in \mathbf{R}, \quad u \in \mathcal{D}(K). \end{aligned}$$

If $\omega_0 > \max_x |c(x)|$, then the spectrum of K is contained into $(-\infty, 0)$ and K generates an analytic semigroup in $X = \mathcal{C}([0,1])$, endowed with the sup-norm; of course, $\mathcal{D}(K)^a \neq X$. Furthermore, if $0 < h < 1, \theta \neq 1/2$, the real interpolation space $(X, \mathcal{D}(K))_{\theta, \infty}$ is characterized by

$$(X, \mathcal{D}(K))_{\theta, \infty} = \begin{cases} \mathcal{C}^{2\theta}([0,1]), & \text{if } 0 < \theta < 1/2, \\ \{f \in \mathcal{C}^{1, 2\theta-1}([0,1]), \alpha_0 f(0) - \beta_0 f'(0) = \alpha_1 f(1) + \beta_1 f'(1) = 0\}, & \\ \text{if } 1/2 < \theta < 1 \end{cases}$$

([2, p. 205]), where the space $\mathcal{C}^\sigma([0,1]) = \mathcal{C}^\sigma([0,1]; \mathbf{C}), 0 < \sigma < 1$.

This being stated, let $\rho > 0$ fixed and consider the operator K previously introduced, with $a(\cdot) = 1, b(\cdot) = c(\cdot) = 0 = \omega_0$.

Let us introduce another operator H by

$$\mathcal{D}(H) = \mathcal{D}(K), \quad (Hu)(x) = a(x)u''(x), \quad u \in \mathcal{D}(H), \quad 0 \leq x \leq 1,$$

where $a \in \mathcal{C}((0,1); \mathbf{R}), \inf_x a(x) > 0$.

Our general statements allow to handle in the space $\mathcal{C}([0,1])$ the mixed problem

$$\left\{ \begin{array}{l} \frac{\partial^2 u}{\partial t^2} - \rho \frac{\partial^3 u}{\partial x^2 \partial t} + a(x) \frac{\partial^4 u}{\partial x^4} = f(t,x), \quad 0 < t \leq T, \quad 0 \leq x \leq 1, \\ u(0,x) = u_0(x), \quad \frac{\partial u}{\partial t}(0,x) = u_1(x), \quad 0 \leq x \leq 1, \\ \alpha_0 u(t,0) - \beta_0 \frac{\partial u}{\partial x}(t,0) = \alpha_1 u(t,1) + \beta_1 \frac{\partial u}{\partial x}(t,1) = 0, \quad 0 < t \leq T, \\ \alpha_0 \frac{\partial^2 u}{\partial x^2}(t,0) - \beta_0 \frac{\partial^3 u}{\partial x^3}(t,0) = \alpha_1 \frac{\partial^2 u}{\partial x^2}(t,1) + \beta_1 \frac{\partial^3 u}{\partial x^3}(t,1) = 0, \quad 0 < t \leq T, \end{array} \right.$$

provided that ρ is sufficiently large

Since the interpolation spaces $(X, \mathcal{D}(K))_{\theta, \infty}$ are well-characterized, it should be a simple matter to translate the conditions given by Theorems 1–7 into precise regularity properties and compatibility relations to ask for the initial and non homogeneous data.

EXAMPLE 7. To begin with, we recall that if $-A$ generates an analytic semigroup in X , with non dense domain $\mathcal{D}(A)$, then the operator $-A_0$ defined in $X_0 = \mathcal{D}(A)^a$ (=closure of $\mathcal{D}(A)$ in X) by

$$\mathcal{D}(A_0) = \{u \in \mathcal{D}(A); Au \in X_0\}, \quad A_0 u = Au, \quad u \in \mathcal{D}(A_0),$$

does generate a bounded analytic semigroup in X_0 that is strongly continuous in $t \geq 0$. Further, by [19, Proposition 1.6],

$$(X, \mathcal{D}(A))_{\theta, \infty} = (X_0, \mathcal{D}(A_0))_{\theta, \infty}, \quad 0 < \theta < 1.$$

In particular, if Ω is a bounded domain in R^q , $q \geq 1$, with a smooth boundary $\partial\Omega$,

$$X = \mathcal{C}(\bar{\Omega}), \quad \mathcal{C}_0(\bar{\Omega}) = \{u \in X; u|_{\partial\Omega} = 0\},$$

and A is given by

$$\mathcal{D}(A) = \{u \in \mathcal{C}_0(\bar{\Omega}); u \in \bar{\Omega}\}, \quad Au = -\Delta u, \quad u \in \mathcal{D}(A),$$

where Δ is understood in the sense of distributions on Ω , then

$$\mathcal{D}(A)^a = \mathcal{C}_0(\bar{\Omega}), \quad \mathcal{D}(A^0) = \{u \in \mathcal{C}_0(\bar{\Omega}); \Delta u \in \mathcal{C}_0(\bar{\Omega})\}, \quad A_0 u = -\Delta u, \quad u \in \mathcal{D}(A_0).$$

It is also well known that A_0 satisfies the spectral assumptions of [7], so that $-A_0^k$, $k \in \mathbb{N}$, do generate analytic semigroups in X_0 .

Now, the space $(X, \mathcal{D}(A))_{\theta, \infty} = (X_0, \mathcal{D}(A_0))_{\theta, \infty}$, $0 < \theta < 1$, $\theta \neq 1/2$, has been characterized by means of Hölder-continuous functions on $\bar{\Omega}$ and, on the other hand, by [23, p. 93],

$$(X_0, \mathcal{D}(A_0^m))_{\theta, \infty} = \{u \in \mathcal{D}(A_0^s); A_0^s u \in (\mathcal{C}(\bar{\Omega}), \mathcal{D}(A))_{\theta m - s, \infty}\},$$

where $s = [\theta m]$ is the integer part of θm and $m \in \mathbf{N}$.

Hence we could consider some mixed problems connected with the equation

$$\frac{\partial^n u}{\partial t^n} + c_1 (-\Delta)^{q_1} \frac{\partial^{n-1} u}{\partial t^{n-1}} + \dots + c_n (-\Delta)^{q_1 + \dots + q_n} u = f(t, x), \quad 0 \leq t \leq T, \quad x \in \Omega,$$

with $c_1, \dots, c_n > 0$, $q_1 > q_2 > \dots > q_n \geq 1$, $q_j \in \mathbf{N}$, taking $X = \mathcal{C}_0(\bar{\Omega})$, $B_1 = c_1 (-\Delta)^{q_1}$, according to the definitions above mentioned, as well as other equations obtained by the preceding one with the help of suitable perturbations that do not change the type of equation.

References

- [1] P. Acquistapace and B. Terreni: *Some existence and regularity results for abstract non-autonomous parabolic equations*. J. Math. Anal. Appl. **99** (1984), 9–64.
- [2] P. Acquistapace and B. Terreni: *Maximal space regularity for abstract linear non-autonomous parabolic equations*, J. Funct. Anal. **60** (1985), 168–210.
- [3] M.K. Balaev: *On evolution equations of arbitrary order of parabolic type*, (Russian), Dokl. Akad. Nauk Azerbaijan Tom **41** (1988), 7–10.
- [4] S. Chen and R. Triggiani: *Proof of extensions of two conjectures on structural damping for elastic systems: the case $\frac{1}{2} \leq \alpha < 1$* , Pacific J. Math. **39** (1989), 15–55.
- [5] S. Chen and R. Triggiani: *Characterization of domains of fractional powers of certain operators arising in elastic systems and applications*. J. Diff. Equations **88** (1990), 279–293.
- [6] G. Da Prato and P. Grisvard: *Sommés d'opérateurs linéaires et équations différentielles opérationnelles*. J. Math. Pures Appl. **54** (1975), 303–387.
- [7] R. de Laubenfels: *Powers of generators of holomorphic semigroups*, Proceed. AMS. **99** (1987), 105–108.
- [8] A. Favini: *Second order parabolic equations in Banach space*, in “Differential equations with applications in biology, physics, and engineering”, ed. by J.A. Goldstein, F. Kappel & W. Schappacher, pp. 129–137, ed. Marcel Dekker, 1991.
- [9] A. Favini and E. Obrecht: *Conditions for parabolicity of second order abstract differential equations*, Diff. & Int. Equations **4** (1991), 1005–1022.
- [10] A. Favini and H. Tanabe: *Linear parabolic equations of higher order in time*, in “Differential Equations in Banach Spaces”, ed. by G. Dore, A. Favini, E. Obrecht & A. Venni. pp. 85–92, ed. Marcel Dekker, 1993.
- [11] T. Kato: *Perturbation theory for linear operators*, Springer, 1966.
- [12] S.G. Krein: *Linear differential equations in Banach space*, Nauka, 1967 (in Russian): English Transl., AMS, 1971.

- [13] A. Lunardi: *Interpolation spaces between domains of elliptic operators and spaces of continuous functions with applications to nonlinear parabolic equations*, Math. Nach. **121** (1985), 295–318.
- [14] A.S. Markus: *Introduction to the spectral theory of polynomial operator pencils*, AMS, 1988.
- [15] E. Obrecht: *Evolution operators for higher order abstract parabolic equations*, Czechoslovak Math. J. **36** (III) (1986), 257–50.
- [16] E. Obrecht: *The Cauchy problem for time dependent abstract parabolic equations of higher order*, J. Math. Anal. Appl. **15** (1987), 508–530.
- [17] E. Obrecht: *Second order abstract parabolic equations*, Semesterbericht Funktional-analysis Tübingen, Sommersemester (1986), 191–206.
- [18] L. Rodman: *An introduction to operator polynomials*, Birkhäuser, 1989.
- [19] E. Sinestrari: *On the abstract Cauchy problem of parabolic type in spaces of continuous functions*, J. Math. Anal. Appl. **107** (1985), 16–66.
- [20] H.B. Stewart: *Generation of analytic semigroups by strongly elliptic operators*, Trans. Amer. Math. Soc. **199** (1974), 141–162.
- [21] H. Tanabe: *Equations of evolution*, Iwanami-Shoten, Tokyo, 1975. (in Japanese); English transl., Pitman, 1979.
- [22] H. Tanabe: *On fundamental solutions of linear parabolic equations of higher order in time and associated Volterra equations*, J. Diff. Equations **73** (1988), 288–308.
- [23] H. Triebel: *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland, 1978.
- [24] S. Ya. Yakubov: *The Cauchy problem for second-order differential equations in Banach space*, Dokl. Acad. Nauka. SSSR **168** (1966), 759–762; English transl. in Soviet Math. Dokl. **7** (1966).
- [25] S. Ya. Yakubov: *A nonlocal boundary value problem for a class of Petrowskii well posed equations*, Math. USSR Sb. **46** (1983), 255–265.

Angelo Favini
Dipartimento di Matematica
Università degli Studi di
Bologna
Piazza di Porta S. Donato, 5
40127 Bologna
Italy

Hiroki Tanabe
Department of Mathematics
Osaka University
Toyonaka, Osaka 560
Japan