

## UNIQUENESS OF THE MOST SYMMETRIC NON-SINGULAR PLANE SEXTICS

HIROSHI DOI, KUNIHIRO IDEI and HITOSHI KANETA

(Received July 18, 1998)

### 0. Introduction

Let  $C$  be a compact Riemann surface of genus  $g \geq 2$ . The order of the holomorphic automorphism group  $\text{Aut}(C)$  takes the value  $84(g-1)$ ,  $48(g-1)$ ,  $40(g-1)$ ,  $36(g-1)$ ,  $30(g-1)$  or less by Hurwitz' theorem ([5, Chap. 6] or [1, Chap. 5]). A homogeneous polynomial  $f \in \mathbf{C}[x, y, z]$  with  $n = \deg f \geq 1$  defines an algebraic curve  $C(f)$  in the projective plane  $\mathbf{P}^2$  over the complex number field  $\mathbf{C}$ . As is well known  $C(f)$  is a compact Riemann surface of genus  $(n-1)(n-2)/2$  if  $C(f)$  is non-singular. Particularly a non-singular plane quartic (resp. sextic) has genus  $g = 3$  (resp.  $g = 10$ ). Let  $\text{Aut}(f)$  be the subgroup of the projectivities  $PGL(3, \mathbf{C})$  of  $\mathbf{P}^2$  consisting of all projectivities  $(A)$  defined by  $A \in GL(3, \mathbf{C})$  such that  $f_A$  is proportional to  $f$ . Here  $f_A(x, y, z) = f((x, y, z)(^t A^{-1}))$  by definition. Clearly  $\text{Aut}(f)$  coincides with the projective automorphism group of  $C(f)$ , if  $f$  is irreducible. It is also known that a holomorphic automorphism of a non-singular curve  $C(f)$  of degree  $n \geq 4$  is induced by a projectivity  $(A) \in PGL(3, \mathbf{C})$  [9, Theorem 5.3.17(3)]. Therefore  $\text{Aut}(C(f)) = \text{Aut}(f)$  if  $C(f)$  is non-singular of degree  $n \geq 4$ . By abuse of terminology we say that a homogeneous polynomial  $f$  is non-singular or singular according as  $C(f)$  is.

As is well known, the Klein quartic  $f_4 = x^3y + y^3z + z^3x$  is the most symmetric in the sense that  $|\text{Aut}(f_4)| = 84 \times (3-1)$ . It is also known that if  $|\text{Aut}(f)| = 168$  for a non-singular plane quartic  $f$ , then  $f$  is projectively equivalent to  $f_4$ . A. Wiman has shown that for the following non-singular sextic

$$f_6 = 27z^6 - 135z^4xy - 45z^2x^2y^2 + 9z(x^5 + y^5) + 10x^3y^3,$$

$\text{Aut}(f_6)$  is isomorphic to the simple group  $A_6 \simeq PSL(2, 3^2)$ [11], as a result  $|\text{Aut}(f_6)| = 40(g-1) = 360$ . We call  $f_6$  the Wiman sextic. He has also shown that the group  $\text{Aut}(f_6)$  acts transitively on the set of 72 flexes of  $C(f_6)$ . We can show even that no three flexes are collinear [6]. Our main results are

**Theorem.** *Let  $f$  be a non-singular plane sextic defined over  $\mathbf{C}$ . Then*

- (1)  $|\text{Aut}(f)| \leq 360$ .
- (2)  $|\text{Aut}(f)| = 360$  if and only if  $f$  is projectively equivalent to the Wiman sextic  $f_6$ .

(1) will be proved in §1 according to [4], while (2) will be shown in §2. We can show that the most symmetric non-singular plane curve of degree 3, 5 or 7 is projectively equivalent to the Fermat curve [7].

We recall a well known fact: Let  $R_A: \mathbf{C}[x, y, z] \rightarrow \mathbf{C}[x, y, z]$  be a mapping defined by  $R_A f = f_A$  for  $A \in GL(3, \mathbf{C})$  and  $f \in \mathbf{C}[x, y, z]$ . Then  $R_A$  is a ring-automorphism of the polynomial ring  $\mathbf{C}[x, y, z]$ . Since  $(f_A)_B = f_{BA}$  for  $A, B \in GL(3, \mathbf{C})$ , the assignment  $A \rightarrow R_A$  is a group homomorphism of  $GL(3, \mathbf{C})$  into  $\text{Aut}(\mathbf{C}[x, y, z])$ .

We write  $a \sim b$  when two quantities  $a$  and  $b$  such as polynomials or matrices are proportional.  $E_3$  stands for the  $3 \times 3$  unite matrix, and  $e_i$  for the  $i$ -th column vector of  $E_3$  ( $1 \leq i \leq 3$ ).

### 1. The maximum order of the automorphism group of non-singular plane sextics

Let  $f$  be a non-singular plane sextic. In this section we will show that the order of the projective automorphism group  $\text{Aut}(f)$  can take the value neither  $84 \times 9$  nor  $48 \times 9$  (Theorem (1)). Otherwise, for some  $f$   $\text{Aut}(f)$  has a subgroup of order  $3^3$  by Sylow's theorem. Thus it suffices to show the following theorem.

**Theorem 1.1.** *Let  $f$  be a non-singular plane sextic. If  $27 \mid |\text{Aut}(f)|$ , then  $|\text{Aut}(f)| < 360$ .*

Our approach is elementary, but involves much computation. There exist exactly five groups of order 27 up to group isomorphism [3, 4.4]. They are three abelian groups and two non-abelian groups: (1)  $\mathbf{Z}_{27}$  (2)  $\mathbf{Z}_9 \times \mathbf{Z}_3$  (3)  $\mathbf{Z}_3 \times \mathbf{Z}_3 \times \mathbf{Z}_3$  (4)  $a^9 = 1, b^3 = 1, b^{-1}ab = a^4$  (5)  $a^3 = 1, b^3 = 1, c^3 = 1, ab = bac, ca = ac, cb = bc$ . The group (5) is isomorphic to the matrix group

$$E(3^3) = \left\{ M(\alpha, \beta, \gamma) = \begin{bmatrix} 1 & \alpha & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{bmatrix}; \alpha, \beta, \gamma \in \mathbf{F}_3 \right\}.$$

We find projective representations of these groups in the projective plane  $\mathbf{P}^2$  defined over  $\mathbf{C}$ , and find a non-singular invariant sextic  $f$ , if any. We can manage to estimate the order of the projective automorphism group  $\text{Aut}(f)$ .

**Lemma 1.2.** *Let  $\varepsilon$  be a primitive 9-th root of 1  $\in \mathbf{C}$ . If  $G_9$  is a subgroup of  $PGL(3, \mathbf{C})$ , isomorphic to  $\mathbf{Z}_9$ , then  $G_9$  is conjugate to one of the following three groups in  $PGL(3, \mathbf{C})$ :*

- (1)  $\langle\langle \text{diag}[1, \varepsilon, \varepsilon] \rangle\rangle$  (2)  $\langle\langle \text{diag}[1, \varepsilon, \varepsilon^2] \rangle\rangle$  (3)  $\langle\langle \text{diag}[1, \varepsilon, \varepsilon^3] \rangle\rangle$ .

Proof. By our assumption  $G_9$  is generated by a projective transformation  $(A)$ , where  $A \in GL(3, \mathbf{C})$  satisfies  $A^9 = E_3$  and  $\text{ord}((A)) = 9$ , namely  $G_9 = \langle (A) \rangle$ . Therefore it is conjugate to  $\langle (\text{diag}[1, \varepsilon^i, \varepsilon^j]) \rangle$  for some  $0 \leq i \leq j \leq 8$  with  $(i, j) \neq (0, 0), (0, 3), (0, 6), (3, 3), (3, 6), (6, 6)$ . If  $(i, j) = (0, j)$  with  $j \not\equiv 0 \pmod 3$  or  $i = j \not\equiv 0 \pmod 3$ , then  $G_9$  is conjugate to (1). If  $1 \leq i < j \leq 8$  with  $(i, j) \not\equiv (0, 0) \pmod 3$ , then  $G_9$  is conjugate to (2) or (3) according as  $(i, j) \in \{(1, 2), (1, 5), (1, 8), (2, 4), (2, 7), (4, 5), (4, 8), (5, 7), (7, 8)\}$  or  $(i, j) \in \{(1, 3), (1, 4), (1, 6), (1, 7), (2, 3), (2, 5), (2, 6), (2, 8), (3, 4), (3, 5), (3, 7), (3, 8), (4, 6), (4, 7), (5, 6), (5, 8), (6, 7), (6, 8)\}$ .  $\square$

**Lemma 1.3.** *Let  $\lambda_j \in \mathbf{C} (1 \leq j \leq n)$  be mutually distinct, and let  $f_{j,A} = \lambda_j f_j$  for some  $A \in GL(3, \mathbf{C})$  and  $f_j \in \mathbf{C}[x, y, z]$ . If  $f = f_1 + \dots + f_n \neq 0$  satisfies  $f_A = \lambda f$  for some  $\lambda \in \mathbf{C}$ , then  $\lambda = \lambda_i$  for some  $i$ , and  $f_j = 0$  for  $j \neq i$ .*

Proof. We have  $\lambda^k f = \lambda_1^k f_1 + \dots + \lambda_n^k f_n$  for  $0 \leq k < n$ . Multiplying the inverse of the Vandermonde matrix, we get  $f_j = c_j f (1 \leq j \leq n)$  for some  $c_j \in \mathbf{C}$ . Thus  $c_j(\lambda_j - \lambda)f = 0$ . Since  $f$  is assumed not to be the zero polynomial, the lemma follows.  $\square$

**Proposition 1.4.** *Let  $f$  be a plane sextic. If  $\text{Aut}(f)$  has a subgroup  $G_9$  isomorphic to  $\mathbf{Z}_9$ , then  $C(f)$  has a singular point.*

Proof. Let  $A_1 = \text{diag}[1, \varepsilon, \varepsilon]$ ,  $A_2 = \text{diag}[1, \varepsilon, \varepsilon^2]$  and  $A_3 = \text{diag}[1, \varepsilon, \varepsilon^3]$ . By Lemma 1.2 we may assume that  $f_{A_j^{-1}} = \lambda_j f$  for some  $\lambda_j \in \mathbf{C} (1 \leq j \leq 3)$ . Since  $A_j^9 = E_3$ , it follows that  $\lambda_j^9 = 1$ . In addition any monomial  $m$  satisfies  $m_{A_j^{-1}} = \varepsilon^i m$  for some  $i$ . Suppose that a homogeneous polynomial  $f'(x, y, z)$  of degree  $d \geq 2$ . Then  $(1, 0, 0)$  is a singular point of  $C(f')$  if and only if  $f'$  contains none of three monomials  $x^d, x^{d-1}y$  and  $x^{d-1}z$ . In the following table we summarize the values  $i$  such that  $m_{A_j^{-1}} = \varepsilon^i m$  for each  $j = 1, 2, 3$  and for special 9 monomials. The proposition is immediate from the table.

	$x^6$	$x^5y$	$x^5z$	$y^6$	$y^5x$	$y^5z$	$z^6$	$z^5x$	$z^5y$
(1)	0	1	1	6	5	6	6	5	6
(2)	0	1	2	6	5	7	3	1	2
(3)	0	1	3	6	5	8	0	6	7

$\square$

**Proposition 1.5.** *No subgroup of  $PGL(3, \mathbf{C})$  is isomorphic to  $\mathbf{Z}_3 \times \mathbf{Z}_3 \times \mathbf{Z}_3$ .*

Proof. Assume that a subgroup  $G$  of  $PGL(3, \mathbf{C})$  is isomorphic to  $\mathbf{Z}_3 \times \mathbf{Z}_3 \times \mathbf{Z}_3$ . Then there exist  $A_1, A_2, A_3 \in GL(3, \mathbf{C})$  such that  $A_1^3 = A_2^3 = A_3^3 = E_3, A_i A_j \sim A_j A_i$  for any  $1 \leq i < j \leq 3$ , and  $G = \langle (A_1), (A_2), (A_3) \rangle$ . Let  $\omega$  be a primitive 3rd root of

1. We may assume that  $G$  contains  $(W)$  of the form  $(\text{diag}[1, 1, \omega])$  or  $(\text{diag}[1, \omega, \omega^2])$ . We will show that the first case implies the second case. Since  $WA_j \sim A_jW$ , the (3,1), (3,2), (1,3) and (2,3) components of  $A_j (j = 1, 2, 3)$  vanish. So we can assume that  $A_1 = \text{diag}[\omega^m, \omega^n, \omega]$  for some  $0 \leq m, n < 3$ . If  $n = m$ , then  $n \neq 1$ , and  $A_2 = \text{diag}[\omega^{m'}, \omega^{n'}, \omega]$  with  $n' \neq m'$ . Thus  $(\text{diag}[1, \omega, \omega^2]) \in G$ . We will show that the assumption  $(A) = (\text{diag}[1, \omega, \omega^2]) \in G$  leads to a contradiction. Let  $P_1 = (1, 0, 0)$ ,  $P_2 = (0, 1, 0)$ , and  $P_3 = (0, 0, 1)$ . Then  $G$  fixes 3-point set  $K = \{P_1, P_2, P_3\}$ , because  $(A)$  and  $(A_j)$  commute. Since some  $A_j$  is not diagonal, the homomorphism  $\varphi$  from  $G$  to the permutation group of  $K$  cannot be trivial. Since  $|G| = 27$ , it cannot be surjective. Thus  $|\varphi(G)| = 3$ , and  $|\text{Ker}\varphi| = 9$ . In other words every projectively  $(\text{diag}[1, \omega^i, \omega^j])$  belongs to  $G$ . Since  $G$  is commutative, any element of  $G$  is induced by a diagonal matrix of order 3. This implies that  $|G| = 9$ , a desired contradiction.  $\square$

We turn to the group  $E(3^3)$ . See the paragraph just below Theorem 1.1 for the definition of the group and its element  $M(\alpha, \beta, \gamma)$ .

**Lemma 1.6.** (1) *Let*

$$B_1 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad B_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega \end{bmatrix}, \quad B_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{bmatrix}.$$

The map  $\phi$  defined by  $\phi(M(\alpha, \beta, \gamma)) = (B_1^\alpha B_2^\beta B_3^{\gamma-\alpha\beta})$  is an isomorphism of  $E(3^3)$  into  $PGL(3, \mathbf{C})$ .

(2) If  $G$  is a subgroup of  $PGL(3, \mathbf{C})$  and isomorphic to  $E(3^3)$ , then  $G$  is conjugate to  $\phi(E(3^3))$ .

*Proof.* (1) Let  $M_1 = M(1, 0, 0)$ ,  $M_2 = M(0, 1, 0)$ ,  $M_3 = M(0, 0, 1)$ . Then  $M(\alpha, \beta, \gamma) = M_1^\alpha M_2^\beta M_3^{\gamma-\alpha\beta}$ . First we will prove that  $\phi$  is a homomorphism by showing  $\phi(M_j M(\alpha, \beta, \gamma)) = \phi(M_j)\phi(M(\alpha, \beta, \gamma))$ . Clearly

$$\begin{aligned} M_1 M(\alpha, \beta, \gamma) &= M(\alpha + 1, \beta, \gamma + \beta) \\ M_2 M(\alpha, \beta, \gamma) &= M(\alpha, \beta + 1, \gamma) \\ M_3 M(\alpha, \beta, \gamma) &= M(\alpha, \beta, \gamma + 1). \end{aligned}$$

On the other hand,  $B_j^3 = E_3$ ,  $B_1 B_2 = B_2 B_1 B_3$ ,  $B_3 B_1 = B_1 B_3$ , and  $B_3 B_2 = B_2 B_3$ . So  $\phi$  is a homomorphism. Since  $B_2$  and  $B_3$  are diagonal, it is easy to see that  $\phi$  is injective. Note that  $\phi(E(3^3))$  does not depend on the choice of  $\omega$ , a primitive 3rd root of 1.

(2) Let  $\phi'$  be an isomorphism of  $E(3^3)$  into  $PGL(3, \mathbf{C})$ , and  $\phi'(M_j) = (B'_j)$ . We may assume  $B'_3 = B_3$  or  $B'_3 = B_2$ . The latter case is impossible. Since  $B'_3 B'_1 \sim B'_1 B'_3$  and  $B'_3 B'_2 \sim B'_2 B'_3$ , we may assume  $B'_1 = \text{diag}[\omega_1, \omega_2, 1]$ , and (1,3), (2,3), (3,1) and (3,2) components of  $B'_2$  are equal to zero. It is not difficult to see that  $B'_1 B'_2 \sim B'_2 B'_1 B'_3$  is

impossible. So let  $B'_3 = B_3$  and let  $e_i$  denote the  $i$ -th unit column vector so that  $E_3 = [e_1, e_2, e_3]$ . A matrix  $B \in GL(3, \mathbf{C})$  satisfies  $BB_3 \sim B_3B$  if and only if either  $B$  is diagonal or takes the form either  $[e_2, e_3, e_1]\text{diag}[a, b, c]$  or  $[e_3, e_1, e_2]\text{diag}[a, b, c]$ . First assume that  $B'_2 = \text{diag}[\omega_1, \omega_2, \omega_3]$ . We may assume  $1 = \omega_1 = \omega_2 \neq \omega_3$  (if necessary, we replace  $\omega$  by  $\omega^2$ ). Furthermore, we may assume  $\omega_3 = \omega$ (if necessary, we replace  $\omega$  by  $\omega^2$ ) so that  $B'_2 = B_2$ . Since  $(B'_2)$  and  $(B'_1)$  do not commute,  $B'_1$  cannot be diagonal. It turns out  $B'_1 = [e_3, e_1, e_2]\text{diag}[a, b, c]$ . By use of a diagonal matrix, we may assume that  $a = b = c$ , namely  $B'_1 = B_1$ . Secondly assume that  $B'_1 = \text{diag}[\omega_1, \omega_2, \omega_3]$ . We note that the map sending  $M(\alpha, \beta, \gamma)$  to  $M(\beta, \alpha, \gamma)$  is an anti-isomorphism. Therefore  $\phi'$  gives an isomorphism  $\phi''(M(\alpha, \beta, \gamma)) = \phi'(M(\beta, \alpha, \gamma)^{-1})$ .  $\phi''$  is an isomorphism whose type we have discussed. Namely,  $\phi'(E(3^3)) = \phi''(E(3^3))$  is conjugate to  $\phi(E(3^3))$ . Thirdly and finally assume that neither  $B'_1$  nor  $B'_2$  is diagonal. Let  $B'_2 = [e_2, e_3, e_1]$ (without loss of generality we may take  $a = b = c = 1$ ). Then we can show that if  $B'_1$  takes the form either  $[e_2, e_3, e_1]\text{diag}[a, b, c]$  or  $[e_3, e_1, e_2]\text{diag}[a, b, c]$  with  $|\{a, b, c\}| = 2$ ,  $ac = b^2\omega$  and  $a^2 = bc\omega^2$ , then  $\phi'(M(\alpha, \beta, \gamma)) = (B_1'^\alpha B_2'^\beta B_3'^{\gamma-\alpha\beta})$  is an isomorphism(if  $|\{a, b, c\}| = 1$  or  $3$ , this  $\phi'$  cannot be an isomorphism). Clearly  $\phi'(E(3^3)) = \phi(E(3^3))$ . The case  $B'_2 = [e_3, e_1, e_2]$  can be reduced to the case  $B'_2 = [e_2, e_3, e_1]$  by use of the matrix  $[e_1, e_3, e_2]$ . □

Let  $f \in \mathbf{C}[x_1, x_2, x_3]$  be a homogeneous polynomial and let  $h$  be its Hessian  $\text{Hess}(f) = \det[f_{jk}]$ , where  $f_{jk} = (\partial^2/\partial x_j \partial x_k)f$ .

**Lemma 1.7.** *Let  $A = [a_{jk}] \in GL(3, \mathbf{C})$ , and let  $f$  be a homogeneous polynomial in  $\mathbf{C}[x_1, x_2, x_3]$  such that  $f_{A^{-1}} = \lambda f$ . Then  $h_{A^{-1}} = \lambda^3(\det A^{-1})^2 h$ , where  $h = \text{Hess}(f)$ .*

*Proof.* Let  $y_j = \sum_{k=1}^3 a_{jk}x_k$ . By our assumption  $\lambda f(x_1, x_2, x_3) = f(y_1, y_2, y_3)$ . Hence

$$\begin{aligned} \lambda f_j(x_1, x_2, x_3) &= \sum_{\ell} f_{\ell}(y_1, y_2, y_3)a_{\ell j} \\ \lambda f_{jk}(x_1, x_2, x_3) &= \sum_{\ell} \sum_{\ell'} f_{\ell\ell'}(y_1, y_2, y_3)a_{\ell'k}a_{\ell j}. \end{aligned}$$

The second equality yields  $\lambda^3 h(x_1, x_2, x_3) = h_{A^{-1}}(x_1, x_2, x_3)(\det A)^2$ . □

**Lemma 1.8.** *Let the marices  $B_j$  be as in Lemma 1.6. A non-singular sextic  $f$  is invariant under all  $(B_j)$  if and only if*

$$f \sim x^6 + y^6\alpha^2 + z^6\alpha + \kappa(x^3y^3 + y^3z^3\alpha^2 + z^3x^3\alpha),$$

where  $\alpha^3 = 1$  with  $(\kappa^2 - 4\alpha^2)(\kappa^3 - 3\alpha\kappa^2 + 4) \neq 0$ .

Proof. First we will show that a non-singular sextic  $f$  invariant under all  $(B_j)$  takes the form as in the lemma. Note that  $f_{B_3^{-1}} = \omega^j f$  and  $f_{B_2^{-1}} = \omega^k f$  for some  $j, k \in \{0, 1, 2\}$ . One can easily see that unless  $(j, k) = (0, 0)$ ,  $f$  is singular. So  $f$  takes the form  $f = a_1x^6 + a_2y^6 + a_3z^6 + a_4x^3y^3 + a_5y^3z^3 + a_6z^3x^3$ . Since  $f_{B_1^{-1}} = a_3x^6 + a_1y^6 + a_2z^6 + a_6x^3y^3 + a_4y^3z^3 + a_5z^3x^3$  must be equal to  $\lambda f$ , where  $\lambda^3 = 1$  (note that  $B_1^3 = E_3$ ), we get  $(a_1, a_2, a_3) = \lambda(a_3, a_1, a_2)$ , and  $(a_4, a_5, a_6) = \lambda(a_6, a_4, a_5)$ . Therefore  $a_2 = \lambda a_1$ ,  $a_3 = \lambda^2 a_1$ ,  $a_5 = \lambda a_4$ ,  $a_6 = \lambda^2 a_4$ . We note that  $a_1 \neq 0$ , because, otherwise,  $f$  is singular.

Let  $f = x^6 + y^6\alpha^2 + z^6\alpha + \kappa(x^3y^3 + y^3z^3\alpha^2 + z^3x^3\alpha)$ , where  $\alpha^3 = 1$ . Obviously  $f$  is invariant under all  $(B_j)$ . We will discuss when  $C(f)$  has a singular point. Simple computation yields

$$\begin{aligned} f_x &= 3x^2(2x^3 + \kappa y^3 + \kappa\alpha z^3) \\ f_y &= 3y^2(\kappa x^3 + 2\alpha^2 y^3 + \alpha^2 \kappa z^3) \\ f_z &= 3z^2(\alpha \kappa x^3 + \alpha^2 \kappa y^3 + 2\alpha z^3). \end{aligned}$$

If  $(a, b, c)$  is a common zero of the three linear forms in  $x^3, y^3, z^3$  above, then the determinant of the coefficient matrix vanishes, namely  $\kappa^3 - 3\alpha\kappa^2 + 4 = 0$ . Conversely, if this determinant vanishes, then  $C(f)$  has clearly a singular point. If the determinant does not vanish and  $C(f)$  has a singular point  $(a, b, c)$ , then one of  $a, b, c$  is equal to zero and  $4\alpha^2 - \kappa^2 = 0$ . It is clear that  $C(f)$  has a singular point if  $4\alpha^2 - \kappa^2 = 0$ . Thus  $C(f)$  has a singular point if and only if  $(\kappa^3 - 3\alpha\kappa^2 + 4)(4\alpha^2 - \kappa^2) = 0$ .  $\square$

**Lemma 1.9.**  $|\text{Aut}(f)| < 360$ , where  $f$  is a non-singular sextic given in Lemma 1.8.

Proof. The Hessian  $h = \text{Hess}(f)$  takes the form  $54h_1h_2$ , where  $h_1 = xyz$  and

$$\begin{aligned} h_2 &= 20\alpha\kappa^2(x^9 + y^9 + z^9) + (-5\alpha\kappa^3 + 20\alpha^2\kappa^2 + 100\kappa)(x^6y^3 + y^6z^3 + z^6x^3) \\ &\quad + (-5\alpha^2\kappa^3 + 20\kappa^2 + 100\alpha\kappa)(x^3y^6 + y^3z^6 + z^3x^6) + (35\kappa^3 - 75\alpha\kappa^2 + 500)x^3y^3z^3. \end{aligned}$$

We consider a set of lines  $L = \{\ell; \ell \text{ is a line such that } \ell \mid h\}$ . By Lemma 1.7  $\text{Aut}(f)$  acts on  $L$  as  $(A)\ell = \{(A)P; P \in \ell\}$ . Denoting the line  $x = 0$  by  $\ell_x$ , let  $G_x = \{(A) \in \text{Aut}(f); (A)\ell_x = \ell_x\}$ . Obviously  $|\text{Aut}(f)\ell_x| \leq |L| \leq 12$ . By the way we remark that  $|L| = 12$  for  $f' = x^6 + y^6 + z^6 - 10(x^3y^3 + y^3z^3 + z^3x^3)$  (Indeed, the  $3 \times 3$  matrix  $B$  whose row vectors are  $[1, 1, 1]$ ,  $[1, \omega, \omega^2]$  and  $[1, \omega^2, \omega]$ ,  $\omega$  being a primitive the third root of 1, satisfies  $f'_{B^{-1}} = -27f'$ ). Assume  $(A) \in G_x$ . Without loss of generality  $A$  takes the form

$$A = \begin{bmatrix} 1 & 0 & 0 \\ a & b & c \\ a' & b' & c' \end{bmatrix} \in GL(3, \mathbf{C}).$$

Putting  $Y = by + cz$  and  $Z = b'y + c'z$ , we get  $f_{A^{-1}} = p_0x^6 + x^5p_1(Y, Z) + x^4p_2(Y, Z) + x^3p_3(Y, Z) + x^2p_4(Y, Z) + xp_5(Y, Z) + p_6(Y, Z)$ . Since this polynomial is proportional to  $f$ ,  $p_5(Y, Z) = 6a\alpha^2Y^5 + 3\kappa\alpha^2(a'Y^3Z^2 + aY^2Z^3) + 6a'\alpha Z^5$  must vanish, namely  $a = a' = 0$ . Now  $f_{A^{-1}} = x^6 + \kappa x^3(Y^3 + Z^3\alpha) + Y^6\alpha^2 + \kappa Y^3Z^3\alpha^2 + Z^6\alpha$ . Assuming first  $\kappa \neq 0$ , we will show that  $|G_x| = 18$  to the effect that  $|\text{Aut}(f)| \leq 18 \times 12 = 216$ . By simple computation  $Y^3 + Z^3\alpha = y^3(b^3 + b^3\alpha) + 3y^2z(b^2c + b^2c'\alpha) + 3yz^2(bc^2 + b'c'^2\alpha) + z^3(c^3 + c'^3\alpha)$ .

Since this must be equal to the polynomial  $y^3 + z^3\alpha$ , it follows that  $b^2c + b'^2c'\alpha = 0$ , and  $bc^2 + b'c'^2\alpha = 0$ . Multiplying  $c$  and  $b$  to each equality and then by subtraction, we get  $b'c'(cb' - bc') = 0$ , namely  $b'c' = 0$ , because  $A$  is non-singular. If  $b' = 0$ , then  $c = 0$ ,  $b^3 = 1$ ,  $c^3 = 1$ . It can be immediately seen that with these values  $(A)$  really belongs to  $G_x$ . If  $c' = 0$ , then  $b = 0$ ,  $b'^3 = \alpha^2$ ,  $c^3 = \alpha$ . It can be also verified that with these values  $(A)$  belongs to  $G_x$ . Thus, if  $\kappa \neq 0$ , then  $|G_x| = 2 \times 9$ . If  $\kappa = 0$ , then  $h = \text{const}x^4y^4z^4$ , in particular,  $L = \{x, y, z\}$ . One can see easily that  $G_x$  consists of  $2 \times 6^2$  points. Since  $\text{Aut}(f)$  acts transitively on  $L$ , we have  $|\text{Aut}(f)| = |L| \times |G_x| = 216$  (see [10, p. 171] or [8] for the automorphism group of the Fermat curves).  $\square$

**2. Uniqueness of sextics with  $|\text{Aut}(f)|=360$**

In the previous section we have shown that  $|\text{Aut}(f)| \leq 360$  for a non-singular plane sextic  $f$ . It is, therefore, reasonable to call a non-singular plane sextic  $f$  satisfying  $|\text{Aut}(f)| = 360$ , the most symmetric. The Wiman sextic

$$f_6 = 27z^6 - 135z^4xy - 45z^2x^2y^2 + 9z(x^5 + y^5) + 10x^3y^3$$

is known to be the most symmetric [11]. The aim of this section is to prove the

**Theorem 2.1.** *The most symmetric sextics are projectively equivalent to the Wiman sextic.*

As a byproduct another proof of  $|\text{Aut}(f_6)| = 360$  will be given (see Proposition 2.22).

There are five groups of order 8 up to isomorphism ([3, chap. 4]):

- 1)  $\mathbf{Z}_8$
- 2)  $\mathbf{Z}_2 \times \mathbf{Z}_4$
- 3)  $\mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{Z}_2$
- 4)  $Q_8$ , which is generated by  $a$  and  $b$  satisfying  $a^4 = 1$ ,  $b^2 = a^2$ , and  $ba = a^{-1}b$
- 5)  $D_8$ , which is generated by  $a$  and  $b$  satisfying  $a^4 = 1$ ,  $b^2 = 1$ , and  $ba = a^{-1}b$ .

In a series of lemmas we will show that if  $f$  is the most symmetric sextic, then the Sylow 2-subgroup of  $\text{Aut}(f)$  is isomorphic to  $D_8$ .

**Lemma 2.2.** *A subgroup  $G_8$  of  $PGL(3, \mathbf{C})$  is isomorphic to  $\mathbf{Z}_8$ , if and only if  $G_8$  is conjugate to one of the following groups:*

- (1)  $\langle\langle \text{diag}[1, 1, \varepsilon] \rangle\rangle$     (2)  $\langle\langle \text{diag}[1, \varepsilon, \varepsilon^2] \rangle\rangle$
- (3)  $\langle\langle \text{diag}[1, \varepsilon, \varepsilon^3] \rangle\rangle$     (4)  $\langle\langle \text{diag}[1, \varepsilon, \varepsilon^4] \rangle\rangle$ .

Proof. Suppose that  $G_8$  and  $Z_8$  are isomorphic. Then there exists an  $A \in GL(3, \mathbf{C})$  such that  $G_8 = \langle\langle A \rangle\rangle$ . Since  $(A)$  is of finite order,  $A$  is diagonalizable;  $T^{-1}AT \sim \text{diag}[1, \varepsilon^i, \varepsilon^j](0 \leq i < j \leq 7)$ , where  $\varepsilon$  is a primitive 8-th root of 1. Clearly  $(i, j) \notin \{(0, 2), (0, 4), (0, 6), (2, 4), (2, 6), (4, 6)\}$ . If  $i = 0$ , then  $G_8$  is conjugate to (1). If  $(i, j) \in \{(1, 2), (1, 7), (2, 5), (3, 5), (3, 6), (6, 7)\}$ , then  $G_8$  is conjugate to (2). If  $(i, j) \in \{(1, 3), (1, 6), (2, 3), (2, 7), (5, 6), (5, 7)\}$ , then  $G_8$  is conjugate to (3). Finally if  $(i, j) \in \{(1, 4), (1, 5), (3, 4), (3, 7), (4, 5), (4, 7)\}$ , then  $G_8$  is conjugate to (4). □

**Lemma 2.3.** *The projective automorphism group  $\text{Aut}(f)$  of a non-singular sextic  $f$  has a subgroup isomorphic to  $Z_8$ , if and only if  $f$  is projectively equivalent to a sextic of the form  $f' = x^6 + Bx^2y^2z^2 + y^5z + yz^5$  with  $B^3 + 27 \neq 0$ .*

Proof. Assume that  $\text{Aut}(f)$  has a subgroup isomorphic to  $Z_8$ . Let  $A$  denote one of the following four matrices;  $\text{diag}[1, 1, \varepsilon]$ ,  $\text{diag}[1, \varepsilon, \varepsilon^2]$ ,  $\text{diag}[1, \varepsilon, \varepsilon^3]$ ,  $\text{diag}[1, \varepsilon, \varepsilon^4]$ , where  $\varepsilon$  is a primitive 8-th root of 1. By Lemma 2.2  $f$  is projectively equivalent to a sextic  $f'$  such that  $f'_{A^{-1}} = \varepsilon^j f'$  for some  $0 \leq j < 8$ . One can easily see that such an  $f'$  is singular except for the case  $(A, j) = (\text{diag}[1, \varepsilon, \varepsilon^3], 0)$  (see the proof of Proposition 1.4). In this exceptional case  $f'$  is a linear combination of monomials  $x^6, x^2y^2z^2, y^5z, yz^5$ . Since  $f'$  is assumed to be non-singular, it takes the form  $x^6 + Bx^2y^2z^2 + (y^5z + yz^5)$  up to projective equivalence. Suppose that  $C(f')$  has a singular point  $(a, b, c)$ . It is immediate that  $abc \neq 0$ . It is a common zero of  $f_1 = 3x^4 + By^2z^2$ ,  $f_2 = 2Bx^2yz + 5y^4 + z^4$  and  $f_3 = 2Bx^2yz + y^4 + 5z^4$ . Being on  $C(f_2)$  and  $C(f_3)$ ,  $(a, b, c)$  satisfies  $Ba^2c + 3b^3 = 0$  and  $Ba^2b + 3c^3 = 0$ , hence  $B^2a^4 = 9b^2c^2$ . Since  $B^2f_1(a, b, c) = 0$ , we get  $(27 + B^3)b^2c^2 = 0$ , namely  $B^3 + 27 = 0$ . Conversely, if  $B^3 + 27 = 0$ , then  $(\sqrt{-3/B}, 1, 1)$  is a singular point of  $C(f')$ . □

We cite two theorems concerning a flex of a plane curve.

**Theorem 2.4** ([2, p. 70]). *A point  $P$  on an irreducible plane curve  $C(f)$  is a simple point if and only if the local ring  $\mathcal{O}_P(f)$  is a discrete valuation ring. In this case, if  $L = ax + by + cz$  is a line through  $P$  different from the tangent to  $C(f)$  at  $P$ , then the image  $\ell$  of  $L$  in  $\mathcal{O}_P(f)$  is a uniformizing parameter for  $\mathcal{O}_P(f)$ .*

**Theorem 2.5** ([2, p. 116]). *Let  $h$  be the Hessian of an irreducible  $f$ .*

- (1)  *$P$  lies both on  $C(h)$  and  $C(f)$ , if and only if  $P$  is a flex or a multiple point of  $f$ .*
- (2) *The intersection number  $I(P, h \cap f)$  is equal to 1 if and only if  $P$  is an ordinary*

*flex.* (Note that if  $P$  is a simple point of  $C(f)$  and  $C(\ell)$  is the tangent at  $P$  to  $C(f)$ , then  $I(P, h \cap f) = \text{ord}_P^f(h)$  [2, p. 81], which is equal to  $I(P, \ell \cap f) - 2 = \text{ord}_P^f(\ell) - 2$  [2, Proof on p. 116].)

The following lemma shows that a Sylow 2-subgroup of  $\text{Aut}(f)$  of the most symmetric sextic  $f$  cannot be isomorphic to  $\mathbf{Z}_8$ .

**Lemma 2.6.** *If  $f' = x^6 + Bx^2y^2z^2 + y^5z + yz^5$  with  $B^3 + 27 \neq 0$ , then  $|\text{Aut}(f')| < 360$ .*

Proof. Since  $f'(x, 1, z) = x^6 + Bx^2z^2 + z + z^5$ ,  $P = (0, 1, 0)$  is a flex of  $C(f')$ . The tangent to  $C(f')$  at  $P$  is  $C(z)$ . Since  $\text{ord}_P^{f'}$  is a discrete valuation of the local ring  $\mathcal{O}_P(f')$ , and  $x$  is a uniformizing parameter of the ring, namely  $\text{ord}_P^{f'}(x) = 1$ , we get  $\text{ord}_P^{f'}(z) = 6$ . Simple calculation yields the Hessian  $h' = \text{Hess}(f')$ , which takes the form  $-360B^2x^8y^2z^2 - 750x^4\{y^8 + z^8 + (10500 + 40B^3)y^4z^4\} - 160b^2x^2(y^7z^3 + y^3z^7) - 50B(y^{10}z^2 + y^2z^{10}) + 700By^6z^6$ . So  $I(P, h' \cap f') = \text{ord}_P^{f'}(h') = 4$ . This value can be obtained as  $\text{ord}_P^{f'}(z) - 2$  by Theorem 2.5 (2). Let  $G_P = \{(A) \in \text{Aut}(f'); (A)P = P\}$ . Since  $(A) \in G_P$  fixes  $P$  as well as the tangent  $C(z)$ , we may assume that

$$A = \begin{bmatrix} a & 0 & c \\ a' & b' & c' \\ 0 & 0 & 1 \end{bmatrix}.$$

The condition  $f'_{A^{-1}} \sim f'$  implies that  $a' = c' = 0$ , because  $5(b'y)^4(a' + c'z)z$  must vanish in  $f'_{A^{-1}}$ . Such an  $(A)$  belongs to  $G_P$  if and only if  $b'^4 = 1$ ,  $a^6 = b'$ , and  $Ba^2b' = B$ . Thus  $|G_P|$  is equal to 8 or 24 according as  $B \neq 0$  or  $B = 0$ . In the case  $B \neq 0$ , we evaluate the order of the group  $\text{Aut}(f')$  as follows:

$$4 \left( \frac{|\text{Aut}(f')|}{|G_P|} \right) = I(P, h' \cap f') \left( \frac{|\text{Aut}(f')|}{|G_P|} \right) \leq \sum_Q I(Q, h' \cap f') = 12 \times 6.$$

Thus  $|\text{Aut}(f')| \leq 144$ .

Suppose  $B = 0$ . In this case  $h' = -750x^4(y^8 - 14y^4z^4 + z^8)$ , and  $h'$  contains 9 linear factors;  $x$  with multiplicity four, and  $\sqrt{-1}^j(7 \pm 4\sqrt{3})y - z$  ( $0 \leq j \leq 3$ ) with multiplicity one. Let  $G_x = \{(A) \in \text{Aut}(f'); (A)\ell_x = \ell_x\}$ , where  $\ell_x$  stands for the line  $C(x)$ . By Lemma 1.7  $G_x = \text{Aut}(f')$ . We shall show that  $|G_x| = 144$ . Assume that  $(A) \in G_x$ .  $(A)$  fixes both  $C(f)$  and  $C(x)$ . Note that each tangent to  $C(f)$  at the intersection  $\in C(f) \cap C(x)$  passes through  $(1, 0, 0)$ . So  $(A)$  fixes  $(1, 0, 0)$  as well. Thus  $A$  takes the form

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & b & c \\ 0 & b' & c' \end{bmatrix}$$

up to constant multiplication. Putting  $Y = by + cz$ ,  $Z = b'y + c'z$ , we write  $f'_{A^{-1}}$  as  $Y^5Z + YZ^5 + x^6$ . Now (A) belongs to  $G_x$  if and only if  $y^5z + yz^5 = Y^5Z + YZ^5$ . The right-hand side takes the form  $y^6(b^5b' + bb'^5) + \dots + z^6(c^5c' + cc'^5)$ . Therefore  $bb'(b^4 + b'^4) = 0$ , and  $cc'(c^4 + c'^4) = 0$ . If  $b = 0$ , then it follows immediately that  $c' = 0$ , and  $c^5b' = cb'^5 = 1$ . The number of such an (A) is equal to 24. Similarly the case  $b' = 0$  gives another 24 elements of  $G_x$ . The case  $cc' = 0$  does not give new (A)  $\in G_x$ . We turn to the case  $bb'cc' \neq 0$ . In this case  $b^4 + b'^4 = c^4 + c'^4 = 0$ . Since the coefficient of  $y^4z^2$  vanishes,  $b^2c^2 + b'^2c'^2 = 0$ . Under these conditions the coefficients of  $y^2z^4$ ,  $y^3z^3$  vanish. The coefficients of  $y^5z$  and  $yz^5$  yield the condition  $1 = -4b^4(bc' - b'c)$  and  $1 = 4c^4(bc' - b'c)$  respectively. In particular  $c^4 = -b^4$ . Therefore if  $bb'cc' \neq 0$ , then (A)  $\in G_x$  if and only if  $b^4 + b'^4 = 0$ ,  $c^4 + c'^4 = 0$ ,  $b^4 + c^4 = 0$ ,  $b^2c^2 + b'^2c'^2 = 0$ , and  $4b^4(-bc' + b'c) = 1$ . Thus  $b' = \sqrt{-1}^j(1 + \sqrt{-1})b/\sqrt{2}$ ,  $c' = \sqrt{-1}^k(1 + \sqrt{-1})c/\sqrt{2}$  with  $0 \leq j, k \leq 3$  and  $j + k \equiv 0 \pmod{2}$ ,  $c = \sqrt{-1}^\ell(1 - \sqrt{-1})b/\sqrt{2}$  with  $0 \leq \ell \leq 3$  such that  $4b^6(\sqrt{-1}^j - \sqrt{-1}^k)\sqrt{-1}^\ell = 1$ . It is easy to see that each  $j$  gives one admissible value of  $k$ , that  $\ell$  can be arbitrary, and that  $b$  can take six values for an admissible  $(j, k, \ell)$ . Consequently there exist  $4 \times 4 \times 6$  (A)  $\in G_x$  such that  $bb'cc' \neq 0$ . Hence  $|G_x| = 24 + 24 + 96 = 144$ . This completes the proof of Lemma 2.6.  $\square$

**Lemma 2.7.** *A subgroup  $G_8$  of  $PGL(3, \mathbf{C})$  is isomorphic to  $\mathbf{Z}_2 \times \mathbf{Z}_4$ , if and only if  $G_8$  is conjugate to one of the following two groups:*

- (1)  $\langle\langle \text{diag}[-1, 1, 1], \text{diag}[1, \sqrt{-1}, \sqrt{-1}] \rangle\rangle$
- (2)  $\langle\langle \text{diag}[-1, 1, 1], \text{diag}[1, \sqrt{-1}, \sqrt{-1}^2] \rangle\rangle$ .

*Proof.* Assume that  $G_8$  is isomorphic to  $\mathbf{Z}_2 \times \mathbf{Z}_4$ . Then there exist commuting (A), and (B) in  $PGL(3, \mathbf{C})$  of order 2 and 4 respectively. We may assume that  $A^2 = E_3$  and B takes the form either  $\text{diag}[1, 1, \sqrt{-1}]$  or  $\text{diag}[1, \sqrt{-1}, \sqrt{-1}^2]$ . First suppose that  $B = \text{diag}[1, 1, \sqrt{-1}]$ . Since  $AB \sim BA$ , (1,3),(2,3),(3,1) and (3,2) components of A vanish. We may assume that (3,3) component of A is equal to 1. Since A is diagonalizable, we may assume that  $A = \text{diag}[-1, 1, 1]$ . Secondly assume that  $B = \text{diag}[1, \sqrt{-1}, \sqrt{-1}^2]$ . Since  $AB \sim BA$ , and A is involutive, it follows that A is diagonal;  $A = \text{diag}[a, b, 1]$ . If  $a = b$ , then  $a = -1$ . There exists a  $T \in GL(3, \mathbf{C})$  such that  $T^{-1}AT \sim \text{diag}[-1, 1, 1]$  and  $T^{-1}BT \sim \text{diag}[1, \sqrt{-1}^3, \sqrt{-1}^2]$ , hence  $T^{-1}B^3T \sim \text{diag}[1, \sqrt{-1}, \sqrt{-1}^2]$ . The case  $a \neq b$  can be dealt with similarly.  $\square$

**Lemma 2.8.** *If a plane sextic is invariant under the group (1) or (2) in Lemma 2.7, then it is singular.*

*Proof.* Let  $A = \text{diag}[-1, 1, 1]$ ,  $B_1 = \text{diag}[1, 1, \sqrt{-1}]$ ,  $B_2 = \text{diag}[1, \sqrt{-1}, \sqrt{-1}^2]$ , and let B denote either  $B_1$  or  $B_2$ . As in the proof of Proposition 1.4 we can show easily that a sextic  $f$  satisfying  $f_{B^{-1}} \sim f$  and  $f_{A^{-1}} \sim f$  is singular. Indeed, if  $f$  contains  $x^6$ , then  $f_{B^{-1}} = f$ , hence three monomials  $z^6, z^5x, z^5y$  or three monomials  $y^6,$

$y^5x, y^5z$  do not appear in  $f$  according as  $B = B_1$  or  $B = B_2$ . Suppose the monomial  $x^6$  does not appear in  $f$ . If  $f$  contains  $x^5y$ , then  $f_{A^{-1}} = -f$  and  $f_{B^{-1}} \sim f$  so that three monomials  $z^6, z^5x, z^5y$  do not appear in  $f$ , namely  $(0, 0, 1)$  is a singular point of  $C(f)$ . If  $f$  contains  $x^5z$ , then  $f_{A^{-1}} = -f$  and  $f_{B^{-1}} \sim f$  so that three monomials  $z^6, z^5x, z^5y$  do not appear in  $f$ . Finally if  $f$  contains none of three monomials  $x^6, x^5y$ , and  $x^5z$ , then  $(1, 0, 0)$  is a singular point of  $C(f)$ .  $\square$

**Lemma 2.9.** *No subgroup of  $PGL(3, \mathbf{C})$  is isomorphic to  $\mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{Z}_2$ .*

*Proof.* Let  $(A)$  and  $(B)$  be mutually distinct commuting involutions. We may assume that  $A = \text{diag}[-1, 1, 1]$ , and  $B = \text{diag}[1, -1, 1]$ . Assume that an involution  $(C)$  commutes with both of them. Then  $C$  is diagonal, hence  $(C) \in \langle (A), (B) \rangle$ . Namely, mutually distinct three commuting involutions in  $PGL(3, \mathbf{C})$  generate a subgroup of order 4.  $\square$

**Lemma 2.10.** *A subgroup  $G_8$  of  $PGL(3, \mathbf{C})$  is isomorphic to  $Q_8$ , if and only if  $G_8$  is conjugate to  $\langle (\text{diag}[1, \sqrt{-1}, \sqrt{-1}^3]), ([e_1, e_3, e_2]\text{diag}[1, \sqrt{-1}, \sqrt{-1}]) \rangle$ , where  $e_i$  is the  $i$ -th column vector of the unit matrix  $E_3$ .*

*Proof.*  $G_8$  is isomorphic to  $Q_8$ , if and only if it is generated by some  $(A)$  of order 4 and  $(B)$  such that  $(B)^2 = (A)^2$  and  $(B)(A) = (A)^{-1}(B)$ . Suppose that  $G_8$  is isomorphic to  $Q_8$ . Since  $(A)$  has order 4, we may assume that  $A$  takes the form either  $\text{diag}[1, 1, \sqrt{-1}]$  or  $\text{diag}[1, \sqrt{-1}, \sqrt{-1}^3]$ , for subgroups  $\langle (\text{diag}[1, \sqrt{-1}^j, \sqrt{-1}^k]) \rangle (0 < j < k < 4)$  are mutually conjugate. If  $A = \text{diag}[1, 1, \sqrt{-1}]$ , we can show easily that no  $B \in GL(3, \mathbf{C})$  satisfies  $ABA \sim B$ . If  $A = \text{diag}[1, \sqrt{-1}, \sqrt{-1}^3]$ , then, up to constant multiplication,  $B = [e_1, e_3, e_2]\text{diag}[1, b, c]$  with  $bc = -1$  alone satisfies  $B^2 \sim A^2$  and  $ABA \sim B$ . Transforming  $B$  by a diagonal matrix we get the lemma.  $\square$

**Lemma 2.11.** *Any  $Q_8$ -invariant sextic is singular.*

*Proof.* Let  $A = \text{diag}[1, \sqrt{-1}, \sqrt{-1}^3]$  and  $B = [e_1, e_3, e_2]\text{diag}[1, \sqrt{-1}, \sqrt{-1}]$ , and  $f$  is a sextic. Suppose  $f_{A^{-1}} = \sqrt{-1}^j f$  for some  $0 \leq j \leq 3$ .  $f$  is a linear combination of monomials  $m$  in  $x, y, z$  satisfying  $m_{A^{-1}} = \sqrt{-1}^j m$ . If  $j = 2$ , then  $f$  contains none of  $x^6, x^5y$ , and  $x^5z$  so that  $(1, 0, 0)$  is a singular point of  $C(f)$ . If  $j \in \{1, 3\}$ , then  $x$  divides  $f$ . Finally if  $j = 0$ , then  $f$  is a linear combination of eight monomials:  $x^6, x^2y^4, x^2y^2z^2, x^2z^4, x^4yz, y^5z, y^3z^3, yz^5$ . Since we also require that  $f_{B^{-1}} \sim f$ ,  $f$  is either a linear combination of the leading four monomials or a linear combination of the remaining four monomials. In either case  $f$  is reducible.  $\square$

We have so far shown that a Sylow 2-subgroup of  $\text{Aut}(f)$  of the most symmetric sextic  $f$  is isomorphic to  $D_8$ . We turn to the study of a Sylow 5-subgroup of  $\text{Aut}(f)$

of the most symmetric sextic  $f$ .

**Lemma 2.12.** *A subgroup  $G_5$  of  $PGL(3, \mathbf{C})$  is isomorphic to  $\mathbf{Z}_5$  if and only if  $G_5$  is conjugate to either  $G_{5,1} = \langle (\text{diag}[1, 1, \varepsilon]) \rangle$  or  $G_{5,2} = \langle (\text{diag}[1, \varepsilon, \varepsilon^2]) \rangle$ , where  $\varepsilon$  is a primitive 5-th root of 1.*

Proof. We can argue as in the proof of Lemma 2.2. □

**Proposition 2.13.** *Let  $f$  be a non-singular sextic. If  $\text{Aut}(f)$  contains a subgroup conjugate to  $G_{5,1}$  in Lemma 2.12, then  $|\text{Aut}(f)| < 360$ .*

Proof. Let a sextic  $f$  satisfy  $f_{A^{-1}} = \varepsilon^j f$ , where  $A = \text{diag}[1, 1, \varepsilon]$ . It turns out that unless  $j = 0$ ,  $f$  is singular. In the case  $j = 0$ ,  $f$  is a linear combination of monomials  $x^{6-k}y^k$  ( $0 \leq k \leq 6$ ),  $xz^5$  and  $yz^5$ . By change of variables  $x' = ax + by$  and  $y' = cx + dy$ , we may assume that

$$f = C_0x^6 + C_1x^5y + C_2x^4y^2 + C_3x^3y^3 + C_4x^2y^4 + C_5xy^5 + C_6y^6 + xz^5,$$

where  $C_6 = 1$ , because if  $C_6 = 0$ , then  $f$  is reducible. So  $P = (0, 0, 1)$  is a flex of  $C(f)$ ,  $C(x)$  is the tangent there to  $C(f)$ ,  $y$  is a uniformizing parameter of  $\mathcal{O}_P(f)$ , and  $\text{ord}_P^f(x) = 6$ . Let  $h = \text{Hess}(f)$ . By Theorem 2.5 (2)  $I(P, h \cap f) = \text{ord}_P^f(x) - 2 = 4$ . Using Bezout's theorem we get  $4|\text{Aut}(f)P| \leq \sum_Q I(Q, h \cap f) = 72$ . Let  $G_P = \{(B) \in \text{Aut}(f); (B)P = P\}$ . If  $|G_P| < 20$ , then  $|\text{Aut}(f)| = |\text{Aut}(f)P||G_P| < 360$ . We will try to show that  $|G_P| < 20$ . Let  $(B) \in G_P$ . Then the first, the second and the third row of  $B$  takes the form  $[a, 0, 0]$ ,  $[b, 1, 0]$ , and  $[a', b', c]$ . Since  $f_{B^{-1}} \sim f$ ,  $a' = b' = 0$ . Now  $f_{B^{-1}}$  is of the following form:

$$\begin{aligned} f_{B^{-1}} = & x^6(C_0a^6 + C_1a^5b + C_2a^4b^2 + C_3a^3b^3 + C_4a^2b^4 + C_5ab^5 + C_6b^6) \\ & + x^5y(C_1a^5 + 2C_2a^4b + 3C_3a^3b^2 + 4C_4a^2b^3 + 5C_5ab^4 + 6b^5) \\ & + x^4y^2(C_2a^4 + 3C_3a^3b + 6C_4a^2b^2 + 10C_5ab^3 + 15b^4) \\ & + x^3y^3(C_3a^3 + 4C_4a^2b + 10C_5ab^2 + 20b^3) \\ & + x^2y^4(C_4a^2 + 5C_5ab + 15b^2) \\ & + xy^5(C_5a + 6b) + y^6 + xz^5ac^5. \end{aligned}$$

This polynomial is proportional to  $f$ , hence, equal to  $f$ . Therefore  $ac^5 = 1$ , and  $b = C_5(1 - a)/6$ . Substituting  $b$  in the coefficients of  $x^2y^4$ , we get  $(a^2 - 1)(C_4 - 5C_5^2/12) = 0$ . If  $C_4 \neq 5C_5^2/12$ , then  $a^2 = 1$ , hence  $|G_P| \leq 10$ . Suppose  $C_4 = 5C_5^2/12$ . Comparing the coefficients of  $x^3y^3$ , we get  $(a^3 - 1)(C_3 - 5C_5^3/54) = 0$ . Suppose  $C_3 = 5C_5^3/54$  (otherwise,  $|G_P| \leq 15$ ). Now

$$f = \left(x \frac{C_5}{6} + y\right)^6 + x^6 \left(1 - \frac{C_5^6}{6^6}\right) + x^5y \left(C_1 - \frac{C_5^5}{6^4}\right) + x^4y^2 \left(C_2 - \frac{C_5^4}{2 \cdot 6^3}\right) + xz^5.$$

By change of variables  $x' = x$ ,  $y' = xC_5/6 + y$ , and  $z' = z$ , we get a projectively equivalent sextic, which will be denoted by  $f$  again:  $f = D_0x^6 + D_1x^5y + D_2x^4y^2 + y^6 + xz^5$ . If  $(B) \in G_P$ , then  $B = \text{diag}[a, 1, c]$ , where

$$D_0a^6 = D_0, \quad D_1a^5 = D_1, \quad D_2a^4 = D_2, \quad \text{and} \quad ac^5 = 1.$$

If  $D_1D_2 \neq 0$ , then  $a = 1$ , hence  $|G_P| = 5$ . If  $D_1 = 0$  and  $D_2 \neq 0$ , then  $D_0 \neq 0$ , hence  $a^2 = 1$  so that  $|G_P| = 10$ . Finally suppose that  $D_1 \neq 0$ ,  $D_2 = 0$  and that  $f$  is non-singular, namely  $6^6D_0^5 \neq 5^5D_1^6$ . Then the line  $C(z)$  intersects  $C(f)$  at distinct six points. Besides  $h = \text{Hess}(f) = 250z^3h'$ , where  $h' = -3y^4z^5 + 24(3D_0x + 2D_1y)x^4y^4 - 2D_1^2x^9$ . Note that  $h'$  has no linear factors. Indeed, none of linear factors  $z - \alpha x - \beta y$ ,  $x - \alpha y$ , and  $y - \beta x$  divides  $h'$ . Let  $G_z = \{(B) \in \text{Aut}(f); (B) \text{ fixes the line } C(z)\}$ . Since  $\text{Aut}(f) \subset \text{Aut}(h)$  by Lemma 1.7,  $(B) \in \text{Aut}(f)$  fixes a line  $C(z)$  and hence the point  $P$  (see the proof of Lemma 2.6). In particular  $G_z = \text{Aut}(f) = G_P$ , and  $B$  takes the form  $\text{diag}[a, 1, c]$ , where  $a^5 = 1$  and  $ac^5 = 1$ . In particular  $|\text{Aut}(f)| = |G_P| \leq 5 \times 5$ . □

**Lemma 2.14.** *Let  $f$  be a non-singular sextic. The automorphism group of  $f$  contains a subgroup conjugate to  $G_{5,2}$ , if and only if  $f$  is projectively equivalent to one of the following forms:*

- (1)  $x^6 + C_1x^3yz^2 + C_2y^2z^4 + C_3x^2y^3z + x(y^5 + z^5)$
- (2)  $z^6 + Bz^4xy + Cz^2x^2y^2 + Dz(x^5 + y^5) + Ex^3y^3$

If  $f$  is the sextic (1), then  $|\text{Aut}(f)| < 360$ .

*Proof.* Let  $A = \text{diag}[1, \varepsilon, \varepsilon^2]$ . Then each of the two sextics (1) and (2), say  $f$ , satisfies  $f_{A^{-1}} \sim f$ . Assume that  $(A) \in \text{Aut}(f)$  for a sextic  $f$ , namely  $f_{A^{-1}} = \varepsilon^j f$  ( $j = 0, 1, 2, 3, 4$ ). If  $j = 3$  or  $j = 4$ ,  $f$  is singular. According as  $j \in \{0, 2\}$  or  $j = 1$ ,  $f$  takes the form (1) or (2) up to projective equivalence. Assuming that  $f$  takes the form (1), we shall show that  $|\text{Aut}(f)| < 360$ .  $P = (0, 1, 0)$  is a flex of  $f$ , and  $C(x)$  is the tangent there. So  $z$  is a uniformizing parameter of  $\mathcal{O}_P(f)$ . Since  $\text{ord}_P^f(x) \geq 4$ , we can estimate the intersection number:  $I(P, h \cap f) = \text{ord}_P^f(x) - 2 \geq 2$ , where  $h$  is the Hessian of  $f$ . Let  $G_P = \{(B) \in \text{Aut}(f); (B)P = P\}$ . If  $(B) \in G_P$ , then the first, the second and the third row of  $B$  takes the form  $[1, 0, 0]$ ,  $[a, b, c]$ , and  $[a', 0, c']$  respectively, because  $(B)$  fixes the line  $C(x)$  (i.e.  $[1, 0, 0]B \sim [1, 0, 0]$ ) and  $(B)P = P$ . Since  $f_{B^{-1}} \sim f$  and  $C_2 \neq 0$ , we get  $c = 0$ ,  $a = 0$ ,  $a' = 0$ ,  $b^5 = 1$  and  $c' = b^2$ . Thus  $|G_P| = 5$ . By Bezout's theorem  $2|\text{Aut}(f)|/|G_P| = 2|\text{Aut}(f)P| \leq \sum_Q I(Q, h \cap f) \leq 72$ , that is,  $|\text{Aut}(f)| \leq 180$ . □

By Lemma 2.14 the most symmetric sextic is projectively equivalent to the following sextic :

$$f = z^6 + Bz^4xy + Cz^2x^2y^2 + Dz(x^5 + y^5) + Ex^3y^3.$$

Let  $I = [e_2, e_1, e_3]$ , where  $E_3 = [e_1, e_2, e_3]$  is the unit matrix. Clearly  $f_I = f$ . If  $f$  is the most symmetric sextic, then any Sylow 2-subgroup of  $\text{Aut}(f)$  is isomorphic to the group  $D_8$ . By Sylow's theorem the involution  $(I)$  belongs to a Sylow 2-subgroup of  $\text{Aut}(f)$ .

**Lemma 2.15.** (1) *If  $g$  is an involution of  $D_8$ , then there exists an involution  $g' \in D_8 \setminus \{g\}$  such  $gg' = g'g$ .*

(2) *Let  $g$  and  $g'$  be mutually distinct commuting involutions of  $D_8$ . Then one of the following cases takes place.*

- 1) *There exists an element  $c \in D_8$  of order 4 such that  $c^2 = g, g'cg' = c^{-1}$ .*
- 2) *There exists an element  $c \in D_8$  of order 4 such that  $c^2 = g', gcg = c^{-1}$ .*
- 3) *There exists an element  $c \in D_8$  of order 4 such that  $c^2 = gg', gcg = c^{-1}$ .*

*Proof.* Let  $a, b$  be generators of  $D_8$  such that  $a^4 = 1, b^2 = 1$  and  $ba = a^{-1}b$ . So  $a$  generates a cyclic group  $H$  of order 4, and  $D_8 = H + bH$ . An element  $g \in D_8$  is an involution if and only if  $g \in \{a^2\} \cup bH$ . (1) If  $g = a^2$ , then we can take  $g' = ba^2$ . If  $g = ba^j$ , we can take  $g' = ba^{j+2}$ . (2) If  $g = a^2$ , then  $g' \in bH$ . So we can take  $c = a$ . If  $g' = a^2$ , then we can take  $c = a$ . Finally if  $g, g' \in bH$ , then  $gg' = a^2$ . So we can take  $c = a$ . □

**Lemma 2.16.** *Assume that  $f = z^6 + Bz^4xy + Cz^2x^2y^2 + Dz(x^5 + y^5) + Ex^3y^3$  is non-singular. If there exists an involution  $(A) \in \text{Aut}(f) \setminus \{(I)\}$  such that  $(A)(I) = (I)(A)$ , then  $A$  takes the form*

$$\begin{bmatrix} \alpha & \beta & \gamma \\ \beta & \alpha & \gamma \\ \lambda & \lambda & 1 \end{bmatrix}, \text{ where } \alpha + \beta + 1 = 0, \alpha\beta + 1 = 0, \gamma\lambda = 2,$$

and

$$(\star) \quad \gamma^2 B = 12 - \gamma^5 D, \quad \gamma^4 C = 48 + \gamma^5 D, \quad \gamma^6 E = 64 - 2\gamma^5 D.$$

*Conversely, if  $(\star)$  holds for some  $\gamma \neq 0$ , then the above matrix  $A$  gives an involution  $(A) \in \text{Aut}(f) \setminus \{(I)\}$  such that  $(A)(I) = (I)(A)$ .*

*Proof.* Suppose that  $\text{Aut}(f)$  contains an involution  $(A) \neq (I)$  commuting with  $(I)$ . Let  $A = [a, b, c]$ , where  $a = [a_j], b = [b_j]$  and  $c = [c_j]$  are column vectors. We claim that  $c_3 \neq 0$ . Otherwise the condition  $AI \sim IA$  yields  $b_1 = \delta a_2, b_2 = \delta a_1, b_3 = \delta a_3$ , and  $c_2 = \delta c_1$ . Since  $A^2 \sim E_3$ , we get  $\delta = 1, a_1 + a_2 = 0$ , and  $c_1 a_3 = 2a_1^2$ . However,  $(A) \notin \text{Aut}(f)$ , because  $f_{A^{-1}} = \sum z^j C_j$  with  $C_1 = 10a_1^7 a_3 D(x+y)(x-y)^4 \not\sim D(x^5 + y^5)$ . Note that  $D \neq 0$  because of non-singularity of  $f$ . Thus we may assume that  $c_3 = 1$ . The condition  $AI \sim IA$  implies that  $a_2 = b_1, a_1 = b_2, a_3 = b_3$  and  $c_2 = c_1$ . We claim that  $c_1 \neq 0$ . If  $c_1 = 0$ , then the condition  $(A) \in \text{Aut}(f)$  yields

$a_3 = 0$  and  $a_1b_1 = 0$ . Besides, by the condition  $A^2 \sim E_3$ , we get  $A \sim E_3$  or  $A \sim I$ . Similarly  $a_3 \neq 0$ . For the sake of simplicity of notation we put  $\alpha = a_1$ ,  $\beta = b_1$ ,  $\gamma = c_1$ , and  $\lambda = a_3$ . Since  $A^2 \sim E_3$ ,  $\alpha + \beta + 1 = 0$ ,  $2\alpha\beta + \gamma\lambda = 0$ , and  $\gamma\lambda \notin \{0, -1/2\}$ . Under these conditions  $A^2 = (2\gamma\lambda + 1)E_3$ . Let  $W = \text{diag}[1, 1, 1/\gamma]$ ,  $A' = W^{-1}AW$ , and  $f_{W^{-1}} = \gamma^{-6}f'$ .  $(A') \in \text{Aut}(f')$ , because  $f'_{A'} = (f_{W^{-1}})_{A'} = f_{A'W^{-1}} = f_{W^{-1}A} = (f_A)_{W^{-1}} = (\text{const } f)_{W^{-1}} = \text{const } f_{W^{-1}} = \text{const } f'$ . By the next lemma  $(A') \in \text{Aut}(f')$  implies  $(\star)$ . Conversely suppose  $(\star)$  holds. Let  $f_{W^{-1}} = \gamma^6 f'$ . By the next lemma there exists an involution  $(A') \in \text{Aut}(f') \setminus \{(I)\}$  such that  $(A')(I) = (I)(A')$ . Since  $f'_W \sim f$ ,  $A = WA'W^{-1}$  gives an involution  $(A) \in \text{Aut}(f) \setminus \{(I)\}$ .  $\square$

**Lemma 2.17.** *Let  $f$  be as in Lemma 2.16, and let*

$$A = \begin{bmatrix} a & b & 1 \\ b & a & 1 \\ d & d & 1 \end{bmatrix}, \text{ where } a + b + 1 = 0, \quad 2ab + d = 0, \quad d \notin \left\{0, -\frac{1}{2}\right\}.$$

Then  $f_{A^{-1}} \sim f$  if and only if

$$d = 2, \quad B = 12 - D, \quad C = 48 + D, \quad E = 64 - 2D.$$

*Proof.* We note that coefficients of  $f_{A^{-1}}$  can be written without using  $a$  and  $b$ . In fact we get the following formula.

$$\begin{aligned} f_{A^{-1}} = & z^5(x + y)\{6d + B(4d - 1) + C(2d - 2) + D(2d - 5) + E(-3)\} \\ & + z^4(x^2 + y^2)\{15d^2 + B(-9/2 + 6d)d + C(1 - 5d + d^2) + D(10 + 5d) \\ & + E(3 - (3/2)d)\} \\ & + z^3(x^3 + y^3)\{20d^3 + B(-8d^2 + 4d^3) + C(3d - 4d^2) + D(-10 - 5d + 10d^2) \\ & + E(-1 + 3d)\} \\ & + z^3(x^2y + xy^2)\{60d^3 + B(4d - 16d^2 + 12d^3) + C(-2 + 9d - 4d^2) \\ & + D(25d - 10d^2) + E(-9 - 3d)\} \\ & + z^2(x^4 + y^4)\{15d^4 + B(-7d^3 + d^4) + C((3 + (1/4))d^2 - d^3) \\ & + D(5 - (25/2)d^2) + E((-3/2)d + (3/4)d^2)\} \\ & + z^2(x^3y + xy^3)\{60d^4 + B(6d^2 - 16d^3 + 4d^4) + C(-5d + 5d^2) \\ & + D(-20d - 10d^2) + E(3 - 3d - 3d^2)\} \\ & + z(x^4y + xy^4)\{30d^5 + B(4d^3 - 7d^4) + C(-4d^2 - (1/2)d^3) \\ & + D((15/2)d + (15/4)d^2 - (15/2)d^3) + E(3d + (9/4)d^2)\} \\ & + z(x^3y^2 + x^2y^3)\{60d^5 + B(12d^3 - 6d^4) + C(2d - 4d^2 - d^3) \\ & + D(-25/2)d^2 + 5d^3) + E(-3 - 3d - (3/2)d^2)\} \\ & + (x^6 + y^6)\{d^6 + B(-(1/2)d^5) + C((1/4)d^4)\} \end{aligned}$$

$$\begin{aligned}
& + D(-1 - (5/2)d - (5/4)d^2)d + E(-(1/8)d^3)\} \\
& +(x^5y + xy^5)\{6d^6 + B(d^4 - d^5) + C(-d^3 - (1/2)d^4) \\
& + D(-d + (5/2)d^3) + E((3/4)d^2 + (3/4)d^3)\} \\
& +(x^4y^2 + x^2y^4)\{15d^6 + B(4d^4 + (1/2)d^5) + C(d^2 - (1/4)d^4) \\
& + D((5/2)d^2 + (5/4)d^3) + E(-(3/2)d - 3d^2 - (15/8)d^3)\} \\
& +z^6\{1 + B + C + 2D + E\} \\
& +z^4xy\{30d^2 + B(1 - 7d + 12d^2) + C(4 - 6d + 2d^2) + D(-30d) + E(9 + 3d)\} \\
& +z^2x^2y^2\{90d^4 + B(12d^2 - 18d^3 + 6d^4) + C(1 - 6d + (15/2)d^2 + 2d^3) \\
& + D(45d^2) + E(9 + 9d + (9/2)d^2)\} \\
& +z(x^5 + y^5)\{6d^5 + B(-3d^4) + C(3/2)d^3 \\
& + D(-1 + (5/2)d + (35/4)d^2 + (5/2)d^3) + E(-3/4)d^2\} \\
& +x^3y^3\{20d^6 + B(6d^4 + 2d^5) + C(2d^2 + 2d^3 + d^4) \\
& + D(-5d^3) + E(1 + 3d + (9/2)d^2 + (5/2)d^3)\}.
\end{aligned}$$

Since  $z^5x$  does not appear in  $f$ , we have  $3E = 6d + B(4d - 1) + C(2d - 2) + D(2d - 5)$ . Since the coefficients of  $z^4x^2$ ,  $z^3x^3$ ,  $z^3x^2y$  vanish, and  $d \neq -1/2$ , we get a system of linear equations on  $B$ ,  $C$ , and  $D$  as follows:

$$\begin{aligned}
B(-2d + 1) + C(1) + D\left(\frac{1}{2}d - 5\right) &= 6d, \\
B\left(4d^2 - 6d + \frac{2}{3}\right) + C\left(-2d + \frac{4}{3}\right) + D\left(12d - \frac{50}{3}\right) &= -20d^2 + 4d, \\
B(12d^2 - 26d + 6) + C(-6d + 8) + D(-12d + 30) &= -60d^2 + 36d.
\end{aligned}$$

The determinant of the coefficient matrix is equal to  $50(4d + 2)(-d + 2)/3$ . We claim that  $d = 2$ . Assume the contrary. Cramer's formula yields  $B = 6d$ ,  $C = 12d^2$ , and  $D = 0$ . On the other hand  $D \neq 0$ , because  $f$  is assumed to be non-singular. Thus  $d = 2$ . The above system of linear equations on  $B$ ,  $C$ , and  $D$ , together with the equality  $3E = 6d + B(4d - 1) + C(2d - 2) + D(2d - 5)$  yields equalities  $B = 12 - D$ ,  $C = 48 + D$ , and  $E = 64 - 2D$ . By easy computation we get  $f_{A^{-1}} = 125f$ .  $\square$

Suppose  $f$  is the most symmetric sextic. By Lemma 2.14 we may assume that  $f$  takes the form given in Lemma 2.16. By Lemma 2.16, we may further assume that  $B = 12 - D$ ,  $C = 48 + D$ ,  $E = 64 - 2D$ .

**Lemma 2.18.** *Let  $f$  be a sextic of the form  $z^6 + Bz^4xy + Cz^2x^2y^2 + Dz(x^5 + y^5) + Ex^3y^3$  with  $B = 12 - D$ ,  $C = 48 + D$ ,  $E = 64 - 2D$ . Let  $M = \text{diag}[1, 1, m](m \neq 0)$ . Then  $f_{M^{-1}}$  is the Wiman sextic*

$$f_6 = 27z^6 - 135z^4xy - 45z^2x^2y^2 + 9z(x^5 + y^5) + 10x^3y^3,$$

if and only if  $[D, 1/m] = [(9 \pm 15\sqrt{15}\sqrt{-1})/2, (-3 \pm \sqrt{15}\sqrt{-1})/12]$ . In particular if  $D^2 - 9D + 864 = 0$ , then  $f$  is projectively equivalent to the Wiman sextic.

Proof. It is evident that  $f$  satisfies the condition if and only if the following 4 equalities hold:

- (1)  $(12 - D)/m^2 = -135/27$
- (2)  $(48 + D)/m^4 = -45/27$
- (3)  $D/m^5 = 9/27$
- (4)  $(64 - 2D)/m^6 = 10/27$ .

The equalities (2) and (3) imply  $(48 + D)m/D = -5$ , while (3) and (4) yield  $(64 - 2D)/Dm = 10/9$ . Thus  $(48 + D)(64 - 2D) + 50D^2/9 = 0$ , namely  $D^2 - 9D + 864 = 0$ .  $m^{-1} = -(48 + D)/(5D)$  gives the value of  $m^{-1}$ . Conversely, since  $m^{-2} = -(1 \pm \sqrt{15}\sqrt{-1})/24$ ,  $m^{-4} = (-7 \pm \sqrt{15}\sqrt{-1})/288$ ,  $12 - D = 15(1 \mp \sqrt{15}\sqrt{-1})/2$ , and  $48 + D = 15(7 \pm \sqrt{15}\sqrt{-1})/2$ , (1) and (2) hold, hence (3) and (4) as well. □

**Lemma 2.19.** *Let  $f$  be as in Lemma 2.18, and let*

$$A = \begin{bmatrix} a & b & 1 \\ b & a & 1 \\ 2 & 2 & 1 \end{bmatrix}, \quad \text{where } a + b + 1 = 0, \text{ and } ab + 1 = 0,$$

$$B = \text{diag}[\delta, \delta^4, 1], \quad \text{where } \delta \text{ is a primitive 5-th root of } 1.$$

Then  $(AB^2) \in \text{Aut}(f)$  and  $\text{ord}((AB^2)) = 3$ .

Proof. Let  $G$  be the subgroup of  $\text{Aut}(f)$  generated by  $(A)$ ,  $(I)$  and  $(B)$ . Let  $P_1 = (1, 0, 0)$ . It is a flex of  $C(f)$ . We can show that the orbit  $GP_1$  consists of  $2+5+5$  points, hence  $|G| = 12 \times 5$ . So it is no wonder that there is an  $(M) \in G$  of order 3. By Lemma 2.17  $(A) \in \text{Aut}(f)$ . Clearly  $(B) \in \text{Aut}(f)$ . We will show that  $c, c\omega, c\omega^2$  are the characteristic roots of  $AB^2$  for some constant  $c$ . Let  $\sqrt{5}$  be a solution to  $x^2 = 5$  (we do not assume  $\sqrt{5} > 0$ ). To get a solution to  $x^4 + x^3 + x^2 + x + 1 = 0$ , put  $y = x + x^{-1}$ . Then  $y^2 + y - 1 = 0$ . So  $y = (-1 \pm \sqrt{5})/2$ , and  $x^2 - yx + 1 = 0$ . Let  $a = (-1 + \sqrt{5})/2$ , and  $b = (-1 - \sqrt{5})/2$ . Let  $\delta$  be a solution of  $x^2 - ax + 1 = 0$ . Then  $\delta^2 = a\delta - 1$ ,  $\delta^3 = -a\delta - a$ ,  $\delta^4 = a - \delta$ , and  $\delta^5 = 1$ .  $AB^2$  now takes the form

$$AB^2 = \begin{bmatrix} a^2\delta - a & \delta + 1 & 1 \\ -\delta - b & -a^2(\delta + 1) & 1 \\ 2(a\delta - 1) & -2a(\delta + 1) & 1 \end{bmatrix}.$$

By careful computation we get  $\det(AB^2 + \sqrt{5}\mu) = 5\sqrt{5}(\mu^3 - 1)$ . As is well known, if  $AB^2 v_j = -\sqrt{5}\omega^j v_j$  and  $v_j \neq 0$ , then  $V = [v_0, v_1, v_2]$  diagonalizes  $AB^2$ ;  $V^{-1}AB^2V =$

$-\sqrt{5}\text{diag}[1, \omega, \omega^2]$ . For example we may take

$$v_0 = \begin{bmatrix} (3 + \sqrt{5})\delta - 1 - \sqrt{5} \\ -(3 + \sqrt{5})\delta \\ 2 \end{bmatrix}, \quad v_1 = \begin{bmatrix} (3 - \sqrt{5})\omega\delta + 2\omega + \sqrt{5} - 1 \\ (-3 + \sqrt{5})\omega\delta + 2(\sqrt{5} - 1)\omega + \sqrt{5} - 1 \\ 4 \end{bmatrix}.$$

Substituting  $\omega^2$  for  $\omega$  in  $v_1$ , we get  $v_2$ . □

**Lemma 2.20.** *Let  $f$  be the sextic in Lemma 2.18, and let  $V = [v_0, v_1, v_2] \in GL(3, \mathbf{C})$  be as in the proof of Lemma 2.19. Set  $U = 2V$ . Then*

$$\begin{aligned} f_{U^{-1}} = & 10240[x^6(-170 - 76\sqrt{5})(-27 + D) + (y^6 + z^6)(100 - 40\sqrt{5})D \\ & + x^3(y^3 + z^3)(-200 - 100\sqrt{5})D + y^3z^3(20 - 8\sqrt{5})(864 - 17D) \\ & + x(y^4z + yz^4)(-75 + 75\sqrt{5})D + x^4yz(75 + 33\sqrt{5})(108 + D) \\ & + x^2y^2z^2(5 + \sqrt{5})(1296 - 63D)]. \end{aligned}$$

Proof. Let  $\lambda = 2\delta$ . Then  $\lambda^2 - (-1 + \sqrt{5})\lambda + 4 = 0$ . So the coefficients of  $f_{U^{-1}}$  are  $\mathbf{Z}$ -linear combinations of  $\sqrt{5}^j \omega^k \lambda^\ell$ . Using computer, we get the reslut. □

REMARK. Let  $f' = f_{U^{-1}}$ . The involution  $(B^{-1}IB) \in \text{Aut}(f)$  gives rise to an involution  $(J) = (U^{-1}B^{-1}IBU) \in \text{Aut}(f')$ , where  $E_3 = [e_1, e_2, e_3]$ ,  $I = [e_2, e_1, e_3]$  and  $J = [e_1, e_3, e_2]$ .

The next lemma completes the proof of Theorem 2.1.

**Lemma 2.21.** *Let  $f$  be the most symmetric sextic of the form in Lemma 2.18. Then  $D^2 - 9D + 864 = 0$ .*

Proof. A Sylow 3-subgroup of  $\text{Aut}(f)$  cannot be isomorphic to  $\mathbf{Z}_9$  by Proposition 1.4. Therefore any Sylow 3-subgroup of  $\text{Aut}(f)$  is isomorphic to  $\mathbf{Z}_3 \times \mathbf{Z}_3$  [3]. By Sylow’s theorem there exists a Sylow 3-subgroup which contains  $(X) = (AB^2)$  in Lemma 2.19. So there exists a  $(Y) \in \text{Aut}(f) \setminus \{(X)\}$  of order 3 such  $(X)(Y) = (Y)(X)$ . Let  $f_{U^{-1}} = 10240f'$ ,  $(X') = (U^{-1}XU)$  (see Lemma 2.20 for the definition of  $U$ ). We may assume that  $X' = \text{diag}[1, \omega, \omega^2]$ . Then there exists a  $(Y') \in \text{Aut}(f') \setminus \{(X')\}$  such that  $X'Y' \sim Y'X'$ , and  $Y'^3 \sim E_3$ . So without loss of generality  $T = Y'$  takes the form either  $\text{diag}[1, 1, \omega]$  or  $[e_2, e_3, e_1]\text{diag}[a, b, 1]$ . The former case is impossible, because  $f'_{T^{-1}} \sim f'$  implies  $f'_{T^{-1}} = f'$  despite the fact that  $f'_{T^{-1}} \neq f'$  (note that  $D \neq 0$ , for  $f$  must be non-singular). Assume the second case for  $T$ . According as the monomial  $x^2y^2z^2$  appears in  $f'$  or not, we proceed as follows.  $[x^jy^zk^\ell]$  denotes the coefficient of  $x^jy^kz^\ell$  in  $f'$ . If  $[x^2y^2z^2] = 0$ , i.e.  $D = 144/7$ , then  $f'$  does not have an automorphism of the form  $(T)$ . Indeed, the assumption  $f'_{T^{-1}} = \text{const}f'$  leads to a contradiction

as follows. Since  $([x^6]x^6)_{T^{-1}} = \text{const}[z^6]z^6$ ,  $\text{const} = [x^6]/[z^6] = (161 + 72\sqrt{5})/32$ . By the two equalities  $a^4b[xy^4z] = \text{const}[x^4yz]$ , and  $ab[x^4yz] = \text{const}[xyz^4]$ , we get  $a^3 = [x^4yz][x^4yz]/([xyz^4][xy^4z]) = 5(161 + 72\sqrt{5})/4^2$ . On the other hand  $a^6[y^6] = \text{const}[x^6]$  gives  $a^6 = \text{const}[x^6]/[y^6] = (161 + 72\sqrt{5})^2/32^2$ . Hence  $a^6 \neq (a^3)^2$ .

Suppose that  $[x^2y^2z^2] \neq 0$ . Then  $f'_{T^{-1}} = a^2b^2f'$ . Equivalently following nine equalities hold:

$$\begin{aligned} a^2b^2[x^6] &= [y^6]a^6, & a^2b^2[x^3y^3] &= [y^3z^3]a^3b^3, & a^2b^2[x^4yz] &= [y^4zx]a^4b \\ a^2b^2[y^6] &= [z^6]b^6, & a^2b^2[y^3z^3] &= [z^3x^3]b^3, & a^2b^2[xy^4z] &= [yz^4x]ab^4 \\ a^2b^2[z^6] &= [x^6], & a^2b^2[z^3x^3] &= [x^3y^3]a^3, & a^2b^2[xyz^4] &= [yzx^4]ab. \end{aligned}$$

The second and the ninth equalities imply

$$0 = [x^3y^3][xyz^4] - [y^3z^3][x^4yz] = -6480(3 + \sqrt{5})(D^2 - 9D + 864).$$

For the sake of completeness we will determine the values of  $a$  and  $b$  in the case  $D^2 - 9D + 864 = 0$ . By the second equality above we get  $ab = [x^3y^3]/[y^3z^3]$ . The eighth equality above yields  $a = b^2$ . So  $b^3 = [x^3y^3]/[y^3z^3] = \{-100(2 + \sqrt{5})D\}/\{(20 - 8\sqrt{5})(864 - 17D)\}$ . Conversely if  $a = b^2$  and  $b^3 = \{-100(2 + \sqrt{5})D\}/\{(20 - 8\sqrt{5})(864 - 17D)\}$  with  $D^2 - 9D + 864 = 0$ , then above nine equalities hold. Clearly the second and the ninth equalities hold. Because  $a^3 = b^6 = ([x^3y^3]/[y^3z^3])^2 = [x^6]/[y^6] = [x^6]/[z^6]$ , the first and the seventh equalities hold. The third and the fifth ones hold too, because  $ab = b^3 = [x^3y^3]/[y^3z^3] = [x^4yz]/[y^4zx] = [z^3x^3]/[y^3z^3]$ . Since  $[y^6] = [z^6]$ ,  $[xy^4z] = [yz^4x]$ , and  $[x^3y^3] = [z^3x^3]$ , the fourth, the sixth and the eighth ones hold.  $\square$

For the sake of completeness we will show the following proposition, which, together with Lemma 2.18, assures us that  $|\text{Aut}(f_6)| = 360$ .

**Proposition 2.22.** *Let  $f$  be a sextic of the form  $z^6 + Bz^4xy + Cz^2x^2y^2 + Dz(x^5 + y^5) + Ex^3y^3$  with  $B = 12 - D$ ,  $C = 48 + D$ ,  $E = 64 - 2D$ , where  $D^2 - 9D + 864 = 0$ . Then  $|\text{Aut}(f)| = 360$ .*

*Proof.* By Lemma 2.14  $|\text{Aut}(f)|$  is a multiple of 5. By the proof of Lemma 2.21  $|\text{Aut}(f)|$  is a multiple of 9. In view of Theorem (1) in the introduction it suffices to show that  $\text{Aut}(f)$  contains a subgroup isomorphic to  $D_8$ . Let  $I = [e_2, e_1, e_3]$  and  $A$  be as in Lemma 2.19. Clearly  $(I) \in \text{Aut}(f)$ , and  $(A) \in \text{Aut}(f)$  by Lemma 2.17. We will show that there exists an  $(M) \in \text{Aut}(f)$  such that  $(M)^2 = (I)$ , and  $(AM)^2 = (E_3)$ (see Lemma 2.15 (2)). It is natural to diagonalize  $A$  and  $I$ . Taking  $a = (-1 + \sqrt{5})/2$ , and  $b = (-1 - \sqrt{5})/2$ , we define

$$U = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad V = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & \sqrt{5} + 1 & \sqrt{5} - 1 \end{bmatrix},$$

and  $W = UV$ . Then  $A'' = W^{-1}AW = \sqrt{5}\text{diag}[1, 1, -1]$ , and  $I'' = W^{-1}IW = \text{diag}[-1, 1, 1]$ . Put  $f' = f_{U^{-1}}$ , and  $f'' = f'_{V^{-1}}$ . We look for an  $M'' \in GL(3, \mathbf{C})$  such that  $M''^2 \sim I''$ ,  $A''M''^2 \sim E_3$  and  $(M'') \in \text{Aut}(f'')$ (see Lemma 2.15(2)). Since  $M''$  and  $I''$  commute due to the first condition, we may assume that the first, the second and the third rows of  $M''$  take the form  $[\sqrt{-1}, 0, 0]$ ,  $[0, a, b]$ , and  $[0, c, d]$  respectively. Either  $a + d = 0$  or  $a + d \neq 0$ ,  $c = d = 0$  due to the condition  $M''^2 \sim I''$ . The second case is impossible, because  $M''$  cannot be diagonal. Now the condition  $A''M''^2 \sim E_3$  yields  $a = d = 0$  and  $bc = 1$ . By careful computaion we get the explicit form of  $f''$ :

$$\begin{aligned}
 f'' &= x^6(-E) \\
 &+ x^4[y^2\{3E + 10(1 + \sqrt{5})D + (6 + 2\sqrt{5})C\} + yz\{-6E - 20D + 8D\} \\
 &\quad + z^2\{3E + 10(1 - \sqrt{5})D + (6 - 2\sqrt{5})C\}] \\
 &+ x^2[y^4\{-3E + 20(1 + \sqrt{5})D - 2(6 + 2\sqrt{5})C - (56 + 24\sqrt{5})B\} \\
 &\quad + y^3z\{12E + 20(-4 - 2\sqrt{5})D + 8(1 + \sqrt{5})C - 16(6 + 2\sqrt{5})B\} \\
 &\quad + y^2z^2\{-18E + 120D + 0C - 96B\} \\
 &\quad + y^3z\{12E + 20(-4 + 2\sqrt{5})D + 8(1 - \sqrt{5})C - 16(6 - 2\sqrt{5})B\} \\
 &\quad + z^4\{-3E + 20(1 - \sqrt{5})D - 2(6 - 2\sqrt{5})C - (56 - 24\sqrt{5})B\}] \\
 &+ x^0[y^6\{E + 2(1 + \sqrt{5})D + (6 + 2\sqrt{5})C + (56 + 24\sqrt{5})B + 16(36 + 16\sqrt{5})\} \\
 &\quad + y^5z\{-6E - 2(6 + 4\sqrt{5})D - 8(2 + \sqrt{5})C - 16(1 + \sqrt{5})B + 192(7 + 3\sqrt{5})\} \\
 &\quad + y^4z^2\{15E + 10(3 + \sqrt{5})D + 10(1 + \sqrt{5})C - 40(1 + \sqrt{5})B + 480(3 + \sqrt{5})\} \\
 &\quad + y^3z^3\{-20E - 40D + 0C + 0B + 1280\} \\
 &\quad + y^2z^4\{15E + 10(3 - \sqrt{5})D + 10(1 - \sqrt{5})C - 40(1 - \sqrt{5})B + 480(3 - \sqrt{5})\} \\
 &\quad + yz^5\{-6E - 2(6 - 4\sqrt{5})D - 8(2 - \sqrt{5})C - 16(1 - \sqrt{5})B + 192(7 - 3\sqrt{5})\} \\
 &\quad + z^6\{E + 2(1 - \sqrt{5})D + (6 - 2\sqrt{5})C + (56 - 24\sqrt{5})B + 16(36 - 16\sqrt{5})\}].
 \end{aligned}$$

We will show that  $(M'') \in \text{Aut}(f'')$  for some  $b$  and  $c$ . The coefficients of  $x^4yz$ ,  $x^2y^3z$ ,  $x^2yz^3$ ,  $y^5z$ ,  $yz^5$  and  $y^3z^3$  in  $f''$  vanish. Note that  $E = 64 - D \neq 0$ , for  $D^2 - 9D + 864 = 0$ . So such  $b$  and  $c$  exist if and only if  $f''_{M''^{-1}} = -f''$ . Let us denote by  $[x^jy^kz^\ell]$  the coefficient of the monomial  $x^jy^kz^\ell$  in  $f''$ . Then the following equalities hold:

(1)  $b^2[x^4y^2] = -[x^4z^2]$  (2)  $b^4[x^2y^4] = [x^2z^4]$  (3)  $b^6[y^6] = -[z^6]$  (4)  $b^2[y^4z^2] = -[y^2z^4]$ .

We can show that the equality (1) implies (2) through (4). To be more precise, assume that  $b$  is a solution to (1) for given  $D$ . (1) gives  $b^4[x^4y^2]^2 = [x^4z^2]^2$ , which implies (2), because  $[x^4y^2]^2[x^2z^4] - [x^4z^2]^2[x^2y^4] = 0$ . (1) and (2) give  $b^6[x^4y^2][x^2y^4] = -[x^4z^2][x^2z^4]$ , which implies (3). (4) is exactly the same condition as (1). This completes the proof of Proposition 2.22. □

ACKNOWLEDGEMENT. We would like to thank Prof. S. Yoshiara at Osaka Kyoiku University who kindly informed us the structure of finite groups. We owe many im-

provements in the representation to the referee, especially in the proof of Proposition 1.4.

---

### References

- [1] H.M. Farkas and I. Kra: *Riemann Surfaces*, 2nd ed., Springer, 1992.
- [2] W. Fulton: *Algebraic Curves*, Addison-Wesely, 1989.
- [3] M. Hall, Jr.: *The Theory of Groups*, Macmillan, New York, 1968.
- [4] K. Idei: *The maximum order of the automorphism groups of non-singular plane sextics*, MS Thesis, Okayama Univ., 1998.
- [5] S. Itaka: *Algebraic Geometry II*(in Japanese), Iwanami, 1977.
- [6] H. Kaneta, S. Marcugini and F. Pambianco: *On arcs and curves with many automorphisms*, preprint.
- [7] H. Kaneta, S. Marcugini and F. Pambianco: *The most symmetric non-singular cubics, quintics and septics*, preprint.
- [8] H. W. Leopoldt: *Über die Automorphismen Gruppe des Fermatkörpers*, *J. Number Theory*, **56** (1996), 256–282.
- [9] M. Namba: *Geometry of Projective Algebraic Curves*, Marcel Dekker, 1984.
- [10] M. Namba: *Geometry of Algebraic Curves*(in Japanese), Gendai-suugaku-sha, 1991.
- [11] A. Wiman: *Übere eine einfache Gruppe von 360 ebenen Collineationen*, *Math. Annalen*, bd. **47** (1896), 531–556.

H. Doi  
 Department of Mathematics  
 Faculty of Science  
 Okayama University  
 Okayama 700-8530, Japan

K. Idei  
 CJK CO., LTD  
 5-3 Kouji-machi  
 Chiyoda-ku  
 Tokyo 102-0083, Japan

H. Kaneta  
 Department of Mathematical Sciences  
 College of Engineering  
 Osaka Prefecture University  
 Sakai, Osaka 599-8531, Japan

