

ON THE DISTANCE BETWEEN TWO SEIFERT SURFACES OF A KNOT

Dedicated to Professor Takao Matumoto on the occasion of his sixtieth birthday

MAKOTO SAKUMA and KENNETH J. SHACKLETON

(Received January 18, 2007, revised November 30, 2007)

Abstract

For a knot K in \mathbb{S}^3 , Kakimizu introduced a simplicial complex whose vertices are all the isotopy classes of minimal genus spanning surfaces for K . The first purpose of this paper is to prove the 1-skeleton of this complex has diameter bounded by a function quadratic in knot genus, whenever K is atoroidal. The second purpose of this paper is to prove the intersection number of two minimal genus spanning surfaces for K is also bounded by a function quadratic in knot genus, whenever K is atoroidal. As one application, we prove the simple connectivity of Kakimizu's complex among all atoroidal genus 1 knots.

1. Introduction

Let K be a knot in the 3-sphere \mathbb{S}^3 . A *Seifert surface* for K is a compact, connected and orientable surface in \mathbb{S}^3 whose boundary is precisely K . Fix a regular neighbourhood $N(K)$ for the knot K , and denote by $E(K)$, or just E , its exterior $\mathbb{S}^3 - \text{int } N(K)$. We say that E , or K , is *atoroidal* if every incompressible torus in E is boundary parallel. We shall say that a properly embedded subsurface of E is a *spanning surface for K* if it is contained in some Seifert surface for K . For any spanning surface or Seifert surface S , we denote its ambient isotopy class by $[S]$. Throughout this paper, we shall assume, unless otherwise stated, that any given pair of Seifert surfaces or spanning surfaces intersects transversely.

To the knot K there is an associated graph $\mathcal{G}(K)$, constructed as follows. The vertex set comprises the isotopy classes of minimal genus spanning surfaces for K , and two distinct vertices are connected by an edge if and only if they can be represented by a pair of disjoint spanning surfaces in E . It is a result of Scharlemann-Thompson (Proposition 5 from [14]) that $\mathcal{G}(K)$ is connected. As it happens, their main argument implies $d(\sigma, \sigma') \leq \iota(\sigma, \sigma') + 1$, where σ and σ' are two isotopy classes of spanning surfaces, d is the path-metric on $\mathcal{G}(K)$ assigning length 1 to each edge, and $\iota(\sigma, \sigma')$ denotes the least number of components $|S \cap S'|$ among all pairs $S \in \sigma, S' \in \sigma'$ of spanning surfaces intersecting transversely. We refer to $\iota(\sigma, \sigma')$ as the *intersection number*

of σ and σ' . It is a result of Jaco-Sedgwick (see Oertel [10]) that $\mathcal{G}(K)$ is a finite graph whenever K has genus at least 2 and is atoroidal. By the work of Hayashi [3] and of Wilson [21] it follows that $\mathcal{G}(K)$ is a finite graph whenever K is atoroidal even if it has genus 1. However, it is a recent result of Tsutsumi [19] that there exists an infinite sequence of atoroidal knots of common genus, in each genus at least 2, with an increasing number of isotopy classes of minimal genus spanning surfaces. In particular, the number of vertices of $\mathcal{G}(K)$ among all atoroidal knots K of any common genus, at least 2, is unbounded.

Our first result offers a uniform bound on the diameter of $\mathcal{G}(K)$, quadratic in knot genus $g(K)$.

Theorem 1.1. *Suppose that K is an atoroidal knot in \mathbb{S}^3 . Then, $\mathcal{G}(K)$ has diameter at most $2g(K)(3g(K) - 2) + 1$.*

The assumption that K be atoroidal is necessary here, for Theorem B of [6] asserts that the graph $\mathcal{G}(K)$ is a bi-infinite line for many composite knots K . Such knots are toroidal, and such graphs are unbounded.

One can go further and place a uniform quadratic bound on the intersection number of any pair of minimal genus Seifert surfaces.

Theorem 1.2. *Suppose that K is an atoroidal knot in \mathbb{S}^3 . Then, $\iota([S], [S'])$ is at most $2(3g(K) - 2)^2$ for any two minimal genus Seifert surfaces S and S' for K .*

We note Theorem 1.2 also offers a second uniform bound on the diameter of $\mathcal{G}(K)$, though less desirable than that already given in Theorem 1.1. A useful consequence of Theorem 1.2 is the following.

Corollary 1.3. *Suppose that K is an atoroidal genus 1 knot in \mathbb{S}^3 . Then, $\mathcal{G}(K)$ has diameter at most 2.*

The better bound on diameter offered by Corollary 1.3 for all atoroidal genus 1 knots is in fact sharp.

Proposition 1.4. *There exists an atoroidal genus 1 knot K in \mathbb{S}^3 such that $\mathcal{G}(K)$ has diameter 2.*

Recall $\mathcal{G}(K)$ is also the 1-skeleton of the corresponding simplicial complex $\mathcal{MS}(K)$ constructed by Kakimizu [6], where a set of pairwise distinct vertices spans a simplex if and only if they may simultaneously be realised disjointly in E . That this complex is flag, so that any inclusion of the boundary of any given simplex extends over the whole simplex, is Proposition 4.9 from [13]. Applications of this complex are found in classifying the incompressible Seifert surfaces of prime knots of at most 10 crossings

(see [7]), using a method that enhances that of Kobayashi's [9]. In [13], the first author proves the contractibility of $\mathcal{MS}(K)$ when K is a special aborescent link. In his joint paper with Hirasawa [4], contractibility when K is a prime, special, alternating link is announced. Together, this partially verifies a challenging conjecture of Kakimizu's [5], asserting among other things that $\mathcal{MS}(K)$ is always contractible. A full statement is given as Conjecture 0.2 in [13].

Applying Theorem 1.2 and Corollary 1.3, we will prove the following.

Proposition 1.5. *Suppose that K is an atoroidal knot of genus 1. Then, the simplicial complex $\mathcal{MS}(K)$ is simply connected.*

We remark the dimension of $\mathcal{MS}(K)$ is at most 6 whenever the knot is atoroidal and has genus 1, as follows from Tsutsumi [18].

The structure of this paper is as follows. In Section 2 we shall recall Kakimizu's characterisation of the metric on $\mathcal{G}(K)$. In Section 3 we use Section 2 to prove Theorem 1.1. In Section 4 we prove Theorem 1.2, and then Corollary 1.3. In Section 5 we prove Proposition 1.5. In each of Section 3, Section 4 and Section 5, we will rely on Theorem 3.1 of Fenley [1] which, among other things, rules out the existence of an accidental peripheral in any minimal genus Seifert surface whenever the knot is atoroidal. To recall, an *accidental peripheral* on a surface S in \mathbb{S}^3 is a simple loop essential on S which is homotopic, in E , to a loop on ∂E . In Section 6 we prove Proposition 1.4. In Section 7 we investigate the growth in diameter of $\mathcal{G}(K)$ with knot genus $g(K)$, proving the following.

Proposition 1.6. *For every positive integer g , there is an atoroidal knot K of genus g such that the diameter of $\mathcal{G}(K)$ is equal to $2g - 1$.*

It seems appropriate to close the introduction by posing the following open question.

QUESTION 1.7. Considering the quadratic upper bound on diameter offered by Theorem 1.1, can this be improved to a linear function of knot genus?

REMARK. A few months after making the first version of this paper publicly available, the authors learned in [11] that Roberto Pelayo has independently since found a version of Theorem 1.1 as part of his PhD thesis, in preparation under the supervision of Danny Calegari.

2. A characterisation of distance

Let us recall Kakimizu's characterisation of the metric on $\mathcal{G}(K)$ before giving a proof to Theorem 1.1. For a knot K in \mathbb{S}^3 let E denote its exterior and consider the infinite cyclic cover $\phi: \tilde{E} \rightarrow E$, denoting by τ a generator for the deck transformation

group. Let S be any minimal genus spanning surface for K and denote by E_0 the closure in \tilde{E} of any ϕ -lift of the complement $E - S$. Set $E_j = \tau^j(E_0)$ and $S_j = E_{j-1} \cap E_j$ for each integer j .

For a second such spanning surface S' we may similarly form E'_0 , the closure in \tilde{E} of any lift of $E - S'$ via ϕ , and then denote by E'_j the translate $\tau^j(E'_0)$ for each integer j . Setting $m_- = \min\{k \in \mathbb{Z}: E_0 \cap E'_k \neq \emptyset\}$ and $m_+ = \max\{k \in \mathbb{Z}: E_0 \cap E'_k \neq \emptyset\}$, we can then define $d_*(S, S') = m_+ - m_-$. Finally, for any two vertices σ and σ' of $\mathcal{G}(K)$ we define $d_*(\sigma, \sigma') = \min\{d_*(S, S'): S \in \sigma, S' \in \sigma'\}$. The following statement combines two key results due to Kakimizu, Proposition 1.4 and Proposition 3.2 (2) of [6].

Proposition 2.1 ([6]). *The function d_* is a metric on the vertex set of $\mathcal{G}(K)$. Moreover, for any two vertices σ and σ' of $\mathcal{G}(K)$, we have $d(\sigma, \sigma') = d_*(\sigma, \sigma')$.*

3. Proof of Theorem 1.1

We begin by proving two statements regarding the intersection of a pair of spanning surfaces.

Lemma 3.1. *Suppose that S and S' are two distinct minimal genus spanning surfaces for the atoroidal knot K . Then, S' is ambient isotopic to a third minimal genus spanning surface S'' such that $S \cap S''$ is a disjoint union of loops and such that $d_*(S, S'') \leq d_*(S, S')$.*

Proof. Suppose $S \cap S'$ contains an arc component, that is ∂S and $\partial S'$ intersect. Then, ∂S and $\partial S'$ bound a bigon on ∂E , because they are isotopic on ∂E . Using this bigon, we deduce S' is isotopic to a second spanning surface S'' such that $|\partial S \cap \partial S''| \leq |\partial S \cap \partial S'| - 2$ while $d_*(S, S'') \leq d_*(S, S')$. A proof can now be completed by induction. \square

Lemma 3.2. *Suppose that S and S' are two distinct minimal genus spanning surfaces for the atoroidal knot K intersecting only in loops. Then, S' is ambient isotopic to a third minimal genus spanning surface S'' such that $S \cap S''$ is a disjoint union of loops, essential both on S and on S'' , and such that $d_*(S, S'') \leq d_*(S, S')$.*

Proof. Suppose for contradiction some component α of $S \cap S'$ is inessential on S , that is α is null-homotopic on S or is boundary parallel on S . If α is null-homotopic on S then, by the incompressibility of S' , it must also be null-homotopic on S' . The irreducibility of E then allows us to reduce the cardinality $|S \cap S'|$ by an isotopy of S or of S' without increasing $d_*(S, S')$. We may thus assume α is boundary parallel on S . By Theorem 3.1 of [1], α must also be boundary parallel on S' .

There exist two Seifert surfaces extending S and S' , respectively, whose intersection is precisely $(S \cap S') \cup K$. Out of convenience, we shall proceed by using S and

S' to respectively denote such a pair of Seifert surfaces. Replacing α with a second component of $S \cap S'$, boundary parallel on S , we may assume α borders an annulus $A \subset S$ such that $A \cap S' = \partial A$, the union $\alpha \cup K$. Let $A' \subset S'$ be the annulus bounded by ∂A . The union $A \cup A'$, denoted by T , is an embedded torus in \mathbb{S}^3 such that $K \subset T$.

Let V be a solid torus in \mathbb{S}^3 bounded by T . Since K is a non-trivial and atoroidal knot, K is isotopic to the core of V and hence the pair (V, A') can be given a product structure $(A' \times [0, 1], A' \times \{0\})$. We deduce S' is isotopic to a second surface S'' such that $|\partial S \cap \partial S''| < |\partial S \cap \partial S'|$ while $d_*(S, S'') \leq d_*(S, S')$. A proof can now be completed by induction. \square

REMARK. We claim that in fact $S' \cap V = A'$, so that the above isotopy of S' can be chosen to fix pointwise the complement of a regular neighbourhood of A' in S' . To see this, let us argue by supposing otherwise. Let S'_* denote the open surface $S' - A'$. The intersection of S'_* with V is then by assumption non-empty and, as S'_* is connected and as $S'_* \cap T = (S' - A') \cap \text{int } A = \emptyset$, so S'_* is entirely contained in the interior of V . However, the closure of S'_* is a surface in V whose only boundary component is α . That is, $[\alpha]$ is trivial in $H_1(V, \mathbb{Z})$ despite α being a longitude for V . We have a contradiction, and we deduce $S' \cap V = A'$.

We shall henceforth denote by η the function $3g - 2$ on the set of all knots, noting $\eta(K)$ is the size of any maximal collection of pairwise disjoint and non-isotopic essential simple loops on any minimal genus spanning or Seifert surface for K . Since an estimate on diameter is easily found for a trivial knot, we shall also assume the genus of K is positive.

Let σ and σ' be any two vertices of $\mathcal{G}(K)$, and take representatives $S \in \sigma$ and $S' \in \sigma'$ so that $d_*(S, S')$ is minimal and so that S and S' intersect transversely in a disjoint union of essential simple loops, as per Lemma 3.1 and Lemma 3.2.

Suppose for contradiction $d(\sigma, \sigma') \geq 2g(K)\eta(K) + 2$. According to Proposition 2.1, we also have $d_*(S, S') \geq 2g(K)\eta(K) + 2$. Let Y_j denote the surface $E_j \cap S'_0$ for each integer j where, perhaps after reindexing, we may assume Y_j is non-empty if and only if $1 \leq j \leq d_*(S, S')$. It should be noted $\chi(Y_j) \leq 0$ for each such index j , so that $|\chi(S'_0)| = \sum_{j=1}^{d_*(S, S')} |\chi(Y_j)|$. Here, $\chi(S)$ denotes the Euler characteristic of a surface S .

Claim. *For the finite sequence $\{Y_j: 2 \leq j \leq d_*(S, S') - 1\}$ of non-empty surfaces, there are $\eta(K)$ consecutive indices whose corresponding surfaces each have zero Euler characteristic. That is, there exists a natural number r , with $1 \leq r \leq d_*(S, S') - \eta(K) - 1$, such that $\chi(Y_j) = 0$ for each j with $r + 1 \leq j \leq r + \eta(K)$.*

Proof. Set $\|K\|$ equal to $2g(K) - 1$, the Thurston norm [16] of either generator of $H_2(E, \partial E)$. In the identity $\|K\| = |\chi(S'_0)| = \sum_{j=1}^{d_*(S, S')} |\chi(Y_j)|$, only at most $\|K\|$ of the summands $|\chi(Y_j)|$ can be non-zero. The proof will now be completed by a pigeonhole-type argument, in the following manner.

We denote by w the string $(w_2, \dots, w_{d_*(S, S')-1})$ of binary digits, where w_j is defined equal to 0 if $\chi(Y_j)$ is 0 or otherwise 1 for each of the indices $j \in \{2, \dots, d_*(S, S')-1\}$. If it should happen that for any $\eta(K)$ consecutive binary digits w_j at least one is always non-zero, we would then have the estimate $|w| \leq (\|K\| + 1)\eta(K) - 1$ on the length $|w|$ of w . This follows from the fact that only at most $\|K\|$ of the binary digits w_j can be non-zero. We can now find an upper bound for $d(\sigma, \sigma')$ as follows:

$$\begin{aligned} d(\sigma, \sigma') &= d_*(S, S') \\ &= |w| + 2 \\ &\leq (\|K\| + 1)\eta(K) + 1 \\ &= 2g(K)\eta(K) + 1. \end{aligned}$$

According to our standing assumption on $d(\sigma, \sigma')$, this is absurd. We deduce that there exist $\eta(K)$ consecutive zeros $w_{r+1}, \dots, w_{r+\eta(K)}$, thus proving the claim. \square

After shifting the indexing E_j by r , we have $|\chi(Y_j)| = 0$ for $1 \leq j \leq \eta(K)$, and the set $Y_1 \cup \dots \cup Y_{\eta(K)}$ is both non-empty and a union of pairwise disjoint annuli. Note that both Y_0 and $Y_{\eta(K)+1}$ are necessarily non-empty. In particular, both $S_1 \cap S'_0$ and $S_{\eta(K)+1} \cap S'_0$ are non-empty. There thus exists a subannulus A of $Y_1 \cup \dots \cup Y_{\eta(K)}$ ending on S_1 and on $S_{\eta(K)+1}$.

Claim. *There exist natural numbers p and q , with $1 \leq p < q \leq \eta(K) + 1$, for which there is a component α of $A \cap S_p$ and a component β of $A \cap S_q$ such that $\phi(\alpha)$ and $\phi(\beta)$ are isotopic loops on S .*

Proof. For each $1 \leq j \leq \eta(K)+1$, $\phi(A \cap S_j)$ is a non-empty collection of essential, pairwise disjoint and pairwise non-isotopic loops on S , and the two sets $\phi(A \cap S_i)$ and $\phi(A \cap S_j)$ are disjoint for distinct i and j . Since any collection of pairwise disjoint and non-isotopic essential simple loops on S has size at most $\eta(K)$, we deduce the claim. \square

Let \mathcal{A} be the family of all those subannuli of A bounded by any pair of loops found in the previous claim. Then, \mathcal{A} is non-empty and we can choose $A' \in \mathcal{A}$ minimal subject to inclusion. The annulus $A'' \subset S$ bounded by $\phi(\partial A')$ has interior disjoint from $\phi(A')$ and so the union $\phi(A') \cup A''$, denoted T , is an embedded torus in E .

Claim. *T is incompressible in E .*

Proof. We shall check the inclusion $e: T \rightarrow E$ induces an injection $e_*: \pi_1(T) \rightarrow \pi_1(E)$ on fundamental groups. Let α be either component of $\partial A'$, and let $p < q$ be such that $\partial A' \subset S_p \cap S_q$. Let the simple loop $\gamma \subset T$ be the union of an arc in $\phi(A')$ and an arc in A'' . Observe that $\pi_1(T)$ is generated by $\phi(\alpha)$ and γ , and that the image

of $\phi(\alpha)$ and the image of $e_*(\gamma)$ in $H_1(E) \cong \mathbb{Z}$ are 0 and $q - p$, respectively. Since $q - p$ is non-zero, it follows $\text{Ker}(e_*)$ is contained in the group $\langle \phi(\alpha) \rangle$. However, $\phi(\alpha)$ is an essential loop on the incompressible surface S in E . Hence $\text{Ker}(e_*)$ is trivial, and T is incompressible in E . \square

Claim. T is essential in E .

Proof. The loop $\phi(\alpha) \subset T$ is essential on S . It follows from Theorem 3.1 [1] that $\phi(\alpha)$ is not isotopic in $E(K)$ to a simple loop on ∂E . Hence T can not be boundary parallel in E . \square

We now have a contradiction, for K is atoroidal, and we deduce $2g(K)\eta(K) + 1$ is an upper bound for the diameter of $\mathcal{G}(K)$. This completes a proof of Theorem 1.1.

4. Proof of Theorem 1.2

We shall once more denote by $\eta(K)$ the number $3g(K) - 2$. An argument similar to that found in the proof of both Lemma 3.1 and Lemma 3.2 permits us to represent any given pair of vertices of $\mathcal{G}(K)$ by a pair of spanning surfaces for K intersecting transversely and minimally, up to isotopy, in loops essential on both surfaces.

Let S and S' be a pair of such spanning surfaces. Suppose for contradiction that $|S \cap S'| \geq 2\eta(K)^2 + 1$. Then, there exist two distinct annuli $A \subset S$ and $A' \subset S'$ such that $|A \cap A'| = 3$ and $\partial A \cup \partial A' \subset A \cap A'$. To see this, consider an $l \times m$ array, with $1 \leq l, m \leq \eta(K)$, whose entries are non-negative integers summing to $2\eta(K)^2 + 1$. It is not so hard to see that at least one of these entries must be at least 3. We may further assume A is minimal subject to inclusion, so that no component of $A \cap (S' - A')$ is isotopic on S' to the core of A' .

The union $A \cup A'$ always separates the 3-sphere into three components, whose closures we denote by X_1, X_2, X_3 and indexed so that $K \subset X_3$. Note it can happen that one of the X_i fails to be a manifold, in which case its frontier, $\text{fr} X_i$, is homeomorphic to an immersed torus whose singular set is the simple loop $\text{int } A \cap \text{int } A'$.

Claim. $\text{int } X_3$ is not an open solid torus.

Proof. Suppose for contradiction $\text{int } X_3$ is an open solid torus. As the knot K is atoroidal, so either K is contained in a compact 3-ball inside $\text{int } X_3$ or K is a core of $\text{int } X_3$, and we rule out both cases separately as follows.

CASE I. K is contained in a compact 3-ball $B \subset \text{int } X_3$. Since S is connected and since $S \cap \text{fr } X_3$ contains a simple loop essential on S , so $S \cap \partial B$ also contains a simple loop essential on S . Thus, there exists a disc D disjoint from the knot K whose boundary ∂D is a non-trivial simple loop on S . However, S is incompressible and we therefore have a contradiction.

CASE II. K is a core of $\text{int } X_3$. Let F denote the open surface $S \cap (\text{int } X_3 - N(K))$. Then, F is necessarily a non-empty disjoint union of open annuli, for the inclusion of F in E descends to a monomorphism on fundamental groups that factors through the abelian group $\pi_1(\text{int } X_3 - N(K)) \cong \mathbb{Z} \oplus \mathbb{Z}$. Thus, S contains an annulus with one boundary component equal to K and the other a component of $S \cap S'$. That is, S and S' intersect in at least one simple loop peripheral on S . However, this is contrary to the standing assumption that S and S' intersect only in loops essential on S (and on S'). \square

To complete a proof of Theorem 1.2 it suffices to rule out the following two mutually exclusive cases. These correspond to the two distinct ways in which A and A' can intersect one another.

CASE I. $\text{fr } X_1, \text{fr } X_2, \text{fr } X_3$ are each tori. Then, at least one of $X_1 \cup X_2$ and X_3 is a solid torus and, according to the claim, it can only be $X_1 \cup X_2$. It follows that both X_1 and X_2 are solid tori, and, using van Kampen's theorem, at least one, say X_1 , has a product structure $(X_1, A'') \cong (A'' \times [0, 1], A'' \times \{0\})$, where $A'' \subseteq \text{fr } X_1 \cap S$ is an annulus. Thus, S is ambient isotopic to a second Seifert surface intersecting S' fewer than $\iota([S], [S'])$ times and this is absurd.

CASE II. Exactly one of $\text{fr } X_1, \text{fr } X_2, \text{fr } X_3$ is not an embedded torus. Then, either $\text{fr } X_3$ is not an embedded torus or exactly one of $\text{fr } X_1$ or $\text{fr } X_2$ is not an embedded torus. We thus need only consider the following two subcases.

SUBCASE II.1. $\text{fr } X_3$ is an immersed singular torus. Then, $N(X_1 \cup X_2)$ is necessarily a solid torus. It follows both X_1 and X_2 are solid tori and at least one, say X_1 , has a product structure $(X_1, A'') \cong (A'' \times [0, 1], A'' \times \{0\})$, where $A'' \subseteq \text{fr } X_1 \cap S$ is an annulus. Once more, we will find S is ambient isotopic to a second Seifert surface intersecting S' fewer than $\iota([S], [S'])$ times and this is absurd.

SUBCASE II.2. $\text{fr } X_3$ is an embedded torus. Then, $X_1 \cup X_2$ is necessarily a solid torus. Let F' denote that component of $S' \cap (X_1 \cup X_2)$ containing $\text{int } A \cap \text{int } A'$. The minimality of A implies $\partial F' - \partial A$ comprises of loops essential on S' , none of which is isotopic on S' to the core of A' . Hence $\pi_1(F')$ is non-abelian and, as S' is incompressible, the inclusion of F' in E descends to a monomorphism on fundamental groups. In particular, it has non-abelian image. However, this monomorphism also factors through the abelian group $\pi_1(X_1 \cup X_2) \cong \mathbb{Z}$ and as such has abelian image, a contradiction.

We thus complete a proof of Theorem 1.2. Let us finish this section by providing a proof of Corollary 1.3.

Proof of Corollary 1.3. Let σ_1 and σ_2 be two vertices of $\mathcal{G}(K)$. Let $S_1 \in \sigma_1$ and $S_2 \in \sigma_2$ be a pair of representative spanning surfaces, together realising intersection number, and such that $S_1 \cap S_2$ is a collection of loops, perhaps empty, essential both on S_1 and on S_2 . Then, since $g(S_1)$ and $g(S_2)$ are both equal to 1, so $S_1 \cap S_2$ comprises only of non-separating loops, parallel on S_1 and on S_2 . Applying to Theorem 1.2, we

have $|S_1 \cap S_2| \leq 2$ and it follows that each lift of S_2 intersects at most one lift of S_1 . Thus, $d_*(S_1, S_2) \leq 2$. By Proposition 2.1, we have $d(\sigma_1, \sigma_2) \leq 2$ as required. \square

5. Proof of Proposition 1.5

We shall need the following criterion for the simple connectivity of a simplicial complex whose 1-skeleton is a metric graph of diameter at most 2, and then a restriction on the intersection number of two genus 1 spanning surfaces.

Lemma 5.1. *Suppose \mathcal{C} is a simplicial complex, whose 1-skeleton can be realised as a metric graph of diameter at most 2, for which every simplicial circuit of length at most 5 is contractible. Then, \mathcal{C} is simply connected.*

Proof. Let $\sigma_1, \dots, \sigma_n$ be the cyclically indexed vertices of a circuit c of length n . Since $d(\sigma_1, \sigma_i) \leq 2$ for $3 \leq i \leq n-1$, so there exists a simplicial path of length at most 2 connecting σ_1 to σ_i for each such i . It follows c can be expressed as a finite sum of simplicial 3-, 4- and 5-circuits. Each such circuit is contractible, by assumption, and so c must also be contractible. Hence, \mathcal{C} is simply connected. \square

By the proof of Corollary 1.3, we have the following lemma.

Lemma 5.2. *Let K be an atoroidal genus 1 knot. Then, for any pair of vertices σ and σ' of $\mathcal{G}(K)$, we have $\iota(\sigma, \sigma') \in \{0, 2\}$.*

In proving the following lemma, we shall make use of a construction that amounts to a special case of the so-called *double curve sum*, after Scharlemann-Thompson [14], and of a construction of Kakimizu's [6].

Lemma 5.3. *Let K be an atoroidal genus 1 knot. For any pair of vertices σ and σ' of $\mathcal{G}(K)$, with $d(\sigma, \sigma') = 2$, there exists a third vertex δ such that $d(\delta, \sigma) = d(\delta, \sigma') = 1$ and such that, for any fourth vertex μ , if $\iota(\mu, \sigma) = \iota(\mu, \sigma') = 0$, then $\iota(\mu, \delta) = 0$.*

Proof. By Lemma 5.2 and arguments given in Section 3, there exist representatives $S \in \sigma$ and $S' \in \sigma'$ such that $S \cap S'$ is a pair of loops essential both on S and on S' . Let $P \subset S$ denote the 3-holed sphere bordered by ∂S and by $S \cap S'$, and let $A' \subset S'$ be the closed annulus bordered by $S \cap S'$. Then, $P \cup A'$ is a genus 1 spanning surface for the knot K and, after a small isotopy, is disjoint from both S and S' . We take δ to be the isotopy class $[P \cup A']$, noting that $\iota(\delta, \sigma) = \iota(\delta, \sigma') = 0$ by construction. Since $d(\sigma, \sigma') = 2$, so $d(\delta, \sigma) = d(\delta, \sigma') = 1$.

Now suppose μ is a fourth vertex, adjacent to both σ and σ' . We claim that $\iota(\delta, \mu) = 0$, and to prove this it suffices to prove the existence of a representative of μ

simultaneously disjoint both from S and from S' . By assumption, $\iota(\mu, \sigma) = 0$ and hence there exists a representative $R \in \mu$ disjoint from S . Perhaps after replacing R with an isotopic surface also disjoint from S , we may further assume that R is transverse to S' .

As $\iota(\mu, \sigma') = 0$, by Proposition 4.8 (2) of [13] there exists a product region V between R and S' such that $V \cap R = \text{fr } V \cap R$ and $V \cap S' = \text{fr } V \cap S'$. Note that, should $S \cap V$ not be empty, then $S \cap V$ is parallel in V to a subsurface of $S' \cap V$, by Corollary 3.2 of [20]. It follows S and S' would share a removable intersection, and this is absurd. Thus, $S \cap V$ is empty. We can therefore use the region V to replace R with an isotopic surface R' such that R' and S are disjoint and such that $|R' \cap S'| \leq |R \cap S'| - 1$.

Continuing inductively, we deduce R is isotopic to a spanning surface simultaneously disjoint both from S and from S' . \square

Lemma 5.4. *Let K be an atoroidal genus 1 knot. Suppose σ, σ_1 and σ_2 are three vertices of $\mathcal{G}(K)$ such that $d(\sigma, \sigma_1) = d(\sigma, \sigma_2) = 2$ and such that $d(\sigma_1, \sigma_2) = 1$. Then, there exist two vertices δ_1 and δ_2 of $\mathcal{G}(K)$ such that $d(\sigma, \delta_i) = d(\delta_i, \sigma_i) = 1$, for $i \in \{1, 2\}$, and such that $d(\delta_1, \delta_2) \leq 1$.*

Proof. Let $S \in \sigma, S_1 \in \sigma_1$ and $S_2 \in \sigma_2$ be such that $S_1 \cap S_2$ is empty and such that S intersects both S_1 and S_2 transversely and in a collection of loops essential on each surface.

Let \tilde{E} denote the infinite cyclic cover of the knot exterior E , with covering map denoted ϕ , and denote by τ either generator of the deck transformation group. Let $\tilde{S}_{1,0}$ denote any lift of S_1 , and let $\tilde{S}_{1,n}$ denote the translate $\tau^n(\tilde{S}_{1,0})$ for each integer $n \in \mathbb{Z}$. We similarly introduce the notation $\tilde{S}_{2,n}$, where $\tilde{S}_{2,0}$ is to separate $\tilde{S}_{1,0}$ and $\tilde{S}_{1,1}$.

The following claim permits us to isotope S so that in addition each lift of S intersects only one lift of S_1 and only one lift of S_2 . Recall the definition of the function d_* from Section 2.

Claim. *There exists an isotopy of S after which $d_*(S, S_1) = d_*(S, S_2) = 2$.*

Proof. Suppose $d_*(S, S_1) + d_*(S, S_2) \geq 5$, and denote by \tilde{S} any lift of S . Then, since $d_*(\sigma, \sigma_1) = d_*(\sigma, \sigma_2) = 2$, so there exists a component R of $\phi^{-1}(S_1 \cup S_2)$ such that $R \cap \tilde{S}$ is not empty and for which there exists an isotopy of S lifting to an isotopy of \tilde{S} after which $R \cap \tilde{S}$ is empty. By Proposition 4.8 (2) of [13], there thus exists a product region V in \tilde{E} between R and \tilde{S} and such that $V \cap R = \text{fr } V \cap R$ and $V \cap \tilde{S} = \text{fr } V \cap \tilde{S}$. As \tilde{S} and each component of $\phi^{-1}(S_1 \cup S_2)$ separate in \tilde{E} , so there exists a subregion $V' \subseteq V$ such that $V' \cap (\tilde{S} \cup \phi^{-1}(S_1 \cup S_2)) = \text{fr } V'$. Let us denote by R' the one component of $\phi^{-1}(S_1 \cup S_2)$ such that $R' \cap V'$ is not empty.

Applying Corollary 3.2 of [20] to the product region V , we find $R' \cap V'$ and $\tilde{S} \cap V'$ are parallel through V' . Note, V' is contained in a single fundamental region. Projecting V' to E then, we can therefore isotope S so as to remove the corresponding

intersection between R' and \tilde{S} and without introducing any new intersections between $\phi^{-1}(S_1 \cup S_2)$ and \tilde{S} .

That is, so long as $d_*(S, S_1) + d_*(S, S_2) \geq 5$, we can successively remove intersections between $\phi^{-1}(S_1 \cup S_2)$ and \tilde{S} via an isotopy of S . There are only finitely many such intersections to begin with, thus in finite time we construct an isotopy of S after which $d_*(S, S_1) + d_*(S, S_2) \leq 4$. The statement of the claim is deduced. \square

Isotope S as indicated by the claim, and denote by \tilde{S} the lift of S intersecting $\tilde{S}_{1,0}$ and $\tilde{S}_{2,0}$. Now let N denote a small regular neighbourhood of $\tilde{S} \cup \tilde{S}_{1,0} \cup \tilde{S}_{2,0}$ in the infinite cyclic cover \tilde{E} , so that $\tau^j(N)$ is disjoint from \tilde{S} , $\tilde{S}_{1,0}$ and $\tilde{S}_{2,0}$ for each non-zero integer j . We define \tilde{R}_1 and \tilde{R}_2 to be the two “outermost” components of $\text{fr } N$, that is \tilde{R}_1 and \tilde{R}_2 bound a region in \tilde{E} containing N and indexed so that \tilde{R}_1 and $\tilde{S}_{2,0}$ are separated by $\tilde{S}_{1,0}$. Note, \tilde{R}_1 and $\tau^{-1}(\tilde{R}_2)$ are disjoint, are both 1-holed tori, and are both contained in the fundamental region bordered by $\tilde{S}_{1,-1}$ and $\tilde{S}_{1,0}$. Thus, \tilde{R}_1 and $\tau^{-1}(\tilde{R}_2)$ project to disjoint genus 1 spanning surfaces, denoted respectively R_1 and R_2 , both of which are disjoint from S_1 and from S_2 .

Finally, we respectively define δ_1 and δ_2 to be the isotopy classes $[R_1]$ and $[R_2]$. This completes a proof of Lemma 5.4. \square

In view of Corollary 1.3 and Lemma 5.1, to prove the simple connectivity of $\mathcal{MS}(K)$, for an atoroidal genus 1 knot K , it suffices to prove the following three claims.

Claim. *Every simplicial 3-circuit in $\mathcal{MS}(K)$ is contractible.*

Proof. This is immediate, for $\mathcal{MS}(K)$ is a flag simplicial complex. That is, any embedding of the 1-skeleton of any given simplex into $\mathcal{G}(K)$ is the restriction of an embedding from the whole simplex into $\mathcal{MS}(K)$. \square

Claim. *Every simplicial 4-circuit in $\mathcal{MS}(K)$ is contractible.*

Proof. Suppose $\sigma_1, \sigma_2, \sigma_3, \sigma_4$ are the cyclically indexed vertices of a simplicial 4-circuit in $\mathcal{MS}(K)$. Assuming $d(\sigma_1, \sigma_3) = 2$, by Lemma 5.3 there exists a vertex δ such that $d(\delta, \sigma_1) = d(\delta, \sigma_3) = 1$ and such that $\iota(\delta, \sigma_2) = \iota(\delta, \sigma_4) = 0$. We deduce δ spans an edge with σ_2 and with σ_4 . Appealing to the previous claim, one may now find an appropriate compressing disc as the union of at most four 2-simplices.

The remaining cases may be similarly treated. \square

Claim. *Every simplicial 5-circuit in $\mathcal{MS}(K)$ is contractible.*

Proof. Suppose $\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5$ are the cyclically indexed vertices of a simplicial 5-circuit in $\mathcal{MS}(K)$. Assuming $d(\sigma_1, \sigma_3) = d(\sigma_1, \sigma_4) = 2$, by Lemma 5.4 there exist

vertices δ_3 and δ_4 of $\mathcal{G}(K)$ such that $d(\sigma_1, \delta_i) = d(\delta_i, \sigma_i) = 1$, for $i \in \{3, 4\}$, and such that $d(\delta_3, \delta_4) \leq 1$. If furthermore δ_3 and δ_4 are distinct, then $(\sigma_1, \delta_3, \delta_4)$ is a 3-circuit and $(\sigma_1, \sigma_2, \sigma_3, \delta_3)$, $(\delta_3, \sigma_3, \sigma_4, \delta_4)$, and $(\sigma_1, \delta_4, \sigma_4, \sigma_5)$ are each circuits of length at most 4. Appealing to the previous two claims respectively, one may now find an appropriate compressing disc as the union of at most four other discs.

The remaining cases may be similarly treated. \square

This completes a proof of Proposition 1.5.

6. A genus 1 knot

The purpose of this section is to prove Proposition 1.4, that is to construct an atoroidal genus 1 knot K whose graph $\mathcal{G}(K)$ has diameter 2.

Let V_0 be a solid torus, and let A_1 and A_2 be annuli on ∂V_0 essential in V_0 such that $\partial A_1 \cap \partial A_2 = \partial A_1 = \partial A_2$ and such that the cyclic group $\pi_1(V_0)$ is not generated by the core of A_1 . Note, $\partial A_1 \cup \partial A_2 = \partial V_0$. Let V be a genus 2 handlebody obtained from V_0 by attaching a 1-handle $D^2 \times [1, 2]$, where $D^2 \times \{i\}$ is identified with a disc in $\text{int } A_i$ for both $i \in \{1, 2\}$. By assumption, the region in V bounded by A_1 and A_2 does not admit a product structure $A_1 \times [0, 1]$. After pushing $\text{int } A_i$ into $\text{int } V$, for both $i \in \{1, 2\}$, we have a pair of annuli properly embedded in V .

Let α, β be the two components of ∂A_1 , with orientation induced by either orientation of A_1 (see Fig. 1). We can choose A_1 and a band sum k of α and β such that k is *complicated* with respect to a preferred maximal meridian system $\{D_1, D_2, D_3\}$ for V , in the sense of Kobayashi [8]. That is, if R_1 and R_2 , both compact 3-holed spheres, denote the two components of $\partial V - \text{int } N(\partial D_1 \cup \partial D_2 \cup \partial D_3)$, then:

- There is no bigon B on ∂V such that ∂B is the union of a subarc of k and a subarc of ∂R_i for some $i \in \{1, 2\}$, and
- For any two boundary components of either 3-holed sphere R_i , there is a subarc of k joining them in R_i .

Appealing to Lemma 6.1 of [8], we have the following.

Lemma 6.1. *$\partial V - \text{int } N(k)$ is incompressible in V .*

Perhaps after an isotopy, we may assume k is disjoint from $\alpha \cup \beta$. Let θ denote any graph with two vertices, connected by three edges, embedded in \mathbb{S}^3 and whose exterior W admits a complete and finite volume hyperbolic metric with totally geodesic boundary. According to Section 3.3 of [17], one may, for instance, take θ to be the Kinoshita theta curve. Let $f: V \rightarrow N(\theta)$ be any homeomorphism, and define K equal to the image $f(k)$. Let $N(K)$ be a regular neighbourhood of K in \mathbb{S}^3 such that $N(K) \cap f(V)$ and $N(K) \cap W$ are regular neighbourhoods of K in $f(V)$ and W respectively. Note, the exterior $E = \mathbb{S}^3 - \text{int } N(K)$ of K is the union of $f(V) - \text{int } N(K)$ and $W - \text{int } N(K)$, with common subsurface $\partial f(V) - \text{int } N(K) = \partial W - \text{int } N(K)$. It should




Fig. 1. The curve k is a band sum of α and β , and is complicated with respect to the indicated maximal meridian system in the sense of Kobayashi.

also be noted that $f(V) - \text{int } N(K)$ is homeomorphic to $f(V)$ via a homeomorphism constant on $\partial f(V) - \text{int } N(K)$, and that $W - \text{int } N(K)$ is homeomorphic to W via a homeomorphism constant on $\partial W - \text{int } N(K)$.

Lemma 6.2. *The surface $\partial W - \text{int } N(K)$ is incompressible in W .*

Proof. Since K is an essential loop on ∂W so the natural inclusion $\partial W - \text{int } N(K) \rightarrow \partial W$ descends to an injection on fundamental groups. As W admits a hyperbolic metric in which ∂W is totally geodesic so ∂W is incompressible in W , and we find the natural homomorphism $\pi_1(\partial W - \text{int } N(K)) \rightarrow \pi_1(W)$ is also injective. It follows $\partial W - \text{int } N(K)$ is incompressible in W . \square

Lemma 6.3. *K is a non-trivial knot in \mathbb{S}^3 .*

Proof. According to Lemma 6.1, the group $\pi_1(\partial V - \text{int } N(K))$ naturally injects into $\pi_1(V)$. According to Lemma 6.2, the same group $\pi_1(\partial V - \text{int } N(K))$ naturally injects into $\pi_1(W)$. The knot group $\pi_1(E)$ is, by using van Kampen's theorem, therefore isomorphic to the amalgamated free product of $\pi_1(V)$ and $\pi_1(W)$ over a common subgroup isomorphic to the fundamental group of a 2-holed torus. Hence, $\pi_1(E)$ is a non-abelian group, and K cannot be a trivial knot. \square

Lemma 6.4. *The pair $(W, \partial W - \text{int } N(K))$ does not contain an essential annulus. That is, suppose A is an essential annulus properly embedded in W , with $\partial A \subset \partial W - \text{int } N(K)$. Then, A is parallel to an annulus in $\partial W - \text{int } N(K)$ or to the annulus $N(K) \cap \partial W$.*

Proof. The pair $(W, \partial W)$ cannot contain an essential annulus, for W admits a hyperbolic metric in which ∂W is totally geodesic. Let A be any incompressible annulus properly embedded in $W - \text{int } N(K)$ so that $\partial A \subset \partial W - \text{int } N(K)$. Then, A is parallel in W to an annulus $A' \subset \partial W$. If $A' \cap K$ is empty then A is also parallel to A' in $W - \text{int } N(K)$. If instead $A' \cap K$ is not empty, then $K \subset A'$ and, as K is essential on ∂W , so K is the core of A' . Thus, A is parallel to the annulus $N(K) \cap \partial W$. \square

Lemma 6.5. *K is an atoroidal knot in \mathbb{S}^3 .*

Proof. Suppose T is an incompressible torus in E . As both $f(V)$ and W are atoroidal, we may assume that T intersects $\partial W - \text{int } N(K)$ only in a collection of loops essential on ∂W and that each component of $T \cap f(V)$ and $T \cap W$ is an incompressible annulus in $f(V)$ and W , respectively.

Let A be a component of $T \cap W$, and consider the dichotomy contained in Lemma 6.4. If A is parallel to an annulus in $\partial W - \text{int } N(K)$, then we can decrease $|T \cap W|$ by an isotopy of T . We may thus assume every component of $T \cap W$ is an annulus parallel in $W - \text{int } N(K)$ to $\partial N(K) \cap W$. In which case, $T \cap \partial W$ consists of loops parallel to K in ∂W .

Now let A denote any component of $T \cap f(V)$. By the preceding argument, both components of ∂A are parallel to K in ∂W . Hence, A is parallel in the handlebody $f(V)$ to the annulus A' on $\partial f(V)$ bounded by ∂A . By the minimality of $|T \cap W|$, so A' necessarily contains K .

We conclude that T is the union of two annuli, one properly embedded in $f(V)$ and the other properly embedded in W and both parallel to A' . It follows that T is necessarily peripheral in E , and hence K is atoroidal. \square

The set $k \cup \alpha \cup \beta$ divides ∂V into a pair of 3-holed spheres, P_1 and P_2 . We now define S_{ij} to be equal to $f(P_i \cup A_j)$, for each $i, j \in \{1, 2\}$. Each is a genus 1 Seifert surface for K and, by Lemma 6.3, each is therefore of minimal genus. Reindexing if need be, we may assume S_{11} and S_{22} intersect transversely along α and along β . Let us abbreviate S_{ii} to S_i for both $i \in \{1, 2\}$. Then, $S_1 \cup S_2$ divides \mathbb{S}^3 into the following three regions:

- W , a hyperbolic 3-manifold;
- The solid torus $f(V_0)$, bounded by $f(A_1)$ and $f(A_2)$, and
- A third region that contains $S_1 \cap S_2$ and that is branched along $S_1 \cap S_2$. In particular, this region is not a 3-manifold.

None of these regions can give a product region between S_1 and S_2 . (Recall $f(V_0)$ does not give a product region between A_1 and A_2 .) It follows from the contrapositive of Proposition 4.8 (2) in [13] that S_1 and S_2 intersect essentially. In particular, $d([S_1], [S_2]) \geq 2$. According to Corollary 1.3, the diameter of $\mathcal{G}(K)$ is at most 2. We conclude that the diameter of $\mathcal{G}(K)$ is exactly 2. This completes a proof of Proposition 1.4.

7. An infinite class of atoroidal knots

The purpose of this section is to prove Proposition 1.6, offering a family of atoroidal knots, parameterised by knot genus, each of whose associated graphs has diameter precisely the modulus of the knot Euler characteristic. In particular, their diameters grow linearly with knot genus.

Given any non-negative integer g , pick a sequence of integers a_1, a_2, \dots, a_g of length g such that $|a_j| \geq 2$ for every j . Let K be the 2-bridge knot whose slope is represented by the continued fraction

$$[2a_1, -2a_2, \dots, 2a_{2g-1}, -2a_{2g}] = \cfrac{1}{2a_1 - \cfrac{1}{2a_2 + \cfrac{1}{2a_{2g-1} - \cfrac{1}{2a_{2g}}}}}$$

Then, the genus of K is precisely g . We show that the diameter of $\mathcal{G}(K)$ is equal to $2g - 1$ by using [13], where the structure of Kakimizu’s complex $\mathcal{MS}(K)$ is explicitly described. To recall, let T be a tree, with $n := 2g$ vertices, whose underlying space is homeomorphic to a closed interval, and let v_1, v_2, \dots, v_n be the vertices of T , lying on the interval in this order. For each vertex v_j we associate an unknotted oriented annulus $F(v_j)$ in S^3 with a_j -right hand full twists. Then, K is equal to the boundary of a surface obtained by successively plumbing the annuli $F(v_1), F(v_2), \dots, F(v_n)$, and this surface is a minimal genus Seifert surface for K . Moreover, every minimal genus Seifert surface of K is obtained in this way (see [2]).

There are 2^{n-1} different ways of successive plumbing, according as A_{j+1} is plumbed to A_j from above or from below with respect to a normal vector field on A_j . Thus, successive plumbing can be represented by an *orientation of T* , directing each edge in one of two ways, by the following rule: If ρ is an orientation of T , then we plumb A_{j+1} to A_j from above or below according as the edge joining v_j and v_{j+1} has initial point v_j or v_{j+1} , respectively, with respect to ρ . See Section 2 of [13] for a more detailed account.

We denote by $S(\rho)$ the Seifert surface of K determined by the orientation ρ . The condition that $|a_j| \geq 2$ for every j guarantees the correspondence $\rho \mapsto S(\rho)$ determines a bijection from the set $\mathcal{O}(T)$ of all orientations of T to the vertex set of $\mathcal{MS}(K)$ (see Theorem 2.3 of [13]).

To describe the structure of $\mathcal{MS}(K)$, we introduce a few definitions. A vertex v_j of T is said to be a *sink* for the orientation ρ of T if every edge of T incident on v_j points towards v_j . If v_j is a sink for ρ , then let $v_j(\rho)$ denote the orientation of T obtained from ρ by reversing the orientations of each edge incident on v_j . A *cycle* in $\mathcal{O}(T)$ is a sequence

$$\rho_1 \xrightarrow{v_{j_1}} \rho_2 \xrightarrow{v_{j_2}} \cdots \xrightarrow{v_{j_{n-1}}} \rho_n \xrightarrow{v_{j_n}} \rho_1,$$

where (j_1, j_2, \dots, j_n) is a permutation of $\{1, 2, \dots, n\}$ and $\rho_1, \rho_2, \dots, \rho_n$ are mutually distinct elements of $\mathcal{O}(T)$ such that $v_{j_k}(\rho_k) = \rho_{k+1}$ for every k , where our indices are considered modulo n . According to Theorem 3.3 of [13], $\mathcal{MS}(K)$ can be described as follows:

- The vertex set of $\mathcal{MS}(K)$ is identified with $\mathcal{O}(T)$, and
- A set of vertices $\{\rho_0, \rho_1, \dots, \rho_k\}$ spans a k -simplex in $\mathcal{MS}(K)$ if and only if it is contained in a cycle of $\mathcal{O}(T)$.

Moreover, $\mathcal{MS}(K)$ gives a triangulation of the cube I^{n-1} whose vertices are all the corners of the cube (see Proposition 3.9 of [13]).

We now show that the diameter of $\mathcal{G}(K)$ is equal to $n - 1$. Identify $\mathcal{O}(T)$ with $\{-, +\}^{(n-1)}$ by identifying $\rho \in \mathcal{O}(T)$ with $(\epsilon_1, \epsilon_2, \dots, \epsilon_{n-1})$, where ϵ_j is $+$ or $-$ according as the initial point of the j -th edge is v_j or v_{j+1} , respectively.

Lemma 7.1. *For any two elements ρ and ρ' of $\mathcal{O}(T)$, we have $d(\rho, \rho') \leq n - 1$, where d denotes the edge-path distance in $\mathcal{G}(K)$.*

Proof. We prove the lemma by inducting on n . Note that if n is odd, so not of the form $2g(K)$, we may still consider a linear tree T with n vertices and a simplicial complex with vertex set $\mathcal{O}(T)$. If $n = 1$, $\mathcal{O}(T)$ consists of a single element and the lemma obviously holds.

Let $\rho = (\epsilon_1, \epsilon_2, \dots, \epsilon_{n-1})$ and $\rho' = (\epsilon'_1, \epsilon'_2, \dots, \epsilon'_{n-1})$ be two elements of $\mathcal{O}(T)$, where T has n vertices. Suppose first that $\epsilon_{n-1} = \epsilon'_{n-1}$. Let T_0 be the sub-tree of T obtained by deleting the last edge. By the inductive hypothesis, the distance between $\rho_0 := (\epsilon_1, \epsilon_2, \dots, \epsilon_{n-2})$ and $\rho'_0 := (\epsilon'_1, \epsilon'_2, \dots, \epsilon'_{n-2})$ in $\mathcal{O}(T_0)$ is at most $n - 2$. Since every edge in $\mathcal{O}(T_0) = \{-, +\}^{(n-2)}$ lifts to an edge in $\{-, +\}^{(n-2)} \times \{\epsilon\} \subset \mathcal{O}(T)$, where $\epsilon = \epsilon_{n-1} = \epsilon'_{n-1}$, we see $d(\rho, \rho')$ is at most $n - 2$.

Suppose next that $\epsilon_{n-1} \neq \epsilon'_{n-1}$. Let ρ'' be the element of $\mathcal{O}(T)$ obtained from ρ' by replacing ϵ'_{n-1} with ϵ_{n-1} . Then, we have

$$d(\rho, \rho') \leq d(\rho, \rho'') + d(\rho'', \rho') \leq n - 2 + 1 = n - 1$$

and this completes a proof of Lemma 7.1. □

Lemma 7.2. *Let $\rho_- = (-, -, \dots, -)$ and $\rho_+ = (+, +, \dots, +)$. Then, $d(\rho_-, \rho_+) \geq n - 1$.*

Proof. Let $w(\rho)$ be the number of $+$ entries of $\rho \in \{-, +\}^{(n-1)}$, so that $w(\rho_-) = 0$ and $w(\rho_+) = n - 1$. The statement of the lemma follows once we prove $|w(\rho) - w(\rho')| \leq 1$ for any edge (ρ, ρ') of $\mathcal{G}(K)$. To prove this, observe that if v_j is a sink for ρ then $w(v_j(\rho))$ is equal to $w(\rho)$, $w(\rho) + 1$, or $w(\rho) - 1$ according as $j \in \{2, 3, \dots, n - 1\}$, $j = 1$ or $j = n$. Let (ρ, ρ') be an edge of $\mathcal{G}(K)$. Then, $\{\rho, \rho'\}$ is contained in the vertex set of a maximal simplex, of $\mathcal{MS}(K)$, which in turn is the set of all orientations for some cycle, say

$$\rho_1 \xrightarrow{v_{j_1}} \rho_2 \xrightarrow{v_{j_2}} \dots \xrightarrow{v_{j_{n-1}}} \rho_n \xrightarrow{v_{j_n}} \rho_1.$$

Since every vertex appears in a cycle, the above observation implies the set $\{w(\rho_1), w(\rho_2), \dots, w(\rho_n)\}$ consists of two successive integers. In particular $|w(\rho) - w(\rho')| \leq 1$, and this completes a proof of Lemma 7.2. \square

By Lemma 7.1 and Lemma 7.2 we see that the diameter of $\mathcal{G}(K)$ is equal to $n - 1$, thus completing a proof of Proposition 1.6.

RESEARCH UPDATE. Roberto Pelayo's thesis [12] became publicly available from April 2007. The upper bound given in Theorem 10.1 of [12] is quadratic in knot genus though is not computed. The argument found therein is based on minimal surface theory, and is quite different from that given here in Section 3.

In August 2007, Jennifer Schultens [15] gave an elegant proof of the simple connectivity of Kakimizu's complex, using PL-minimal surface theory. In fact her argument can be extended to prove that the second homotopy group of Kakimizu's complex is also trivial.

The Kakimizu conjecture remains open.

ACKNOWLEDGEMENTS. The authors wish to thank Osamu Kakimizu, for two particularly illuminating conversations, and to thank Tom Fleming, David Futer, François Guéritaud, Mikami Hirasawa, Kazuhiro Ichihara, Toshio Saito, and Yukihiro Tsutsumi for many interesting and helpful conversations. This project began life while the second author was visiting Osaka University. The second author wishes to thank Osaka University for its warm hospitality.

The first author is supported by JSPS Grants-in-Aid 18340018, and is partially supported by JSPS Core-to-Core Program 18005. The second author was partially supported by a short-term JSPS Post-Doctoral Fellowship, number PE05043, and by a long-term JSPS Post-Doctoral Fellowship, number P06034. Both authors wish to thank the JSPS for its financial support.

References

- [1] S.R. Fenley: *Quasi-Fuchsian Seifert surfaces*, Math. Z. **228** (1998), 221–227.
- [2] A.E. Hatcher and W.P. Thurston: *Incompressible surfaces in 2-bridge knot complements*, Invent. Math. **79** (1985), 225–246.
- [3] C. Hayashi: *The finiteness of the number of minimal Seifert surfaces up to homeomorphism*, Kobe J. Math. **10** (1993), 79–105.
- [4] M. Hirasawa and M. Sakuma: *Minimal genus Seifert surfaces for alternating links*; in Proceedings of Knots 96, 1997, 383–394.
- [5] O. Kakimizu: *Talk at the meeting “Knot theory and related topics” held in Osaka*, 1989.
- [6] O. Kakimizu: *Finding disjoint incompressible spanning surfaces for a link*, Hiroshima Math. J. **22** (1992), 225–236.
- [7] O. Kakimizu: *Classification of the incompressible spanning surfaces for prime knots of 10 or less crossings*, Hiroshima Math. J. **35** (2005), 47–92.
- [8] T. Kobayashi: *Casson-Gordon’s rectangle condition of Heegaard diagrams and incompressible tori in 3-manifolds*, Osaka J. Math. **25** (1988), 553–573.
- [9] T. Kobayashi: *Uniqueness of minimal genus Seifert surfaces for links*, Topology and its Appl. **33** (1989), 265–279.
- [10] U. Oertel: *On the existence of infinitely many essential surfaces of bounded genus*, Pacific J. Math. **202** (2002), 449–458.
- [11] R.C. Pelayo: *E-mail communication*, October 2006.
- [12] R.C. Pelayo: *Diameter bounds on the complex of minimal genus Seifert surfaces for hyperbolic knots*, Ph.D. thesis, April 2007, available online from the Caltech Library System at <http://resolver.caltech.edu/CaltechETD:etd-06042007-015951>.
- [13] M. Sakuma: *Minimal genus Seifert surfaces for special aborescent links*, Osaka J. Math. **31** (1994), 861–905.
- [14] M.G. Scharlemann and A.A. Thompson: *Finding disjoint Seifert surfaces*, Bull. London Math. Soc. **20** (1988), 61–64.
- [15] J.C. Schultens: *Covering spaces and the Kakimizu complex*, arXiv:0707.3926v3.
- [16] W.P. Thurston: *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. **59** (1986), i–vi, 99–130.
- [17] W.P. Thurston: *Three-Dimensional Geometry and Topology*, Princeton University Press, 1997.
- [18] Y. Tsutsumi: *Universal bounds for genus one Seifert surfaces for hyperbolic knots and surgeries with non-trivial JSJT-decompositions*; in Proceedings of the Winter Workshop of Topology/Workshop of Topology and Computer (Sendai, 2002/Nara, 2001), Interdiscip. Inform. Sci. **9**, 2003, 53–60.
- [19] Y. Tsutsumi: *Hyperbolic knots with a large number of disjoint minimal genus Seifert surfaces*, Tokyo J. Math. **31** (2008), 253–258.
- [20] F. Waldhausen: *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. **87** (1968), 56–88.
- [21] R.T. Wilson: *Knots with infinitely many incompressible Seifert surfaces*, arXiv:math.GT/0604001.

Makoto Sakuma
Department of Mathematics
Graduate School of Science
Osaka University
Toyonaka, Osaka 560-0043
Japan

Current address:
Department of Mathematics
Graduate School of Science
Hiroshima University
1-3-1 Kagamiyama
Higashi-Hiroshima
739-8526
Japan
e-mail: sakuma@math.sci.hiroshima-u.ac.jp
URL: <http://www.math.sci.hiroshima-u.ac.jp/~sakuma>

Kenneth J. Shackleton
Institute for the Physics and Mathematics of the Universe
The University of Tokyo
5-1-5 Kashiwanoha Kashiwa-shi
Chiba 277-8568
Japan
e-mail: kenneth.shackleton@ipmu.jp
e-mail: kjs2006@alumni.soton.ac.uk
URL: <http://member.ipmu.jp/kenneth.shackleton/>