

## LOCALLY CONFORMAL KÄHLER STRUCTURES ON COMPACT SOLVMANIFOLDS

HIROSHI SAWAI

(Received June 21, 2010, revised March 28, 2011)

### Abstract

Let  $(M, g, J)$  be a compact Hermitian manifold and  $\Omega$  the fundamental 2-form of  $(g, J)$ . A Hermitian manifold  $(M, g, J)$  is said to be locally conformal Kähler if there exists a closed 1-form  $\omega$  such that  $d\Omega = \omega \wedge \Omega$ . The purpose of this paper is to investigate a relation between a locally conformal Kähler structure and the adapted differential operator on compact solvmanifolds.

### Introduction

Let  $(M, g, J)$  be a  $2n$ -dimensional compact Hermitian manifold. We denote by  $\Omega$  the fundamental 2-form, that is, the 2-form defined by  $\Omega(X, Y) = g(X, JY)$ . A Hermitian manifold  $(M, g, J)$  is said to be *locally conformal Kähler* if there exists a closed 1-form  $\omega$  such that  $d\Omega = \omega \wedge \Omega$ . The closed 1-form  $\omega$  is called *Lee form*. In the case  $\omega = df$ ,  $(M, e^{-f}g, J)$  is Kähler. The main non-Kähler examples of locally conformal Kähler manifolds are Hopf manifolds [15], Inoue surfaces [14] and generalized Kodaira–Thurston manifolds [5] (cf. [6]). Note that Inoue surfaces and generalized Kodaira–Thurston manifolds have a structure of a compact solvmanifold. In this paper, we investigate locally conformal Kähler structures on a compact solvmanifold  $\Gamma \backslash G$  with a left-invariant complex structure, where  $G$  is a simply-connected solvable Lie group and  $\Gamma$  is a lattice of  $G$ , that is, a discrete co-compact subgroup.

An  $n$ -dimensional complex manifold  $M$  is said to be *complex parallelizable* if it admits holomorphic vector fields  $\{X_1, \dots, X_n\}$  which are linearly independent at every point. Wang [16] proved that a compact complex parallelizable manifold  $M$  is biholomorphic to a homogeneous space  $D \backslash G$ , where  $G$  is a simply connected complex Lie group and  $D$  is a discrete subgroup of  $G$ . Abbena–Grassi [1] proved that a non-toral compact complex parallelizable manifold has no locally conformal Kähler structures. The author in [12] proved that if a compact nilmanifold  $M$  with a left-invariant complex structure has a locally conformal Kähler structure, then  $M$  is biholomorphic to a Kodaira–Thurston manifold or a generalized Kodaira–Thurston manifold.

A locally conformal Kähler manifold  $(M, g, J)$  is said to be a *generalized Hopf manifold* if the Lee form  $\omega$  is parallel with respect to metric  $g$ . Vaisman [15] proved

that a generalized Hopf manifold has a structure of a principal  $S^1$ -bundle over a compact Sasakian manifold. Hopf manifolds and generalized Kodaira–Thurston manifolds are generalized Hopf manifolds. Tricerri [14] constructed a locally conformal Kähler structure on Inoue surfaces with non-parallel Lee form  $\omega$ . In Section 3, we prove that

**Main Theorem.** *Let  $(\Gamma \backslash G, g, J)$  be a locally conformal Kähler solvmanifold of  $\dim \Gamma \backslash G \geq 4$  with a left-invariant complex structure and  $\omega$  the Lee form. We assume that  $(\Gamma \backslash G, g, J)$  satisfies the following conditions:*

1. *There exists a left-invariant closed 1-form  $\omega_0$  such that the Lee form  $\omega$  is cohomologous to  $\omega_0$ .*
2. *The fundamental 2-form  $\Omega$  of  $(g, J)$  can be written as  $\Omega = -\omega \wedge \eta + d\eta$ , where  $\eta = -\omega \circ J$ .*

*Then  $\Gamma \backslash G$  has a left-invariant locally conformal Kähler structure with the parallel Lee form  $\omega_0$ , in particular,  $\Gamma \backslash G$  is a generalized Hopf manifold.*

In Section 4, we see that Kodaira–Thurston manifolds and secondary Kodaira–Thurston manifolds are examples of Main Theorem.

Hasegawa [7] proved that a complex structure on a 4-dimensional compact solvmanifold is induced from a left-invariant complex structure on the solvable Lie group and classified 4-dimensional compact solvmanifolds with a complex structure. By this classification, we have a classification of 4-dimensional locally conformal Kähler solvmanifolds in Section 4.

## 1. Preliminaries

Let  $(M = \Gamma \backslash G, g, J)$  be a compact locally conformal Kähler solvmanifold such that  $J$  is left-invariant. In this section, we see that the locally conformal Kähler metric  $g$  induces a left-invariant locally conformal Kähler metric and consider its fundamental 2-form on the Lie algebra  $\mathfrak{g}$  of  $G$ .

Let  $\Omega$  be the fundamental 2-form of  $(g, J)$ . There exists a closed 1-form  $\omega$  such that  $d\Omega = \omega \wedge \Omega$ . We now assume that the associated Lee form satisfies the condition 1 in Main Theorem. Then there exists a left-invariant closed 1-form  $\omega_0$  such that  $\omega_0 - \omega = df$ . This assumption holds for completely solvmanifolds [9]. Then we define a left-invariant 2-form  $\Omega_0$  by

$$\Omega_0(X, Y) = \int_M (e^f \Omega)(X, Y) d\mu$$

for left-invariant vector fields  $X, Y$ , where  $d\mu$  is the volume element induced by a bi-invariant volume element on  $G$ . Belgun [3] proved that  $\Omega_0$  is  $J$ -invariant and it is the fundamental 2-form of a Hermitian structure  $(\langle \cdot, \cdot \rangle, J)$  on the solvable Lie algebra  $\mathfrak{g}$  of  $G$  such that  $d\Omega_0 = \omega_0 \wedge \Omega_0$ .

DEFINITION 1.1. Let  $\mathfrak{g}$  be a Lie algebra and  $\alpha$  a closed 1-form on  $\mathfrak{g}$ . We define the adapted algebraic complex  $(\wedge \mathfrak{g}^*, d_\alpha)$  with the differential operator  $d_\alpha$ :

$$d_\alpha \beta := \alpha \wedge \beta + d\beta$$

for  $\beta \in \wedge^p \mathfrak{g}^*$ . Note that  $d_\alpha^2 = 0$  because  $\alpha$  is closed. A  $p$ -form  $\beta$  is called  $\alpha$ -closed if  $d_\alpha \beta = 0$ . It is called  $\alpha$ -exact if there exists a  $(p-1)$ -form  $\gamma$  such that  $\beta = d_\alpha \gamma$ .

On the above Hermitian structure  $(\langle \cdot, \cdot \rangle, J)$  on  $\mathfrak{g}$ , since  $\omega_0$  is a closed 1-form, the fundamental 2-form  $\Omega_0$  is  $-\omega_0$ -closed. We say that  $(\langle \cdot, \cdot \rangle, J)$  on  $\mathfrak{g}$  is *locally conformal Kähler*. In Section 4, we consider a relation between a property of Lee form and the adapted differential operator on 4-dimensional solvmanifolds.

We next show that  $\Omega_0$  is  $-\omega_0$ -exact if the second condition in Main Theorem also holds true.

**Proposition 1.2** (cf. [3]). *Let  $\Omega_0$  be the 2-form on  $\mathfrak{g}$  above. If  $\Omega = -\omega \wedge \eta + d\eta$ , then  $\Omega_0$  is  $-\omega_0$ -exact. In addition, if  $\eta = -\omega \circ J$ , then  $\eta_0 = -k\omega_0 \circ J$ , where  $k = \int_M e^f d\mu$ .*

Proof. Since  $\Omega = -\omega \wedge \eta + d\eta$  and  $\omega = \omega_0 - df$ , we see that

$$\begin{aligned} e^f \Omega &= e^f (-\omega \wedge \eta + d\eta) \\ &= e^f (-\omega_0 \wedge \eta + df \wedge \eta + d\eta) \\ &= -\omega_0 \wedge e^f \eta + d(e^f) \wedge \eta + e^f d\eta \\ &= -\omega_0 \wedge e^f \eta + d(e^f \eta). \end{aligned}$$

Then we get

$$\begin{aligned} \Omega_0(X, Y) &= \int_M (-\omega_0 \wedge e^f \eta + d(e^f \eta))(X, Y) d\mu \\ &= -\omega_0(X) \int_M (e^f \eta)(Y) d\mu + \omega_0(Y) \int_M (e^f \eta)(X) d\mu + \int_M d(e^f \eta)(X, Y) d\mu, \end{aligned}$$

for  $X, Y \in \mathfrak{g}$ .

Now, we define a 1-form  $\eta_0$  on  $\mathfrak{g}$  by  $\eta_0(X) = \int_M (e^f \eta)(X) d\mu$  for  $X \in \mathfrak{g}$ . Since  $d\mu$  is the right-invariant volume element, its Lie derivative  $L_X d\mu$  along a left-invariant vector fields  $X$  is zero. Then, for any function  $F$  on  $M$ , we see that

$$(XF) d\mu = (L_X F) d\mu = L_X(F d\mu) = di(X)F d\mu + i(X) d(F d\mu) = di(X)F d\mu,$$

where  $i(X)$  is the interior product with the vector field  $X$ . Therefore, by Stokes's theorem, we have

$$\begin{aligned} & \int_M d(e^f \eta)(X, Y) d\mu \\ &= \int_M X((e^f \eta)(Y)) d\mu - \int_M Y((e^f \eta)(X)) d\mu - \int_M (e^f \eta)([X, Y]) d\mu \\ &= \int_M di(X)(e^f \eta(Y)) d\mu - \int_M di(Y)(e^f \eta(X)) d\mu - \eta_0([X, Y]) \\ &= d\eta_0(X, Y). \end{aligned}$$

Thus we see that

$$\begin{aligned} \Omega_0(X, Y) &= -\omega_0(X) \int_M (e^f \eta)(Y) d\mu + \omega_0(Y) \int_M (e^f \eta)(X) d\mu + \int_M d(e^f \eta)(X, Y) d\mu \\ &= -\omega_0(X)\eta_0(Y) + \omega_0(Y)\eta_0(X) + d\eta_0(X, Y) \\ &= (-\omega_0 \wedge \eta_0 + d\eta_0)(X, Y). \end{aligned}$$

Similarly, if  $\eta = -\omega \circ J$ , then we have

$$\begin{aligned} \eta_0(X) &= \int_M (e^f \eta)(X) d\mu = - \int_M (e^f \omega)(JX) d\mu = - \int_M (e^f \omega_0 - e^f df)(JX) d\mu \\ &= -\omega_0(JX) \int_M e^f d\mu + \int_M d(e^f)(JX) d\mu \\ &= -\omega_0(JX) \int_M e^f d\mu + \int_M (JX)(e^f) d\mu \\ &= -\left(\int_M e^f d\mu\right)\omega_0(JX). \end{aligned}$$

Thus we see that  $\eta_0 = -k\omega_0 \circ J$ , where  $k = \int_M e^f d\mu$ . □

By Proposition 1.2, we see that  $\Omega_0 = -\omega_0 \wedge k\eta'_0 + k d\eta'_0 = k(-\omega_0 \wedge \eta'_0 + d\eta'_0) = kd_{-\omega_0}\eta'_0$ , where  $\eta'_0 = (1/k)\eta_0 = -\omega_0 \circ J$ . Then the locally conformal Kähler structure  $(\Omega_0, J)$  induces the locally conformal Kähler structure  $(\Omega'_0 = d_{-\omega_0}(-\omega_0 \circ J), J)$  on the solvable Lie algebra  $\mathfrak{g}$ . Therefore, by replacing  $\Omega_0$  by  $\Omega'_0$ , in order to prove Main Theorem, it is enough to show that if  $(\mathfrak{g}, \Omega_0 = d_{-\omega_0}(-\omega_0 \circ J), J)$  is a locally conformal Kähler solvable Lie algebra, then the Lee form  $\omega_0$  is parallel with respect to  $\langle , \rangle$  (see Section 3).

### 2. The adjoint operators and the inner product

Let  $(\langle , \rangle, J)$  be the locally conformal Kähler structure as we consider in Section 1 on solvable Lie algebra  $\mathfrak{g}$ , namely  $\Omega_0$  can be written as  $\Omega_0 = d_{-\omega_0}\eta_0 = -\omega_0 \wedge \eta_0 + d\eta_0$ ,

where  $\omega_0$  is a closed 1-form on  $\mathfrak{g}$  and  $\eta_0 = -\omega_0 \circ J$ . In this section, we investigate the properties of  $(\mathfrak{g}, \langle \cdot, \cdot \rangle, J)$ .

Let  $\gamma$  be the canonical isomorphism from  $\mathfrak{g}^*$  to  $\mathfrak{g}$  induced by the inner product  $\langle \cdot, \cdot \rangle$ . Put  $A = \gamma(\omega_0)$ . By this normalization, we may assume that  $\langle A, A \rangle = 1$ .

We easily see that an abelian Lie algebra of dimension equal to or more than 4 has no locally conformal Kähler structures. From now on, we assume that  $\mathfrak{g}$  is not abelian.

Since  $\mathfrak{g}$  is solvable,  $[\mathfrak{g}, \mathfrak{g}]$  is nilpotent. We take the descending central series for  $[\mathfrak{g}, \mathfrak{g}]$ :  $\mathfrak{n} = [\mathfrak{g}, \mathfrak{g}] \supset \mathfrak{n}^{(1)} = [\mathfrak{n}, \mathfrak{n}] \supset \mathfrak{n}^{(2)} = [\mathfrak{n}, \mathfrak{n}^{(1)}] \supset \dots \supset \mathfrak{n}^{(r)} \supset \mathfrak{n}^{(r+1)} = 0$ , where  $\mathfrak{n}^{(i+1)} = [\mathfrak{n}, \mathfrak{n}^{(i)}]$  ( $i \geq 1$ ) and  $\mathfrak{n}^{(r)} \neq 0$ . We easily see that  $\mathfrak{n}^{(r)}$  is contained in the center  $Z(\mathfrak{n})$  of  $\mathfrak{n}$ . Note that, for  $X \in \mathfrak{g}$ ,  $\text{ad}(X)\mathfrak{n}^{(i)} \subset \mathfrak{n}^{(i)}$  for each  $i$ . Then we get

**Lemma 2.1.**  $J\mathfrak{n}^{(r)} \subset [\mathfrak{g}, \mathfrak{g}]^\perp$ , where  $[\mathfrak{g}, \mathfrak{g}]^\perp$  is the orthogonal component of  $[\mathfrak{g}, \mathfrak{g}]$ .

Proof. For  $Z \in \mathfrak{n}^{(r)}$  and  $X \in [\mathfrak{g}, \mathfrak{g}]$ , we see that

$$\langle JZ, X \rangle = \Omega_0(X, Z) = (-\omega_0 \wedge \eta_0 + d\eta_0)(X, Z) = 0,$$

because  $\omega_0$  is closed and  $Z \in Z(\mathfrak{n})$ . □

**Lemma 2.2.** For  $Z, Z' \in \mathfrak{n}^{(r)}$ ,  $[JZ, Z'] = [JZ', Z]$  and  $[JZ, JZ'] = 0$ .

Proof. The Nijenhuis tensor  $\mathbf{N}_J$  vanishes, because  $J$  is integrable. Since  $Z, Z' \in \mathfrak{n}^{(r)}$ , we see that

$$\begin{aligned} 0 &= \mathbf{N}_J(Z, Z') = [Z, Z'] + J[JZ, Z'] + J[Z, JZ'] - [JZ, JZ'] \\ &= J\{[JZ, Z'] + [Z, JZ']\} - [JZ, JZ']. \end{aligned}$$

Note that  $[JZ, Z'], [Z, JZ'] \in \mathfrak{n}^{(r)}$ . It follows that  $J\{[JZ, Z'] + [Z, JZ']\} \in [\mathfrak{g}, \mathfrak{g}]^\perp$  by Lemma 2.1. Thus we have  $[JZ, Z'] + [Z, JZ'] = 0$  and  $[JZ, JZ'] = 0$ . □

By Lemmas 2.1 and 2.2, we have

**Proposition 2.3.** For  $U, V, W \in \mathfrak{n}^{(r)}$ ,

$$\langle \text{ad}(JU)V, W \rangle + \omega_0(JV)\langle U, W \rangle = \langle V, \text{ad}(JU)W \rangle + \omega_0(JW)\langle U, V \rangle.$$

Proof. Since  $\Omega_0 = -\omega_0 \wedge \eta_0 + d\eta_0$  and  $\omega_0$  is closed 1-form, we see that

$$\begin{aligned} \langle \text{ad}(JU)V, W \rangle &= \Omega_0(J \text{ad}(JU)V, W) = -\Omega_0(\text{ad}(JU)V, JW) \\ (2.1) \quad &= (\omega_0 \wedge \eta_0 - d\eta_0)(\text{ad}(JU)V, JW) \\ &= -\omega_0(JW)\eta_0(\text{ad}(JU)V) - d\eta_0(\text{ad}(JU)V, JW). \end{aligned}$$

Since  $\eta_0 = -\omega_0 \circ J$ , we get

$$\begin{aligned}
 (2.2) \quad -\omega_0(JW)\eta_0(\text{ad}(JU)V) &= \omega_0(JW) d\eta_0(JU, V) \\
 &= \omega_0(JW)\{\Omega_0(JU, V) + \omega_0 \wedge \eta_0(JU, V)\} \\
 &= \omega_0(JW)\langle U, V \rangle - \omega_0(JW)\omega_0(JU)\omega_0(JV).
 \end{aligned}$$

From the derivation conditions and Lemma 2.2, we get

$$\begin{aligned}
 d\eta_0(\text{ad}(JU)V, JW) &= d\eta_0(JU, [V, JW]) - d\eta_0(V, [JU, JW]) \\
 &= d\eta_0(JU, [V, JW]) \\
 &= d\eta_0(JU, [W, JV]) \\
 &= d\eta_0([JU, W], JV) + d\eta_0(W, [JU, JV]) \\
 &= d\eta_0([JU, W], JV) = -d\eta_0(JV, [JU, W]).
 \end{aligned}$$

It follows that

$$\begin{aligned}
 (2.3) \quad &-d\eta_0(\text{ad}(JU)V, JW) \\
 &= d\eta_0(JV, [JU, W]) \\
 &= \Omega_0(JV, [JU, W]) + \omega_0 \wedge \eta_0(JV, [JU, W]) \\
 &= \langle V, [JU, W] \rangle - \omega_0(JV) d\eta_0(JU, W) \\
 &= \langle V, [JU, W] \rangle - \omega_0(JV)\{\Omega_0(JU, W) + \omega_0 \wedge \eta_0(JU, W)\} \\
 &= \langle V, [JU, W] \rangle - \omega_0(JV)\langle U, W \rangle + \omega_0(JV)\omega_0(JU)\omega_0(JW).
 \end{aligned}$$

From (2.1) and (2.2)), (2.3), we have

$$\begin{aligned}
 \langle \text{ad}(JU)V, W \rangle &= \omega_0(JW)\langle U, V \rangle - \omega_0(JW)\omega_0(JU)\omega_0(JV) \\
 &\quad + \langle V, [JU, W] \rangle - \omega_0(JV)\langle U, W \rangle + \omega_0(JV)\omega_0(JU)\omega_0(JW), \\
 \langle \text{ad}(JU)V, W \rangle + \omega_0(JV)\langle U, W \rangle &= \langle V, \text{ad}(JU)W \rangle + \omega_0(JW)\langle U, V \rangle. \quad \square
 \end{aligned}$$

It is well-known that if a solvable Lie group  $G$  admits a lattice  $\Gamma$ , then the solvable Lie algebra  $\mathfrak{g}$  of  $G$  is unimodular. We define

**DEFINITION 2.4.** A solvable Lie algebra  $\mathfrak{g}$  is called strongly unimodular if, for  $X \in \mathfrak{g}$ ,  $\text{tr ad}(X)|_{\mathfrak{n}^{(i)}} = 0$  for each  $i$ .

Benson–Gordon [4] proved that if a solvable Lie group  $G$  admits a lattice  $\Gamma$ , then the solvable Lie algebra  $\mathfrak{g}$  of  $G$  is strongly unimodular.

We take an orthonormal frame  $\{Z_1, \dots, Z_m\}$  of  $\mathfrak{n}^{(r)}$  and consider the strongly unimodular conditions of  $\text{ad}(JZ_i)$  from  $\mathfrak{n}^{(r)}$  to  $\mathfrak{n}^{(r)}$  for each  $i$ .

For each  $i$ , let

$$\text{ad}(JZ_i)(Z_1, \dots, Z_m) = (Z_1, \dots, Z_m) \begin{pmatrix} a_{11}(i) & \cdots & a_{1m}(i) \\ \vdots & \vdots & \vdots \\ a_{m1}(i) & \cdots & a_{mm}(i) \end{pmatrix}.$$

By Proposition 2.3, we see that

$$\begin{aligned} \langle \text{ad}(JZ_i)Z_j, Z_k \rangle + \omega_0(JZ_j)\langle Z_i, Z_k \rangle &= \langle Z_j, \text{ad}(JZ_i)Z_k \rangle + \omega_0(JZ_k)\langle Z_i, Z_j \rangle, \\ a_{kj}(i) + \delta_{ik}\omega_0(JZ_j) &= a_{jk}(i) + \delta_{ij}\omega_0(JZ_k). \end{aligned}$$

It follows that, in the case of  $j, k \neq i$ ,

$$a_{kj}(i) = a_{jk}(i)$$

and, in the case of  $j = i, k \neq i$ ,

$$a_{ki}(i) = a_{ik}(i) + \omega_0(JZ_k).$$

Then we get

$$(2.4) \quad (a_{jk}(i)) = A_i + \begin{pmatrix} \text{line } i & & \\ \omega_0(JZ_1) & & \\ \vdots & & \\ \omega_0(JZ_m) & & \mathbf{0} \end{pmatrix},$$

where  $A_i$  is an  $(m \times m)$ -symmetric matrix.

Moreover, put

$$A_i = \text{row } i \begin{pmatrix} & \text{line } i & \\ & a_1^i & \\ * & \vdots & * \\ a_1^i & \cdots & a_i^i & \cdots & a_m^i \\ & \vdots & & & \\ * & a_m^i & & & * \end{pmatrix},$$

for each  $i$ . Since  $[JZ_i, Z_j] = [JZ_j, Z_i]$ , we get

$$a_{jj}(i) = \langle \text{ad}(JZ_i)Z_j, Z_j \rangle = \langle \text{ad}(JZ_j)Z_i, Z_j \rangle = a_i^j$$

for  $j \neq i$ . Then  $A_i$  can be written as

$$A_i = \text{row } i \begin{pmatrix} & \text{line } i & & & \\ a_i^1 & a_1^i & & * & \\ & \ddots & \vdots & & \\ a_1^i & \cdots & a_i^i & \cdots & a_m^i \\ & & \vdots & \ddots & \\ & * & a_m^i & & a_i^m \end{pmatrix}.$$

Thus (2.4) can be expressed as follows:

$$(2.5) \quad (a_{jk}(i)) = \text{row } i \begin{pmatrix} & \text{line } i & & & \\ a_i^1 & a_1^i + \omega_0(JZ_1) & & & \\ & * & & \vdots & * \\ & \ddots & & \vdots & \\ a_1^i * & \cdots & a_i^i + \omega_0(JZ_i) & \cdots & a_m^i \\ & & \vdots & \ddots & * \\ & * & a_m^i + \omega_0(JZ_m) & & a_i^m \end{pmatrix}.$$

From the strongly unimodular condition of  $\text{ad}(JZ_i)|_{\mathfrak{n}^\varphi}$ , we have

**Proposition 2.5.** For each  $i$ ,  $(\sum_j a_i^j) + \omega_0(JZ_i) = 0$ .

Then we have

**Corollary 2.6.**  $\sum_i \omega_0(JZ_i)^2 = 1$ .

Proof. For each  $i$ , we see that

$$\begin{aligned} 1 &= \langle Z_i, Z_i \rangle = \Omega_0(JZ_i, Z_i) \\ &= (-\omega_0 \wedge \eta_0 + d\eta_0)(JZ_i, Z_i) \\ &= -\omega_0(JZ_i)\eta_0(Z_i) - \eta_0([JZ_i, Z_i]) \\ &= \omega_0(JZ_i)^2 + \omega_0(J[JZ_i, Z_i]), \end{aligned}$$



because  $\eta_0 = -\omega_0 \circ J$ . From (2.5),

$$\begin{aligned}
 1 &= \langle Z_i, Z_i \rangle = \omega_0(JZ_i)^2 + \omega_0(J[JZ_i, Z_i]), \\
 (2.6) \quad 1 &= \omega_0(JZ_i)^2 + \omega_0 \circ J \left( \sum_j (a_j^i + \omega_0(JZ_j))Z_j \right), \\
 1 &= \omega_0(JZ_i)^2 + \sum_j a_j^i \omega_0(JZ_j) + \sum_j \omega_0(JZ_j)^2.
 \end{aligned}$$

Take the sum of (2.6) for  $i$ , we get

$$\begin{aligned}
 \sum_i 1 &= \sum_i \omega_0(JZ_i)^2 + \sum_{i,j} a_j^i \omega_0(JZ_j) + \sum_{i,j} \omega_0(JZ_j)^2, \\
 m &= \sum_i \omega_0(JZ_i)^2 + \sum_i \left( \sum_j a_j^i \right) \omega_0(JZ_i) + m \sum_i \omega_0(JZ_i)^2.
 \end{aligned}$$

By Proposition 2.5, we have

$$\begin{aligned}
 m &= \sum_i \omega_0(JZ_i)^2 + \sum_i (-\omega_0(JZ_i))\omega_0(JZ_i) + m \sum_i \omega_0(JZ_i)^2, \\
 m &= \sum_i \omega_0(JZ_i)^2 - \sum_i \omega_0(JZ_i)^2 + m \sum_i \omega_0(JZ_i)^2, \\
 m &= m \sum_i \omega_0(JZ_i)^2,
 \end{aligned}$$

which implies that  $\sum_i \omega_0(JZ_i)^2 = 1$ . □

**Corollary 2.7.**  $A = \sum_i \omega_0(JZ_i)JZ_i \in J\mathfrak{n}^{(r)}$ .

Proof. Since  $A = \gamma(\omega_0)$ , it can be given by

$$A = \sum_i \omega_0(JZ_i)JZ_i + B,$$

where  $B \in (J\mathfrak{n}^{(r)})^\perp \cap [\mathfrak{g}, \mathfrak{g}]^\perp$ . From  $\langle A, A \rangle = 1$ , we have

$$1 = \langle A, A \rangle = \sum_i \omega_0(JZ_i)^2 \langle JZ_i, JZ_i \rangle + \langle B, B \rangle = \sum_i \omega_0(JZ_i)^2 + \langle B, B \rangle.$$

By Corollary 2.6, we have  $\langle B, B \rangle = 0$ , which implies that  $B = 0$ . □

### 3. Structure on the solvable Lie algebra

We use same notation introduced in Section 2. In this section, we prove that  $JA$  is in the center  $Z(\mathfrak{g})$  of  $\mathfrak{g}$  and have Main Theorem.

We see  $JA = -\sum_i \omega_0(JZ_i)Z_i \in \mathfrak{n}^{(r)}$  by Corollary 2.7. Put  $Z_0 = JA$ . Note that  $\eta_0(Z_0) = -\omega_0 \circ J(JA) = 1$  and  $\eta_0(A) = -\omega_0 \circ J(A) = 0$ . We get

**Lemma 3.1.** *For  $X \in \mathfrak{g}$ ,  $d\eta_0(A, X) = 0$  and  $d\eta_0(Z_0, X) = 0$ .*

*Proof.* Since  $\eta_0 = -\omega_0 \circ J$ , we see that

$$\begin{aligned} d\eta_0(A, X) &= \Omega_0(A, X) + \omega_0 \wedge \eta_0(A, X) = \langle A, JX \rangle + \eta_0(X) = \omega_0(JX) + \eta_0(X) \\ &= 0. \end{aligned}$$

Similarly, we get

$$\begin{aligned} d\eta_0(Z_0, X) &= \Omega_0(Z_0, X) + \omega_0 \wedge \eta_0(Z_0, X) = -\langle X, JZ_0 \rangle - \omega_0(X) = \langle X, A \rangle - \omega_0(X) \\ &= 0. \end{aligned} \quad \square$$

Then we see that

**Proposition 3.2.** *For  $U \in \mathfrak{n}^{(r)}$ ,  $\text{ad}(A)U = 0$ .*

*Proof.* By a straightforward computation, we see that

$$\begin{aligned} \langle \text{ad}(A)U, \text{ad}(A)U \rangle &= \Omega_0(J \circ \text{ad}(A)U, \text{ad}(A)U) \\ &= (-\omega_0 \wedge \eta_0 + d\eta_0)(J \circ \text{ad}(A)U, \text{ad}(A)U) \\ &= -\omega_0(J \circ \text{ad}(A)U)\eta_0(\text{ad}(A)U) + d\eta_0(J \circ \text{ad}(A)U, \text{ad}(A)U). \end{aligned}$$

By Lemma 3.1 and the derivation conditions,

$$\begin{aligned} \langle \text{ad}(A)U, \text{ad}(A)U \rangle &= \omega_0(J \circ \text{ad}(A)U) d\eta_0(A, U) - d\eta_0(\text{ad}(A)U, J \circ \text{ad}(A)U) \\ &= -d\eta_0(\text{ad}(A)U, J \circ \text{ad}(A)U) \\ &= -d\eta_0(A, [U, J \circ \text{ad}(A)U]) + d\eta_0(U, [A, J \circ \text{ad}(A)U]) \\ &= d\eta_0(U, [A, J \circ \text{ad}(A)U]). \end{aligned}$$

Now, since  $JA = Z_0$  and  $\text{ad}(A)U \in \mathfrak{n}^{(r)}$ , we see that

$$[A, J \circ \text{ad}(A)U] = [-JZ_0, J \circ \text{ad}(A)U] = 0$$

by Lemma 2.2. It follows that  $d\eta_0(U, [A, J \circ \text{ad}(A)U]) = 0$ , which implies that  $\text{ad}(A)U = 0$ . □

Therefore we have

**Theorem 3.3.**  $Z_0 \in Z(\mathfrak{g})$ .

Proof. Let  $X \in \mathfrak{g}$ . By Lemma 3.1, we see that

$$\begin{aligned} \langle \text{ad}(X)Z_0, \text{ad}(X)Z_0 \rangle &= \Omega_0(J \circ \text{ad}(X)Z_0, \text{ad}(X)Z_0) \\ &= (-\omega_0 \wedge \eta_0 + d\eta_0)(J \circ \text{ad}(X)Z_0, \text{ad}(X)Z_0) \\ &= -\omega_0(J \circ \text{ad}(X)Z_0)\eta_0(\text{ad}(X)Z_0) + d\eta_0(J \circ \text{ad}(X)Z_0, \text{ad}(X)Z_0) \\ &= \omega_0(J \circ \text{ad}(X)Z_0) d\eta_0(X, Z_0) + d\eta_0(J \circ \text{ad}(X)Z_0, \text{ad}(X)Z_0) \\ &= d\eta_0(J \circ \text{ad}(X)Z_0, \text{ad}(X)Z_0). \end{aligned}$$

Moreover, from the derivation conditions and  $Z_0 \in Z([\mathfrak{g}, \mathfrak{g}])$ ,

$$\begin{aligned} \langle \text{ad}(X)Z_0, \text{ad}(X)Z_0 \rangle &= d\eta_0(J \circ \text{ad}(X)Z_0, \text{ad}(X)Z_0) = -d\eta_0(\text{ad}(X)Z_0, J \circ \text{ad}(X)Z_0) \\ &= -d\eta_0(X, [Z_0, J \circ \text{ad}(X)Z_0]) + d\eta_0(Z_0, [X, J \circ \text{ad}(X)Z_0]) \\ &= -d\eta_0(X, [Z_0, J \circ \text{ad}(X)Z_0]). \end{aligned}$$

Now, since  $\text{ad}(X)Z_0 \in \mathfrak{n}^{(r)}$ , we see that

$$[Z_0, J \circ \text{ad}(X)Z_0] = [\text{ad}(X)Z_0, JZ_0] = [A, \text{ad}(X)Z_0] = 0$$

by Lemma 2.2 and Proposition 3.2. It follows that  $d\eta_0(X, [Z_0, J \circ \text{ad}(X)Z_0]) = 0$ , which implies that  $\text{ad}(X)Z_0 = 0$  for  $X \in \mathfrak{g}$ .  $\square$

From Theorem 3.3, we have

**Corollary 3.4.**  $\text{ad}(A) \circ J = J \circ \text{ad}(A)$ .

Proof. Let  $X \in \mathfrak{g}$ . Since  $J$  is integrable, we see that

$$\begin{aligned} 0 &= \mathbf{N}_J(A, X) = [A, X] + J[JA, X] + J[A, JX] - [JA, JX] \\ 0 &= [A, X] + J[Z_0, X] + J[A, JX] - [Z_0, JX]. \end{aligned}$$

From  $Z_0 \in Z(\mathfrak{g})$ ,  $[A, X] + J[A, JX] = 0$ . Then we have our claim.  $\square$

**Corollary 3.5.** *Lee form  $\omega_0$  is parallel, if and only if  $\Omega_0$  is  $\text{ad}(A)$ -invariant.*

Proof. Let  $\nabla$  be the Riemannian connection of  $(\cdot, \cdot)$ . For  $X, Y \in \mathfrak{g}$ , we see that

$$\begin{aligned} 2(\nabla_X \omega_0)(Y) &= -2\omega_0(\nabla_X Y) = -2\langle A, \nabla_X Y \rangle \\ &= -\langle [A, X], Y \rangle - \langle X, [A, Y] \rangle - \langle A, [X, Y] \rangle \\ &= -\Omega_0(J[A, X], Y) - \Omega_0(JX, [A, Y]) - \omega_0([X, Y]) \\ &= -\Omega_0(J[A, X], Y) - \Omega_0(JX, [A, Y]), \end{aligned}$$

because  $\omega_0$  is a closed 1-form. From Corollary 3.4, we get

$$2(\nabla_X \omega_0)(Y) = -\Omega_0([A, JX], Y) - \Omega_0(JX, [A, Y]) = (\text{ad}^*(A)\Omega_0)(JX, Y).$$

Thus we have our claim.  $\square$

**Proof of Main Theorem.** By Corollary 3.5, to prove Main Theorem, it is enough to show that  $\Omega_0$  is  $\text{ad}(A)$ -invariant.

Let  $X, Y \in \mathfrak{g}$ . By a straightforward computation, we see that

$$\begin{aligned} (\text{ad}^*(A)\Omega_0)(X, Y) &= -\Omega_0([A, X], Y) - \Omega_0(X, [A, Y]) \\ &= (\omega_0 \wedge \eta_0 - d\eta_0)([A, X], Y) + (\omega_0 \wedge \eta_0 - d\eta_0)(X, [A, Y]) \\ &= -\omega_0(Y)\eta_0([A, X]) - d\eta_0([A, X], Y) \\ &\quad + \omega_0(X)\eta_0([A, X]) - d\eta_0(X, [A, Y]). \end{aligned}$$

By Lemma 3.1, we get  $\eta_0([A, X]) = -d\eta_0(A, X) = 0$  and  $\eta_0([A, Y]) = 0$ . Moreover, from the derivation conditions,

$$d\eta_0([A, X], Y) + d\eta_0(X, [A, Y]) = d\eta_0(A, [X, Y]) = 0,$$

which implies that  $\Omega_0$  is  $\text{ad}(A)$ -invariant. This completes the proof of Main Theorem.  $\square$

#### 4. Examples

In this section, we give examples of Main Theorem and consider locally conformal Kähler structures on 4-dimensional compact solvmanifolds. Note that a compact Kähler solvmanifold is a finite quotient of a complex torus which has a structure of a complex torus bundle over a complex torus ([2], [8]).

Hasegawa [7] classified a 4-dimensional compact solvmanifold and proved that any complex structure on such solvmanifold is induced from a left-invariant complex structure on Lie group. By this classification, we see that a 4-dimensional locally conformal Kähler solvmanifold is biholomorphic to Kodaira–Thurston manifold, Secondary Kodaira–Thurston manifold or Inoue surfaces.

We see that Kodaira–Thurston manifolds and secondary Kodaira–Thurston manifolds are examples of Main Theorem:

EXAMPLE 4.1 (Kodaira–Thurston manifold [5]). Let  $G$  be a 3-dimensional nilpotent Lie group given by

$$G = \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} : x, y, z \in \mathbb{R} \right\}.$$

The Lie group  $G$  admits a lattice  $\Gamma$ . Let  $\mathfrak{g}$  be the nilpotent Lie algebra corresponding to  $G$ .  $\mathfrak{g} \times \mathbb{R}$  is given by

$$\mathfrak{g} \times \mathbb{R} = \text{span}\{X, Y, Z, A : [X, Y] = Z\}.$$

Let  $\{x, y, z, \omega\}$  be the dual base of  $\{X, Y, Z, A\}$ :

$$dx = dy = d\omega = 0, \quad dz = -x \wedge y.$$

We define a left-invariant metric  $\langle \cdot, \cdot \rangle$  on  $\Gamma \backslash G \times S^1$  such that  $\{X, Y, Z, A\}$  is an orthonormal frame and a left-invariant complex structure  $J$  by  $JA = Z, JX = Y$ . Then  $(\Gamma \backslash G \times S^1, \langle \cdot, \cdot \rangle, J)$  is a locally conformal Kähler manifold with the fundamental 2-form given by

$$\Omega_0 = -\omega \wedge z - x \wedge y = d_{-\omega}z = d_{-\omega}(-\omega \circ J).$$

Hence the fundamental 2-form  $\Omega_0$  is  $-\omega$ -exact. We easily see that Lee form  $\omega$  is parallel.

EXAMPLE 4.2 (Secondary Kodaira–Thurston manifold (cf. [15])). Let  $G$  be a 4-dimensional solvable Lie group given by

$$G = \left\{ \begin{pmatrix} 1 & -\frac{1}{2}(x \sin t + y \cos t) & \frac{1}{2}(x \cos t - y \sin t) & z \\ 0 & \cos t & \sin t & x \\ 0 & -\sin t & \cos t & y \\ 0 & 0 & 0 & 1 \end{pmatrix} : t, x, y, z \in \mathbb{R} \right\}.$$

The Lie group  $G$  admits a lattice  $\Gamma$ :

$$\Gamma = \left\{ \begin{pmatrix} 1 & -\frac{1}{2}(u \sin 2\pi s + v \cos 2\pi s) & \frac{1}{2}(u \cos 2\pi s - v \sin 2\pi s) & w \\ 0 & \cos 2\pi s & \sin 2\pi s & u \\ 0 & -\sin 2\pi s & \cos 2\pi s & v \\ 0 & 0 & 0 & 1 \end{pmatrix} : s, u, v, w \in \mathbb{Z} \right\}.$$

Let  $\mathfrak{g}$  be the solvable Lie algebra corresponding to  $G$ :

$$\mathfrak{g} = \text{span}\{A, X, Y, Z : [A, X] = Y, [A, Y] = -X, [X, Y] = Z\}.$$

Let  $\{\omega, x, y, z\}$  be the dual base of  $\{A, X, Y, Z\}$ :

$$d\omega = 0, \quad dx = -\omega \wedge y, \quad dy = \omega \wedge x, \quad dz = -x \wedge y.$$

We define a left-invariant metric  $\langle \cdot, \cdot \rangle$  on  $\Gamma \backslash G$  such that  $\{A, X, Y, Z\}$  is an orthonormal frame and a left-invariant complex structure  $J$  by  $JA = Z, JX = Y$ . Then  $(\Gamma \backslash G, \langle \cdot, \cdot \rangle, J)$  is a locally conformal Kähler manifold with the fundamental 2-form given by

$$\Omega_0 = -\omega \wedge z - x \wedge y = d_{-\omega}z = d_{-\omega}(-\omega \circ J).$$

Hence the fundamental 2-form  $\Omega_0$  is  $-\omega$ -exact. We easily see that Lee form  $\omega$  is parallel.

A locally conformal Kähler structure on Inoue surfaces are different from one on solvmanifolds as Main Theorem:

EXAMPLE 4.3 (Inoue surface  $S^0$  [11], [14]). Let  $B \in \text{SL}(3, \mathbb{Z})$  be a unimodular matrix with eigenvalues  $\alpha, \beta, \bar{\beta}$  such that  $\beta \neq \bar{\beta}$ , and eigenvectors  $(a_1, a_2, a_3), (b_1, b_2, b_3)$  of  $\alpha, \beta$ , respectively. Then we define a structure of the group on  $\mathbb{H} \times \mathbb{C} = \{(x + \sqrt{-1}\alpha^t, z) : x, t \in \mathbb{R}, z \in \mathbb{C}\}$  as follows:

$$(x + \sqrt{-1}\alpha^t, z) \cdot (x' + \sqrt{-1}\alpha^{t'}, z') = (\alpha^t x' + x + \sqrt{-1}\alpha^{t+t'}, \beta^t z' + z).$$

It can be expressed by

$$G = \left\{ \left( \begin{array}{cccc} \alpha^t & 0 & 0 & x \\ 0 & \beta^t & 0 & z \\ 0 & 0 & \bar{\beta}^t & \bar{z} \\ 0 & 0 & 0 & 1 \end{array} \right) : t, x \in \mathbb{R}, z \in \mathbb{C} \right\}.$$

Thus we easily see that the Lie group  $G$  is solvable and it admits a lattice  $\Gamma$ :

$$\Gamma = \left\{ \left( \begin{array}{cccc} \alpha^s & 0 & 0 & P \begin{pmatrix} u \\ w_1 \\ w_2 \end{pmatrix} \\ 0 & \beta^s & 0 & \\ 0 & 0 & \bar{\beta}^s & \\ 0 & 0 & 0 & 1 \end{array} \right) : s, u, w_1, w_2 \in \mathbb{Z} \right\},$$

where  $P = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ \bar{b}_1 & \bar{b}_2 & \bar{b}_3 \end{pmatrix} \in \text{GL}(3, \mathbb{C})$ . Let  $\mathfrak{g}$  be the solvable Lie algebra corresponding to  $G$ :

$$\mathfrak{g} = \text{span}\{A, X, Y_1, Y_2 : [A, X] = -2rX, [A, Y_1] = rY_1 + \theta Y_2, [A, Y_2] = rY_2 - \theta Y_1\},$$

where  $\alpha = e^{-2r}$  and  $\beta = e^{r+\sqrt{-1}\theta}$ . Let  $\{\omega, x, y_1, y_2\}$  be the dual base of  $\{A, X, Y_1, Y_2\}$ :

$$d\omega = 0, \quad dx = 2r\omega \wedge x, \quad dy_1 = -r\omega \wedge y_1 + \theta\omega \wedge y_2, \quad dy_2 = -r\omega \wedge y_2 - \theta\omega \wedge y_1.$$

We define a left-invariant metric  $\langle \cdot, \cdot \rangle$  on  $\Gamma \backslash G$  such that  $\{A, X, Y_1, Y_2\}$  is an orthonormal frame and a left-invariant complex structure  $J$  by  $JA = X, JY_1 = Y_2$ . Then  $(\Gamma \backslash G, \langle \cdot, \cdot \rangle, J)$  is a locally conformal Kähler manifold with the fundamental 2-form given by

$$\Omega = -\omega \wedge x - y_1 \wedge y_2.$$

Note that the fundamental 2-form  $\Omega$  is not  $-2r\omega$ -exact. We see that the Lee form  $\omega$  is not parallel:  $(\nabla_X\omega)X \neq 0$ .

EXAMPLE 4.4 (Inoue surface  $S^+$  [5], [11], [14]). Let  $G$  be a 4-dimensional solvable Lie group given by

$$G = \left\{ \left( \begin{array}{cccc} 1 & -\frac{1}{2}y & \frac{1}{2}x & z \\ 0 & e^t & 0 & x \\ 0 & 0 & e^{-t} & y \\ 0 & 0 & 0 & 1 \end{array} \right) : t, x, y, z \in \mathbb{R} \right\}.$$

We can construct a lattice  $\Gamma$  on  $G$  (cf. [13]). Let  $\mathfrak{g}$  be the solvable Lie algebra corresponding to  $G$ :

$$\mathfrak{g} = \text{span}\{A, X, Y, Z : [A, X] = X, [A, Y] = -Y, [X, Y] = Z\}.$$

Let  $\{\omega, x, y, z\}$  be the dual base of  $\{A, X, Y, Z\}$ :

$$d\omega = 0, \quad dx = -\omega \wedge x, \quad dy = \omega \wedge y, \quad dz = -x \wedge y.$$

We define a left-invariant metric  $\langle \cdot, \cdot \rangle$  on  $\Gamma \backslash G$  such that  $\{A, X, Y, Z\}$  is an orthonormal frame and a left-invariant complex structure  $J$  by  $JA = Y, JZ = X$ . Then  $(\Gamma \backslash G, \langle \cdot, \cdot \rangle, J)$  is a locally conformal Kähler manifold with the fundamental 2-form given by

$$\Omega_0 = -\omega \wedge y - z \wedge x.$$

Note that the fundamental 2-form  $\Omega$  is not  $-\omega$ -exact. We see that the Lee form  $\omega$  is not parallel:  $(\nabla_Y\omega)Y \neq 0$ .

We mention that an Inoue surface  $S^-$  is not of the form  $\Gamma \backslash G$ , but it is a double covering space of an Inoue surface  $S^+$  (cf. [10]). Then an Inoue surface  $S^-$  has a locally conformal Kähler structure.

ACKNOWLEDGEMENTS. The author would like to express his deep appreciation to Professor Yusuke Sakane and Professor Takumi Yamada for their thoughtful guidance and encouragement during the completion of this paper. The author also thanks Professor Tomonori Noda for several advices.

---

### References

- [1] E. Abbena and A. Grassi: *Hermitian left invariant metrics on complex Lie groups and cosymplectic Hermitian manifolds*, Boll. Un. Mat. Ital. A (6) **5** (1986), 371–379.
- [2] D. Arapura and M. Nori: *Solvable fundamental groups of algebraic varieties and Kähler manifolds*, Compositio Math. **116** (1999), 173–188.
- [3] F.A. Belgun: *On the metric structure of non-Kähler complex surfaces*, Math. Ann. **317** (2000), 1–40.
- [4] C. Benson and C.S. Gordon: *Kähler structures on compact solvmanifolds*, Proc. Amer. Math. Soc. **108** (1990), 971–980.
- [5] L.A. Cordero, M. Fernández and M. de León: *Compact locally conformal Kähler nilmanifolds*, Geom. Dedicata **21** (1986), 187–192.
- [6] S. Dragomir and L. Ornea: *Locally Conformal Kähler Geometry*, Birkhäuser Boston, Boston, MA, 1998.
- [7] K. Hasegawa: *Complex and Kähler structures on compact solvmanifolds*, J. Symplectic Geom. **3** (2005), 749–767.
- [8] K. Hasegawa: *A note on compact solvmanifolds with Kähler structures*, Osaka J. Math. **43** (2006), 131–135.
- [9] A. Hattori: *Spectral sequence in the de Rham cohomology of fibre bundles*, J. Fac. Sci. Univ. Tokyo Sect. I **8** (1960), 289–331.
- [10] M. Inoue: *On surfaces of class VII<sub>0</sub>*, Invent. Math. **24** (1974), 269–310.
- [11] Y. Kamishima: *Note on locally conformal Kähler surfaces*, Geom. Dedicata **84** (2001), 115–124.
- [12] H. Sawai: *Locally conformal Kähler structures on compact nilmanifolds with left-invariant complex structures*, Geom. Dedicata **125** (2007), 93–101.
- [13] H. Sawai: *A construction of lattices on certain solvable Lie groups*, Topology Appl. **154** (2007), 3125–3134.
- [14] F. Tricerri: *Some examples of locally conformal Kähler manifolds*, Rend. Sem. Mat. Univ. Politec. Torino **40** (1982), 81–92.
- [15] I. Vaisman: *Locally conformal Kähler manifolds with parallel Lee form*, Rend. Mat. (6) **12** (1979), 263–284.
- [16] H.-C. Wang: *Complex parallisable manifolds*, Proc. Amer. Math. Soc. **5** (1954), 771–776.

Numazu National College of Technology  
 3600 Ooka, Numazu  
 Japan  
 e-mail: sawai@numazu-ct.ac.jp