

# COMPLEX STRUCTURES AND NON-DEGENERATE CLOSED 2-FORMS OF COMPACT REAL PARALLELIZABLE NILMANIFOLDS

TAKUMI YAMADA

(Received May 21, 2015, revised January 5, 2016)

## Abstract

In this paper, we consider a relation of non-degenerate closed 2-forms and complex structures on compact real parallelizable nilmanifolds.

## 1. Introduction

Let  $\Gamma \backslash G$  be a compact complex parallelizable manifold, where  $G$  is a simply connected complex Lie group, and  $\Gamma$  is a lattice in  $G$ . H. C. Wang [9] has proven that if  $\Gamma \backslash G$  has a Kähler structure, then  $G$  is abelian, in particular,  $\Gamma \backslash G$  is biholomorphic to a complex torus. If  $(\Gamma \backslash G, J)$  has a pseudo-Kählerian structure, then  $G$  is 2-step solvable ([8], [11, 12]). Dorfmeister-Guan [6] have proven that a compact homogeneous pseudo-Kähler manifold is biholomorphic to the product of a homogeneous rational manifold and a complex torus. Thus, we are interested in the case of compact non-homogeneous complex manifolds which have a non-degenerate closed 2-form. By a compact *real* parallelizable manifold we mean a compact manifold of the form  $\Gamma \backslash G$ , where  $G$  is a simply connected *real* Lie group, and  $\Gamma$  is a lattice in  $G$ , that is, a discrete co-compact subgroup of  $G$ . It is known due to Auslander [2] that a solvmanifold has a solvmanifold of the form  $\Gamma \backslash G$  as a finite covering, where  $G$  is a simply connected solvable Lie group, and  $\Gamma$  is a discrete subgroup.

In this paper, we consider a relation of non-degenerate closed 2-forms and complex structures on compact real parallelizable nilmanifolds, and we prove following result.

**Theorem 1.1.** *There exists a compact real parallelizable nilmanifold  $\Gamma \backslash N$  and its complex structures  $J_1, J_2$  such that  $(\Gamma \backslash N, J_1)$  has a pseudo-Kähler structure and no holomorphic symplectic structures, however  $(\Gamma \backslash N, J_2)$  has a holomorphic symplectic structure and no pseudo-Kähler structures.*

## 2. Non-degenerate closed 2-forms and complex structures

In this section, we consider a relation of non-degenerate closed 2-forms and complex structures on compact real parallelizable manifolds.

**DEFINITION 2.1.** Let  $(M, J)$  be a complex manifold. A *pseudo-Kähler structure* on  $(M, J)$  is

---

2010 Mathematics Subject Classification. Primary 53C30; Secondary 53D05.

a real closed non-degenerate  $(1, 1)$ -form. Let  $\dim_{\mathbb{C}} M = 2m$ . Then a *holomorphic symplectic structure* is a closed holomorphic 2-form on  $(M, J)$  of maximal rank  $2m$ .

Note that if  $\Omega$  is a holomorphic symplectic form on  $(M, J)$ , then  $\Omega + \bar{\Omega}$  is a symplectic form on  $M$ .

**Proposition 2.2** ([5]). *Let  $\Gamma \backslash N$  be a compact complex parallelizable nilmanifold. If  $\Gamma \backslash N$  has a pseudo-Kähler structure, then  $\Gamma \backslash N$  is a complex torus.*

Let  $N$  be a simply connected real nilpotent Lie group, and  $\mathfrak{n}$  its Lie algebra. It is well known that  $N$  has a lattice if and only if  $\mathfrak{n}$  has a rational structure, i.e., there exists a rational Lie subalgebra  $\mathfrak{n}_{\mathbb{Q}}$  such that  $\mathfrak{n} \cong \mathfrak{n}_{\mathbb{Q}} \otimes \mathbb{R}$ .

Let us consider following nilpotent Lie groups  $N_1, N_2$  with left-invariant complex structures:

$$N_1 = \left\{ \begin{pmatrix} 1 & w_1 & w_3 \\ 0 & 1 & w_2 \\ 0 & 0 & 1 \end{pmatrix} \middle| w_i \in \mathbb{C} \right\} \times \left\{ \begin{pmatrix} 1 & w_4 \\ 0 & 1 \end{pmatrix} \middle| w_4 \in \mathbb{C} \right\},$$

$$N_2 = \left\{ \begin{pmatrix} 1 & \bar{z}_1 & z_3 \\ 0 & 1 & z_2 \\ 0 & 0 & 1 \end{pmatrix} \middle| z_i \in \mathbb{C} \right\} \times \left\{ \begin{pmatrix} 1 & \bar{z}_4 \\ 0 & 1 \end{pmatrix} \middle| z_4 \in \mathbb{C} \right\}.$$

Let  $\Gamma_1, \Gamma_2$  be lattices in  $N_1, N_2$ , respectively. Then,

$$W_1 = \frac{\partial}{\partial w_1}, W_2 = \frac{\partial}{\partial w_2} + w_1 \frac{\partial}{\partial w_3}, W_3 = \frac{\partial}{\partial w_3}, W_4 = \frac{\partial}{\partial w_4}$$

is a basis of left-invariant vector fields of type  $(1, 0)$  on  $N_1$ , and

$$\tau_1 = dw_1, \tau_2 = dw_2, \tau_3 = dw_3 - w_1 dw_2, \tau_4 = dw_4$$

is its dual basis. Thus,

$$d\tau_1 = d\tau_2 = d\tau_4 = 0, d\tau_3 = -\tau_1 \wedge \tau_2.$$

Hence,  $\Gamma_1 \backslash N_1$  has the following two holomorphic symplectic structures  $\Omega_1, \Omega_2$ , which are not cohomologous:

$$\Omega_1 = \tau_1 \wedge \tau_3 + \tau_2 \wedge \tau_4,$$

$$\Omega_2 = \tau_1 \wedge \tau_4 + \tau_2 \wedge \tau_3.$$

On the other hand,

$$Z_1 = \frac{\partial}{\partial z_1}, Z_2 = \frac{\partial}{\partial z_2} + \bar{z}_1 \frac{\partial}{\partial z_3}, Z_3 = \frac{\partial}{\partial z_3}, Z_4 = \frac{\partial}{\partial z_4}$$

is a basis of left-invariant vector fields of type  $(1, 0)$  on  $N_2$ , and

$$\omega_1 = dz_1, \omega_2 = dz_2, \omega_3 = dz_3 - \bar{z}_1 dz_2, \omega_4 = dz_4$$

is its dual basis. Thus,

$$d\omega_1 = d\omega_2 = d\omega_4 = 0, d\omega_3 = -\bar{\omega}_1 \wedge \omega_2.$$

Hence,  $\Gamma_2 \setminus N_2$  has the following pseudo-Kähler structure  $\omega$ , and holomorphic symplectic structure  $\Omega$ :

$$\omega = \operatorname{Re}(\omega_1 \wedge \bar{\omega}_3 + \omega_2 \wedge \bar{\omega}_4),$$

$$\Omega = \omega_1 \wedge \omega_4 + \omega_2 \wedge \omega_3,$$

where  $\operatorname{Re}(\omega_1 \wedge \bar{\omega}_3 + \omega_2 \wedge \bar{\omega}_4)$  denotes the real part of  $\omega_1 \wedge \bar{\omega}_3 + \omega_2 \wedge \bar{\omega}_4$ .

We can easily see that  $N_1$  and  $N_2$  are isomorphic as *real* Lie groups. Indeed, let  $W_i = U_i + \sqrt{-1}V_i$ ,  $Z_i = X_i + \sqrt{-1}Y_i$  ( $i = 1, 2, 3$ ), where  $U_i, V_i, X_i$ , and  $Y_i$  are real vector fields. Then the structures of Lie algebras of  $N_1$  and  $N_2$  are following:

$$\begin{aligned} [U_1, U_2] &= \frac{1}{2}U_3, & [-V_1, V_2] &= \frac{1}{2}U_3, & [-V_1, U_2] &= -\frac{1}{2}V_3, & [U_1, V_2] &= \frac{1}{2}V_3, \\ [X_1, X_2] &= \frac{1}{2}X_3, & [Y_1, Y_2] &= \frac{1}{2}X_3, & [Y_1, X_2] &= -\frac{1}{2}Y_3, & [Y_1, Y_2] &= \frac{1}{2}Y_3, \end{aligned}$$

and the other brackets being zero. In particular, we can assume that  $\Gamma_1 = \Gamma_2$ .

**REMARK 2.3.** Let us consider the following subgroups of  $N_1$

$$\begin{aligned} \Gamma_1 &= \left\{ \begin{pmatrix} 1 & \mu_1 & \mu_3 \\ 0 & 1 & \mu_2 \\ 0 & 0 & 1 \end{pmatrix} \middle| \mu_i \in \mathbb{Z}[\sqrt{-1}] \right\} \times \left\{ \begin{pmatrix} 1 & \mu_4 \\ 0 & 1 \end{pmatrix} \middle| \mu_4 \in \mathbb{Z}[\sqrt{-1}] \right\} \\ H_1 &= \left\{ \begin{pmatrix} 1 & w_1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \middle| w_1 \in \mathbb{C} \right\} \times \left\{ \begin{pmatrix} 1 & w_4 \\ 0 & 1 \end{pmatrix} \middle| w_4 \in \mathbb{C} \right\}. \end{aligned}$$

Then  $\Gamma_1 \cap H_1 \setminus H_1$  is a holomorphic Lagrangian submanifold of  $(\Gamma_1 \setminus N_1, \Omega)$ .

We generalize our examples above to the other Lie groups. Let us consider the following Lie algebra over  $\mathbb{R}$ :

$$\mathfrak{g} = \mathfrak{a} \ltimes \mathfrak{b},$$

where  $\mathfrak{a}$  is abelian, and  $\mathfrak{b}$  is an ideal. Take bases of Lie subalgebras  $\mathfrak{a}$  and  $\mathfrak{b}$ :

$$\begin{aligned} \mathfrak{a} &= \operatorname{span}_{\mathbb{R}}\{U_1^1, \dots, U_p^1\}, \\ \mathfrak{b} &= \operatorname{span}_{\mathbb{R}}\{V_1^1, \dots, V_q^1\}. \end{aligned}$$

Consider the complexification  $\mathfrak{g}^{\mathbb{C}}$  of  $\mathfrak{g}$ . Since  $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g} + \sqrt{-1}\mathfrak{g}$ ,  ${}_{\mathbb{R}}(\mathfrak{g}^{\mathbb{C}})$  has the following basis:

$$\{U_1^1, \dots, U_p^1, U_1^2, \dots, U_p^2, V_1^1, \dots, V_q^1, V_1^2, \dots, V_q^2\},$$

where  $U_i^2 = \sqrt{-1}U_i^1$ ,  $V_j^2 = \sqrt{-1}V_j^1$ . Let  $J$  be a complex structure on  ${}_{\mathbb{R}}(\mathfrak{g}^{\mathbb{C}})$  defined by

$$JU_i^1 = U_i^2 (JU_i^2 = -U_i^1), \quad JV_j^1 = V_j^2 (JV_j^2 = -V_j^1)$$

for  $i = 1, \dots, p$ ,  $j = 1, \dots, q$ . Note that  $({}_{\mathbb{R}}(\mathfrak{g}^{\mathbb{C}}), J)$  is a complex Lie algebra.

Let  ${}_{\mathbb{R}}(G^{\mathbb{C}})$  be the simply connected real Lie group corresponding to  ${}_{\mathbb{R}}(\mathfrak{g}^{\mathbb{C}})$ . We have the following.

**Proposition 2.4** ([7], [10]). *If  $\mathfrak{g}$  has a non-degenerate closed 2-form, then  $({}_{\mathbb{R}}(G^{\mathbb{C}}), J)$  has a left-invariant holomorphic symplectic form.*

We define other complex structure on  ${}_{\mathbb{R}}(\mathfrak{g}^{\mathbb{C}})$  by the following:

$$\tilde{J}U_i^1 = -U_i^2 \ (\tilde{J}U_i^2 = U_i^1), \quad \tilde{J}V_j^1 = V_j^2 \ (\tilde{J}V_j^2 = -V_j^1)$$

for  $i = 1, \dots, p, j = 1, \dots, q$ .

**Proposition 2.5.**  $\tilde{J}$  is integrable on  ${}_{\mathbb{R}}(G^{\mathbb{C}})$ .

Proof. We show that the Nijenhuis tensor  $N_{\tilde{J}}$  vanishes. Let  $X \in \mathfrak{a}, Y \in \mathfrak{b}$ . Note that  $J \circ ad(Z) = ad(Z) \circ J$  for  $Z \in {}_{\mathbb{R}}(\mathfrak{g}^{\mathbb{C}})$ . Then,

$$\begin{aligned} N_{\tilde{J}}(X, Y) &= [X, Y] + \tilde{J}[\tilde{J}X, Y] + \tilde{J}[X, \tilde{J}Y] - [\tilde{J}X, \tilde{J}Y] \\ &= [X, Y] + J[-JX, Y] + J[X, JY] - [-JX, JY] \\ &= 2([X, Y] + [JX, JY]) \\ &= 2([X, Y] + J^2[X, Y]) = 0. \end{aligned}$$

The other cases are similar, and hence omitted.  $\square$

Moreover, assume that  $\mathfrak{b}$  is a direct sum:  $\mathfrak{b}_1 + \mathfrak{b}_2$ , where  $\mathfrak{b}_1$  is abelian and  $[\mathfrak{a}, \mathfrak{b}] \subset \mathfrak{b}_2$ , and that  $\dim \mathfrak{a} = \dim \mathfrak{b}_2$ . We take bases of the Lie subalgebras  $\mathfrak{b}_1$  and  $\mathfrak{b}_2$ :

$$\begin{aligned} \mathfrak{b}_1 &= \text{span}_{\mathbb{R}}\{V_1^1, \dots, V_r^1\}, \\ \mathfrak{b}_2 &= \text{span}_{\mathbb{R}}\{W_1^1, \dots, W_p^1\}. \end{aligned}$$

Let

$$\{\alpha_1^1, \dots, \alpha_p^1, \beta_1^1, \dots, \beta_r^1, \gamma_1^1, \dots, \gamma_p^1, \alpha_1^2, \dots, \alpha_p^2, \beta_1^2, \dots, \beta_r^2, \gamma_1^2, \dots, \gamma_p^2\}$$

be the dual basis of

$$\{U_1^1, \dots, U_p^1, V_1^1, \dots, V_r^1, W_1^1, \dots, W_p^1, U_1^2, \dots, U_p^2, V_1^2, \dots, V_r^2, W_1^2, \dots, W_p^2\}.$$

Put

$$\lambda_i = \alpha_i^1 + \sqrt{-1}(-\alpha_i^2), \quad \mu_j = \beta_j^1 + \sqrt{-1}\beta_j^2, \quad \nu_k = \gamma_k^1 + \sqrt{-1}\gamma_k^2$$

for  $i, k = 1, \dots, p, j = 1, \dots, r$  (note that  $\tilde{J}U_i^1 = -U_i^2$ ). Then, we can take a basis of left-invariant  $(1, 0)$ -forms of  $({}_{\mathbb{R}}(G^{\mathbb{C}}), \tilde{J})$  as follows:

$$\{\lambda_i, \mu_j, \nu_k\}_{i,k=1,\dots,p,j=1,\dots,r}.$$

**Proposition 2.6.** If  $\mathfrak{a} + \mathfrak{b}_2$  has a non-degenerate 2-form  $\omega_{\mathfrak{ab}_2} = \sum_{k,h} P_{kh} \alpha_k^1 \wedge \gamma_h^1 \in \mathfrak{a}^* \wedge \mathfrak{b}_2^*$  which is closed on  $\mathfrak{g}$ . Then,  $({}_{\mathbb{R}}(G^{\mathbb{C}}), \tilde{J})$  has a left-invariant pseudo-Kähler structure.

Proof. Let

$$\tau = \sum_{k,h} P_{kh} (\bar{\lambda}_k \wedge \nu_h + \lambda_k \wedge \bar{\nu}_h) = 2 \sum_{k,h} P_{kh} (\alpha_k^1 \wedge \gamma_h^1 - \alpha_k^2 \wedge \gamma_h^2).$$

Since  $d\omega_{\mathfrak{ab}_2} = 0$ , and  $\tau(X, \tilde{J}Y) = 0$  for  $X, Y \in \mathfrak{g}$ , we see

$$d\tau(X, Y, Z) = d\tau(\tilde{J}X, \tilde{J}Y, \tilde{J}Z) = d\tau(\tilde{J}X, Y, Z) = 0$$

for  $X, Y, Z \in \mathfrak{g}$ . Let  $X, Y \in \mathfrak{b}, Z \in \mathfrak{g}$ . Since  $\tau(JX, JY) = -\tau(X, Y)$ ,

$$\begin{aligned} d\tau(\tilde{J}X, \tilde{J}Y, Z) &= -\tau([\tilde{J}X, \tilde{J}Y], Z) + \tau([\tilde{J}X, Z], \tilde{J}Y) - \tau([\tilde{J}Y, Z], \tilde{J}X) \\ &= \tau([X, Y], Z) - \tau([X, Z], Y) + \tau([Y, Z], X) \\ &= -2d\omega_{\mathfrak{ab}_2}(X, Y, Z) = 0. \end{aligned}$$

Similarly, we see

$$\begin{aligned} d\tau(\tilde{J}X, \tilde{J}Y, Z) &= 2d\omega_{\mathfrak{ab}_2}(X, Y, Z) = 0 \quad \text{for } X \in \mathfrak{a}, Y \in \mathfrak{b}, Z \in \mathfrak{g} \\ d\tau(\tilde{J}X, \tilde{J}Y, Z) &= -2d\omega_{\mathfrak{ab}_2}(X, Y, Z) = 0 \quad \text{for } X, Y \in \mathfrak{a}, Z \in \mathfrak{g}. \end{aligned}$$

Thus,

$$\omega = \tau + \sqrt{-1} \sum_{j=1}^r \mu_j \wedge \bar{\mu}_j$$

is a pseudo-Kähler structure on  $(\mathbb{R}(G^\mathbb{C}), \tilde{J})$ .  $\square$

**EXAMPLE 2.7.** Let  $H(1, n)$  be the nilpotent Lie group defined by

$$H(1, n) = \left\{ \begin{pmatrix} I_n & \mathbf{x} & \mathbf{z} \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mid \mathbf{x}, \mathbf{z} \in \mathbb{R}^n, y \in \mathbb{R} \right\},$$

where  $I_n$  denotes the identity  $n \times n$  matrix.  $H(1, n)$  is called a *generalized Heisenberg group*. Let

$$N = H(1, n) \times \left\{ \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} \mid w \in \mathbb{R} \right\}.$$

Then,

$$X_i = \frac{\partial}{\partial x_i}, \quad Y = \frac{\partial}{\partial y} - \sum_{i=1}^n x_i \frac{\partial}{\partial z_i}, \quad Z_i = \frac{\partial}{\partial z_i}, \quad W = \frac{\partial}{\partial w}$$

is a basis of left-invariant holomorphic vector fields on  $N$ , and

$$\alpha_i = dx_i, \beta = dy, \gamma_k = dz_k - x_k dy, \delta = dw \quad (i, k = 1, \dots, n)$$

is its dual basis. Then,

$$\sum_{i=1}^n \alpha_i \wedge \gamma_i + \beta \wedge \delta$$

is a symplectic structure. Let

$$\mathfrak{a} = \text{span}\{X_1, \dots, X_n\}, \mathfrak{b}_1 = \text{span}\{Y, Z, W\}, \mathfrak{b}_2 = \{Z_1, \dots, Z_n\}.$$

Then,  $[\mathfrak{a}, \mathfrak{b}_1 + \mathfrak{b}_2] \subset \mathfrak{b}_2$ , and

$$\sum_{i=1}^n \alpha_i \wedge \gamma_i \in \mathfrak{a}^* \wedge \mathfrak{b}_2^*$$

is non-degenerate 2-form on  $\mathfrak{a} + \mathfrak{b}_2$  which is closed on the Lie algebra  $\mathfrak{n}$ .

Thus we see that complex structures  $J, \tilde{J}$  of  $\mathbb{R}(N^\mathbb{C})$  have the properties that the complex manifold  $(\mathbb{R}(N^\mathbb{C}), J)$  admits a left-invariant holomorphic symplectic structure, and the complex manifold  $(\mathbb{R}(N^\mathbb{C}), \tilde{J})$  admits a left-invariant pseudo-Kähler structure. Indeed, let

$$N_2 = (\mathbb{R}(N^{\mathbb{C}}), \tilde{J}) = \left\{ \begin{pmatrix} I_n & \bar{\mathbf{v}} & \mathbf{w} \\ 0 & 1 & u \\ 0 & 0 & 1 \end{pmatrix} \mid \mathbf{v}, \mathbf{w} \in \mathbb{C}^n, u \in \mathbb{C} \right\} \times \left\{ \begin{pmatrix} 1 & \bar{v} \\ 0 & 1 \end{pmatrix} \mid v \in \mathbb{C} \right\}.$$

Then,

$$\zeta_i = dv_i, \lambda_i = dw_i - \bar{v}_i du \quad (i = 1, \dots, n), \mu = du, \nu = dv$$

is a basis of left-invariant  $(1, 0)$ -forms on  $N_2$ . Thus,

$$d\zeta_i = d\mu = dv = 0, \quad d\lambda_i = -\bar{\zeta}_i \wedge \mu \quad (i = 1, \dots, n).$$

Hence,

$$\omega = \operatorname{Re} \left( \sum_{i=1}^n \zeta_i \wedge \bar{\lambda}_i + \mu \wedge \bar{v} \right)$$

is a pseudo-Kähler structure.

By a straightforward computation, we see that  $(\mathbb{R}(N^{\mathbb{C}}), \tilde{J})$  has no left-invariant holomorphic symplectic structures for  $n \geq 2$  and thus  $(\Gamma \backslash \mathbb{R}(N^{\mathbb{C}}), \tilde{J})$  has no holomorphic symplectic structures. Indeed, if  $(\Gamma \backslash \mathbb{R}(N^{\mathbb{C}}), \tilde{J})$  has a holomorphic symplectic structure, then  $(\Gamma \backslash \mathbb{R}(N^{\mathbb{C}}), \tilde{J})$  has an invariant holomorphic symplectic structure by results of Console-Fino [3], and Angella [1].

Next, considering other complex structure  $J$  on  $\mathbb{R}(N^{\mathbb{C}})$ ,

$$N_1 = (\mathbb{R}(N^{\mathbb{C}}), J) = \left\{ \begin{pmatrix} I_n & \mathbf{z} & \mathbf{w} \\ 0 & 1 & u \\ 0 & 0 & 1 \end{pmatrix} \mid \mathbf{z}, \mathbf{w} \in \mathbb{C}^n, u \in \mathbb{C} \right\} \times \left\{ \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \mid v \in \mathbb{C} \right\}.$$

Then,

$$\zeta_i = dz_i, \lambda_i = dw_i - z_i du \quad (i = 1, \dots, n), \mu = du, \nu = dv$$

is a basis of left-invariant holomorphic 1-forms on  $N_1$ . Thus,

$$d\zeta_i = d\mu = dv = 0, \quad d\lambda_i = -\zeta_i \wedge \mu \quad (i = 1, \dots, n).$$

Hence,

$$\Omega = \sum_{i=1}^n \zeta_i \wedge \lambda_i + \mu \wedge \nu$$

is a left-invariant holomorphic symplectic structure. By Proposition 2.2, we see that  $N_1$  has no left-invariant pseudo-Kähler structures. We also see that  $\mathfrak{h}((\Gamma \backslash \mathbb{R}(N^{\mathbb{C}}), \tilde{J})) = n + 2$ , and  $\mathfrak{h}((\Gamma \backslash \mathbb{R}(N^{\mathbb{C}}), J)) = 2n + 2$ , where  $\mathfrak{h}(M)$  is the Lie algebra of holomorphic vector fields on a complex manifold  $M$ .

Hence, we have the following main result.

**Theorem 2.8.** *There exists a compact real parallelizable nilmanifold  $\Gamma \backslash N$  and its complex structures  $J_1, J_2$  such that  $(\Gamma \backslash N, J_1)$  has a pseudo-Kähler structure and no holomorphic symplectic structures, however  $(\Gamma \backslash N, J_2)$  has a holomorphic symplectic structure and no pseudo-Kähler structures.*

**REMARK 2.9.** There exist other examples which satisfy the properties in Theorem 2.8. For example, let  $H(q, p)$  be the nilpotent Lie group defined by

$$H(q, p) = \left\{ \begin{pmatrix} I_p & A & C \\ 0 & I_q & B \\ 0 & 0 & I_q \end{pmatrix} \mid A, C \text{ are } p \times q \text{ matrices, } B \text{ is a diagonal matrix} \right\}.$$

We denote by  $\mathfrak{h}(q, p)$  the Lie algebra of  $H(q, p)$ . Put  $N = H(q_1, p_1) \times H(q_2, p_2)$ . Then,  $N$  is a generalization of the example above (the example above is  $N = H(1, n) \times H(1, 0)$ ). Then

$$\mathfrak{h}(q_1, p_1) \times \mathfrak{h}(q_2, p_2) = \text{span}\{X_{ij}, Y_j, Z_{ij}, X'_{st}, Y'_t, Z'_{st}\}_{i=1, \dots, p_1, j=1, \dots, q_1, s=1, \dots, p_2, t=1, \dots, q_2}$$

with nontrivial equations

$$[X_{ij}, Y_j] = Z_{ij}, [X'_{st}, Y'_t] = Z'_{st}$$

for  $i = 1, \dots, p_1, j = 1, \dots, q_1, s = 1, \dots, p_2, t = 1, \dots, q_2$ .

Consider the Lie subalgebras  $\mathfrak{a}$  and  $\mathfrak{b}$  defined by

$$\begin{aligned} \mathfrak{a} &= \text{span}\{X_{ij}, X'_{st}\}_{i=1, \dots, p_1, j=1, \dots, q_1, s=1, \dots, p_2, t=1, \dots, q_2}, \\ \mathfrak{b} &= \text{span}\{Y_j, Y'_t, Z_{ij}, Z'_{st}\}_{i=1, \dots, p_1, j=1, \dots, q_1, s=1, \dots, p_2, t=1, \dots, q_2}. \end{aligned}$$

Assume that  $q_1 = q_2$ . Then  $N$  has a left-invariant symplectic structure which satisfies the condition in Proposition 2.6 (cf.[4]). Thus, we see that  $(\mathbb{R}(N^{\mathbb{C}}), J)$  admits a left-invariant holomorphic symplectic structure and no left-invariant pseudo-Kähler structures, and  $(\mathbb{R}(N^{\mathbb{C}}), \tilde{J})$  admits a left-invariant pseudo-Kähler structure. By a straightforward computation, we see that  $(\mathbb{R}(N^{\mathbb{C}}), \tilde{J})$  has no left-invariant holomorphic symplectic structures for  $p_1 \geq 2$ .

**ACKNOWLEDGEMENTS.** The author would like to express his deep appreciation to Professor Yusuke Sakane for valuable advice and encouragement during his preparation of the paper. The author also thanks Professor Ryushi Goto for several advice, in particular, Remark 2.3 is due to Professor Ryushi Goto.

---

## References

- [1] D. Angella: *The cohomologies of the Iwasawa manifold and its small deformations*, J. Geom. Anal. **23** (2013), 1355–1378.
- [2] L. Auslander: *Discrete uniform subgroups of solvable Lie group*, Trans. Amer. Math. Soc. **99** (1961), 398–402.
- [3] S. Console and A. Fino: *Dolbeault cohomology of compact nilmanifolds*, Transform. Groups **6** (2001), 111–124.
- [4] L.A. Cordero, M. Fernández and A. Gray: *Symplectic manifolds with no Kahler structure*, Topology **25** (1986), 375–380.
- [5] L.A. Cordero, M. Fernández and L. Ugarte: *Lefschetz complex conditions for complex manifolds*, Ann. Global Anal. Geom. **22** (2002), 355–373.
- [6] J. Dorfmeister and Z.-D. Guan: *Classification of compact homogeneous pseudo-Kähler manifolds*, Comment. Math. Helv. **67** (1992), 499–513.

- [7] D. Guan: *Examples of compact holomorphic symplectic manifolds which admits no manifolds Kähler structures*, Geometry and analysis on complex manifolds, World Sci. Publishing, River Edge, NJ, 63–74, 1994.
- [8] D. Guan: *Classification of compact complex homogeneous manifolds with pseudo-kähler structures*, J. Algebra **324** (2010), 2010–2024.
- [9] H.C. Wang: *Complex parallelisable manifolds*, Proc. Amer. Math. Soc. **5** (1954), 771–776.
- [10] T. Yamada: *A construction of compact pseudo-Kähler solvmanifolds with no Kähler structures*, Tsukuba J. Math. **29** (2005), 79–109.
- [11] T. Yamada: *A pseudo-Kähler structure on a compact non-toral complex parallelizable solvmanifolds*, Geom. Dedicata **112** (2005), 115–122.
- [12] T. Yamada: *A structure theorem of compact complex parallelizable pseudo-Kähler solvmanifolds*, Osaka J. Math. **43** (2006), 923–933.

Department of Mathematics  
Shimane University  
Nishikawatsu-cho 1060  
Matsue 690-8504  
Japan  
e-mail: t\_yamada@riko.shimane-u.ac.jp