

DIAGONALIZING PROJECTIONS IN MULTIPLIER ALGEBRAS AND IN MATRICES OVER A C^* -ALGEBRA

SHUANG ZHANG

Assume that \mathcal{A} is a C^* -algebra with the FS property ([3] and [16]). We prove that every projection in $M_n(\mathcal{A})$ ($n \geq 1$) or in $L(\mathcal{H}_{\mathcal{A}})$ is homotopic to a projection whose diagonal entries are projections of \mathcal{A} and off-diagonal entries are zeros. This yields partial answers for Questions 7 and 8 raised by M. A. Rieffel in [18]. If \mathcal{A} is σ -unital but non-unital, then every projection in the multiplier algebra $M(\mathcal{A})$ is unitarily equivalent to a diagonal projection, and homotopic to a block-diagonal projection with respect to an approximate identity of \mathcal{A} consisting of an increasing sequence of projections. The unitary orbits of self-adjoint elements of \mathcal{A} and $M(\mathcal{A})$ are also considered.

0. Introduction. It is well known that a projection in $M_n(\mathbb{C})$ or in $L(\mathcal{H})$ is homotopic to a diagonal projection whose diagonal entries are either 1 or 0, where $M_n(\mathbb{C})$ is the algebra consisting of $n \times n$ scalar matrices and $L(\mathcal{H})$ is the algebra consisting of bounded operators on a separable Hilbert space \mathcal{H} . The following natural question comes up: if \mathbb{C} is replaced by a C^* -algebra \mathcal{A} , is every projection in $M_n(\mathcal{A})$ or $L(\mathcal{H}_{\mathcal{A}})$ homotopic to a diagonal projection whose diagonal entries are projections of \mathcal{A} and off-diagonal entries are zeros? Here $M_n(\mathcal{A})$ is the C^* -algebra of $n \times n$ matrices over \mathcal{A} and $L(\mathcal{H}_{\mathcal{A}})$ can be regarded as bounded infinite matrices over \mathcal{A} whose adjoints exist (see §1 for a more precise description). Certainly, diagonalizing projections of $M_n(\mathcal{A})$ for $n \geq 1$ would yield information about $K_0(\mathcal{A})$ (here diagonalizing projections in the sense of Murray-von Neumann is enough for this purpose).

Concerning the matrix algebra $M_n(\mathcal{A})$, R. V. Kadison proved ([13] and [14]) that if \mathcal{A} is a von Neumann algebra, then every normal element in $M_n(\mathcal{A})$ is unitarily equivalent to a diagonal normal matrix over \mathcal{A} . Consequently, every projection in $M_n(\mathcal{A})$ is homotopic to a diagonal projection, since the unitary group of a von Neumann algebra is connected. In general, we certainly do not expect a positive answer for the question if \mathcal{A} is an arbitrary C^* -algebra. K. Grove and

G. K. Pedersen have pointed out ([11, 1.3]) that if \mathcal{A} is the algebra $C(S^2)$, the algebra of complex-valued continuous functions on S^2 , then there exists a projection in $M_2(\mathcal{A})$ which is not unitarily equivalent to any diagonal projection. However, we do expect a positive answer for a large class of C^* -algebras.

The author has proved ([22]) that if \mathcal{A} is a C^* -algebra with FS, then every projection in $M_n(\mathcal{A})$ or in $L(\mathcal{K}_{\mathcal{A}})$ is Murray-von Neumann equivalent to a diagonal projection. In this note, we will strengthen the previous results to unitary equivalence or homotopy. We prove that if \mathcal{A} is a C^* -algebra with FS (not necessarily σ -unital), and if p is a projection of the multiplier algebra $M(\mathcal{A})$, then every projection q of \mathcal{A} is homotopic to a projection $q' = p_1 + p_2$, where p_1 is a projection of $p\mathcal{A}p$ and p_2 is a projection of $(1-p)\mathcal{A}(1-p)$. As a special case, by induction we conclude that every projection in $M_n(\mathcal{A})$ is homotopic to a diagonal projection. This yields partial answers for Questions 7 and 8 raised by M. A. Rieffel in [18]. If \mathcal{A} is σ -unital and $\{e_n\}$ is a fixed sequence of mutually orthogonal projections of \mathcal{A} such that $\sum_{n=1}^{\infty} e_n = 1$, we prove that every projection in $M(\mathcal{A})$ is unitarily equivalent to a diagonal projection and homotopic to a block-diagonal projection with respect to the decomposition $\sum_{n=1}^{\infty} e_n = 1$. As a consequence, every projection in $L(\mathcal{K}_{\mathcal{A}})$ is unitarily equivalent (and hence homotopic) to a diagonal projection. In addition, the unitary orbits of self-adjoint elements of \mathcal{A} or $M(\mathcal{A})$ are considered.

The class of C^* -algebras with FS includes many interesting subclasses of C^* -algebras. Obviously, AF algebras, the Calkin algebra, von Neumann algebras and AW^* -algebras have FS. The Bunce-Deddens algebras have FS ([2]). All purely infinite, simple C^* -algebras have FS ([24, Part I (1.3)] and [25]); in particular, the Cuntz algebras \mathcal{O}_n and \mathcal{O}_A , where $2 \leq n \leq \infty$ and A is an irreducible scalar matrix, have FS. Certain irrational rotation C^* -algebras have FS ([9]). Many corona and multiplier algebras have FS ([5], [24, Part I] and [24, Part IV]). L. G. Brown and G. K. Pedersen have recently proved ([5]) that a C^* -algebra \mathcal{A} has FS if and only if $M_n(\mathcal{A})$ has FS for all $n \geq 1$; and \mathcal{A} has FS if and only if \mathcal{A} has real rank zero. In [21], [22], [23] and [24] the author has investigated the multiplier and corona algebras of C^* -algebras with FS from various angles.

1. Notations. If \mathcal{A} is a C^* -algebra, we denote the Banach space double dual of \mathcal{A} by \mathcal{A}^{**} and the multiplier algebra of \mathcal{A} by $M(\mathcal{A})$; where $M(\mathcal{A}) = \{m \in \mathcal{A}^{**} : xm, mx \in \mathcal{A} \ \forall x \in \mathcal{A}\}$ ([1], [7], [15], among others).

Let $\mathcal{H}_{\mathcal{A}} = \{\{a_i\}: a_i \in \mathcal{A} \text{ and } \sum_{i=1}^{\infty} a_i^* a_i \text{ converges in norm}\}$. Then $\mathcal{H}_{\mathcal{A}}$ becomes a Hilbert \mathcal{A} -module with the \mathcal{A} -valued inner product

$$\langle \{a_i\}, \{b_i\} \rangle = \sum_{i=1}^{\infty} a_i^* b_i \text{ for all } \{a_i\}, \{b_i\} \in \mathcal{H}_{\mathcal{A}}.$$

We denote by $L(\mathcal{H}_{\mathcal{A}})$ the set of all bounded module maps with an adjoint and by $K(\mathcal{H}_{\mathcal{A}})$ a closed ideal of $L(\mathcal{H}_{\mathcal{A}})$ called the “compact maps”; more precisely, $K(\mathcal{H}_{\mathcal{A}})$ is the norm closure of the set of all “finite rank” module maps, $\{\sum_{i=1}^n \theta_{x_i, y_i}: x_i, y_i \in \mathcal{H}_{\mathcal{A}} \text{ and } n \in \mathbb{N}\}$. Here for any pair of elements x and y in $\mathcal{H}_{\mathcal{A}}$, $\theta_{x, y}$ is defined by $\theta_{x, y}(a) = x \langle y, a \rangle \in \mathcal{H}_{\mathcal{A}}$ for all $a \in \mathcal{H}_{\mathcal{A}}$ ([15]). It was proved ([15]) that

$$L(\mathcal{H}_{\mathcal{A}}) \cong M(\mathcal{A} \otimes \mathcal{K}) \text{ and } K(\mathcal{H}_{\mathcal{A}}) \cong \mathcal{A} \otimes \mathcal{K}$$

as C^* -algebras, where \mathcal{K} is the algebra consisting of compact operators on \mathcal{H} . The formulation of $L(\mathcal{H}_{\mathcal{A}})$ and $K(\mathcal{H}_{\mathcal{A}})$ are closely analogous to those of $L(\mathcal{H})$ and \mathcal{K} .

If \mathcal{A} is a unital C^* -algebra, we will denote the unitary group of $M_n(\mathcal{A})$ by $U_n(\mathcal{A})$ and the path component of $U_n(\mathcal{A})$ containing the identity by $U_n^0(\mathcal{A})$. In particular, we will denote $U_1^0(\mathcal{A})$ by $U_0(\mathcal{A})$.

If p and q are projections in \mathcal{A} , $p \sim q$ means that p and q are equivalent in the sense of Murray-von Neumann, and $p \approx q$ means that p and q are homotopic, i.e., in the same norm path component of projections in \mathcal{A} . It is well known that $p \approx q$ if and only if there exists a unitary element v in $U_0(\mathcal{A})$ such that $vpv^* = q$. We denote the matrix units of \mathcal{K} by $\{e_{ij}\}$.

2. Key Lemmas. The following technical lemmas are the key of this paper:

2.1. **LEMMA.** *Suppose that \mathcal{A} is a C^* -algebra with FS (not necessarily σ -unital) and p is a projection in $M(\mathcal{A})$. If q is a projection in \mathcal{A} , then for any $\varepsilon_0 > 0$ there exists a projection q' in \mathcal{A} such that both $pq'p$ and $(1-p)q'(1-p)$ have finite spectra and $\|q - q'\| < \varepsilon_0$. More precisely, the projection q' has the following form:*

$$q' = \begin{pmatrix} f_0 & 0 & 0 \\ 0 & a_0 & b_0 \\ 0 & b_0^* & c_0 \end{pmatrix},$$

where f_0 and the range of a_0 are mutually orthogonal subprojections of p . Consequently $q' \approx q$ if $\varepsilon_0 < 1$.

Proof. Let $q = \begin{pmatrix} a & b \\ b^* & c \end{pmatrix}$ be the decomposition of q with respect to $p + (1-p) = 1$. It follows that $a - a^2 = bb^*$, $c - c^2 = b^*b$, $ab + bc = b$, $0 \leq a \leq p$ and $0 \leq c \leq 1 - p$. (Actually these conditions are also sufficient for q to be a projection.) We will start with the idea in [6] and then go further to construct a projection $q' = \begin{pmatrix} a' & b' \\ b'^* & c' \end{pmatrix}$ such that both $\sigma(a')$ and $\sigma(c')$ are finite sets, and q' is close to q in norm.

Let $0 < \delta < 1$ be a fixed positive number and ε be another positive number such that $3\varepsilon < \delta$. Since \mathcal{A} has FS, there exists a positive element c_1 in $(1-p)\mathcal{A}(1-p)$ with a finite spectrum such that

$$(1) \quad \|c - c_1\| < \varepsilon.$$

Set $e = \chi_{(\delta, \infty)}(c_1 - c_1^2)$. If δ_1 is the smaller root of $t^2 - t + \delta = 0$, then $e = \chi_{(\delta_1, 1 - \delta_1)}(c_1)$ which is a projection in $(1-p)\mathcal{A}(1-p)$.

Set $c_0 = c_1e + \chi_{(1 - \delta_1, 1]}(c_1)$. Then $\sigma(c_0)$ is a finite set, $c_0 - c_0^2 = e(c_1 - c_1^2)e \in e\mathcal{A}e$ and $\|c_0 - c_1\| \leq \delta_1$. It follows that

$$(2) \quad \|c_0 - c\| \leq \varepsilon + \delta_1 < \varepsilon + \sqrt{\delta}.$$

Set $v = (eb^*be)^{-1/2}(eb^*)$, of course where $(eb^*be)^{-1}$ is taken in $e\mathcal{A}e$. Since $e(c_1 - c_1^2)e \geq \delta e$ and hence $eb^*be \geq (\delta - 3\varepsilon)e$, $(eb^*be)^{-1/2}$ exists. It is clear that $vv^* = e$.

Set $b_0 = v^*(c_0 - c_0^2)^{1/2}$. Then $b_0^*b_0 = c_0 - c_0^2$.

Set $a_0 = v^*(e - c_0)v$. Then $a_0 - a_0^2 = b_0b_0^*$ and $a_0b_0 + b_0c_0 = b_0$.

If we first fix δ small enough, then we choose ε small enough and c_1 satisfying (1) such that $\|c - c_0\|$, $\|b - b_0\|$ and $\|(a - a^2) - (a_0 - a_0^2)\|$ are all smaller than any preassigned positive number. However, $\|a - a_0\|$ can be equal to one. Here we give details for further reference.

It is obvious that

$$(3) \quad \|b^*b - (c_1 - c_1^2)\| \leq 3\|c - c_1\| < 3\varepsilon.$$

Since $\|(1 - e)b^*b(1 - e) - (1 - e)(c_1 - c_1^2)(1 - e)\| \leq 3\varepsilon$ and $\|(1 - e)(c_1 - c_1^2)(1 - e)\| \leq \delta$, it is easily seen that

$$(4) \quad \|b(1 - e)\| \leq \sqrt{3\varepsilon + \delta}.$$

Since $eb^*be \geq (\delta - 3\varepsilon)e$, then

$$(5) \quad \|(eb^*be)^{-1}\| \leq (\delta - 3\varepsilon)^{-1}.$$

By [12, 126] and (3), we can choose ε small enough such that

$$(6) \quad \|(eb^*be)^{1/2} - [e(c_1 - c_1^2)e]^{1/2}\| < \delta.$$

By (4) and (6) we can choose ε small enough such that

$$(7) \quad \begin{aligned} \|b_0 - b\| &\leq \|v^*(c_0 - c_0^2)^{1/2} - v^*(eb^*be)^{1/2}\| + \|b(1 - e)\| \\ &\leq \| [e(c_1 - c_1^2)e]^{1/2} - (eb^*be)^{1/2} \| + \sqrt{3\varepsilon + \delta} \\ &< \delta + \sqrt{3\varepsilon + \delta}. \end{aligned}$$

Consequently,

$$(8) \quad \begin{aligned} \|(a - a^2) - (a_0 - a_0^2)\| &= \|bb^* - b_0b_0^*\| \\ &\leq 2\|b_0 - b\| < 2\delta + 2\sqrt{3\varepsilon + \delta}. \end{aligned}$$

It is clear from construction that $q_0 = \begin{pmatrix} a_0 & b_0 \\ b_0^* & c_0 \end{pmatrix}$ is a projection. By Lemma (2.4) of [21], $\sigma(a_0) \setminus \{0, 1\} = \sigma(1 - c_0) \setminus \{0, 1\}$, and hence $\sigma(a_0)$ is also a finite set. The idea of constructing the projection q_0 is due L. G. Brown ([6]) for different purpose.

We will go further to adjust q_0 to a projection $q' = \begin{pmatrix} a' & b' \\ b'^* & c' \end{pmatrix}$ so that $\|a - a'\|$ is small, too. Set $f = v^*v$. Then f is a subprojection of p and $fa_0 = a_0f = a_0$. We claim that $\|faf - a_0\|$ can be arbitrarily small if δ , ε and c_1 are properly chosen. To prove this claim, we need the following estimates.

$$(9) \quad \begin{aligned} \|e(b^*b)^{1/2}(1 - e)\| &= \|e[(b^*b)^{1/2} - (c_1 - c_1^2)^{1/2}](1 - e)\| \\ &\leq \|(b^*b)^{1/2} - (c_1 - c_1^2)^{1/2}\|. \end{aligned}$$

Then by [12, 126] and

$$[(eb^*be)^{1/2}]^2 = eb^*be = [e(b^*b)^{1/2}e]^2 + e(b^*b)^{1/2}(1 - e)(b^*b)^{1/2}e,$$

for a fixed $\delta > 0$ we can choose ε small enough (by (3)) such that

$$(10) \quad \|(b^*b)^{1/2} - (c_1 - c_1^2)^{1/2}\| < \frac{\delta^2}{2} \quad \text{and}$$

$$(11) \quad \|(eb^*be)^{1/2} - e(b^*b)^{1/2}e\| < \delta\sqrt{\frac{\delta}{2}}.$$

Since

$$\begin{aligned} f(a - a_0)f &= v^*ev(a - a_0)v^*ev \\ &= v^*e(vav^* - va_0v^*)ev \\ &= v^*[ec_0e - v(p - a)v^*]v, \end{aligned}$$

then

$$(12) \quad \begin{aligned} \|f(a - a_0)f\| &\leq \|ec_0e - ece\| + \|ece - v(p - a)v^*\| \\ &< \varepsilon + \|ece - v(p - a)v^*\|. \end{aligned}$$

Since $(1-a)b = bc$, $p(1-a)b = bp(c)$ for any polynomial $p(t)$. Approximating by polynomials, we obtain that $\sqrt{1-ab} = b\sqrt{c}$, and hence

$$b^*(1-a)b = c^2 - c^3 = (b^*b)^{1/2}c(b^*b)^{1/2}.$$

It follows that

$$\begin{aligned} v(p-a)v^* &= (eb^*be)^{-1/2}eb^*(p-a)be(eb^*be)^{-1/2} \\ &= (eb^*be)^{-1/2}e[b^*b - b^*ab]e(eb^*be)^{-1/2} \\ &= (eb^*be)^{-1/2}[e(b^*b)^{1/2}c(b^*b)^{1/2}e](eb^*be)^{-1/2} \\ &= (eb^*be)^{-1/2}[h_1 + h_2](eb^*be)^{-1/2}, \end{aligned}$$

where

$$\begin{aligned} h_1 &= e(b^*b)^{1/2}ece(b^*b)^{1/2}e \\ &= (eb^*be)^{1/2}c(eb^*be)^{1/2} + [e(b^*b)^{1/2}e - (eb^*be)^{1/2}]c(eb^*be)^{1/2} \\ &\quad + (eb^*be)^{1/2}c[e(b^*b)^{1/2}e - (eb^*be)^{1/2}] \\ &\quad + [e(b^*b)^{1/2}e - (eb^*be)^{1/2}]c[e(b^*b)^{1/2}e - (eb^*be)^{1/2}], \\ h_2 &= e(b^*b)^{1/2}(1-e)ce(b^*b)^{1/2}e \\ &\quad + e(b^*b)^{1/2}ec(1-e)(b^*b)^{1/2}e \\ &\quad + e(b^*b)^{1/2}(1-e)c(1-e)(b^*b)^{1/2}e. \end{aligned}$$

If δ is first fixed small enough, and ε and c_1 can be chosen such that $6\varepsilon < \delta$ and

$$\begin{aligned} (13) \quad &\|(eb^*be)^{-1/2}h_1(eb^*be)^{-1/2} - ece\| \\ &\leq 2\|(eb^*be)^{-1/2}\| \|e(b^*b)^{1/2}e - (eb^*be)^{1/2}\| \|c\| \\ &\quad + \|(eb^*be)^{-1/2}\|^2 \|e(b^*b)^{1/2}e - (eb^*be)^{1/2}\|^2 \|c\| \\ &\leq 2\frac{\delta\sqrt{\delta/2}}{\sqrt{\delta-3\varepsilon}} + \left[\frac{\delta\sqrt{\delta/2}}{\sqrt{\delta-3\varepsilon}}\right]^2 < \delta^2 + 2\delta, \end{aligned}$$

(where using (5), (10) and (11)) and

$$\begin{aligned} (14) \quad &\|(eb^*be)^{-1/2}h_2(eb^*be)^{-1/2}\| \\ &\leq 2\|(eb^*be)^{-1/2}\|^2 \|e(b^*b)^{1/2}(1-e)\| \|c\| \|(b^*b)^{1/2}\| \\ &\quad + \|(eb^*be)^{-1/2}\|^2 \|e(b^*b)^{1/2}(1-e)\|^2 \|c\| \\ &< (\delta-3\varepsilon)^{-1} \left[\delta^2 + \frac{\delta^4}{4}\right] < 2\delta + \delta^2, \end{aligned}$$

where we used $\delta-3\varepsilon > \delta/2$. Consequently,

$$\begin{aligned} \|v(p-a)v^* - ece\| &\leq 4\delta + 2\delta^2, \quad \text{and so} \\ \|f(a-a_0)f\| &< \varepsilon + 4\delta + 2\delta^2 \quad \text{by (12)}. \end{aligned}$$

If δ is fixed small enough and ε is chosen small enough, then $\|faf - a_0\|$ can be arbitrarily small if c_1 satisfies (1).

Moreover, by properly choosing $\delta > 0$, ε and c_1 in a similar way we can require that $\|(p - f)af\|$ is less than any preassigned positive number. This can be done as follows.

Since $a - a^2 = bb^*$ and the spectral mapping theorem, it is clear $\|b\| \leq 1/2$. Since $(1 - a)b = bc$, we have

$$\begin{aligned} -(1 - f)av^* &= (1 - f)(1 - a)be(eb^*be)^{-1/2} \\ &= bce(eb^*be)^{-1/2} - be(eb^*be)^{-1}eb^*bce(eb^*be)^{-1/2} \\ &= bce(eb^*be)^{-1/2} - be(eb^*be)^{-1}eb^*bece(eb^*be)^{-1/2} \\ &\quad - be(eb^*be)^{-1}eb^*b(1 - e)ce(eb^*be)^{-1/2} \\ &= b(1 - e)ce(eb^*be)^{-1/2} \\ &\quad - be(eb^*be)^{-1}eb^*b(1 - e)ce(eb^*be)^{-1/2}. \end{aligned}$$

It follows that

$$\begin{aligned} (15) \quad \|(1 - f)af\| &\leq \|(1 - f)av^*\| \\ &\leq \|b\| \|(1 - e)ce\| \|(eb^*be)^{-1/2}\| \\ &\quad + \|b\| \|(eb^*be)^{-1}\| \|e(b^*b)(1 - e)\| \|c\| \|(eb^*be)^{-1/2}\| \\ &< \frac{\varepsilon}{2} \left[\frac{1}{\sqrt{\delta - 3\varepsilon}} \right] + \frac{1}{2} \left[\frac{1}{\delta - 3\varepsilon} \right] (3\varepsilon) \left[\frac{1}{\sqrt{\delta - 3\varepsilon}} \right] \\ &< \left[\frac{\varepsilon}{2} \right] \sqrt{\frac{2}{\delta}} + \left[\frac{3\varepsilon}{2} \right] \left[\frac{2}{\delta} \right]^{3/2}, \end{aligned}$$

where we use (1), (3), (5) and the facts:

$$\begin{aligned} \|(1 - e)ce\| &= \|(1 - e)(c - c_1)e\| \leq \|c - c_1\|, \quad \text{and} \\ \|eb^*b(1 - e)\| &= \|e[b^*b - (c_1 - c_1^2)](1 - e)\| \leq \|b^*b - (c_1 - c_1^2)\|. \end{aligned}$$

As a consequence of the last estimate and (8), for any $0 < \lambda < 1/2$, we can fix δ small enough and then choose ε small enough such that $\sigma((p - f)a(p - f)) \subset [0, \lambda] \cup [1 - \lambda, 1]$. This is because of the following estimates:

$$\begin{aligned} (p - f)[(a - a^2) - (a_0 - a_0^2)](p - f) &= (p - f)(a - a^2)(p - f) \\ &= (p - f)a(p - f) - [(p - f)a(p - f)]^2 - (p - f)afa(p - f), \end{aligned}$$

$$\begin{aligned} &\|(p - f)a(p - f) - [(p - f)a(p - f)]^2\| \\ &\leq \|(p - f)[(a - a^2) - (a_0 - a_0^2)](p - f)\| + \|(1 - f)af\|^2 \\ &\leq \|(a - a^2) - (a_0 - a_0^2)\| + \|(p - f)af\|^2. \end{aligned}$$

Set $f_0 = \chi_{[1/2, 1]}((p-f)a(p-f))$. Then f_0 is a projection in $(p-f)\mathcal{A}(p-f)$ such that $f_0a_0 = a_0f_0 = 0$ and $\|f_0 - (p-f)a(p-f)\| \leq \lambda$. Set $a' = a_0 + f_0$, $b' = b_0$ and $c' = c_0$. Then $q' = \begin{pmatrix} a' & b' \\ b' & c' \end{pmatrix}$ is a projection in \mathcal{A} such that

$$(16) \quad \begin{aligned} \|q' - q\| &\leq \|(f_0 + a_0) - a\| + 2\|b_0 - b\| + \|c_0 - c\| \\ &\leq \|f(a - a_0)f\| + 2\|fa(p-f)\| \\ &\quad + \|f_0 - (p-f)a(p-f)\| + 2\|b_0 - b\| + \|c_0 - c\|. \end{aligned}$$

Combining all above estimates, we first fix λ small enough, then fix δ small enough, and then choose ε small enough and c_1 satisfying (1) so that each term on the right-hand side of (16) is small. Then $\|q - q'\|$ is small. It is clear that $\sigma(pq'p) = \sigma(f_0 + a_0)$ is a finite set. The last sentence in the statement of this lemma is well known. \square

2.2. LEMMA. *Suppose that \mathcal{A} is a C^* -algebra (not necessarily σ -unital) and p is a projection in $M(\mathcal{A})$. If q is a projection in \mathcal{A} such that $\sigma(pqp) \neq [0, 1]$, then there exist two projections q_1 and q_2 in \mathcal{A} such that $q_1 \leq p$, $q_2 \leq 1 - p$ and $q \approx q_1 + q_2$.*

Proof. Let $q = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$ be the composition of q with respect to $p + (1-p) = 1$. Then $a = pqp$, $c = (1-p)q(1-p)$ and $b = pq(1-p)$. By [21, 2.4], $\sigma(a) \setminus \{0, 1\} = \sigma(1-c) \setminus \{0, 1\}$.

If $b = 0$, then $q_1 = a$ and $q_2 = c$ are as desired. Assume that $b \neq 0$. If $1 \notin \sigma(c)$, then $\|c\| < 1$. By the argument of [8, 1], q is path connected to a subprojection q_1 of p . We can assume that $1 \in \sigma(c)$. Since $\sigma(c) \neq [0, 1]$ and 0 is always in $\sigma(c)$, there is a λ in $(0, 1) \setminus \sigma(c)$. Then there exists a positive number ε such that $(\lambda - \varepsilon, \lambda + \varepsilon) \cap \sigma(c) = \emptyset$. Since $b \neq 0$, we can assume that $\sigma(c) \cap (\lambda + \varepsilon, 1) \neq \emptyset$ (Otherwise, $\sigma(a) \cap (\lambda + \varepsilon, 1) \neq \emptyset$, we consider a instead.) We will use a variation of [8, 1] to construct a path of projections for our purpose.

Define a family of continuous positive functions $\{f_t\}_{t \in [0, 1]}$ from $[0, 1]$ to $[0, 1]$ with the following properties:

- (1) $\lim_{t \rightarrow t_0} \|f_t - f_{t_0}\|_\infty = 0$ for any t_0 in $[0, 1]$;
- (2) $f_1(s) = s$ for all s in $[0, 1]$;
- (3)

$$f_0(s) = \begin{cases} 1, & \text{if } \lambda \leq s \leq 1, \\ \text{linear}, & \text{if } \lambda - \varepsilon < s < \lambda, \\ 0, & \text{if } 0 \leq s \leq \lambda - \varepsilon; \end{cases}$$

(4) For all t in $(0, 1)$, $f_t(s) \leq s$ if $s \in [0, \lambda - \varepsilon]$ and $f_t(s) \geq s$ if $s \in [\lambda, 1]$.

Since q is a projection, $bc = (1 - a)b$. Approximating by polynomials, we obtain that $bg(c) = g(1 - a)b$ for any continuous function g on $[0, 1]$. Set

$$\begin{aligned} c_t &= f_t(c), \\ b_t &= b \left[\frac{f_t(c) - f_t(c)^2}{c - c^2} \right]^{1/2}, \\ a_t &= p - f_t(p - a). \end{aligned}$$

Then b_t and c_t are well defined elements in \mathcal{A} by the properties of f_t . Although $p - a$ is not in $p\mathcal{A}p$ if p is in $M(\mathcal{A}) \setminus \mathcal{A}$, $p - f_t(p - a)$ is in $p\mathcal{A}p$ for $t \in [0, 1]$. To see this, first, $f_t(p - a)$ is well defined for each $t \in [0, 1]$ since $\sigma(p - a) \setminus \{0, 1\} = \sigma(c) \setminus \{0, 1\}$. Second, if we denote by π the canonical map from $(p\mathcal{A}p)^+$ to $(p\mathcal{A}p)^+ / p\mathcal{A}p$, where $(p\mathcal{A}p)^+$ is the C^* -algebra obtained by joining an identity to $p\mathcal{A}p$, then $p - f_t(p - a) \in p\mathcal{A}p$, since $\pi(p - f_t(p - a)) = \pi(p) - f_t(\pi(p)) = 0$. It is easily verified that

$$\begin{aligned} a_t - a_t^2 &= b_t b_t^*, \\ a_t b_t &= b_t (1 - c_t), \\ c_t - c_t^2 &= b_t^* b_t. \end{aligned}$$

Thus $q(t) = \begin{pmatrix} a_t & b_t \\ b_t^* & c_t \end{pmatrix}$ is a projection in \mathcal{A} for each t in $[0, 1]$. By the property (1) of $\{f_t\}$, $\{q(t)\}_{t \in [0, 1]}$ is contained in the same path component of projections in \mathcal{A} . Then $q(0) \approx q(1) = q$. Since $(\lambda - \varepsilon, \lambda) \cap \sigma(c) = \emptyset$, $c_0 = f_0(c) = \chi_{[\lambda, 1]}(c)$ is a projection of $(1 - p)\mathcal{A}(1 - p)$. It is obvious that

$$q(0) = \begin{pmatrix} a_0 & b_0 \\ b_0^* & c_0 \end{pmatrix} = \begin{pmatrix} a_0 & 0 \\ 0 & c_0 \end{pmatrix}.$$

Consequently, a_0 is a projection of $p\mathcal{A}p$. Set $q_1 = a_0$ and $q_2 = c_0$, as desired. □

Roughly speaking, with respect to a fixed sequential increasing approximate identity of \mathcal{A} a block-diagonal projection of $M(\mathcal{A})$ whose blocks are with the same size is homotopic to a diagonal projection. More precisely, we have the following lemma:

2.3. LEMMA. *Suppose that \mathcal{A} is a σ -unital, non-unital C^* -algebra with FS and $\sum_{i=1}^{\infty} (s_{i1} + s_{i2} + \dots + s_{in}) = 1$, where $\{s_{ij} : i \geq 1, 1 \leq j \leq n\}$ are mutually orthogonal projections in \mathcal{A} and the sum converges*

in the strict topology. If p is a projection in $M(\mathcal{A})$ with the form $\sum_{i=1}^{\infty} p_i$, where p_i is a projection in $(s_{i1} + s_{i2} + \cdots + s_{in})\mathcal{A}(s_{i1} + s_{i2} + \cdots + s_{in})$ for $i \geq 1$, then $p \approx \sum_{i=1}^{\infty} (p_{i1} + p_{i2} + \cdots + p_{in})$, where p_{ij} is a projection in $s_{ij}\mathcal{A}s_{ij}$ for $i \geq 1$ and $1 \leq j \leq n$.

Proof. It suffices to prove the case if $n = 2$. If $n > 2$, we simply employ the same proof recursively $n - 1$ times by induction to reach the conclusion.

We write

$$p_i = \begin{pmatrix} a_i^* & b_i \\ b_i^* & c_i \end{pmatrix}$$

with respect to $s_{i1} + s_{i2}$. By Lemma (2.1), for each $i \geq 1$ we can find a projection

$$p'_i = \begin{pmatrix} f_i & 0 & 0 \\ 0 & a'_i & b'_i \\ 0 & b_i^* & c'_i \end{pmatrix}$$

in $(s_{i1} + s_{i2})\mathcal{A}(s_{i1} + s_{i2})$ such that $\|p'_i - p_i\| < 1/4$, and both a'_i and c'_i have finite spectra. Here we use the proof of Lemma (2.1) to properly choose a positive number δ_i and a positive element c'_{1i} in $s_{i2}\mathcal{A}s_{i2}$ with a finite spectrum, then we have that

$$\begin{aligned} e_i &= \chi_{(\delta_i, 1-\delta_i)}(c'_{1i}), & c'_i &= c'_{1i}e_i + \chi_{(1-\delta_i, 1)}(c'_{1i}), \\ v_i &= (e_i b_i^* b_i e_i)^{-1/2} (e_i b_i^*), & b'_i &= v_i^* (c'_i - c'^2_{ii})^{1/2}, \\ a'_i &= v_i^* (e_i - c'_{1i}) v_i \end{aligned}$$

and f_i is a projection of $s_{i1}\mathcal{A}s_{i1}$ orthogonal to the range projection of a'_i .

Let $p' = \sum_{i=1}^{\infty} p'_i$. Then $\|p' - p\| < 1/4$, and hence $p \approx p'$.

Let $\sigma(c'_i) = \{\lambda_{i1}, \lambda_{i2}, \dots, \lambda_{il_i}\}$ for each $i \geq 1$. It follows from the construction or [21, 2.4] that $\sigma(a'_i) = \{1 - \lambda_{i1}, 1 - \lambda_{i2}, \dots, 1 - \lambda_{il_i}\}$. We can write $c'_i = \sum_{j=1}^{l_i} \lambda_{ij} r_{ij}$, where $\{r_{ij} : 1 \leq j \leq l_i\}$ is a set of mutually orthogonal projections in $s_{i2}\mathcal{A}s_{i2}$. Let λ be any number in the open interval $(\frac{1}{2}, \frac{3}{4})$ but not in $\bigcup_{i=1}^{\infty} \sigma(c'_i)$. Let $\varepsilon = \min\{\lambda - \frac{1}{2}, \frac{3}{4} - \lambda\}$. For $i \geq 1$, if λ_{ij} is in the open interval $(\lambda - \varepsilon, \lambda)$, we replace λ_{ij} by $\lambda'_{ij} = \lambda - \varepsilon$, and if λ_{ij} is in $(\lambda, \lambda + \varepsilon)$, we replace λ_{ij} by $\lambda'_{ij} = \lambda + \varepsilon$. If λ_{ij} is not in $(\lambda - \varepsilon, \lambda + \varepsilon)$, then we let $\lambda'_{ij} = \lambda_{ij}$. Set $c''_i = \sum_{j=1}^{l_i} \lambda'_{ij} r_{ij}$ for $i \geq 1$, and correspondingly set $b''_i = v_i^* (c''_i - c''^2_{ii})^{1/2}$ and $a''_i = v_i^* (e_i - c''_i) v_i$. Then

$$\begin{aligned} \|a'_i - a''_i\| &\leq \|c'_i - c''_i\| < \varepsilon \quad \text{and} \\ \|b'_i - b''_i\| &\leq \|(c'_i - c'^2_{ii})^{1/2} - (c''_i - c''^2_{ii})^{1/2}\| < \frac{1}{8}. \end{aligned}$$

It follows that

$$p_i'' = \begin{pmatrix} f_i & 0 & 0 \\ 0 & a_i'' & b_i'' \\ 0 & b_i''^* & c_i'' \end{pmatrix}$$

is a projection in $(s_{i1} + s_{i2})\mathcal{A}(s_{i1} + s_{i2})$ such that $\|p_i' - p_i''\| \leq 2\varepsilon + \frac{1}{4} < 1$. Define $p'' = \sum_{i=1}^{\infty} p_i''$. Then $\|p' - p''\| < 1$, and hence $p' \approx p''$. The remaining job is to prove that p'' is homotopic to a desired diagonal projection.

Let $\{f_i\}_{t \in [0, 1]}$ be the family of continuous functions defined in the proof of Lemma (2.2). Since $\sigma(c_i'')$ does not intersect with the open interval $(\lambda - \varepsilon, \lambda + \varepsilon)$ for $i \geq 1$, we can define

$$\begin{aligned} c_i(t) &= f_i(c_i''), \\ b_i(t) &= b_i'' \left[\frac{f_t(c_i'') - f_t(c_i'')^2}{c_i'' - c_i''^2} \right]^{1/2}, \\ a_i(t) &= p - f_i(p - a_i'' - f_i). \end{aligned}$$

Then $a_i(t)$, $b_i(t)$ and $c_i(t)$ are well defined elements in $(s_{i1} + s_{i2})\mathcal{A}(s_{i1} + s_{i2})$ for each t in $[0, 1]$ and $i \geq 1$ by the properties of f_i . Thus for each t in $[0, 1]$

$$p_i(t) = \begin{pmatrix} a_i(t) & b_i(t) \\ b_i(t)^* & c_i(t) \end{pmatrix}$$

is a projection in $(s_{i1} + s_{i2})\mathcal{A}(s_{i1} + s_{i2})$. It is easily seen that

$$p_i(1) = p_i'' \quad \text{and} \quad p_i(0) = \begin{pmatrix} a_i(0) & 0 \\ 0 & c_i(0) \end{pmatrix},$$

where $a_i(0)$ is a projection of $s_{i1}\mathcal{A}s_{i1}$ and $c_i(0)$ is a projection of $s_{i2}\mathcal{A}s_{i2}$. Define $p(t) = \sum_{i=1}^{\infty} p_i(t)$ for each t in $[0, 1]$. Then $\{p(t)\}_{t \in [0, 1]}$ is a path of projection in $M(\mathcal{A})$. It is obvious that

$$p(1) = p'' \quad \text{and} \quad p(0) = \sum_{i=1}^{\infty} \begin{pmatrix} a_i(0) & 0 \\ 0 & c_i(0) \end{pmatrix}.$$

Since the choice of $\{f_i\}_{t \in [0, 1]}$ does not depend on i , the path $\{p(t) : t \in [0, 1]\}$ is continuous in the norm topology.

Set $p_{i1} = a_i(0)$, $p_{i2} = c_i(0)$ for $i \geq 1$. Then

$$p \approx p' \approx p'' \approx p(0) = \sum_{i=1}^{\infty} (p_{i1} + p_{i2}), \quad \text{as desired.} \quad \square$$

3. Diagonalizing projections in \mathcal{A} and in $M_n(\mathcal{A})$. Since we will frequently employ the following well-known fact in this paper, we state it as a lemma.

3.1. **LEMMA.** *If \mathcal{A} is a C^* -algebra, and if p and q are two mutually orthogonal projections in \mathcal{A} , then $p \sim q$ if and only if $p \approx q$.*

Proof. Let v be a partial isometry in \mathcal{A} such that $vv^* = p$ and $v^*v = q$. Define $w = v + v^* + (1 - p - q)$. Then w is a self-adjoint unitary in $M(\mathcal{A})$ such that $w^*pw = q$. It is well known that $w \in U_0(\mathcal{A})$. It follows that $p \approx q$. \square

3.2. **THEOREM.** *Suppose that \mathcal{A} is a C^* -algebra with FS and p_1, p_2, \dots, p_n ($n \geq 1$) are mutually orthogonal projections in $M(\mathcal{A})$ such that $\sum_{i=1}^n p_i = 1$. If p is a projection in \mathcal{A} , then $p \approx \sum_{i=1}^n q_i$, where q_i is a projection in \mathcal{A} such that $q_i \leq p_i$ for $1 \leq i \leq n$.*

Proof. Recursively using Lemma (2.1) and Lemma (2.2), we reach the conclusion. \square

The following theorem can be regarded as an analogue of the well-known fact: Every projection in $M_n(\mathbb{C})$ is homotopic to a diagonal projection whose entries are either 1 or 0.

3.3. **THEOREM.** *Assume that \mathcal{A} is a C^* -algebra with FS and $n \geq 1$. If p is a projection in $M_n(\mathcal{A})$, then $p \approx \sum_{i=1}^n p_i \otimes e_{ii}$, where $\{p_i\}$ is a set of projections in \mathcal{A} such that*

$$p_1 \leq p_2 \leq \dots \leq p_{n-1} \leq p_n.$$

Proof. It has been recently proved ([5]) that $\mathcal{A} \otimes \mathcal{K}$ has FS if and only if \mathcal{A} has FS. By Theorem (3.2) we have $p \approx \sum_{i=1}^n p'_i \otimes e_{ii}$, where $\{p'_i\}$ is a set of projections in \mathcal{A} . The remaining work is to adjust $\{p'_i\}$. We use induction on n .

If $n = 2$, $p \approx p'_1 \otimes e_{11} + p'_2 \otimes e_{22}$, where p'_1 and p'_2 are projections in \mathcal{A} . Combining Lemma (2.1) and Lemma (2.2), we obtain that $p'_1 \approx q_1 + q_2$ in \mathcal{A} , where q_1 and q_2 are two projections in \mathcal{A} such that $q_1 \leq p'_2$ and $q_2 \leq 1 - p'_2$. It follows that $p \approx (q_1 + q_2) \otimes e_{11} + p'_2 \otimes e_{22}$. Working in the hereditary C^* -subalgebra of $M_n(\mathcal{A})$ generated by $(1 - q_1) \otimes e_{11} + 1 \otimes e_{22}$, we have $q_2 \otimes e_{11} + p'_2 \otimes e_{22} \approx (p'_2 + q_2) \otimes e_{22}$ by Lemma (3.1). It follows that $p \approx q_1 \otimes e_{11} + (p'_2 + q_2) \otimes e_{22}$. Let $p_1 = q_1$ and $p_2 = q_2 + p'_2$.

Assume that $p \approx \sum_{i=1}^n p'_i \otimes e_{ii}$ such that $p'_2 \leq p'_3 \leq \dots \leq p'_n$. Applying Lemma (2.1) and Lemma (2.2) to p'_1 , and p'_n , we have $p'_1 \approx q_n + q'_n$, where q_n and q'_n are projections in \mathcal{A} such that $q_n \leq 1 - p'_n$ and $q'_n \leq p'_n$. By the same argument as in the last paragraph

we have that $p \approx q'_n \otimes e_{11} + \sum_{i=2}^{n-1} p'_i \otimes e_{ii} + (p'_n + q_n) \otimes e_{nn}$. Repeating this argument to q'_n and p'_{n-1} , we have that $q'_n \approx q'_{n-1} + q_{n-1}$, where q'_{n-1} and q_{n-1} are two projections in \mathcal{A} such that $q_{n-1} \leq p'_n - p'_{n-1}$ and $q'_{n-1} \leq p'_{n-1}$. It follows that $p \approx q'_{n-1} \otimes e_{11} + \sum_{i=2}^{n-2} p'_i \otimes e_{ii} + (p'_{n-1} + q_{n-1}) \otimes e_{n-1, n-1} + (p'_n + q_n) \otimes e_{nn}$.

Proceeding in this way, we write $p'_1 = \sum_{i=1}^n q_i$, where $\{q_i\}$ is a set of mutually orthogonal projections in \mathcal{A} such that $q_i \leq p'_{i+1} - p'_i$ for $2 \leq i \leq n$ (where $p'_{n+1} = 1$), $q_1 \leq p'_2$, and $p \approx q_1 \otimes e_{11} + \sum_{i=2}^n (p'_i + q_i) \otimes e_{ii}$. Let $p_1 = q_1$ and $p_i = p'_i + q_i$ for $2 \leq i \leq n$. Then $p_1 \leq p_2 \leq \dots \leq p_n$ and $p \approx \sum_{i=1}^n p_i \otimes e_{ii}$. \square

M. A. Rieffel raised a question in [18, 7]: If \mathcal{A} is a unital C^* -algebra with cancellation, and if two projections p and q in $M_n(\mathcal{A})$ represent the same class in $K_0(\mathcal{A})$, are p and q in the same path component of projections in $M_n(\mathcal{A})$? Since \mathcal{A} has cancellation, $[p] = [q]$ in $K_0(\mathcal{A})$ if and only if $p \sim q$ ([3] or [4]). Hence, Rieffel's question is equivalent to whether two Murray-von Neumann equivalent projections in $M_n(\mathcal{A})$ are in the same path component of projections in $M_n(\mathcal{A})$. The following corollary provides a partial answer for his question in the case that \mathcal{A} has FS:

3.4. COROLLARY. *If \mathcal{A} is a unital C^* -algebra with FS and cancellation, and if p and q are two projections in $M_n(\mathcal{A})$, then $p \sim q$ if and only if $p \approx q$.*

Proof. Of course we need only to show that $p \sim q$ implies $p \approx q$. Since $M_n(\mathcal{A})$ has FS, by Theorem (3.2) we have $p \approx q_1 + q_2$, where q_1 is a subprojection of q and q_2 is a subprojection of $1 - q$. Since \mathcal{A} has cancellation and $p \sim q$, $q_2 \sim q - q_1$. Working in $(1 - q_1)M_n(\mathcal{A})(1 - q_1)$, by Lemma (3.1) we can find a unitary v in $U_0((1 - q_1)M_n(\mathcal{A})(1 - q_1))$ such that $vq_2v^* = q - q_1$. Set $u = q_1 + v$. Then u is a unitary in $U_0(M_n(\mathcal{A}))$ such that $uq_1 = q_1u$. Thus $p \approx q_1 + q_2 \approx q$. \square

Concerning the unitary orbit of elements in $M_n(\mathcal{A})$, we have the following corollary:

3.5. COROLLARY. *If \mathcal{A} is a C^* -algebra with FS and x is a normal element in $M_n(\mathcal{A})$ with finite spectrum, then there is a unitary element u in $U_n^0(\mathcal{A})$ such that $uxu^* = \sum_{j=1}^n [\sum_{i=1}^n \lambda_i p_{ij}] \otimes e_{jj}$, where $\{p_{ij}\}$ is a set of projections in \mathcal{A} such that $p_{i,j} \perp p_{i,j}$ in $\mathcal{A} \otimes e_{jj}$ if $i_1 \neq i_2$.*

Proof. By operator calculus we write $x = \sum_{i=1}^m \lambda_i p_i$, where $\{\lambda_i\}$ is a set of complex numbers and $\{p_i\}$ is a set of mutually orthogonal projections in $M_n(\mathcal{A})$. By Theorem (3.2) we can find a unitary element u_1 in $U_n^0(\mathcal{A})$ such that $u_1 p_1 u_1^* = \sum_{j=1}^n p_{1j} \otimes e_{jj} (= q_1)$ for some projections $\{p_{1j}\}$ in \mathcal{A} . Working in $(I_n - q_1)M_n(\mathcal{A})(I_n - q_1)$ and repeating the same argument, we can find a unitary u_2' in $U_0[(I_n - q_1)M_n(\mathcal{A})(I_n - q_1)]$ such that $u_2'(u_1 p_2 u_1^*) u_2'^2 = \sum_{j=1}^n p_{2j} \otimes e_{jj}$ for some projections $\{p_{2j}\}$ in \mathcal{A} . It follows from $p_1 p_2 = 0$ that $p_{1j} p_{2l} = 0$ for $1 \leq j < l \leq n$. Set $u_2 = q_1 + u_2'$. Then u_2 is a unitary in $U_n^0(\mathcal{A})$ and $u_2 u_1 (p_1 + p_2) u_1^* u_2^* = \sum_{i=1}^2 \sum_{j=1}^n p_{ij} \otimes e_{jj} = \sum_{j=1}^n (\sum_{i=1}^2 p_{ij}) \otimes e_{jj}$.

Proceeding in this way we can find unitary elements $\{u_i: 1 \leq i \leq m\}$ in $U_n^0(\mathcal{A})$ such that

$$\begin{aligned} & u_m u_{m-1} \cdots u_1 (p_1 + p_2 + \cdots + p_m) u_1^* \cdots u_{m-1}^* u_m^* \\ &= \sum_{i=1}^m \left[\sum_{j=1}^n p_{ij} \otimes e_{jj} \right] = \sum_{j=1}^n \left[\sum_{i=1}^m p_{ij} \right] \otimes e_{jj}. \end{aligned}$$

Let $u = u_m \cdots u_2 u_1$. It is obvious that u is in $U_n^0(\mathcal{A})$ and

$$u x u^* = \sum_{j=1}^n \left[\sum_{i=1}^m \lambda_i p_{ij} \right] \otimes e_{jj}. \quad \square$$

It is well known that the unitary orbit of a self-adjoint matrix in $M_n(\mathbb{C})$ contains a diagonal self-adjoint matrix. If \mathbb{C} is replaced by a unital C^* -algebra with FS, we have the following weaker analogue:

3.6. COROLLARY. *If \mathcal{A} is a C^* -algebra with FS and x is a self-adjoint element in $M_n(\mathcal{A})$ ($n \geq 1$), then for any $\varepsilon > 0$ there exist a unitary element u in $U_n^0(\mathcal{A})$ and elements a_i in \mathcal{A} with finite spectra such that*

$$\left\| u x u^* - \sum_{i=1}^n a_i \otimes e_{ii} \right\| < \varepsilon.$$

Proof. Since $M_n(\mathcal{A})$ has FS, there is a self-adjoint element h in $M_n(\mathcal{A})$ with finite spectrum such that $\|x - h\| < \varepsilon$. By the same argument as in the proof of Corollary (3.5) we can find a unitary element u in $U_n^0(\mathcal{A})$ such that $u h u^* = \sum_{i=1}^n a_i \otimes e_{ii}$, where $\{a_i\}$ is a set of self-adjoint elements in \mathcal{A} with finite spectra. Therefore,

$$\left\| u x u^* - \sum_{i=1}^n a_i \otimes e_{ii} \right\| = \|x - h\| < \varepsilon. \quad \square$$

3.7. **REMARK.** Concerning the computation of K_0 -groups of a C^* -algebra, M. A. Rieffel raised a question in [18, 8]: What is the smallest n such that the projections in $M_n(\mathcal{A})$ generate $K_0(\mathcal{A})$? Theorem (3.3) provides a partial answer for his question for the class of C^* -algebras with FS (actually it has been given in [22] although it was not mentioned there). In fact, if \mathcal{A} is a C^* -algebra with FS, then the smallest such an integer is $n = 1$; in other words, $K_0(\mathcal{A})$ is generated by the set of Murray-von Neumann equivalence classes of projections in \mathcal{A} .

4. Diagonalizing projections in $M(\mathcal{A})$.

4.1. **THEOREM.** *Assume that \mathcal{A} is a σ -unital C^* -algebra with FS and $\{e_n\}$ is a fixed increasing sequential approximate identity consisting of projections. If p is a projection in $M(\mathcal{A})$, then the following hold:*

(i) *There is a unitary u in $M(\mathcal{A})$ connected to the identity by a path of unitaries, where the path is continuous in the strict topology, such that $upu^* = \sum_{i=1}^{\infty} p_i$, where $p_i \leq e_i$ for $i \geq 1$; in other words, each strict path component of projections in $M(\mathcal{A})$ contains a diagonal projection with respect to $\{e_n\}$.*

(ii) *There exist a unitary v in $U_0(M(\mathcal{A}))$ and a subsequence $\{e_{m_i}\}$ of $\{e_n\}$ such that $vpv^* = \sum_{i=1}^{\infty} p'_i$, where p'_i is a projection of $(e_{m_i} - e_{m_{i-1}})\mathcal{A}(e_{m_i} - e_{m_{i-1}})$ for $i \geq 1$; in other words, each norm path component of projections in $M(\mathcal{A})$ contains a block-diagonal projection with respect to $\{e_n\}$.*

Before proving this theorem, we state the following corollary, which can be regarded as an analogue of the well known fact that a projection on a separable Hilbert space is unitarily equivalent to a diagonal projection whose diagonal entries are either 1 or 0.

4.2. **COROLLARY.** *If \mathcal{A} is a σ -unital C^* -algebra with FS, and if p is a projection in $L(\mathcal{H}_{\mathcal{A}})$, then there is a unitary u in $L(\mathcal{H}_{\mathcal{A}})$ such that $upu^* = \sum_{i=1}^{\infty} p_i \otimes e_{ii}$, where $\{p_i\}$ is a sequence of projections in \mathcal{A} . Consequently, $p \approx \sum_{i=1}^{\infty} p_i \otimes e_{ii}$ (by [8]).*

Proof of Theorem (4.1).

Case 1. If p is a projection of \mathcal{A} .

Choose $n \geq 1$ large enough such that $\|p(1 - e_n)p\|$ is small. Then Lemma (2.1) of [10] applies. We find a unitary u in $U_0(M(\mathcal{A}))$ such

that $upu^* \leq e_n$. By Theorem (3.2), $p \approx upu^* \approx \sum_{i=1}^n p_i$, where $p_i \leq e_i - e_{i-1}$ for $1 \leq i \leq n$. Hence both (i) and (ii) hold.

Case 2. If p is a projection in $M(\mathcal{A}) \setminus \mathcal{A}$.

Let $\{q_n\}$ and $\{q'_n\}$ be two increasing sequences of projections in \mathcal{A} such that $q_n \nearrow p$ and $q'_n \nearrow 1 - p$ in the strict topology. Set $f_n = q_n + q'_n$. Then $\{f_n\}$ is an increasing sequential approximate identity of \mathcal{A} consisting of projections. By the argument of [10, 2.4] we find a unitary element v in $U_0(M(\mathcal{A}))$ such that

$$e_{m_1} \leq v f_{n_1} v^* \leq e_{m_2} \leq v f_{n_2} v^* \leq e_{m_3} \leq \cdots,$$

where $\{n_i\}$ and $\{m_i\}$ are increasing sequences. It is clear that

$$v p v^* = \sum_{i=1}^{\infty} v p (f_{n_i} - f_{n_{i-1}}) v^* = \sum_{i=1}^{\infty} v (q_{n_i} - q_{n_{i-1}}) v^*$$

and $v(q_{n_i} - q_{n_{i-1}})v^* \leq v(f_{n_i} - f_{n_{i-1}})v^* = (v f_{n_i} v^* - e_{m_i}) + (e_{m_i} - v f_{n_{i-1}} v^*)$ (where $q_{n_0} = 0$ and $f_{n_0} = 0$).

We first prove (i). By Theorem (3.2) we find a unitary w_i in $U_0(\mathcal{A}_i)$, where $\mathcal{A}_i = [v(f_{n_i} - f_{n_{i-1}})v^*]\mathcal{A}[v(f_{n_i} - f_{n_{i-1}})v^*]$, such that $w_i v(q_{n_i} - q_{n_{i-1}})v^* w_i^* = r_i + r'_i$, where $r_i \leq v f_{n_i} v^* - e_{m_i}$ and $r'_i \leq e_{m_i} - v f_{n_{i-1}} v^*$. Set $w = \sum_{i=1}^{\infty} w_i$. Then w is a unitary in $M(\mathcal{A})$ such that w is path connected (in the strict topology) to the identity and

$$w v p v^* w^* = \sum_{i=1}^{\infty} (r_i + r'_i) \leq \sum_{i=1}^{\infty} [(v f_{n_i} v^* - e_{m_i}) + (e_{m_i} - v f_{n_{i-1}} v^*)].$$

Since $r_i + r'_{i+1} \leq e_{m_{i+1}} - e_{m_i}$, we can apply Theorem (3.2) again to get a unitary w'_i in $U_0(\mathcal{B}_i)$, where $\mathcal{B}_i = (e_{m_{i+1}} - e_{m_i})M(\mathcal{A})(e_{m_{i+1}} - e_{m_i})$ such that

$$w'_i (r_i + r'_{i+1}) w_i^* = \sum_{j=m_i+1}^{m_{i+1}} p_j,$$

where p_j is in $(e_j - e_{j-1})\mathcal{A}(e_j - e_{j-1})$ for $m_i < j \leq m_{i+1}$.

Define $w' = \sum_{i=1}^{\infty} w'_i$. Then w' is a unitary in $M(\mathcal{A})$ such that w' is path connected in the strict topology to the identity and $w' w v p v^* w^* w'^* = \sum_{i=1}^{\infty} p_i$. Set $u = w' w v$, as (i) desired.

To prove (ii), we start with $p \approx v p v^* = \sum_{i=1}^{\infty} v(q_{n_i} - q_{n_{i-1}})v^*$, where $s_i = v(q_{n_i} - q_{n_{i-1}})v^* \leq v(f_{n_i} - f_{n_{i-1}})v^* = (v f_{n_i} v^* - e_{m_i}) + (e_{m_i} - v f_{n_{i-1}} v^*)$ for each $1 \geq 1$ and $q_{n_0} = 0$ and $f_{n_0} = 0$. With respect to

$$v(f_{n_i} - f_{n_{i-1}})v^* = (v f_{n_i} v^* - e_{m_i}) + (e_{m_i} - v f_{n_{i-1}} v^*),$$

we can write

$$s_i = \begin{pmatrix} a_i & b_i \\ b_i^* & c_i \end{pmatrix} \quad \text{for } i \geq 1.$$

By Lemma (2.3),

$$vpv^* \approx \sum_{i=1}^{\infty} (s_i + s'_i),$$

where s_i is a projection in $(vf_{n_i}v^* - e_{m_i})\mathcal{A}(vf_{n_i}v^* - e_{m_i})$ and s'_i is a projection in $(e_{m_i} - vf_{n_{i-1}}v^*)\mathcal{A}(e_{m_i} - vf_{n_{i-1}}v^*)$. Let $p'_i = s'_i + s_{i-1}$ for $i \geq 1$, where $s_0 = 0$, as desired. \square

The following theorem asserts that the unitary orbit of each self-adjoint element of $M(\mathcal{A})$ contains an “almost” diagonal form, which is a natural analogue of the classical Weyl-von Neumann theorem.

4.3. THEOREM. *Assume that \mathcal{A} is a σ -unital C^* -algebra with FS and also $M(\mathcal{A})$ has FS. If $\{e_n\}$ is a fixed increasing approximate identity of \mathcal{A} consisting of projections and h is a self-adjoint element in $M(\mathcal{A})$, then there exist a unitary u in $M(\mathcal{A})$, an element a in \mathcal{A} , some mutually orthogonal subprojection p_{ij} ($1 \leq j \leq n_i$) of $e_i - e_{i-1}$ for each $i \geq 1$ and a real bounded scalar sequence $\{\lambda_{ij}\}$ such that*

$$\sum_{ij} p_{ij} = 1, \quad \text{and} \quad uhu^* = \sum_{i=1}^{\infty} \left[\sum_{j=1}^{l_i} \lambda_{ij} p_{ij} \right] + a,$$

where a can be chosen such that $\|a\|$ is arbitrarily small. Moreover, u is connected to the identity by a path of unitaries in $M(\mathcal{A})$, where the path is continuous in the strict topology.

4.4. COROLLARY. *If \mathcal{A} is a unital C^* -algebra with FS and $L(\mathcal{K}_{\mathcal{A}})$ has FS also, then for any self-adjoint element h in $L(\mathcal{K}_{\mathcal{A}})$ there are a unitary u in $L(\mathcal{K}_{\mathcal{A}})$, an element a in $K(\mathcal{K}_{\mathcal{A}})$, a sequence of projections $\{p_{ij}\}$ in \mathcal{A} and a real bounded scalar sequence $\{\lambda_{ij}\}$ such that*

$$\sum_{i=1}^{\infty} \left(\sum_{j=1}^{l_i} p_{ij} \right) \otimes e_{ii} = 1 \quad \text{and} \quad uhu^* = \sum_{i=1}^{\infty} \left[\sum_{j=1}^{l_i} \lambda_{ij} p_{ij} \right] \otimes e_{ii} + a,$$

where p_{ij} ($i \leq j \leq l_i$) are mutually orthogonal for each fixed i , and a can be chosen with an arbitrarily small norm.

Proof of Theorem (4.3). Since \mathcal{A} is σ -unital and both \mathcal{A} and $M(\mathcal{A})$ have FS, by [21, 3.1] we can find mutually orthogonal projections p_i in \mathcal{A} with $\sum_{i=1}^{\infty} p_i = 1$, a real bounded scalar sequence

$\{\lambda_i\}$ and an element b in \mathcal{A} with arbitrarily small norm such that $h = \sum_{i=1}^{\infty} \lambda_i p_i + b$. Let $f_n = \sum_{i=1}^n p_i$. Then $\{f_n\}$ is an increasing approximate identity consisting of projections. By the same argument as in [10, 2.4] we can find a unitary v in $M(\mathcal{A})$ such that $v \sim 1$, and

$$e_{m_1} \leq v f_{n_1} v^* \leq e_{m_2} \leq v f_{n_2} v^* \leq e_{m_3} \leq \cdots,$$

where $\{n_i\}$ and $\{m_i\}$ are increasing sequences. Since

$$v \left(\sum_{j=n_{i-1}+1}^{n_i} p_j \right) v^* = (v f_{n_i} v^* - e_{m_i}) + (e_{m_i} - v f_{n_{i-1}} v^*)$$

(where $f_{n_0} = 0$), by the same arguments in the proof of Theorem (4.1) we can find a unitary w_i of $[v(f_{n_i} - f_{n_{i-1}})v^*]M(\mathcal{A})[v(f_{n_i} - f_{n_{i-1}})v^*]$ path connected to the identity $v(f_{n_i} - f_{n_{i-1}})v^*$ such that

$$w_i v \left(\sum_{j=n_{i-1}+1}^{n_i} p_j \right) v^* w_i^* = \sum_{j=n_{i-1}+1}^{n_i} w_i v p'_j v^* w_i^* + \sum_{j=n_{i-1}+1}^{n_i} w_i v p''_j v^* w_i^*,$$

where

$$p'_i + p''_i = p_i, \quad r_i = \sum_{j=n_{i-1}+1}^{n_i} w_i v p'_j v^* w_i^* = v f_{n_i} v^* - e_{m_i} \quad \text{and}$$

$$r'_i = \sum_{j=n_{i-1}+1}^{n_i} w_i v p''_j v^* w_i^* = e_{m_i} - v f_{n_{i-1}} v^*$$

Let $w = \sum_{i=1}^{\infty} w_i$. Then w is a unitary in $M(\mathcal{A})$ such that w is connected to the identity by a path of unitaries, where the path is continuous in the strict topology. Since $r_j + r'_{j+1} \leq e_{m_{j+1}} - e_{m_j}$, by the same arguments in the proof of Theorem (4.1), we obtain a unitary w'_j of $(e_{m_{j+1}} - e_{m_j})M(\mathcal{A})(e_{m_{j+1}} - e_{m_j})$ path connected to the identity $e_{m_{j+1}} - e_{m_j}$ such that

$$w'_j (r_j + r'_{j+1}) w_j^* = \sum_{i=m_j+1}^{m_{j+1}} \sum_{j=1}^{l_i} p_{ij},$$

where $\{p_{ij}: 1 \leq j \leq l_i\}$ is a set of mutually orthogonal subprojections in $(e_i - e_{i-1})\mathcal{A}(e_i - e_{i-1})$.

Define $w' = \sum_{i=1}^{\infty} w'_i$. Then w' is a unitary in $M(\mathcal{A})$ such that w' is path connected to the identity, where the path is continuous in the strict topology. Set $u = w' w v$. Then u is path connected to the

identity, where the path is continuous in the strict topology. It is easily verified that uhu^* has a desired form. (Notice that $\{\lambda_i\}$ is equal to $\{\lambda_{ij}\}$ as sets.) \square

4.5. REMARKS. (i) The condition “ $M(\mathcal{A})$ has FS” in the hypotheses of Theorem (4.3) and Corollary (4.4) has been studied in [5], [21] and [24]. Actually many multiplier algebras have the FS property.

(ii) Several applications of the results in this note have been given in the author’s subsequent papers [24, Part II, III, IV].

Acknowledgment. The author wishes to thank the referee for pointing out an inaccuracy in Theorem (4.1) in the primary manuscript of this paper.

REFERENCES

- [1] C. A. Akemann, G. K. Pedersen and J. Tomiyama, *Multipliers of C^* -algebras*, J. Funct. Anal., **13** (1973), 277–301.
- [2] J. Bunce and J. Deddens, *A family of simple C^* -algebras related to weighted shift operators*, J. Funct. Anal., **19** (1975), 13–24.
- [3] B. Blackadar, *Notes on the structure of projections in simple C^* -algebras*, Semesterbericht Funktionalanalysis, W82, Tübingen, March 1983.
- [4] ———, *K -Theory for Operator Algebras*, Springer-Verlag, New York, Berlin, Heidelberg, London, Paris, Tokyo, 1987.
- [5] L. G. Brown and G. K. Pedersen, *C^* -algebras of real rank zero*, preprint, 1989.
- [6] L. G. Brown, *Extensions of AF algebras: the projection lifting problem*, Operator Algebras and Applications, Proc. Symp. Pure Math., **38** (1981), Part I, 175–176.
- [7] R. Busby, *Double centralizers and extensions of C^* -algebras*, Trans. Amer. Math. Soc., **132** (1968), 79–99.
- [8] J. Cuntz and N. Higson, *Kuiper’s theorem for Hilbert modules*, Operator Alg. and Math. Phys., Vol. 62, Proc. of a Summer Conference June 17–21, 1985.
- [9] M.-D. Choi and G. A. Elliott, *Density of the self-adjoint elements with finite spectrum in an irrational rotation C^* -algebra*, Math. Scand., (to appear).
- [10] G. A. Elliott, *Derivations of matroid C^* -algebras II*, Ann. of Math., **100** (1974), 407–422.
- [11] K. Grove and G. K. Pedersen, *Diagonalizing matrices over $C(X)$* , J. Funct. Anal., **59** (1984), 65–89.
- [12] P. R. Halmos, *A Hilbert Space Problem Book*, second edition, Springer-Verlag New York, Heidelberg, Berlin, 1982.
- [13] R. V. Kadison, *Diagonalizing matrices over operator algebras*, Bull. Amer. Math. Soc., **8** (1983), 84–86.
- [14] ———, *Diagonalizing matrices*, Amer. J. Math., **106** (1984), 1451–1468.
- [15] G. Kasparov, *Hilbert C^* -modules: Theorems of Stinespring and Voiculescu*, J. Operator Theory, **4** (1980), 133–150.
- [16] G. K. Pedersen, *The linear span of projections in simple C^* -algebras*, J. Operator Theory, **4** (1980), 289–296.
- [17] M. A. Reiffel, *The homotopy groups of the unitary groups of non-commutative tori*, J. Operator Theory, **17** (1987), 237–254.

- [18] M. A. Reiffel, *Non-stable K -theory and non-commutative tori*, Contemporary Math., **62** (1987), 267–279.
- [19] —, *Dimension and stable rank in the K -theory of C^* -algebras*, Proc. London Math. Soc., **46** (1983), 301–333.
- [20] H. Weyl, *Über beschränkte quadratischen formen deren differenz vollstetig ist*, Rend. Circ. Mat. Palermo, **27** (1909), 373–392.
- [21] S. Zhang, *K_1 -groups, quasidiagonality and interpolation by multiplier projections*, Trans. Amer. Math. Soc., to appear.
- [22] —, *A Riesz decomposition property and ideal structure of multiplier algebras*, J. Operator Theory, (to appear).
- [23] —, *On the structure of projections and ideals of corona algebras*, Canad. J. Math., **41** (1989), 721–742.
- [24] —, *C^* -algebras with real rank zero and the internal structure of their corona and multiplier algebras*, Part I, II, IV, preprints. Part III to appear in Canad. J. Math.
- [25] —, *A property of purely infinite simple C^* -algebras*, Proc. Amer. Math. Soc., to appear.

Received November 1, 1988 and in revised form September 28, 1989.

UNIVERSITY OF KANSAS
LAWRENCE, KS 66045-2142