

ENDOSCOPIC GROUPS AND BASE CHANGE \mathbf{C}/\mathbf{R}

D. SHELSTAD

We consider a real reductive group G with complex points $G(\mathbf{C})$, Galois automorphism σ , and real points $G(\mathbf{R}) = \{g \in G(\mathbf{C}) : \sigma(g) = g\}$. In general, an irreducible admissible representation Π of $G(\mathbf{C})$ equivalent to its Galois conjugate $\Pi \circ \sigma$ need not be a lift from $G(\mathbf{R})$, even if G is quasi-split over \mathbf{R} . Following the results of L -indistinguishability we might expect this phenomenon to be related to the fact that σ -twisted conjugacy on $G(\mathbf{C})$ need not be “stable”, and therefore attempt to match the various “unstable” combinations of σ -twisted orbital integrals on $G(\mathbf{C})$ with stable orbital integrals on certain groups $H(\mathbf{R})$. The principle of functoriality in the L -group would then suggest, with reservations in the nontempered case, a relation between the σ -twisted characters of representations of $G(\mathbf{C})$ fixed up to equivalence by σ and the “dual lifts” to $G(\mathbf{C})$ of stable characters on the groups $H(\mathbf{R})$.

In this paper we define the relevant groups $H \dots$ they turn out to be the endoscopic groups from L -indistinguishability... and prove a matching theorem for orbital integrals. As a preliminary to the proposed dual liftings of characters we also study the “factoring” of Galois-invariant Langlands parameters for $G(\mathbf{C})$.

1. Introduction. We begin with two simple examples. Let $G(\mathbf{C}) = \mathbf{C}^\times$ and $\sigma(z) = \bar{z}^{-1}$, $z \in \mathbf{C}^\times$, so that $G(\mathbf{R}) = \{g \in G(\mathbf{C}) : \sigma(g) = g\}$ is the unit circle in \mathbf{C}^\times . A quasicharacter on \mathbf{C}^\times fixed by σ , i.e., trivial on the positive reals, need not be of the form $z \rightarrow \chi(z\sigma(z)) = \chi(z/\bar{z})$, with χ a character on the unit circle. At the same time $z \in \mathbf{C}^\times$ is stably σ -conjugate to $-z$, but not σ -conjugate (see [Sh6] for definitions). Let $f \in C_c^\infty(\mathbf{C}^\times)$ and write $f(r, \theta)$ for $f(re^{i\theta})$. Set $H_1 = H_2 = G$, so that $H_1(\mathbf{R}) = S^1$. Let

$$f_1(e^{i\theta}) = \frac{1}{2} \int_0^\infty (f(r, \theta/2) + f(r, \theta/2 + \pi)) dr/r$$

and

$$f_2(e^{i\theta}) = \frac{1}{2} e^{i\theta/2} \int_0^\infty (f(r, \theta/2) - f(r, \theta/2 + \pi)) dr/r$$

for $-\pi < \theta < \pi$. Then both f_1 and f_2 extend smoothly to S^1 . If χ is a character on S^1 then $f \rightarrow \int_{-\pi}^\pi \chi(e^{i\theta}) f_1(e^{i\theta}) d\theta$ is a distribution on \mathbf{C}^\times representing the usual lift of χ to $G(\mathbf{C})$, i.e., representing the quasicharacter $z \rightarrow \chi(z\sigma(z))$. On the other hand, $f \rightarrow \int_{-\pi}^\pi \chi(e^{i\theta}) f_2(e^{i\theta}) d\theta$ lifts χ to the quasicharacter $z = re^{i\theta} \rightarrow \chi(z\sigma(z))e^{i\theta}$. We have therefore recovered the remaining Galois-invariant quasicharacters on \mathbf{C}^\times .

For a general group, however, there are difficulties more akin to those for L -indistinguishability. Consider $G = \mathrm{SL}_2$. Let

$$H(\mathbf{R}) = \left\{ r(\theta) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \right\}.$$

Note that if $\theta \not\equiv 0 \pmod{\pi}$ then $r(\theta)$ and $r(\theta + \pi)$ are stably σ -conjugate in $G(\mathbf{C})$ but not σ -conjugate (see [Sh6, Lemma 2.5.2]). For $f \in C_c^\infty(\mathrm{SL}_2(\mathbf{C}))$, define

$$f_H(r(\theta)) = e^{i\theta/2}(e^{i\theta} - e^{-i\theta})(\Phi_f^\sigma(\theta/2) + \Phi_f^\sigma(\theta/2 + \pi)),$$

for $-\pi < \theta < \pi$, where

$$\Phi_f^\sigma(\theta) = \int_{G(\mathbf{C})/H(\mathbf{R})} f(\sigma(g)r(\theta)g^{-1}) \frac{dg}{d\theta},$$

dg denoting a Haar measure on $G(\mathbf{C}) = \mathrm{SL}_2(\mathbf{C})$. It can be shown that f_H extends to a C^∞ function on $H(\mathbf{R})$. Then $f \rightarrow \int_{H(\mathbf{R})} \chi f_H$ is a distribution on $\mathrm{SL}_2(\mathbf{C})$ (see [Sh6, §5.4] for an explicit formula). L. Clozel has shown that this distribution is, up to a constant, the twisted character of a Galois-fixed equivalence class of representations of $\mathrm{SL}_2(\mathbf{C})$. It is easily verified that all such classes of (irreducible, admissible) representations of $\mathrm{SL}_2(\mathbf{C})$ which are not lifts from $\mathrm{SL}_2(\mathbf{R})$ are lifts in this way.

Returning to the general problem, we find it convenient to consider $G(\mathbf{C})$ as the group of real points on a group \tilde{G} , and σ as the restriction to $\tilde{G}(\mathbf{R})$ of an algebraic automorphism α of \tilde{G} (cf. §2). Also, since (\tilde{G}, α) is our starting point, rather than G itself, we may as well assume that G is quasi-split over \mathbf{R} .

In this paper we will be concerned with the matchings for α -twisted orbital integrals on $\tilde{G}(\mathbf{R})$; this includes the problem of determining what it is they should match. Theorem 7.1 is our main result, and §§2 to 6 are preparation for it. Also, as both a check on our definitions and a preliminary to the proposed dual liftings, we will consider the question of “factoring” Galois-invariant Langlands parameters for $G(\mathbf{C})$ or, equivalently [L1] α -invariant parameters for $\tilde{G}(\mathbf{R})$. Theorem 8.1 is the main result.

In [Sh6] we started a study of the matching problem for α -twisted orbital integrals. We found that, despite various “technical” difficulties, the jump formulas for twisted orbital integrals on $\tilde{G}(\mathbf{R})$ are closely related to those for ordinary orbital integrals on $G(\mathbf{R})$. Making convenient technical assumptions, we then put together a matching theorem involving the endoscopic groups from L -indistinguishability. In this paper we start afresh, making none of the technical assumptions of [Sh6]. We first define

the notion of *endoscopic group* for (\tilde{G}, α) . This turns out to be the same as the notion of endoscopic group in L -indistinguishability [L3], [Sh4]. However, there is new information in the data for an endoscopic group H for (\tilde{G}, α) and it is this information which allows us to formulate a matching theorem without the assumption (4.3.2) of [Sh6]. Moreover in relating the embeddings ${}^L H \hookrightarrow {}^L \tilde{G}$ relevant to our present problem to the embeddings ${}^L H \hookrightarrow {}^L G$ from L -indistinguishability we find a remarkable quasicharacter on $\tilde{H}(\mathbf{R}) \simeq H(\mathbf{C})$ which allows us to dispense with the “cross-section for the norm” in [Sh6] (cf. Lemma 6.4).

As always, the twisted orbital integrals must be normalized. The normalization factors will be written in a form suitable for global applications [L3] and, more specifically, in a form to reflect the connection with L -indistinguishability for real groups. The proof of Theorem 7.1 itself relies heavily on the proof of the matching theorem for L -indistinguishability (see [Sh5] for an outline of the latter proof).

We will follow the notation of [Sh1]–[Sh7] as closely as possible, especially with respect to L -group data. However, we now write $G(\mathbf{C})$ and $G(\mathbf{R})$ in place of \mathbf{G} and G . The definitions in this paper may be presented in greater generality (cf. [Sh7]); in the general case there is no such intimate tie with L -indistinguishability.

2. The groups G , \tilde{G} and the automorphism α . Let G be a connected reductive linear algebraic group defined over \mathbf{R} . Assume that G is quasi-split over \mathbf{R} . In fixing the usual L -group data, we take G itself for G^* , a quasi-split inner form of G , and the identity map for ψ , an inner twist from G to G^* . Then B^* will be a Borel subgroup over \mathbf{R} in G , and T^* a maximal torus over \mathbf{R} in B^* . We form the dual $({}^L G^0, {}^L B^0, {}^L T^0, \{X_r\})$ with $r \in \Sigma({}^L B^0, {}^L T^0)$, the set of simple roots of ${}^L T^0$ in ${}^L B^0$. In fact it will be convenient to have fixed a root vector X_r , for any root r of ${}^L T^0$ in ${}^L G^0$. We therefore fix a Chevalley basis and take for $\{X_r, r \in \Sigma({}^L B^0, {}^L T^0)\}$ the vectors so provided. Then ${}^L G = {}^L G^0 \rtimes W$, with σ_G denoting the action of $1 \times \sigma \in W$ on ${}^L G^0$. See [Sh 3, 4, or 5] for further explanation of the notation.

Let \tilde{G} be the group obtained from G by restriction of scalars from \mathbf{C} to \mathbf{R} . We realize \tilde{G} as $G \times G$ with Galois automorphism $\sigma_{\tilde{G}}: (x, y) \rightarrow (\sigma_G(y), \sigma_G(x))$. Then $\tilde{B}^* = B^* \times B^*$ will be the distinguished Borel subgroup defined over \mathbf{R} and $\tilde{T}^* = T^* \times T^*$. We realize the L -group ${}^L \tilde{G}$ of \tilde{G} as follows. Set ${}^L \tilde{G}^0 = {}^L G^0 \times {}^L G^0$, ${}^L \tilde{B}^0 = {}^L B^0 \times {}^L B^0$, ${}^L \tilde{T}^0 = {}^L T^0 \times {}^L T^0$, $X_{(r,r')} = (X_r, X_{r'})$ for all roots r, r' of ${}^L T^0$ in ${}^L G^0$, and define $\sigma_{\tilde{G}}: {}^L \tilde{G}^0 \rightarrow {}^L \tilde{G}^0$ by $\sigma_{\tilde{G}}(g, h) = (\sigma_G(h), \sigma_G(g))$, $g, h \in {}^L G^0$. Then ${}^L \tilde{G} = {}^L \tilde{G}^0 \rtimes W$, with $\mathbf{C}^x \times 1$ acting trivially and $1 \times \sigma$ by $\sigma_{\tilde{G}}$.

Let $\alpha: \tilde{G} \rightarrow \tilde{G}$ be the automorphism $(x, y) \rightarrow (y, x)$. We take the standard dual automorphism (cf. [Sh7]) of α , and denote it by α also. Thus:

$$\alpha((g, h) \times w) = (h, g) \times w, \quad g, h \in {}^L G^0, w \in W.$$

3. Endoscopic groups for (\tilde{G}, α) . The following is a special case of the definitions in [Sh7]. Let $s \in {}^L \tilde{G}^0$. Then we set $N(s) = s\alpha(s)$, $\text{Cent}(N(s), {}^L \tilde{G}^0) = \{g \in {}^L \tilde{G}^0: g^{-1}N(s)g = N(s)\}$ and $\text{Cent}_\alpha(s, {}^L \tilde{G}^0) = \{g \in {}^L \tilde{G}^0: g^{-1}s\alpha(g) = s\}$. Call s α -semisimple if $\text{Cent}_\alpha(s, {}^L \tilde{G}^0)$ is reductive. In §4 we will observe that s is α -semisimple if and only if $N(s)$ is semisimple (cf. Lemma 4.2). Let \tilde{Z}^W be the group of W -invariants in the center of ${}^L \tilde{G}^0$. Thus $\tilde{Z}^W = {}^L \tilde{G}^0 \cap \text{Center}({}^L \tilde{G}) = \{(g, \sigma_G(g)) \times 1 \times 1: g \in \text{Center}({}^L G^0)\}$. Also

$$\text{Cent}_\alpha(sz, {}^L \tilde{G}^0) = \text{Cent}_\alpha(s, {}^L \tilde{G}^0), \quad s \in {}^L \tilde{G}^0, z \in \tilde{Z}^W.$$

We will now use s to denote a coset of \tilde{Z}^W in ${}^L \tilde{G}^0$ and $\text{Cent}_\alpha(s, {}^L \tilde{G}^0)$ to denote $\text{Cent}_\alpha(a, {}^L \tilde{G}^0)$ for a in the coset s . Following [Sh7], we consider tuples

$$(s, {}^L H_s^0, {}^L B_s^0, {}^L T_s^0, \{Y\}, \rho_s)$$

where

- (i) $s \in {}^L \tilde{G}^0$ is a coset of \tilde{Z}^W consisting of α -semisimple elements,
- (ii) ${}^L H_s^0 = (\text{Cent}_\alpha(s, {}^L \tilde{G}^0))^0$,
- (iii) ${}^L B_s^0$ is a Borel subgroup of ${}^L H_s^0$,
- (iv) ${}^L T_s^0 \subset {}^L B_s^0$ is a maximal torus in ${}^L H_s^0$,
- (v) $\{Y\}$ is a set of root vectors for the simple roots of ${}^L T_s^0$ in ${}^L B_s^0$,
- (vi) $\rho_s: W \rightarrow \text{Aut}({}^L H_s^0, {}^L B_s^0, {}^L T_s^0, \{Y\})$ is a homomorphism which factors through $\text{Gal}(\mathbf{C}/\mathbf{R})$ and is “realized in $\text{Cent}_\alpha(s, {}^L \tilde{G})$ ”, i.e. $\rho_s(w) = \text{ad } n(w)|_{{}^L H_s^0}$, $w \in W$, for some $n(w) \in {}^L \tilde{G}^0 \times w$ such that $n(w)^{-1}\alpha n(w) = a$ for each a in the coset s .

Let ${}^L H_s = {}^L H_s^0 \rtimes W$, the action of W on ${}^L H_s^0$ being that defined by ρ_s . Often we will write σ_s for the automorphism $\rho_s(1 \times \sigma)$, and abbreviate $(s, {}^L H_s^0, \dots, \rho_s)$ by $(s, {}^L H_s)$.

Two tuples

$$(s, {}^L H_s^0, {}^L B_s^0, {}^L T_s^0, \{Y\}, \rho_s) \quad \text{and} \quad (s', {}^L H_{s'}^0, {}^L B_{s'}^0, {}^L T_{s'}^0, \{Y'\}, \rho_{s'})$$

are *equivalent* if there exists $g \in {}^L \tilde{G}^0$ such that ${}^L H_{s'}^0 = g^{-1}{}^L H_s^0 g$, ${}^L B_{s'}^0 = g^{-1}{}^L B_s^0 g$, ${}^L T_{s'}^0 = g^{-1}{}^L T_s^0 g$, $\{Y'\} = \{\text{Ad } g^{-1}(Y)\}$ and if $n(w) \in \text{Cent}_\alpha(s, {}^L \tilde{G})$ realizes $\rho_s(w)$ then $g^{-1}n(w)g$ lies in $\text{Cent}_\alpha(s', {}^L \tilde{G})$ and realizes $\rho_{s'}(w)$, $w \in W$. The set of all equivalence classes will be denoted

$\mathfrak{S}(\tilde{G}, \alpha)$. Using the results of the next section and Lemma 2.3.3 of [Sh4], we may show that $\mathfrak{S}(\tilde{G}, \alpha)$ is a finite set. Since this fact will not be needed we omit the proof.

Finally, we call a quasi-split group H over \mathbf{R} an *endoscopic group* for (\tilde{G}, α) if some ${}^L H_s$ as above is an L -group for H .

4. The relation between endoscopic groups for (\tilde{G}, α) and endoscopic groups for G . By the endoscopic groups for G we mean the groups “ H ” of [Sh4], i.e. essentially the groups of [L1]. The set $\mathfrak{S}(G)$, or $\mathfrak{S}(G, 1)$ in the more general notation of [Sh7], and the tuples used in its definition will be taken from [Sh4] (... there is a small difference in the definitions of [L3]).

We embed ${}^L G$ “diagonally” in ${}^L \tilde{G}$, i.e. by the map $g \times w \rightarrow (g, g) \times w$, $g \in {}^L G^0$, $w \in W$, and will frequently *identify ${}^L G$ with its image in ${}^L \tilde{G}$* . As in [Sh4], Z^W will denote the set of W -invariants in the center of ${}^L G^0$.

By an α -conjugacy class in ${}^L \tilde{G}^0$, we will mean a set $\{g^{-1} \alpha(g); g \in {}^L \tilde{G}^0\}$, where $a \in {}^L \tilde{G}^0$.

LEMMA 4.1.

- (i) *Each α -conjugacy class in ${}^L \tilde{G}^0$ contains an element of the form $(x, 1)$, $x \in {}^L G^0$.*
- (ii) *For $x \in {}^L G^0$, $\text{Cent}_\alpha((x, 1), {}^L \tilde{G}^0) = \text{Cent}(x, {}^L G^0)$.*

Here, of course, $\text{Cent}(x, {}^L G^0)$ has been identified with its image in ${}^L \tilde{G}$ under the diagonal map.

Proof. Let $a = (g_1, g_2) \in {}^L \tilde{G}^0$, $g = (1, g_2)$. Then $g^{-1} \alpha(g) = (1, g_2^{-1})(g_1, g_2)(g_2, 1) = (g_1 g_2, 1)$, so that (i) is proved. (ii) is also a simple calculation.

LEMMA 4.2. *$a \in {}^L \tilde{G}^0$ is α -semisimple if and only if $N(a) = \alpha(a)$ is semisimple.*

Proof. Let $a \in {}^L \tilde{G}^0$. Choose $g \in {}^L \tilde{G}^0$ such that $g^{-1} \alpha(g) = (x, 1)$, for suitable $x \in {}^L G^0$. Then

$$\text{Cent}_\alpha(a, {}^L \tilde{G}^0) = g \text{Cent}_\alpha((x, 1), {}^L \tilde{G}^0) g^{-1} = g \text{Cent}(x, {}^L G^0) g^{-1}.$$

On the other hand, $N(a) = g(x, x)g^{-1}$, so that

$$\text{Cent}(N(a), {}^L \tilde{G}^0) = g(\text{Cent}(x, {}^L G^0) \times \text{Cent}(x, {}^L G^0))g^{-1}.$$

The lemma then follows from standard facts.

LEMMA 4.3. *Let s be a coset of \tilde{Z}^W in ${}^L\tilde{G}^0$ consisting of α -semisimple elements. Then there exists $g \in {}^L\tilde{G}^0$ such that $s' = g^{-1}s\alpha(g)$ has the property that $\{\alpha\alpha(a) : a \in s'\}$ is contained in ${}^LG^0$. Then $\{\alpha\alpha(a) : a \in s'\}$ is contained in a unique coset of Z^W in ${}^LG^0$. This coset, to be denoted $N(s')$, consists of semisimple elements.*

Proof. Let $a \in s$. Choose $g \in {}^L\tilde{G}^0$ such that $g^{-1}a\alpha(g) = (x, 1)$, where $x \in {}^LG^0$ is semisimple. Let $s' = (x, 1)\tilde{Z}^W$. Then if $b \in s'$, $b\alpha(b) = (x, x)(z\sigma_G(z), z\sigma_G(z))$, for some $z \in \text{Cent}({}^LG^0)$. Thus, with our identifications, $b\alpha(b) \in xZ^W$, a coset of Z^W in ${}^LG^0$ consisting of semisimple elements. The rest is clear.

LEMMA 4.4. *Each element of $\mathfrak{S}(\tilde{G}, \alpha)$ has a representative $(s, {}^LH_s)$ such that $(N(s), {}^LH_s)$ is a representative for an element of $\mathfrak{S}(G)$ i.e. such that $\{\alpha\alpha(a) : a \in s\}$ is contained in ${}^LG^0$ (...so that $N(s)$ is defined), ${}^LH_s^0$ coincides with $(\text{Cent}(N(s), {}^LG^0))^0$, and ρ_s is “realized in $\text{Cent}(N(s), {}^LG)$.”*

Proof. We may take $s = (x, 1)\tilde{Z}^W$, some $x \in {}^LT^0$. Then $N(s) = xZ^W$ and $\text{Cent}_\alpha(s, {}^L\tilde{G}^0) = \text{Cent}(N(s), {}^LG^0)$. We may also assume that ${}^LT_s^0 = {}^LT^0$, ${}^LB_s^0 = {}^LB^0 \cap {}^LH_s^0$ (... ${}^LT^0$ and ${}^LB^0$ being identified with their images in ${}^L\tilde{G}^0$) and that $\{Y\} = \{X_r : r \in \Sigma({}^LB^0 \cap {}^LH_s^0, {}^LT^0)\}$. Then ρ_s is a homomorphism of W into $\text{Aut}({}^LH_s^0, {}^LB^0 \cap {}^LH_s^0, {}^LT^0, \{Y\})$. Suppose that $\rho_s(w) = \text{ad } n(w) |_{\mathfrak{L}_{H_s^0}}$, where $n(w) \in {}^L\tilde{G}^0 \times w$ satisfies $n(w)^{-1}(x, 1)\alpha(n(w)) = (x, 1)$ (cf. (vi) in §3). Then $n(w)^{-1}(x, x)n(w) = (x, x)$. Also, if $n(w) = (n_1(w), n_2(w)) \times w$ then calculation shows that for $w \in \mathbf{C}^x \times 1$ we have $n_1(w) = n_2(w)$ lies in the center of ${}^LH_s^0$ and for $w = 1 \times \sigma$ we have $n_1(w) = xn_2(w)$. Thus for all $w \in W$, $\rho_s(w) = \text{ad } m(w) |_{\mathfrak{L}_{H_s^0}}$ where $m(w) = (n_1(w), n_1(w)) \times w \in {}^LG$. Also, $m(w)$ centralizes (x, x) . Thus ρ_s is “realized in $\text{Cent}(N(s), {}^LG)$ ” and the lemma is proved.

LEMMA 4.5. *The correspondence in Lemma 4.4 induces a map*

$$\mathfrak{N} : \mathfrak{S}(\tilde{G}, \alpha) \rightarrow \mathfrak{S}(G).$$

Proof. We have to show that if $(s, {}^LH_s)$ and $(s', {}^LH_{s'})$ are as in Lemma 4.4, representing the same element of $\mathfrak{S}(\tilde{G}, \alpha)$, then the 5-tuples defining LH_s and ${}^LH_{s'}$ are conjugate under ${}^LG^0$. They are conjugate under ${}^L\tilde{G}^0$, by definition. It is easily checked that this conjugation may be replaced by one from ${}^LG^0$.

The map \mathcal{U} need not be injective, as the example that G is a compact torus shows. However \mathcal{U} does have finite fibers (which implies that $\mathfrak{S}(\tilde{G}, \alpha)$ is finite, as asserted in the last section). Reversing the construction in the proof of Lemma 4.4 shows that \mathcal{U} is surjective.

5. Allowed embeddings of ${}^L H_s$ in ${}^L \tilde{G}$. Fix an element of $\mathfrak{S}(\tilde{G}, \alpha)$, with representative $(s, {}^L H_s)$ chosen as in the proof of Lemma 4.4. In particular, $s = (x, 1)\tilde{Z}^W$, $x \in {}^L T^0$, and ${}^L H_s^0 = (\text{Cent}_{\alpha}(s, {}^L \tilde{G}^0))^0 = (\text{Cent}(N(s), {}^L G^0))^0$. We may further assume that ${}^L H_s^0$ is in standard position (cf. [Sh3, §2.2, Ex. 4.3.1]).

Suppose that $\xi: {}^L H_s \hookrightarrow {}^L G$ is an admissible embedding, as in L -indistinguishability [L1], [Sh3]. Here we regard ${}^L H_s^0$ as a subgroup of ${}^L G^0$ yet to be embedded diagonally in ${}^L \tilde{G}^0$, and assume that $\xi|_{{}^L H_s^0}$ is the inclusion map. The “diagonal” embedding of ${}^L G$ in ${}^L \tilde{G}$ then yields an embedding of ${}^L H_s$ in ${}^L \tilde{G}$, again denoted ξ . Explicitly, ξ is of the form:

$$\begin{aligned} \xi(h \times 1 \times 1) &= (h, h) \times 1 \times 1, & h \in {}^L H_s^0, \\ \xi(1 \times z \times 1) &= (\xi_0(z), \xi_0(z)) \times z \times 1, & z \in \mathbf{C}^x, \end{aligned}$$

where $\xi_0: \mathbf{C}^x \rightarrow \text{Cent}({}^L H_s^0)$ is a homomorphism satisfying $\xi_0(\bar{z}) = \sigma_s(\xi_0(z))$, $z \in \mathbf{C}^x$, and

$$\xi(1 \times 1 \times \sigma) = (n_0, n_0) \times 1 \times \sigma,$$

where $n_0 \in {}^L G^0$ normalizes ${}^L T^0$, $n_0 \sigma_G(n_0) = \xi_0(-1)$ and $n_0 \times 1 \times \sigma \in {}^L G$ acts on ${}^L H_s^0$ as $\sigma_s = \rho_s(1 \times \sigma)$. It follows immediately that $\xi({}^L H_s) \subset \text{Cent}(N(s), {}^L G)$. However, our present problem dictates (cf. §8) that we consider embeddings for which the image of ${}^L H_s$ is contained in $\text{Cent}_{\alpha}(s, {}^L \tilde{G})$. That this is a quite different condition is indicated even by the example that G is a compact torus.

DEFINITION 5.1. Let $(s, {}^L H_s)$ be a representative for an element of $\mathfrak{S}(\tilde{G}, \alpha)$. Then $\tilde{\xi}: {}^L H_s \hookrightarrow {}^L \tilde{G}$ is an allowed embedding if:

- (i) $\tilde{\xi}$ is an admissible homomorphism, i.e. $\tilde{\xi}$ is a homomorphism such that $\tilde{\xi}({}^L H_s^0 \times w) \subset {}^L \tilde{G}^0 \times w$, $w \in W$,
- (ii) on ${}^L H_s^0$, $\tilde{\xi}$ is the inclusion mapping, and
- (iii) $\tilde{\xi}({}^L H_s) \subset \text{Cent}_{\alpha}(s, {}^L \tilde{G})$.

We return to our choice $s = (x, 1)\tilde{Z}^W$, etc. Once again it is more convenient to regard ${}^L H_s^0$ as a subgroup of ${}^L G^0$ yet to be embedded

diagonally in ${}^L\tilde{G}^0$. Then an allowed embedding $\tilde{\xi}: {}^LH_s \hookrightarrow {}^L\tilde{G}$ is of the form:

$$\begin{aligned} \tilde{\xi}(h \times 1 \times 1) &= (h, h) \times 1 \times 1, & h \in {}^LH_s^0, \\ \tilde{\xi}(1 \times z \times 1) &= (\tilde{\xi}_0(z), \tilde{\xi}_0(z)) \times z \times 1, & z \in \mathbf{C}^\times, \end{aligned}$$

where $\tilde{\xi}_0$ satisfies the same conditions as ξ_0 earlier, and

$$\tilde{\xi}(1 \times 1 \times \sigma) = (xm_0, m_0) \times 1 \times \sigma$$

where $m_0 \in {}^LG^0$ normalizes ${}^LT^0$, $xm_0\sigma_G(m_0) = \tilde{\xi}_0(-1)$, and $m_0 \times 1 \times \sigma \in {}^LG$ acts on ${}^LH_s^0$ as σ_s (... then also $xm_0 \times 1 \times \sigma$ acts on ${}^LH_s^0$ as σ_s , as we have already used in the proof of Lemma 4.2).

Let ${}^L\tilde{H}_s^0 = {}^LH_s^0 \times {}^LH_s^0$. We of course regard ${}^L\tilde{H}_s^0$ as a subgroup of ${}^L\tilde{G}^0$. Define an action of W on ${}^L\tilde{H}_s^0$ by requiring $\mathbf{C}^\times \times 1$ to act trivially and $1 \times \sigma$ to act by the automorphism $(h_1, h_2) \rightarrow (\sigma_s(h_2), \sigma_s(h_1))$. If LH_s is the L -group of H then ${}^L\tilde{H}_s$ is the L -group of $\tilde{H} = \text{Res}_{\mathbf{R}}^{\mathbf{C}} H$.

LEMMA 5.2. *Let $\tilde{\xi}$ be an allowed embedding of LH_s in ${}^L\tilde{G}$ and ξ be an admissible embedding of LH_s in ${}^LG \subset {}^L\tilde{G}$. Then*

$$\tilde{\xi}(h \times w) = a(w)\xi(h \times w), \quad h \in {}^LH_s^0, w \in W,$$

where $a(w)$ is a 1-cocycle of W in $\text{Cent}({}^L\tilde{H}_s^0)$.

Proof. This follows easily from our explicit description of $\tilde{\xi}$ and ξ . The details are omitted.

Suppose that $\tilde{\xi}, \tilde{\xi}'$ are both allowed embeddings of LH_s in ${}^L\tilde{G}$. Then $\tilde{\xi}'(w) = b(w)\tilde{\xi}(w)$, $w \in W$, where $w \rightarrow b(w)$ is a 1-cocycle of W in the center of ${}^LH_s^0$ embedded diagonally in ${}^L\tilde{G}^0$. We conclude then that the image of LH_s under an allowed embedding is independent of the choice of embedding; we write thus simply “Image LH_s .” Suppose next that $(s, {}^LH_s)$ and $(s', {}^LH_{s'})$ are equivalent in the sense of §3. Fix $g \in {}^L\tilde{G}^0$ as in the definition. Suppose that $\tilde{\xi}$ is an allowed embedding of LH_s in ${}^L\tilde{G}$. Then $ad g$ and $\tilde{\xi}$ determine an allowed embedding of ${}^LH_{s'}$ in ${}^L\tilde{G}$. We conclude then that there is an allowed embedding of LH_s in ${}^L\tilde{G}$ if and only if there is such an embedding of ${}^LH_{s'}$. Moreover, when embeddings exist we have $g^{-1}(\text{Image } {}^LH_s)g = \text{Image } {}^LH_{s'}$.

We defer a study of the existence of allowed embeddings. Recall, however, that if the center of ${}^LG^0$ is connected then LH_s embeds admissibly in LG [L1]. The proof of this result can be used to show also that there is an allowed embedding of LH_s in ${}^L\tilde{G}$.

6. Ingredients for the matching theorem. Fix an element of $\mathfrak{S}(\tilde{G}, \alpha)$ with representative $(s, {}^LH_s)$ satisfying $s = (x, 1)\tilde{Z}^W$, etc., as in the last section. We assume that $\tilde{\xi}: {}^LH_s \hookrightarrow {}^L\tilde{G}$ is an allowed embedding. The main

purpose of this section is to attach to $\tilde{\xi}$ normalizing factors to appear in the matching theorem of the next section. We will assume also that there is an admissible embedding of ${}^L H_s$ in ${}^L G$, say ξ . The choice of ξ will not affect the normalization factors (cf. Lemma 6.2), but we write individual terms in the factors in a way that involves ξ , in order to make clear the relation with the factors from L -indistinguishability.

Let H be an endoscopic group for (\tilde{G}, α) with L -group ${}^L H_s$. We fix a Borel subgroup B_H over \mathbf{R} containing the maximal torus T_H over \mathbf{R} , and assume that $X^*(T_H) = X_*({}^L T^0) = X^*(T^*)$ and that $\Sigma(B_H, T_H)$ is the dual of $\Sigma({}^L B_s^0, {}^L T^0)$. The group $\tilde{H} = \text{Res}_{\mathbf{R}}^{\mathbf{C}} H$ will also play a role. We set $\tilde{B}_H = B_H \times B_H$ and $\tilde{T}_H = T_H \times T_H$; ${}^L \tilde{H}_s$, which appeared in the last section, is an L -group for \tilde{H} .

Since H is also an endoscopic group for G we may invoke many of the definitions from L -indistinguishability (cf. [L1], [Sh4]). Let T be a maximal torus over \mathbf{R} in G . A pseudodiagonalization (p.d.) η of T is a map from T to T^* of the form $T \xrightarrow{\text{ad } x} T_0 \xrightarrow{\text{ad } m} T^*$, where $x \in \mathfrak{A}(T)$ [L1], $T_0 = xTx^{-1}$ is standard (i.e. the maximal R -split torus in T_0 lies in T^*) and m belongs to the Levi group attached to T_0 . Then $\sigma_{(T, \eta)}$ denotes the transfer, by η , of the Galois action on T to T^* , and to $X^*(T^*) = X_*({}^L T^0)$, $X_*(T^*) = X^*({}^L T^0)$ and ${}^L T^0 = X_*({}^L T^0) \otimes \mathbf{C}^x$.

The set $\mathfrak{T}_H(G) = \{(T, \eta) : \sigma_{(T, \eta)} \in \Omega({}^L H_s^0, {}^L T^0)\sigma_s\}$, where $\Omega({}^L H_s^0, {}^L T^0)$ denotes the Weyl group of $({}^L H_s^0, {}^L T^0)$, is the starting point for the definitions of [Sh4, §2.4]. We will use it again. First, because G is quasi-split over \mathbf{R} , for each maximal torus T' over \mathbf{R} in H there exists $h \in H(\mathbf{C})$ and $(T, \eta) \in \mathfrak{T}_H(G)$ such that $hT'h^{-1} = T_H$ and

$$X^*(T') \xrightarrow{\text{ad } h} X^*(T_H) = X^*(T^*) \xrightarrow{\eta^{-1}} X^*(T)$$

lifts to an isomorphism $i(h, \eta) : T' \rightarrow T$ over \mathbf{R} . We say that $\gamma' \in H(\mathbf{R})$ originates from $\gamma \in G(\mathbf{R})$ via (T, η) if γ' is the preimage of γ under some such map $i(h, \eta)$.

Recall that $s = (x, 1)\tilde{Z}^W$. Any element of this coset is of the form $a = (xz, \sigma_G(z))$, where z is in the center of ${}^L G^0$. But $a\alpha(a) = (xz\sigma_G(z), xz\sigma_G(z))$, an element of ${}^L T^0 = \text{Hom}(X^*({}^L T^0), \mathbf{C}^x)$. Also $\sigma_s(x) = x$. Thus $\{a\alpha(a) : a \in s\}$ defines a family of quasicharacters on $X^*({}^L T^0)$, each invariant under $\sigma_{(T, \eta)}$, for any $(T, \eta) \in \mathfrak{T}_H(G)$. Fix $(T, \eta) \in \mathfrak{T}_H(G)$. Then, on transfer to T via η , we get a family of quasicharacters on $X_*(T)$, each invariant under σ_T . On $X_*(T_{\text{sc}})$, the span of the coroots of T in G , these quasicharacters all coincide and so we have defined a single quasicharacter of the type used in L -indistinguishability (cf. [L1], also [Sh4, §2.4]). Moreover on $\{\lambda^\vee \in X_*(T) : \sigma_T \lambda^\vee = -\lambda^\vee\}$, the quasicharacters

coincide again. We therefore obtain a single character on

$$\left\{ \lambda^\vee \in X_*(T) : \sigma_T \lambda^\vee = -\lambda^\vee \right\} / \left\{ \mu^\vee - \sigma_T \mu^\vee : \mu^\vee \in X_*(T) \right\}$$

and thence by Tate-Nakayama duality, a character on $H^1(T) = H^1(\text{Gal}(\mathbf{C}/\mathbf{R}), T(\mathbf{C}))$. Unless otherwise indicated, κ will denote both the quasicharacter on $X_*(T_{\text{sc}})$ and the character on $H^1(T)$ attached to s and the pair $(T, \eta) \in \mathfrak{T}_H(G)$.

With G embedded diagonally in \tilde{G} , we have $\tilde{T} = \text{Res}_{\mathbf{R}}^{\mathbf{C}} T$ naturally embedded in \tilde{G} as $\text{Cent}(T, \tilde{G}) = T \times T$, for any maximal torus T over \mathbf{R} in G . The norm from \tilde{T} to T is obtained from the map $\tilde{T}(\mathbf{R}) \rightarrow T(\mathbf{R})$ defined by $\delta = (t, \sigma_G(t)) \rightarrow \delta\alpha(\delta) = (t\sigma_G(t), t\sigma_G(t))$. As in [Sh6] we regard the norm from \tilde{G} to G (... or from \tilde{T} to T) as an (injective) map from the set of stable regular α -semisimple twisted conjugacy classes in $\tilde{G}(\mathbf{R})$ (... or in $\tilde{T}(\mathbf{R})$) to the set of stable regular semisimple conjugacy classes in $G(\mathbf{R})$ (... or to $T(\mathbf{R})$). by Lemma 2.4.3(ii) of [Sh6] this norm from \tilde{G} to G can be recovered from the norms from \tilde{T} to T , as T ranges over the maximal tori over \mathbf{R} in G .

Note that if $\eta: T \rightarrow T^*$ is a p.d., then so is $\eta \times \eta: \tilde{T} \rightarrow \tilde{T}^*$. Thus we can use η to transfer data from \tilde{T} to \tilde{T}^* or from \tilde{T}^* to \tilde{T} .

We come then to the normalizing factors. The admissible embedding $\xi: {}^L H_s \hookrightarrow {}^L G$ has been fixed, and ${}^L H_s$ chosen to satisfy the conditions of [Sh3, Sh4]. We may therefore write $\xi = \xi(\mu^*, \lambda^*)$, for suitable $\mu^*, \lambda^* \in X_*({}^L T^0) \otimes \mathbf{C}$, and define the attached correction (quasi) characters $\Lambda_{(T, \eta)}$ on $T(\mathbf{R})$, for $(T, \eta) \in \mathfrak{T}_H(G)$. Although the notation does not reflect it, $\Lambda_{(T, \eta)}$ depends on the choice of ξ .

Since $\tilde{\xi}: {}^L H_s \hookrightarrow {}^L \tilde{G}$ has also been fixed, we have the 1-cocycle $a(w)$ of W in $\text{Center}({}^L \tilde{H}_s^0)$ from Lemma 5.2. A procedure in [L2] attaches to $a(w)$ a quasicharacter on $\tilde{H}(\mathbf{R})$. This quasicharacter determines a pair $(\tilde{\mu}_0, \tilde{\lambda}_0)$ of elements from $X^*(\tilde{T}_H) \otimes \mathbf{C} = X_*({}^L \tilde{T}^0) \otimes \mathbf{C}$. We may also recover $(\tilde{\mu}_0, \tilde{\lambda}_0)$ directly from the 1-cocycle $a(w)$. Thus define $\tilde{\mu}_0, \tilde{\lambda}_0$ by

$$\lambda^\vee(a(z \times 1)) = z^{\langle \tilde{\mu}_0, \lambda^\vee \rangle} \bar{z}^{\langle \sigma_s \tilde{\mu}_0, \lambda^\vee \rangle}, \quad z \in \mathbf{C}^x,$$

$$\lambda^\vee(a(1 \times \sigma)) = e^{2\pi i \langle \tilde{\lambda}_0, \lambda^\vee \rangle}$$

for $\lambda^\vee \in X^*({}^L \tilde{T}^0)$. Then $\tilde{\mu}_0$ is uniquely determined and $\tilde{\lambda}_0$ is uniquely determined modulo

$$X_*({}^L \tilde{T}^0) + \{ \tilde{\lambda} - \tilde{\sigma}_s \tilde{\lambda} : \tilde{\lambda} \in X_*({}^L \tilde{T}^0) \otimes \mathbf{C} \}.$$

Also

$$\tilde{\mu}_0 - \tilde{\sigma}_s \tilde{\mu}_0 \in X_*({}^L \tilde{T}^0), \quad 1/2(\tilde{\mu}_0 - \tilde{\sigma}_s \tilde{\mu}_0) \equiv \tilde{\lambda}_0 + \tilde{\sigma}_s \tilde{\lambda}_0 \pmod{X_*({}^L \tilde{T}^0)},$$

and

$$\langle \tilde{\mu}_0, \lambda^\vee \rangle = 0, \quad \langle \tilde{\lambda}_0, \lambda^\vee \rangle \in \mathbf{Z}$$

whenever λ^\vee lies in the span of the roots of ${}^L\tilde{T}^0$ in ${}^L\tilde{H}_s^0$ (cf. §9.1 of [Sh3]). Here we have used $\tilde{\sigma}_s$ to denote the action of $1 \times 1 \times \sigma \in {}^L\tilde{H}_s$.

Let $(T, \eta) \in \mathfrak{G}_H(G)$. Then on transferring $(\tilde{\mu}_0, \tilde{\lambda}_0)$ to \tilde{T} using η we obtain the data also denoted $(\tilde{\mu}_0, \tilde{\lambda}_0)$ for a quasicharacter on $\tilde{T}(\mathbf{R})$ (cf. [Sh3, §4.1]). This quasicharacter will be denoted $a_{(T,\eta)}$.

LEMMA 6.1.

$a_{(T,\eta)}$ is α -invariant.

Proof. We describe $a_{(T,\eta)}$ explicitly. Let $\delta = (t, \sigma_T(t)) \in \tilde{T}(\mathbf{R})$. Write t as $\exp X$, $X \in \text{Lie}(T(\mathbf{C})) = X_*(T) \otimes \mathbf{C}$. Then $\sigma_T(t) = \exp \sigma_T(\bar{X})$, where if $X = \sum_{i=1}^n \lambda_i^\vee \otimes z_i$ then $\sigma_T(\bar{X}) = \sum_{i=1}^n \sigma_T(\lambda_i^\vee) \otimes \bar{z}_i$. Because $a(\mathbf{C}^\times \times 1)$ lies in the diagonal subgroup of $\text{Center}({}^L\tilde{H}_s^0)$, as is evident from the form of the embeddings ξ and $\tilde{\xi}$ (cf. last section), we must have $\tilde{\mu}_0$ lying in the diagonal subspace of $X_*({}^L\tilde{T}^0) \otimes \mathbf{C} = (X_*({}^LT^0) \otimes \mathbf{C}) \times (X_*({}^LT^0) \otimes \mathbf{C})$. Thus we write $\tilde{\mu}_0$ as (μ_0, μ_0) , $\mu_0 \in X_*({}^LT^0) \otimes \mathbf{C}$. As usual, we transfer μ_0 to $X^*(T) \otimes \mathbf{C}$ via η without change in notation. Then

$$a_{(T,\eta)}(\delta) = e^{\mu_0(X + \sigma_T(\bar{X}))}.$$

Since $\alpha(\delta) = (\exp \sigma_T(\bar{X}), \exp X)$ it is now clear that $a_{(T,\eta)}(\alpha(\delta)) = a_{(T,\eta)}(\delta)$, and the lemma is proved.

Note that $a_{(T,\eta)}$ is uniquely determined by the class of $a(w)$ in $H^1(W, \text{Center}({}^L\tilde{H}_s^0))$, but is affected by a change in ξ or $\tilde{\xi}$. The dependence on $\tilde{\xi}$ of our normalization factors is to be expected; the dependence on ξ is not.

LEMMA 6.2. Fix $(T, \eta) \in \mathfrak{G}_H(G)$ and $\delta \in \tilde{T}(\mathbf{R})$. Then $\alpha_{(T,\eta)}(\delta)\Lambda_{(T,\eta)}(\delta\alpha(\delta))$ depends on $\tilde{\xi}$ alone.

Proof. The embedding ξ may be replaced only by $h \times w \rightarrow a_0(w)\xi(h \times w)$, where $a_0(w)$ is a 1-cocycle of W in the center of ${}^LH_s^0$ embedded diagonally in the center of ${}^L\tilde{H}_s^0$. Then $a(w)$ is replaced by $a_0(w)^{-1}a(w)$. The cocycle $a_0(w)$ defines first a quasicharacter χ on $H(\mathbf{R})$ and second a quasicharacter $\tilde{\chi}$ on $\tilde{H}(\mathbf{R})$. As before, we use η to transfer data and define quasicharacters $\chi_{(T,\eta)}$ on $T(\mathbf{R})$ and $\tilde{\chi}_{(T,\eta)}$ on $\tilde{T}(\mathbf{R})$. Since $\Lambda_{(T,\eta)}$ is replaced by $\chi_{(T,\eta)}\Lambda_{(T,\eta)}$ and $a_{(T,\eta)}$ by $\tilde{\chi}_{(T,\eta)}^{-1}a_{(T,\eta)}$, we have only to show that $\tilde{\chi}_{(T,\eta)}(\delta) = \chi_{(T,\eta)}(\delta\alpha(\delta))$. Define parameters $\mu_1, \lambda_1 \in X_*({}^LT^0) \otimes \mathbf{C}$ for χ as usual; use the same symbols for their transfer to $X^*(T) \otimes \mathbf{C}$

via η . For $\tilde{\chi}$ we can use parameters $\tilde{\mu}_1 = (\mu_1, \mu_1)$, $\tilde{\lambda}_1 = (\lambda_1, \lambda_1)$ in $X_*({}^L\tilde{T}^0) \otimes \mathbf{C}$ (...or $X^*(\tilde{T}) \otimes \mathbf{C}$, after transfer). Since $\tilde{\chi}$ is clearly α -invariant (see the last proof), we may take $\delta = (\exp X, \exp X)$, $X \in \text{Lie}(T(\mathbf{R}))$. Then $\tilde{\chi}(\delta) = e^{\langle 2\mu_1, X \rangle}$ and $\chi(\delta\alpha(\delta)) = \chi(\delta^2) = e^{\langle \mu_1, 2X \rangle}$, so that the lemma is proved.

The next lemma is simple but very useful (cf. proof of Lemma 6.4). Each element of $H^1(T)$ can be represented by a cocycle $\sigma \rightarrow \exp i\pi\lambda^\vee$, where $\lambda^\vee \in X_*(T)$ and $\sigma_T\lambda^\vee = -\lambda^\vee$. We will use $\exp i\pi\lambda^\vee$ to denote this cocycle *and* its class in $H^1(T)$; of course, $\exp i\pi\lambda^\vee$ also denotes an element of $T(\mathbf{R}) \subset \tilde{T}(\mathbf{R})$. Recall that to $(T, \eta) \in \mathfrak{S}_H(G)$ and our fundamental datum $s = (x, 1)\tilde{Z}^W$ we have attached a character κ on $H^1(T)$.

LEMMA 6.3.

$$a_{(T,\eta)}(\exp i\pi\lambda^\vee) = \kappa(\exp i\pi\lambda^\vee)$$

for all $\lambda^\vee \in X_*(T)$ such that $\sigma_T\lambda^\vee = -\lambda^\vee$.

Note that the left side alone appears to depend on the choice of ξ and $\tilde{\xi}$. However a quasicharacter $\tilde{\chi}$ as in the last proof annihilates $\exp i\pi\lambda^\vee$, if $\lambda^\vee \in X_*(T)$ and $\sigma_T\lambda^\vee = -\lambda^\vee$. Indeed we then have $i\pi\lambda^\vee \in \text{Lie}(T(\mathbf{R}))$, so that $\tilde{\chi}(\exp i\pi\lambda^\vee) = e^{2\pi i\langle \mu_1, \lambda^\vee \rangle} = 1$, since $\frac{1}{2}(\mu_1 - \sigma_T\mu_1) \equiv (\lambda_1 + \sigma_T\lambda_1) \pmod{X^*(T)}$ implies that $\langle \frac{1}{2}(\mu_1 - \sigma_T\mu_1), \lambda^\vee \rangle = \langle \mu_1, \lambda^\vee \rangle$ lies in \mathbf{Z} . It then follows that neither side of the formula depends on ξ or $\tilde{\xi}$.

Proof of Lemma 6.3. First we evaluate the right side. The cocycle $\sigma \rightarrow \exp i\pi\lambda^\vee$ corresponds under the Tate-Nakayama isomorphism to the coset of λ^\vee in

$$\begin{aligned} & H^{-1}(X_*(T)) \\ &= \{ \mu^\vee \in X_*(T) : \sigma_T\mu^\vee = -\mu^\vee \} / \{ \nu^\vee - \sigma_T\nu^\vee : \nu^\vee \in X_*(T) \}. \end{aligned}$$

Thus $\kappa(\exp i\pi\lambda^\vee) = \lambda^\vee(x)$, where $s = (x, 1)\tilde{Z}^W$ was used to define κ . Note that we have transferred λ^\vee to ${}^L T^0$ via η .

For the left side, we write $a(z \times 1) = (a_0(z), a_0(z))$, $z \in \mathbf{C}^\times$, and $a(1 \times \sigma) = (xb_0, b_0)$, where $a_0(z), b_0$ lie in the center of ${}^L H_s^0$. Since $i\pi\lambda^\vee \in \text{Lie}(T(\mathbf{R}))$, we have $a_{(T,\eta)}(\exp i\pi\lambda^\vee) = e^{2\pi i\langle \mu_0, \lambda^\vee \rangle} = \lambda^\vee(a_0(-1))$, where again we have transferred λ^\vee to ${}^L T^0$ without change in notation (cf. proof of Lemma 6.1). On the other hand, $a(1 \times \sigma)\tilde{\sigma}_s(a(1 \times \sigma)) = a(-1)$ implies that $a_0(-1) = xb_0\sigma_s(b_0) = xb_0\sigma_{(T,\eta)}(b_0)$. Since $\sigma_{(T,\eta)}\lambda^\vee = -\lambda^\vee$, we have that $\lambda^\vee(a_0(-1)) = \lambda^\vee(x)$, and the lemma is proved.

We continue with $(T, \eta) \in \mathfrak{T}_H(G)$ and associated character κ on $H^1(T)$. Fix a set $\{u = \exp i\pi\lambda^\vee : \lambda^\vee \in X_*(T), \sigma_T\lambda^\vee = -\lambda^\vee\}$ such that the cocycles $\sigma \rightarrow \exp i\pi\lambda^\vee$ form a complete set of (noncohomologous) representatives for the elements of $H^1(T)$.

For $f \in C_c^\infty(\tilde{G}(\mathbf{R}))$, and Haar measures dt on $T(\mathbf{R})$, $d\tilde{g}$ on $\tilde{G}(\mathbf{R})$ form (cf. [Sh6]):

$$\Phi_f^{(T, \alpha, \kappa)}(\delta, dt, d\tilde{g}) = \sum_u \kappa(u) \int_{\tilde{G}(\mathbf{R})/T(\mathbf{R})} f(\alpha(\tilde{g})u\delta\tilde{g}^{-1}) \frac{d\tilde{g}}{dt},$$

for $\delta \in \tilde{T}(\mathbf{R})$ such that $\delta\alpha(\delta)$ is regular. Note that for all $\delta \in \tilde{T}(\mathbf{R})$, $\delta\alpha(\delta)$ lies in $T(\mathbf{R})^0$, the identity component of $T(\mathbf{R})$.

LEMMA 6.4.

$$\gamma \rightarrow a_{(T, \eta)}(\delta)\Phi_f^{(T, \alpha, \kappa)}(\delta, dt, d\tilde{g}),$$

if $\delta\alpha(\delta) = \gamma$, $\gamma \in T(\mathbf{R})_{\text{reg}}^0 = T(\mathbf{R})^0 \cap G_{\text{reg}}$, is a well-defined function on $T(\mathbf{R})_{\text{reg}}^0$.

Proof. By Lemma 6.3,

$$a_{(T, \eta)}(\delta)\Phi_f^{(T, \alpha, \kappa)}(\delta, dt, d\tilde{g}) = \sum_u a_{(T, \eta)}(u\delta) \int_{\tilde{G}(\mathbf{R})/T(\mathbf{R})} f(\alpha(\tilde{g})u\delta\tilde{g}^{-1}) \frac{d\tilde{g}}{dt}$$

which we will write as $\Phi(\delta)$. If $\delta\alpha(\delta) = \delta'\alpha(\delta')$ then $\delta' = v\delta$, where $v\alpha(v) = 1$, $v \in \tilde{T}(\mathbf{R})$. Then it is easily seen that $v = t^{-1}\alpha(t)u$ for some $t \in \tilde{T}(\mathbf{R})$ and u as in the summation. Since $a_{(T, \eta)}$ is α -invariant we then have $\Phi(\delta') = \Phi(v\delta) = \Phi(u\delta)$ which clearly coincides with $\Phi(\delta)$. Thus the lemma is proved.

Finally, suppose that $(T, \eta) \in \mathfrak{T}_H(G)$ and that $i(h, \eta): T' \rightarrow T$ is defined over \mathbf{R} . Then the Haar measure dt on $T(\mathbf{R})$ is transported via $i(h, \eta)$ to a Haar measure dt' on $T'(\mathbf{R})$; dt' is independent of the choice of h . Also, we say that $\gamma' \in T'(\mathbf{R})_{\text{reg}}$ is not a norm if it is not in the image of the norm map from $\tilde{T}' = \text{Res}_{\mathbf{R}}^{\mathbf{C}} T'$ to T' , i.e. γ' does not lie in the identity component of $T'(\mathbf{R})$. Then if γ' originates from $\gamma \in T(\mathbf{R})_{\text{reg}}$ via (T, η) , γ is not in the image of the norm from \tilde{T} to T (and conversely...).

We have not assumed that ξ or $\tilde{\xi}$ is of “unitary type” [Sh3]. It is easily checked that there is a quasicharacter χ on $H(\mathbf{R})$ such that $|\chi(\gamma')\Lambda_{(T, \eta)}(\gamma)a_{(T, \eta)}(\delta)| = 1$ if γ' originates from $\gamma = \delta\alpha(\delta)$ via (T, η) . We then define $\mathcal{C}_{\tilde{\xi}}(H(\mathbf{R}))$ to be the set of functions f on $H(\mathbf{R})$ such that $f\chi$ belongs to $\mathcal{C}(H(\mathbf{R}))$, the Schwartz space of $H(\mathbf{R})$. As the notation indicates, this space does not depend on the choice of χ . For $f \in \mathcal{C}_{\tilde{\xi}}(H(\mathbf{R}))$ the

stable orbital integrals $\Phi_f^{(T',1)}(\gamma', dt', dh)$, $\gamma' \in T'(\mathbf{R}) \cap H_{\text{reg}}$ (cf. [Sh4] etc.) are well-defined.

It remains now to recall the factor $\Delta_{(T,\eta)}$ from L -indistinguishability. Thus

$$\Delta_{(T,\eta)} = (-1)^{q(G,H)} \varepsilon(T, \eta) \Lambda_{(T,\eta)} \prime \Delta_{(T,\eta)},$$

where $q(G, H)$ is an integer, $(-1)^{q(G,H)}$ being inserted only for convenience, $\varepsilon(T, \eta) = \pm 1$ is defined implicitly, $\Lambda_{(T,\eta)}$ is as earlier in this section and $\prime \Delta_{(T,\eta)}$ is a discriminant function (see [Sh4, §3] for further details).

7. The matching theorem.

THEOREM 7.1. *Let H be an endoscopic group for (\tilde{G}, α) , with L -group ${}^L H_s$ chosen as earlier. Suppose that $\tilde{\xi}: {}^L H_s \hookrightarrow {}^L \tilde{G}$ is an allowed embedding and that $\xi: {}^L H_s \hookrightarrow {}^L G$ is admissible (for L -indistinguishability). Then for each $f \in C_c^\infty(\tilde{G}(\mathbf{R}))$ there exists $f_H \in \mathcal{C}_{\tilde{\xi}}(H(\mathbf{R}))$ such that:*

$$\Phi_{f_H}^{(T',1)}(\gamma', dt', dh) = \begin{cases} \Delta_{(T,\eta)}(\gamma) a_{(T,\eta)}(\delta) \Phi_f^{(T,\alpha,\kappa)}(\delta, dt, d\tilde{g}), \\ \text{if } \gamma' \text{ originates from } \gamma = \delta\alpha(\delta) \\ \text{via } (T, \eta) \in \mathfrak{T}_H(G), \\ 0 \text{ if } \gamma \text{ is not a norm.} \end{cases}$$

Here it is assumed that γ' originates from *regular* elements in $G(\mathbf{R})$. Then γ' is regular in $H(\mathbf{R})$ [Sh2]; T' is the maximal torus containing γ' . Recall that $\Delta_{(T,\eta)}$ depends on ξ alone, that $a_{(T,\eta)}$ depends on both ξ and $\tilde{\xi}$, and that $\Delta_{(T,\eta)}(\gamma) a_{(T,\eta)}(\delta)$ depends on $\tilde{\xi}$ alone... as long as (T, η) and δ are fixed.

REMARK. We have used $C_c^\infty(\tilde{G}(\mathbf{R}))$ instead of the more natural $\mathcal{C}(\tilde{G}(\mathbf{R}))$ since the necessary analysis of “twisted F_f ” (cf. [Sh6]), for f a Schwartz function, has not been carried out. Work of L. Clozel now in progress should settle this matter and allow us to replace $C_c^\infty(\tilde{G}(\mathbf{R}))$ by $\mathcal{C}(\tilde{G}(\mathbf{R}))$.

Proof of the theorem. Let $\gamma' \in H(\mathbf{R})$. Suppose that γ' originates from $\gamma \in G_{\text{reg}}$ via (T, η) and from $\bar{\gamma}$ via $(\bar{T}, \bar{\eta})$. Choose δ so that $\delta\alpha(\delta) = \gamma$. Write $\bar{\gamma}$ as $y\gamma y^{-1}$ and $\bar{\eta}$ as $\omega_H \circ \eta \circ \text{ad } y^{-1}$, where $\omega_H \in \Omega(H, T_H) \subset \Omega(G, T^*)$ and $y \in \mathfrak{A}(T)$ (cf. [Sh4, §3]). Then for $\bar{\delta}$ such that $\bar{\delta}\alpha(\bar{\delta}) = \bar{\gamma}$ we may take $y\delta y^{-1}$, where $y \in G$ has been identified with its image in \tilde{G} under the diagonal embedding. With this choice of $\bar{\delta}$ we have $a_{(\bar{T},\bar{\eta})}(\bar{\delta}) = a_{(T,\eta)}(\delta)$. The relation between $\Delta_{(\bar{T},\bar{\eta})}(\bar{\gamma})$ and $\Delta_{(T,\eta)}(\gamma)$ is described in [Sh4, §3].

For fixed $(T, \eta) \in \mathfrak{T}_H(G)$ the function

$$\gamma' \rightarrow a_{(T,\eta)}(\delta)\Delta_{(T,\eta)}(\gamma)\Phi_f^{(T,\alpha,\kappa)}(\delta, dt, d\tilde{g}),$$

if γ' originates from $\gamma = \delta\alpha(\delta)$ via (T, η) , is well-defined and invariant under $\mathfrak{A}(T')$. To prove this we invoke [Sh4, Propositions 2.4.5 and 3.1.2] and [Sh6, Lemma 4.3.2]. These results show that we have only to check that $a_{(T,\eta)}(\delta^\omega) = a_{(T,\eta)}(\delta)$ for ω an element of the Weyl group $\Omega(G, T)$ of (G, T) which commutes with the Galois action on \dot{T} and “comes from H ” (i.e. $\omega \in \Omega_0(G, T) \cap \Omega^{(\kappa)}(G, T)$) as in [Sh4, Proposition 2.4.5]). But this invariance of $a_{(T,\eta)}$ follows easily from the fact that $\langle \tilde{\mu}_0, \lambda^\vee \rangle = 0$ for λ^\vee in the span of the roots of ${}^L\tilde{T}^0$ in ${}^L\tilde{H}_s^0$ (see the proof of Lemma 6.1).

Suppose now that we fix a “framework of Cartan subgroups [Sh3], [Sh4, §3.2]. Thus we have specified certain pairs $(T_n, \eta_n) \in \mathfrak{T}_H(G)$ and embeddings $i_n = i(h_n, \eta_n): T'_n \rightarrow T_n$ over \mathbf{R} ; the set $\{T'_n(\mathbf{R})\}$ provides a complete family of representatives, without redundancy, for the conjugacy classes of Cartan subgroups of $H(\mathbf{R})$. Given $\gamma' \in T'_n(\mathbf{R})$, set $\gamma = i_n(\gamma')$, and choose any δ such that $\delta\alpha(\delta) = \gamma$. Call γ' G -regular if γ is regular. Then for each n we may consider the function on the G -regular elements of $T'_n(\mathbf{R})$ given by

$$\Phi_n(\gamma', dt', dh) = \begin{cases} \varepsilon_n \hat{\Delta}_{(T_n, \eta_n)}(\gamma) a_{(T_n, \eta_n)}(\delta) \Phi_f^{(T_n, \alpha, \kappa_n)}(\gamma, dt, d\tilde{g}) \\ \text{if } \gamma' \in T'_n(\mathbf{R})^0, \\ 0 \text{ if } \gamma' \notin T'_n(\mathbf{R})^0, \end{cases}$$

where $\varepsilon_n = \pm 1$ (to be chosen), $\hat{\Delta}_{(T,\eta)} = \varepsilon(T_n, \eta_n) \Delta_{(T_n, \eta_n)}$ (i.e. $\hat{\Delta}_{(T,\eta)}$ is $\Delta_{(T,\eta)}$ with the $\varepsilon(T, \eta)$ removed), and κ_n is the “ κ ” associated to (T_n, η_n) . Note that $\{\kappa_n |_{X_*(T_n)_{sc}}\}$ is exactly the set $\{\kappa_n\}$ from [Sh2, §7] and [Sh3, §2].

Suppose that we are able to show that there exists $f_H \in \mathcal{C}_{\tilde{z}}(H(\mathbf{R}))$ such that

$$(*) \quad \Phi_{f_H}^{(T'_n, 1)}(\gamma', dt', dh) = \begin{cases} \Phi_n(\gamma', dt', dh) & \text{if } \gamma' \in T'_n(\mathbf{R})^0, \\ 0 & \text{if } \gamma' \notin T'_n(\mathbf{R})^0, \end{cases}$$

for all G -regular γ' in $T'_n(\mathbf{R})$ and for all n provided $\varepsilon_m \varepsilon_n = \varepsilon(m, n)$ whenever $T'_m(\mathbf{R})$ and $T'_n(\mathbf{R})$ are adjacent Cartan subgroups. Here $\varepsilon(m, n)$ is as defined in [Sh4, §3.5] (cf. [Sh2]). Then we shall take $\varepsilon_n = \varepsilon(T_n, \eta_n)$, so that by the results of L -indistinguishability (exp. [Sh4, §3.5]) there does exist f_H satisfying (*). It is then routine to verify that f_H satisfies the statement of our theorem (see the first paragraph of this proof; similar arguments for L -indistinguishability are given in [Sh4, §3]).

Returning to the condition on the existence of f_H , we have only to show that our family $\{\Phi_n(\cdot, \cdot, \cdot)\}$ behaves like the family $\{\Phi_n\}$ of [Sh2, §9] (cf. [Sh4, §3.2]). The invariance and growth requirements being satisfied (clearly), only the “jump conditions” remain. Thus we need the jump formulas for the functions $\Psi_{(T,\eta)}$:

$$\gamma \rightarrow \begin{cases} a_{(T,\eta)}(\delta) \hat{\Delta}_{(T,\eta)}(\gamma) \Phi^{(T,\alpha,\kappa)}(\delta, dt, d\tilde{g}) & \text{if } \gamma \in T(\mathbf{R})_{\text{reg}}^0, \\ 0 & \text{if } \gamma \in T(\mathbf{R})_{\text{reg}} - T(\mathbf{R})^0. \end{cases}$$

These are contained essentially in the analysis of §§4 and 5 of [Sh6]. To be more precise, we seek analogues of Lemmas 5.2.2 and 5.2.5 of [Sh6], when “ $\Delta_T \Phi^\tau$ ” is replaced by the function above (with the necessary adjustment in the choice of positive system for the imaginary roots of T used to define the factor $\hat{\Delta}_{(T,\eta)}$). The proof of the analogue of Lemma 5.2.2 is straightforward; because of notational complications we omit further details. Note that the “ κ -signature” [Sh2] which appears depends only on $\kappa|_{X_*(T_{\text{sc}})}$, i.e. the jump is indeed like that from L -indistinguishability. The analogue of Lemma 5.2.5 will be stronger than the original statement, because we no longer need the assumption “ $\kappa(\alpha^\vee) = 1$ if (5.2.3) holds.” We now have the exact analogue of [Sh2, Proposition 9.1] from L -indistinguishability. Indeed, let γ_0 be a semiregular element in $T(\mathbf{R})$ such that $\lambda(\gamma_0) = 1$, where λ is an imaginary root such that $\kappa(\lambda^\vee) = -1$. We wish to show that $\Psi_{(T,\eta)}$ is smooth on some neighborhood of γ_0 . We may assume that $\gamma_0 \in T(\mathbf{R})^0$. Fix $\delta_0 \in T(\mathbf{R})^0$ such that $\delta_0^2 = \gamma_0$. For γ close to γ_0 choose δ close to δ_0 such that $\delta^2 = \gamma$. It will be sufficient to show that $\delta \rightarrow \Psi_{(T,\eta)}(\delta^2)$ is smooth near δ_0 . This follows immediately from Lemma 4.3.3 of [Sh6]. Note that this type of argument could not be used in the proof of Lemma 5.2.5 of [Sh6] because the “cross-section for the norm” was not smooth near γ_0 .

We now complete the proof of Theorem 7.1 by the arguments already indicated.

8. The dual lifting. Again we fix an element of $\mathfrak{S}(\tilde{G}, \alpha)$ and choose a convenient representative $(s, {}^L H_s)$ for this element, as in §5. Let H_s be the corresponding endoscopic group. Since H_s is, by definition, quasi-split over \mathbf{R} , the set $\Phi(H_s)$ [L2] consists of all equivalence classes of admissible homomorphisms $\phi: W \rightarrow {}^L H_s$. Suppose that $\tilde{\xi}: {}^L H_s \hookrightarrow {}^L \tilde{G}$ is an allowed embedding. Then $\tilde{\xi}$ induces a map, also to be denoted $\tilde{\xi}$, from $\Phi(H_s)$ to $\Phi(\tilde{G})$; the image of the class of $\phi: W \rightarrow {}^L H_s$ is the class of $\tilde{\phi} = \tilde{\xi} \circ \phi: W \rightarrow {}^L \tilde{G}$. It is easily checked that the image of $\Phi(H_s)$ in $\Phi(\tilde{G})$ is independent of the choice for $\tilde{\xi}$. By the remarks at the end of §5 it is also independent of the choice for $(s, {}^L H_s)$.

On the other hand, the automorphism α of \tilde{G} has a standard dual [Sh7], again denoted α :

$$\alpha((g, h) \times w) = (h, g) \times w, \quad h, g \in {}^L G^0, w \in W.$$

If $\phi: W \rightarrow {}^L \tilde{G}$ is admissible then so is $\alpha \circ \phi: W \rightarrow {}^L \tilde{G}$. We write $\{\phi\}$ for the class of ϕ and $\{\phi\}^\alpha$ for the class of $\alpha \circ \phi$. Then $\Phi(\tilde{G})^\alpha = \{\{\phi\} \in \Phi(\tilde{G}): \{\phi\}^\alpha = \{\phi\}\}$.

For each element of $\mathfrak{S}(\tilde{G}, \alpha)$ we fix a representative $(s, {}^L H_s)$ as before, and assume that each ${}^L H_s$ has an allowed embedding $\tilde{\xi}$ in ${}^L \tilde{G}$. Also, we will use \bigcup_{H_s} to denote a union over the corresponding endoscopic groups.

THEOREM 8.1.

$$\Phi(\tilde{G})^\alpha = \bigcup_{H_s} \tilde{\xi}(\Phi(H_s)).$$

Proof. Let $\phi: W \rightarrow {}^L H_s$ be admissible. Set $\tilde{\phi} = \tilde{\xi} \circ \phi$. We may assume that $\phi(\mathbf{C}^x \times 1) \subset {}^L T^0 \times \mathbf{C}^x \times 1$. Then clearly $\tilde{\phi}$ and $\alpha \circ \tilde{\phi}$ coincide on $\mathbf{C}^x \times 1$. We write $\phi(1 \times \sigma)$ as $n_H \times 1 \times \sigma \in {}^L H_s$, and $\tilde{\xi}(1 \times 1 \times \sigma)$ as $(xm_0, m_0) \times 1 \times \sigma$ (cf. §5). Then $\tilde{\phi}(1 \times \sigma) = (xn_H m_0, n_H m_0) \times 1 \times \sigma$ and

$$\begin{aligned} (\alpha \circ \tilde{\phi})(1 \times \sigma) &= (n_H m_0, xn_H m_0) \times 1 \times \sigma \\ &= (x^{-1}, x)\tilde{\phi}(1 \times \sigma) = g^{-1}\tilde{\phi}(1 \times \sigma)g, \end{aligned}$$

where $g = (x, 1)$. Then clearly $\alpha \circ \tilde{\phi} = \text{ad } g^{-1} \circ \tilde{\phi}$, and so $\tilde{\xi}(\Phi(H)) \subset \Phi(\tilde{G})^\alpha$.

Suppose now that $\tilde{\phi}: W \rightarrow {}^L \tilde{G}$ is admissible and that $\{\tilde{\phi}\}^\alpha = \{\tilde{\phi}\}$. Then it is sufficient to show that $\tilde{\phi}$ factors through some ${}^L H_s$ (not necessarily among our fixed representatives) embedded (via an allowed embedding) in ${}^L \tilde{G}$.

Let $S_{\tilde{\phi}}^\alpha = \{a \in {}^L \tilde{G}^0: a\tilde{\phi}(w)a^{-1} = (\alpha \circ \tilde{\phi})(w), w \in W\}$. Then $S_{\tilde{\phi}}^\alpha$ is nonempty. If a_0 lies in $S_{\tilde{\phi}}^\alpha$ then so does $a_0 z$, for $z \in \tilde{Z}^W$. In fact, then $S_{\tilde{\phi}}^\alpha = a_0 S_{\tilde{\phi}}$, where $S_{\tilde{\phi}}$ is the centralizer of $\tilde{\phi}(W)$ in ${}^L \tilde{G}^0 \dots$ recall that the results of [Sh4], with a little extra argument for the case $\tilde{\phi}$ unbounded, show that $S_{\tilde{\phi}} = S_{\tilde{\phi}}^0 \tilde{Z}^W$, $S_{\tilde{\phi}}^0$ denoting the identity component in $S_{\tilde{\phi}}$. Choose $s = a_0 \tilde{Z}^W$ contained in $S_{\tilde{\phi}}^\alpha$. Assume that s consists of α -semisimple elements (\dots we will prove below that such an s exists). Then set ${}^L H_s^0 = (\text{Cent}_{\alpha}(s, {}^L \tilde{G}^0))^0$, and select ${}^L B_s^0$, ${}^L T_s^0$ and $\{Y\}$ as in §3. To define a suitable action of W on ${}^L H_s^0$ we have just to give a homomorphism of

$\text{Gal}(\mathbf{C}/\mathbf{R})$ into $\text{Aut}({}^L H_s^0, {}^L B_s^0, {}^L T_s^0, \{Y\})$ such that σ_s , the image of σ , is “realized in $\text{Cent}_\alpha(s, {}^L \tilde{G}) = \text{Cent}_\alpha(a_0, {}^L \tilde{G})$ ”. But

$$\tilde{\phi}(1 \times \sigma)^{-1} a_0 \alpha(\tilde{\phi}(1 \times \sigma)) = a_0.$$

Thus $\tilde{\phi}(1 \times \sigma)$ normalizes ${}^L H_s^0$. We may write $\text{ad } \tilde{\phi}(1 \times \sigma)|_{{}^L H_s^0}$ as $\omega \sigma_s$, where ω is an inner automorphism of ${}^L H_s^0$ and $\sigma_s \in \text{Aut}({}^L H_s^0, {}^L B_s^0, {}^L T_s^0, \{Y\})$. Note that $\sigma_s^2 = 1$ and is “realized in $\text{Cent}_\alpha(s, {}^L \tilde{G})$ ”. Using the associated W -action we form ${}^L H_s$ and so obtain a representative $(s, {}^L H_s)$ for an element of $\mathfrak{S}(\tilde{G}, \alpha)$. We claim that $\tilde{\phi}$ factors through ${}^L H_s$. Thus, suppose that $\tilde{\xi}: {}^L H_s \hookrightarrow {}^L \tilde{G}$ is an allowed embedding. Then for each $w \in W$, $\tilde{\phi}(w)$ lies in $\text{Cent}_\alpha(s, {}^L \tilde{G})$ and acts on ${}^L H_s^0 = (\text{Cent}_\alpha(s, {}^L \tilde{G}^0))^0$ as an element $n(w) \times w$ of the image of ${}^L H_s$ in ${}^L \tilde{G}$. By definition, $n(w) \times w \in \text{Cent}_\alpha(s, {}^L \tilde{G})$. Thus $\tilde{\phi}(w) = a(w)(n(w) \times w)$, where $a(w) \in \text{Cent}_\alpha(s, {}^L \tilde{G}^0)$ centralizes ${}^L H_s^0$. But then $a(w)$ lies in the center of ${}^L H_s^0$. Hence $\tilde{\phi}$ factors through ${}^L H_s$.

It remains now to show that S_ϕ^α contains an α -semisimple element. If we replace $\tilde{\phi}$ by $\text{ad } g \circ \tilde{\phi}$, $g \in {}^L \tilde{G}^0$, then we must replace S_ϕ^α by $\alpha(g) S_\phi^\alpha g^{-1}$. Therefore we may assume that S_ϕ^α contains an element $(x^{-1}, 1)$, $x \in {}^L G^0$ (cf. Lemma 4.1). Then we write $\tilde{\phi}(w)$ as $(\phi_1(w), \phi_2(w)) \times w$ and obtain from $(x^{-1}, 1)\tilde{\phi}(w)(x, 1) = \alpha(\tilde{\phi}(w))$, $w \in W$, that $\phi_1(z \times 1) = \phi_2(z \times 1)$, $z \in \mathbf{C}^x$, and $\phi_1(1 \times \sigma) = x\phi_2(1 \times \sigma)$; also, x lies in the centralizer S_0 of the image of the homomorphism $\hat{\phi}_1: w \rightarrow \phi_1(w) \times w$ of W into ${}^L G$. Write $x = x_u x_s$, where $x_u \in S_0$ is unipotent and $x_s \in S_0$ is semisimple. Then with the same ϕ_1 and with x_s in place of x we can use the formulas above for $\tilde{\phi}$ to define $\tilde{\phi}_0: W \rightarrow {}^L \tilde{G}$ such that $S_{\tilde{\phi}_0}^\alpha$ contains $(x_s^{-1}, 1)$. But $\tilde{\phi}_0$ is easily seen to be equivalent to $\tilde{\phi}$ because x_u^{-1} , being unipotent and fixed by $\hat{\phi}_1(W)$, can be written as $v(\hat{\phi}_1(1 \times \sigma)v)^{-1}$, $v \in \text{Cent}(\phi_1(\mathbf{C}^x \times 1), {}^L G^0)$. Since $(x_s^{-1}, 1)$ is α -semisimple our proof of Theorem 8.1 is complete.

According to Langlands’ functoriality principle this factoring of the α -fixed parameters $\{\tilde{\phi}\}$ should be reflected in character theory. Let $\tilde{\phi} \in \Phi(\tilde{G})$ be α -fixed (we now drop the $\{ \}$ from the notation for parameters). Then the L -packet $\Pi_{\tilde{\phi}}$ consists of a single infinitesimal equivalence class of irreducible admissible representations fixed by the automorphism $\alpha: \tilde{G}(\mathbf{R}) \rightarrow \tilde{G}(\mathbf{R})$ (... this is easily checked, see also [C1]). Thus the twisted character $\chi_{\tilde{\phi}}^\alpha$ of $\Pi_{\tilde{\phi}}$ is well-defined up to sign (see [C1] for a detailed discussion, especially concerning the question of signs). Assume that $\tilde{\phi}$ is bounded, i.e. if $\tilde{\phi}(w) = \tilde{\phi}_0(w) \times w$, $w \in W$, then $\tilde{\phi}_0(W)$ is bounded. Then $\chi_{\tilde{\phi}}^\alpha$ is tempered [C1, Theorem 5.12]. On the other hand, suppose that $\tilde{\phi}$ is the lift of $\phi \in \Phi(H)$, in the sense afforded by Theorem

8.1. Then ϕ is essentially bounded, so that the L -packet Π_ϕ consists of essentially tempered (equivalence classes of) representations. Thus $\chi_\phi = \sum_{\pi \in \Pi_\phi} \chi_\pi$, χ_π denoting the ordinary character of π , is a stable essentially tempered distribution on $H(\mathbf{R})$ [Sh1, Lemma 5.2].

Theorem 7.1 provides a correspondence (f, f_H) between $C_c^\infty(G(\mathbf{R}))$ and $\mathcal{C}_{\tilde{\xi}}(H(\mathbf{R}))$. As mentioned already, an adequate analysis of the “twisted F_f transform” would provide a correspondence between $\mathcal{C}(\tilde{G}(\mathbf{R}))$ and $\mathcal{C}_{\tilde{\xi}}(H(\mathbf{R}))$; it would also give a dual lifting of stable tempered distributions on $H(\mathbf{R})$ to twisted-invariant tempered distributions on $\tilde{G}(\mathbf{R})$, with eigendistributions mapping to eigendistributions (see [Sh4, §4] for the analogous arguments in the case of L -indistinguishability). Nevertheless, with the correspondence of Theorem 7.1 we can define $(\text{Lift } \chi_\phi)(f) = \chi_\phi(f_H)$, $f \in C_c^\infty(\tilde{G}(\mathbf{R}))$. Writing $\chi_\phi(f_H)$ as $\int_{H(\mathbf{R})} f_H(h) \chi_\phi(h) dh$, and applying the Weyl Integration Formula, the matching theorem and the twisted analogue of the Weyl Integration Formula, we find that $\text{Lift } \chi_\phi$ is a twisted-invariant distribution on $\tilde{G}(\mathbf{R})$ represented by a function explicitly computed in terms of χ_ϕ . Moreover, this function transforms under the center of the universal enveloping algebra of $\tilde{G}(\mathbf{C})$ according to the infinitesimal character of χ_ϕ^α . We may therefore ask if $\text{Lift } \chi_\phi$ coincides with χ_ϕ^α up to a constant (depending only on G and H , once the sign for χ_ϕ^α has been suitably fixed). According to [C1] with some minor additional arguments, this is true if $H = G$; recall that we are assuming that ϕ is bounded, so that ϕ is an essentially bounded parameter. Work of L. Clozel now in progress should provide the answer to our question for the case $H \neq G$.

REFERENCES

[C1] L. Clozel, *Changement de base pour les représentations tempérées des groupes réductifs réels*, Ann. Sci. Ec. Norm. Sup., 4^e série, t. 15 (1982), 45–115.
 [L1] R. Langlands, *Stable conjugacy: definitions and lemmas* Canad. J. Math., XXXI, No. 4 (1979), 700–725.
 [L2] ———, *On the classification of irreducible representations of real algebraic groups*, preprint (1973).
 [L3] ———, *Les débuts d’une formule des traces stables*, Publ. Math. Univ. Paris VII, vol. 13 (1983).
 [Sh1] D. Shelstad, *Characters and inner forms of a quasi-split group over R* , Compositio Math., 39 (1979), 11–45.
 [Sh2] ———, *Orbital integrals and a family of groups attached to a real reductive group*, Ann. Sci. Ec. Norm. Sup., 4^e série, t. 12 (1979), 1–31.
 [Sh3] ———, *Embeddings of L -groups*, Canad. J. Math., 33 (1981), 513–558.
 [Sh4] ———, *L -indistinguishability for real groups*, Math. Ann., 259 (1982), 385–430.
 [Sh5] ———, *Orbital integrals, endoscopic groups and L -indistinguishability for real groups*, to appear in Publ. Math. Univ. Paris VII.

[Sh6] ———, *Base Change and a Matching Theorem for Real Groups*, in Springer Lecture Notes, Vol. **880** (1982), pp. 425–482.

[Sh7] ———, *A generalization of endoscopic groups*, (in preparation).

Received April 20, 1982. Supported by NSF Grants MCS 79-01712, MCS 81-02392 and MCS 81-08814(A01).

RUTGERS UNIVERSITY
NEWARK, NJ 07102