

ON EMBEDDING SEMIFLOWS INTO A RADIAL FLOW ON l_2

ROGER C. McCANN

Let π be a semiflow on a separable metric space X such that the negative escape time function is lower semicontinuous and $x \rightarrow x\pi t$ is a one-to-one mapping for each $t \in R^+$. If π has a globally uniformly asymptotically stable critical point, then π can be embedded into a radial flow on l_2 . This generalizes known results on embedding flows or semiflows into radial flows on l_2 .

1. Introduction. In [3] L. Janos showed that a semiflow π on a compact metric space X satisfying

(i) $\cdot\pi t$ is one-to-one for every $t \in R^+$

(ii) there is a $p \in X$ such that $\cap \{X\pi t: t \geq 0\} = \{p\}$ can be embedded into a radial flow on l_2 . In [2] M. Edelstein generalized this result to

THEOREM I. *Let π be a semiflow on a separable metric space X satisfying*

(a) *for each $t \in R^+$, $x \rightarrow x\pi t: X \rightarrow X$ is a homeomorphism, of X onto a closed subset of X ,*

(b) *there is a $p \in X$ such that for each neighborhood U of p there is a $T \in R^+$ such that $X\pi t \subset U$ for all $t \geq T$.*

Then π can be embedded into a radial flow on l_2 .

Evidently properties (a) and (b) generalize properties (i) and (ii) respectively. Note that property (b) imposes a type of compactness on the semiflow. For example, a radial flow on l_2 can be embedded into itself trivially, but such a flow does not have property (b).

In this paper we further generalize properties (a) and (b) to

(c) $x \rightarrow x\pi t$ is one-to-one for each $t \in R^+$,

(d) the negative escape time function is lower semicontinuous,

(e) π has a globally uniformly asymptotically stable critical point p .

We will show (Corollary 8) that property (a) implies properties (c) and (d). Evidently property (b) implies property (e). Property (e) imposes a type of local compactness on the semiflow. Notice that a radial flow on l_2 does satisfy property (e).

The principal result of this paper, Theorem 7, generalizes every other result known to the author concerning embedding flows or semiflows into radial flows on l_2 .

2. Notation and definitions. Throughout this paper R and R^+ will denote the reals and nonnegative reals respectively. A flow on a topological space X is a continuous mapping $\pi: X \times R \rightarrow X$ such that (where $x\pi t = \pi(x, t)$) $x\pi 0 = x$ for all $x \in X$ and $(x\pi t)\pi s = x\pi(t + s)$ for all $x \in X$ and $t, s \in R$. If R is replaced by R^+ in the previous sentence, then π is called a semiflow. A point p of X is called a critical point of π if $p\pi t = p$ for all $t \in R$ (or $t \in R^+$ if π is a semiflow). A compact subset M of X is said to be stable with respect to π if for any neighborhood U of M there is a neighborhood V of M such that $V\pi R^+ \subset U$. A compact subset M of X is said to be a global attractor if for any neighborhood U of M and any $x \in X$ there is a $d \in R^+$ such that $x\pi[d, \infty) \subset U$. The compact set M is called a global uniform attractor if it is a global attractor and if there is a neighborhood U of M such that for any neighborhood $V \subset U$ of M there is a $c \in R^+$ such that $U\pi[c, \infty) \subset V$. A stable global (uniform) attractor is said to be globally (uniformly) asymptotically stable.

A continuous function $L: X \rightarrow R^+$ is called a Liapunov function for a compact subset M of X if $L(x\pi t) < L(x)$ for every $x \in X - M$ and $0 < t$, $L(x\pi t) \rightarrow 0$ as $t \rightarrow \infty$ for every $x \in X$, and $L(x) = 0$ if $x \in M$. Let M be a compact asymptotically stable subset of X . A straightforward argument shows that if $x \in X - M$ and if U is any neighborhood of M , then there is a neighborhood V of x and a $T > 0$ such that $V\pi[T, \infty) \subset U$. With this observation the proof of the following theorem is essentially identical with that of Theorem 10 in [1].

THEOREM II. *A compact subset M of a metric space X is globally asymptotically stable with respect to a semiflow π if and only if there is Liapunov function for M .*

Let X and Y be topological spaces on which are defined flows (semiflows) π and ρ respectively. We say that π can be embedded into ρ if there is a homeomorphism h of X onto a subset of Y such that $h(x\pi t) = h(x)\rho t$ for every $x \in X$ and $t \in R$ ($t \in R^+$).

The set of all sequences $x = \{x_1, x_2, \dots, x_n, \dots\}$ of real numbers such that $\sum_{n=1}^{\infty} x_n^2$ converges is denoted by l_2 . If addition and scalar multiplication are defined coordinatewise and if a norm is defined by $\|x\| = (\sum_{n=1}^{\infty} x_n^2)^{1/2}$, then l_2 is a real Banach space. A flow ρ on l_2 is called a radial flow if there is a $c \in (0, 1)$ such that $x\rho t = c^t x$ for every $(x, t) \in l_2 \times R$.

Let π be a semiflow on X . The function $\alpha: X \rightarrow [-\infty, 0]$ defined by $\alpha(x) = \inf\{-t: \text{there exists } y \in X \text{ with } y\pi t = x\}$ is called the negative escape time function. Throughout this paper we shall

assume that α is lower semicontinuous, i.e., $\alpha(x) \leq \lim_{y \rightarrow x} \inf \alpha(y)$. It is an elementary exercise to show that $\alpha(x\pi t) = \alpha(x) - t$ for all $t \geq 0$ and $x \in X$.

3. The embedding. Henceforth, π shall denote a semiflow on a separable metric space X satisfying

- (1) $x \rightarrow x\pi t$ is one-to-one for each $t \in R^+$,
- (2) the negative escape time function is lower semicontinuous,
- (3) π has a globally uniformly asymptotically stable critical point p .

Also, U shall denote a neighborhood of p such that for any neighborhood $V \subset U$ of p , there is a $T > 0$ such that $U\pi[T, \infty) \subset V$.

Let $t < 0$ and $x \in X$. Since $\cdot\pi(-t)$ is one-to-one there is at most one $y \in X$ with $y\pi(-t) = x$. If such a y exists then we shall denote this y by $x\pi t$. It is a straightforward exercise to show that if $s, t \in R$ and $x \in X$, then $(x\pi t)\pi s = x\pi(t + s)$ whenever each side of the equality is defined. Suppose that $\{x_i\}$ and $\{t_i\}$ are sequences in X and R converging to $x \in X$ and $t \in R$ respectively. Using property 2 it is easy to show that if $x_i\pi t_i$ is defined for each i , then $x\pi t$ is defined and $x_i\pi t_i \rightarrow x\pi t$ as $i \rightarrow \infty$.

LEMMA 1. *If $x\pi(\alpha(x), 0] \subset U$, then $-\infty < \alpha(x)$.*

Proof. Let $V \subset U$ be a neighborhood of p such that $V\pi R^+ = V$ and $x \notin V$. Then $x\pi(\alpha(x), 0] \cap V = \emptyset$. Let $T > 0$ be such that $U\pi T \subset V$. Then $x\pi(\alpha(x) + T, \infty) \subset V$. In order that this be consistent with $x\pi(\alpha(x), 0] \cap V = \emptyset$, we must have $\alpha(x) \neq -\infty$.

LEMMA 2. *Let σ be a semiflow on a metric space Z . If*

- (i) *the negative escape time function γ is lower semicontinuous,*
- (ii) *each trajectory contains a start point, i.e., for each $x \in Z$ there is a $y \in Z$ such that $y\sigma(-\gamma(x)) = x$,*
then $Z\pi t$ is a closed subset of Z for each $t \geq 0$.

Proof. Let $t \geq 0$ and let $\{x_i\}$ be a sequence in Z such that $x_i\sigma t \rightarrow y$ for some $y \in Z$. Then $\gamma(x_i\sigma t) \leq -t$ for every i so that $\gamma(y) \leq -t$. By (ii) there is a $z \in Z$ such that $z\sigma(-\gamma(y)) = y$. Then $y = (z\pi(-\gamma(y) - t))\sigma t \in Z\sigma t$. It follows that $Z\sigma t$ is a closed subset of Z .

Let L be a Liapunov function for p (Theorem II) and let λ be any number in the range of L such that $L^{-1}([0, \lambda]) \subset U$. Set

$$Y = \{x \in L^{-1}([0, \lambda]): \alpha(x) \leq -1 \text{ and } x\pi(-1, \infty) \subset L^{-1}([0, \lambda])\}$$

and let σ denote the semiflow obtained by restricting π to $Y \times R^+$. Let β denote the negative escape time function with respect to σ .

We will show that σ satisfies the hypotheses of Theorem I. Hence, σ can be embedded into a radial flow on l_2 . We will then extend this embedding to an embedding of π into a radial flow.

LEMMA 3. *For every $x \in Y$ there is a $y \in Y$ such that $y\sigma(-\beta(x)) = x$.*

Proof. There are two cases to consider: $x\pi(\alpha(x), \infty) \subset L^{-1}([0, \lambda])$ and $x\pi(\alpha(x), \infty) \cap L^{-1}(\lambda) \neq \emptyset$. In the latter case there is a unique $z \in x\pi(\alpha(x), \infty) \cap L^{-1}(\lambda)$ and a unique $t \in R$ such that $z\pi t = x$. Since $x \in Y$ we must have $1 \leq t$. Then $\beta(x) = -t + 1$. Set $y = z\pi 1$. Then $y \in Y$ and $y\sigma(-\beta(x)) = y\pi(-\beta(x)) = (z\pi 1)\pi(t - 1) = z\pi t = x$. Now suppose $x\pi(\alpha(x), \infty) \subset L^{-1}([0, \lambda])$. Then $x\pi(\alpha(x), \infty) \subset U$ so that, by Lemma 1, $-\infty < \alpha(x)$. Since $x \in Y$ we must have $\alpha(x) \leq -1$. Let $y \in x\pi(\alpha(x), \infty)$ be such that $y\pi(-\alpha(x) - 1) = x$. Since $\alpha(x) = \alpha(y\pi(-\alpha(x) + 1)) = \alpha(y) + \alpha(x) + 1$ we have $\alpha(y) = -1$. If $y = z\pi t$ for some $t > 0$ then $-1 = \alpha(y) = \alpha(z\pi t) = \alpha(z) - t$ so that $\alpha(z) = t - 1 > -1$. It follows that $\beta(x) = \alpha(x) + 1$ and that $y\sigma(-\beta(x)) = x$. This completes the proof.

LEMMA 4. *Let $\{x_i\}$ be a sequence such that $x_i \rightarrow x$ for some $x \in X$. If there exists a $t \in R$ such that $x\pi t \in L^{-1}(\lambda)$, then either $t \leq \liminf \alpha(x_i)$ or there are a subsequence $\{x_j\}$ of $\{x_i\}$ and a sequence $\{t_j\}$ in R such that $x_j\pi t_j \in L^{-1}(\lambda)$. In the latter case $t_j \rightarrow t$.*

Proof. Suppose $\liminf \alpha(x_i) < t$. Let $\{x_j\}$ be a subsequence of $\{x_i\}$ such that $\alpha(x_j) \rightarrow \liminf \alpha(x_i)$. For any $\delta \in (0, t - \liminf \alpha(x_i))$ eventually $\alpha(x_j) < t - \delta$. Also $\alpha(x) \leq t - \delta$ because $\alpha(x) \leq \liminf \alpha(x_i)$. Since $L(x\pi(t - \delta)) > L(x\pi t) = \lambda > L(x\pi(t + \delta))$ we have $L(x_j\pi(t - \delta)) > \lambda > L(x_j\pi(t + \delta))$ eventually. Hence, there are $t_j \in (t - \delta, t + \delta)$, eventually, such that $L(x_j\pi t_j) = \lambda$. Since δ can be chosen arbitrarily small we must have $t_j \rightarrow t$.

LEMMA 5. *β is lower semicontinuous.*

Proof. Let $x \in Y$ and let $\{x_i\}$ be a sequence in Y such that $x_j \rightarrow x$. Let $\{x_j\}$ be any subsequence of $\{x_i\}$ such that $\beta(x_j) \rightarrow \beta$ for some $\beta \in [-\infty, 0]$. There are two cases to consider: $x\pi t \in L^{-1}(\lambda)$ for some $t \in R$ and $x\pi(\alpha(x), \infty) \subset L^{-1}([0, \lambda])$. If $x\pi t \in L^{-1}(\lambda)$ for some t , then by Lemma 4 either $\alpha(x) \leq t \leq \liminf \alpha(x_j)$ or there are a subsequence $\{x_k\}$ of $\{x_j\}$ and a sequence $\{t_k\}$ in R such that $t_k \rightarrow t$ and $x_k\pi t_k \in$

$L^{-1}(\lambda)$. If $t \leq \liminf \alpha(x_j)$, then $\beta(x) = t + 1$ and $\beta(x_j) = \alpha(x_j) + 1$ so that $\beta(x) \leq \liminf \beta(x_j) = \beta$. If there are a subsequence $\{x_k\}$ of $\{x_j\}$ and a sequence $\{t_k\}$ in R such that $t_k \rightarrow t$ and $x_k \pi t_k \in L^{-1}(\lambda)$, then $\beta(x) = t + 1$ and $\beta(x_k) = t_k + 1$. Then $\beta(x) = \lim \beta(x_k) = \beta$. Thus if $x \pi t \in L^{-1}(\lambda)$, then $\beta(x) \leq \beta$. It follows that $\beta(x) \leq \liminf \beta(x_i)$ whenever $x \pi t \in L^{-1}(\lambda)$ for some $t \in R$. Now suppose that $x \pi(\alpha(x), \infty) \subset L^{-1}([0, \lambda])$. Then $\beta(x) = \alpha(x) + 1$. Again there are two cases to consider: $x_i \pi(\alpha(x_i), \infty) \subset L^{-1}([0, \lambda])$ for every i and there exist a subsequence $\{x_n\}$ of $\{x_i\}$ and a sequence $\{s_n\}$ in R such that $x_n \pi s_n \in L^{-1}(\lambda)$ for every n . In the former case we have $\beta(x_i) = \alpha(x_i) + 1$ and $\beta(x) \leq \liminf \beta(x_i)$ since α is lower semicontinuous. In the latter case, let $V \subset U$ be a neighborhood of p such that $x \notin \overline{V \pi R^+}$ and let $T > 0$ be such that $U \pi [T, \infty) \subset V$. Then $L^{-1}(\lambda) \pi [T, \infty) \subset V$ and we must have $s_n \in [0, T]$ for all n sufficiently large. Let s be any accumulation point of $\{s_n\}$ and let $\{s_j\}$ be a subsequence of $\{s_n\}$ such that $s_j \rightarrow s$. Then $x_j \pi s_j \in L^{-1}(\lambda)$ and $x_j \pi s_j \rightarrow x \pi s$. Hence, $x \pi s \in L^{-1}(\lambda)$ which contradicts our assumption that $x \pi(\alpha(x), \infty) \subset L^{-1}([0, \lambda])$. It follows that $\beta(x) \leq \liminf \beta(x_i)$ whenever $x \pi(\alpha(x), \infty) \subset L^{-1}([0, \lambda])$. Combining this with the result $\beta(x) \leq \liminf \beta(x_i)$ whenever $x \pi t \in L^{-1}(\lambda)$ for some $t \in R$ obtained earlier in the proof, we conclude that β is lower semicontinuous.

Collecting together the above results we have that

- (i) σ is a semiflow on the separable metric space Y ,
- (ii) if V is a neighborhood in Y of p , then there is a $T > 0$ such that $Y \sigma [T, \infty) \subset V$, (This follows directly from the facts that $Y \subset U$ and σ is a restriction of π .)
- (iii) $Y \sigma t$ is a closed subset of Y for every $t \geq 0$ (Lemmas 3, 5, and 2).

In light of Theorem I the semiflow σ on Y can be embedded into a radial flow ρ on l_2 . Let $c \in (0, 1)$ be such that $x \rho t = c^t x$ and let $h: Y \rightarrow l_2$ be a homeomorphism of Y onto $h(Y)$ such that $h(x \sigma t) = h(x) \rho t$ for every $(x, t) \in Y \times R^+$. Since σ is a restriction of π we have $h(x \pi t) = h(x) \rho t$ for every $(x, t) \in Y \times R^+$. Now define a mapping $H: X \rightarrow l_2$ by

$$H(x) = h(x \pi t) \rho(-t)$$

where $t \in R^+$ is such that $x \pi t \in Y$. (H will be shown to be well defined in the following lemma.)

LEMMA 6. H is a homeomorphism of X onto $H(X)$.

Proof. We will first show that H is well defined. Clearly for

every $x \in X$, there is a $t \geq 0$ such that $x\pi t \in Y$. Moreover, if $x\pi t \in Y$, then $x\pi(t+s) \in Y$ for every $s \geq 0$. In order to show that H is well defined it suffices to show that $h(x\pi t)\rho(-t) = h(x\pi(t+s))\rho(-t-s)$ whenever $x\pi t \in Y$ and $s \geq 0$. Since $x\pi t \in Y$ and $s \geq 0$ we have $h(x\pi(t+s)) = h((x\pi t)\pi s) = h(x\pi t)\rho s$. Hence $h(x\pi(t+s))\rho(-t-s) = (h(x\pi t)\rho s)\rho(-t-s) = h(x\pi t)\rho(-t)$. The mapping H is well defined. We will now show that H is one-to-one. Suppose that $H(x) = h(x\pi t)\rho(-t)$, $H(y) = h(y\pi s)\rho(-s)$, and $H(x) = H(y)$. Without loss of generality we may assume that $t \geq s$. Then $H(y) = h(y\pi t)\rho(-t)$ since $y\pi t \in Y$ whenever $y\pi s \in Y$ and $s \leq t$. Since $H(x) = H(y)$ we must have $h(x\pi t) = h(y\pi t)$. Recalling that h is a homeomorphism we have $x\pi t = y\pi t$ so that $x = y$ since $\cdot\pi t$ is one-to-one. The mapping H is one-to-one. Next we will show that H is continuous. Let $x \in X$ and let $\{x_i\}$ be a sequence in X such that $x_i \rightarrow x$. Let $t \in R^+$ be such that $L(x\pi t) < \lambda$. Then $x\pi(t+1) \in Y$. Also for all i sufficiently large $L(x_i\pi t) < \lambda$ and $x_i\pi(t+1) \in Y$. Then $H(x_i) = h(x_i\pi(t+1))\rho(-t-1) \rightarrow h(x\pi(t+1))\rho(-t-1) = H(x)$. Hence, H is continuous. Finally we will prove that H^{-1} is continuous. Let $y \in X$ and let $\{y_i\}$ be a sequence in X such that $H(y_i) \rightarrow H(y)$. Then there exist $t, t_i \in R^+$ such that $H(y_i) = h(y_i\pi t_i)\rho(-t_i)$ and $H(y) = h(y\pi t)$. Let $s_i = \inf\{s \in R^+ : y_i\pi s \in Y\}$. We will show that $\{s_i\}$ is bounded. Suppose not. Then there is a subsequence $\{s_j\}$ of $\{s_i\}$ such that $s_j \rightarrow \infty$. If $y_i \in L^{-1}([0, \lambda])$, then $s_i \leq 1$. Hence, we may assume $1 \leq s_j$ and $y_j \notin L^{-1}([0, \lambda])$ for every j . Then $y_j\pi(s_j-1) \in L^{-1}(\lambda)$. Note that $H(y) \leftarrow H(y_j) = h(y_j\pi s_j)\rho(-s_j) = c^{-s_j}h(y_j\pi s_j)$. Since $s_j \rightarrow \infty$ and $c \in (0, 1)$ we have $c^{-s_j} \rightarrow \infty$. In order that $c^{-s_j}h(y_j\pi s_j) \rightarrow H(y)$ we must also have $h(y_j\pi s_j) \rightarrow \bar{0}$ where $\bar{0}$ is the origin in l_2 . Since h is a homeomorphism $y_j\pi s_j \rightarrow p$ so that $y_j\pi(s_j-1) \rightarrow p$. This is impossible because $y_j\pi(s_j-1) \in L^{-1}(\lambda)$ and $L(p) = 0$. Hence $\{s_i\}$ must be bounded. Without loss of generality we may suppose that $0 \leq s_i \leq t$ for every i . Then $H(y_i) = h(y_i\pi t)\rho(-t) \rightarrow h(y\pi t)\rho(-t) = H(y)$ so that $h(y_i\pi t) \rightarrow h(y\pi t)$. Since h is a homeomorphism, $y_i\pi t \rightarrow y\pi t$ and we have $y_i \rightarrow y$. Hence, H^{-1} is continuous and H is a homeomorphism of X onto $H(X) \subset l_2$.

THEOREM 7. *Let π be a semiflow on a separable metric space X such that the negative escape time function is lower semicontinuous and $\cdot\pi t$ is one-to-one for each $t \in R^+$. If π has a globally uniformly asymptotically stable critical point, then π can be embedded into a radial flow on l_2 .*

Proof. In light of Lemma 6, we need only show that $H(x\pi s) = H(x)\rho s$ for every $(x, s) \in X \times R^+$. Let $x \in X$ and $t \geq 0$ be such that $x\pi t \in Y$. Then $(x\pi s)\pi t = x\pi(t+s) \in Y$ and we have $H(x\pi s) =$

$$h((x\pi s)\pi t)\rho(-t) = h((x\pi t)\pi s)\rho(-t) = (h(x\pi t)\rho s)\rho(-t) = (h(x\pi t)\rho(-t))\rho s = H(x)\rho s.$$

COROLLARY 8. ([2, Theorem I].) *Let π be a semiflow on a separable metric space having the properties*

(i) *$x \rightarrow x\pi t$ is a homeomorphism of X onto a closed subset of X for each $t \in R^+$,*

(ii) *there is a $p \in X$ such that for any neighborhood U of p there is a $T \in R^+$ with $X\pi t \subset U$ for all $t \geq T$.*

Then π can be embedded into a radial flow on l_2 .

Proof. Clearly (i) and (ii) imply that $\cdot\pi t$ is one-to-one for all $t \in R^+$ and p is globally uniformly asymptotically stable respectively. It remains to show that (i) implies that the negative escape time function α is lower semicontinuous. Suppose that α is not lower semicontinuous. Then there exist $x \in X$, $\delta > 0$, and a sequence $\{x_i\}$ in X such that $x_i \rightarrow x$ and $\alpha(x_i) < \alpha(x) - \delta$ for every i . Thus $x_i\pi(\alpha(x) - \delta)$ is defined for every i . Then $(x_i\pi(\alpha(x) - \delta))\pi(-\alpha(x) + \delta) = x_i \rightarrow x$ so that $x \in \overline{X\pi(-\alpha(x) + \delta)} = X\pi(-\alpha(x) + \delta)$ since $X\pi t$ is closed for every $t \geq 0$. Then there exists $z \in X$ such that $z\pi(\alpha(x) - \delta) = x$. This is impossible because $\alpha(x) - \delta < \alpha(x)$ and $\alpha(x) = \inf\{-t: \text{there exists } y \in X \text{ with } y\pi t = x\}$. Therefore, we must have that α is lower semicontinuous. The desired result now follows from Theorem 7.

In the proof of Corollary 8 we showed that if $X\pi t$ is a closed subset of X for all $t \in R^+$ then the negative escape time function α is lower continuous. The converse of this is not valid. Let $X = [0, 1)$ and define $\pi: X \times R \rightarrow X$ by $x\pi t = e^{-t}x$. Evidently π is a semiflow on X . The negative escape time function is defined by

$$\alpha(x) = \begin{cases} \ln x & \text{if } x \neq 0 \\ -\infty & \text{if } x = 0. \end{cases}$$

Thus α is lower semicontinuous. However $X\pi 1 = [0, e^{-1})$ is not a closed subset of $[0, 1)$. Thus the lower semicontinuity of α does not imply that $X\pi t$ is a closed subset of X for every $t \in R^+$.

COROLLARY 9. (Theorem 5 of [4].) *Let π be a flow on a separable metric space which has a globally asymptotically stable critical point p . Then π can be embedded into a radial flow on l_2 if and only if p is globally uniformly asymptotically stable.*

Proof. Since π is a flow $x \rightarrow x\pi t$ is one-to-one for every $t \in R^+$ and $\alpha(x) = -\infty$ for every $x \in X$. If p is globally uniformly asymp-

totically stable, then, by Theorem 7, π can be embedded into a radial flow on ℓ_2 . The converse follows easily since the origin in ℓ_2 is globally uniformly asymptotically stable with respect to a radial flow.

REFERENCES

1. J. Auslander and P. Seibert, *Prolongations and stability in dynamical systems*, Ann. Inst. Fourier, Grenoble, **14** (1964), 237-268.
2. M. Edelstein, *On the homomorphic and isomorphic embeddings of a semiflow into a radial flow*, Pacific J. Math., **91** (1980), 281-291.
3. L. Janos, *A linearization of semiflows in the Hilbert space l_2* , in Topology Proceedings (Baton Rouge, La.), **2** (1977), 219-232.
4. R. McCann, *Embedding asymptotically stable dynamical systems into radial flows in l_2* , Pacific J. Math., **90** (1980), 425-429.

Received June 12, 1980 and in revised form September 2, 1980.

MISSISSIPPI STATE UNIVERSITY
MISSISSIPPI STATE, MS 39762