

FREE PRODUCTS AND ELEMENTARY EQUIVALENCE

PHILIP OLIN

It is shown that the free product operation on two groupoids preserves both elementary equivalence and elementary subsystem. An example is given showing the above results for semigroups false, thus answering in the negative a question of Feferman and Vaught.

In their important paper [3], Feferman and Vaught show, as a consequence of a stronger result, that many of the usual product operations preserve elementary equivalence. For example, if $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ are structures such that \mathfrak{A}_1 and \mathfrak{B}_1 have the same elementary first order properties (denoted $\mathfrak{A}_1 \equiv \mathfrak{B}_1$) and similarly $\mathfrak{A}_2 \equiv \mathfrak{B}_2$, then for the direct products we have $\mathfrak{A}_1 \times \mathfrak{A}_2 \equiv \mathfrak{B}_1 \times \mathfrak{B}_2$. In the footnote on page 76 of that paper they state that their methods do not apply to free products or tensor products, and they ask if these two operations preserve elementary equivalence. The answer is known to be negative for tensor products (see [2], [4]). We show here that for free products the answer is also negative for both elementary equivalence and elementary subsystem. In our counterexample $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ are semigroups, and the idea used is similar to the idea in Example 1.3 of [5].

In that same footnote, Feferman and Vaught mention a method due to Fraïssé and later developed by Ehrenfeucht [1], and they ask if this method might be applied to the problem of preserving elementary equivalence. We show here, using this method of games of Fraïssé-Ehrenfeucht, that the free product operation on groupoids preserves both elementary equivalence and elementary subsystem. A groupoid is simply a nonempty set with a binary function.

Also explicitly mentioned in that footnote is the question whether free products of groups preserve elementary equivalence. We have been unable to answer this question.

It should be noted that the definition of free product depends on the class of structures considered, so that if \mathfrak{A} and \mathfrak{B} are semigroups then their free product as semigroups is different from the free product formed with them by thinking of them as groupoids. At the end of the paper we attempt in a short space to give some motivation for the example and the proof.

The results of this paper were announced in [6].

We denote the sentence φ being true in the model \mathfrak{A} by $\mathfrak{A} \models \varphi$. If for every elementary first order sentence φ we have $\mathfrak{A} \models \varphi$ iff $\mathfrak{B} \models \varphi$ then \mathfrak{A} and \mathfrak{B} are said to be elementarily equivalent and

we write $\mathfrak{A} \equiv \mathfrak{B}$. If \mathfrak{A} is a submodel of \mathfrak{B} and if for every such φ with possibly constants from \mathfrak{A} we have $\mathfrak{A} \models \varphi$ iff $\mathfrak{B} \models \varphi$ then we say \mathfrak{A} is an elementary subsystem of \mathfrak{B} and we write $\mathfrak{A} < \mathfrak{B}$. The free product operation (over a given, understood class) is denoted by $*$. For the result for groupoids we assume some familiarity with the method of games (see [1] or [2]).

EXAMPLE. We consider the class of semigroups and now construct semigroups $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ such that $\mathfrak{A}_1 \equiv \mathfrak{B}_1, \mathfrak{A}_2 \equiv \mathfrak{B}_2$ and yet $\mathfrak{A}_1 * \mathfrak{A}_2 \not\equiv \mathfrak{B}_1 * \mathfrak{B}_2$. In fact we will have $\mathfrak{A}_1 \cong \mathfrak{B}_1$, the trivial semigroup; and $\mathfrak{A}_2 < \mathfrak{B}_2$, both being denumerably infinite.

Let \mathfrak{B} be the semigroup $\langle B, \cdot \rangle$ where B is the one-element set $\{b\}$ and $b \cdot b = b$. Of course $\mathfrak{B} < \mathfrak{B}$. We will now define a semigroup $\mathfrak{A} = \langle A, \cdot \rangle$. The generators of \mathfrak{A} will be the members (all distinct) of the set $G = \{a_i\}_{i < \omega} \cup \{c_j^i\}_{i, j < \omega}$. For each $i < \omega$ let $\{S_j^i\}_{j < \omega}$ be a list of all the subsets of G of cardinality i . Let R be the following set of relations:

$$\{c_j^i \cdot a_i \cdot y = y \cdot c_j^i \cdot a_i \mid i, j < \omega, y \in S_j^i\}.$$

Then \mathfrak{A} is obtained by starting with the free semigroup on the set G of generators and then introducing the relations in R .

Since the only relations which we have added to the free semigroup on G are of a "commuting" nature and in particular introduce no cancellation or reduction in the length of words, several properties follow. First, the indivisible members of \mathfrak{A} are exactly the members of G ; i.e., the formula $\sim(\exists x_1)(\exists x_2)(x_1 \cdot x_2 = x)$, denoted by $\psi(x)$, is satisfied in \mathfrak{A} by, and only by, the members of G . Also, every member of \mathfrak{A} is either indivisible or can be written as an indivisible times some other element; i.e.,

$$\mathfrak{A} \models (\forall y)(\exists z)[\psi(z) \wedge (z = y \vee (\exists w)(y = z \cdot w))].$$

Let $\varphi_n(x)$ denote the formula

$$(\forall y_1) \cdots (\forall y_n)(\exists z)[\psi(x) \wedge \psi(z) \wedge (\bigwedge_{i=1}^n (\psi(y_i) \rightarrow z \cdot x \cdot y_i = y_i \cdot z \cdot x))].$$

The formula "says" of x that it is indivisible and for any y_1, \dots, y_n there is an indivisible z such that, for each i , if y_i is indivisible then $z \cdot x \cdot y_i = y_i \cdot z \cdot x$. \mathfrak{A} was constructed in such a way that $\mathfrak{A} \models \varphi_n(a_n)$, the desired z being any c_j^n where S_j^n contains all the indivisible y_i 's. Furthermore, if $g_1, g_2, g_3 \in G$ then $g_1 \cdot g_2 \cdot g_3 = g_3 \cdot g_1 \cdot g_2$ in \mathfrak{A} iff this relation is already in R or $g_1 = g_2 = g_3$. We omit a proof of this; such a proof would first note that because there is no cancellation involved in the relations in R , words of length other than three could not be used to derive such a new relation, and then,

by considering the various cases as to whether g_1 is some a_i or some c_k^i and so on, one could eliminate the different possibilities.

Hence for any $g \in G$ there is a positive integer m and $g_1, \dots, g_m \in G$ such that for any $g' \in G$ there is at least one $i, 1 \leq i \leq m$, such that $g' \cdot g \cdot g_i \neq g_i \cdot g' \cdot g$. If $g = a_i$ then $m = i + 1$ and g_1, \dots, g_m all different will suffice. But the set of formulas $T = \{\varphi_n(x)\}_{n < \omega}$ is finitely satisfiable in \mathfrak{A} because, given any finite subset T' of T , if n is the largest integer such that $\varphi_n \in T'$ then clearly a_n satisfies all the members of T' .

By the Compactness Theorem for elementary first order logic there is a semigroup \mathfrak{A}' such that \mathfrak{A}' is denumerably infinite, $\mathfrak{A}' > \mathfrak{A}$, and there is some $\bar{a} \in \mathfrak{A}'$ such that for each $n < \omega$, $\mathfrak{A}' \models \varphi_n(\bar{a})$.

We will now show that $\mathfrak{B} * \mathfrak{A}$ and $\mathfrak{B} * \mathfrak{A}'$ are not elementarily equivalent and hence $\mathfrak{B} * \mathfrak{A}$ is not an elementary subsystem of $\mathfrak{B} * \mathfrak{A}'$. Let θ be the sentence

$$\begin{aligned} & (\exists v)(\exists x)(\forall y)(\exists z)(\forall u_1)(\forall u_2)(\forall u_3)\{\psi(x) \wedge \psi(z) \wedge v \cdot v = v \\ & \wedge [(y = u_1 \cdot v \cdot u_2 \cdot v \cdot u_3 \wedge \sim(\exists u_4)(\exists u_5)(u_2 = u_4 \cdot v \cdot u_5)) \\ & \rightarrow (\exists w)(\exists r)((u_2 = w \vee u_2 = w \cdot r) \wedge \psi(w) \wedge z \cdot x \cdot w = w \cdot z \cdot x)]\}. \end{aligned}$$

This sentence θ , as it will be applied below, says roughly the following: There is an idempotent (which will have to be b) and an indivisible x such that for any word y we can find an indivisible z such that for any way of writing y as $u_1 \cdot b \cdot u_2 \cdot b \cdot u_3$ with $u_2 \in A$ or $\in A'$ (as the case may be), there is a left-most indivisible factor w of u_2 such that $z \cdot x \cdot w = w \cdot z \cdot x$. We are using here the fact that $\mathfrak{B} * \mathfrak{A} \models \sim(\exists u_4)(\exists u_5)(u = u_4 \cdot b \cdot u_5)$ iff $u \in A$, and similarly with A' in place of A .

We claim $\mathfrak{B} * \mathfrak{A}' \models \theta$ and $\mathfrak{B} * \mathfrak{A} \models \sim \theta$. First, why $\mathfrak{B} * \mathfrak{A}' \models \theta$? Let v be $b \in \mathfrak{B}$ and let x be the $\bar{a} \in \mathfrak{A}'$. $\psi(\bar{a})$ holds in $\mathfrak{B} * \mathfrak{A}'$ because \bar{a} satisfies $\varphi_1(x)$ in \mathfrak{A}' . Now suppose $y \in \mathfrak{B} * \mathfrak{A}'$ is given. We can assume y is of the form $h_1 \cdot b \cdot t_1 \cdot b \cdot t_2 \cdot b \cdot \dots \cdot b \cdot t_m \cdot b \cdot h_2$ where $m \geq 1$, each $t_i \in \mathfrak{A}'$ and h_1, h_2 are each either b or in \mathfrak{A}' ; otherwise the antecedent of $[\dots \rightarrow \dots]$ in θ could not be satisfied and we would easily finish. So for each t_i there is an indivisible $w_i \in \mathfrak{A}'$ such that either $t_i = w_i$ or $t_i = w_i \cdot r_i$ for some $r_i \in \mathfrak{A}'$. (Recall every member of \mathfrak{A} has this property and $\mathfrak{A} \equiv \mathfrak{A}'$.) We know $\mathfrak{A}' \models \varphi_m(\bar{a})$. So replacing y_i in this formula by w_i we get an indivisible $z \in \mathfrak{A}'$ such that for all $i, 1 \leq i \leq m, z \cdot \bar{a} \cdot w_i = w_i \cdot z \cdot \bar{a}$. Now for any u_1, u_2, u_3 in $\mathfrak{B} * \mathfrak{A}'$, if the antecedent of $[\dots \rightarrow \dots]$ is satisfied, it must be the case that for some $i, u_2 = t_i$. So let w be w_i , let r be r_i and we are done.

We now wish to show $\mathfrak{B} * \mathfrak{A} \models \sim \theta$. Suppose not. So we get v and x . Since $v \cdot v = v$ we must have $v = b$ since b is the only idempotent in $\mathfrak{B} * \mathfrak{A}$. Since $\mathfrak{B} * \mathfrak{A} \models \psi(x)$ we must have $x = g \in G$.

As remarked above, there is an m and $g_1, \dots, g_m \in G$ such that for any $g' \in G$, $\mathfrak{A} \models \sim \bigwedge_{i=1}^m (g' \cdot g \cdot g_i = g_i \cdot g' \cdot g)$. Let $y = b \cdot g_1 \cdot b \cdot g_2 \cdot \dots \cdot b \cdot g_m \cdot b$. So we then get z . Since $\mathfrak{B} * \mathfrak{A} \models \psi(z)$, we must have $z \in G$; say $z = g'$. Say i_0 is such that $g' \cdot g \cdot g_{i_0} \neq g_{i_0} \cdot g' \cdot g$. If $1 < i_0 < m$, let u_2 be g_{i_0} , let u_1 be $b \cdot g_1 \cdot b \cdot \dots \cdot b \cdot g_{i_0-1}$ and let u_3 be $g_{i_0+1} \cdot b \cdot \dots \cdot b \cdot g_m \cdot b$. If $i_0 = 1$, let u_2 be g_{i_0} , let u_1 be b and let u_3 be as above. The case $i_0 = m$ is similar. Clearly the antecedent of $[\dots \rightarrow \dots]$ in θ is satisfied. So we get w and r . It must be the case that $w = u_2 (= g_{i_0})$. But $z \cdot x \cdot w (= g' \cdot g \cdot g_{i_0}) \neq w \cdot z \cdot x (= g_{i_0} \cdot g' \cdot g)$, contradicting $\mathfrak{B} * \mathfrak{A} \models \theta$. So $\mathfrak{B} * \mathfrak{A} \models \sim \theta$.

We remark that in the logical hierarchy of formulas, θ is a Σ_6 -sentence. It seems likely that an example showing that free products do not preserve elementary equivalence could be constructed in which the sentence θ is Σ_4 , or perhaps Σ_3 . It also seems likely that Σ_3 or at least Σ_2 equivalence is preserved by free products, and the method of games of the next result should suffice to show it.

A groupoid $\mathfrak{A} = \langle A, \cdot \rangle$ is a nonempty set A and a function from $A \times A$ into A ; and $*$ is now in the class of groupoids. We wish to show that if $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ are groupoids, $\mathfrak{A}_1 \equiv \mathfrak{B}_1, \mathfrak{A}_2 \equiv \mathfrak{B}_2$, then $\mathfrak{A}_1 * \mathfrak{A}_2 \equiv \mathfrak{B}_1 * \mathfrak{B}_2$. The method to be used is the method of games [1], [2], and the winning strategy for player II is similar to that which is used in showing that, as linearly ordered sets, $\omega \equiv \omega + \omega^* + \omega$. We wish to thank the referee whose questions and comments led to, among other things, improvement in the proof of the following theorem.

THEOREM. *If $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ are groupoids and $\mathfrak{A}_1 \equiv \mathfrak{B}_1, \mathfrak{A}_2 \equiv \mathfrak{B}_2$ then $\mathfrak{A}_1 * \mathfrak{A}_2 \equiv \mathfrak{B}_1 * \mathfrak{B}_2$.*

Proof. We can assume $A_1 \cap A_2 = B_1 \cap B_2 = \emptyset$. Let n be a fixed positive integer. We will describe a winning strategy for player II in the game $G_n(\mathfrak{A}_1 * \mathfrak{A}_2, \mathfrak{B}_1 * \mathfrak{B}_2)$. We can assume that the nonlogical constants of the language are only $=$ (interpreted always as identity) and a three-place predicate P , where $P(a, b, c)$ means $a \cdot b = c$. This allows us to avoid considering terms. In the game, ${}_i x, {}_j x, \dots, {}_n x$ will be chosen from $\mathfrak{A}_1 * \mathfrak{A}_2$ and ${}_i y, {}_j y, \dots, {}_n y$ from $\mathfrak{B}_1 * \mathfrak{B}_2$. II wins iff for all $1 \leq i, j, k \leq n$

$${}_i x \cdot {}_j x = {}_k x \quad \text{iff} \quad {}_i y \cdot {}_j y = {}_k y$$

and

$${}_i x = {}_j x \quad \text{iff} \quad {}_i y = {}_j y .$$

We shall need II's winning strategy in $G_m(\mathfrak{A}_1, \mathfrak{B}_1)$ and in $G_m(\mathfrak{A}_2, \mathfrak{B}_2)$, where $m = \sum_{i=1}^n 2^{(3^i-2)}$. We require some notation. For each ${}_i x$ in

$\mathfrak{A}_1 * \mathfrak{A}_2$, let ${}_i x = {}_i x_1^1$. Then by induction, for $j \geq 1$ and $k \geq 1$, if ${}_i x_k^j \in \mathfrak{A}_1 * \mathfrak{A}_2 - (\mathfrak{A}_1 \cup \mathfrak{A}_2)$, let ${}_i x_k^j = {}_i x_{2k-1}^{j+1} \cdot {}_i x_{2k}^{j+1}$. This decomposition is unique because of the definition of free product of groupoids. We say that each ${}_i x_k^j$ is a factor of ${}_i x_{k'}^{j'}$ if $j \geq j'$ and $2^{(j-j')(k'-1)} + 1 \leq k \leq 2^{(j-j')(k')}$. And we say ${}_i x_k^j$ has depth j in ${}_i x$. Similar notation is adopted for each ${}_i y$. The m defined above is the largest number of members of, say, A_1 which can appear among ${}_i x, \dots, {}_n x$ at a depth $\leq 3^n$ in ${}_i x$ or $\leq 3^{n-1}$ in ${}_2 x$ or \dots or ≤ 3 in ${}_n x$. It is these members which are, in some sense, directly threatened by player I in the game. It will be convenient to assume first that A_1, A_2, B_1, B_2 are all infinite. We now begin to describe II's strategy.

First of all, if I chooses ${}_i x$ then the ${}_i y$ that II will choose will have exactly the same "form" as ${}_i x$ — i.e., ${}_i y_k^j \in B_1$ iff ${}_i x_k^j \in A_1$, and ${}_i y_k^j \in B_2$ iff ${}_i x_k^j \in A_2$. Similarly if I chooses ${}_i y$.

Secondly, say I chooses ${}_i x$ and ${}_i x_{k_1}^{j_1}, \dots, {}_i x_{k_p}^{j_p}$ is a list of those factors of ${}_i x$ which are at a depth $\leq 3^n$ in ${}_i x$ (i.e., each $j_i \leq 3^n$) and which are in A_1 . Then II chooses ${}_i y_{k_1}^{j_1}, \dots, {}_i y_{k_p}^{j_p}$ from B_1 according to his winning strategy in $G_m(\mathfrak{A}_1, \mathfrak{B}_1)$. Note that $p \leq 2^{(3^n-2)}$. A similar procedure is followed for such factors from A_2 and B_2 . If I had chosen ${}_i y$, again the procedure is similar. The method II uses for completing his choice of factors of ${}_i y$ (if ${}_i y$ is not already completely defined) will be specified later. It will not affect some parts of his later strategy, which it is convenient to give now. Say I chooses ${}_2 y$. Say ${}_2 y_{k_1}^{j_1}, \dots, {}_2 y_{k_r}^{j_r}$ is a list of those factors of ${}_2 y$ which are in B_1 and which are at a depth $\leq 3^{n-1}$ in ${}_2 y$. Note $r \leq 2^{(3^{n-1}-2)}$. Then II chooses ${}_2 x_{k_1}^{j_1}, \dots, {}_2 x_{k_r}^{j_r}$ from A_1 by using his winning strategy in $G_m(\mathfrak{A}_1, \mathfrak{B}_1)$ and taking into account the choices made in this game when ${}_i x$ and ${}_i y$ were discussed above. (So this involves choices number $p+1$ to $p+r$ in $G_m(\mathfrak{A}_1, \mathfrak{B}_1)$.) Similarly for factors of depth $\leq 3^{n-1}$ in ${}_i x$ and ${}_i y$ which are from A_2 and B_2 , and also similarly if I had chosen ${}_2 x$. Player II continues in this way for the rest of the game, choosing factors which are to be in $A_1 \cup A_2$ or $B_1 \cup B_2$ and at depth $\leq 3^{n-d+1}$ in ${}_i x$ according to his given winning strategies in the games G_m and in the light of earlier choices in these games. Note that the choice of m ensures that II has enough "room to work in". Note also that the winning strategy for II in these games $G_m(\mathfrak{A}_1, \mathfrak{B}_1)$ and $G_m(\mathfrak{A}_2, \mathfrak{B}_2)$ certainly includes maintaining equalities and inequalities — i.e., for those choices of factors specified above, we have ${}_i x_a^c = {}_i x_a^c$ iff ${}_i y_a^c = {}_i y_a^c$. This will be required later. Furthermore, at the end of the game, since the above winning strategies were used, we will have: If $i, j, k \leq n$ then ${}_i x, {}_j x, {}_k x$ are all in A_1 iff ${}_i y, {}_j y, {}_k y$ are all in B_1 , and in this case ${}_i x = {}_j x$ iff ${}_i y = {}_j y$ and ${}_i x \cdot {}_j x = {}_k x$ iff ${}_i y \cdot {}_j y = {}_k y$. Similarly for A_2 and B_2 .

We define for each $r, 0 \leq r \leq n$, conditions $K_r(1), \dots, K_r(4)$ as follows:

$K_r(1)$: Suppose $1 \leq s < r$ and $j_1 + j_2 - 1 \leq 3^{n-r+1}$. Suppose ${}_r x_{k_1}^{j_1}$ and ${}_s x_{k_2}^{j_2}$ are defined and equal. Then ${}_r y_{k_1}^{j_1}$ and ${}_s y_{k_2}^{j_2}$ are also defined and equal.

$K_r(2)$: Suppose $j_1 + j_2 - 1 \leq 3^{n-r+1}$ and suppose ${}_r x_{k_1}^{j_1}$ and ${}_r x_{k_2}^{j_2}$ are defined and equal. Then ${}_r y_{k_1}^{j_1}$ and ${}_r y_{k_2}^{j_2}$ are also defined and equal.

$K_r(3)$: Replace "equal" in $K_r(1)$ by "unequal".

$K_r(4)$: Replace "equal" in $K_r(2)$ by "unequal". Clearly $K_0(1), \dots, K_0(4)$ are vacuously satisfied.

Assume that $1 \leq t \leq n$ and that ${}_1 x, \dots, {}_{t-1} x$ and ${}_1 y, \dots, {}_{t-1} y$ have been completely specified, following that part of II's strategy already indicated above and, for the rest (if any) of the factors in these elements, in such a way that for each $r, 0 \leq r \leq t-1, K_r(1), \dots, K_r(4)$ are all satisfied. Suppose I chooses ${}_i x$. Then for those factors of ${}_i y$ which are to be in $B_1 \cup B_2$ and at a depth $\leq 3^{n-(t-1)} = 3^{n-t+1}$ in ${}_i y$, II specifies these factors according to that part of his strategy already given above. We now wish to show that II can complete his definition of ${}_i y$ so that $K_t(1), \dots, K_t(4)$ are all satisfied.

Consider those factors ${}_i y_k^j$ of ${}_i y$ which are in $B_1 \cup B_2$ and with $j \leq 3^{n-t+1}$ — i.e., those just specified by II. Conditions $K_t(1)$ and $K_t(3)$ might require some ${}_i y_k^j$ equal or unequal to some ${}_r y_d^c, r < t$ and $a + c - 1 \leq 3^{n-t+1}$; and this in turn might mean that there is a factor ${}_i y_k^j$ of ${}_i y_k^j$, which is a member of $B_1 \cup B_2$, has $j \leq 3^{n-t+1}$ (and thus was specified by II already) and which, if we are to have ${}_i y_k^j$ equal or unequal to ${}_r y_d^c$, will have to be equal or unequal (as the case may be) to ${}_r y_{k_1}^{j_1}$, which is a factor of ${}_r y_d^c$ and a member of $B_1 \cup B_2$. Is this equality or inequality, needed for $K_t(1)$ or $K_t(3)$, satisfied? Assume it is equality we need. We have $a + c - 1 \leq 3^{n-t+1}, j \leq 3^{n-t+1}$ and $c - j_1 = a - j$. So $j_1 = c + j - a \leq c + j$. But $c \leq c + a - 1 \leq 3^{n-t+1}$. So $j_1 \leq 3^{n-t+1} + 3^{n-t+1} \leq 3^{n-(t-1)+1} \leq 3^{n-r+1}$. Hence ${}_r x_{k_1}^{j_1}$ and ${}_r y_{k_1}^{j_1}$ were "earlier moves" in the games G_m being played and so, since ${}_i x_k^j = {}_r x_d^c$ and thus ${}_i x_k^j = {}_r x_{k_1}^{j_1}$, player II, as required by his winning strategies in the games G_m , chose ${}_i y_k^j = {}_r y_{k_1}^{j_1}$. We have shown that the part of II's strategy already given does not conflict with conditions $K_t(1)$ and $K_t(3)$. The check that there is also no conflict with $K_t(2)$ and $K_t(4)$ is simpler and we do not give it.

We will now show how the rest (if any) of ${}_i y$ is to be defined by II. Let ${}_i x_{k_1}^{j_1}, \dots, {}_i x_{k_p}^{j_p}$ be a list of those factors of ${}_i x$ which satisfy the hypothesis of condition $K_t(1)$ and such that no member of this list is a factor in ${}_i x$ of any other (and hence there is no "overlap" among them at all, in the sense that they have no factors in common). It follows that every ${}_i x_k^j$ which (together with some ${}_s x_{k'}^{j'}, s < t$) satisfies the hypothesis of $K_t(1)$ is a factor of one of the members of this list.

Say ${}_i x_{k_1}^{j_1} = {}_r x_u^i, r < t, j_1 + i - 1 \leq 3^{n-t+1}$. For those ${}_i x_{k'}^{j'}$ which are

factors of ${}_i x_{k_1}^{j_1}$, are in $A_1 \cup A_2$ and satisfy $j' > 3^{n-t+1}$, II defines ${}_i y_{k_1}^{j_1}$, so that ${}_i y_{k_1}^{j_1} = {}_r y_u^i$. There is no difficulty in doing this. However, suppose ${}_i x_k^j$ is a factor of ${}_i x_{k_1}^{j_1}$ and suppose ${}_i x_k^j$ and ${}_s y_b^a$ ($s < t$) satisfy the hypothesis of $K_t(1)$. Since ${}_i y_k^j$ has just been defined (since it is a piece of ${}_i y_{k_1}^{j_1}$), does it satisfy the conclusion of $K_t(1)$? We have $j + a - 1 \leq 3^{n-t+1}$. We require ${}_i y_k^j = {}_s y_b^a$. Since ${}_i x_{k_1}^{j_1} = {}_r x_u^i$ and since they have exactly the same form, let ${}_i x_u^{i'}$ be that factor of ${}_r x_u^i$ which corresponds to ${}_i x_k^j$ under the correspondence given by this "sameness of form". Of course ${}_r x_u^{i'} = {}_i x_k^j$ and $i' - i = j - j_1$. In the same way, ${}_i y_k^j = {}_r y_u^{i'}$. But $a + j_1 + (j - j_1) - 1 = a + j - 1 \leq 3^{n-t+1}$ and this implies $a + (j - j_1) \leq 3^{n-t+1}$. Also $j_1 + i - 1 \leq 3^{n-t+1}$ implies $i \leq 3^{n-t+1}$. Hence $a + i' = a + (i' - i) + i = a + (j - j_1) + i \leq 2 \cdot 3^{n-t+1} \leq 3^{n-(t-1)+1} \leq 3^{n-\max(s,r)+1}$. So by the induction hypothesis for $K_{\max(s,r)}(1)$ if $r \neq s$ or for $K_s(2)$ if $r = s$, we get ${}_s y_b^a = {}_r y_u^{i'}$. So ${}_s y_b^a = {}_i y_k^j$.

Suppose ${}_i x_{k_1}^{j_1} = {}_r x_u^i$ (as above) and in addition ${}_i x_{k_1}^{j_1} = {}_v x_d^c$ with $v < t$ and $j_1 + c - 1 \leq 3^{n-t+1}$. Then by an argument similar to the above, we could show ${}_v y_d^c = {}_r y_u^i$ and so no conflict arises here.

Player II now repeats the above procedure for ${}_i x_{k_1}^{j_1}$ on ${}_i x_{k_2}^{j_2}, \dots, {}_i x_{k_p}^{j_p}$. Since, as remarked above, no two of these overlap there is no difficulty in making the definitions to satisfy $K_t(1)$; and again as above, factors of members of this list are automatically taken care of.

We now wish to consider condition $K_t(2)$. Suppose ${}_i x_{k_1}^{j_1}$ and ${}_i x_{k_2}^{j_2}$ satisfy the hypothesis of $K_t(2)$ (i.e., they are equal and $j_1 + j_2 - 1 \leq 3^{n-t+1}$). Suppose further that there is no pair of factors of ${}_i x$ satisfying the hypothesis of $K_t(2)$ and with either of these factors being factors of ${}_i x_{k_1}^{j_1}$ or of ${}_i x_{k_2}^{j_2}$ — i.e., this latter pair is "minimal" with respect to the hypothesis of $K_t(2)$. If there are any pairs satisfying the hypothesis of $K_t(2)$ then there is a minimal pair because: If ${}_i x_{k_1}^{j_1}, {}_i x_{k_2}^{j_2}$ and ${}_i x_{k_3}^{j_3}, {}_i x_{k_4}^{j_4}$ are different pairs satisfying the hypothesis of $K_t(2)$ and if ${}_i x_{k_1}^{j_1}$ is a factor of ${}_i x_{k_3}^{j_3}$ then if ${}_i x_{k_4}^{j_4}$ were a factor of ${}_i x_{k_2}^{j_2}$, we would get ${}_i x_{k_2}^{j_2}$ equal to a proper factor of itself and this is impossible in a free product. So we can "work our way down in depth" and consider a minimal pair ${}_i x_{k_1}^{j_1}, {}_i x_{k_2}^{j_2}$. Player II must arrange ${}_i y_{k_1}^{j_1} = {}_i y_{k_2}^{j_2}$. There are several ways in which the parts of ${}_i y$ defined in satisfying $K_t(1)$ might conflict with this desired result.

Suppose ${}_i y_{k'}^{j'} = {}_r y_u^{i'}$, $i' + j' - 1 \leq 3^{n-t+1}$, $r < t$, and ${}_i y_{k''}^{j''} = {}_s y_u^{i''}$, $j'' + i'' - 1 \leq 3^{n-t+1}$, $s < t$ were arranged in satisfying $K_t(1)$. Say ${}_i y_{k_1}^{j_1}$ is a factor of ${}_i y_{k'}^{j'}$ and ${}_i y_{k_2}^{j_2}$ is a factor of ${}_i y_{k''}^{j''}$; hence ${}_i y_{k_1}^{j_1}$ and ${}_i y_{k_2}^{j_2}$ have already been completely defined. Let ${}_r y_b^a$ be the factor of ${}_r y_u^{i'}$ which corresponds (under the sameness-of-form correspondence between ${}_r y_u^{i'}$ and ${}_i y_{k'}^{j'}$) to ${}_i y_{k_1}^{j_1}$. Similarly for ${}_s y_b^a$ and ${}_i y_{k_2}^{j_2}$. It suffices to show ${}_r y_b^a = {}_s y_b^a$. We have $i' - a = j' - j_1$, $i'' - c = j'' - j_2$, $i' + j' - 1 \leq 3^{n-t+1}$, $i'' + j'' - 1 \leq 3^{n-t+1}$, and $j_1 + j_2 - 1 \leq 3^{n-t+1}$. So $i' \leq 3^{n-t+1}$, $i'' \leq 3^{n-t+1}$ and $(j_1 - j') + j' + (j_2 - j'') - j'' - 1 \leq 3^{n-t+1}$. Hence $(j_1 - j') +$

$(j_2 - j'') - 1 \leq 3^{n-t+1}$. But then $a + c - 1 = i' + j_1 - j' + i'' + j_2 - j'' - 1 \leq 3 \cdot 3^{n-t+1} = 3^{n-(t-1)+1} \leq 3^{n-\max(r,s)+1}$. So by the induction hypothesis, using $K_{\max(r,s)}(1)$ if $s \neq r$ and $K_s(2)$ if $s = r$, we get ${}_r y_s^a = {}_s y_s^c$.

Now suppose things are as above, except ${}_t y_{k'}^{j'}$ is a factor of ${}_t y_{k_2}^{j_2}$ instead of the other way round. By computations very similar to those above we would get ${}_t y_{k'}^{j'}$ equal to its corresponding (under the correspondence determined by ${}_t x_{k_1}^{j_1}$ being equal to ${}_t x_{k_2}^{j_2}$) image in ${}_t y_{k_1}^{j_1}$. Player II then defines those ${}_t y_s^a$ which are factors of ${}_t y_{k_2}^{j_2}$, which are required to be in $B_1 \cup B_2$, and which are not already defined, by making them equal to the corresponding (same correspondence) factors in ${}_t y_{k_1}^{j_1}$, all of which were defined earlier.

Now suppose things are as above, except ${}_t y_{k'}^{j'}$ is a factor of ${}_t y_{k_1}^{j_1}$ and ${}_t y_{k'}^{j'}$ is a factor of ${}_t y_{k_2}^{j_2}$. There are now several subcases to consider, depending on how these factors overlap under the correspondence determined by ${}_t x_{k_1}^{j_1}$ being equal to ${}_t x_{k_2}^{j_2}$. All of them involve computations similar to the one given above, and we omit them. Once the definitions made in satisfying $K_t(1)$ are seen not to conflict with ${}_t y_{k_1}^{j_1}$ being equal to ${}_t y_{k_2}^{j_2}$, II can define those factors of, say, ${}_t y_{k_1}^{j_1}$ which are to be in $B_1 \cup B_2$ and which are supposed to equal factors already defined in ${}_t y_{k_2}^{j_2}$. One possibility remains. A certain factor of ${}_t y_{k_1}^{j_1}$ is to be a member of $B_1 \cup B_2$ and equal to its corresponding factor in ${}_t y_{k_2}^{j_2}$ — but neither has been defined by any of the above considerations. In this case we use the assumption that B_1 and B_2 are each infinite, and II chooses any member (from B_1 or B_2 , whichever is needed so that ${}_t y$ and ${}_t x$ will have the same form) which is completely new — i.e., which appears nowhere in ${}_t x, \dots, {}_t x, {}_t y, \dots, {}_{t-1} y$ and that part of ${}_t y$ so far defined. This completes the definition of ${}_t y_{k_1}^{j_1}$ and ${}_t y_{k_2}^{j_2}$.

Other minimal pairs satisfying the hypothesis of $K_t(2)$ are handled similarly. We then consider pairs which are minimal in the sense above, but with respect only to those pairs not yet considered. The arguments are analogous, and II proceeds to define as much of ${}_t y$ as is required to satisfy $K_t(2)$.

We have thus defined part of ${}_t y$ and at the same time shown that $K_t(1)$ and $K_t(2)$ are satisfied. For the remaining (if any) factors of ${}_t y$ which are to be members of $B_1 \cup B_2$ we again use the assumption that B_1 and B_2 are infinite and II chooses completely new elements. It follows that $K_t(3)$ and $K_t(4)$ are thus satisfied.

If player I had picked ${}_t y$ then $K_t(1), \dots, K_t(4)$ and II's strategy are gotten by interchanging x and y .

Now suppose not all of A_1, A_2, B_1, B_2 are infinite. If A_1 is finite, since $\mathfrak{A}_1 \cong \mathfrak{B}_1$, we get $f: \mathfrak{A}_1 \cong \mathfrak{B}_1$. If in addition A_2 is finite, we get $\mathfrak{A}_1 * \mathfrak{A}_2 \cong \mathfrak{B}_1 * \mathfrak{B}_2$. So assume A_2 (and hence also B_2) is infinite. Then player II's winning strategy is modified so that if I chooses ${}_t x$ and

${}_i x_i^j \in A_1$ then II defines ${}_i y_k^j = f({}_i x_k^j)$, and similarly if I chooses ${}_i y$. It can be shown that all of the conclusions above are obtained also in this situation.

Now suppose the game is over and ${}_1 x, \dots, {}_n x, {}_1 y, \dots, {}_n y$ have been chosen. We want to show, for $1 \leq i, j, k \leq n$, that ${}_i x \cdot {}_j x = {}_k x$ iff ${}_i y \cdot {}_j y = {}_k y$ and ${}_i x = {}_j x$ iff ${}_i y = {}_j y$. As noted above, if ${}_i x, {}_j x, {}_k x$ are all in A_1 then ${}_i y, {}_j y, {}_k y$ are all in B_1 , and conversely, and the result then follows from II's winning strategy in $G_m(\mathfrak{A}_1, \mathfrak{B}_1)$. Similarly for A_2 and B_2 . So now assume this is not the case. There are now several cases to consider; we discuss two of them.

(i) Suppose $i < k < j$ and ${}_i x \cdot {}_j x = {}_k x$. So ${}_k x_1^2 = {}_i x_1^1$ and ${}_k x_2^2 = {}_j x_1^1$. Since $j \leq n$ and $k \leq n$ we have $3^{n-k+1} \geq 2$ and $3^{n-j+1} \geq 2$. And so $2 + 1 - 1 \leq 3^{n-k+1}$ and $2 + 1 - 1 \leq 3^{n-j+1}$. Thus conditions $K_k(1), K_k(3), K_j(1), K_j(3)$ ensure that ${}_k y_1^2 = {}_i y_1^1$ and ${}_k y_2^2 = {}_j y_1^1$, and thus ${}_k y = {}_i y \cdot {}_j y$.

(ii) Suppose $i > k$ and ${}_i x \cdot {}_i x = {}_k x$. So ${}_k x_1^2 = {}_i x_1^2 = {}_i x$. Again $k \leq n$, so $3 \leq 3^{n-k+1}$, so $2 + 2 - 1 \leq 3^{n-k+1}$. Thus condition $K_k(2)$ and $K_k(4)$ ensured that, when ${}_k x$ and ${}_k y$ were chosen, we had ${}_k x_1^2 = {}_k x_2^2$ iff ${}_k y_1^2 = {}_k y_2^2$. Then, as in case (i) above, conditions $K_i(1)$ and $K_i(3)$ ensure ${}_i x = {}_k x_1^2 = {}_k x_2^2$ iff ${}_i y_1^2 = {}_k y_2^2 = {}_i y$.

The other cases are no more difficult.

REMARK. For any positive integer n , let m be defined (as a function of n) as in the proof of the above theorem. Let \equiv_p mean equivalence with respect to sentences with at most p variables. Then in fact the above proof shows that:

(1) if $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ are all infinite,

$$\mathfrak{A}_1 \equiv_m \mathfrak{B}_1, \mathfrak{A}_2 \equiv_m \mathfrak{B}_2, \text{ then } \mathfrak{A}_1 * \mathfrak{A}_2 \equiv_n \mathfrak{B}_1 * \mathfrak{B}_2.$$

(2) if $\mathfrak{A}_1 \cong \mathfrak{B}_1, \mathfrak{A}_2$ and \mathfrak{B}_2 are infinite, and

$$\mathfrak{A}_2 \equiv_m \mathfrak{B}_2 \text{ then } \mathfrak{A}_1 * \mathfrak{A}_2 \equiv_n \mathfrak{B}_1 * \mathfrak{B}_2.$$

It seems likely that these last results could be strengthened — in particular by weakening the hypotheses.

COROLLARY. If $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{B}_1, \mathfrak{B}_2$ are groupoids, $\mathfrak{A}_2 < \mathfrak{B}_2, \mathfrak{A}_1 < \mathfrak{B}_1$, then $\mathfrak{A}_1 * \mathfrak{A}_2 < \mathfrak{B}_1 * \mathfrak{B}_2$.

The proof of the corollary is essentially the same as that for the theorem, except that we start with ${}_1 x = {}_1 y, \dots, {}_p x = {}_p y$ for some fixed $p < n$, and the first part of II's strategy is modified to use II's winning strategy gotten from the games appropriate for $\mathfrak{A}_1 < \mathfrak{B}_1$ and $\mathfrak{A}_2 < \mathfrak{B}_2$.

What follows is a short attempt to motivate intuitively the

above results. The major reason the preservation result is true for groupoids is that, because of the lack of an associative law, factors of an $x \in \mathfrak{A} * \mathfrak{B} - (\mathfrak{A} \cup \mathfrak{B})$ which are inside a sufficient number of brackets cannot be "connected" with x in a game with only n rounds. For example, to "state" that ${}_1x_i^j$ is a factor of ${}_1x$, player I would need at least j rounds. If G_n is being played and $n < j$, player II knows that I cannot do it. So when ${}_1x$, in the game G_n , has been chosen, only members of the original groupoids at a depth $\leq 3^n$ in ${}_1x$ are "threatened" by I; for the others it suffices that II maintain certain equalities and inequalities.

However, for semigroups an element, say $a_1 \cdot b_1 \cdot a_2 \cdot b_2 \cdot \dots \cdot a_n \cdot b_n$ in $\mathfrak{A} * \mathfrak{B}$, does not depend on the bracketing. And the b_i 's are all equally and quickly "accessible". Thus in round 1, player II commits himself to some choice and in round 2 player I can then present II with an arbitrarily large finite subset of elements, any one of which is accessible in 2 or 3 more rounds. The above counterexample for semigroups takes advantage of this, as well as the idea in Example 1.3 of [5].

REFERENCES

1. A. Ehrenfeucht, *An application of games to the completeness problem for formalized theories*, Fund. Math., **49** (1961), 129-141.
2. S. Feferman, *Infinitary Properties, Local Functors, and Systems of Ordinal Functions*, Conference in Mathematical Logic — London 1970, Lecture Notes in Mathematics, no. 255, Springer-Verlag, 1972, 63-97.
3. S. Feferman and R. L. Vaught, *The first order properties of products of algebraic systems*, Fund. Math., **47** (1959), 57-103.
4. P. Olin, *Direct multiples and powers of modules*, Fund. Math., **73** (1971), 113-124.
5. ———, *Products of two-sorted structures*, J. Symbolic Logic, **37** (1972), 75-80.
6. ———, *Free products and elementary equivalence*, Notices of Amer. Math. Soc., **20** (1973), A-284.

Received January 15, 1973. This work was partially supported by the National Research Council of Canada.

YORK UNIVERSITY, TORONTO, CANADA