

## EXCEPTIONAL 3/2-TRANSITIVE PERMUTATION GROUPS

D. S. PASSMAN

**Solvable 3/2-transitive permutation groups were previously classified to within a finite number of exceptions. In this paper the exceptional groups are determined. They have degrees  $3^2$ ,  $5^2$ ,  $7^2$ ,  $11^2$ ,  $17^2$  and  $3^4$ . In addition, these groups are shown to have no transitive extensions.**

There are three families of groups which play a special role here. Let  $q$  be a prime. We let  $\mathcal{S}(q^n)$  denote the set of all semilinear transformations on the finite field  $GF(q^n)$ . Thus  $\mathcal{S}(q^n)$  consists of all transformations

$$x \rightarrow ax^\sigma + b$$

with  $a, b \in GF(q^n)$ ,  $a \neq 0$  and  $\sigma$  a field automorphism. Clearly this is a solvable group, doubly transitive on  $GF(q^n)$ .

We let  $\mathcal{S}_0(q^n)$  be the group acting on a 2-dimensional space over  $GF(q^n)$  which contains the transformations

$$(x, y) \rightarrow (x, y) \begin{pmatrix} a & 0 \\ 0 & \pm a^{-1} \end{pmatrix} + (b, c)$$

and

$$(x, y) \rightarrow (x, y) \begin{pmatrix} 0 & a \\ \pm a^{-1} & 0 \end{pmatrix} + (b, c)$$

with  $a, b, c \in GF(q^n)$  and  $a \neq 0$ . We see easily that  $\mathcal{S}_0(q^n)$  is solvable and if  $q \neq 2$  then it acts 3/2-transitively on the 2-dimensional space.

Finally we let  $\Gamma(q^n)$  denote the set of all functions of the form

$$x \rightarrow \frac{ax^\sigma + b}{cx^\sigma + d}$$

with  $a, b, c, d \in GF(q^n)$ ,  $ad - bc \neq 0$  and  $\sigma$  a field automorphism. These functions permute the set  $GF(q^n) \cup \{\infty\}$  and  $\Gamma(q^n)$  is triply transitive. Clearly  $\Gamma(q^n)_\infty = \mathcal{S}(q^n)$  is solvable. Let  $\bar{\Gamma}(q^n)$  denote the subgroup of  $\Gamma(q^n)$  consisting of these functions of the form

$$x \rightarrow \frac{ax + b}{cx + d}$$

with  $ad - bc$  a nonzero square in  $GF(q^n)$ .

The following results are proved here.

**THEOREM A.** *Let  $\mathfrak{G}$  be a linear group acting on vector space  $\mathfrak{V}$  of order  $q^n$ . Suppose that  $\mathfrak{G}$  acts half-transitively but not semi-regularly on  $\mathfrak{V}^\#$ . If  $\mathfrak{G}$  is primitive as a linear group then*

- (i)  $O_p(\mathfrak{G})$  is cyclic for  $p > 2$ .
- (ii) The Frattini subgroup  $\Phi(O_2(\mathfrak{G}))$  is cyclic and

$$[O_2(\mathfrak{G}) : \Phi(O_2(\mathfrak{G}))] \leq 2^8.$$

**THEOREM B.** *Let  $\mathfrak{G}$  be a solvable 3/2-transitive permutation group. Then with suitable identification,  $\mathfrak{G}$  satisfies one of the following.*

- (i)  $\mathfrak{G}$  is a Frobenius group.
- (ii)  $\mathfrak{G} \subseteq \mathcal{S}(q^n)$
- (iii)  $\mathfrak{G} = \mathcal{S}_0(q^n)$  or
- (iv)  $\mathfrak{G}$  has degree  $3^2, 5^2, 7^2, 11^2, 17^2$  or  $3^4$ .

The exceptions of (iv) above do in fact exist. If  $\deg \mathfrak{G} \neq 17^2$  then we can take  $\mathfrak{G}$  to be an exceptional solvable doubly transitive group, while if  $\deg \mathfrak{G} = 17^2$  then we construct this group explicitly and show that it has order  $96 \cdot 17^2$ .

**THEOREM C.** *Let  $\mathfrak{G}$  be a 5/2-transitive permutation group and suppose that the stabilizer of a point is solvable. Then with suitable identification we have one of the following*

- (i)  $\mathfrak{G}$  is a Zassenhaus group or
- (ii)  $\Gamma(q^n) \cong \mathfrak{G} > \bar{\Gamma}(q^n)$ .

The main result here is Theorem B. Theorem A isolates that part of the proof in which solvability is not assumed. Theorem C follows immediately from the results of [8] and the fact that these exceptional groups have no transitive extensions.

**1. Preliminaries.** We will be concerned here with linear groups  $\mathfrak{G}$  which act half-transitively but not semi-regularly on the set  $\mathfrak{V}^\#$  of nonzero vectors. This implies (see [11], Th. 10.4) that  $\mathfrak{G}$  acts irreducibly on  $\mathfrak{V}$ . There are thus two possibilities according to whether  $\mathfrak{G}$  is primitive or imprimitive as a linear group. The latter case is completely classified in Theorem 4.2 of [7] which we restate below for convenience.

**THEOREM 1.1.** *Let  $\mathfrak{G}$  act faithfully on vector space  $\mathfrak{V}$  over  $GF(q)$  and let  $\mathfrak{G}$  act half-transitively but not semi-regularly on  $\mathfrak{V}^\#$ . If  $\mathfrak{G}$  is imprimitive as a linear group, then  $\mathfrak{G}$  satisfies one of the following*

- (i)  $\mathfrak{G} = \mathcal{S}_0(q^n)$  with  $q \neq 2$  and  $n$  an integer.

- (ii)  $|\mathfrak{B}| = 3^4$  and  $\mathfrak{G}$  is isomorphic to a central product of the dihedral and quaternion groups of order 8.
- (iii)  $|\mathfrak{B}| = 2^6$  and  $\mathfrak{G}$  is isomorphic to the dihedral group of order 18 with cyclic Sylow 3-subgroup.

Here  $\mathcal{S}_0(q^n)$  is the stabilizer in  $\mathcal{S}_0(q^n)$  of the zero vector and hence we know all these groups explicitly. Thus we need only consider the primitive case here.

Let  $\mathfrak{G}$  be a primitive linear group and let  $\mathfrak{B}$  be a normal  $p$ -subgroup of  $\mathfrak{G}$ . Since every normal abelian subgroup of  $\mathfrak{G}$  is cyclic (see for example [9], Lemma 1) it follows that every characteristic abelian subgroup of  $\mathfrak{B}$  is cyclic. Hence by definition  $\mathfrak{B}$  is a group of symplectic type. A characterization of these groups can be found in [1]. In particular for  $p > 2$ ,  $\mathfrak{B}$  is a central product of one cyclic  $p$ -group and any number of nonabelian groups of order  $p^3$  and period  $p$ . If  $p = 2$ , then  $\mathfrak{B}$  is a central product of either a cyclic 2-group or a 2-group of maximal class (that is, a dihedral, semidihedral or quaternion group) and any number of nonabelian groups of order 8. A special case of these are groups of type  $E(p, m)$ .

We say  $\mathfrak{G}$  is a group of type  $E(p, m)$  with  $m \neq 0$  if  $\mathfrak{G}$  has the following structure. If  $p > 2$ , then  $\mathfrak{G}$  is a central product of  $m$  nonabelian groups of order  $p^3$  and period  $p$ . If  $p = 2$ , then  $\mathfrak{G}$  is a central product of a cyclic group of order 2 or 4, and  $m$  nonabelian groups of order 8. Thus in both cases  $|\mathfrak{G}'| = p$ ,  $\mathbf{Z}(\mathfrak{G})$ , the center of  $\mathfrak{G}$ , is cyclic and  $[\mathfrak{G} : \mathbf{Z}(\mathfrak{G})] = p^{2m}$ . Moreover  $|\mathbf{Z}(\mathfrak{G})| = p$  for  $p > 2$  and  $|\mathbf{Z}(\mathfrak{G})| = 2$  or 4 for  $p = 2$ . We call  $m$  the width of  $\mathfrak{G}$ .

Again let  $\mathfrak{B}$  be of symplectic type. If  $p > 2$ , then  $\Omega_1(\mathfrak{B})$ , the subgroup generated by all elements of order  $p$ , is either cyclic (if  $\mathfrak{B}$  is) or of type  $E(p, m)$ . If  $p = 2$ , then the Frattini subgroup  $\Phi(\mathfrak{B})$  is cyclic, and  $\Omega_2(C_{\mathfrak{B}}\Phi(\mathfrak{B}))$  is either cyclic or of type  $E(2, m)$ . The latter group is cyclic only if  $\mathfrak{B}$  is cyclic or  $|\mathfrak{B}| \geq 16$  and  $\mathfrak{B}$  is maximal class. Thus modulo the above mentioned exceptions  $\mathfrak{B}$  contains a characteristic subgroup  $\mathfrak{G}$  of type  $E(p, m)$  with  $m \neq 0$ .

If  $p > 2$ , then for each  $(p, m)$  there is precisely one group of type  $E(p, m)$ . On the other hand, if  $p = 2$ , then for each  $m$  there are three isomorphism classes for  $E(2, m)$  and we describe these now. For convenience we will use the following notation throughout this paper:  $\mathfrak{D}$  denotes the dihedral group of order 8,  $\mathfrak{Q}$  denotes the quaternion group of order 8, and  $\mathfrak{B}$  denotes a cyclic group of order 4. Furthermore any product of these written as  $\mathfrak{D}\mathfrak{D}$ ,  $\mathfrak{B}\mathfrak{D}\mathfrak{D}$ , etc. will indicate a central product. Now we have easily  $\mathfrak{D}\mathfrak{D} \cong \mathfrak{Q}\mathfrak{Q}$  and  $\mathfrak{B}\mathfrak{D} \cong \mathfrak{B}\mathfrak{Q}$ . Hence if  $\mathfrak{G}$  is type  $E(2, m)$  then  $\mathfrak{G}$  is isomorphic to one of the following three groups.

$$\begin{aligned}
 \text{iso I: } \mathfrak{G} &\cong \underbrace{\mathfrak{D}\mathfrak{D} \cdots \mathfrak{D}}_{m-1} \\
 \text{iso II: } \mathfrak{G} &\cong \mathfrak{D} \underbrace{\mathfrak{D} \cdots \mathfrak{D}}_{m-1} \\
 \text{iso III: } \mathfrak{G} &\cong \mathfrak{Z} \underbrace{\mathfrak{D}\mathfrak{D} \cdots \mathfrak{D}}_{m-1} .
 \end{aligned}$$

We will see below that these three groups are nonisomorphic.

For any group  $\mathfrak{G}$  we let  $I(\mathfrak{G})$  denote the number of its noncentral involutions.

LEMMA 1.2. *Let  $\mathfrak{G}$  be a group of type  $E(2, m)$ . Then*

$$\begin{aligned}
 I(\mathfrak{G}) &= 2^{2m} + (-2)^m - 2 \quad \text{if } \mathfrak{G} = \text{iso I} \\
 &= 2^{2m} - (-2)^m - 2 \quad \text{if } \mathfrak{G} = \text{iso II} \\
 &= 2^{2m+1} - 2 \quad \text{if } \mathfrak{G} = \text{iso III} .
 \end{aligned}$$

*In particular these three groups are nonisomorphic. Moreover with the exception of  $\mathfrak{G} = \mathfrak{D}$ ,  $\mathfrak{G}$  is generated by all its noncentral involutions.*

*Proof.* Let  $I^*(\mathfrak{G})$  denote the number of elements  $G \in \mathfrak{G}$  with  $G^2 = 1$ . Then  $I(\mathfrak{G}) = I^*(\mathfrak{G}) - 2$ . Suppose  $\mathfrak{G}$  is iso I or II and write  $\mathfrak{G} = \mathfrak{G}_1\mathfrak{D}$  where  $\mathfrak{G}_1$  is type  $E(2, m - 1)$ . Clearly

$$I^*(\mathfrak{G}) = 3(|\mathfrak{G}_1| - I^*(\mathfrak{G}_1)) + I^*(\mathfrak{G}_1) .$$

Thus if  $I^*(\mathfrak{G}_1) = 2^{2(m-1)} + \delta(-2)^{m-1}$  then  $I^*(\mathfrak{G}) = 2^{2m} + \delta(-2)^m$ . Hence the first two results follow easily. If  $\mathfrak{G} = \text{iso III}$ , let  $\mathbf{Z}(\mathfrak{G}) = \langle \mathbf{Z} \rangle$ . Then the map  $X \rightarrow X\mathbf{Z}$  yields a one to one correspondence between the elements of  $\mathfrak{G}$  with square 1 and those of order 4. Hence clearly  $I^*(\mathfrak{G}) = 1/2 |\mathfrak{G}| = 2^{2m+1}$ .

Now any such  $\mathfrak{G}$  can be written as  $\mathfrak{G}_1\mathfrak{D}\mathfrak{D} \cdots \mathfrak{D}$  and of course  $\mathfrak{D}$  is generated by its noncentral involutions. Since the same is easily seen to be true for  $\mathfrak{G}_1 = \mathfrak{D}, \mathfrak{D}\mathfrak{D}$  or  $\mathfrak{Z}\mathfrak{D}$ , the result follows.

Let  $\mathfrak{G}$  be type  $E(p, m)$  and let  $\mathfrak{B} = \mathfrak{G}/\mathbf{Z}(\mathfrak{G})$ . Then  $\mathfrak{B}$  is elementary abelian of order  $p^{2m}$  and we view this additively as a  $2m$ -dimensional vector space over  $GF(p)$ . If  $p = 2$  we say  $W \in \mathfrak{B}$  is an involution vector if the coset corresponding to  $W$  in  $\mathfrak{G}$  contains an involution of  $\mathfrak{G}$ . Here we let  $i(\mathfrak{B})$  denote the number of such involution vectors.

LEMMA 1.3. *Let  $\mathfrak{S}$  be a group of automorphisms of group  $\mathfrak{G}$  of type  $E(p, m)$  which centralizes  $\mathbf{Z}(\mathfrak{G})$  and let  $\mathfrak{R}$  be the subgroup of  $\mathfrak{S}$  consisting of those elements which centralize  $\mathfrak{B}$ . Then*

- (i)  $\mathfrak{R}$  is isomorphic to a subgroup of the direct product of

$Z(\mathfrak{G})$  taken  $2m$  times.

(ii) The commutator map  $(,)$  of  $\mathfrak{G}$  induces a nonsingular skew-symmetric bilinear form on  $\mathfrak{B}$ . As such  $\mathfrak{S}/\mathfrak{R}$  is contained isomorphically in the symplectic group  $Sp(2m, p)$ .

(iii) If  $p = 2$ , then in addition  $\mathfrak{S}/\mathfrak{R}$  permutes the  $i(\mathfrak{B})$  involution vectors of  $\mathfrak{B}$ . Here

$$\begin{aligned} i(\mathfrak{B}) &= 2^{2m-1} - (-2)^{m-1} - 1 && \text{if } \mathfrak{G} = \text{iso I} \\ &= 2^{2m-1} + (-2)^{m-1} - 1 && \text{if } \mathfrak{G} = \text{iso II} \\ &= 2^{2m} - 1 && \text{if } \mathfrak{G} = \text{iso III.} \end{aligned}$$

*Proof.* (i) Let  $E_1, \dots, E_{2m}$  be a set of coset representatives of  $Z(\mathfrak{G})$  in  $\mathfrak{G}$ . We define  $\theta : \mathfrak{R} \rightarrow \prod Z(\mathfrak{G})$  ( $2m$  times) by  $\theta(K) = \prod E_i^K E_i^{-1}$ . This is easily seen to be a monomorphism.

(ii) and (iii) If  $W$  is an involution vector then we see easily that the coset of  $W$  contains precisely two noncentral involutions of  $\mathfrak{G}$ . Hence  $i(\mathfrak{B}) = 1/2I(\mathfrak{G})$ . The result now follows easily.

We now consider the action of  $\mathfrak{G}$  on a vector space  $\mathfrak{B}$ .

**LEMMA 1.4.** *Let group  $\mathfrak{G}$  of type  $E(p, m)$  act on vector space  $\mathfrak{B}$  of order  $q^n$ . Suppose further that  $\mathfrak{G}'$  acts without fixed points on  $\mathfrak{B}^\#$ . Then*

(i)  $sp^m \mid n$  where  $s$  is the smallest positive integer with  $|Z(\mathfrak{G})| \mid q^s - 1$ .

(ii) If  $T \in \mathfrak{G} - Z(\mathfrak{G})$  has order  $p$  then  $|C_{\mathfrak{B}}(T)| = q^{n/p}$ .

(iii) If  $x \in \mathfrak{B}^\#$ , then  $\mathfrak{G}_x$ , the stabilizer of  $x$  in  $\mathfrak{G}$  is elementary abelian.

*Proof.* (i) Since  $\mathfrak{G}'$  acts without fixed points  $q \neq p$ . By complete reducibility we can assume that  $\mathfrak{G}$  acts irreducibly on  $\mathfrak{B}$ . Let  $\chi$  be the character of an absolutely irreducible constituent of  $\mathfrak{G}$ . From the representation of  $\mathfrak{G}$  as a homomorphic image of a direct product of nonabelian group of order  $p^3$  (and possibly a cyclic group of order 4 if  $p = 2$ ) we see easily that  $\deg \chi = p^m$  and  $\chi$  vanishes off  $Z(\mathfrak{G})$ . Hence by definition of  $s$ ,  $GF(q)(\chi) = GF(q^s)$  and  $\mathfrak{B}$  contains as absolutely irreducible constituents the  $s$  algebraic conjugates of the representation affording  $\chi$ . Thus (i) follows.

(ii) We wish to show here that  $\dim C_{\mathfrak{B}}(T) = n/p$ . This dimension is clearly invariant under field extension so by complete reducibility we can assume  $\mathfrak{B}$  is absolutely irreducible. If  $\theta$  is the corresponding complex character then  $\theta(T)$  is a sum of  $p$ th roots of unity (including 1) and  $\theta(T) \neq 0$ . Hence all eigenvalues occur with the same multiplicity  $n/p$  and (ii) follows.

(iii) This is clear since  $\Phi(\mathfrak{G})$  acts semiregularly on  $\mathfrak{B}^\#$ .

LEMMA 1.5. *Let group  $\mathfrak{G}$  of type  $E(p, m)$  act on vector space  $\mathfrak{B}$  of order  $q^n$  and let  $T \in \mathfrak{G} - Z(\mathfrak{G})$  have order  $p$ . Suppose further that  $\mathfrak{G}$  acts without fixed points on  $\mathfrak{B}$ . Then*

(i) *There exists  $x \in \mathfrak{B}^\#$  with  $\mathfrak{G}_x = \langle 1 \rangle$  with the following exceptions which occur for  $p = 2$ : (a)  $q^n = 3^2$ ,  $\mathfrak{G} = \mathfrak{D}$ , (b)  $q^n = 5^2$ ,  $\mathfrak{G} = \mathfrak{B}\mathfrak{D}$ , (c)  $q^n = 3^4$ ,  $\mathfrak{G} = \mathfrak{D}\mathfrak{D}$ . In each of these exceptions  $|\mathfrak{G}_x| = 2$  for all  $x \in \mathfrak{B}^\#$ .*

(ii) *There exists  $x \in \mathfrak{B}^\#$  with  $\mathfrak{G}_x = \langle T \rangle$  with the following exceptions which occur for  $p = 2$ : (a)  $q^n = 3^4$ ,  $\mathfrak{G} = \mathfrak{D}\mathfrak{D}$ , (b)  $q^n = 5^4$ ,  $\mathfrak{G} = \mathfrak{B}\mathfrak{D}\mathfrak{D}$ , (c)  $q^n = 3^8$ ,  $\mathfrak{G} = \mathfrak{D}\mathfrak{D}\mathfrak{D}$ . In each of these exceptions  $|\mathfrak{G}_x| = 4$  or  $1$  for all  $x \in \mathfrak{B}^\#$ .*

*Proof.* (i) We first note that by [4] Theorem II (a), (b) and (c) are in fact exceptions. Suppose now that  $\mathfrak{G}_x \neq \langle 1 \rangle$  for all  $x \in \mathfrak{B}^\#$ . Then every element of  $\mathfrak{B}^\#$  is centralized by a noncentral element  $P \in \mathfrak{G}$  of order  $p$ . Thus

$$\mathfrak{B} = \bigcup_p C_{\mathfrak{B}}(P)$$

where the union is over representatives of the  $N$  noncentral subgroups of  $\mathfrak{G}$  of order  $p$ . By Lemma 1.4 we have

$$q^n = |\mathfrak{B}| \leq Nq^{n/p}$$

and  $q^{n(1-1/p)} \leq N$ .

Let  $p > 2$ . Then  $N < p^{2m+1}/(p-1)$  and  $n \geq sp^m$ . Furthermore  $p \mid q^s - 1$  so  $q^s \geq p + 1$ . Thus

$$\begin{aligned} p^{p^m - p^{m-1}} &< (p + 1)^{p^m - p^{m-1}} \leq q^{s(p^m - p^{m-1})} \\ &\leq q^{n(1-1/p)} \leq N < p^{2m+1}/(p-1) < p^{2m+1}. \end{aligned}$$

This yields  $p^{m-1}(p-1) < 2m+1$  and since  $p > 2$  we have  $p = 3$ ,  $m = 1$  here. However with  $p = 3$ ,  $m = 1$  the equation

$$(p + 1)^{p^m - p^{m-1}} < p^{2m+1}/(p-1)$$

is not satisfied so  $p > 2$  cannot occur here.

Now let  $p = 2$  so that  $N = I(\mathfrak{G})$ . Suppose first that  $|Z(\mathfrak{G})| = 4$ . Then  $4 \mid q^s - 1$  and  $I(\mathfrak{G}) < 2^{2m+1}$ . Thus

$$5^{2m-1} \leq q^{s2^{m-1}} \leq q^{n(1-1/p)} \leq N \leq 2^{2m+1}.$$

This yields  $5^{2m-1} < 2^{2m+1}$  so  $m = 1$  or  $2$ . If  $m = 1$ , then  $q^{n/2} < 8$  and  $4 \mid q^s - 1$  yields  $q^n = 5^2$  and we have exception (b). If  $m = 2$ , then  $q^{n/2} < 32$ ,  $4 \mid q^s - 1$  and  $4s \mid n$  yields  $q^n = 5^4$ . We show now that this possibility does not occur. Let  $x \in \mathfrak{B}^\#$  and suppose that  $\mathfrak{G}_x \neq \langle 1 \rangle$ .

Choose  $P \in \mathbb{G}_x^\#$ . Since  $\mathbb{G}_x$  is abelian  $\mathbb{G}_x \cong C_{\mathbb{G}}(P) = \langle P \rangle \times \bar{\mathbb{G}}$  where  $\bar{\mathbb{G}} \cong 3\Omega$ . Now  $x \in C_{\mathbb{G}}(P)$ ,  $|C_{\mathbb{G}}(P)| = 5^2$  and  $\bar{\mathbb{G}}$  acts on this subspace. Since this action yields the exceptional case (b) we have  $|\bar{\mathbb{G}}_x| = 2$  and hence  $|\mathbb{G}_x| = 4$ . Thus for all  $x \in \mathbb{B}^\#$ ,  $|\mathbb{G}_x| = 1$  or  $4$ . This, by the way, is the exceptional case (b) of part (ii). If  $\mathbb{B} = \bigcup C_{\mathbb{G}}(P)$  then since each  $\mathbb{G}_x$  is elementary abelian, we see that this union covers  $\mathbb{B}$  three times. Thus

$$5^4 - 1 = |\mathbb{B}^\#| \leq \frac{1}{3}I(\mathbb{G}) \cdot (5^2 - 1) < \frac{1}{3} \cdot 2^5(5^2 - 1)$$

a contradiction.

Now let  $|Z(\mathbb{G})| = 2$  so  $I(\mathbb{G}) \leq 2^{2m} + 2^m - 2$ . Since  $q^s \geq 3$  we have

$$3^{2m-1} \leq q^{s2^{m-1}} \leq q^{n(1-1/p)} \leq N \leq 2^{2m} + 2^m - 2.$$

This yields  $3^{2m-1} < 2^{2m} + 2^m$  so  $m = 1$  or  $2$ . If  $m = 1$  then  $q^{n/2} \leq 4$  so  $q^n = 3^2$ . Clearly  $\mathbb{G} \neq \Omega$  so we have exception (a) here. If  $m = 2$ , then  $4 | n$  and  $q^{n/2} \leq 18$  yields  $q^n = 3^4$ . If  $\mathbb{G} \cong \mathbb{D}\Omega$  we have exception (c). We show finally that  $\mathbb{G} \neq \Omega\Omega$ . Let  $x \in \mathbb{B}^\#$  and suppose  $\mathbb{G}_x \neq \langle 1 \rangle$ . Choose  $P \in \mathbb{G}_x^\#$  and let  $C_{\mathbb{G}}(P) = \langle P \rangle \times \bar{\mathbb{G}}$ . Here  $\bar{\mathbb{G}}$  is nonabelian of order 8. Since  $C_{\mathbb{G}}(\bar{\mathbb{G}})$  contains  $P$  we see that  $C_{\mathbb{G}}(\bar{\mathbb{G}}) \cong \mathbb{D}$  and hence  $\mathbb{G} \cong \mathbb{D}\bar{\mathbb{G}}$ . Thus  $\bar{\mathbb{G}} \cong \mathbb{D}$ . This implies as above that  $|\bar{\mathbb{G}}_x| = 2$  and  $|\mathbb{G}_x| = 4$ , thereby yielding exception (a) of part (ii). Again if  $\mathbb{B} = \bigcup C_{\mathbb{G}}(P)$ , then  $\mathbb{B}$  is triply covered so

$$3^4 - 1 = |\mathbb{B}^\#| \leq \frac{1}{3}I(\mathbb{G})(3^2 - 1) < \frac{1}{3}20(3^2 - 1),$$

a contradiction. This completes the proof of (i).

(ii) If  $m = 1$ , then any abelian subgroup of  $\mathbb{G}$  of order 4 meets  $Z(\mathbb{G})$ . Since  $Z(\mathbb{G})$  acts semiregularly, we conclude that for all  $x \in \mathbb{B}^\#$ ,  $|\mathbb{G}_x| = 1$  or  $2$ . Thus the result follows here.

Let  $m \geq 2$ . Then  $C_{\mathbb{G}}(T) = \langle T \rangle \times \bar{\mathbb{G}}$  where  $\bar{\mathbb{G}}$  is type  $E(p, m - 1)$ . Note that  $\mathbb{G} = \bar{\mathbb{G}}C_{\mathbb{G}}(\bar{\mathbb{G}})$  and  $T \in C_{\mathbb{G}}(\bar{\mathbb{G}})$ . Thus if  $p = 2$  then  $C_{\mathbb{G}}(\bar{\mathbb{G}}) \cong \mathbb{D}$  and the isomorphism class of  $\bar{\mathbb{G}}$  is uniquely determined by  $\mathbb{G} \cong \bar{\mathbb{G}}\mathbb{D}$ . Now  $\bar{\mathbb{G}}$  acts on  $C_{\mathbb{G}}(T)$  a subspace of size  $q^{n/2}$  and hence if this is not one of the exceptions of part (i), then there exists  $x \in C_{\mathbb{G}}(T)^\#$  with  $\bar{\mathbb{G}}_x = \langle 1 \rangle$ . Since  $T \in \mathbb{G}_x$  and  $\mathbb{G}_x$  is abelian, it then follows that  $\mathbb{G}_x = \langle T \rangle$ . The result now clearly follows.

We now turn to a variant of an argument used in [2] (§ 2.5).

LEMMA 1.6. *Let  $\mathbb{G} = \mathbb{G}\mathfrak{J}$  where  $\mathbb{G}$  is type  $E(p, m)$ ,  $\mathbb{G} \triangle \mathbb{G}$  and  $\mathfrak{J} = \langle J \rangle$  is cyclic of order  $j$ . Suppose  $\mathbb{G}$  acts on  $F$ -vector space  $\mathbb{B}$  in such a way that the restriction to  $\mathbb{G}$  is faithful and absolutely*

irreducible. If further the characteristic of  $F$  is prime to  $|\mathfrak{G}|$ , then there exists nonnegative integers  $a_0, a_1, \dots, a_{j-1}$  satisfying

- (i)  $a_0 + a_1 + \dots + a_{j-1} = p^m$
- (ii)  $a_0^2 + a_1^2 + \dots + a_{j-1}^2 \leq N$  and
- (iii)  $a_0 = \dim_F C_{\mathfrak{B}}(J)$

where  $N$  is the number of orbits in  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$  under the action of  $\mathfrak{S}$ .

*Proof.* Since  $\dim_F C_{\mathfrak{B}}(J)$  is clearly invariant under field extension, we can assume  $F$  is algebraically closed. Let  $\varepsilon \in F$  be a primitive  $j$ th root of unity and suppose that  $\varepsilon^i$  occurs as an eigenvalue of  $J$  with multiplicity  $a_i$  for  $i = 0, 1, \dots, j - 1$ . If  $\Sigma$  denotes the enveloping algebra of this representation then clearly

$$\begin{aligned} a_0 + a_1 + \dots + a_{j-1} &= \dim_F \mathfrak{B} \\ a_0^2 + a_1^2 + \dots + a_{j-1}^2 &= \dim_F C_{\Sigma}(J) \\ a_0 &= \dim_F C_{\mathfrak{B}}(J) . \end{aligned}$$

Now  $\mathfrak{B}$  is a faithful absolutely irreducible  $\mathfrak{G}$ -module so  $\dim_F \mathfrak{B} = p^m$ . Hence (i) and (iii) follow. In addition the group ring  $F(\mathfrak{G})$  maps onto  $\Sigma$  in the obvious manner. Under this map  $Z(\mathfrak{G})$  is sent into the field of scalars so the image of  $F[\mathfrak{G}]$  is spanned by  $p^{2m}$  coset representatives of  $Z(\mathfrak{G})$  in  $\mathfrak{G}$ . But  $\dim_F \Sigma = p^{2m}$  so these must in fact form a basis of  $\Sigma$ . With this choice of basis we see clearly that  $\dim_F C_{\Sigma}(J)$  is at most equal to the number of orbits of  $\mathfrak{S}$  on  $\mathfrak{G}/Z(\mathfrak{G})$  so the result follows.

The following two results enable us to use inductive methods in our study of half-transitive linear groups.

LEMMA 1.7. *Let  $\mathfrak{G}$  be a half-transitive permutation group and let  $\mathfrak{N} \triangleleft \mathfrak{G}$ . Suppose that either  $\mathfrak{N} = \langle 1 \rangle$  or  $\mathfrak{N}$  acts half-transitively. Let  $\mathfrak{G} \cong \mathfrak{S} \cong \mathfrak{N}$  where  $\mathfrak{S}/\mathfrak{N}$  is a normal Hall subgroup of  $\mathfrak{G}/\mathfrak{N}$ . Then  $\mathfrak{S}$  acts half-transitively.*

*Proof.* See Lemma 2.1 of [5].

LEMMA 1.8. (*Reduction Lemma*). *Let  $\mathfrak{G}$  be a linear group on  $GF(q)$ -vector space  $\mathfrak{B}$  and suppose that  $\mathfrak{G}$  acts half-transitively but not semiregularly on  $\mathfrak{B}^*$ . Let  $\mathfrak{C}$  be a group of type  $E(p, m)$  with  $\mathfrak{C} \triangleleft \mathfrak{G}$ . Then there exists a linear group  $\overline{\mathfrak{G}}$  acting on  $GF(q)$ -vector space  $\mathfrak{U}$  and a normal subgroup  $\overline{\mathfrak{C}}$  of  $\overline{\mathfrak{G}}$  satisfying*

- (i)  $\overline{\mathfrak{G}}$  acts half-transitively on  $\mathfrak{U}$ .
- (ii)  $\overline{\mathfrak{C}} \cong \mathfrak{C}$  and  $\overline{\mathfrak{C}}$  acts irreducibly on  $\mathfrak{U}$ .
- (iii) If  $\mathfrak{G}$  is solvable so is  $\overline{\mathfrak{G}}$ .



- (iv) If  $\mathcal{G} \neq \mathcal{D}$ , then  $\bar{\mathcal{G}}$  does not act semiregularly on  $\mathcal{U}$ .
- (v) Suppose that either  $p > 2$  or  $p = 2$  and  $m \geq 2$ . Then either  $\bar{\mathcal{G}} = \bar{\mathcal{G}} \cong \mathcal{D}\mathcal{D}$  with  $q = 3$  or  $\bar{\mathcal{G}}$  is primitive as a linear group.

*Proof.* Since  $\mathcal{G}$  does not act semiregularly, it acts irreducibly on  $\mathcal{B}$ . By Clifford's theorem all irreducible  $\mathcal{G}$  constituents of  $\mathcal{B}$  are conjugate and hence  $\mathcal{G}$  acts faithfully on each. Let  $\mathcal{U}$  be an irreducible  $\mathcal{G}$ -submodule of  $\mathcal{B}$  and let  $\mathcal{N} = \{G \in \mathcal{G} \mid \mathcal{U}G = \mathcal{U}\}$ . Suppose  $x \in \mathcal{U}^\#$ . Since  $\mathcal{G} \triangleleft \bar{\mathcal{G}}$

$$(x\mathcal{G})\mathcal{G}_x = (x\mathcal{G}_x)\mathcal{G} = x\mathcal{G}$$

and hence  $\mathcal{G}_x$  normalizes  $x\mathcal{G}$ . Moreover  $\mathcal{G}$  acts irreducibly on  $\mathcal{U}$  so  $\mathcal{U}$  is the linear span of  $x\mathcal{G}$  and hence  $\mathcal{G}_x \subseteq \mathcal{N}$ . If  $\mathcal{R}$  is the kernel of the action of  $\mathcal{N}$  on  $\mathcal{U}$ , then clearly  $\bar{\mathcal{G}} = \mathcal{N}/\mathcal{R}$  acts semiregularly on  $\mathcal{U}^\#$ . Since  $\mathcal{G}$  acts faithfully on  $\mathcal{U}$ ,  $\bar{\mathcal{G}} = \mathcal{G}\mathcal{R}/\mathcal{R} \cong \mathcal{G}$ . Also  $\bar{\mathcal{G}} \triangleleft \bar{\mathcal{G}}$  and  $\bar{\mathcal{G}}$  acts irreducibly on  $\mathcal{U}$  so (i), (ii) and (iii) follow.

We have  $\bar{\mathcal{G}} \cong \mathcal{G}$ . Thus if  $\mathcal{G} \neq \mathcal{D}$  then  $\bar{\mathcal{G}}$ , and hence  $\bar{\mathcal{G}}$ , cannot act semiregularly. This yields (iv). Finally suppose that either  $p > 2$  or  $p = 2$  and  $m \geq 2$ . Then  $\mathcal{G} \neq \mathcal{D}$  so  $\bar{\mathcal{G}}$  does not act semiregularly. Hence if  $\bar{\mathcal{G}}$  is imprimitive as a linear group, then the structure of  $\bar{\mathcal{G}}$  is given in Theorem 1.1. In both (i) and (iii) of that theorem  $\bar{\mathcal{G}}$  has a normal abelian subgroup of index 2 and hence  $\bar{\mathcal{G}}$  could not possibly contain  $\bar{\mathcal{G}}$ . Thus only (ii) of that theorem can occur here and since  $m \geq 2$  this yields  $\bar{\mathcal{G}} = \bar{\mathcal{G}} \cong \mathcal{D}\mathcal{D}$  and  $|\mathcal{U}| = 3^4$ . This completes the proof of the lemma.

We close this section by offering a precise statement of Lemma 6 of [4]. The proof is the same and will not be repeated.

**LEMMA 1.9.** *Let  $\mathcal{G}$  act faithfully on vector space  $\mathcal{B}$  and half-transitively on  $\mathcal{B}^\#$ . Suppose that for all  $x \in \mathcal{B}^\#$ ,  $|\mathcal{G}_x| = 2$ . If  $\mathcal{G}$  has a central involution, then  $|\mathcal{B}| = q^{2r}$  with  $q \neq 2$  and  $q^r + 1 = I(\mathcal{G})$ .*

**2. Theorem A.** The following assumptions hold throughout this section.

**ASSUMPTIONS.** Group  $\mathcal{G}$  acts faithfully on vector space  $\mathcal{B}$  of order  $q^n$  and half-transitively but not semiregularly on  $\mathcal{B}^\#$ .  $\mathcal{G}$  is a group of type  $E(p, m)$  with  $\mathcal{G} \triangleleft \bar{\mathcal{G}}$ . In addition  $\mathcal{G}$  acts irreducibly on  $\mathcal{B}$  and  $\bar{\mathcal{G}}$  is primitive as a linear group.

It is convenient to keep track of four separate possibilities.

**DEFINITION.** We define the type of  $\mathcal{G}$  as follows.

- type I:  $p > 2$   
 type II:  $p = 2, |Z(\mathfrak{G})| = 2$   
 type III:  $p = 2, |Z(\mathfrak{G})| = 4, Z(\mathfrak{G}) \subseteq Z(\mathfrak{G})$   
 type IV:  $p = 2, |Z(\mathfrak{G})| = 4, Z(\mathfrak{G}) \not\subseteq Z(\mathfrak{G})$ .

LEMMA 2.1. *Let  $s \geq 1$  be minimal with  $|Z(\mathfrak{G})| \mid q^s - 1$ . Let  $\mathfrak{M}$  be any subgroup of  $\mathfrak{G}$  with  $\mathfrak{G} \subseteq \mathfrak{M} \subseteq C_{\mathfrak{G}}(Z(\mathfrak{G}))$ . Then  $\mathfrak{M} \subseteq GL(p^m, q^s)$  and this representation of  $\mathfrak{M}$  is absolutely irreducible. Furthermore  $n = sp^m$  and we have the following*

- type I:  $s \mid (p - 1)$   
 type II:  $s = 1$   
 type III:  $s = 1$  or  $2$

type IV:  $s = 2$ , and if  $\bar{\mathfrak{M}}$  is a  $q'$ -subgroup of  $\mathfrak{G}$  with  $\mathfrak{G} \subseteq \bar{\mathfrak{M}}$  and  $\bar{\mathfrak{M}} \not\subseteq C_{\mathfrak{G}}(Z(\mathfrak{G}))$ , then  $\bar{\mathfrak{M}} \subseteq GL(p^{m+1}, q)$  and this is an absolutely irreducible representation.

*Proof.* If  $s$  is defined as above then  $GF(q^s)$  is clearly the minimal splitting field of the representation of  $\mathfrak{G}$ . Hence  $n = sp^m$  since we are dealing with finite fields here and since the absolutely irreducible constituents of  $\mathfrak{G}$  have degree  $p^m$ .

Now  $\mathfrak{G}$  is primitive as a linear group so by Lemma 1.1 of [5],  $C_{\mathfrak{G}}(Z(\mathfrak{G})) \subseteq GL(p^m, q^s)$ . Let  $\mathfrak{M}$  be a subgroup of  $\mathfrak{G}$  with

$$\mathfrak{G} \subseteq \mathfrak{M} \subseteq C_{\mathfrak{G}}(z(\mathfrak{G}))$$

so that  $\mathfrak{M} \subseteq GL(p^m, q^s)$ . Since  $\mathfrak{M} \supseteq \mathfrak{G}$  and the degree of this representation is  $p^m$ , the representation is clearly absolutely irreducible.

The results on the value of  $s$  for types I, II and III are clear. Let  $\mathfrak{G}$  be type IV. Then certainly  $s = 1$  or  $2$ . If  $s = 1$ , then since  $\mathfrak{G}$  is primitive,  $Z(\mathfrak{G})$  consists of scalar matrices and is therefore central in  $\mathfrak{G}$ , a contradiction. Thus  $s = 2$ . Let  $\bar{\mathfrak{M}}$  be given with  $\mathfrak{G} \subseteq \bar{\mathfrak{M}}$ ,  $\bar{\mathfrak{M}} \not\subseteq C_{\mathfrak{G}}(Z(\mathfrak{G}))$ . Since  $s = 2$ ,  $\bar{\mathfrak{M}} \subseteq GL(p^{m+1}, q)$ . Clearly  $\bar{\mathfrak{M}}$  is either absolutely irreducible or it has two absolutely irreducible constituents of degree  $p^m$ . In the latter case,  $Z(\mathfrak{G})$  would be central in each such constituent and hence in  $\bar{\mathfrak{M}}$ , a contradiction.

LEMMA 2.2. *Let  $\mathfrak{M}$  be a  $p$ -group acting faithfully and absolutely irreducibly on  $F$ -vector space  $\mathfrak{B}$ . Let  $\dim_F \mathfrak{B} = k$ . Then there exists subgroups  $\mathfrak{N}$  and  $\mathfrak{R}$  of  $\mathfrak{M}$  and an  $\mathfrak{N}$ -subspace  $\mathfrak{U}$  of  $\mathfrak{B}$  with the representation of  $\mathfrak{M}$  on  $\mathfrak{B}$  induced from that of  $\mathfrak{N}$  on  $\mathfrak{U}$ . Furthermore  $\mathfrak{R} = C_{\mathfrak{M}}(\mathfrak{U})$  and either*

- (i)  $[\mathfrak{M} : \mathfrak{N}] = k$ ,  $\dim \mathfrak{U} = 1$  and  $\mathfrak{N}/\mathfrak{R}$  is cyclic, or  
 (ii)  $[\mathfrak{M} : \mathfrak{N}] = k/2$ ,  $\dim \mathfrak{U} = 2$ ,  $\mathfrak{N}/\mathfrak{R}$  is dihedral, semidihedral or quaternion and  $p = 2$ .

*Proof.* The result is trivial if  $\text{char } F = p$  so assume this is not the case. Applying Roquette's theorem ([9]) repeatedly we can find  $\mathfrak{N}, \mathfrak{R}$  and  $\mathfrak{U}$  as above with  $\mathfrak{N}/\mathfrak{R}$  cyclic, dihedral, semidihedral or quaternion. Since  $\mathfrak{M}$  is absolutely irreducible so is the action of  $\mathfrak{N}/\mathfrak{R}$  on  $\mathfrak{U}$ . Thus  $\dim \mathfrak{U} = 1$  if  $\mathfrak{N}/\mathfrak{R}$  is cyclic and  $\dim \mathfrak{U} = 2$  otherwise.

LEMMA 2.3. *Let  $w$  denote the period of a Sylow  $p$ -subgroup of  $C_{\mathfrak{G}}(\mathbf{Z}(\mathfrak{G}))$ . Then for all  $x \in \mathfrak{B}^{\#}$  we have*

- type I:  $[\mathfrak{G} : \mathfrak{G}_x]_p \leq p^m \min \{w, |q^s - 1|_p\}$
- type II:  $[\mathfrak{G} : \mathfrak{G}_x]_p \leq p^{m+1} \min \{w, |q^2 - 1|_p\}$
- type III:  $[\mathfrak{G} : \mathfrak{G}_x]_p \leq p^m \min \{w, |q^2 - 1|_p\}$
- type IV:  $[\mathfrak{G} : \mathfrak{G}_x]_p \leq p^{m+1} \min \{w, |q^2 - 1|_p\}$ .

*Proof.* We consider types I, II and III first. Let  $\mathfrak{B}$  be a Sylow  $p$ -subgroup of  $\mathfrak{G}$ . Then  $\mathfrak{B} \supseteq \mathfrak{G}$  and  $\mathbf{Z}(\mathfrak{G})$  is central in  $\mathfrak{B}$ . By Lemma 2.1 we can view  $\mathfrak{B}$  as a subgroup of  $GL(p^m, q^s)$  and this representation is absolutely irreducible. Let  $\mathfrak{N}, \mathfrak{R}$  and  $\mathfrak{U}$  be as in the preceding lemma with  $\mathfrak{M} = \mathfrak{B}$ . Note that for  $y \in \mathfrak{U}^{\#}$ ,  $\mathfrak{B}_y \supseteq \mathfrak{R}$ . If  $\mathfrak{N}/\mathfrak{R}$  is cyclic, then  $[\mathfrak{B} : \mathfrak{N}] = p^m$ ,  $[\mathfrak{N} : \mathfrak{R}] \leq \min \{w, |q^s - 1|_p\}$  so

$$[\mathfrak{B} : \mathfrak{B}_y] \leq p^m \min \{w, |q^s - 1|_p\}.$$

Suppose that  $\mathfrak{N}/\mathfrak{R}$  is not cyclic. Then  $p = 2$ . Now it is clear that  $\mathbf{Z}(\mathfrak{G}) \subseteq \mathbf{Z}(\mathfrak{B}) \subseteq \mathfrak{N}$  and  $\mathbf{Z}(\mathfrak{G}) \cap \mathfrak{R} = \langle 1 \rangle$ . Thus since 2-groups of maximal class have centers of order 2,  $\mathfrak{G}$  must be type II. Here  $[\mathfrak{B} : \mathfrak{N}] = p^{m-1}$  and  $[\mathfrak{N} : \mathfrak{R}] \leq p \min \{w, |q^{2s} - 1|_p\}$  since  $\mathfrak{N}/\mathfrak{R}$  has a cyclic subgroup of index  $p = 2$  which has a faithful irreducible representation in  $GF(q^{2s})$ . Note that  $s = 1$  here. Now by half-transitivity, for all  $x \in \mathfrak{B}^{\#}$

$$[\mathfrak{G} : \mathfrak{G}_x]_p = [\mathfrak{G} : \mathfrak{G}_y]_p \leq [\mathfrak{B} : \mathfrak{B}_y].$$

Thus the first three results follow.

Now let  $\mathfrak{G}$  be type IV and again let  $\mathfrak{B}$  be a Sylow  $p$ -subgroup of  $\mathfrak{G}$ . Let  $\mathfrak{M} = C_{\mathfrak{B}}(\mathbf{Z}(\mathfrak{G}))$  so that  $\mathfrak{B} > \mathfrak{M} \supseteq \mathfrak{G}$  and  $[\mathfrak{B} : \mathfrak{M}] = 2$ . By Lemma 2.1,  $\mathfrak{B}$  is absolutely irreducible as a subgroup of  $GL(p^{m+1}, q)$ . We extend the field now to  $GF(q^s) = GF(q^2)$ . Thus we let  $\mathfrak{B}$  act on  $\mathfrak{B} \otimes GF(q^2)$  and this representation is again absolutely irreducible. If the restriction to  $\mathfrak{M}$  were irreducible, then since  $4 | q^s - 1$ ,  $\mathbf{Z}(\mathfrak{G})$  which is central in  $\mathfrak{M}$  would consist of scalar matrices and hence it would be central in  $\mathfrak{B}$ , a contradiction. Thus the representation of  $\mathfrak{B}$  is induced from one of  $\mathfrak{M}$ . Let  $\mathfrak{N}, \mathfrak{R}$  and  $\mathfrak{U} \subseteq \mathfrak{B} \otimes GF(q^s)$  be as in the preceding lemma with  $\mathfrak{M} \subseteq \mathfrak{M}$ . Since  $\mathbf{Z}(\mathfrak{G}) \subseteq \mathfrak{N}$  and  $|\mathbf{Z}(\mathfrak{G})| = 4$  we see that  $\mathfrak{N}/\mathfrak{R}$  is cyclic. Hence  $[\mathfrak{N} : \mathfrak{R}] \leq \min \{w, |q^s - 1|_p\}$ . Moreover  $[\mathfrak{B} : \mathfrak{N}] = p^{m+1}$  so

$$[\mathfrak{B} : \mathfrak{R}] \leq p^{m+1} \min \{w, |q^s - 1|_p\}.$$

Now all elements of  $\mathfrak{R}$  have a common nonzero fixed point in  $\mathfrak{B} \otimes GF(q^s)$ . This means that a certain set of simultaneous linear equations over  $GF(q)$  has a nonzero solution over  $GF(q^s)$ . Thus there is a nonzero solution over  $GF(q)$  and hence there exists  $y \in \mathfrak{B}^\#$  with  $\mathfrak{B}_y \cong \mathfrak{R}$ . The result now follows as above.

LEMMA 2.4. *Let  $\mathfrak{A} = C_{\mathfrak{G}}(\mathfrak{G})$ . Then  $\mathfrak{A}$  is a normal cyclic subgroup of  $\mathfrak{G}$  which is central in  $C_{\mathfrak{G}}(\mathfrak{Z}(\mathfrak{G}))$  and acts semiregularly on  $\mathfrak{B}^\#$ . Suppose that  $m \geq 3$  if  $p = 2$ . Then there exists  $x \in \mathfrak{B}^\#$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$  and hence  $[\mathfrak{G} : \mathfrak{G}_x]_p \geq |\mathfrak{A}_p| p^{2m}$  where  $\mathfrak{A}_p$  is the normal Sylow  $p$ -subgroup of  $\mathfrak{A}$ . This yields*

- type I:  $w \geq p^m |\mathfrak{A}_p|, \quad |q^s - 1|_p \geq p^{m+1}$
- type II:  $w \geq p^{m-1} |\mathfrak{A}_p|, \quad |q^2 - 1|_p \geq p^m$
- type III:  $w \geq p^m |\mathfrak{A}_p|, \quad |q^2 - 1|_p \geq p^{m+2}$
- type IV:  $w \geq p^{m-1} |\mathfrak{A}_p|, \quad |q^2 - 1|_p \geq p^{m+1}$ .

*Proof.* Since  $\mathfrak{G}$  is irreducible, Schur's lemma guarantees that  $\mathfrak{A}$  is cyclic and acts semiregularly. Clearly  $\mathfrak{A} \subseteq C_{\mathfrak{G}}(\mathfrak{Z}(\mathfrak{G}))$ . By Lemma 2.1,  $\mathfrak{G} \subseteq C_{\mathfrak{G}}(\mathfrak{Z}(\mathfrak{G})) \subseteq GL(p^m, q^s)$  and this is an absolutely irreducible representation of  $\mathfrak{G}$ . Since  $\mathfrak{A}$  centralizes  $\mathfrak{G}$ ,  $\mathfrak{A}$  consists of scalar matrices here and hence  $\mathfrak{A}$  is central in  $C_{\mathfrak{G}}(\mathfrak{Z}(\mathfrak{G}))$ .

If  $p > 2$  set  $\mathfrak{G}^* = \mathfrak{G}$  while if  $p = 2$  we set  $\mathfrak{G}^* = \mathfrak{A}^*\mathfrak{G}$  where  $\mathfrak{A}^* = \{A \in \mathfrak{A} \mid A^4 = 1\}$ . Then  $\mathfrak{G}^*$  is also of type  $E(p, m)$  and every subgroup of  $\mathfrak{A}\mathfrak{G}$  of order  $p$  is in  $\mathfrak{G}^*$ . With the additional assumption that  $m \geq 3$  if  $p = 2$ , Lemma 1.5 applied to  $\mathfrak{G}^*$  guarantees the existence of a point  $x \in \mathfrak{B}^\#$  with  $\mathfrak{G}_x \cap \mathfrak{G}^* = \langle 1 \rangle$ . This clearly yields  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$ .

Now  $\mathfrak{A}_p\mathfrak{G} \trianglelefteq \mathfrak{G}$  and  $|\mathfrak{A}_p\mathfrak{G}| = |\mathfrak{A}_p| p^{2m}$ . If  $x$  is as above then

$$|\mathfrak{G}|_p \geq |\mathfrak{G}_x \mathfrak{A}_p \mathfrak{G}|_p = |\mathfrak{G}_x|_p |\mathfrak{A}_p \mathfrak{G}| = |\mathfrak{G}_x|_p |\mathfrak{A}_p| p^{2m}$$

and hence  $[\mathfrak{G} : \mathfrak{G}_x]_p \geq |\mathfrak{A}_p| p^{2m}$ . By half-transitivity this holds for all  $x \in \mathfrak{B}^\#$ . Combining this with the results of Lemma 2.3 and noting that  $|\mathfrak{A}_p| \geq p$  for type I and II groups and  $|\mathfrak{A}_p| \geq p^2$  for type III and IV groups, we clearly obtain our result.

LEMMA 2.5. *Let  $\mathfrak{H} = C_{\mathfrak{G}}(\mathfrak{Z}(\mathfrak{G}))$ . Then  $\mathfrak{G}$  has the following structure.*

- (i)  $\mathfrak{G}/\mathfrak{H}$  is cyclic
- (ii)  $\mathfrak{H}/\mathfrak{A}\mathfrak{G}$  acts faithfully on  $\mathfrak{B} = \mathfrak{G}/\mathfrak{Z}(\mathfrak{G})$  and as a linear group on  $\mathfrak{B}$  we have  $\mathfrak{H}/\mathfrak{A}\mathfrak{G} \subseteq Sp(2m, p)$
- (iii)  $\mathfrak{A}\mathfrak{G}/\mathfrak{A}$  is elementary abelian of order  $p^{2m}$
- (iv)  $\mathfrak{A}$  is cyclic.

*Proof.* All results but (ii) are clear. Let  $\mathfrak{B} = C_{\mathfrak{H}}(\mathfrak{B})$ . Clearly  $\mathfrak{B} \cong \mathfrak{A}\mathfrak{G}$ . The result will follow from Lemma 1.3 if we show that

$\mathfrak{B} = \mathfrak{A}\mathfrak{C}$ .

Suppose first that  $|\mathbf{Z}(\mathfrak{C})| = p$  so  $\mathfrak{C}$  is type I or II. By Lemma 1.3,  $\mathfrak{B}/\mathfrak{A} \subseteq \mathbf{Z}(\mathfrak{C}) \times \mathbf{Z}(\mathfrak{C}) \times \cdots \times \mathbf{Z}(\mathfrak{C})$  ( $2m$  times). Hence  $[\mathfrak{B}:\mathfrak{A}] \leq p^{2m}$ . Since  $[\mathfrak{A}\mathfrak{C}:\mathfrak{A}] = p^{2m}$  we have  $\mathfrak{B} = \mathfrak{A}\mathfrak{C}$  here. Now let  $|\mathbf{Z}(\mathfrak{C})| = p^2$  so  $p = 2$  and  $\mathfrak{C}$  is type III or IV. As above  $\mathfrak{B}/\mathfrak{A} \subseteq \mathbf{Z}(\mathfrak{C}) \times \mathbf{Z}(\mathfrak{C}) \times \cdots \times \mathbf{Z}(\mathfrak{C})$  ( $2m$  times) so  $\mathfrak{B}/\mathfrak{A}$  is a 2-group. Since  $\mathfrak{A}$  is central in  $\mathfrak{G}$ ,  $\mathfrak{B}$  is nilpotent with Sylow 2-subgroup  $\mathfrak{B}_2$ . Now  $\mathfrak{G}$  is primitive and  $\mathfrak{B}_2 \triangleleft \mathfrak{G}$  so  $\mathfrak{B}_2$  is of symplectic type. Clearly  $\mathbf{Z}(\mathfrak{B}_2) = \mathfrak{A}_2$  and  $|\mathfrak{A}_2| \geq 4$  here. Hence  $\mathfrak{B}_2$  is the central product of  $\mathfrak{A}_2$  and a group of type  $E(2, r)$ . Thus  $\mathfrak{B}_2/\mathfrak{A}_2$  has period 2 and we can conclude again that  $[\mathfrak{B}:\mathfrak{A}] \leq p^{2m}$ . The result follows.

LEMMA 2.6. *We must have one of the following.*

- type I:  $p = 3, m \leq 2$
- type II:  $p = 2, m \leq 6$
- type III:  $p = 2, m \leq 3$
- type IV:  $p = 2, m \leq 5$ .

*Proof.* We first show the following.

- type I:  $w \leq p(2m - 1)|\mathfrak{A}_p|$   
 $w \leq |\mathfrak{A}_p|$  for  $m = 1, p > 3$
- type II:  $w \leq p^2(2m - 1)|\mathfrak{A}_p|$
- type III:  $w \leq p(2m - 1)|\mathfrak{A}_p|$
- type IV:  $w \leq p(2m - 1)|\mathfrak{A}_p|$ .

Now the  $p$ -period of  $Sp(2m, p)$  is clearly at most  $(2m - 1)p$ . If  $\mathfrak{C}$  is type I, III or IV, then the period of  $\mathfrak{A}_p\mathfrak{C}$  is  $|\mathfrak{A}_p|$ . If  $\mathfrak{C}$  is type II, then the period of  $\mathfrak{A}_p\mathfrak{C}$  is at most  $p|\mathfrak{A}_p|$ . Combining these facts with the structure given in the preceding lemma yields all the above facts except for the one concerning  $p > 3, m = 1$ .

Now let  $m = 1$  and  $p > 3$ . Let  $\mathfrak{P}$  be a Sylow  $p$ -subgroup of  $C_{\mathfrak{G}}(\mathbf{Z}(\mathfrak{C}))$  and hence a Sylow  $p$ -subgroup of  $\mathfrak{G}$ . Thus  $\mathfrak{P} \supseteq \mathfrak{A}_p\mathfrak{C}$  and since  $|Sp(2, p)|_p = p$  we have  $[\mathfrak{P}:\mathfrak{A}_p\mathfrak{C}] \leq p$ . Since  $\mathfrak{C}$  does not act semiregularly we have  $p \mid |\mathfrak{G}_x|$  for all  $x \in \mathfrak{B}^{\#}$ . As we have seen, there exists  $x \in \mathfrak{B}^{\#}$  with  $\mathfrak{G}_x \cap \mathfrak{A}_p\mathfrak{C} = \langle 1 \rangle$ . Let  $\bar{\mathfrak{P}}$  be a subgroup of  $\mathfrak{G}_x$  of order  $p$ . By taking a suitable conjugate of  $\mathfrak{P}$  if necessary we can assume that  $\bar{\mathfrak{P}} \subseteq \mathfrak{P}$ . Then  $\mathfrak{P} = \mathfrak{A}_p(\mathfrak{C}\bar{\mathfrak{P}})$ . Now  $|\mathfrak{C}\bar{\mathfrak{P}}| = p^4$  and this group is generated by elements of order  $p$ . Hence if  $p > 3$ , then  $\mathfrak{C}\bar{\mathfrak{P}}$  has period  $p$ . Since  $\mathfrak{A}_p$  is central in  $\mathfrak{P}$  we see that  $\mathfrak{P}$  has period  $|\mathfrak{A}_p|$  and the above follows.

Combining the above with the lower bound for  $w$  given in Lemma 2.4 yields the following equations.

- type I:  $p^m|\mathfrak{A}_p| \leq p(2m - 1)|\mathfrak{A}_p|$   
 $p|\mathfrak{A}_p| \leq |\mathfrak{A}_p|$  for  $m = 1, p > 3$
- type II:  $p^{m-1}|\mathfrak{A}_p| \leq p^2(2m - 1)|\mathfrak{A}_p|$
- type III:  $p^m|\mathfrak{A}_p| \leq p(2m - 1)|\mathfrak{A}_p|$

type IV :  $p^{m-1} | \mathfrak{A}_p | \leq p(2m - 1) | \mathfrak{A}_p |$ .

Note that the equations for types II, III and IV hold only for  $m \geq 3$ . The result now follows easily.

We note that the above yields a stronger result than Proposition 2.1 of [6] and the proof is considerably less computational. We now strengthen the above argument to eliminate additional cases. We first eliminate  $p = 3$ .

LEMMA 2.7.  $p = 3, m = 1$  does not occur.

*Proof.* Suppose  $p = 3$  and  $m = 1$ . Then  $\mathfrak{G}$  has the structure described in Lemma 2.5. In addition,  $[\mathfrak{G} : C_{\mathfrak{G}}(\mathbf{Z}(\mathfrak{G}))] = 1$  or 2 and  $Sp(2m, p) = SL(2, p)$ . By Lemma 2.4,  $\mathfrak{A}$  is central in  $C_{\mathfrak{G}}(\mathbf{Z}(\mathfrak{G})) = \mathfrak{G}$ .

Suppose that  $\mathfrak{G}/\mathfrak{A}\mathfrak{C}$  has a normal Sylow 3-subgroup  $\mathfrak{B}/\mathfrak{A}\mathfrak{C}$ . Then  $\mathfrak{B}/\mathfrak{A}$  is a normal Sylow 3-subgroup of  $\mathfrak{G}/\mathfrak{A}$ . Now both  $\mathfrak{G}$  and  $\mathfrak{A}$  act half-transitively so by Lemma 1.7  $\mathfrak{B}$  acts half-transitively on  $\mathfrak{B}^*$ . Since  $\mathfrak{A}$  is central in  $\mathfrak{B}$ ,  $\mathfrak{B}$  is nilpotent and hence its normal Sylow 3-subgroup  $\mathfrak{B}_3$  acts half-transitively. By Theorem II of [4],  $\mathfrak{B}_3$  is cyclic, a contradiction since  $\mathfrak{B}_3 \cong \mathfrak{C}$ . Hence  $\mathfrak{G}/\mathfrak{A}\mathfrak{C}$  is a subgroup of  $SL(2, 3)$  which does not have a normal Sylow 3-subgroup. This implies that  $\mathfrak{G}/\mathfrak{A}\mathfrak{C} \cong SL(2, 3)$ , a group of order 24.

We show now that we cannot have  $8 \mid |\mathfrak{G}_x|$  for all  $x \in \mathfrak{B}^*$ . Assume by way of contradiction that this is the case. Let  $\mathfrak{P}$  be a subgroup of  $\mathfrak{C}$  of order 3 having a fixed point  $y \neq 0$ . Since  $\mathfrak{A}_y = \langle 1 \rangle$  we see that  $8 \mid |\mathfrak{A}\mathfrak{G}_y/\mathfrak{A}|$  so  $4 \mid |\mathfrak{A}\mathfrak{G}_y/\mathfrak{A}|$ . Now a Sylow 2-subgroup of  $\mathfrak{G}/\mathfrak{A}$  is quaternion of order 8 so  $\mathfrak{G}_y$  has an element  $B$  of order 4. Since  $B^2 \in \mathfrak{A}\mathfrak{C}$ ,  $B$  does not normalize  $\mathfrak{P}\mathbf{Z}(\mathfrak{C})/\mathbf{Z}(\mathfrak{C})$ . Thus  $\mathfrak{C} = \langle \mathfrak{P}, \mathfrak{P}^B \rangle \subseteq \mathfrak{G}_y$ , a contradiction since  $\mathbf{Z}(\mathfrak{C})$  acts semiregularly.

Let  $\mathfrak{P}$  be a subgroup of  $\mathfrak{G}$  of order 3. We show that  $\dim C_{\mathfrak{B}}(\mathfrak{P}) = 0$  or  $s$ . Since  $\mathfrak{P} \cong GL(3, q^s)$  we see that  $\dim C_{\mathfrak{B}}(\mathfrak{P}) = 0, s$  or  $2s$ . Suppose the dimension is  $2s$ . By Lemma 1.4,  $\mathfrak{P} \not\subseteq \mathfrak{A}\mathfrak{C}$ . Since  $\mathfrak{G}/\mathfrak{A}\mathfrak{C} \cong SL(2, 3)$  there exists  $G \in \mathfrak{G}$  such that  $\mathfrak{P}$  and  $\mathfrak{P}^G$  generate this quotient. Now  $\mathfrak{B}$  is 3-dimensional over  $GF(q^s)$  and  $C_{\mathfrak{B}}(\mathfrak{P})$  and  $C_{\mathfrak{B}}(\mathfrak{P}^G)$  are 2-dimensional subspaces. Thus there exists  $x \in \mathfrak{B}^*$  with  $\mathfrak{P}, \mathfrak{P}^G \subseteq \mathfrak{G}_x$ . This implies that  $24 \mid |\mathfrak{G}_x|$  and this contradicts the comments of the preceding paragraph.

We now proceed to count. The group  $\mathfrak{G}/\mathfrak{A}$  is easily seen to contain at most 40 subgroups of order 3. If  $\mathfrak{P}$  is a group of order 3 in  $\mathfrak{G}$ , then  $\mathfrak{P}\mathfrak{A}$  being abelian has at most 3 subgroups of order 3 other than  $\mathbf{Z}(\mathfrak{C})$ . Hence  $\mathfrak{G}$  has at most  $3 \cdot 40 = 120$  subgroups of order 3 other than  $\mathbf{Z}(\mathfrak{C})$ . Each such  $\mathfrak{P}$  fixes at most  $q^s - 1$  points of  $\mathfrak{B}^*$  so since clearly  $3 \mid |\mathfrak{G}_x|$  we have

$$120(q^s - 1) \geq |\mathfrak{B}^*| = q^{3s} - 1$$

and

$$120 \geq q^{2s} + q^s + 1 .$$

Thus  $q^s \leq 10$ . However by Lemma 2.4,  $3^2 \mid (q^s - 1)$  so  $q^s$  being a prime power is at least 19, a contradiction. Thus  $p = 3$ ,  $m = 1$  does not occur.

LEMMA 2.8.  $p = 3$ ,  $m = 2$  does not occur.

*Proof.* The equation obtained in the proof of Lemma 2.6 is an equality at  $p = 3$ ,  $m = 2$ . Thus all inequalities used in obtaining it must also be equalities. Thus from Lemma 2.4 we must have  $w = p^m \mid \mathfrak{A}_p$ . Furthermore if  $x \in \mathfrak{B}^*$  with  $\mathfrak{G}_x \cap \mathfrak{A}_p \mathfrak{G} = \langle 1 \rangle$ , then  $|\mathfrak{G}|_p = |\mathfrak{G}_x|_p \mid \mathfrak{A}_p \mathfrak{G}$ .

The latter fact implies that  $\mathfrak{A}_p \mathfrak{G}$  has a complement  $\mathfrak{Z}$  in  $\mathfrak{P}$  a Sylow  $p$ -subgroup of  $\mathfrak{G}$ . Since  $\mathfrak{Z} \subseteq Sp(4, 3)$ ,  $\mathfrak{Z}$  has period at most  $(2m - 1)p = 9$  and thus  $\mathfrak{G}\mathfrak{Z}$  has period at most  $3 \cdot 9 = 27$ . Since  $\mathfrak{P} = \mathfrak{A}_p(\mathfrak{G}\mathfrak{Z})$  and  $\mathfrak{A}_p$  is central here, we have clearly  $w \leq \max \{ |\mathfrak{A}_p|, p^3 \}$ . But  $w = p^2 \mid \mathfrak{A}_p$  so we must have  $|\mathfrak{A}_p| = p$  and  $\mathfrak{Z}$  has period 9.

Let  $\mathfrak{J} = \langle J \rangle$  be a subgroup of order 9 with  $\mathfrak{J} \cap \mathfrak{A}_p \mathfrak{G} = \langle 1 \rangle$ . We see clearly that the Jordan form of the matrix of  $J$  with respect to its action on  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$  is

$$\begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} .$$

Thus  $\mathfrak{J}$  has

$$(3^4 - 3^3)/3^2 + (3^3 - 3)/3 + 3 = 17$$

orbits on  $\mathfrak{B}$ . Note that  $\mathfrak{G}\mathfrak{J} \subseteq GL(p^m, q^s)$  and the restriction to  $\mathfrak{G}$  is absolutely irreducible. Thus if  $a_0 = \dim_{GF(q^s)} C_{\mathfrak{B}}(J)$  then by Lemma 1.6

$$\begin{aligned} a_0 + a_1 + \dots + a_8 &= p^m = 9 . \\ a_0^2 + a_1^2 + \dots + a_8^2 &\leq 17 . \end{aligned}$$

These yield easily  $a_0 \leq 3$  and hence  $\dim_{GF(q)} C_{\mathfrak{B}}(J) = sa_0 \leq 3s$ .

Let  $\mathcal{N}$  denote the set of subgroups of  $\mathfrak{G}$  of order 3 together with the set of cyclic subgroup  $\mathfrak{J}$  of order 9 with  $\mathfrak{J} \cap \mathfrak{A}_p \mathfrak{G} = \langle 1 \rangle$ . By the above and Lemma 1.4, if  $\mathfrak{N} \in \mathcal{N}$  then  $\dim C_{\mathfrak{B}}(\mathfrak{N}) \leq 3s$ . We have also shown above that for all  $y \in \mathfrak{B}$  there exists  $\mathfrak{N} \in \mathcal{N}$  with  $y \in C_{\mathfrak{B}}(\mathfrak{N})$ , since in that argument, if  $\mathfrak{G}_x \cap \mathfrak{A}_p \mathfrak{G} = \langle 1 \rangle$  then  $\mathfrak{J} = \mathfrak{Z} \subseteq \mathfrak{G}_x$ . Hence  $\mathfrak{B} = \bigcup_{\mathfrak{N} \in \mathcal{N}} C_{\mathfrak{B}}(\mathfrak{N})$ . If  $|\mathcal{N}| = N$ , then this yields

$$q^{9s} = |\mathfrak{B}| \leq Nq^{3s}$$

or  $q^{6s} \leq N$ . On the other hand by Lemma 2.5,

$$\begin{aligned} |\mathfrak{G}| &\leq 2 |\mathfrak{A}| p^4 |Sp(4, p)| \\ &\leq 2 |\mathfrak{A}| p^4 p^4 (p^4 - 1)(p^2 - 1) \leq 2 |\mathfrak{A}| p^{14}. \end{aligned}$$

Since  $\mathfrak{A}$  is central in the absolutely irreducible representation  $\mathfrak{A}\mathfrak{G} \cong GL(p^m, q^s)$  we have  $|\mathfrak{A}| < q^s$ . Thus

$$N \leq |\mathfrak{G}|/2 < q^s p^{14}.$$

Combining this with the lower bound we previously obtained for  $N$  yields  $q^{5s} < p^{14}$ . Finally by Lemma 2.4,  $q^s \geq p^3$  so  $p^{15} < q^{5s} < p^{14}$ , a contradiction. Thus  $p = 3, m = 2$  does not occur.

We now consider special cases with  $p = 2$ .

LEMMA 2.9. *The cases type II,  $m = 6$ , type III,  $m = 3$  and type IV,  $m = 5$  do not occur.*

*Proof.* If we consider the inequalities obtained in the proof of Lemma 2.6, we see that any of the above mentioned cases would be eliminated if a strengthening of the inequalities by a factor of  $p = 2$  could be obtained. Let us suppose that one of the above occurs.

The results of Lemma 2.4 concerning  $[\mathfrak{G} : \mathfrak{G}_x]_p$  and  $w$  must be equalities. In particular this implies that for given  $x \in \mathfrak{B}^*$  with  $\mathfrak{G}_x \cap \mathfrak{A}_p \mathfrak{G} = \langle 1 \rangle$  we must have  $|\mathfrak{G}|_p = |\mathfrak{G}_x \mathfrak{A}_p \mathfrak{G}|_p$ . Thus  $\mathfrak{A}_p \mathfrak{G}$  has a complement in a Sylow  $p$ -subgroup of  $\mathfrak{G}$  and thus also in  $\mathfrak{P}$ , a Sylow  $p$ -subgroup of  $\mathfrak{G}$ . Let  $\mathfrak{P} = \mathfrak{A}_p \mathfrak{G} \mathfrak{L}$  where  $\mathfrak{L} \cap \mathfrak{A}_p \mathfrak{G} = \langle 1 \rangle$  and let  $w^*$  denote the period of the group  $\mathfrak{G} \mathfrak{L} / \mathfrak{G}'$ . Since  $\mathfrak{A}_p$  is central in  $\mathfrak{P}$  we have

$$w \leq \max \{ |\mathfrak{A}_p|, 2w^* \} \leq \begin{cases} |\mathfrak{A}_p| w^* & \text{type II} \\ \frac{1}{2} |\mathfrak{A}_p| w^* & \text{types III, IV.} \end{cases}$$

We consider  $w^*$ . Let  $\mathfrak{W}^* = \mathfrak{G} / \mathfrak{G}'$  so  $\mathfrak{W}^*$  is elementary abelian of order  $p^{2m}$  or  $p^{2m+1}$ . Since  $\mathfrak{L}$  acts faithfully on  $\mathfrak{G} / \mathfrak{Z}(\mathfrak{G})$ , it also acts faithfully on  $\mathfrak{W}^*$ . If  $\mathfrak{L}$  has period  $p^d$ , then  $w^* = p^d$  or  $p^{d+1}$ . Note that  $\mathfrak{L} \cong GL(2m + 1, p)$ . If  $w^* = p^d$ , then since  $p^d \leq p(2m)$  we have  $w^* \leq p(2m)$ . If  $w^* = p^{d+1}$ , then there must exist an element  $L \in \mathfrak{L}$  of order  $p^d$  whose minimal polynomial in  $GL(2m + 1, p)$  has degree  $p^d$ . Thus we must have  $p^d \leq 2m + 1$  and  $w^* \leq p(2m + 1)$ . The latter bound being the larger of the two holds in all cases. Now  $w^*$  is a power of 2 and in the three cases we are considering neither  $2m + 1$  nor  $2m$  is a power of 2. Hence we have  $w^* \leq p(2m - 1)$  and

$$\begin{aligned} w &\leq p(2m - 1) |\mathfrak{A}_p| && \text{for type II} \\ w &\leq (2m - 1) |\mathfrak{A}_p| && \text{for types III, IV.} \end{aligned}$$



This therefore improves the bounds on  $w$  given in the proof of Lemma 2.6 by a factor of  $p = 2$  and, as we mentioned above, this yields a contradiction.

LEMMA 2.10. *The case type IV,  $m = 4$  does not occur.*

*Proof.* We see that in the inequalities obtained in the proof of Lemma 2.6, a strengthening by a factor of  $p = 2$  would eliminate this possibility. Hence if this case occurs, then we must have the following. If  $x \in \mathfrak{B}^*$ , then either  $x$  is fixed by a subgroup of  $\mathfrak{G}$  of order 2 or a cyclic subgroup  $\mathfrak{F} \subseteq \mathfrak{G}$  of order 8 with  $\mathfrak{F} \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$ . Let  $\mathcal{N}$  denote collection of such subgroups of both types.

We show now that if  $\mathfrak{F} \in \mathcal{N}$  then  $\dim C_{\mathfrak{B}}(\mathfrak{F}) \leq n/2$ . We know this to be the case if  $\mathfrak{F} \subseteq \mathfrak{G}$  so suppose  $\mathfrak{F} = \langle J \rangle$  has order 8. Then  $\mathfrak{F}$  acts faithfully on  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$ . Since  $|\mathfrak{F}| = 8$  we see that in its action on  $\mathfrak{B}$ ,  $J$  must have one Jordan block of rank at least 5. This implies easily that  $\mathfrak{F}$  has at most

$$\frac{2^8 - 2^7}{8} + \frac{2^7 - 2^5}{4} + \frac{2^5 - 2^4}{2} + 2^4 = 2^6$$

orbits on  $\mathfrak{B}$ . We apply Lemma 1.6 to each of the two absolutely irreducible constituents of  $\mathfrak{G}\mathfrak{F}$  on  $\mathfrak{B} \otimes GF(q^2)$ . Hence

$$a_0^2 + a_1^2 + \dots + a_7^2 \leq 2^6.$$

Thus  $a_0 \leq 8$  and since  $\dim C_{\mathfrak{B}}(\mathfrak{F})$  is invariant under field extension we have  $\dim C_{\mathfrak{B}}(\mathfrak{F}) \leq 2a_0 \leq n/2$ . Now

$$\mathfrak{B} = \bigcup_{\mathfrak{F} \in \mathcal{N}} C_{\mathfrak{B}}(\mathfrak{F})$$

and if  $N = |\mathcal{N}|$ , then  $q^n = |\mathfrak{B}| \leq Nq^{n/2}$  and  $q^{n/2} \leq N$ . By Lemma 2.5

$$|\mathfrak{G}| \leq |\mathfrak{A}| 2^{2m} |Sp(2m, 2)|.$$

Since  $|\mathfrak{A}| \leq q^s$  and  $|Sp(8, 2)| \leq 2^{36}$  we have  $N \leq |\mathfrak{G}| \leq q^s \cdot 2^{44}$ . With  $n = 2^m s = 16s$  this yields

$$q^{8s} = q^{n/2} \leq N \leq q^s \cdot 2^{44}$$

or  $q^{7s} \leq 2^{44}$ . Now  $s = 2$  and by Lemma 2.4,  $2^5 = 2^{m+1}$  divides  $q^2 - 1 = q^s - 1$ . Since  $s = 2$  and  $q^s > 9$  it follows (see for example Lemma 4 of [4]) that  $q^s - 1$  cannot be a power of 2. Hence  $q^s > q^s - 1 \geq 3 \cdot 2^5$  so  $q^{7s} > 3^7 \cdot 2^{35}$ . Combining this with the above yields  $3^7 \cdot 2^{35} < q^{7s} \leq 2^{44}$  or  $3^7 < 2^9$ , a contradiction. Therefore this case does not occur.

LEMMA 2.11. *The case type II,  $m = 5$  does not occur.*

*Proof.* In the inequality in the proof of Lemma 2.6 for type II,  $m = 5$  we see that a strengthening by a factor of  $p^2 = 4$  will yield a contradiction. Hence if  $x \in \mathfrak{B}^*$  is such that  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{C} = \langle 1 \rangle$  and if  $\mathfrak{P}$  is a Sylow 2-subgroup of  $\mathfrak{G}$  extending one of  $\mathfrak{G}_x$ , then either (a)  $\mathfrak{P}_x\mathfrak{C} = \mathfrak{P}$  and  $w \geq 32$  or (b)  $[\mathfrak{P} : \mathfrak{P}_x\mathfrak{C}] = 2$  and  $w \geq 64$ . In the latter case  $\mathfrak{P}_x\mathfrak{C} \triangleleft \mathfrak{P}$  so in both cases  $\mathfrak{P}_x\mathfrak{C}$  has period  $\geq 32$  and  $\mathfrak{P}_x$  has period  $\geq 8$ . Note  $|\mathfrak{A}_2| = 2$  here by Lemma 2.6.

Let  $\mathfrak{S} = \langle J \rangle$  be a cyclic subgroup of  $\mathfrak{P}_x$  of order 8 and let  $a_0 = \dim_{GF(q)} C_{\mathfrak{B}}(J)$ . Since  $\mathfrak{S}$  acts faithfully on  $\mathfrak{B} = \mathfrak{C}/\mathfrak{Z}(\mathfrak{C})$  and  $|\mathfrak{S}| = 8$  we see that  $J$  must have one Jordan block of rank at least 5. This implies easily that  $\mathfrak{S}$  has at most

$$\frac{2^{10} - 2^9}{8} + \frac{2^9 - 2^7}{4} + \frac{2^7 - 2^5}{2} + 2^5 = 2^8$$

orbits on  $\mathfrak{B}$ . Hence by Lemma 1.6

$$\begin{aligned} a_0 + a_1 + \dots + a_7 &= p^m = 32 \\ a_0^2 + a_1^2 + \dots + a_7^2 &\leq 2^8. \end{aligned}$$

Thus  $a_0 < 2^4 = 16$  and  $|C_{\mathfrak{B}}(J)| = q^{a_0} \leq q^{15}$ .

Now if  $\mathfrak{X}$  is a subgroup of  $\mathfrak{A}\mathfrak{C}$  of order 2 then  $|C_{\mathfrak{B}}(\mathfrak{X})| \leq q^{n/2} = q^{16}$ . We have shown that with the above notation

$$\mathfrak{B} = \bigcup_{\mathfrak{S}} C_{\mathfrak{B}}(\mathfrak{S}) \cup \bigcup_{\mathfrak{X}} C_{\mathfrak{B}}(\mathfrak{X}).$$

Now  $\mathfrak{A}$  is cyclic and central and by Lemma 2.6,  $4 \nmid |\mathfrak{A}|$ . Hence the number of choices for  $\mathfrak{X}$  is at most  $|\mathfrak{C}| = 2^{11}$  and the number of choices for  $\mathfrak{S}$  is at most  $1/4 |\mathfrak{G}/\mathfrak{A}_2|$ . Here  $\mathfrak{A}_2$  is the normal 2-complement of  $\mathfrak{A}$  and the  $1/4$  factor comes from the fact that  $\mathfrak{S}$  has four distinct generators. Since  $|\mathfrak{G}/\mathfrak{A}_2| \leq |\mathfrak{C}| |Sp(10, 2)| \leq 2^{66}$ , the above union yields

$$q^{32} = |\mathfrak{B}| \leq 2^{66}q^{15}/4 + 2^{11}q^{16}.$$

Putting  $q^{15} < q^{16}/2$  in the above we have

$$q^{32} < (2^{63} + 2^{11})q^{16} < 2^{64}q^{16}$$

so  $q^{16} < 2^{94}$  and  $q < 2^4 = 16$ . On the other hand by Lemma 2.4  $2^5 | q^2 - 1$  so  $16 | q \pm 1$ . This yields  $q \geq 17$ , a contradiction.

The following partial result will be completed later under the additional assumption of solvability.

**LEMMA 2.12.** *In the case type II,  $m = 4$  we have  $q \geq 7$  and  $|\mathfrak{G}/\mathfrak{A}\mathfrak{C}| > 10^4$ .*

*Proof.* In the inequalities of the proof of Lemma 2.6 for type II,  $m = 4$ , we see that a strengthening by a factor of  $p^2 = 4$  will yield a contradiction. Suppose  $x \in \mathfrak{B}^*$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$  and let  $\mathfrak{P}$  be a Sylow 2-subgroup of  $\mathfrak{G}$  extending one of  $\mathfrak{G}_x$ . Using the same argument as in the preceding lemma we conclude that  $\mathfrak{P}_x\mathfrak{G}$  has period  $\geq 16$  and hence  $\mathfrak{P}_x\mathfrak{G}/\mathfrak{Z}(\mathfrak{G})$  has period  $\geq 8$ .

Suppose first that  $\mathfrak{P}_x$  has a cyclic subgroup  $\mathfrak{J} = \langle J \rangle$  of order 8. Then in its action on  $\mathfrak{B} = \mathfrak{G}/\mathfrak{Z}(\mathfrak{G})$ ,  $J$  has a Jordan block of rank at least 5 so  $\mathfrak{J}$  has at most

$$\frac{2^8 - 2^7}{8} + \frac{2^7 - 2^5}{4} + \frac{2^5 - 2^4}{2} + 2^4 = 64$$

orbits on  $\mathfrak{B}$ . By Lemma 1.6 if  $a_0 = \dim C_{\mathfrak{B}}(J)$  then

$$a_0^2 + a_1^2 + \dots + a_7^2 \leq 64$$

and  $a_0 \leq 8$ .

Now suppose  $\mathfrak{P}_x$  has period 4. Then since  $\mathfrak{P}_x\mathfrak{G}/\mathfrak{Z}(\mathfrak{G})$  has period 8,  $\mathfrak{P}_x$  must contain an element  $J$  of order 4 with a Jordan block of rank 4. If  $\mathfrak{J} = \langle J \rangle$ , then  $\mathfrak{J}$  has at most

$$\frac{2^8 - 2^6}{4} + \frac{2^6 - 2^5}{2} + 2^5 = 96$$

orbits on  $\mathfrak{B}$ . By Lemma 1.6 if  $a_0 = \dim C_{\mathfrak{B}}(J)$  then

$$\begin{aligned} a_0 + a_1 + a_2 + a_3 &= p^m = 16 \\ a_0^2 + a_1^2 + a_2^2 + a_3^2 &\leq 96. \end{aligned}$$

It is easy to see that the possibility  $a_0 = 9$  is excluded and hence in both cases  $a_0 \leq 8$ .

We have

$$\mathfrak{B} = \bigcup C_{\mathfrak{B}}(\mathfrak{J}) \cup \bigcup C_{\mathfrak{B}}(\mathfrak{I})$$

where the subgroups  $\mathfrak{J}$  are as above and the subgroups  $\mathfrak{I}$  have order 2 and are contained in  $\mathfrak{G}$ . This follows since  $4 \nmid |\mathfrak{A}|$  by Lemma 2.6. The number of choices for  $\mathfrak{J}$  or  $\mathfrak{I}$  is clearly at most  $|\mathfrak{G}/\mathfrak{A}_2|$  where  $\mathfrak{A}_2$  is the normal 2-complement of  $\mathfrak{A}$ . Since  $4 \nmid |\mathfrak{A}|$  we have  $|\mathfrak{G}/\mathfrak{A}_2| = 2^9 |\mathfrak{G}/\mathfrak{A}\mathfrak{G}|$ . Therefore the above union yields

$$q^{16} = |\mathfrak{B}| \leq q^8 2^9 |\mathfrak{G}/\mathfrak{A}\mathfrak{G}|$$

since  $|C_{\mathfrak{B}}(\mathfrak{J})|$  and  $|C_{\mathfrak{B}}(\mathfrak{I})|$  are both at most  $q^8$ . Thus  $|\mathfrak{G}/\mathfrak{A}\mathfrak{G}| \geq q^8/2^9$ . By Lemma 2.4,  $2^4 |q^2 - 1|$  so  $q \geq 7$ . This yields

$$|\mathfrak{G}/\mathfrak{A}\mathfrak{G}| \geq 7^8/2^9 = (2401)^2/2^9 > 10^4$$

and the result follows.

We now temporarily drop the assumptions stated at the beginning of this section and prove the first of our three theorems.

*Proof of Theorem A.* Let  $\mathfrak{G}$  be a linear group acting on vector space  $\mathfrak{V}$  of order  $q^n$  and suppose that  $\mathfrak{G}$  acts half-transitively but not semiregularly on  $\mathfrak{V}^*$ . Let  $\mathfrak{P} = O_p(\mathfrak{G})$  be the maximal normal  $p$ -subgroup of  $\mathfrak{G}$ . By assumption  $\mathfrak{G}$  is primitive so  $\mathfrak{P}$  is of symplectic type. Suppose first that  $p > 2$ . If  $\mathfrak{P}$  is not cyclic, then  $\mathfrak{P}$  contains a characteristic subgroup  $\mathfrak{C}$  of type  $E(p, m)$ . By the Reduction Lemma (Lemma 1.8) and Lemmas 2.6, 2.7 and 2.8 we have a contradiction.

Now let  $p = 2$  so that  $\Phi(\mathfrak{P})$  is cyclic. Suppose  $[\mathfrak{P} : \Phi(\mathfrak{P})] > 2^8$ . Then  $\mathfrak{P}$  has a characteristic subgroup  $\mathfrak{C}$  of type  $E(2, m)$  with  $m > 3$ . Thus by the Reduction Lemma and Lemmas 2.6 through 2.11 we see that  $m = 4$  and  $|Z(\mathfrak{C})| = 2$ . But then  $|\Phi(\mathfrak{P})| = 2$  so  $\mathfrak{P} = \mathfrak{C}$  and  $[\mathfrak{P} : \Phi(\mathfrak{P})] \leq 2^8$  here also. This completes the proof.

**3. Solvable cases,  $m = 1$ .** We have seen in the preceding section that if  $\mathfrak{C}$  is a group of type  $E(p, m)$  normal in a half-transitive linear group  $\mathfrak{G}$ , then  $p = 2$  and  $m \leq 4$ . We will consider these cases in the next few sections under the additional assumption that  $\mathfrak{G}$  is solvable.

For convenience we restate Lemmas 1.3 and 1.4 of [5].

**LEMMA 3.1.** *Suppose  $\mathfrak{G}$  is an irreducible linear group of degree  $n$  over  $GF(q)$  and  $\mathfrak{A} = \langle \mathfrak{A} \rangle$  is a cyclic normal subgroup all of whose irreducible constituents are similar. Let  $\zeta$  be an eigenvalue of  $A$  with  $GF(q)(\zeta) = GF(q^r)$  and  $n/r = k$ . Let  $p$  be a prime and suppose that for all vectors  $x$ ,  $p \mid |\mathfrak{G}_x|$ . Consider those subgroups  $\mathfrak{P}/\mathfrak{A}$  of  $\mathfrak{G}/\mathfrak{A}$  of order  $p$  for which there exists an  $x \neq 0$  with  $\mathfrak{P} \cap \mathfrak{G}_x \neq \langle 1 \rangle$ . If  $\lambda_1$  of the  $\mathfrak{P}$  are contained in  $C_{\mathfrak{G}}(\mathfrak{A})$  and  $\lambda_2$  are not, then*

- (i)  $\frac{q^{kr} - 1}{q^r - 1} \leq \lambda_1 \left\{ 1 + \frac{q^{r(k-1)} - 1}{q^r - 1} \right\} + \lambda_2 \left\{ \frac{q^{rk/p} - 1}{q^{r/p} - 1} \right\}$
- (ii)  $q^r + 1 \leq 2\lambda_1 + \lambda_2(q^{r/p} + 1)$  for  $k = 2$
- (iii)  $q^r < 2(\lambda_1 + \lambda_2)$  for  $k > 2$ .

This is a very coarse statement which we will have to strengthen at times. The following assumptions hold throughout the remainder of this section.

**ASSUMPTIONS.** Group  $\mathfrak{G}$  acts faithfully on vector space  $\mathfrak{V}$  of order  $q^n$  and half-transitively but not semiregularly on  $\mathfrak{V}^*$ .  $\mathfrak{C}$  is a group of type  $E(2, 1)$  which is normal in  $\mathfrak{G}$  and acts irreducibly on  $\mathfrak{V}$ .

Note that we do not assume that  $\mathfrak{G}$  is primitive here. The

reason for this, is that part (v) of the Reduction Lemma does not guarantee primitivity in this case.

LEMMA 3.2. *Let  $\mathfrak{G} \cong \mathfrak{D}$  (that is,  $\mathfrak{G} = \text{iso } I$ ). Then  $q^n = 3^2, 5^2, 7^2, 11^2$  or  $17^2$ .*

*Proof.* Clearly  $q^n = q^2$  and hence  $C_{\mathfrak{G}}(\mathfrak{G})$  consists of scalar matrices so  $C_{\mathfrak{G}}(\mathfrak{G}) = Z(\mathfrak{G})$ . Note that  $\text{Aut } \mathfrak{G} \cong \text{Sym}_4$ , the symmetric group of degree 4.

Suppose first that  $3 \nmid |\mathfrak{G}/Z(\mathfrak{G})|$ . Then  $|\mathfrak{G}/Z(\mathfrak{G})| = 4$  or 8 and hence  $\mathfrak{G}$  is nilpotent. Thus  $\mathfrak{G}_2 = O_2(\mathfrak{G})$  is half-transitive. Since  $O_2(\mathfrak{G}) \subseteq Z(\mathfrak{G})$  acts semiregularly, we conclude that  $\mathfrak{G}_2$  is not semi-regular. Hence  $\mathfrak{G}_2 > \mathfrak{G}$  and since  $[\mathfrak{G}_2 : Z(\mathfrak{G}_2)] = 4$  or 8 we have by Theorem II of [4],  $q^n = 3^2, 5^2$  or  $7^2$ .

We assume now that  $3 \mid |\mathfrak{G}/Z(\mathfrak{G})|$ . We consider the possibility  $3 \mid |\mathfrak{G}_x|$  first. If  $\mathfrak{L}$  is a subgroup of  $\mathfrak{G}$  of order 3 fixing a vector  $x$ , then  $|C_{\mathfrak{G}}(\mathfrak{L})| = q$  clearly. Also either  $q = 3$  or by complete reducibility  $3 \mid q - 1$ . Now  $\mathfrak{G}/Z(\mathfrak{G})$  has at most 4 subgroups of order 3 and since  $Z(\mathfrak{G})$  is cyclic, we see that  $\mathfrak{G}$  contains at most  $4 \cdot 3 = 12$  subgroups of order 3 not contained in  $Z(\mathfrak{G})$ . From  $\mathfrak{B} = \bigcup C_{\mathfrak{G}}(\mathfrak{L})$  we obtain easily

$$q^2 - 1 = |\mathfrak{B}^\#| \leq 12(q - 1)$$

so  $q + 1 \leq 12$ . Since either  $q = 3$  or  $3 \mid q - 1$  we have  $q = 3$  or 7 here.

We now assume that  $3 \nmid |\mathfrak{G}_x|$ . If  $\mathfrak{G}_x \cap Z(\mathfrak{G})\mathfrak{G} \neq \langle 1 \rangle$  for all  $x \in \mathfrak{B}^\#$ , then by Lemma 1.5  $q^n = 3^2$  or  $5^2$ . Thus we can suppose that some  $x \in \mathfrak{B}^\#$ ,  $\mathfrak{G}_x \cap Z(\mathfrak{G})\mathfrak{G} = \langle 1 \rangle$ . This yields  $|\mathfrak{G}_x| = 2$  and by Lemma 1.9,  $I(\mathfrak{G}) = q + 1$ . We have actually shown above that  $|\mathfrak{G}/Z(\mathfrak{G})|$  is divisible by  $3 \cdot 8$  so  $\mathfrak{G}/Z(\mathfrak{G}) \cong \text{Sym}_4$  and this group has two conjugacy classes of involutions,  $\mathfrak{C}_1$  of size 3 and  $\mathfrak{C}_2$  of size 6. If  $\bar{T} \in \mathfrak{G}/Z(\mathfrak{G})$  is an involution then since  $Z(\mathfrak{G})$  is cyclic of even order and central, the coset corresponding to  $\bar{T}$  will contain either 0 or 2 noncentral involutions of  $\mathfrak{G}$  and this number is the same for all conjugates of  $\bar{T}$ . Thus we have

$$q + 1 = I(\mathfrak{G}) = \delta_1 \cdot 2 \cdot 3 + \delta_2 \cdot 2 \cdot 6$$

where  $\delta_1, \delta_2 = 0$  or 1. Moreover since for some  $x \in \mathfrak{B}^\#$ ,  $\mathfrak{G}_x \cap Z(\mathfrak{G})\mathfrak{G} = \langle 1 \rangle$  we have  $\delta_2 = 1$ . Thus  $q + 1 = 6\delta_1 + 12$  and  $q = 11$  or 17. This completes the proof.

LEMMA 3.3. *Let  $\mathfrak{G} \cong \mathfrak{D}$  (that is,  $\mathfrak{G} = \text{iso } II$ ). Then  $q^n = 3^2, 5^2$  or  $7^2$ .*

*Proof.* Clearly  $q^n = q^2$  so  $C_{\mathfrak{G}}(\mathfrak{G}) = Z(\mathfrak{G})$  consists of scalar matrices.

Now  $|\text{Aut } \mathbb{G}| = 8$  so  $[\mathbb{G} : \mathbf{Z}(\mathbb{G})] = 4$  or  $8$  and hence  $\mathbb{G}$  is nilpotent. Then  $\mathbb{G}_2 = O_2(\mathbb{G})$  is half-transitive but not semiregular and  $[\mathbb{G}_2 : \mathbf{Z}(\mathbb{G}_2)] = 4$  or  $8$ . By Theorem II of [4],  $q^n = q^2 = 3^2, 5^2$  or  $7^2$ .

LEMMA 3.4. *Let  $\mathbb{G} \cong 3\mathcal{Q}$  (that is,  $\mathbb{G} = \text{iso III}$ ). Then  $q^n = 5^2, 17^2$  or  $\mathbb{G}$  is imprimitive and  $q^n = 3^4$ .*

*Proof.* Here  $q^n = q^2$  if  $q \equiv 1 \pmod{4}$  and  $q^n = q^4$  if  $q \equiv -1 \pmod{4}$ . Say  $q^n = q^{2r}$ .

Suppose first that  $\mathbb{G}$  is imprimitive. Here we can apply Theorem 1.1. Note that if  $q^r - 1$  is not a power of 2 then  $O_2(\mathcal{F}_0(q^r))$  is abelian. Hence by Theorem 1.1 either  $q^n = 3^4$  or  $\mathbb{G} = \mathcal{F}_0(q)$  for Fermat prime  $q$ . Here  $q \equiv 1 \pmod{4}$  so  $q \geq 5$ . Let  $\mathfrak{B}$  be the diagonalized subgroup of  $\mathbb{G}$  of index 2 so  $\mathfrak{B}$  is abelian. Then  $\mathbb{G} = \mathfrak{B}\mathbb{G}$  and  $\mathbb{G}' = (\mathfrak{B}, \mathbb{G}) \subseteq \mathbb{G}$ . Since  $\mathbb{G}'$  is cyclic of order  $(q - 1)/2$  we have  $(q - 1)/2 \leq 4$  so  $q \leq 9$  and hence since  $q$  is a Fermat prime  $q = 5$  and  $q^n = 5^2$ .

Now we assume that  $\mathbb{G}$  is primitive and we use the notation of Lemma 2.5. Then  $[\mathbb{G} : \mathfrak{H}] = 1$  or  $2$  where  $\mathfrak{H} = C_{\mathbb{G}}(\mathbf{Z}(\mathbb{G}))$  and  $\mathfrak{H}/\mathfrak{A}\mathbb{G} \subseteq Sp(2, 2) = SL(2, 2)$ , a group of order 6. Now  $\mathbb{G}$  has precisely 3 abelian subgroups of order 8 and these are not cyclic. Since  $\mathbb{G}$  is primitive none of these groups is normal. Hence  $\mathbb{G}$  permutes these transitively so  $3 \mid |\mathfrak{H}/\mathfrak{A}\mathbb{G}|$ . Now  $\mathbb{G}/\mathfrak{A}\mathbb{G}$  also acts on  $\mathbb{G}/\mathbb{G}'$  and this action is clearly faithful on  $\mathfrak{H}/\mathfrak{A}\mathbb{G}$ . If  $\mathfrak{F} = \mathfrak{F}/\mathfrak{A}\mathbb{G}$  is the normal 3-subgroup of  $\mathfrak{H}/\mathfrak{A}\mathbb{G}$  then  $\mathfrak{F}$  centralizes  $\mathbf{Z}(\mathbb{G})/\mathbb{G}'$  and acts faithfully on the commutator  $\mathbb{G}_0/\mathbb{G}'$ , a 2-dimensional complement. Clearly  $\mathbb{G}_0 \cong \mathcal{Q}$  and  $\mathbb{G}_0 \triangleleft \mathbb{G}$ . If  $n = 2$ , then by Lemma 3.2 and the fact that  $q \equiv 1 \pmod{4}$  we have  $q^n = 5^2$  or  $17^2$ .

Let  $n = 4$  so  $q \equiv -1 \pmod{4}$ .  $\mathbb{G}/\mathfrak{A}$  acts on  $\mathbb{G}_0$  and the kernel acts faithfully on  $\mathbf{Z}(\mathbb{G})$ . Thus we see that either  $\mathbb{G}/\mathfrak{A} \subseteq \text{Aut } \mathbb{G}_0 = \text{Sym}_4$  or  $\mathbb{G}/\mathfrak{A} \subseteq \mathfrak{H}/\mathfrak{A} \times \mathfrak{F}/\mathfrak{A} \subseteq \text{Sym } 4 \times \mathfrak{F}$  where  $|\mathfrak{F}| = |\mathfrak{F}/\mathfrak{A}| = 2$ . We apply Lemma 3.1 with  $p = 2$ . We have clearly  $\lambda_1 \leq 9, \lambda_2 \leq 10$  and since  $r = 2, k = 2, n = 4$  we obtain

$$q^2 + 1 \leq 18 + 10(q + 1)$$

or  $q(q - 10) \leq 27$  so  $q < 13$ . Since  $q \equiv 3 \pmod{4}$  we have  $q \equiv 3, 7$  or  $11$ . Suppose  $3 \mid |\mathbb{G}_x|$ . Let  $T$  be a noncentral involution of  $\mathbb{G}$ . By Lemma 1.5 there exists a point  $x \in \mathfrak{B}^*$  with  $\mathbb{G}_x = \langle T \rangle$ . Let  $\mathfrak{L}$  be a subgroup of  $\mathbb{G}_x$  of order 3. Then  $\mathfrak{L} \cap \mathfrak{A}\mathbb{G} = \langle 1 \rangle$ ,  $\mathfrak{L} \subseteq \mathfrak{H}$  and  $\mathfrak{L}$  normalizes  $\mathbb{G}_x \cap \mathbb{G} = \mathbb{G}_x$ , a contradiction since  $\mathfrak{L}$  acts irreducibly on  $\mathbb{G}/\mathbf{Z}(\mathbb{G})$ . Hence  $3 \nmid |\mathbb{G}_x|$  and since  $3 \mid |\mathbb{G}|$  we conclude that  $q \neq 3$ .

Let  $q = 7$  or  $11$ . By Lemma 1.5 there exists a point  $x \in \mathfrak{B}^*$  with  $\mathbb{G}_x \cap \mathfrak{A}\mathbb{G} = \langle 1 \rangle$ . Since  $3 \nmid |\mathbb{G}_x|$  we see that  $|\mathbb{G}_x| = 2$  or  $4$ . Suppose  $|\mathbb{G}_x| = 4$ . Then certainly  $2 \mid |\mathfrak{H}_x|$  for all  $x \in \mathfrak{B}^*$  and Lemma 3.1

applies to  $\mathfrak{G}$ . Here  $\lambda_1 \leq 9$ ,  $\lambda_2 = 0$ ,  $r = 2$ ,  $n = 4$ ,  $k = 2$  so

$$q^2 + 1 \leq 2 \cdot 9 + 0,$$

a contradiction. Thus  $|\mathfrak{G}_x| = 2$  and by Lemma 1.9,  $I(\mathfrak{G}) = q^2 + 1$ . Let  $\mathfrak{S}$  be a Sylow 3-subgroup of  $\mathfrak{G}$ . Then  $\mathfrak{S}$  permutes by conjugation the noncentral involutions of  $\mathfrak{G}$ . Since  $3 \nmid (q^2 + 1)$ ,  $\mathfrak{S}$  must centralize such an involution. Now subgroups of  $\text{Sym}_4$  of order 3 are self-centralizing so this implies that  $\mathfrak{G}/\mathfrak{A} \not\subseteq \text{Sym}_4$ . Hence  $\mathfrak{G}/\mathfrak{A} \cong \mathfrak{S}/\mathfrak{A} \times \mathfrak{T}/\mathfrak{A}$  where  $\mathfrak{S}/\mathfrak{A} \cong \text{Sym}_4$  and  $|\mathfrak{T}/\mathfrak{A}| = 2$ . Clearly  $\mathfrak{S}/\mathfrak{A} \cong \text{Alt}_4$  and if  $\mathfrak{S}/\mathfrak{A} \cong \text{Alt}_4$  then in the notation of Lemma 3.1 with  $p = 2$ ,  $\lambda_1 \leq 3$ ,  $\lambda_2 \leq 4$  and

$$(q^2 + 1) \leq 2\lambda_1 + (q + 1)\lambda_2 \leq 6 + 4(q + 1)$$

a contradiction for  $q = 7, 11$ . Hence  $\mathfrak{S}/\mathfrak{A} \cong \text{Sym}_4$  and  $\mathfrak{G}/\mathfrak{A}$  has five classes  $\mathfrak{C}_i$  of involutions. These satisfy  $\mathfrak{C}_1, \mathfrak{C}_2 \subseteq \mathfrak{S}/\mathfrak{A}$  with  $|\mathfrak{C}_1| = 3$ ,  $|\mathfrak{C}_2| = 6$  and  $\mathfrak{C}_3, \mathfrak{C}_4, \mathfrak{C}_5 \not\subseteq \mathfrak{S}/\mathfrak{A}$  with  $|\mathfrak{C}_3| = 1$ ,  $|\mathfrak{C}_4| = 3$ ,  $|\mathfrak{C}_5| = 6$ .

Let  $\bar{T}$  be an involution of  $\mathfrak{G}/\mathfrak{A}$ . If the coset of  $\bar{T}$  contains  $\alpha$  involutions, then the same is true for all conjugates of  $\bar{T}$ . If  $\bar{T} \in \mathfrak{S}/\mathfrak{A}$  then certainly  $\alpha = 0$  or  $2$ . If  $\bar{T} \notin \mathfrak{S}/\mathfrak{A}$ , then by Lemma 1.1 of [5]  $\bar{T}$  acts on  $\mathfrak{A}$  like a field automorphism of  $GF(q^2)$  of order 2 (that is, the map  $x \rightarrow x^q$ ). Suppose the coset contains an involution  $T$ . Then for  $B \in \mathfrak{A}$ ,  $BT$  is an involution if and only if  $B^q = B^T = B^{-1}$ . Hence  $\alpha = 0$  or the number  $N$  of elements of  $\mathfrak{A}$  of order dividing  $q + 1$ . Note that since  $|Z(\mathfrak{G})| = 4$  we have  $N = 4$  or  $8$  for  $q = 7$  and  $N = 4$  or  $12$  for  $q = 11$ . Now if  $\delta_i = 1$  or  $0$  according to whether the coset of  $\bar{T} \in \mathfrak{C}_i$  contains an involution of  $\mathfrak{G}$  then we obtain

$$q^2 + 1 = I(\mathfrak{G}) = 6\delta_1 + 12\delta_2 + N(\delta_3 + 3\delta_4 + 6\delta_5).$$

Considering this modulo 3 we have

$$2 \equiv q^2 + 1 \equiv N\delta_3 \pmod{3}.$$

This shows that  $q \neq 11$ . If  $q = 7$  then  $N = 8$  so  $8 \mid |\mathfrak{A}|$  and  $\delta_3 = 1$ . Furthermore  $\delta_5 = 0$  and then  $\delta_1 = \delta_2 = \delta_3 = \delta_4 = 1$ .

Since  $\delta_2 = 1$  we can find an involution  $T \in \mathfrak{S}$  corresponding to a transposition in  $\mathfrak{S}/\mathfrak{A} \cong \text{Sym}_4$ . Now  $T$  normalized  $\mathfrak{C}_0 \cong \Omega$  as mentioned before and  $T$  does not fix  $\mathfrak{C}_0/\mathfrak{C}'_0$  since  $T$  does not fix  $\mathfrak{C}/Z(\mathfrak{G})$ . Thus  $\langle \mathfrak{C}_0, T \rangle$  is a maximal class group of order 16 and hence this group has a cyclic subgroup  $\mathfrak{B}$  of order 8. The group  $\mathfrak{A}_2\mathfrak{B}$  is abelian and has period  $|\mathfrak{A}_2|$  since  $\mathfrak{B} \subseteq \mathfrak{S}$  and  $|\mathfrak{A}_2| \geq |\mathfrak{B}|$ . Also  $|\mathfrak{B} \cap \mathfrak{A}_2| = 2$  so  $|\mathfrak{A}_2\mathfrak{B}| = 4|\mathfrak{A}_2|$ . Let  $\mathfrak{U} \subseteq \mathfrak{B}$  be an irreducible  $\mathfrak{A}_2\mathfrak{B}$ -submodule and let  $\mathfrak{K} \subseteq \mathfrak{A}_2\mathfrak{B}$  be the kernel. Then  $\mathfrak{A}_2\mathfrak{B}/\mathfrak{K}$  is cyclic so  $|\mathfrak{A}_2\mathfrak{B}/\mathfrak{K}| \leq |\mathfrak{A}_2|$  and

hence  $|\mathfrak{R}| \geq 4$ . If  $x \in \mathfrak{U}^\#$ , then  $\mathfrak{G}_x \cong \mathfrak{R}$  and  $|\mathfrak{G}_x| \geq 4$ , a contradiction. This completes the proof.

EXAMPLES. The examples with  $q^n = 3^2, 5^2, 7^2$  and  $11^2$  can occur as transitive groups and these are given in [3]. We consider the case  $q^n = 17^2$ . Let  $SL(2, 17)^*$  denote the subgroup of  $GL(2, 17)$  consisting of those matrices with determinant  $\pm 1$ . Let  $\mathfrak{G} = \mathfrak{Q}\mathfrak{B}$  where  $\mathfrak{Q}$  is the quaternion group of order 8,  $\mathfrak{Q} \triangle \mathfrak{G}$  and  $\mathfrak{B} \cong \text{Sym}_3$  acts faithfully on  $\mathfrak{Q}/\mathfrak{Q}'$ . Clearly  $\mathfrak{G}' = \mathfrak{Q}\mathfrak{B}' \cong SL(2, 3)$ . This group has a unique faithful irreducible rational character of degree 2. Hence  $\mathfrak{G}$  has a faithful character  $\chi$  of degree 2 with  $\chi|_{\mathfrak{G}'}$  rational. Now all elements of  $\mathfrak{G} - \mathfrak{G}'$  are 2-elements and a Sylow 2-subgroup of  $\mathfrak{G}$  has period 8. Thus  $Q(\chi) \subseteq Q(\varepsilon)$  where  $\varepsilon$  is a primitive 8th root of unity. Since  $8 \nmid |GF(17)^*|$ , this representation of  $\mathfrak{G}$  is realizable over  $GF(17)$  and hence we can assume  $\mathfrak{G} \subseteq GL(2, 17)$ . All subgroups of  $\mathfrak{G}$  of order 3 are contained in  $SL(2, 17)$  since  $3 \nmid |GF(17)^*|$  so  $\mathfrak{G}' \subseteq SL(2, 17)$  and  $\mathfrak{G} \subseteq SL(2, 17)^*$ . Let  $i = \sqrt{-1} \in GF(17)$  and let  $\mathfrak{Z} = \langle \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix} \rangle$ . Then  $\mathfrak{Z}$  is cyclic of order 4,  $\mathfrak{Z} \subseteq SL(2, 17)^*$  and  $\mathfrak{Z}$  is central in  $GL(2, 17)$ . Set  $\mathfrak{G} = \mathfrak{Z}\mathfrak{G}$  so  $\mathfrak{G} \subseteq SL(2, 17)^*$ .

We show first that  $\mathfrak{G}$  has precisely  $17 + 1 = 18$  noncentral involutions. Now  $|\mathfrak{Z}| = 4$  and  $\mathfrak{G}/\mathfrak{Z} \cong \text{Sym}_4$ . This quotient group has two classes of involutions  $\mathfrak{C}_1, \mathfrak{C}_2$  with  $|\mathfrak{C}_1| = 3$ ,  $|\mathfrak{C}_2| = 6$ . If  $\bar{T} \in \mathfrak{C}_i$  and the coset of  $\bar{T}$  contains an involution of  $\mathfrak{G}$ , then the same is true for all conjugates of  $\bar{T}$ . Moreover the coset would then clearly contain precisely two such involutions. Thus if  $\delta_i = 0, 1$  has the obvious meaning, then

$$I(\mathfrak{G}) = 2\delta_1 |\mathfrak{C}_1| + 2\delta_2 |\mathfrak{C}_2| = 6\delta_1 + 12\delta_2.$$

Let  $W \subseteq \mathfrak{B}$  have order 2. Then  $\bar{W} \in \mathfrak{C}_2$  so  $\delta_2 = 1$ . Let  $Q \in \mathfrak{Q}$  have order 4 and let  $\mathfrak{Z} = \langle Z \rangle$ . Then  $QZ$  has order 2 and  $\bar{QZ} \in \mathfrak{C}_1$ . Hence  $\delta_1 = 1$  and  $I(\mathfrak{G}) = 18$ .

Let  $\mathfrak{B}$  be a 2-dimensional  $GF(17)$ -vector space and let  $x \in \mathfrak{B}^\#$ . Since  $|\mathfrak{G}|$  is prime to 17 we can write  $\mathfrak{G}_x \subseteq \left\{ \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \mid a \in GF(17)^* \right\}$  by taking a suitable basis. Now  $\mathfrak{G} \subseteq SL(2, 17)^*$  and  $\det \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} = a$  so we see that  $|\mathfrak{G}_x| = 1$  or 2. If  $T$  is a noncentral involution of  $\mathfrak{G}$ , then  $\mathfrak{B} > C_{\mathfrak{B}}(T) > \{0\}$  and hence  $|C_{\mathfrak{B}}(T)^\#| = 17 - 1$ . By the above the centralizer spaces for the involutions are disjoint. Hence

$$\begin{aligned} |\bigcup_T C_{\mathfrak{B}}(T)^\#| &= I(\mathfrak{G})(17 - 1) = (17 + 1)(17 - 1) \\ &= 17^2 - 1 = |\mathfrak{B}^\#|. \end{aligned}$$

Thus  $\bigcup_T C_{\mathfrak{B}}(T) = \mathfrak{B}$  and so for all  $x \in \mathfrak{B}^\#$ ,  $|\mathfrak{G}_x| \geq 2$ . This yields  $|\mathfrak{G}_x| = 2$  and  $\mathfrak{G}$  is half-transitive but not semiregular. Finally



$\mathfrak{G} \not\subseteq \mathcal{S}(17^2)$ , the semilinear transformations, since  $\mathfrak{G}$  does not have a cyclic subgroup of index 2.

We close this section with some additional information about the degree  $17^2$  group.

LEMMA 3.5. *If  $q^n = 17^2$ , then  $|\mathfrak{G}| = 96$ .*

*Proof.* These groups occur in Lemmas 3.2 and 3.4. However the latter case was deduced from the former so we can assume  $\mathfrak{G}$  is as described in the proof of Lemma 3.2. We showed there that  $|\mathfrak{G}_x| = 2$ ,  $\mathfrak{G}/\mathbf{Z}(\mathfrak{G}) \cong \text{Sym}_4$  and  $\delta_1 = \delta_2 = 1$ . The latter says that if  $\bar{T}$  is any involution of  $\mathfrak{G}/\mathbf{Z}(\mathfrak{G})$ , then its coset contains an involution of  $\mathfrak{G}$ .

Now  $\mathfrak{X} = \mathbf{Z}(\mathfrak{G})$  has order dividing  $|GF(17)^2| = 16$ . If  $|\mathfrak{X}| = 2$ , then an involution  $T$  in the four groups of  $\text{Sym}_4$  would not have an involution of  $\mathfrak{G}$  in its coset. We assume that  $|\mathfrak{X}| \geq 8$  and derive a contradiction. Let  $T$  be an involution of  $\mathfrak{G}$  corresponding to a transposition of  $\text{Sym}_4$ . Then  $\langle \mathfrak{G}, T \rangle$  is a maximal class group of order 16 and this group has a cyclic subgroup  $\mathfrak{B}$  of order 8. We see that  $|\mathfrak{X} \cap \mathfrak{B}| = 2$  so  $|\mathfrak{X}\mathfrak{B}| = 4|\mathfrak{X}|$  and  $\mathfrak{X}\mathfrak{B}$  has period  $|\mathfrak{X}|$  since  $|\mathfrak{X}| \geq |\mathfrak{B}| = 8$ . As in the last paragraph of the proof of the preceding lemma, this implies that  $|\mathfrak{G}_x| \geq 4$ , a contradiction. Thus  $|\mathfrak{X}| = 4$  and since  $\mathfrak{G}/\mathfrak{X} \cong \text{Sym}_4$  we have  $|\mathfrak{G}| = 4 \cdot 24 = 96$ . This completes the proof of the lemma.

4. Solvable case,  $m = 2$ . In this and the next section the following assumptions hold.

ASSUMPTIONS. Group  $\mathfrak{G}$  acts faithfully on vector space  $\mathfrak{B}$  of order  $q^n$  and half-transitively but not semiregularly on  $\mathfrak{B}^*$ .  $\mathfrak{G}$  is a group of type  $E(2, m)$  with  $\mathfrak{G} \triangleleft \mathfrak{G}$ . In addition  $\mathfrak{G}$  acts irreducibly on  $\mathfrak{B}$ ,  $\mathfrak{G}$  is primitive as a linear group and  $\mathfrak{G}$  is solvable.

We will use the notation of Lemma 2.5. Moreover set  $\bar{\mathfrak{G}} = \mathfrak{G}/\mathfrak{X}\mathfrak{G}$  so that  $\bar{\mathfrak{G}}$  is a solvable subgroup of  $Sp(2m, 2)$ . We let  $\bar{\mathfrak{F}} = F(\bar{\mathfrak{G}})$ , the Fitting subgroup of  $\bar{\mathfrak{G}}$ , and for each prime  $p$  we let  $\bar{\mathfrak{F}}_p$  be the normal Sylow  $p$ -subgroup of  $\bar{\mathfrak{F}}$ . By Fitting's theorem,  $C_{\bar{\mathfrak{G}}}(\bar{\mathfrak{F}}) \subseteq \bar{\mathfrak{F}}$ . Recall the possible isomorphism classes for  $\mathfrak{G}$  namely: iso I if  $\mathfrak{G} \cong \mathfrak{D}\mathfrak{D} \cdots \mathfrak{D}$ , iso II if  $\mathfrak{G} \cong \mathfrak{D}\mathfrak{D}\mathfrak{D} \cdots \mathfrak{D}$  and iso III if  $\mathfrak{G} \cong \mathfrak{Z}\mathfrak{D}\mathfrak{D}\mathfrak{D} \cdots \mathfrak{D}$ .

LEMMA 4.1. *Suppose  $\bar{\mathfrak{F}}_2 \neq \langle 1 \rangle$ . Then  $|\bar{\mathfrak{F}}_2| = 2$ ,  $\mathfrak{G} = \text{iso I or II}$  and  $\mathfrak{G}$  has a normal subgroup  $\mathfrak{G}_0$  of type  $E(2, m - 1)$  with  $\mathfrak{G}_0 = \text{iso III}$ .*

*Proof.* Let  $\mathfrak{S}$  be the complete inverse image of  $\bar{\mathfrak{F}}_2$  in  $\mathfrak{G}$  so  $\mathfrak{S}/\mathfrak{X}\mathfrak{G} = \bar{\mathfrak{F}}_2$ . Then  $\mathfrak{S}/\mathfrak{X}$  is a 2-group and since  $\mathfrak{X}$  is central in  $\mathfrak{G}$ ,  $\mathfrak{S}$

is nilpotent. If  $\mathcal{C}_2$  is the normal Sylow 2-subgroup of  $\mathcal{C}$ , then  $\mathcal{C}_2 \cong \mathcal{C}$  and  $\mathcal{C}_2 \triangleleft \mathcal{G}$ . Since  $\mathcal{G}$  is primitive,  $\mathcal{C}_2$  is of symplectic type. Suppose  $4 \mid |\mathcal{U}_2|$ . Then since  $\mathcal{U}_2$  is central in  $\mathcal{C}_2$ ,  $\mathcal{C}_2$  has a center of order at least 4 and hence  $\mathcal{C}_2$  is the central product of  $Z(\mathcal{C}_2)$  with a number of nonabelian groups of order 8. Note that since  $\mathcal{C} \subseteq \mathcal{C}_2$ ,  $Z(\mathcal{C}_2) \subseteq C_{\mathcal{G}}(\mathcal{C}) = \mathcal{U}$  so that  $Z(\mathcal{C}_2) = \mathcal{U}_2$ . Since  $|\bar{\mathcal{F}}_2| > 1$ ,  $\mathcal{C}_2 \neq \mathcal{U}_2\mathcal{C}$  and thus  $\mathcal{C}_2 \cong \mathcal{U}_2\mathcal{C}\mathcal{B}$  where  $|\mathcal{B}| = 8$ ,  $\mathcal{B} \not\subseteq \mathcal{U}_2$  and  $\mathcal{B} \subseteq C_{\mathcal{G}}(\mathcal{C})$ , a contradiction. Thus  $|\mathcal{U}_2| = 2$  and hence  $|Z(\mathcal{C})| = 2$ . This implies that  $\dim \mathcal{B} = 2^m$  and since  $\mathcal{C}_2$  acts faithfully on  $\mathcal{B}$ ,  $\mathcal{C}_2$  has at most  $m$  nonabelian factors. Since  $|Z(\mathcal{C}_2)| = |\mathcal{U}_2| = 2$  we see that  $\mathcal{C}_2 = \mathcal{B}_0\mathcal{B}_1 \cdots \mathcal{B}_{m-1}$ , a central product of nonabelian groups with  $|\mathcal{B}_i| = 8$  if  $i > 0$  and  $\mathcal{B}_0$  a maximal class group. Now  $\mathcal{B}_0 \cap \mathcal{C}$  is a 2-generator subgroup of  $\mathcal{C}$  so  $|\mathcal{B}_0 \cap \mathcal{C}| \leq 8$ . Thus  $|\mathcal{B}_0\mathcal{C}| \geq |\mathcal{B}_0| \cdot |\mathcal{C}|/8 = |\mathcal{B}_0| \cdot 2^{2(m-1)} = |\mathcal{C}_2|$ . Hence we have equality throughout and  $|\mathcal{B}_0 \cap \mathcal{C}| = 8$ . Now  $\mathcal{B}_0 \cap \mathcal{C} \triangleleft \mathcal{B}_0$  and  $\mathcal{B}_0 \cap \mathcal{C}$  is noncyclic. As is well known this implies that  $[\mathcal{B}_0 : \mathcal{B}_0 \cap \mathcal{C}] \leq 2$  so  $|\mathcal{B}_0| \leq 16$  and  $[\mathcal{C}_2 : \mathcal{C}] \leq 2$ . If  $|\bar{\mathcal{F}}_2| \geq 1$ , then  $|\bar{\mathcal{F}}_2| = 2$ . Finally  $\mathcal{O}(\mathcal{C}_2)$  is cyclic of order 4 and from  $\mathcal{C}_2 = \mathcal{B}_0\mathcal{C}$  we see that  $\mathcal{C}_0 = C_{\mathcal{G}}(\mathcal{O}(\mathcal{C}_2))$  has the appropriate properties. Thus the result follows.

We assume throughout the remainder of this section that  $m = 2$ . Since  $\bar{\mathcal{G}} \subseteq Sp(4, 2)$  here, we make some comments about this latter group. Suppose  $Sp(4, 2)$  acts on symplectic space  $\mathcal{W}$ . If  $\mathcal{U}$  is an isotropic subspace of  $\mathcal{W}$  of dimension 2, then the symplectic form restricted to  $\mathcal{U}$  is trivial. We see easily that  $\mathcal{W}$  contains 15 such subspaces. Note that  $|Sp(4, 2)| = 2^4 \cdot 3^2 \cdot 5$ .

Let  $\bar{\mathcal{K}}$  be a Sylow 3-subgroup of  $Sp(4, 2)$ . Then  $\bar{\mathcal{K}}$  is abelian of type (3, 3) and contains the four subgroups  $\bar{\mathcal{K}}_1, \bar{\mathcal{K}}_2, \bar{\mathcal{K}}_3, \bar{\mathcal{K}}_4$  of order 3. We can take (see [10]) the following concrete realization for  $\bar{\mathcal{K}}$ . Write  $\mathcal{W} = \mathcal{W}_1 \oplus \mathcal{W}_2$ , a direct sum of two nonisotropic 2-dimensional subspaces and then let  $\bar{\mathcal{K}}_1$  centralize  $\mathcal{W}_2$  and act irreducibly on  $\mathcal{W}_1$  and  $\bar{\mathcal{K}}_2$  centralize  $\mathcal{W}_1$  and act irreducibly on  $\mathcal{W}_2$ .

Let  $\bar{\mathcal{L}} = \bar{\mathcal{K}}_1$  or  $\bar{\mathcal{K}}_2$ . Then  $\mathcal{W} = C_{\mathcal{W}}(\bar{\mathcal{L}}) \oplus (\mathcal{W}, \bar{\mathcal{L}})$  a direct sum of 2-dimensional subspaces. Let  $\mathcal{U}$  be a 2-dimensional  $\bar{\mathcal{L}}$ -subspace of  $\mathcal{W}$ . If  $\mathcal{U} \cap (\mathcal{W}, \bar{\mathcal{L}}) = \{0\}$ , then certainly  $\mathcal{U} \subseteq C_{\mathcal{W}}(\bar{\mathcal{L}})$  so  $\mathcal{U} = C_{\mathcal{W}}(\bar{\mathcal{L}})$ . If  $\mathcal{U} \cap (\mathcal{W}, \bar{\mathcal{L}}) \neq \{0\}$ , then since  $\bar{\mathcal{L}}$  acts irreducibly on  $(\mathcal{W}, \bar{\mathcal{L}})$  we have  $\mathcal{U} \cong (\mathcal{W}, \bar{\mathcal{L}})$  so  $\mathcal{U} = (\mathcal{W}, \bar{\mathcal{L}})$ . Thus  $\mathcal{U} = \mathcal{W}_1$  or  $\mathcal{W}_2$ . In particular  $\bar{\mathcal{K}}_1$  and  $\bar{\mathcal{K}}_2$  do not normalize a 2-dimensional isotropic subspace of  $\mathcal{W}$ . If  $\mathcal{U}$  is a 1-dimensional  $\bar{\mathcal{L}}$ -subspace, then certainly  $\mathcal{U} \subseteq C_{\mathcal{W}}(\bar{\mathcal{L}})$  so  $\mathcal{U} \subseteq \mathcal{W}_1$  or  $\mathcal{W}_2$ .

Now let  $\bar{\mathcal{L}} = \bar{\mathcal{K}}_3$  or  $\bar{\mathcal{K}}_4$ . Then  $\bar{\mathcal{L}}$  acts irreducibly on both  $\mathcal{W}_1$  and  $\mathcal{W}_2$  so  $\bar{\mathcal{L}}$  has no 1-dimensional invariant subspace. Let  $\mathcal{U}$  be a 2-dimensional  $\bar{\mathcal{L}}$ -invariant subspace. If  $\mathcal{U} = \mathcal{W}_1$  or  $\mathcal{W}_2$ , then  $\mathcal{U}$  is nonisotropic.

Suppose  $\mathfrak{U} \neq \mathfrak{B}_1$  or  $\mathfrak{B}_2$  and  $w_1 + w_2 \in \mathfrak{U}$  with  $w_i \in \mathfrak{B}_i$ . Clearly  $w_1, w_2 \neq 0$ . It is now easy to see that we get precisely three subspaces  $\mathfrak{U}$  and since  $\mathfrak{B}_1$  and  $\mathfrak{B}_2$  are orthogonal each such  $\mathfrak{U}$  is isotropic. Thus  $\bar{\mathfrak{G}}$  normalizes two nonisotropic 2-dimensional subspaces and three isotropic ones.

If  $\bar{\mathfrak{F}}$  is a subgroup of  $Sp(4, 2)$  of order 5, then  $\bar{\mathfrak{F}}$  acts irreducibly on  $\mathfrak{B}$ . Then  $|C(\bar{\mathfrak{F}})| \mid 2^4 - 1$  so  $|C(\bar{\mathfrak{F}})| = 5$  or 15. In the latter case let  $\bar{\mathfrak{E}}$  be a subgroup of order 3 centralizing  $\bar{\mathfrak{F}}$ . Then  $\bar{\mathfrak{F}}$  permutes the two 2-dimensional nonisotropic subspaces normalized by  $\bar{\mathfrak{E}}$  and hence  $\bar{\mathfrak{F}}$  normalizes each, a contradiction. Thus  $Sp(4, 2)$  has no elements of order 10 or 15.

LEMMA 4.2.  $\bar{\mathfrak{F}}$  is a normal 2-complement of  $\bar{\mathfrak{G}}$  and  $|\bar{\mathfrak{F}}| = 3, 5$  or  $\bar{\mathfrak{F}}$  is abelian of type (3,3).

*Proof.* Suppose first that  $\bar{\mathfrak{F}}_2 \neq \langle 1 \rangle$ . By Lemma 4.1,  $\mathfrak{G}$  has a normal subgroup  $\mathfrak{C}_0 \cong \mathfrak{B}\mathfrak{D}$  and moreover  $4 \nmid |Z(\mathfrak{G})|$ . By the Reduction Lemma and Lemma 3.4 we have  $q = 3, 5$  or 17. Suppose  $q = 3$ . Since  $|Z(\mathfrak{G})| = 2$  and  $\mathfrak{G}$  acts irreducibly,  $q^n = 3^4$  and thus  $\mathfrak{C}_0$  also acts irreducibly. By Lemma 4.1  $\mathfrak{G}$  is imprimitive, a contradiction. Let  $q = 5$  or 17. Then  $4 \mid q - 1$  and since  $\mathfrak{G}$  is primitive and  $Z(\mathfrak{C}_0) \triangle \mathfrak{G}$  with  $|Z(\mathfrak{C}_0)| = 4$  we conclude that  $Z(\mathfrak{C}_0)$  consists of scalar matrices and  $4 \mid |Z(\mathfrak{G})|$ , a contradiction.

Now suppose  $\bar{\mathfrak{F}} = \langle 1 \rangle$ . Then  $\bar{\mathfrak{G}} = \langle 1 \rangle$ . If  $Z(\mathfrak{G})$  is central then  $\mathfrak{G} = \mathfrak{A}\mathfrak{G}$  is nilpotent so  $\mathfrak{G}_2 \cong \mathfrak{G}$  is half-transitive. By Theorem II of [4],  $\mathfrak{G}_2 \cong \mathfrak{D}\mathfrak{D}$  and  $q^n = 3^4$ . Then  $|\mathfrak{A}| \mid q - 1$  so  $\mathfrak{G} = \mathfrak{G}_2 \cong \mathfrak{D}\mathfrak{D}$  and this group is imprimitive, a contradiction. Thus  $Z(\mathfrak{G})$  is not central and in particular  $|Z(\mathfrak{G})| = 4$ . Since  $|\mathfrak{G}/\bar{\mathfrak{G}}| = 2$  we see that  $\mathfrak{G}$  normalizes a hyperplane in  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$ , say  $\mathfrak{B}_0 = \mathfrak{C}_0/Z(\mathfrak{G})$ . Then  $\mathfrak{C}_0 \triangle \mathfrak{G}$  and  $\mathfrak{C}_0$  has period 4. Since  $\mathfrak{G}$  is primitive  $Z(\mathfrak{C}_0)$  is cyclic so  $Z(\mathfrak{C}_0) = Z(\mathfrak{G})$  and then  $\mathfrak{C}_0/Z(\mathfrak{C}_0)$  has odd dimension, a contradiction.

Using the fact that  $Sp(4, 2)$  has no elements of order 15 we conclude that  $\bar{\mathfrak{F}}$  is one of the three possibilities mentioned in the statement of the lemma. Since  $\bar{\mathfrak{F}}$  is abelian,  $\bar{\mathfrak{G}}/\bar{\mathfrak{F}} \cong \text{Aut } \bar{\mathfrak{F}}$  and from this we see easily that  $\bar{\mathfrak{F}}$  is a normal 2-complement.

LEMMA 4.3.  $\mathfrak{G} = \text{iso I}$  does not occur.

*Proof.* Suppose  $\mathfrak{G} \cong \mathfrak{D}\mathfrak{D} \cong \mathfrak{D}\mathfrak{D}$ . Then  $\bar{\mathfrak{G}} = \mathfrak{G}$  and  $\bar{\mathfrak{G}}$  permutes the involution vectors of  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$ . By Lemma 1.3,  $i(\mathfrak{B}) = 9$  and this clearly implies that  $|\bar{\mathfrak{F}}| \neq 5$ . Thus  $\bar{\mathfrak{F}}$  is abelian of type (3) or (3, 3). Since  $\mathfrak{G} \cong \mathfrak{D}\mathfrak{D}$  we see easily that  $\mathfrak{G}$  contains an abelian subgroup  $\mathfrak{B}$  of type (2, 2, 2). If  $\mathfrak{B}_1$  is an irreducible  $\mathfrak{B}$ -submodule of  $\mathfrak{B}$  then by Schur's lemma,  $[\mathfrak{B} : C_{\mathfrak{B}}(\mathfrak{B}_1)] \leq 2$  so for  $x \in \mathfrak{B}_1^*$ ,  $4 \mid |\mathfrak{B}_x|$  and hence  $4 \mid |\mathfrak{G}_x|$ . Moreover since  $\mathfrak{G}_x$  is abelian and  $\mathfrak{G}_x \cap Z(\mathfrak{G}) = \langle 1 \rangle$  we

see easily that  $\mathbb{G}_x = \mathbb{B}_x$ . Suppose  $|\bar{\mathfrak{F}}| = 3$ . By Lemma 1.5 there exists  $y \in \mathbb{B}^*$  with  $\mathbb{G}_y \cap \mathfrak{A}\mathbb{G} = \langle 1 \rangle$ . Since  $\mathbb{G} = \mathfrak{H}$  and  $|\bar{\mathfrak{H}}| = 3$  or  $6$  by Fitting's theorem, we have  $|\mathbb{G}_y| \mid 6$ , a contradiction. Thus  $\bar{\mathfrak{F}}$  is abelian of type  $(3, 3)$ .

First suppose  $q = 3$ . Then a Sylow 3-subgroup of  $\mathbb{G}$  has a fixed point in  $\mathbb{B}^*$  and thus by half-transitivity  $\mathbb{G}_x \cong \mathfrak{R}$  where  $x$  is the above mentioned point and  $\mathfrak{R}$  is a Sylow 3-subgroup of  $\mathbb{G}$ . Note that if  $\bar{\mathfrak{R}}$  is the image of  $\mathfrak{R}$  in  $\bar{\mathfrak{H}}$  then  $\bar{\mathfrak{R}} = \bar{\mathfrak{F}}$ . Since  $\mathbb{G}_x = \mathbb{G} \cap \mathbb{G}_x \triangle \mathbb{G}_x$  we see that  $\bar{\mathfrak{R}}$  normalizes  $Z(\mathbb{G})\mathbb{G}_x/Z(\mathbb{G}) = \mathbb{B}/Z(\mathbb{G})$  a 2-dimensional isotropic subspace of symplectic space  $\mathbb{B}$ . This contradicts our preceding remarks about  $Sp(4, 2)$  since the subgroup  $\bar{\mathfrak{L}}_1$  of  $\bar{\mathfrak{R}}$  normalizes no such subspaces. Thus  $q \neq 3$ .

Now  $\bar{\mathfrak{F}}$  acts on  $\mathbb{B} = \mathbb{G}/Z(\mathbb{G})$  and let  $\mathbb{B} = \mathbb{B}_1 \oplus \mathbb{B}_2$  be the decomposition of  $\mathbb{B}$  given in our earlier discussion of  $Sp(4, 2)$ . If  $Z(\mathbb{G}) \subseteq \mathbb{G}_i \subseteq \mathbb{G}$  with  $\mathbb{G}_i/Z(\mathbb{G}) = \mathbb{B}_i$ , then  $\mathbb{G}_i$  is nonabelian since  $\mathbb{B}_i$  is nonisotropic, and since  $\mathbb{G}_i$  admits an automorphism of order 3 we have  $\mathbb{G}_i \cong \Omega$ . Hence we can find a noncentral involution  $T \in \mathbb{G} - (\mathbb{G}_1 \cup \mathbb{G}_2)$ . By Lemma 1.5 there exists  $x \in \mathbb{B}^*$  with  $\mathbb{G}_x = \langle T \rangle$ . Now a Sylow 3-subgroup of  $\mathbb{G}$  is not cyclic, since  $\bar{\mathfrak{F}}$  is not cyclic and hence it cannot act semiregularly. By half-transitivity  $\mathbb{G}_x$  contains a subgroup  $\mathfrak{L}$  of order 3. Then  $\mathfrak{L} \cap \mathfrak{A}\mathbb{G} = \langle 1 \rangle$  so if  $\bar{\mathfrak{L}}$  denotes the image of  $\mathfrak{L}$  in  $\bar{\mathfrak{H}}$ , then  $|\bar{\mathfrak{L}}| = 3$ . Since  $\langle T \rangle = \mathbb{G}_x = \mathbb{G} \cap \mathbb{G}_x \triangle \mathbb{G}_x$  we see that  $\bar{\mathfrak{L}}$  normalizes the 1-dimensional subspace  $\mathbb{G}_x Z(\mathbb{G})/Z(\mathbb{G}) = \mathfrak{U}$ . Now  $T$  was chosen in such a way that  $\mathfrak{U} \not\subseteq \mathbb{B}_1$  or  $\mathbb{B}_2$ . Hence in the notation of our discussion of  $Sp(4, 2)$  we see that  $\bar{\mathfrak{L}} \neq \bar{\mathfrak{L}}_1$  or  $\bar{\mathfrak{L}}_2$ . On the other hand  $\bar{\mathfrak{L}}_3$  and  $\bar{\mathfrak{L}}_4$  do not normalize 1-dimensional subspaces. Hence  $\bar{\mathfrak{L}} \neq \bar{\mathfrak{L}}_1, \bar{\mathfrak{L}}_2, \bar{\mathfrak{L}}_3$  or  $\bar{\mathfrak{L}}_4$ , a contradiction.

LEMMA 4.4. *If  $\mathbb{G} = \text{iso II}$ , then  $q^n = 3^4$ .*

*Proof.* Let us assume that  $q^n \neq 3^4$ . Since  $\mathbb{G}$  acts irreducibly on  $\mathbb{B}$  we have  $|\mathbb{B}| = q^n = q^4$  so  $q \geq 5$ . We consider the possibilities for  $\bar{\mathfrak{F}}$ . Suppose  $\bar{\mathfrak{F}}$  is abelian of type  $(3, 3)$ . Then  $\bar{\mathfrak{F}}$  is a Sylow 3-subgroup of  $Sp(4, 2)$  and we can write  $\mathbb{B} = \mathbb{B}_1 \oplus \mathbb{B}_2$ , the corresponding decomposition of  $\mathbb{G}/Z(\mathbb{G}) = \mathbb{B}$ . If  $\mathbb{G}_i/Z(\mathbb{G}) = \mathbb{B}_i$ , then since  $\mathbb{B}_i$  is nonisotropic,  $\mathbb{G}_i$  is nonabelian of order 8. Now  $\mathbb{G}_i$  admits an automorphism of order 3 so  $\mathbb{G}_i \cong \Omega$  and  $\mathbb{G} \cong \Omega\Omega$ , a contradiction. Thus  $|\bar{\mathfrak{F}}| = p$  for  $p = 3$  or  $5$ .

Note that  $\mathfrak{H} = \mathbb{G}$  and  $|\bar{\mathfrak{H}}/\bar{\mathfrak{F}}| \mid (p - 1)$ . Thus  $\bar{\mathfrak{H}}/\bar{\mathfrak{F}}$  is a cyclic 2-group. Suppose  $p \mid |\mathbb{G}_x|$  for all  $x \in \mathbb{B}^*$ . Let  $T$  be a noncentral involution of  $\mathbb{G}$ . Since  $q \neq 3$  there exists by Lemma 1.5 an  $x \in \mathbb{B}^*$  with  $\mathbb{G}_x = \langle T \rangle$ . Let  $\mathfrak{L}$  be a subgroup of  $\mathbb{G}_x$  of order  $p$ . Since

$\mathfrak{B} \cap \mathfrak{A}\mathfrak{C} = \langle 1 \rangle$ ,  $\bar{\mathfrak{B}}$ , the image of  $\mathfrak{B}$  in  $\bar{\mathfrak{G}}$ , has order  $p$  so  $\bar{\mathfrak{B}} = \bar{\mathfrak{F}}$ . Since  $\langle T \rangle = \mathfrak{C}_x = \mathfrak{G}_x \cap \mathfrak{C}$  we see that  $\bar{\mathfrak{F}}$  centralizes the involution vector in  $\mathfrak{B}$  corresponding to  $T$ . By Lemma 1.2,  $\bar{\mathfrak{F}}$  centralizes  $\mathfrak{B}$ , a contradiction. Thus  $p \nmid |\mathfrak{G}_x|$  and in particular  $p \neq q$ .

Suppose  $p = 3$ . By Lemma 1.5 there exists  $x \in \mathfrak{B}^*$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{C} = \langle 1 \rangle$ . Hence  $|\mathfrak{G}_x| \mid |\bar{\mathfrak{G}}|$ . Since  $|\bar{\mathfrak{G}}| = 6$  we conclude that  $|\mathfrak{G}_x| = 2$ . We note now that  $4 \nmid |\mathfrak{A}|$ . Otherwise  $\mathfrak{A}\mathfrak{C}$  contains  $\mathfrak{C}^* \cong \mathfrak{B}\mathfrak{D}\mathfrak{D}$  and this group contains an abelian subgroup of type  $(2, 2, 2)$ . This easily implies that  $4 \mid |\mathfrak{G}_x|$ , a contradiction. Let  $\mathfrak{B}$  be a Sylow 3-subgroup of  $\mathfrak{G}$ . Since  $\bar{\mathfrak{B}}/\bar{\mathfrak{F}}$  acts faithfully on  $\bar{\mathfrak{F}}$  we see by the above that if  $T$  is a noncentral involution of  $\mathfrak{G}$  and  $T \subseteq C_{\bar{\mathfrak{B}}}(\bar{\mathfrak{F}})$  then  $T \in \mathfrak{C}$ . Now  $\bar{\mathfrak{B}} = \bar{\mathfrak{F}}$  permutes faithfully the  $i(\mathfrak{B}) = 5$  involution vectors of  $\mathfrak{B}$ . Thus  $\bar{\mathfrak{B}}$  moves 3 such and fixes 2 such. Since each involution vector corresponds to two noncentral involutions of  $\mathfrak{C}$  we see that  $\mathfrak{B}$  centralizes precisely four noncentral involutions of  $\mathfrak{G}$ . Thus clearly  $I(\mathfrak{G}) \equiv 4 \pmod 3$ . On the other hand by Lemma 1.9 we have  $I(\mathfrak{G}) = 1 + q^2$ . Thus  $q^2 \equiv 0 \pmod 3$ , a contradiction since  $q \neq 3$ .

We consider  $p = 5$  so  $q \geq 7$ . Let  $\bar{I}$  denote the number of involutions of  $\mathfrak{G}/\mathfrak{A}$ . Since  $\mathfrak{A}$  is cyclic and central in  $\mathfrak{G}$ , each involution of  $\mathfrak{G}/\mathfrak{A}$  corresponds to at most two noncentral involutions of  $\mathfrak{G}$  so  $I(\mathfrak{G}) \leq 2\bar{I}$ . Now  $\mathfrak{B}\mathfrak{F} \trianglelefteq \mathfrak{G}/\mathfrak{A}$  where  $\mathfrak{B}$  is elementary abelian of order  $2^4$ ,  $|\mathfrak{F}| = 5$  and  $\mathfrak{F}$  acts irreducibly on  $\mathfrak{B}$ . Furthermore  $(\mathfrak{G}/\mathfrak{A})/(\mathfrak{B}\mathfrak{F})$  is a cyclic 2-group which acts faithfully on  $(\mathfrak{B}\mathfrak{F})/\mathfrak{B}$ . Hence we see easily that  $\bar{I} \leq 15 + 5 \cdot 4 = 35$  and  $I(\mathfrak{G}) \leq 70$ .

Let  $T$  be a noncentral involution of  $\mathfrak{G}$ . If  $T \in \mathfrak{A}\mathfrak{C}$  then certainly  $|C_{\mathfrak{B}}(T)| = q^2$ . Suppose  $T \notin \mathfrak{A}\mathfrak{C}$ . From the structure of  $\bar{\mathfrak{G}}$  we see that for some  $F \in \mathfrak{G}$ ,  $\langle T, T^F \rangle \cong \bar{\mathfrak{F}}$ . Since  $5 \nmid |\mathfrak{G}_x|$  we see that  $C_{\mathfrak{B}}(T) \cap C_{\mathfrak{B}}(T^F) = \{0\}$ . Hence  $|C_{\mathfrak{B}}(T)| \leq q^2$  here also. Now every element of  $\mathfrak{B}^*$  is fixed by some noncentral involution of  $\mathfrak{G}$  so  $\mathfrak{B}^* = \bigcup_T C_{\mathfrak{B}}(T)^*$  and hence

$$q^4 - 1 = |\mathfrak{B}^*| \leq I(\mathfrak{G})(q^2 - 1)$$

or  $q^2 + 1 \leq I(\mathfrak{G}) \leq 70$ . Since  $q > 5$ , we have  $q = 7$ .

For  $q = 7$  the argument is somewhat involved. Since  $|\mathfrak{A}| \mid q - 1$  we have  $|\mathfrak{A}| = 2$  or  $6$ . Now  $O_3(\mathfrak{A})$  is central in  $\mathfrak{G}$  and is a Sylow 3-subgroup of  $\mathfrak{G}$ . Thus  $\mathfrak{G}$  has a normal 3-complement. Since this group is also half-transitive we see that it suffices to assume that  $O_3(\mathfrak{A}) = \langle 1 \rangle$  and hence  $|\mathfrak{A}| = 2$ ,  $\mathfrak{A}\mathfrak{C} = \mathfrak{C}$ .

We can now get a tighter count on  $I(\mathfrak{G})$ . Let  $\bar{I} = \bar{I}_1 + \bar{I}_2$  where  $\bar{I}_1$  counts the number of involutions of  $\mathfrak{G}/\mathfrak{A}$  and  $\bar{I}_2$  counts those of  $\mathfrak{G}/\mathfrak{A}$  not in  $\mathfrak{C}/\mathfrak{A}$ . We have as before  $\bar{I}_1 = 15$ ,  $\bar{I}_2 \leq 20$ . If  $I(\mathfrak{G}) = I_1 + I_2$  is the corresponding break up of  $I(\mathfrak{G})$ , then  $I_2 \leq 2\bar{I}_2 \leq 40$  and

$I_1 = I(\mathfrak{G}) = 10$ . Hence  $I(\mathfrak{G}) \leq 50$  here. As above  $\mathfrak{B}^\sharp = \bigcup_T C_{\mathfrak{B}}(T)$  yields  $50 = q^2 + 1 \leq I(\mathfrak{G}) \leq 50$ . Thus we must have equality throughout and hence  $\bigcup_T C_{\mathfrak{B}}(T)$  is a disjoint union. This implies that every element  $x \in \mathfrak{B}^\sharp$  is centralized by precisely one involution so  $\mathfrak{G}_x$  has a unique involution.

Let  $\mathfrak{R}$  be the subgroup of  $\mathfrak{G}$  with  $\mathfrak{R} \supseteq \mathfrak{G}$  and  $[\bar{\mathfrak{R}} : \bar{\mathfrak{G}}] = 2$ . Since  $\bar{\mathfrak{G}}/\bar{\mathfrak{G}}$  is cyclic,  $\mathfrak{R}$  contains all the involutions of  $\mathfrak{G}$ . We study the group  $\mathfrak{R}$ . Note that  $\mathfrak{R}$  is dihedral of order 10 and  $\bar{\mathfrak{G}}$  acts irreducibly on  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$ . Let  $\mathfrak{Q}$  be a Sylow 5-subgroup of  $\mathfrak{R}$  so that  $|\mathfrak{Q}| = 5$  and let  $\mathfrak{R} = N_{\mathfrak{R}}(\mathfrak{Q})$ . From the above we see that  $\mathfrak{R}/Z(\mathfrak{G})$  is dihedral of order 10. Let  $\mathfrak{Q} = \langle L \rangle$  and let  $N \in \mathfrak{R} - Z(\mathfrak{G})$  be a 2-element. Then  $L^N = L^{-1}$ .

Now  $\mathfrak{R}$  permutes the 10 noncentral involutions of  $\mathfrak{G}$  and the corresponding five involution vectors of  $\mathfrak{B}$ . Using  $(( \ ))$  to denote cyclic permutations, it is clear that we can label the involutions by  $X_i, Y_i, i = 1, 2, \dots, 5$  such that  $Y_i = -X_i$  and as a permutation

$$L = ((X_1, X_2, X_3, X_4, X_5))((Y_1, Y_2, Y_3, Y_4, Y_5)) .$$

Here for convenience we denoted the central involution of  $\mathfrak{G}$  by  $-1$ . We consider  $N$ . As a permutation, it has order 2. Since  $N$  acts on the five involution vectors of  $\mathfrak{B}$ ,  $N$  must fix at least one such, say the one corresponding to  $\{X_1, Y_1\}$ . Then either  $N$  fixes both  $X_1$  and  $Y_1$  or  $N$  interchanges the two. Since  $L^N = L^{-1}$  this completely determines the cycle structure of  $N$  and we have either

- (a)  $N = ((X_1))((X_2, X_5))((X_3, X_4))((Y_1))((Y_2, Y_5))((Y_3, Y_4))$  or
- (b)  $N = ((X_1, Y_1))((X_2, Y_5))((X_3, Y_4))((X_4, Y_3))((X_5, Y_2))$  .

Note that it is easy to see that for  $i \neq j, (X_i, X_j) = (Y_i, Y_j) = -1$ . Now the sum of the five involution vectors of  $\mathfrak{B}$  is  $L$  invariant and hence must be 0. Thus  $Z = X_1 X_2 X_3 X_4 X_5 \in Z(\mathfrak{G})$ . If  $N$  acts like (b) above, then

$$\begin{aligned} Z &= Z^N = (X_1 X_2 X_3 X_4 X_5)^N = Y_1 Y_5 Y_4 Y_3 Y_2 \\ &= -X_1 (X_5 X_4 X_3 X_2) = -Z^{-1} . \end{aligned}$$

Thus  $Z^2 = -1$ , a contradiction and hence  $N$  must act like (a) above.

Suppose  $N$  has order 2. Then  $\langle N, X_1, Y_1 \rangle$  is elementary abelian of order 8. This yields as usual an element  $x \in \mathfrak{B}^\sharp$  such that  $\mathfrak{G}_x$  contains a subgroup of type  $(2, 2)$  and this contradicts our preceding remarks. Hence  $N^2 = -1$ .

Now  $\mathfrak{S} = \langle \mathfrak{G}, N \rangle$  is a Sylow 2-subgroup of  $\mathfrak{R}$ . We show that every involution of  $\mathfrak{S}$  is contained in  $\mathfrak{G}$ . This will imply that  $\mathfrak{G}$  contains only 10 noncentral involutions and this will yield the required contradiction. Suppose  $T \in \mathfrak{S} - \mathfrak{G}$  is an involution. Then  $T = NE$  for some  $E \in \mathfrak{G}$ . Since  $N^2 = -1$  we have

$$1 = T^2 = NENE = -E^N E$$

so  $E^N = -E^{-1}$ . In particular the image of  $E$  in  $\mathfrak{B} = \mathfrak{G}/Z(\mathfrak{G})$  is centralized by  $N$ . Now  $C_{\mathfrak{B}}(N)$  is a 2-dimensional subspace which is clearly spanned by the images in  $\mathfrak{B}$  of  $X_1$  and  $X_2X_5$ . Note that  $X_1$  and  $X_2X_5$  commute and  $X_2X_5$  has order 4. Hence  $E \in \langle X_1, X_2X_5 \rangle = \mathfrak{B}$ . We have  $X_1^N = X_1 = X_1^{-1}$  and  $(X_2X_5)^N = X_5X_2 = (X_2X_5)^{-1}$  so since  $\mathfrak{B}$  is abelian,  $N$  acts in a dihedral manner on  $\mathfrak{B}$ . Thus  $E^N = E^{-1}$  which contradicts the previous relation  $E^N = -E^{-1}$ . This implies that  $T$  does not exist and the proof is complete.

If  $q^n = 3^4$  above then  $\mathfrak{G} = F(\mathfrak{G})$  is half-transitive. Thus these groups are given in [5] where uniqueness was proved. Since  $\mathfrak{G}$  is primitive, we see that  $\mathfrak{G}$  is transitive and hence it is one of the groups given in [3].

LEMMA 4.5.  $\mathfrak{G} = \text{iso III}$  does not occur.

*Proof.* Suppose  $\mathfrak{G} \cong 3\Omega\Omega$ . Since  $|Z(\mathfrak{G})| = 4$  and  $\mathfrak{G}$  acts irreducibly we see that  $|\mathfrak{B}| = q^4$  if  $q \equiv 1 \pmod 4$  and  $|\mathfrak{B}| = q^8$  if  $q \equiv -1 \pmod 4$ . If  $\mathfrak{H} = C_{\mathfrak{B}}(Z(\mathfrak{G}))$ , then  $[\mathfrak{G} : \mathfrak{H}] = 1$  or  $2$ . Moreover if  $[\mathfrak{G} : \mathfrak{H}] = 2$  then  $q \equiv -1$ .

We consider  $\bar{\mathfrak{H}}$ . Suppose  $|\bar{\mathfrak{H}}| = 5$  or  $9$  so that  $C_{\mathfrak{B}}(\bar{\mathfrak{H}}) = \langle 1 \rangle$ . Clearly  $\bar{\mathfrak{H}}$  acts faithfully on  $\mathfrak{G}/\mathfrak{G}'$  and centralizes  $Z(\mathfrak{G})/\mathfrak{G}'$ . Let  $\mathfrak{G}_0$  be the commutator subgroup of  $\mathfrak{G}\bar{\mathfrak{H}}$ . Then clearly  $|\mathfrak{G}_0/\mathfrak{G}'| = 2^4$ ,  $\mathfrak{G}_0 \triangle \mathfrak{G}$  and  $\mathfrak{G}_0 = \text{iso I}$  or  $\text{II}$ . By the Reduction Lemma and the previous two lemmas,  $\mathfrak{G}_0 = \text{iso II}$  and  $q = 3$ . Since as we have seen, this group does not admit an automorphism group of type  $(3, 3)$  we must have  $|\bar{\mathfrak{H}}| = 5$ . Since  $q = 3$ ,  $q^n = 3^8$ .

Now  $\mathfrak{G}$  has an abelian subgroup of type  $(2, 2, 2)$  so it follows that  $4 \mid |\mathfrak{G}_x|$  and hence  $2 \mid |\mathfrak{H}_x|$  for all  $x \in \mathfrak{B}^*$ . As in the proof of the previous lemma we see that  $5 \nmid |\mathfrak{G}_x|$  and hence if  $T$  is a noncentral involution of  $\mathfrak{H}$ , then  $|C_{\mathfrak{B}}(T)| \leq 3^4$ . Now  $\mathfrak{H}/\mathfrak{A}$  contains at most  $15 + 5 \cdot 4 = 35$  involutions and hence since  $\mathfrak{A}$  is central and cyclic we have  $I(\mathfrak{H}) \leq 2 \cdot 35 = 70$ . Since  $\mathfrak{B} = \bigcup_T C_{\mathfrak{B}}(T)$  we have

$$3^8 = |\mathfrak{B}| \leq 3^4 I(\mathfrak{H}) \leq 3^4 \cdot 70$$

or  $3^4 \leq 70$ , a contradiction.

Finally let  $|\bar{\mathfrak{H}}| = 3$ . As above we see that  $4 \mid |\mathfrak{G}_x|$ . Since by Lemma 1.5 there exists  $x \in \mathfrak{B}^*$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$ , we conclude that  $4 \mid |\mathfrak{G}/\mathfrak{A}\mathfrak{G}|$ . Hence  $|\bar{\mathfrak{H}}| = 6$  and  $[\mathfrak{G} : \mathfrak{H}] = 2$  so  $q \equiv -1 \pmod 4$ ,  $q^n = q^8$  and  $q \neq 5$ . By Lemma 1.5, if  $T$  is a noncentral involution of  $\mathfrak{G}$  then for some  $x \in \mathfrak{B}^*$ ,  $\mathfrak{G}_x = \langle T \rangle$ . Hence if  $3 \mid |\mathfrak{G}_x|$ , then  $\bar{\mathfrak{H}}$  fixes all involution vectors of  $\mathfrak{B}$  and  $\bar{\mathfrak{H}}$  centralizes  $\mathfrak{B}$ , a contradiction. Thus

$3 \nmid |\mathbb{G}_x|$  and this implies easily that if  $T$  is an involution of  $\mathfrak{G}$ , then  $|C_{\mathfrak{G}}(T)| \leq q^4$ . Also  $q \neq 3$  so  $q \geq 7$ . We have clearly  $I(\mathfrak{G}) \leq 2 \cdot 2 \cdot 16 \cdot 3 = 192$  and since  $\mathfrak{B} = \bigcup_r C_{\mathfrak{G}}(T)$  we have

$$q^8 = |\mathfrak{B}| \leq q^4 I(\mathfrak{G}) \leq 192q^4.$$

Thus  $7^4 \leq q^4 \leq 192$ , a contradiction. This completes the proof of the lemma.

5. Solvable case,  $m = 3$  and 4. We continue with the assumptions of the preceding section except that  $m = 3$  or 4 here. First let  $m = 3$ . Now  $|Sp(2m, 2)| = 2^9 \cdot 3^4 \cdot 5 \cdot 7$ . We consider the possibilities for  $\bar{\mathfrak{G}}$ .

LEMMA 5.1.  $\bar{\mathfrak{G}}$  is a 3-group.

*Proof.* If  $p$  is a prime, we let  $\bar{\mathfrak{G}}_p$  denote the normal Sylow  $p$ -subgroup of  $\bar{\mathfrak{G}}$ . We show here that  $\bar{\mathfrak{G}}_2 = \bar{\mathfrak{G}}_5 = \bar{\mathfrak{G}}_7 = \langle 1 \rangle$ .

Suppose  $\bar{\mathfrak{G}}_2 \neq \langle 1 \rangle$ . By Lemma 4.1  $\mathfrak{G}$  has a normal subgroup  $\mathfrak{G}_0 \cong 3\mathcal{D}\mathcal{D}$ . By the Reduction Lemma and Lemma 4.5 this does not occur.

Suppose  $\bar{\mathfrak{G}}_7 \neq \langle 1 \rangle$ . Then  $|\bar{\mathfrak{G}}_7| = 7$  and  $\bar{\mathfrak{G}}_7$  acts irreducibly on  $\mathfrak{B}$ . By Schur's lemma,  $C_{\bar{\mathfrak{G}}}(\bar{\mathfrak{G}}_7)$  is a cyclic group of odd order and  $[\bar{\mathfrak{G}} : C_{\bar{\mathfrak{G}}}(\bar{\mathfrak{G}}_7)] \mid 6$ . Hence if  $\mathfrak{G} = \text{iso I or II}$  then  $4 \nmid [\mathfrak{G} : \mathfrak{A}\mathfrak{G}]$  while if  $\mathfrak{G} = \text{iso III}$ , then  $8 \nmid [\mathfrak{G} : \mathfrak{A}\mathfrak{G}]$ . Now if  $\mathfrak{G} = \text{iso I or II}$  then  $\mathfrak{G}$  has an abelian subgroup of type  $(2, 2, 2)$  so for some  $y \in \mathfrak{B}^*$ ,  $4 \mid |\mathfrak{G}_y|$ . If  $\mathfrak{G} = \text{iso III}$ , then  $\mathfrak{G}$  has an abelian subgroup of type  $(2, 2, 2, 2)$  so  $8 \mid |\mathfrak{G}_y|$ . Finally by Lemma 1.5 there exists  $x \in \mathfrak{B}^*$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$  so  $|\mathfrak{G}_x| \mid [\mathfrak{G} : \mathfrak{A}\mathfrak{G}]$ . Since  $|\mathfrak{G}_x| = |\mathfrak{G}_y|$  we have a contradiction.

Suppose  $\bar{\mathfrak{G}}_5 \neq \langle 1 \rangle$ . Then  $|\bar{\mathfrak{G}}_5| = 5$  and we can write  $\mathfrak{B} = \mathfrak{B}_1 \oplus \mathfrak{B}_2$  where  $|\mathfrak{B}_1| = 2^2$ ,  $|\mathfrak{B}_2| = 2^4$ , both these spaces are  $\bar{\mathfrak{G}}_5$  invariant and  $\mathfrak{B}_1 = C_{\mathfrak{B}}(\bar{\mathfrak{G}}_5)$ . Let  $\mathfrak{G} \supseteq \mathfrak{G}_i \supseteq Z(\mathfrak{G})$  with  $\mathfrak{G}_i/Z(\mathfrak{G}) = \mathfrak{B}_i$ . Clearly  $\mathfrak{G}_i \triangleleft \mathfrak{G}$  and since  $\mathfrak{G}$  is primitive each  $\mathfrak{G}_i$  is of symplectic type. By the Reduction Lemma applied to  $\mathfrak{G}_2$  and Lemmas 4.3, 4.4 and 4.5 we have  $q = 3$  and  $\mathfrak{G}_2 \cong \mathcal{D}\mathcal{D}$ . Hence  $|Z(\mathfrak{G})| = 2$  so  $\mathfrak{G} \neq \text{iso III}$ .

Now  $\mathfrak{B}_1$  and  $\mathfrak{B}_2$  are nonisotropic and we know that  $\bar{\mathfrak{G}}_5$  is self-centralizing in its action on  $\mathfrak{B}_2$ . Write  $C_{\bar{\mathfrak{G}}}(\bar{\mathfrak{G}}_5) = \bar{\mathfrak{B}} \times \bar{\mathfrak{G}}_5$  where  $\bar{\mathfrak{B}} \triangleleft \bar{\mathfrak{G}}$ . Then  $\bar{\mathfrak{B}}$  acts faithfully on  $\mathfrak{B}_1$  so since  $\bar{\mathfrak{G}}_2 = \langle 1 \rangle$ , either  $\bar{\mathfrak{B}} = \langle 1 \rangle$  or  $\bar{\mathfrak{B}}$  has a normal 3-subgroup of order 3 which is clearly  $\bar{\mathfrak{G}}_3$ . Suppose  $\bar{\mathfrak{B}} = \langle 1 \rangle$ . Then  $\bar{\mathfrak{G}}/\bar{\mathfrak{G}}_5$  is a 2-group which acts on  $\mathfrak{B}_1$  and hence there is a 1-dimensional  $\bar{\mathfrak{G}}$ -invariant subspace  $\mathfrak{B}_0$  of  $\mathfrak{B}_1$ . Note that  $\bar{\mathfrak{G}} = \mathfrak{G}$  since  $\mathfrak{G} \neq \text{iso III}$  and thus if  $\mathfrak{G} \supseteq \mathfrak{G}_3 \supseteq Z(\mathfrak{G})$  with  $\mathfrak{G}_3/Z(\mathfrak{G}) = \mathfrak{B}_0 \oplus \mathfrak{B}_2$  then  $\mathfrak{G}_3 \triangleleft \mathfrak{G}$ . By the Reduction Lemma and Lemma 4.5 we have a contradiction since clearly  $\mathfrak{G}_3 \cong 3\mathcal{D}\mathcal{D}$ .



Thus  $\bar{\mathfrak{B}} \cong \bar{\mathfrak{F}}_3$  and  $|\bar{\mathfrak{F}}_3| = 3$ . Since  $q = 3$  we see that the Sylow 3-subgroups of  $\mathfrak{G}$  have order 3. Now  $\bar{\mathfrak{F}}_3$  centralizes  $\mathfrak{W}_2$  so clearly  $\mathfrak{G}$  contains precisely four Sylow 3-subgroups say  $\mathfrak{S}_i$  for  $i = 1, 2, 3, 4$ . Since  $q = 3$  each  $\mathfrak{S}_i$  has a fixed point on  $\mathfrak{V}^\#$  so by half-transitivity  $\mathfrak{V} = \bigcup_i C_{\mathfrak{V}}(\mathfrak{S}_i)$ . Hence since the  $\mathfrak{S}_i$  are all conjugate in  $\mathfrak{G}$  we see that each  $C_{\mathfrak{V}}(\mathfrak{S}_i)$  has codimension 1 in  $\mathfrak{V}$ . But  $|\mathfrak{V}| = 3^8$  so  $\mathfrak{V}_0 = \bigcap C_{\mathfrak{V}}(\mathfrak{S}_i) \neq \{0\}$ . Since  $\mathfrak{V}_0$  is clearly a proper  $\mathfrak{G}$ -invariant subspace of  $\mathfrak{V}$  we have a contradiction.

LEMMA 5.2.  $\bar{\mathfrak{F}}$  is not cyclic and  $q \neq 3$ .

*Proof.* We have shown that  $\bar{\mathfrak{F}} = \bar{\mathfrak{F}}_3$ . If  $\bar{\mathfrak{F}}$  is cyclic (including the possibility that  $\bar{\mathfrak{F}} = \langle 1 \rangle$ ) then clearly  $4 \nmid |\bar{\mathfrak{G}}|$ . If  $\mathfrak{G} = \text{iso I or II}$  then  $\mathfrak{G} = \mathfrak{H}$  so  $4 \nmid |\mathfrak{G}/\mathfrak{A}(\mathfrak{G})|$ . If  $\mathfrak{G} = \text{iso III}$  then  $8 \nmid |\mathfrak{G}/\mathfrak{A}(\mathfrak{G})|$ . If  $\mathfrak{G} = \text{iso I or II}$ , then  $\mathfrak{G}$  has an abelian subgroup of type  $(2, 2, 2)$  so we see that  $4 \mid |\mathfrak{G}_x|$ . If  $\mathfrak{G} = \text{iso III}$ , then  $\mathfrak{G}$  has an abelian subgroup of type  $(2, 2, 2, 2)$  so  $8 \mid |\mathfrak{G}_x|$ . Now by Lemma 1.5 there exists  $y \in \mathfrak{V}^\#$  with  $\mathfrak{G}_y \cap \mathfrak{A}(\mathfrak{G}) = \langle 1 \rangle$ . Hence  $|\mathfrak{G}_y| \mid |\mathfrak{G} : \mathfrak{A}(\mathfrak{G})|$ , a contradiction.

Let  $q = 3$  so that for all  $x \in \mathfrak{V}^\#$ ,  $\mathfrak{G}_x$  contains a Sylow 3-subgroup of  $\mathfrak{G}$ . Let  $\bar{\mathfrak{F}}$  be the complete inverse image of  $\bar{\mathfrak{F}}$  in  $\mathfrak{G}$ . For any  $x \in \mathfrak{V}^\#$ , let  $\mathfrak{S}$  be a Sylow 3-subgroup of  $\bar{\mathfrak{F}}_x$ . Then clearly  $\bar{\mathfrak{S}} = \mathfrak{S}\mathfrak{A}(\mathfrak{G})/\mathfrak{A}(\mathfrak{G}) = \bar{\mathfrak{F}}$  and since  $\mathfrak{G}_x = \mathfrak{G} \cap \mathfrak{G}_x \triangle \mathfrak{G}_x$  we see that  $\bar{\mathfrak{F}}$  normalizes  $\mathfrak{G}_x\mathfrak{Z}(\mathfrak{G})/\mathfrak{Z}(\mathfrak{G})$ . If  $\mathfrak{G} = \text{iso II or III}$ , then by Lemma 1.5 if  $T$  is any noncentral involution of  $\mathfrak{G}$  then for some  $x \in \mathfrak{V}^\#$ ,  $\mathfrak{G}_x = \langle T \rangle$ . This implies that  $\bar{\mathfrak{F}}$  fixes all involution vectors and  $\bar{\mathfrak{F}} = \langle 1 \rangle$ , a contradiction. If  $\mathfrak{G} = \text{iso I}$  then by Lemma 1.5,  $|\mathfrak{G}_x| = 1$  or 4. However here it is easy to see that for each such  $T$  we can find two points  $x_1, x_2 \in \mathfrak{V}^\#$  with  $\langle T \rangle = \mathfrak{G}_{x_1} \cap \mathfrak{G}_{x_2}$ . This again implies that  $\bar{\mathfrak{F}}$  fixes all involution vectors and the result follows.

LEMMA 5.3.  $\mathfrak{G} = \text{iso I}$  does not occur.

*Proof.* Here  $\mathfrak{G} \cong \Omega\Omega\Omega$  and we see easily that  $\text{Aut } \mathfrak{G}$  contains  $\bar{\mathfrak{F}} \sim \bar{\mathfrak{F}}$  where  $|\bar{\mathfrak{F}}| = 3$  and this is a full Sylow 3-subgroup of  $Sp(6, 2)$ . Then any 3-group acting on  $\mathfrak{G}$  can be embedded in this Sylow 3-subgroup. Let  $\bar{\mathfrak{K}}$  be a Sylow 3-subgroup of  $\text{Aut } \mathfrak{G}$ . Then  $\bar{\mathfrak{K}}$  acts faithfully on  $\mathfrak{W} = \mathfrak{G}/\mathfrak{Z}(\mathfrak{G})$ . As a Sylow 3-subgroup of  $Sp(6, 2)$  we know that it has the following structure. We can write  $\mathfrak{W} = \mathfrak{W}_1 \oplus \mathfrak{W}_2 \oplus \mathfrak{W}_3$ , a direct sum of orthogonal 2-dimensional nonisotropic subspaces.  $\bar{\mathfrak{K}}$  has a subgroup  $\bar{\mathfrak{N}}$  of index 3 with  $\bar{\mathfrak{N}} = \bar{\mathfrak{S}}_1 \times \bar{\mathfrak{S}}_2 \times \bar{\mathfrak{S}}_3$ . Here  $|\bar{\mathfrak{S}}_i| = 3$  and  $\bar{\mathfrak{S}}_i$  acts irreducibly on  $\mathfrak{W}_i$  and centralizes the remaining  $\mathfrak{W}_j$ . Further, any element of  $\bar{\mathfrak{K}} - \bar{\mathfrak{N}}$  permutes these three subspaces. Now let  $\mathfrak{G}_i$

be the subgroup of  $\mathcal{G}$  with  $\mathcal{G}_i/Z(\mathcal{G}) = \mathfrak{W}_i$ . Then  $\mathcal{G}_i$  is nonabelian of order 8 and admits an automorphism of order 3. Thus  $\mathcal{G}_i \cong \mathcal{Q}$ . Suppose  $T = T_1T_2T_3$  is a noncentral involution of  $\mathcal{G}$  with  $T_i \in \mathcal{G}_i$ . Since  $\mathcal{G}_i \cong \mathcal{Q}$  we see that precisely one of the  $T_i$  is contained in  $Z(\mathcal{G})$ , say for example  $T_1$ . Then we can write  $T = T_2T_3$ . If some subgroup  $\bar{\mathfrak{X}}$  of  $\bar{\mathfrak{K}}$  centralizes the involution vector corresponding to  $T$  then clearly  $\bar{\mathfrak{X}}$  normalizes  $\mathfrak{W}_1$ . Thus  $\bar{\mathfrak{X}} \subseteq \bar{\mathfrak{N}}$  so  $\bar{\mathfrak{X}}$  normalizes  $\mathfrak{W}_2$  and  $\mathfrak{W}_3$ . This clearly implies that  $\bar{\mathfrak{X}}$  centralizes  $\mathfrak{W}_2$  and  $\mathfrak{W}_3$  and thus  $\bar{\mathfrak{X}} = \bar{\mathfrak{X}}_i$ . Hence the only subgroups of  $\bar{\mathfrak{K}}$  which centralize involution vectors are  $\bar{\mathfrak{X}}_1, \bar{\mathfrak{X}}_2$  and  $\bar{\mathfrak{X}}_3$ .

Now  $\bar{\mathfrak{X}}$  is not cyclic and hence a Sylow 3-subgroup of  $\mathcal{G}$  is not cyclic. Thus  $3 \mid |\mathcal{G}_x|$  for all  $x \in \mathfrak{B}^*$ . By the preceding lemma again  $q \neq 3$ . Hence if  $T \in \mathcal{G}$  is an involution, then by Lemma 1.5 there exists  $x \in \mathfrak{B}^*$  with  $\mathcal{G}_x = \langle T \rangle$ . Let  $\mathfrak{X}$  be a Sylow 3-subgroup of  $\mathcal{G}_x$  so  $|\mathfrak{X}| \geq 3$  and  $\mathfrak{X} \cap \mathcal{A}\mathcal{G} = \langle 1 \rangle$ . Then  $\bar{\mathfrak{X}}$  acts faithfully on  $\mathcal{G}$  so we can extend  $\mathfrak{X}$  to  $\bar{\mathfrak{K}}$  as above. Since  $\bar{\mathfrak{X}}$  normalizes the involution vector corresponding to  $T$  we see that  $\bar{\mathfrak{X}} = \bar{\mathfrak{X}}_i$  for some  $i$ . Thus  $|\bar{\mathfrak{X}}| = 3$  and  $9 \nmid |\mathcal{G}_x|$ .

Suppose  $\bar{\mathcal{G}} = \mathcal{G}/\mathcal{A}\mathcal{G}$  contains a copy of  $\bar{\mathfrak{N}} \subseteq \bar{\mathfrak{K}}$ . Then let  $\mathcal{C}$  be a 3-subgroup of  $\mathcal{G}$  with  $\mathcal{C}\mathcal{A}\mathcal{G}/\mathcal{A}\mathcal{G} = \bar{\mathfrak{N}}$ . Certainly  $\mathcal{C}' \subseteq \mathcal{A}$ . Now  $\mathcal{C}$  acts on  $\mathfrak{B}$ , a vector space of dimension  $n = 2^4$ . Since  $\mathcal{C}$  is a 3-group we conclude that  $\mathcal{C}'$  is in the kernel of some irreducible constituent and hence  $\mathcal{C}'$  has a fixed point in  $\mathfrak{B}^*$ . Since  $\mathcal{C}' \subseteq \mathcal{A}$  we see that  $\mathcal{C}' = \langle 1 \rangle$  and  $\mathcal{C}$  is abelian. Now  $\mathcal{C}/\mathcal{C}' \cap \mathcal{A}$  is abelian of type  $(3, 3, 3)$  and hence  $\mathcal{C}$  contains a subgroup of type  $(3, 3, 3)$ . But this implies that  $9 \mid |\mathcal{G}_x|$ , a contradiction. In particular we see that a Sylow 3-subgroup of  $\bar{\mathcal{G}}$  has order  $\leq 3^3$ .

Let  $T$  and  $\mathfrak{X}$  be as above and set  $\bar{\mathfrak{X}} = \mathfrak{X}\mathcal{A}\mathcal{G}/\mathcal{A}\mathcal{G}$ . This time embed 3-group  $\bar{\mathfrak{X}}\bar{\mathfrak{X}}$  in  $\bar{\mathfrak{K}}$ . Again  $\bar{\mathfrak{X}} = \bar{\mathfrak{X}}_i$  for some  $i$ . Now  $\bar{\mathfrak{K}}$  is generated by  $\bar{\mathfrak{X}}_i$  and any element outside  $\bar{\mathfrak{N}}$ . Since  $\bar{\mathfrak{X}}\bar{\mathfrak{X}} < \bar{\mathfrak{K}}$  we must have  $\bar{\mathfrak{X}} \subseteq \bar{\mathfrak{N}}$  and hence  $\bar{\mathfrak{X}}\bar{\mathfrak{X}} \subseteq \bar{\mathfrak{N}}$ . Since  $\bar{\mathfrak{X}}$  centralizes  $\bar{\mathfrak{X}}$  we have  $\bar{\mathfrak{X}} \subseteq \bar{\mathfrak{X}}$ .

Now embed  $\bar{\mathfrak{X}}$  alone in  $\bar{\mathfrak{K}}$ . We have shown that for each involution vector of  $\mathfrak{B}$ ,  $\bar{\mathfrak{X}}$  contains a subgroup of order 3 centralizing it. Thus  $\bar{\mathfrak{X}} \supseteq \bar{\mathfrak{X}}_1, \bar{\mathfrak{X}}_2, \bar{\mathfrak{X}}_3$  and  $\bar{\mathfrak{X}} \supseteq \bar{\mathfrak{N}}$ , a contradiction since  $\bar{\mathcal{G}} \not\supseteq \bar{\mathfrak{N}}$ . This completes the proof of this result.

LEMMA 5.4.  $\mathcal{G} = \text{iso II and III do not occur.}$

*Proof.* Suppose  $C_{\mathfrak{B}}(\bar{\mathfrak{X}}) = \mathfrak{W}_1 \neq \langle 1 \rangle$ . Then  $\mathfrak{B} = \mathfrak{W}_1 \oplus \mathfrak{W}_2$  where  $\mathfrak{W}_2 = (\mathfrak{B}, \bar{\mathfrak{X}})$ . Since  $\mathfrak{W}_2$  has even dimension (the nonprincipal irreducible representations of a 3-group over  $GF(2)$  have even dimension) so does  $\mathfrak{W}_1$ . One of these two subspaces, say  $\mathfrak{W}_1$ , has dimension equal to 4.

Let  $\mathbb{C}_i$  be the subgroup of  $\mathbb{C}$  with  $\mathbb{C}_i/Z(\mathbb{C}) = \mathbb{W}_i$ . Then  $\mathbb{C}_i \triangle \mathbb{G}$  and  $\mathbb{G}$  is primitive so  $\mathbb{C}_i$  is of symplectic type. By the Reduction Lemma and Lemmas 4.3, 4.4 and 4.5 we have  $q = 3$ , a contradiction by Lemma 5.2.

Now let  $\mathbb{C} = \text{iso II}$ . By Lemma 1.3,  $\bar{\mathbb{F}}$  permutes the  $i(\mathbb{W}) = 35$  involution vectors. Hence  $\bar{\mathbb{F}}$  must fix one of these and  $C_{\mathbb{W}}(\bar{\mathbb{F}}) \neq \langle 1 \rangle$ , a contradiction.

Having already eliminated  $\mathbb{C} = \text{iso I}$  and II we now eliminate iso III.  $\bar{\mathbb{F}}$  acts on  $\mathbb{C}/\mathbb{C}' = \mathbb{U}$  and centralizes  $Z(\mathbb{C})/\mathbb{C}'$ . Since  $C_{\mathbb{W}}(\bar{\mathbb{F}}) = \langle 1 \rangle$  we see that  $\mathbb{U} = \mathbb{U}_1 \oplus \mathbb{U}_2$  where  $\mathbb{U}_1 = C_{\mathbb{W}}(\bar{\mathbb{F}})$ ,  $\mathbb{U}_2 = (\mathbb{U}, \bar{\mathbb{F}})$ ,  $|\mathbb{U}_1| = 2$ ,  $|\mathbb{U}_2| = 2^4$ . Let  $\mathbb{C}_2$  be a subgroup of  $\mathbb{C}$  with  $\mathbb{C}_2/\mathbb{C}' = \mathbb{U}_2$ . Then  $\mathbb{C}_2 \triangle \mathbb{G}$  and  $\mathbb{C}_2$  is type  $E(2, 3)$  and iso I or II. By the Reduction Lemma and the above we have a contradiction.

We now consider  $m = 4$ . Here we have partial results in Lemmas 2.6, 2.10 and 2.12. Thus  $\mathbb{C} \neq \text{iso III}$ ,  $q \geq 7$  and  $|\mathbb{G}/\mathbb{W}\mathbb{C}| > 10^4$ . We consider  $\bar{\mathbb{F}}$ .

LEMMA 5.5. *All irreducible constituents of  $\bar{\mathbb{F}}_p$  on  $\mathbb{W}$  have the same degree. Thus  $\bar{\mathbb{F}}_2 = \langle 1 \rangle$ ,  $\bar{\mathbb{F}}_p = \langle 1 \rangle$  if  $p \nmid i(\mathbb{W})$  and  $\bar{\mathbb{F}}_3$  is elementary abelian.*

*Proof.* Suppose  $\bar{\mathbb{F}}_2 \neq \langle 1 \rangle$ . Then by Lemma 4.1,  $\mathbb{G}$  has a normal subgroup  $\mathbb{C}_0$  of type  $E(2, 3)$  and iso III. By the Reduction Lemma and Lemma 5.4 this is a contradiction.

If  $p \neq 2$  then  $\bar{\mathbb{F}}_p$  acts in a completely reducible manner on  $\mathbb{W}$ . If all its irreducible constituents do not have the same degree, then certainly we can write  $\mathbb{W} = \mathbb{W}_1 \oplus \mathbb{W}_2$  where  $\mathbb{W}_i \neq \langle 1 \rangle$  and  $\mathbb{W}_i$  is  $\bar{\mathbb{G}}$  invariant. One of these two, say  $\mathbb{W}_1$ , has dimension at least 4. If  $\mathbb{C}_1/Z(\mathbb{C}) = \mathbb{W}_1$  then  $\mathbb{C}_1 \triangle \mathbb{G}$  and since  $\mathbb{G}$  is primitive,  $\mathbb{C}_1$  is type  $E(2, m')$  with  $m' = 2$  or 3. Since  $q \geq 7$ , the Reduction Lemma and the  $m = 2$  and 3 results yield a contradiction. Now if  $p \nmid i(\mathbb{W})$ , then certainly  $\bar{\mathbb{F}}_p$  has a 1-dimensional constituent so they are all 1-dimensional and over  $GF(2)$  this implies that  $\bar{\mathbb{F}}_p$  centralizes  $\mathbb{W}$  so  $\bar{\mathbb{F}}_p = \langle 1 \rangle$ .

Finally we consider  $\bar{\mathbb{F}}_3$ . If  $\bar{\mathbb{F}}_3$  is nonabelian then the degree of an irreducible representation of  $\bar{\mathbb{F}}_3$ , with  $\bar{\mathbb{F}}'_3$  not in the kernel is divisible by 3. Since  $3 \nmid \dim \mathbb{W}$ ,  $\bar{\mathbb{F}}'_3$  is in the kernel of all constituents so  $\bar{\mathbb{F}}'_3 = \langle 1 \rangle$  and  $\bar{\mathbb{F}}_3$  is abelian. Let  $\mathbb{W}_0$  be an irreducible  $\bar{\mathbb{F}}_3$ -constituent of  $\mathbb{W}$  with dimension  $j$ . Then  $j \mid \dim \mathbb{W}$  so  $j = 1, 2, 4$  or 8. In all these cases  $9 \nmid 2^j - 1$  and hence clearly  $\bar{\mathbb{F}}_3$  is elementary abelian.

LEMMA 5.6.  $\mathbb{C} = \text{iso I}$  does not occur.

*Proof.* Here by Lemma 1.3,  $i(\mathbb{W}) = 3^3 \cdot 5$  so only  $\bar{\mathbb{F}}_5$  and  $\bar{\mathbb{F}}_3$  can

be nontrivial. We show first that  $\bar{\mathfrak{F}}_5 = \langle 1 \rangle$ . Note that a Sylow 5-subgroup of  $Sp(8, 2)$  is abelian of type (5, 5).

Suppose first that  $|\bar{\mathfrak{F}}_5| = 5^2$ . Then  $\bar{\mathfrak{F}}_5$  is elementary abelian and a Sylow 5-subgroup of  $\mathfrak{G}$ . We can write  $\mathfrak{B} = \mathfrak{B}_1 \oplus \mathfrak{B}_2$ ,  $\bar{\mathfrak{F}}_5 = \bar{\mathfrak{F}}_1 \bar{\mathfrak{F}}_2$  where  $\dim \mathfrak{B}_i = 4$ ,  $|\bar{\mathfrak{F}}_i| = 5$  and  $\bar{\mathfrak{F}}_i$  acts irreducibly on  $\mathfrak{B}_i$  and centralizes the other  $\mathfrak{B}_j$ . Now a Sylow 5-subgroup of  $\mathfrak{G}$  is not cyclic so  $5 \mid |\mathfrak{G}_x|$  for all  $x \in \mathfrak{B}^\#$ . We have  $i(\mathfrak{B}) = 135$  and  $|\mathfrak{B}_1 \cup \mathfrak{B}_2| = 31$ . Hence we can find a noncentral involution  $T \in \mathfrak{G}$  with  $TZ(\mathfrak{G})/Z(\mathfrak{G}) \not\subseteq \mathfrak{B}_1 \cup \mathfrak{B}_2$ . By Lemma 1.5 there exists  $x \in \mathfrak{B}^\#$  with  $\mathfrak{G}_x = \langle T \rangle$  and if  $\mathfrak{B} \subseteq \mathfrak{G}_x$  has order 5, then  $\mathfrak{B}$  normalizes  $\mathfrak{G}_x \cap \mathfrak{G} = \langle T \rangle$ . Thus  $\bar{\mathfrak{B}} = \mathfrak{B}\mathfrak{G}/\mathfrak{A}\mathfrak{G} \subseteq \bar{\mathfrak{F}}_5$  centralizes the involution vector corresponding to  $T$ . Since  $C_{\mathfrak{B}}(\bar{\mathfrak{F}}_1) = \mathfrak{B}_2$  and  $C_{\mathfrak{B}}(\bar{\mathfrak{F}}_2) = \mathfrak{B}_1$  we see by our choice of  $T$  that  $\bar{\mathfrak{B}} \neq \bar{\mathfrak{F}}_1$  or  $\bar{\mathfrak{F}}_2$ . But then  $\bar{\mathfrak{B}}$  acts irreducibly on  $\mathfrak{B}_1$  and  $\mathfrak{B}_2$  so by the Jordan-Holder Theorem,  $C_{\mathfrak{B}}(\bar{\mathfrak{B}}) = \langle 1 \rangle$ , a contradiction.

Now let  $|\bar{\mathfrak{F}}_5| = 5$ . By the preceding lemma  $\bar{\mathfrak{F}}_5$  is abelian. Since the irreducible nonprincipal representations of  $\bar{\mathfrak{F}}_5$  over  $GF(2)$  have degree 4 we see that either  $\bar{\mathfrak{F}}_5$  is irreducible or it has two irreducible constituents of dimension 4. Thus  $\bar{\mathfrak{F}}_5$  has two generators and  $\bar{\mathfrak{F}}_5$  is abelian of type (5), (3, 5) or (3, 3, 5). Hence

$$|\bar{\mathfrak{G}}| \leq 3^2 |GL(2, 3)| \cdot 5 \cdot 4 = 8640 < 10^4,$$

a contradiction.

Thus  $\bar{\mathfrak{F}}_5 = \bar{\mathfrak{F}}_3$  is elementary abelian. If  $|\bar{\mathfrak{F}}_3| \leq 3^2$ , then

$$|\bar{\mathfrak{G}}| \leq 3^2 |GL(2, 3)| = 432 < 10^4,$$

a contradiction. If  $|\bar{\mathfrak{F}}_3| = 3^3$ , then  $|\bar{\mathfrak{G}}|$  divides both  $|\bar{\mathfrak{F}}_3| |GL(3, 3)| = 2^5 \cdot 3^6 \cdot 13$  and  $|Sp(8, 2)| = 2^{16} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 17$  so  $|\bar{\mathfrak{G}}|$  divides  $2^5 \cdot 3^5 = 7776 < 10^4$ , a contradiction. Since the Sylow 3-subgroup of  $Sp(8, 2)$  is nonabelian of order  $3^5$  this leaves only  $|\bar{\mathfrak{F}}_3| = 3^4$ .

Let  $\mathfrak{S}$  be a 3-subgroup of  $\mathfrak{G}$  with  $\mathfrak{S}\mathfrak{A}\mathfrak{G}/\mathfrak{A}\mathfrak{G} = \bar{\mathfrak{F}}_3$ . Clearly  $\mathfrak{S}' \subseteq \mathfrak{A}$ . The action of  $\mathfrak{S}$  on  $\mathfrak{B}$  is completely reducible since  $q \neq 3$  and since  $\dim \mathfrak{B} = 2^4$  is not divisible by 3 it follows that  $\mathfrak{S}'$  is in the kernel of some constituent so  $\mathfrak{S}'$  has a fixed point in  $\mathfrak{B}^\#$ . Since  $\mathfrak{A}$  acts semi-regularly,  $\mathfrak{S}' = \langle 1 \rangle$ . Now  $\mathfrak{S}$  is abelian and  $\mathfrak{S}/(\mathfrak{S} \cap \mathfrak{A})$  is abelian of type (3, 3, 3, 3). Thus  $\mathfrak{S}$  contains a subgroup of type (3, 3, 3, 3) and hence  $3^3 \mid |\mathfrak{G}_x|$ .

Now  $\mathfrak{G} \cong \mathfrak{D}\mathfrak{D}\mathfrak{D}\mathfrak{D}$  so it is clear that the automorphism group of  $\mathfrak{G}$  contains  $\bar{\mathfrak{R}} = \bar{\mathfrak{F}} \times (\bar{\mathfrak{F}} \sim \bar{\mathfrak{F}})$  where  $|\bar{\mathfrak{F}}| = 3$ . This group is a Sylow 3-subgroup of  $Sp(8, 2)$  and hence is a Sylow 3-subgroup of  $\text{Aut } \mathfrak{G}$ . We describe it more precisely. Write  $\mathfrak{G} = \mathfrak{G}_0 \mathfrak{G}_1 \mathfrak{G}_2 \mathfrak{G}_3$  where each  $\mathfrak{G}_i \cong \mathfrak{D}$ . Then  $\bar{\mathfrak{R}}$  has an elementary abelian subgroup  $\bar{\mathfrak{R}}$  of index 3 with

$\bar{\mathfrak{N}} = \bar{\mathfrak{G}}_0 \bar{\mathfrak{G}}_1 \bar{\mathfrak{G}}_2 \bar{\mathfrak{G}}_3$ . Here  $\bar{\mathfrak{G}}_i$  acts nontrivially on  $\mathfrak{G}_i$  and centralizes the remaining  $\mathfrak{G}_j$ . Every element of  $\bar{\mathfrak{N}} - \bar{\mathfrak{N}}$  normalizes  $\mathfrak{G}_0$  and cyclically permutes  $\mathfrak{G}_1, \mathfrak{G}_2$  and  $\mathfrak{G}_3$ . Let  $\mathfrak{B} = \mathfrak{B}_0 \oplus \mathfrak{B}_1 \oplus \mathfrak{B}_2 \oplus \mathfrak{B}_3$  be the corresponding decomposition of  $\mathfrak{B}$ .

Let  $T$  be a noncentral involution of  $\mathfrak{G}$ . Then there exists  $x \in \mathfrak{B}^\#$  by Lemma 1.5 with  $\mathfrak{G}_x = \langle T \rangle$ . Since  $3^3 \mid |\mathfrak{G}_x|$  let  $\mathfrak{X}$  be a subgroup of  $\mathfrak{G}_x$  of order  $3^3$ . Then  $\mathfrak{X}$  normalizes  $\mathfrak{G}_x \cap \mathfrak{G} = \mathfrak{G}_x = \langle T \rangle$ . Since  $\mathfrak{X} \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$ ,  $\mathfrak{X}$  acts faithfully on  $\mathfrak{G}$ . Thus a suitable conjugate  $\bar{\mathfrak{X}}$  of  $\mathfrak{X}$  in  $\text{Aut } \mathfrak{G}$  is contained in  $\bar{\mathfrak{N}}$  and clearly  $\bar{\mathfrak{X}}$  also centralizes an involution vector of  $\mathfrak{B}$ . Let  $W = W_0 + W_1 + W_2 + W_3 \in \mathfrak{B}$  with  $W_i \in \mathfrak{B}_i$ . Then we see easily that  $W$  is an involution vector if and only if either none or two of the  $W_i$  are zero. Suppose two of the  $W_i$  are zero. Then clearly  $C_{\bar{\mathfrak{N}}}(W) \subseteq \bar{\mathfrak{N}}$  and then  $|C_{\bar{\mathfrak{N}}}(W)| \leq 3^2$ . If none of the  $W_i$  are zero, then  $C_{\bar{\mathfrak{N}}}(W) \cap \bar{\mathfrak{N}} = \langle 1 \rangle$  so  $|C_{\bar{\mathfrak{N}}}(W)| \leq 3$ . This contradicts the fact that  $|\bar{\mathfrak{X}}| = 3^3$  and  $\bar{\mathfrak{X}}$  fixes an involution vector.

LEMMA 5.7.  $\mathfrak{G} = \text{iso II}$  does not occur.

*Proof.* Here  $i(\mathfrak{B}) = 7 \cdot 17$  by Lemma 1.3. Hence only  $\bar{\mathfrak{F}}_7$  and  $\bar{\mathfrak{F}}_{17}$  can be nontrivial. If  $\bar{\mathfrak{F}}_7 \neq \langle 1 \rangle$  then since  $7^2 \nmid |Sp(8, 2)|$ ,  $|\bar{\mathfrak{F}}_7| = 7$ . But the nonprincipal irreducible representations of this group over  $GF(2)$  all have degree 3. Since  $3 \nmid \dim \mathfrak{B}$  we have a contradiction. Then  $\bar{\mathfrak{F}} = \bar{\mathfrak{F}}_{17}$  has order 1 or 17 and  $|\bar{\mathfrak{G}}| \leq 17 \cdot 16 < 10^4$ , a contradiction.

We have therefore shown in this section that if  $\mathfrak{G}$  is solvable then  $m = 3$  and 4 do not occur.

6. Theorem B. The following assumption holds throughout this section.

ASSUMPTION. Group  $\mathfrak{G}$  acts faithfully on vector space  $\mathfrak{B}$  of order  $q^n$ ,  $q$  a prime, and acts half-transitively but not semiregularly on  $\mathfrak{B}^\#$ . Further  $\mathfrak{G}$  is primitive as a linear group and  $\mathfrak{G}$  is solvable.

Let  $\mathcal{S}(q^n)$  denote the group of all semilinear transformations on  $GF(q^n)$  of the form  $x \rightarrow ax^\sigma$  where  $a \in GF(q^n)^\#$  and  $\sigma$  is a field automorphism. Thus  $\mathcal{S}(q^n)$  is the stabilizer in the permutation group  $\mathcal{S}(q^n)$  of the point 0.

LEMMA 6.1. Let  $\bar{\mathfrak{F}} = F(\mathfrak{G})$  and set  $\mathfrak{A} = Z(C_{\bar{\mathfrak{F}}}(\Phi(\bar{\mathfrak{F}})))$ . Then  $\mathfrak{A}$  is a normal cyclic subgroup of  $\mathfrak{G}$

(i) If  $\mathfrak{A} = C_{\bar{\mathfrak{F}}}(\Phi(\bar{\mathfrak{F}}))$ , then with suitable identification we have  $\mathfrak{G} \subseteq \mathcal{S}(q^n)$ .

(ii) If  $\mathfrak{A} \neq C_{\bar{\mathfrak{F}}}(\Phi(\bar{\mathfrak{F}}))$ , then  $C_{\bar{\mathfrak{F}}}(\Phi(\bar{\mathfrak{F}})) = \mathfrak{A}\mathfrak{G}$  where  $\mathfrak{G}$  is a group of type  $E(2, m)$  and  $\mathfrak{G} \triangleleft \mathfrak{G}$ . Moreover  $m = 1$  or 2.

(iii) *In the above if  $m = 1$  and  $4 \nmid |\mathfrak{A}|$ , then either  $\mathfrak{G} \cong \mathcal{S}(q^n)$  or  $q^n = 3^2, 7^2$  or  $11^2$ .*

*Proof.* Let  $\mathfrak{F}_p$  be the normal Sylow  $p$ -subgroup of  $\mathfrak{F}$ . By Theorem A  $\mathfrak{F}_p$  is cyclic for  $p > 2$  and  $\mathfrak{F}_2$  is a group of symplectic type. Since  $\mathfrak{A} = Z(C_{\mathfrak{F}}(\Phi(\mathfrak{F})))$  is a normal abelian subgroup of a primitive group it is cyclic.

From the structure of 2-groups of symplectic type we see that if  $\mathfrak{A} = C_{\mathfrak{F}}(\Phi(\mathfrak{F}))$ , then  $\mathfrak{F}_2$  is either cyclic or maximal class of order at least 16. Now  $\mathfrak{F} = \mathfrak{A}\mathfrak{F}_2$  so  $C_{\mathfrak{G}}(\mathfrak{A})/Z(\mathfrak{F})$  acts faithfully on  $\mathfrak{F}_2$ . Since  $\text{Aut } \mathfrak{F}_2$  is a 2-group and  $Z(\mathfrak{F}) \subseteq \mathfrak{A}$  we see that  $C_{\mathfrak{G}}(\mathfrak{A})$  is a normal nilpotent subgroup of  $\mathfrak{G}$  and hence  $C_{\mathfrak{G}}(\mathfrak{A}) \subseteq \mathfrak{F}$ . This yields easily  $C_{\mathfrak{G}}(\mathfrak{A}) = \mathfrak{A}$ . By Proposition 1.2 of [5] we see that  $\mathfrak{G} \cong \mathcal{S}(q^n)$  and (i) follows.

Suppose  $\mathfrak{A} \neq C_{\mathfrak{F}}(\Phi(\mathfrak{F}))$ . Then as we pointed out in § 1,  $C_{\mathfrak{F}}(\Phi(\mathfrak{F})) = \mathfrak{A}\mathfrak{C}$  where  $\mathfrak{C}$  is a group of type  $E(2, m)$  and  $\mathfrak{C} \triangleleft \mathfrak{G}$ . By Theorem A and the results of § 5,  $m = 1$  or  $2$ .

Let  $m = 1$  and suppose  $4 \nmid |\mathfrak{A}|$ . Then  $\mathfrak{F}_2 \cong \mathfrak{D}$  or  $\mathfrak{Q}$ . If  $\mathfrak{F}_2 \cong \mathfrak{D}$  then  $\mathfrak{F}$  has a characteristic cyclic subgroup  $\mathfrak{B}$  of index 2. Since  $\text{Aut } \mathfrak{D}$  is a 2-group, the above argument yields  $\mathfrak{G} \cong \mathcal{S}(q^n)$  again. If  $\mathfrak{F}_2 \cong \mathfrak{Q}$ , then by Proposition 1.10 of [5]  $q^n = 3^2, 7^2$  or  $11^2$ . This completes the proof.

We assume now that  $\mathfrak{A} \neq C_{\mathfrak{F}}(\Phi(\mathfrak{F}))$ .

**LEMMA 6.2.** *Let  $\mathfrak{B} = C_{\mathfrak{G}}(\mathfrak{A})/\mathfrak{A}\mathfrak{C}$ . Then  $O_2(\mathfrak{B}) = \langle 1 \rangle$ ,  $\mathfrak{B}$  acts faithfully on  $\mathfrak{C}/Z(\mathfrak{C})$  and  $\mathfrak{B} \cong Sp(2m, 2)$ .*

*Proof.* Let  $\mathfrak{L}/\mathfrak{A}\mathfrak{C} = O_2(\mathfrak{B})$ . Since  $\mathfrak{A}$  is central in  $\mathfrak{L}$  and  $\mathfrak{L}/\mathfrak{A}$  is a 2-group, we see that  $\mathfrak{L}$  is a normal nilpotent subgroup of  $\mathfrak{G}$  and hence  $\mathfrak{L} \subseteq \mathfrak{F}$ . Now  $\Phi(\mathfrak{F}) \subseteq \mathfrak{A}$  and  $C_{\mathfrak{F}}(\Phi(\mathfrak{F})) = \mathfrak{A}\mathfrak{C}$ . Hence

$$\mathfrak{L} \subseteq C_{\mathfrak{F}}(\mathfrak{A}) \subseteq C_{\mathfrak{F}}(\Phi(\mathfrak{F})) = \mathfrak{A}\mathfrak{C}$$

so  $\mathfrak{L} = \mathfrak{A}\mathfrak{C}$  and  $O_2(\mathfrak{B}) = \langle 1 \rangle$ .

Let  $\mathfrak{H} = C_{\mathfrak{G}}(\mathfrak{A})$  and let  $\mathfrak{R} = C_{\mathfrak{H}}(\mathfrak{B})$  where  $\mathfrak{B} = \mathfrak{C}/Z(\mathfrak{C})$ . We have of course  $\mathfrak{R} \supseteq \mathfrak{A}\mathfrak{C}$ . First  $\mathfrak{R}$  centralizes  $O_2(\mathfrak{F}) \subseteq \mathfrak{A}$ . If  $\mathfrak{F}_2 = O_2(\mathfrak{F})$ , then since clearly  $[\mathfrak{F}_2 : \mathfrak{A}_2\mathfrak{C}] = 2$ , where  $\mathfrak{A}_2 = \mathfrak{A} \cap \mathfrak{F}_2$ , we see that  $\mathfrak{R}$  stabilizes the chain  $\mathfrak{F}_2 \supseteq \mathfrak{A}_2\mathfrak{C} \supseteq \mathfrak{A}_2 \supseteq \langle 1 \rangle$ . Thus  $\mathfrak{R}/C_{\mathfrak{R}}(\mathfrak{F})$  is a 2-group. Since  $\mathfrak{R} \supseteq Z(\mathfrak{F})$ ,  $C_{\mathfrak{R}}(\mathfrak{F}) = Z(\mathfrak{F})$  and hence  $\mathfrak{R}/Z(\mathfrak{F})$  is a 2-group. But  $Z(\mathfrak{F}) \subseteq \mathfrak{A}$  and  $\mathfrak{A}$  is central in  $\mathfrak{R}$  so  $\mathfrak{R}$  is a normal nilpotent subgroup of  $\mathfrak{G}$  and  $\mathfrak{R} \subseteq \mathfrak{F}$ . This yields easily  $\mathfrak{R} = \mathfrak{A}\mathfrak{C}$  and thus  $\mathfrak{B} = \mathfrak{H}/\mathfrak{R}$  acts faithfully on  $\mathfrak{B}$ . It now follows immediately that  $\mathfrak{B} \cong Sp(2m, 2)$ .

**LEMMA 6.3.** *Let  $\mathfrak{A} = \langle A \rangle$  and let  $\zeta$  be an eigenvalue of  $A$  with  $GF(q)(\zeta) = GF(q^r)$ . Then*

- (i)  $C_{\mathfrak{G}}(\mathfrak{A}) \cong GL(n/r, q^r)$ ,  $|\mathfrak{A}| \mid (q^r - 1)$
- (ii)  $\mathfrak{G}/C_{\mathfrak{G}}(\mathfrak{A})$  is cyclic of order dividing  $r$ .
- (iii)  $n = w2^m r$  for some integer  $w$ .

*Proof.* Parts (i) and (ii) follow from Lemma 1.1 of [5]. Now all irreducible constituents of  $\mathfrak{G}$  are faithful and the same is clearly true if we view  $\mathfrak{G} \cong GL(n/r, q^r)$ . Thus  $n/r$  is divisible by  $2^m$ , the degree of the nonlinear absolutely irreducible representations of  $\mathfrak{G}$ .

LEMMA 6.4. *If  $m = 1$  and  $4 \mid |\mathfrak{A}|$ , then  $q^n = 5^2$  or  $17^2$ .*

*Proof.* We can assume that  $|Z(\mathfrak{G})| = 4$  so  $\mathfrak{G} \cong 3\mathfrak{D}$ . By the Reduction Lemma and Lemma 3.4,  $q = 3, 5$  or  $17$ . Set  $\mathfrak{H} = C_{\mathfrak{G}}(\mathfrak{A})$ . Then by the above  $\mathfrak{H}/\mathfrak{A}\mathfrak{H} = \mathfrak{B}$  is contained isomorphically in  $Sp(2, 2) = SL(2, 2) \cong \text{Sym}_3$ . Since  $O_2(\mathfrak{B}) = \langle 1 \rangle$ ,  $|\mathfrak{B}| = 1, 3$  or  $6$ .

Suppose  $|\mathfrak{B}| = 1$ . Now  $2 \mid |\mathfrak{G}_x|$  so we can apply Lemma 3.1 with  $p = 2$ . Note that  $\mathfrak{G}/\mathfrak{A}\mathfrak{G}$  is cyclic of order dividing  $r$  and  $k = n/r = 2w$ . If  $r$  is odd, then  $\lambda_1 \leq 3$ ,  $\lambda_2 = 0$  so by Lemma 3.1, (ii) and (iii), we have  $q^r < 6$  so  $r = 1$ . If  $r$  is even, then  $\lambda_1 \leq 3$ ,  $\lambda_2 \leq 4$  so we get easily  $q^{r/2} \leq 5$  and hence  $r = 2$ . Now  $\mathfrak{G}$  has precisely three normal abelian subgroups of type  $(2, 2)$ . Since  $\mathfrak{G}/\mathfrak{A}\mathfrak{G}$  is a 2-group one of these three abelian groups will be normal in  $\mathfrak{G}$ , a contradiction since  $\mathfrak{G}$  is primitive. Thus  $|\mathfrak{B}| = 3$  or  $6$ .

Suppose  $3 \mid |\mathfrak{G}_x|$ . We again apply Lemma 3.1. If  $3 \nmid r$  then  $\lambda_1 \leq 4$ ,  $\lambda_2 = 0$  while if  $3 \mid r$ , then  $\lambda_1 \leq 4$  and we see easily that  $\lambda_2 \leq 9$ . Let  $3 \nmid r$  so by Lemma 3.1 we have  $q^r < 8$ . Since  $4 \mid q^r - 1$ ,  $q^r = 5$  and then by Lemma 3.1 (i) we have  $k = 2$  and  $n = 2$ . But  $3 \nmid q - 1$  so no element of  $GL(2, 5)$  of order 3 can have a nonzero fixed point, a contradiction. Let  $3 \mid r$ . Then Lemma 3.1, (ii) and (iii), yields  $q^{r/3} < 4$  so  $q^r = 3^3$ . This is a contradiction since  $4 \nmid (3^3 - 1)$ . Now  $3 \mid |\mathfrak{G}|$  so we see also that  $q \neq 3$  and thus  $q = 5$  or  $17$ . We assume that  $q^n \neq 5^2$  or  $17^2$  and derive a contradiction.

Suppose first that  $r$  is odd. We apply Lemma 3.1 with  $p = 2$ . Then  $\lambda_1 \leq 9$ ,  $\lambda_2 = 0$  so we have  $q^r < 18$ . Thus  $q^r = 5$  or  $17$  and  $r = 1$ . By Lemma 1.5, there exists  $x \in \mathfrak{B}^{\#}$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{G} = \langle 1 \rangle$ . Since  $r = 1$  and  $3 \nmid |\mathfrak{G}_x|$  we have  $|\mathfrak{G}_x| = 2$ . Hence by Lemma 1.9,  $I(\mathfrak{G}) = q^{n/2} + 1$ . Now  $\mathfrak{A}$  is central in  $\mathfrak{G}$  and cyclic so each involution of  $\mathfrak{G}/\mathfrak{A}$  corresponds to at most two noncentral involutions of  $\mathfrak{G}$ . Thus

$$q^{n/2} + 1 = I(\mathfrak{G}) \leq 2 \cdot 9 = 18$$

so  $q^n = 5^2$  or  $17^2$ , a contradiction.

Now let  $r$  be even. We have easily  $\lambda_1 \leq 9$ ,  $\lambda_2 \leq 10$ . Thus if  $k > 2$  then Lemma 3.1 (iii) yields  $q^r = 5^2$  and then by Lemma 3.1 (i)

with  $k > 2$  we have a contradiction. Thus  $k = 2$  and by Lemma 3.1 (ii),  $q^r + 1 \leq 18 + 10(q^{r/2} + 1)$  so  $q^{r/2} < 13$ . Since  $r$  is even  $q^r = 5^2$ . By Lemma 1.5 there exists  $x \in \mathfrak{B}^\#$  with  $\mathfrak{G}_x \cap \mathfrak{A}\mathfrak{C} = \langle 1 \rangle$  and hence since  $3 \nmid |\mathfrak{G}_x|$  we have  $|\mathfrak{G}_x| = 2$  or  $4$ .

Suppose  $|\mathfrak{G}_x| = 4$ . Since  $[\mathfrak{G} : \mathfrak{G}] = 2$  where  $\mathfrak{G} = C(\mathfrak{A})$  we see that  $2 \mid |\mathfrak{G}_x|$  for all  $x \in \mathfrak{B}^\#$ . Clearly  $\mathfrak{G}$  acts irreducibly on  $\mathfrak{B}$  so by Lemma 3.1 applied to  $\mathfrak{G}$  with  $p = 2$  we have  $\lambda_1 \leq 9$ ,  $\lambda_2 = 0$  so  $25 = q^r < 18$ , a contradiction. Thus  $|\mathfrak{G}_x| = 2$ .

Now here  $n = kr = 4$ . By Lemma 1.9, we have  $I(\mathfrak{G}) = 1 + q^{n/2} = 26$ . Let  $\mathfrak{L}$  be a Sylow 3-subgroup of  $\mathfrak{G}$ . Since  $3 \nmid |\mathfrak{G}_x|$ ,  $\mathfrak{L}$  is cyclic and acts semiregularly so  $|\mathfrak{L}| \mid 5^4 - 1$  and  $|\mathfrak{L}| = 3$ . Since  $3 \mid |\mathfrak{B}|$  we have  $\mathfrak{L} \cap \mathfrak{A}\mathfrak{C} = \langle 1 \rangle$ . Now  $\mathfrak{L}$  permutes by conjugation the noncentral involutions of  $\mathfrak{G}$  and since  $3 \nmid I(\mathfrak{G})$  we see that  $\mathfrak{L}$  centralizes a noncentral involution of  $\mathfrak{G}$ . The group  $\mathfrak{G}/\mathfrak{A}\mathfrak{C}$  acts on  $\mathfrak{B} = \mathfrak{C}/Z(\mathfrak{C})$ . If the action is faithful then clearly  $\mathfrak{G}/\mathfrak{A}\mathfrak{C} \cong \text{Sym}_4$ . Since subgroups of order 3 of  $\text{Sym}_4$  are self-centralizing we have a contradiction. Hence the action is not faithful so say  $\mathfrak{R}/\mathfrak{A}\mathfrak{C}$  is the kernel with  $\mathfrak{R} > \mathfrak{A}\mathfrak{C}$ . Now  $\mathfrak{G}/\mathfrak{A}\mathfrak{C}$  does act faithfully so  $[\mathfrak{R} : \mathfrak{A}\mathfrak{C}] = 2$ . Note that  $\mathfrak{R} \triangleleft \mathfrak{G}$ . Also  $3 \nmid |\mathfrak{A}|$  and  $|\mathfrak{A}| \mid 5^2 - 1$  implies  $\mathfrak{A}$  is a 2-group and hence  $\mathfrak{R}$  is a 2-group. Since  $\mathfrak{G}$  is primitive,  $\mathfrak{R}$  is of symplectic type. Moreover  $Z(\mathfrak{C}) \triangleleft \mathfrak{G}$ ,  $|Z(\mathfrak{C})| = 4$  and  $4 \mid q - 1$ . Hence  $Z(\mathfrak{C})$  is central in  $\mathfrak{G}$  so  $\mathfrak{R}$  must be the central product of a cyclic group with a nonabelian group of order 8. Now  $\mathfrak{R} \cong \mathfrak{F}$  and since  $[\mathfrak{F} : \mathfrak{A}\mathfrak{C}] \leq 2$  we have  $\mathfrak{R} = \mathfrak{F}$ . Then  $\Phi(\mathfrak{F})$  is central in  $\mathfrak{F}$  and  $\mathfrak{F} = C_{\mathfrak{F}}(\Phi(\mathfrak{F})) = \mathfrak{A}\mathfrak{C}$ , a contradiction. This completes the proof of the lemma.

LEMMA 6.5. *If  $m = 2$ , then  $q^n = 3^4$ .*

*Proof.* By the Reduction Lemma and Lemmas 4.3, 4.4 and 4.5 we have  $q = 3$  and  $\mathfrak{C} \cong \mathfrak{D}\Omega$ . Hence  $4 \nmid |\mathfrak{A}|$ . We consider  $\bar{\mathfrak{R}} = F(\mathfrak{G}/\mathfrak{A}\mathfrak{C})$ . By Lemma 6.2,  $\bar{\mathfrak{R}}_2 = O_2(\bar{\mathfrak{R}}) = \langle 1 \rangle$ . Suppose  $\bar{\mathfrak{R}}_3 = O_3(\bar{\mathfrak{R}}) \neq \langle 1 \rangle$ . Since  $q = 3$ , a Sylow 3-subgroup of  $\mathfrak{G}$  has a fixed point in  $\mathfrak{B}^\#$  and hence by half-transitivity  $\mathfrak{G}_x$  contains a Sylow 3-subgroup of  $\mathfrak{G}$  for all  $x \in \mathfrak{B}^\#$ . Let  $T$  be a noncentral involution of  $\mathfrak{C}$ . By Lemma 1.5 there exists  $x \in \mathfrak{B}^\#$  with  $\mathfrak{C} = \langle T \rangle$ . Now we can find 3-subgroup  $\mathfrak{L}$  of  $\mathfrak{G}_x$  such that  $\bar{\mathfrak{L}} = \mathfrak{L}\mathfrak{A}\mathfrak{C}/\mathfrak{A}\mathfrak{C} = \bar{\mathfrak{R}}_3$ . Since  $\mathfrak{L}$  normalizes  $\mathfrak{C} \cap \mathfrak{G}_x = \mathfrak{C}_x$  we see that  $\bar{\mathfrak{R}}_3$  centralizes the involution vector corresponding to  $T$ . Thus  $\bar{\mathfrak{R}}_3$  centralizes all the involution vectors of  $\mathfrak{B} = \mathfrak{C}/Z(\mathfrak{C})$  so by Lemma 6.3,  $\bar{\mathfrak{R}}_3 = \langle 1 \rangle$ .

Now  $\mathfrak{B} \cong Sp(4, 2)$  and  $|Sp(4, 2)| = 2^4 \cdot 3^2 \cdot 5$ . Since  $\bar{\mathfrak{R}} = O_3(\mathfrak{B})$  by the above we have  $|\bar{\mathfrak{R}}| = 1$  or  $5$  and hence  $|\mathfrak{B}| \leq 20$  and  $|\mathfrak{G}/\mathfrak{A}\mathfrak{C}| \leq 16 \cdot 20 = 320$ . We use Lemma 3.1 with  $p = 2$ . Note that  $k = n/r \geq 2^m = 4$  so Lemma 3.1 (iii) always applies. Certainly  $\lambda_2 \leq 320$ . From the



structure of  $\bar{\mathfrak{G}} = \mathfrak{G}/\mathfrak{A}$  we see that  $\lambda_1 \leq 15 + 5 \cdot 4 = 35$ . Hence

$$q^r < 2(\lambda_1 + \lambda_2) = 710 .$$

Since  $q = 3$ , this yields  $r \leq 5$ . However if  $r = 5$ , then  $[\mathfrak{G} : \mathfrak{G}]$  is odd so  $\lambda_2 = 0$  and then  $q^r < 2\lambda_1 = 70$ , a contradiction. Thus  $r \leq 4$ .

Since  $r \leq 4$  we see that  $\bar{\mathfrak{K}}$  is a Sylow 5-subgroup of  $\mathfrak{G}/\mathfrak{A}\mathfrak{C}$ . Hence if  $5 \mid |\mathfrak{G}_x|$ , then as in the preceding argument with  $\bar{\mathfrak{R}}_3$  we conclude that  $\bar{\mathfrak{K}}$  fixes all involution vectors of  $\mathfrak{B} = \mathfrak{C}/\mathfrak{Z}(\mathfrak{C})$  and thus  $\bar{\mathfrak{K}} = \langle 1 \rangle$ . This certainly contradicts  $5 \mid |\mathfrak{G}_x|$ . Hence  $5 \nmid |\mathfrak{G}_x|$ . Let  $T$  be a noncentral involution of  $\mathfrak{G}$ . We show that  $|C_{\mathfrak{B}}(T)| \leq q^{n/2}$ . This is certainly the case if  $T \in \mathfrak{A}\mathfrak{C}$ . Let  $T \in \mathfrak{G} - \mathfrak{A}\mathfrak{C}$ . Then  $|\bar{\mathfrak{K}}| = 5$  since  $\mathfrak{G}/\mathfrak{A}\mathfrak{C} \neq \langle 1 \rangle$ . Clearly there exists  $K \in \mathfrak{G}$  so that the image of  $\langle T, T^K \rangle$  in  $\mathfrak{G}/\mathfrak{A}\mathfrak{C}$  contains  $\bar{\mathfrak{K}}$ . Since  $5 \nmid |\mathfrak{G}_x|$  we see that  $C_{\mathfrak{B}}(T) \cap C_{\mathfrak{B}}(T^K) = \{0\}$ . Thus the result follows here. Finally if  $T \in \mathfrak{G} - \mathfrak{G}$ , then there exists  $A \in \mathfrak{A}$  with  $\langle T, T^A \rangle \cap \mathfrak{A} \neq \langle 1 \rangle$ . Since  $\mathfrak{A}$  acts semiregularly the result follows.

We show that  $r$  is not even. If  $r$  is even, then  $r = 2$  or  $4$ . If  $r = 2$  then  $|\mathfrak{A}| \mid q^r - 1$  and  $q^r - 1 = 8$ . Since  $4 \nmid |\mathfrak{A}|$  we have  $|\mathfrak{A}| = 2$  and  $|\mathfrak{A}| \mid q - 1$ . This violates the definition of  $r$  and hence  $r = 4$ . Here  $|\mathfrak{A}| \mid q^r - 1$  and  $q^r - 1 = 2^4 \cdot 5$  so  $|\mathfrak{A}| \mid 10$ . Since  $|\mathfrak{A}| \leq 10$  each involution of  $(\mathfrak{G} - \mathfrak{G})/\mathfrak{A}$  comes from at most 10 of  $\mathfrak{G} - \mathfrak{G}$ . Thus

$$I(\mathfrak{G}) \leq 2 \cdot 35 + 10 \cdot 320 = 3270 .$$

Since  $2 \mid |\mathfrak{G}_x|$  we have  $\mathfrak{B} = \bigcup C_{\mathfrak{B}}(T)$  over involutions  $T$  and hence

$$q^n = |\mathfrak{B}| \leq I(\mathfrak{G})q^{n/2} \leq 3270q^{n/2}$$

so  $q^{n/2} \leq 3270$ . Thus  $n < 16$ . But  $r = 4$  and  $n \geq 2^m r = 16$  so we have a contradiction. Thus  $r$  is odd.

Since  $r$  is odd, all involutions of  $\mathfrak{G}$  are contained in  $\mathfrak{G}$ . Now  $\mathfrak{A}$  is cyclic and central in  $\mathfrak{G}$  so each involution of  $\mathfrak{G}/\mathfrak{A}$  comes from at most two of  $\mathfrak{G}$ . Hence  $I(\mathfrak{G}) \leq 2 \cdot 35 = 70$  and since  $2 \mid |\mathfrak{G}_x|$  we have

$$q^n = |\mathfrak{B}| \leq I(\mathfrak{G})q^{n/2} \leq 70q^{n/2}$$

or  $q^{n/2} \leq 70$ . Since  $4 \mid n$  we have  $n = 4$  and thus  $r = 1$ . This completes the proof of the lemma.

Combining Lemmas 6.1, 6.4 and 6.5 we obtain

**THEOREM 6.6.** *Let  $\mathfrak{G}$  act faithfully on vector space  $\mathfrak{B}$  of order  $q^n$  and let  $\mathfrak{G}$  act half-transitively but not semiregularly on  $\mathfrak{B}^*$ . If  $\mathfrak{G}$  is primitive as a linear group and if  $\mathfrak{G}$  is solvable, then  $\mathfrak{G}$  satisfies one of the following.*

- (i)  $\mathfrak{G} \cong \mathcal{S}(q^n)$ .  
(ii)  $q^n = 3^2, 5^2, 7^2, 11^2, 17^2$  or  $3^4$ .

The proof of the main theorem now follows easily.

*Proof of Theorem B.* Let  $\mathfrak{G}$  be the given solvable 3/2-transitive permutation group and assume that  $\mathfrak{G}$  is not a Frobenius group. By Theorem 10.4 of [11],  $\mathfrak{G}$  is primitive. Let  $\mathfrak{B}$  be a minimal normal subgroup of  $\mathfrak{G}$ . Since  $\mathfrak{G}$  is solvable,  $\mathfrak{B}$  is elementary abelian of order  $q^n$ . Since  $\mathfrak{G}$  is primitive,  $\mathfrak{B}$  is transitive and hence regular. If  $\alpha$  is a point being permuted, then by Theorem 11.2 of [11],  $\mathfrak{G}_\alpha$  is an automorphism group of  $\mathfrak{B}$  which acts half-transitively but not semiregularly on  $\mathfrak{B}^\#$ . By Theorems 1.1 and 6.6 we have  $\mathfrak{G}_\alpha = \mathcal{S}_0(q^{n/2})$ ,  $\mathfrak{G}_\alpha \cong \mathcal{S}(q^n)$  or  $q^n = 3^2, 5^2, 7^2, 11^2, 17^2, 3^4$ . Note that the exception of Theorem 1.1 of degree  $2^6$  is a subgroup of  $\mathcal{S}(2^6)$ . Since  $\deg \mathfrak{G} = q^n$  and  $\mathfrak{G} = \mathfrak{B}\mathfrak{G}_\alpha$ , the result follows.

7. Theorem C. We can now obtain several easy corollaries.

**COROLLARY 7.1.** *Let  $\mathfrak{G}$  be a solvable 3/2-transitive permutation group. Then for all points  $\alpha \neq \beta$  the stabilizers  $\mathfrak{G}_{\alpha\beta}$  are isomorphic. In fact if  $q^n \neq 3^2$ , then  $\mathfrak{G}_{\alpha\beta}$  is cyclic, while if  $q^n = 3^2$ , then  $\mathfrak{G}_{\alpha\beta} \cong \text{Sym}_3$ .*

*Proof.* The result is clear if  $\mathfrak{G}$  is a Frobenius group,  $\mathfrak{G} \cong \mathcal{S}(q^n)$  or  $\mathfrak{G} = \mathcal{S}_0(q^n)$ . Thus we need only consider the exceptions. Here  $\mathfrak{G}_\alpha$  acts on  $\mathfrak{B}$  and  $\mathfrak{G}_{\alpha\beta}$  is the stabilizer of  $\beta \in \mathfrak{B}^\#$ . Suppose  $q^n = 5^2, 7^2, 11^2$  or  $17^2$ . Since we see easily that  $|\mathfrak{G}_\alpha|$  is prime to  $q$  it follows by complete reducibility that  $\mathfrak{G}_{\alpha\beta}$  has a faithful 1-dimensional representation and hence is cyclic. Suppose  $q^n = 3^2$ . Since  $\mathfrak{G}_\alpha \cong \mathfrak{G} \cong \mathfrak{D}$  we see that  $\mathfrak{G}_\alpha$  is transitive on  $\mathfrak{B}^\#$ . Also  $\mathfrak{G}_\alpha/\mathfrak{C} \cong \text{Sym}_3$  and  $\mathfrak{G}_{\alpha\beta} \cap \mathfrak{C} = \langle 1 \rangle$  so the result follows here. Finally let  $q^n = 3^4$  so that  $\mathfrak{C} \trianglelefteq \mathfrak{G}_\alpha$  with  $\mathfrak{C} \cong \mathfrak{D}\mathfrak{D}$ . Then  $\mathfrak{C} = O_2(\mathfrak{G}_\alpha)$ . If  $\mathfrak{C} = \mathfrak{G}_{\alpha\beta}$  then  $|\mathfrak{G}_{\alpha\beta}| = 2$ . If  $\mathfrak{G}_\alpha > \mathfrak{C}$  then as we have seen  $5 \mid |\mathfrak{G}_\alpha/\mathfrak{C}|$ . This implies that  $\mathfrak{G}_\alpha$  acts transitively on  $\mathfrak{B}^\#$ . The result now follows by Lemma 2.4 of [5].

**COROLLARY 7.2.** *Let  $\mathfrak{G}$  be a solvable linear group acting on  $GF(q)$ -vector space  $\mathfrak{B}$ . Suppose  $\mathfrak{G}$  acts half-transitively on  $\mathfrak{B}^\#$ . If  $q \neq 2$  and  $|\mathfrak{G}|$  is even, then  $\mathfrak{G}$  has a central involution.*

*Proof.* The result is well known if  $\mathfrak{G}$  acts semi-regularly and obvious in all of the remaining cases with the exception of  $\mathfrak{G} \cong \mathcal{S}(q^n)$ . Here the argument of Step 1 of the proof of Proposition 2.7 of [8] yields the result.

Finally we consider the transitive extensions of these exceptional 3/2-transitive groups.

*Proof of Theorem C.* Let  $\mathcal{G}$  be a 5/2-transitive permutation group on the set  $\Omega$  and assume that  $\mathcal{G}$  is not a Zassenhaus group. Let  $\infty, 0 \in \Omega$  and assume that  $\mathcal{G}_\infty$  is solvable. Thus  $\mathcal{G}_\infty$  is a solvable 3/2-transitive group which is not a Frobenius group. If  $\mathcal{G}_\infty \cong \mathcal{S}(q^n)$  or  $\mathcal{G}_\infty = \mathcal{S}_0(q^{n/2})$  then by the results of [8],  $\bar{\Gamma}(q^n) < \mathcal{G} \cong \Gamma(q^n)$ . Hence we need only consider the exceptional groups. We show that these have no transitive extensions.

Set  $\mathcal{H} = \mathcal{G}_{\infty 0}$  so that  $\mathcal{G}_\infty = \mathcal{H}\mathfrak{B}$  where  $\mathfrak{B}$  is a regular normal elementary abelian subgroup of order  $q^n$ . Let  $Z$  denote the central involution of  $\mathcal{H}$ . Then  $Z$  fixes 0 and  $\infty$  and moves all the rest. Since  $\mathcal{G}$  is doubly transitive we can find a suitable conjugate  $T$  of  $Z$  with  $T = ((0, \infty)) \dots$ . Thus  $T$  normalizes  $\mathcal{H}$ . By Lemma 1.3 of [8],  $|\mathcal{H}| \geq (q^n - 1)/2$ . If  $q^n = 17^2$ , then by Lemma 3.5

$$96 = |\mathcal{H}| \geq (17^2 - 1)/2 .$$

a contradiction.

We will use results of § 3 and § 4 about these exceptional groups which were not explicitly stated. Let  $\mathcal{C} = O_2(\mathcal{H})$  so that  $T$  normalizes  $\mathcal{C}$ . Suppose  $T$  fixes the point  $\alpha$ . Since  $T$  centralizes  $Z$  we see that  $(\alpha Z)T = \alpha TZ = \alpha Z$  so  $T$  also fixes  $\beta = \alpha Z$  and these must be the two points of  $\Omega$  fixed by  $T$ . Since  $T$  is conjugate to  $Z$  and  $Z$  is central in  $\mathcal{G}_{\infty 0}$  we see that  $T$  is central in  $\mathcal{G}_{\alpha\beta}$ . Thus  $T$  centralizes  $\mathcal{H}_{\alpha\beta}$ . Note that  $\mathcal{H}_{\alpha\beta} = \mathcal{H}_\alpha = \mathcal{H}_\beta$  since  $\alpha Z = \beta$ . Conversely let  $T$  centralize  $H \in \mathcal{H}$ . Then  $(\alpha H)T = \alpha TH = \alpha H$  so  $\alpha H = \alpha$  or  $\beta$ . Hence  $H \in \langle Z, \mathcal{H}_\alpha \rangle$  and hence  $C_{\mathcal{H}}(T) = \langle Z, \mathcal{H}_\alpha \rangle$ .

Suppose  $3 \mid |\mathcal{H}_x|$  for  $x \in \mathfrak{B}^*$ . This implies easily that  $q^n = 3^2$  or  $7^2$  and  $\mathcal{C} \cong \Omega$ . Since  $\mathcal{C}$  acts semiregularly on  $\mathfrak{B}^*$ ,  $C_{\mathcal{C}}(T) = \langle Z \rangle$  and thus  $T$  acts nontrivially on  $\mathcal{C}/Z(\mathcal{C})$ . Let  $\mathfrak{F}$  be a subgroup of  $\mathcal{H}_x$  of order 3. Then  $\langle T, \mathfrak{F} \rangle$  is cyclic of order 6 and acts faithfully on  $\mathcal{C}/Z(\mathcal{C})$ , a contradiction. Thus  $|\mathcal{H}_x|$  is a cyclic 2-group. Note that if  $q^n = 3^2$ , then  $3 \nmid |\mathcal{H}|$  so clearly  $\mathcal{H} \cong \mathcal{S}(3^2)$  and  $\mathcal{G}_\infty$  is not exceptional.

Set  $\mathfrak{R} = \mathcal{G} \cap \text{Alt } \Omega$ . Since  $\mathfrak{R} \cong \mathfrak{B}$ ,  $T, Z \in \mathfrak{R}$  is doubly transitive and  $\mathfrak{R}_{\infty 0}$  has a central involution. Also  $[\mathcal{G} : \mathfrak{R}] \leq 2$ . Let  $q^n = 7^2$  or  $11^2$ . Then  $|\mathcal{H}_x| \mid q - 1$  so clearly  $|\mathcal{H}_x| = 2$ . If  $H$  is a noncentral involution of  $\mathcal{H}$  then  $H$  moves  $q^2 - q$  points and hence  $H$  is a product of  $q(q - 1)/2$  transpositions. Thus with  $q = 7$  or  $11$ ,  $H \notin \mathfrak{R}$  and therefore  $\mathfrak{R}$  is a Zassenhaus group. Since  $\mathfrak{R}_{\infty 0}$  has a central involution the results of [12] yield  $\mathfrak{R} \cong \mathcal{S}(q^2)$  and hence  $\mathfrak{R}_{\infty 0}$  has a normal Sylow 3-subgroup, a contradiction. This leaves only  $q^n = 5^2$  and  $3^4$ .

Let  $q^n = 5^2$ . Suppose  $H \in \mathcal{H}$  has order 4 and fixes a point of  $\mathfrak{B}^*$ .

Since  $H$  and  $H^2$  fix the same set of points here, we see that  $H$  is a product of  $(5^2 - 5)/4 = 5$  4-cycles. Thus  $H \in \mathfrak{R}$ . Now  $\mathfrak{G}_\infty$  is exceptional so  $3 \mid |\mathfrak{G}_\infty|$  and hence by the above remarks  $|\mathfrak{R}_\infty| = 16 \cdot 3 = 48$ . Thus  $|\mathfrak{R}| = 26 \cdot 25 \cdot 48$ . Let  $\mathfrak{P}$  be a Sylow 13-subgroup of  $\mathfrak{R}$ . Then  $[\mathfrak{R} : \mathfrak{P}] = 2 \cdot 25 \cdot 48 \equiv 8 \pmod{13}$ . If  $\mathfrak{N} = N_{\mathfrak{R}}(\mathfrak{P})$ , then by Sylow's theorem,  $[\mathfrak{N} : \mathfrak{P}] \equiv 8 \pmod{13}$ . We see easily that  $\mathfrak{P}$  has two orbits of size 13. If  $\mathfrak{A}$  is an abelian subgroup of  $\mathfrak{R}$  containing  $\mathfrak{P}$ , then either  $\mathfrak{A}$  has two orbits and then  $\mathfrak{A} = \mathfrak{P}$  or  $\mathfrak{A}$  is transitive. In the latter case  $\mathfrak{A}$  is regular so if  $A \in \mathfrak{A}$  has order 2, then  $A$  is a product of 13 transpositions and  $A \in \text{Alt } \Omega$ , a contradiction. Hence  $\mathfrak{A} = \mathfrak{P}$  and  $\mathfrak{P} = C_{\mathfrak{R}}(\mathfrak{P})$ . Thus  $\mathfrak{N}/\mathfrak{P} \subseteq \text{Aut } \mathfrak{P}$  so  $[\mathfrak{N} : \mathfrak{P}] \mid 12$ . Since  $[\mathfrak{N} : \mathfrak{P}] \equiv 8 \pmod{13}$ , we have a contradiction.

Finally let  $q^n = 3^4$  so that  $\mathfrak{G}$  has degree  $3^4 + 1 = 2 \cdot 41$ . Now  $|\mathfrak{G}| \geq (q^n - 1)/2 = 40$  so we cannot have  $\mathfrak{G} \cong \mathfrak{D}\Omega$ . Hence we must have  $5 \mid |\mathfrak{G}|$  so  $\mathfrak{G}$  is transitive on  $\mathfrak{B}^2$  and we thus see easily that  $\mathfrak{R}$  is triply transitive. Now  $|\mathfrak{G}_x| = 2, 4$  or  $8$  so write  $|\mathfrak{R}_x| = 2 \cdot 2^{\delta}$  where  $2^{\delta} = 1, 2$  or  $4$ . Then

$$|\mathfrak{R}| = 82(82 - 1)(82 - 2) \cdot 2 \cdot 2^{\delta}.$$

Let  $\mathfrak{P}$  be a Sylow 41-subgroup of  $\mathfrak{R}$  so that  $[\mathfrak{R} : \mathfrak{P}] \equiv 8 \cdot 2^{\delta} \pmod{41}$ . Hence if  $\mathfrak{N} = N_{\mathfrak{R}}(\mathfrak{P})$ , then  $[\mathfrak{N} : \mathfrak{P}] \equiv 8 \cdot 2^{\delta} \pmod{41}$ . As in the  $q^n = 5^2$  case we see easily that  $\mathfrak{P}$  is self-centralizing so  $\mathfrak{N}/\mathfrak{P} \subseteq \text{Aut } \mathfrak{P}$  and  $[\mathfrak{N} : \mathfrak{P}] \mid 40$ . Since  $2^{\delta} \leq 4$  this yields  $2^{\delta} = 1$  and  $[\mathfrak{N} : \mathfrak{P}] = 8$ .

The fact that  $2^{\delta} = 1$  implies that  $\mathfrak{G} \cong \mathfrak{D}\Omega$  is normal in  $\mathfrak{R}_\infty$  and  $[\mathfrak{R}_\infty : \mathfrak{G}] = 5$ . Since  $\mathfrak{N}/\mathfrak{P}$  is cyclic, let  $\mathfrak{L} = \langle L \rangle$  be a subgroup of  $\mathfrak{N}$  of order 8.  $\mathfrak{L}$  permutes the two orbits of  $\mathfrak{P}$ . If it fixes each then  $L$  clearly has fixed points in each orbit. Thus some conjugate of  $L$  is contained in  $\mathfrak{R}_\infty$ , a contradiction since  $\mathfrak{G} \cong \mathfrak{D}\Omega$  has period 4. Thus  $\mathfrak{L}$  interchanges the two orbits. This implies easily that  $L$  is a product of ten 8-cycles and one transposition. Hence  $L$  is an odd permutation, a contradiction. This completes the proof of the theorem.

#### REFERENCES

1. D. Gorenstein, *On a theorem of Philip Hall*, Pacific J. Math. **19** (1966), 77-80.
2. P. Hall and G. Higman, *On the  $p$ -length of  $p$ -soluble groups and reduction theorems for Burnside's problem*, Proc. London Math. Soc. (3) **6** (1956), 1-42.
3. B. Huppert, *Zweifach transitive, auflösbare Permutationsgruppen*, Math. Z. **68** (1957), 126-150.
4. I. M. Isaacs and D. S. Passman, *Half transitive automorphism groups*, Canad. J. Math. **18** (1966), 1243-1250.
5. D. S. Passman, *Solvable half-transitive automorphism groups*, J. Algebra **6** (1967), 285-304.
6. ———, *Solvable 3/2-transitive permutation groups*, J. Algebra **7** (1967), 192-207.
7. ———,  *$p$ -Solvable doubly transitive permutation groups*, Pacific J. Math. **26** (1968), 555-577.

8. ———, *Some 5/2-transitive permutation groups*, Pacific J. Math. **28** (1969), 157-171.
9. P. Roquette, *Realisierung von Darstellungen endlichen nilpotenter Gruppen*, Arch. Math, **9** (1958), 241-250.
10. A. Weir, *Sylow  $p$ -subgroups of the classical groups over finite fields with characteristic prime to  $p$* , Proc. Amer. Math. Soc. **6** (1955), 529-533.
11. H. Wielandt, *Finite Permutation Groups*, Academic Press, New York, 1964.
12. H. Zassenhaus, *Kennzeichnung endlichen linearer Gruppen als Permutationsgruppen*, Abh. Math. Sem. Univ. Hamburg **11** (1936), 17-40.

Received January 23, 1968. This research partially supported by Army Contract SAR/DA-31-124-ARO (D) 336.

YALE UNIVERSITY

