

CONJUGACY IN HOUGHTON'S GROUPS

Y. ANTOLÍN, J. BURILLO, AND A. MARTINO

Abstract: Let $n \in \mathbb{N}$. Houghton's group H_n is the group of permutations of $\{1, \dots, n\} \times \mathbb{N}$, that eventually act as a translation in each copy of \mathbb{N} . We prove the solvability of the conjugacy problem and conjugator search problem for H_n , $n \geq 2$.

2010 Mathematics Subject Classification: Primary: 20F10, 20B99.

Key words: Houghton group, permutation groups, conjugacy problem.

1. Introduction

As Bourbaki intended, we let \mathbb{N} denote the set of finite cardinals, $\{0, 1, 2, \dots\}$.

Fix a positive integer n . We use X_n to denote the set $\{1, \dots, n\} \times \mathbb{N}$. Elements of X_n will be usually denoted as pairs (i, m) , $i \in \{1, \dots, n\}$, $m \in \mathbb{N}$.

Throughout, $\text{Sym} = \text{Sym}(X_n)$ will denote the group of all permutations of the set X_n , and $\text{FSym} = \text{FSym}(X_n)$ the subgroup of permutations with finite support.

In [3] Houghton introduced the group H_n of permutations of X_n that are “eventually translations” in each copy of \mathbb{N} . More precisely, an element g of Sym lies in H_n if and only if there exists a positive integer z and integers $t_1(g), \dots, t_n(g)$ such that:

$$(1) \quad (i, m)g = (i, m + t_i(g)), \quad \forall m \geq z.$$

The groups H_n , $n = 3, 4, \dots$ are known as *Houghton's groups* and they are finitely generated. This is not at all obvious from our definition, but it is clear from the original definition of Houghton, who defined H_n as the subgroups of Sym generated by g_2, \dots, g_n , where g_i acts on the first copy of \mathbb{N} by pushing elements one step to infinity and acts on the elements of the i th copy of \mathbb{N} by pushing elements one step towards the origin. See Figure 1. In cycle notation we have that

$$(2) \quad g_i = \dots (i, 2), (i, 1), (i, 0), (1, 0), (1, 1), (1, 2), \dots$$

In [4], Johnson found a presentation for H_3 . One of the main interesting properties of Houghton's groups was discovered by Brown in [1]. Brown proved that, for $n \geq 1$, H_n is FP_{n-1} but not FP_n . In particular H_1 is not finitely generated, H_2 is finitely generated but not finitely presented, and for $n > 2$, H_n is finitely presented.

Presentations of Houghton's groups and an alternative proof of Brown's results can be found in [5]. In [8], St. John-Green studied Bredon finiteness conditions for Houghton's groups. Finally, we remark that Röver [7], showed that for all $n \geq 1$, $r \geq 2$, $m \geq 1$, H_n embeds in Higman's groups $G_{r,m}$ (defined in [2]), and in particular all Houghton's groups are subgroups of Thompson's group V .

Definition 1.1. Let $\mathcal{P} = \langle S \mid R \rangle$ be a group presentation, and G the corresponding group.

The presentation has *solvable conjugacy problem* if there exists an algorithm that given two words over S , the algorithm decides whether or not these two words correspond to conjugate elements in the group.

The solvability of the conjugacy problem is in fact an algebraic property of the group, i.e. it is independent of the presentation of the group and clearly it implies the solvability of the word problem. See [6].

In this paper we investigate the conjugacy problem for Houghton's groups H_n , $n \geq 2$.

Our main theorem is the following:

Theorem 1.2. *Let $n \geq 2$. The conjugacy problem for Houghton's group H_n is solvable. Moreover, given two conjugate elements $g, h \in H_n$, there is an algorithm that finds $x \in H_n$ such that $g^x = x^{-1}gx = h$.*

As far as we know, Houghton's groups do not belong to a family for which the conjugacy problem is already known. For example, Houghton's groups are not Gromov hyperbolic or biautomatic, since by Brown's result the $K(\pi, 1)$ for H_n cannot have finitely many cells in dimensions greater than n . Also, these groups are not conjugacy separable, since they contain the infinite (finite support) alternating group, which is an infinite simple group.

The structure of the paper is as follows. In Section 2 we describe basic computations that we can do given a word representing an element of H_n , in particular, we show that the word problem is solvable and that for each $g \in H_n$, the numbers appearing in (1) are computable. In Section 3, we prove that we can decide if two elements of H_n are conjugate by an element of FSym , and in that event, find a conjugator in FSym . Finally, in Section 4, we show how to reduce our original problem, to the problem about conjugators in FSym .

2. Computations in H_n

From now on, we fix $n \in \mathbb{N}$, $n \geq 2$. We assume that $H = H_n$ and $X = X_n$. We also fix S , a finite generating set, by setting $S = \{g_2, \dots, g_n\}$ if $n \geq 3$, and $S = \{g_2, ((1,0)(2,0))\}$ if $n = 2$. Here the generators g_i are the permutations described in (2).

The next lemma is straightforward, however we include the proof to justify that all the computations in the paper can be actually done.

Lemma 2.1. *Let w be a word over $S^{\pm 1}$ and suppose that w represents $g \in H$. Let $|w|$ denote the length of w over S . Then one can compute $t(g) = (t_1(g), \dots, t_n(g))$, $(i, m)g$ for all $(i, m) \in X$. Also, one has that $|t_i(g)| < |w|$ for $i = 1, \dots, n$, and moreover*

$$(i, m)g = (i, m + t_i(g)), \quad \forall m > |w|, i \in \{1, \dots, n\}.$$

Proof: We know that the image of g_i under t is $(1, 0, \dots, 0, -1, 0, \dots, 0)$, where the -1 is in the i th coordinate. Also the image of $((1,0)(2,0))$ under t is equal to the zero vector. Thus, given any word w over $S^{\pm 1}$ representing g one can compute $t(g)$. Clearly $|t_i(g)| < |w|$.

Similarly, given that we know the action of the generators on the elements of X , we can compute $(i, m)g$ for all $(i, m) \in X$.

We prove the last claim of the lemma only in the case when $n \geq 3$ and we leave the case $n = 2$ as exercise for the reader. We argue by induction on the length of w . If $|w| = 0$, then $g = 1$ and the claim holds. For the general case, $w = w'g_j^\varepsilon$ for some $j \in \{2, \dots, n\}$, w' a word over S of length $|w| - 1$, $g_j \in S$, $\varepsilon \in \{-1, +1\}$. Let g' be the element of H represented by w' . By induction hypothesis $(i, m)g' = (i, m + t_i(g'))$, for all $i \in \{1, \dots, n\}$ and all $m > |w'|$.

Now $(i, m)g = (i, m)g'g_j^\varepsilon$ and we have two cases to consider. If $i \notin \{1, j\}$, then $(i, m)g = (i, m)g'g_j^\varepsilon = (i, m + t_i(g') + t_i(g_j^\varepsilon)) = (i, m + t_i(g'))$, and this holds for $m > |w'|$, and in particular for $m > |w|$. If $i = j$, then $(j, m)g = (j, m)g'g_j^\varepsilon = (i, m + t_i(g'))g_j^\varepsilon$. If $m > |w|$, then $m > |w'| + 1$ and in particular, $m + t_i(g) > 0$. Thus, $(j, m)g'g_j^\varepsilon = (j, m + t_j(g') - \varepsilon) = (j, m + t_j(g))$.

The case $i = 1$ is similar. □

Given an element of H as a word w over $S^{\pm 1}$ we can compute its image under t and check if it acts trivially or not on the set $\{(i, m) \in X : m \leq |w|\}$. In particular, we can decide whether the element is trivial or not. So we have proved the following.

Corollary 2.2. *The word problem for H is solvable.*

Remark 2.3. We note that the solvability of the word problem for H was already known, and can be proved using a variety of techniques. It may be of interest to observe that the approach we outline here gives a quadratic time algorithm for the word problem.

As a consequence of Lemma 2.1, it is worth noting that a number z satisfying equation (1) is computable, for instance, it is enough to take $z = |w|$, where w is a word representing g . We note that this number is not uniquely defined from equation (1). However, for our purposes it will be sufficient to find any number for which equation (1) is valid.

The support of an element of $g \in H$ is the set $\{x \in X : xg \neq x\}$, and it is denoted by $\text{supp}(g)$. Notice that, in general, elements in H do not have finite support. However, we will show that they do have finitely many orbits of length at least 2 (there are finitely many non-trivial finite orbits by definition, and Corollary 2.8 shows that there are finitely many infinite orbits).

From the computability of the $t_i(g)$ and z of equation (1) it is straightforward to determine the orbits of g . Infinite orbits of g will consist of pairs of the sets codified by arithmetic progressions except for finitely many points (where the cutoff is determined by the z from equation (1)).

We now show that there are finitely many infinite orbits, and hence g is a product of finitely many disjoint cycles.

Definition 2.4. Given an element $g \in H$, for which $t_i(g) \neq 0$, we denote by $X_{i,r}(g)$ the following subset of X :

$$\{(i, m) : m \equiv r \pmod{|t_i(g)|}\}.$$

Note that these sets depend on g , or more precisely, the numbers $t_i(g)$.

When the element $g \in H$ is clear from the context, we shall only write $X_{i,r}$ for $X_{i,r}(g)$.

Also note that for a given g there are finitely many such sets, as we can (and shall) always assume that when writing $X_{i,r}$ we have $0 \leq r < |t_i(g)|$.

One can then essentially read off the infinite orbits of g using the sets $X_{i,r}$, and therefore the infinite cycles in the cycle decomposition of g . We say “essentially” as generally an orbit which meets some $X_{i,r}$ will contain all but finitely many points of it.

Definition 2.5. We shall say that two sets, A, B , are almost equal if their symmetric difference is finite.

The following is then clear.

Proposition 2.6. *Let $g \in H$ and $(i, m) \in X$. If the set $\{(i, m)g^k : k \geq 0\}$ is infinite, then it is almost equal to $X_{i,r}$ for some i, r and $t_i(g)$ must be positive. Conversely, if $t_i(g)$ is positive, then $X_{i,r}$ is almost equal to $\{(i, m)g^k : k \geq 0\}$ for some $m \in \mathbb{N}$.*

Similarly, if $\{(j, m)g^k : k \leq 0\}$ is infinite, then it is almost equal to some $X_{j,s}$ and $t_j(g)$ is negative.

Corollary 2.7. *Every infinite orbit of $g \in H$ is almost equal to $X_{i,r} \cup X_{j,s}$, for some i, j, r, s with $i \neq j$.*

Corollary 2.8. *Given $g \in H_n$ there are exactly $\frac{1}{2}(\sum |t_i(g)|)$ distinct infinite orbits, and hence the same number of infinite cycles in the cycle decomposition of g .*

Given the results above, one can determine the infinite orbits of g and hence the infinite cycles, and from there the entire cycle decomposition of g .

Remark 2.9. Note that in writing an element of H as a product of disjoint cycles, the cycles which appear may not themselves be elements of H .

Given such a decomposition as a product of disjoint cycles, it is then straightforward to decide if two elements of H are conjugate in Sym ; they are conjugate if and only if they have the same cycle type. However, this will not be the same as conjugacy in H in general.

3. Conjugacy by elements of FSym

In this section we will prove the following.

Proposition 3.1. *There is an algorithm that, given $a, b \in H$, decides if there exists $x \in \text{FSym} \leq H$ such that $b = xax^{-1}$ and produces such an element in the case it exists.*

Before proving the proposition, we need a lemma.

Lemma 3.2. *Let $a, b \in H$. If there is $x \in \text{FSym} \leq H$ such that $b = xax^{-1}$ then*

$$(3) \quad |\text{supp}(a) - (\text{supp}(a) \cap \text{supp}(b))| = |\text{supp}(b) - (\text{supp}(a) \cap \text{supp}(b))|.$$

Proof: Notice that x restricts to a bijection on any subset of X containing $\text{supp}(x)$, which is finite. Also, x restricts to a bijection from $\text{supp}(a)$ to $\text{supp}(b)$. Thus x restricts to a bijection on the set $\text{supp}(a) \cup \text{supp}(b) \cup \text{supp}(x)$ which we express as the disjoint union of four sets: I, C_a, C_b, C_x , where $I = \text{supp}(a) \cap \text{supp}(b)$, $C_a = \text{supp}(a) - I$, $C_b = \text{supp}(b) - I$, $C_x = \text{supp}(x) - (\text{supp}(a) \cup \text{supp}(b))$.

Notice that as a and b are conjugate by $x \in \text{FSym}$, $\text{supp}(a)$ and $\text{supp}(b)$ are almost equal and $\text{supp}(x)$ is finite. Thus C_a , C_b , and C_x are finite sets. Moreover, x maps $\text{supp}(a)$ to $\text{supp}(b)$ and so maps $C_a \sqcup I$ to $C_b \sqcup I$. Since x restricts to a bijection of $C_a \sqcup C_b \sqcup C_x \sqcup I$, then x must map $C_a \sqcup C_x$ to $C_b \sqcup C_x$. Thus C_a and C_b have the same number of elements, as required. \square

Now we are ready to prove the proposition.

Proof of Proposition 3.1: First we notice that necessary conditions for a and b to be conjugate are that $t(a) = t(b)$, and by Lemma 3.2 that (3) holds. We notice that both conditions are decidable. We assume that both conditions hold.

Suppose for some $i \in \{1, \dots, n\}$, that $t_i(a) = t_i(b) < 0$. Then there exists a z such that $(i, m)a = (i, m + t_i(a))$ and $(i, m)b = (i, m + t_i(b))$ for all $m \geq z$ (this z is computable and depends only on a and b). Let $x \in \text{FSym}$ be a potential conjugator. There exists a z' such that

$$(4) \quad (i, m)x = (i, m) \text{ for all } m \geq z'.$$

Let us suppose this z' is minimal. We claim that $z' \leq z - t_i(a)$ (recall that $t_i(a)$ is negative).

To do this we argue by contradiction and suppose that $z' > z - t_i(a)$, and in particular, $z' > z$. Then for any $m \geq 0$ we have that,

$$(i, z' + m)x^{-1}ax = (i, z' + m)b = (i, z' + m + t_i(b)) = (i, z' + m + t_i(a))$$

since $z' + m$ is larger than both z and z' . On the other hand,

$$(i, z' + m)x^{-1}ax = (i, z' + m)ax = (i, z' + m + t_i(a))x,$$

and from here we deduce that, for all $m \geq 0$,

$$(i, z' + m + t_i(a))x = (i, z' + m + t_i(a))$$

which contradict the minimality of z' in (4) (recall that $t_i(a)$ is negative). A similar argument (replacing a , b by their inverses) holds when $t_i(a) > 0$. Note that any potential conjugator must send $\text{supp}(a)$ bijectively to $\text{supp}(b)$, and the argument above shows that outside of a finite set whose size only depends on a and b (can be computed from t and z), a potential conjugator is the identity. In fact, a potential conjugator x will be the identity on a computable set contained in the intersection of $\text{supp}(a)$ and $\text{supp}(b)$ which is almost equal to both $\text{supp}(a)$ and $\text{supp}(b)$. Therefore, we can decide if there exists a bijection, x_0 , from $\text{supp}(a)$ to $\text{supp}(b)$ which satisfies $x_0^{-1}ax_0 = b$ for all elements of $\text{supp}(b)$. All that remains is to decide whether such an x_0 may be extended to a bijection on the whole of X . So let us suppose that such an x_0 exists.

Note that *any* extension of x_0 to a bijection on X will be a valid conjugator in Sym , though not necessarily in H . Recall that we are assuming that (3) holds, and hence, there is a bijection between the finite sets $\text{supp}(a) - I$ and $\text{supp}(b) - I$, where $I = \text{supp}(a) \cap \text{supp}(b)$. In this case, it is clear that we can extend x_0 to a bijection on $\text{supp}(a) \cup \text{supp}(b) = \text{supp}(a) \sqcup (\text{supp}(b) - I) = \text{supp}(b) \sqcup (\text{supp}(a) - I)$, and then we can extend it by the identity on the rest of X . Such a bijection is clearly an element of FSym . \square

4. Reducing the problem to finding a conjugator in FSym

We now turn to the conjugacy problem for H . To this we consider two elements, $a, b \in H$ and try to decide whether there exists a conjugator, $x \in H$ such that $x^{-1}ax = a^x = b$.

Note that a necessary (but not sufficient) condition for conjugacy is that $t(a) = t(b)$. In this situation, the sets $X_{i,r}(a)$ and $X_{i,r}(b)$ of Definition 2.4, when defined, are the same, as these sets only depend on $t(a)$ and $t(b)$ respectively. We shall henceforth always mean these particular sets, based on a or b . Throughout this section we will use I to denote the set $\{i \in \{1, \dots, n\} \mid t_i(a) \neq 0\}$, that is the subset of branches that are not almost fixed by a (and by b).

Our first step is to reduce the problem to the case where $(X_{i,r})x$ is almost equal to $X_{i,r}$. This is equivalent to requiring that $t_i(x) \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$.

The following lemma is a useful observation.

Lemma 4.1. *Let C be a subset of G . Suppose that $C = C_0 \sqcup C_1 \sqcup \dots \sqcup C_k$ and that there exist $z_1, \dots, z_k \in G$ such that $C_i z_i = C_0$.*

Suppose that there is an algorithm that decides if given two elements g, h of G they are conjugate by an element of C_0 and finds this conjugator. Then there is an algorithm that decides if given two elements g, h of G they are conjugate by an element of C and finds this conjugator.

Proof: Set $z_0 = 1$. Given two elements g, h of G , the algorithm to decide conjugacy in C consists in running the algorithm for conjugacy in C_0 for all the pairs (g, h^{z_i}) for $i = 0, 1, \dots, k$. If there are no conjugators for any of the pairs, then there is no conjugator for g and h in C either, because if $g^x = h$ with x in some C_i , then g and h^{z_i} are conjugate by xz_i , which is in C_0 by hypothesis.

And if there is a conjugator x in C_0 for some pair (g, h^{z_i}) , namely, $x^{-1}gx = z_i^{-1}hz_i$, then g and h are conjugate by xz_i^{-1} , which is in C_i , using our hypothesis, but now written as $C_i = C_0 z_i^{-1}$. \square

Proposition 4.2. *Suppose that there is an algorithm which, given two elements $a, b \in H$, can determine whether there exists an $x \in H$ such that $a^x = b$ with $t_i(x) \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$, and find this x in case it exists.*

Then, there is an algorithm such that given two elements of H , decides if they are conjugate in H , and in the event they are, the algorithm finds a conjugator.

Proof: Let R be the set of tuples $\{r_i\}_{i \in I} \in \mathbb{Z}^{|I|}$ with $0 \leq r_i < |t_i(a)|$. Observe that $|R| < \infty$. Let $C = H$, and for $r \in R$, let $C_r = \{g \in H : t_i(g) \equiv r_i \pmod{|t_i(a)|}\}$. We denote by C_0 the set $C_{(0,0,\dots,0)}$. Given a tuple $w = (w_1, \dots, w_n) \in \mathbb{Z}^n$ and $r \in R$, it is easy to decide whether or not $w_i \equiv r_i \pmod{|t_i(a)|}$ for all $i \in I$. Hence, we can decide whether or not a given element $g \in H$ belongs to C_r .

Moreover, given $r \in R$ with $\sum r_i = 0$, we can easily construct $x_r \in H$ such that $t_i(x_r) = r_i$ for all i .

We now use Lemma 4.1. Observe that we have $C_r x_r^{-1} = C_0$ and all the hypotheses of the lemma are satisfied. This ensures the desired conclusion. \square

We shall now attempt to find an algorithm satisfying the hypotheses of Proposition 4.2. As noted, for any potential conjugator x , $(X_{i,r})x$ is almost equal to $X_{i,r}$ for all i, r . Since any infinite orbit of a is almost equal to $X_{i,r} \cup X_{j,s}$ for some i, r, j, s this means that if a and b are to be conjugate via such an x , then every infinite orbit of a must be almost equal to some infinite orbit of b and vice versa.

The key technical result is the following.

Proposition 4.3. *Let $a, b, x \in H$ and $l_1, \dots, l_n \in \mathbb{Z}$. Suppose that $a^x = b$ with $t_i(x) = l_i |t_i(a)|$ for all $i \in I$. Further suppose that $X_{i,r} \cup X_{j,s}$ is almost equal to some infinite orbit of a (and hence of b). Then there is a (computable) constant $K = K(a, b)$, depending only on a and b , such that $||l_i| - |l_j|| \leq K$.*

Proof: We may algorithmically find integers z, z' such that

$$\begin{aligned} (k, m)a &= (k, m + t_k(a)), & k = i, j, & \forall m \geq z, \\ (k, m)b &= (k, m + t_k(b)), & k = i, j, & \forall m \geq z, \\ (k, m)x &= (k, m + t_k(x)), & k = i, j, & \forall m \geq z'. \end{aligned}$$

For convenience for what follows, since we can increase arbitrarily the number z' , we shall further assume that $z' \geq z + |t_k(x)| \geq z$ for $k = i, j$, and that $z' \equiv z$ modulo both $|t_i(a)|$ and $|t_j(a)|$.

Notice that $(X_{i,r})x$ is almost equal to $X_{i,r}$ and $(X_{j,s})x$ is almost equal to $X_{j,s}$.

Let \mathcal{O}_a be the infinite orbit of a which is almost equal to $X_{i,r} \cup X_{j,s}$ and \mathcal{O}_b be the corresponding orbit for b . Then $X_{i,r} \cap \mathcal{O}_a$ contains $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, m \geq z + \epsilon\}$ for some ϵ such that $|\epsilon| \leq |t_i(a)|$ chosen such that $z + \epsilon \equiv r \pmod{|t_i(a)|}$. Similarly, $X_{i,r} \cap \mathcal{O}_b$ also contains $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, m \geq z + \epsilon\}$ for the same ϵ . We do the same for j , so that both $X_{j,s} \cap \mathcal{O}_a$ and $X_{j,s} \cap \mathcal{O}_b$ contain $\{(j, m) : m \equiv s \pmod{|t_j(a)|}, m \geq z + \delta\}$ for some number δ such that $z + \delta \equiv s \pmod{|t_j(a)|}$ and $|\delta| \leq |t_j(a)|$.

We can then write \mathcal{O}_a as the (disjoint) union of $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, m \geq z + \epsilon\}$, $\{(j, m) : m \equiv s \pmod{|t_j(a)|}, m \geq z + \delta\}$ and some finite set S . Similarly, \mathcal{O}_b is the (disjoint) union of $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, m \geq z + \epsilon\}$, $\{(j, m) : m \equiv s \pmod{|t_j(a)|}, m \geq z + \delta\}$ and some finite set T .

The hypotheses imply that x restricts to a bijection from \mathcal{O}_a to \mathcal{O}_b and eventually acts as translation at both ends, of amplitudes l_i and l_j . As $z' \equiv z \pmod{|t_i(a)|}$, we know that x restricts to a bijection from $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, m \geq z' + \epsilon\}$ to $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, m \geq z' + \epsilon + l_i|t_i(a)|\}$. Similarly, x also restricts to a bijection from $\{(j, m) : m \equiv s \pmod{|t_j(a)|}, m \geq z' + \delta\}$ to $\{(j, m) : m \equiv s \pmod{|t_j(a)|}, m \geq z' + \delta + l_j|t_j(a)|\}$.

Hence x must also restrict to a bijection between the following finite sets: $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, z' + \epsilon > m \geq z + \epsilon\} \cup \{(j, m) : m \equiv s \pmod{|t_j(a)|}, z' + \delta > m \geq z + \delta\} \cup S$ and $\{(i, m) : m \equiv r \pmod{|t_i(a)|}, z' + \epsilon + l_i|t_i(a)| > m \geq z + \epsilon\} \cup \{(j, m) : m \equiv s \pmod{|t_j(a)|}, z' + \delta + l_j|t_j(a)| > m \geq z + \delta\} \cup T$. Comparing cardinalities, we get that

$$2(z' - z) + |S| = 2(z - z') + |T| + l_i + l_j.$$

Hence, $|l_i| \leq |S| + |T| + |l_j|$ and $|l_j| \leq |S| + |T| + |l_i|$. \square

In order to exploit Proposition 4.3, we shall define the following equivalence relation.

Definition 4.4. Given $g \in H$ we define an equivalence relation \sim_g on $\{1, \dots, n\}$, as the one generated by setting $i \sim_g j$ if and only if some infinite orbit of g is almost equal to $X_{i,r} \cup X_{j,s}$, for some r, s .

Remark 4.5. This equivalence relation depends on g . However, if $g^x = h$ with $t_i(x) \equiv 0 \pmod{|t_i(g)|}$ for all i , then \sim_g and \sim_h are the same equivalence relation.

The strategy will now be to repeatedly use Proposition 4.3 to show that the translation lengths $t_i(x)$ for all $i \in I$ within an equivalence class are within bounded distance of each other. However, in order to obtain a solution to our problem, we will need to bound these lengths absolutely. Of course, we cannot obtain this bound for any x , but we can prove that if the elements are conjugate, there always is a conjugator within a computable bound. To find that, we can multiply x on the right by any element of the centraliser of g without changing its properties and we shall bound the translation lengths of x after multiplying by an appropriate element.

In order to do this, we need to define certain elements of the centraliser.

Lemma 4.6. *Let $g \in H$, with $g = \sigma_1 \dots \sigma_p$ being the expression of g as a product of disjoint cycles. Choose $i \in \{1, 2, \dots, n\}$ and let $[i]$ be its equivalence class with respect to \sim_g . Let g_i be the product of all cycles among the $\sigma_1, \dots, \sigma_p$ which are infinite and whose support is almost equal to $X_{j,r} \cup X_{k,s}$ for some $j, k \in [i]$. Then $g_i \in H$ and is in the centraliser of g .*

Proof: The fact that g_i commutes with g is clear. To see that $g_i \in H$ note that if the support of g_i meets X_j in an infinite set, then j must be equivalent to i and every infinite cycle whose support meets X_j in an infinite set appears in g_i . Hence g_i agrees with g on all but finitely many points of X_j . \square

Observe that by construction, the element g_i produced in this lemma satisfies that $t_j(g_i) = t_j(g)$ for all $j \in [i]$.

Proposition 4.7. *Let $a, b, x \in H$ and suppose that $a^x = b$ with $t_i(x) \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$. Let i_1, \dots, i_k be a set of representatives of the equivalence classes in I under the equivalence relation \sim_a (or \sim_b). Then there exists an $x' \in H$ such that $a^{x'} = b$ with $t_i(x') \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$ and $t_{i_1}(x') = \dots = t_{i_k}(x') = 0$.*

Proof: Apply Lemma 4.6 to a and to all the equivalence classes for \sim_a . For the class $[i_1]$, we construct an element a_{i_1} such that $t_j(a_{i_1}) = t_j(a)$ for all $j \in [i_1]$, and which is in the centraliser of a . Recall that the translation for x satisfies $t_{i_1}(x) = l_{i_1} t_{i_1}(a)$, so we can replace x by $xa_{i_1}^{-l_{i_1}}$, and this new element still conjugates a into b , and has t_{i_1} equal to zero.

The element x' will then be $xa_{i_1}^{-l_{i_1}} \dots a_{i_k}^{-l_{i_k}}$. \square

Corollary 4.8. *Let $a, b, x \in H$ and suppose that $a^x = b$ with $t_i(x) \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$. Then there exists an $x' \in H$ such that $a^{x'} = b$*

and such that $\sum_{i \in I} |t_i(x')|$ is bounded by a number that depends only on a and b and is computable.

Proof: Repeated use of Proposition 4.3 applied to the element produced by Proposition 4.7 will produce the desired result. Fixed an equivalence class $[i]$ of rays, there is one index $i \in [i]$ such that $t_i(x') = 0$, i.e. setting $l_i = 0$, $t_i(x') = l_i |t_i(a)|$. Then, for $j \in [i]$, $t_j(x') = l_j |t_j(a)|$, and $||l_i| - |l_j|| = |l_j| \leq K$, where K is the constant of Proposition 4.3. So the total translation $\sum_{i \in I} |t_i(x')| \leq n \cdot K \cdot M$, where $M = \max\{|t_i(a)| : i \in I\}$. \square

We need a last lemma dealing with branches outside I .

Lemma 4.9. *Let $a, b, x \in H$ and suppose that $a^x = b$ $t_i(x) \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$. Then there exists an $x' \in H$ such that $a^{x'} = b$ and such that $\sum_{i=1}^n |t_i(x')|$ is bounded by a number that depends only on a and b and is computable.*

Proof: By Corollary 4.8, we can assume that there exists a computable constant K depending only on a and b such that $\sum_{i \in I} |t_i(x)| < K$.

Let I^c denote $\{1, \dots, n\} - I = \{i_1, \dots, i_s\}$. If $s = 1$, then since $\sum_{i=1}^n t_i(x) = 0$, we get that $|t_{i_1}(x)| < K$ and hence that $\sum_{i=1}^n |t_i(x)| < 2K$ and we are done.

Otherwise, for each $j \neq 1$, define y_j to be an element of H_n such that $\text{supp}(y_j) \cap \text{supp}(a) = \emptyset$ and such that

$$\begin{aligned} t_{i_1}(y_j) &= t_{i_j}(x), \\ t_{i_j}(y_j) &= -t_{i_j}(x), \\ t_i(y_j) &= 0, \quad \text{otherwise.} \end{aligned}$$

Such an element clearly exists, since a fixes almost every point of the rays i_1 and i_j .

Now let y be the product of the y_j , and let $x' = yx$. Since each y_j commutes with a we have that,

$$a^{x'} = a^x = b.$$

Moreover, $t_i(x') = t_i(x)$ for all $i \in I$ and $t_{i_j}(x') = 0$ for $j = 2, \dots, s$. Repeating the argument for when $s = 1$, we get that $\sum_{i=1}^n |t_i(x')| < 2K$. \square

We are now ready to prove the main theorem of this note.

Proof of Theorem 1.2: By Proposition 4.2, we need to show that there is an algorithm which, given two elements $a, b \in H$, the algorithm determines whether there exists $x \in H$ such that $a^x = b$ with $t_i(x) \equiv 0$

mod $|t_i(a)|$ for all $i \in I$. By Lemma 4.9, there is a computable number N , depending on a and b such that if such conjugator x exists, then $\sum |t_i(x)| < N$.

Let S denote the finite family of tuples $(w_1, \dots, w_n) \in \mathbb{Z}^n$ satisfying $\sum |w_i| < N$, $\sum w_i = 0$, and $w_i \equiv 0 \pmod{|t_i(a)|}$ for all $i \in I$. For each tuple $s \in S$, we can produce an element $z_s \in H$ such that $t(z_s) = s$. We now let $C = \{x \in H : t_i(x) \equiv 0 \pmod{|t_i(a)|}, \sum |t_i(x)| < N\}$ and for $s \in S$, $C_s = \{x \in H : t(x) = s\}$. Set $C_0 = C_{(0,0,\dots,0)}$ and observe that $C_s z_s^{-1} = C_0$. We now use again Lemma 4.1, and we conclude the desired algorithm exists, if there exists an algorithm that can decide, for any $g, h \in H$, if there exists an $x \in H$ such that $g^x = h$ and $t_i(x) = 0$ for all i , and find x in case it exists. But observe that if all translations are zero, then $x \in \text{FSym}$ and the algorithm to find it is exactly the one provided by Proposition 3.1 in Section 3. \square

Acknowledgments. Part of this work was done at Centre de Recerca Matemàtica (CRM), Bellaterra in the Fall semester of 2012, during the research program “Automorphisms of Free Groups: Algo., Geom. and Dyn.”. The authors are grateful to CRM for its support and hospitality.

The first and second author were supported by Micinn (Spain) through project MTM2011-25955. The first author is also supported by the swiss SNF grant: FN 200020-137696/1.

References

- [1] K. S. BROWN, Finiteness properties of groups, in: “*Proceedings of the Northwestern conference on cohomology of groups*” (Evanston, Ill., 1985), *J. Pure Appl. Algebra* **44(1–3)** (1987), 45–75. DOI: 10.1016/0022-4049(87)90015-6.
- [2] G. HIGMAN, “*Finitely presented infinite simple groups*”, Notes on Pure Mathematics **8**, Department of Pure Mathematics, Department of Mathematics, I.A.S. Australian National University, Canberra, 1974.
- [3] C. H. HOUGHTON, The first cohomology of a group with permutation module coefficients, *Arch. Math. (Basel)* **31(1)** (1978/79), 254–258. DOI: 10.1007/BF01226445.
- [4] D. L. JOHNSON, Embedding some recursively presented groups, in: “*Groups St. Andrews 1997 in Bath, II*”, London Math. Soc. Lecture Note Ser. **261**, Cambridge Univ. Press, Cambridge, 1999, pp. 410–416. DOI: 10.1017/CB09780511666148.004.
- [5] S. R. LEE, Geometry of Houghton’s groups, Preprint (2012), arXiv:1212.0257.

- [6] C. F. MILLER, III, Decision problems for groups—survey and reflections, in: “*Algorithms and classification in combinatorial group theory*” (Berkeley, CA, 1989), Math. Sci. Res. Inst. Publ. **23**, Springer, New York, 1992, pp. 1–59. DOI: 10.1007/978-1-4613-9730-4-1.
- [7] C. E. RÖVER, Subgroups of finitely presented simple groups, PhD. Thesis, University of Oxford (1999).
- [8] S. ST. JOHN-GREEN, Centralisers in Houghton’s groups, Preprint (2012), arXiv:1207.1597v1.
- [9] J. WIEGOLD, Transitive groups with fixed-point-free permutations. II, *Arch. Math. (Basel)* **29(6)** (1977), 571–573. DOI: 10.1007/BF01220455.

Yago Antolín:
University of Neuchâtel
Mathematics Department
Rue Emile-Argand 11
CH-2000 Neuchâtel
Switzerland
E-mail address: yago.anpi@gmail.com
URL: <https://sites.google.com/site/yagoanpi/>

José Burillo:
Departament de Matemàtica Aplicada IV
Escola Politècnica Superior de Castelldefels
Universitat Politècnica de Catalunya
C/ Esteve Torrades 5
08860 Castelldefels, Barcelona
Spain
E-mail address: burillo@ma4.upc.edu
URL: <http://www-ma4.upc.edu/~burillo/>

Armando Martino:
School of Mathematics
University of Southampton
University Road
Southampton SO17 1BJ
UK
E-mail address: A.Martino@soton.ac.uk
URL: <http://www.personal.soton.ac.uk/am1t07/>

Primera versió rebuda el 6 de juny de 2013,
darrera versió rebuda el 13 de juny de 2014.