

Brian S. Thomson, Mathematics Department, Simon Fraser University, B.C.,  
Canada V5A 1S6. email: thomson@sfu.ca

## CHARACTERIZATIONS OF AN INDEFINITE RIEMANN INTEGRAL

Dedicated to Stefan Schwabik (1941–2009).

### Abstract

What are necessary and sufficient conditions in order that a function may be an indefinite integral in the Riemann sense? The problem has been explicitly posed in a short note [3] published by Erik Talvila in 2008 in THIS EXCHANGE. Since neither Erik nor I have been able to find a solution in the literature I propose the following solution which is the sole subject of the paper.

The easiest route to a conjecture that might work for this problem is to compare it to a similar problem solved by Riesz [2] for functions of bounded variation. In order for a function  $F : [a, b] \rightarrow \mathbb{R}$  to be represented in the form

$$F(x) = C + \int_a^x f(t) dt \quad (a \leq x \leq b)$$

for some constant  $C$  and for some function  $f$  that has bounded variation on  $[a, b]$  it is necessary and sufficient that there is a constant  $K$  so that

$$\sum_{i=1}^n \left| \frac{F(\xi_i) - F(x_{i-1})}{\xi_i - x_{i-1}} - \frac{F(x_i) - F(\xi'_i)}{x_i - \xi'_i} \right| \leq K \quad (1)$$

for every subdivision  $a = x_0 < x_1 < x_2 < \cdots < x_n = b$  and every choice of points  $x_{i-1} < \xi_i \leq \xi'_i < x_i$ . This property has been labeled *bounded slope*

---

Mathematical Reviews subject classification: Primary: 26A42, 26A16

Key words: Riemann integral, Lebesgue integral, indefinite integral, bounded slope variation

Received by the editors January 31, 2010

Communicated by: Paul Humke

*variation* and has received some attention by later authors. This is more often expressed by placing a bound on the sums

$$\sum_{i=1}^{n-1} \left| \frac{F(x_{i+1}) - F(x_i)}{x_{i+1} - x_i} - \frac{F(x_i) - F(x_{i-1})}{x_i - x_{i-1}} \right| \quad (2)$$

but the equivalent formulation here makes many computations more transparent. For details connecting the two expressions (1) and (2), see Ene [1, p. 719].

This characterization of Riesz, along with Riemann's own characterization of integrability, suggests a solution to the problem. Note that our condition (3) is easily implied by the stronger condition (1).

**Theorem 1.** *A necessary and sufficient condition in order for a function  $F : [a, b] \rightarrow \mathbb{R}$  to be representable in the form*

$$F(x) = C + \int_a^x f(t) dt \quad (a \leq x \leq b)$$

*for some constant  $C$  and for some Riemann integrable function  $f$  on  $[a, b]$  is that, for all  $\epsilon > 0$ , a positive  $\delta$  can be found so that*

$$\sum_{i=1}^n \left| \frac{F(\xi_i) - F(x_{i-1})}{\xi_i - x_{i-1}} - \frac{F(x_i) - F(\xi'_i)}{x_i - \xi'_i} \right| (x_i - x_{i-1}) < \epsilon \quad (3)$$

*for every subdivision  $a = x_0 < x_1 < x_2 < \cdots < x_n = b$  that is finer than  $\delta$  and every choice of associated points  $x_{i-1} < \xi_i \leq \xi'_i < x_i$ .*

PROOF. For the proof that the condition is necessary let us suppose that  $F$  is the indefinite integral of a Riemann integrable function  $f$ . Let  $\epsilon > 0$  and choose  $\delta > 0$  so that

$$\sum_{i=1}^n \omega_f([x_{i-1}, x_i]) (x_i - x_{i-1}) < \epsilon$$

for every subdivision  $a = x_0 < x_1 < x_2 < \cdots < x_n = b$  that is finer than  $\delta$ . Here

$$\omega_f([c, d]) = \sup\{|f(x) - f(y)| : x, y \in [c, d]\}$$

is used to denote the oscillation of the function  $f$  on a closed interval  $[c, d]$ . Since  $f$  is Riemann integrable this is possible (indeed it is one of Riemann's own characterizations of integrability).

Observe that, if  $s \leq f(x) \leq t$  on an interval  $[c, d]$ , then

$$s - t \leq \frac{F(\xi) - F(c)}{\xi - c} - \frac{F(d) - F(\xi')}{d - \xi'} \leq t - s$$

for every  $c < \xi \leq \xi' < d$ . It follows that

$$\left| \frac{F(\xi) - F(c)}{\xi - c} - \frac{F(d) - F(\xi')}{d - \xi'} \right| \leq \omega_f([c, d]).$$

Consequently, using a subdivision  $a = x_0 < x_1 < x_2 < \cdots < x_n = b$  that is finer than  $\delta$ ,

$$\begin{aligned} & \sum_{i=1}^n \left| \frac{F(\xi_i) - F(x_i)}{\xi_i - x_i} - \frac{F(x_i) - F(\xi'_i)}{x_i - \xi'_i} \right| (x_i - x_{i-1}) \\ & \leq \sum_{i=1}^n \omega_f([x_{i-1}, x_i]) (x_i - x_{i-1}) < \epsilon \end{aligned}$$

proving (3) for any choice of associated points  $x_{i-1} < \xi_i \leq \xi'_i < x_i$ .

In the opposite direction we suppose  $\epsilon > 0$  and that  $\delta > 0$  has been chosen so that the condition (3) is satisfied for such subdivisions.

First we claim that  $F$  is Lipschitz. The argument that bounded slope variation implies Lipschitz is classical (cf. [1, p. 721]); this is closely related but requires some different details. We note that  $F$  must be bounded, even continuous, otherwise the condition (3) is easily violated. Suppose then that  $|F(x)| < K$  for all  $x \in [a, b]$ .

Fix a number  $0 < t < \delta$ . We work in the interval  $[a, b - t]$ . For any  $x \in [a, b - t]$  we use the interval  $[x, x + t]$  and observe, for any  $0 < h < t/2$ , that

$$\left| \frac{F(x+h) - F(x)}{h} - \frac{F(x+t) - F(x+t/2)}{t/2} \right| (x+t-x) < \epsilon$$

because of the condition (3). Consequently

$$\left| \frac{F(x+h) - F(x)}{h} \right| < \frac{4K + \epsilon}{t}.$$

This imposes a bound on all the right-hand derived numbers of the continuous function  $F$  in the interval  $[a, b - t]$ . It follows that this bound also serves as a Lipschitz constant for  $F$  in  $[a, b - t]$ . By identical arguments, working on the left side, we can show that this same bound is a Lipschitz constant for  $F$  on the interval  $[a + t, b]$ . It follows that  $F$  is Lipschitz on  $[a, b]$ .

Since  $F$  is Lipschitz the derivative  $F'(x)$  is a bounded function that exists at all points  $x$  in a set  $D$  having full measure in  $[a, b]$  and  $F$  is an indefinite integral for  $F'$  in the Lebesgue sense. We define  $f(x) = F'(x)$  for  $x \in D$  and, at points  $x$  not in  $D$ , we write

$$f(x) = \inf_{t>0} \sup\{F'(y) : y \in D, |x - y| < t\}.$$

Certainly

$$F(x) = C + \int_a^x f(t) dt \quad (a \leq x \leq b) \quad (4)$$

for some constant  $C$ ,  $f$  is bounded and Lebesgue integrable. It remains only for us to prove that  $f$  is in fact a Riemann integrable function. To prove this we shall show that  $f$  is continuous at almost every point of  $[a, b]$ . It is enough to check that  $f$  is continuous at almost every point of the set  $D$  since the remaining points form a set of measure zero.

Let  $\omega_f(x)$  denote the oscillation of the function  $f$  at a point  $x$ ; i.e.,

$$\omega_f(x) = \inf_{t>0} \sup\{|f(x+h) - f(x)| : x+h \in [a, b], |h| < t\}.$$

The function  $f$  is continuous at a point  $x$  if and only if  $\omega_f(x) = 0$ . Thus the collection of discontinuity points of  $f$  can be expressed as the union of an increasing sequence of sets  $\{E_m\}$  where

$$E_m = \{x \in [a, b] : \omega_f(x) > 1/m\} \quad (m = 1, 2, 3, \dots).$$

We show that each  $|E_m| = 0$ ; i.e., that each is a set of Lebesgue measure zero.

For each  $x \in D \cap E_m$  we may choose a sequence of nonzero numbers  $h_n \rightarrow 0$  so that

$$|f(x + h_n) - f(x)| \geq 1/(2m).$$

By the way in which  $f$  was defined we may select these points so that  $x + h_n$  are in  $D$ .

Thus for each point  $x$  that is in  $D \cap E_m$  we may collect all the intervals of the form  $[x, y]$  or  $[y, x]$  with length smaller than  $\delta$  and for which  $y \in D$  and

$$|f(y) - f(x)| \geq 1/(2m).$$

This must form a Vitali cover of  $D \cap E_m$ .

By Vitali's theorem there is a disjoint collection  $[x_1, y_1], [x_2, y_2], \dots, [x_p, y_p]$  chosen from the cover with the property that

$$|D \cap E_m| \leq \sum_{k=1}^p (y_k - x_k) + \epsilon$$

and  $0 < y_k - x_k < \delta$  and

$$|f(y_k) - f(x_k)| \geq 1/(2m) \quad (k = 1, 2, \dots, p).$$

For each  $k = 1, 2, \dots, p$  select points  $\xi_k, \xi'_k$  with  $x_k < \xi_k \leq \xi'_k < y_k$  in such a way that

$$\left| \frac{F(\xi_k) - F(x_k)}{\xi_k - x_k} - F'(x_k) \right| < \epsilon$$

and

$$\left| \frac{F(y_k) - F(\xi'_k)}{y_k - \xi'_k} - F'(y_k) \right| < \epsilon.$$

Now observe that

$$\begin{aligned} \frac{1}{2m}(y_k - x_k) &\leq |f(y_k) - f(x_k)|(y_k - x_k) \leq \\ &\left| \frac{F(\xi_k) - F(x_k)}{\xi_k - x_k} - F'(x_k) \right| (y_k - x_k) + \left| \frac{F(y_k) - F(\xi'_k)}{y_k - \xi'_k} - F'(y_k) \right| (y_k - x_k) \\ &+ \left| \frac{F(\xi_k) - F(x_k)}{\xi_k - x_k} - \frac{F(y_k) - F(\xi'_k)}{y_k - \xi'_k} \right| (y_k - x_k). \end{aligned}$$

But

$$\sum_{k=1}^p \left| \frac{F(\xi_k) - F(x_k)}{\xi_k - x_k} - \frac{F(y_k) - F(\xi'_k)}{y_k - \xi'_k} \right| (y_k - x_k) < \epsilon$$

by the assumed condition (3). (This isn't a full subdivision of  $[a, b]$  but the sum remains smaller than  $\epsilon$ .)

The other inequalities we have imposed then show that

$$|D \cap E_m| \leq \sum_{k=1}^p (y_k - x_k) + \epsilon \leq (2m)\epsilon[2 + 2(b - a)].$$

As this argument works for any  $\epsilon > 0$  it verifies the claim that  $|D \cap E_m| = 0$  for each  $m$ . Thus the set of discontinuities of  $f$  in  $D$  have been expressed as the union of a sequence of sets of measure zero.

In particular we now know that  $f$  is continuous at almost every point of  $D$  and hence at almost every point of  $[a, b]$ . It is certainly bounded since  $F'$  is bounded by the Lipschitz constant for  $F$ . It follows that  $f$  is Riemann integrable and the representation in (4) can be interpreted in the Riemann sense.  $\square$

## References

- [1] V. Ene, *Riesz type theorems for general integrals*, Real Anal. Exchange, **22(2)** (1997), 714–733.
- [2] F. Riesz, *Sur certain systèmes singulier d'equations intégrales*, Annales de L'École Norm. Sup., Paris (3) 28 (1911), 33–62.
- [3] E. Talvila, *Characterizing integrals of Riemann integrable functions*, Real Anal. Exchange, **33(2)** (2007), 487–488.