

RAAGs in Diffeos

Sang-hyun Kim and Thomas Koberda

Abstract.

We survey embeddability results related to RAAGs (right-angled Artin groups) and various automorphism groups of manifolds. We give two different methods of embedding a RAAG to into another, and deduce that every RAAG embeds into some braid groups. This gives the unsolvability of the isomorphism problem for finitely presented subgroups of braid groups. Also, we prove that every RAAG is a quasi-isometrically embedded subgroup of the symplectomorphism groups of the disk and the sphere, given with suitable L^p metrics. Finally, we embed RAAGs in the smooth diffeomorphism group of the real line. These results reveal many closed hyperbolic manifold subgroups of diffeomorphism groups of manifolds.

§1. Right-angled Artin groups

Let Γ be a finite graph. We define the *RAAG* (*Right-Angled Artin Group*) on Γ as the group presentation

$$A(\Gamma) = \langle V(\Gamma) \mid [a, b] = 1 \text{ if } \{a, b\} \in E(\Gamma) \rangle.$$

We will also adopt an opposite notation; that is, we define

$$G(\Gamma) = \langle V(\Gamma) \mid [a, b] = 1 \text{ if } \{a, b\} \notin E(\Gamma) \rangle.$$

For example, $A(\bullet) \cong G(\bullet) \cong \mathbb{Z}$, $A(\Delta) \cong G(\{\text{three points}\}) \cong \mathbb{Z}^3$ and $A(\{\text{three points}\}) \cong G(\Delta) \cong F_3$. RAAGs are linear [27], residually torsion-free nilpotent [21] (hence bi-orderable), and have solvable word, conjugacy and isomorphism problems. We refer the readers to [15] and bibliography therein for standard facts on RAAGs.

Received January 31, 2015.

Revised July 1, 2015.

Key words and phrases. right-angled Artin group, braid group, cancellation theory, hyperbolic manifold, quasi-isometry.

Charney pointed out that RAAGs “interpolate” between free groups and free abelian groups [15]. Note that subgroups of free and free abelian groups are again free and free abelian, respectively. In general, subgroups of RAAGs enjoy strong group theoretic restrictions.

- Theorem 1.** (1) [26] *Every non-trivial subgroup of a RAAG subjects onto \mathbb{Z} .*
- (2) [1] *If the fundamental group of a closed aspherical 3-manifold M embeds into a RAAG, then M virtually fibers.*

Despite the simplicity of presentations, RAAGs are known to have strikingly rich isomorphism types of subgroups. Haglund and Wise discovered an effective way of embedding groups into RAAGs, using codimension-one subgroups [25]. A group G *virtually* embeds into another group H , if a finite-index subgroup of G embeds into H . Many word-hyperbolic groups are shown to virtually embed into RAAGs.

- Theorem 2.** (1) [20, 18] *The fundamental group of every closed hyperbolic surface with Euler characteristic not equal to -1 embeds into a RAAG.*
- (2) [2] *If a $K(\pi, 1)$ of a word-hyperbolic group G is a locally $CAT(0)$ cube complex, then G virtually embeds into a RAAG.*
- (3) [2, 40] *Every closed (Agol) or finite-volume (Wise) hyperbolic 3-manifold group virtually embeds into a RAAG.*
- (4) [8] *For each $n \geq 2$, there exists a closed hyperbolic n -manifold M such that $\pi_1(M)$ embeds into a RAAG.*

In particular, (3) in the above theorem relies on the Surface Subgroup Theorem by Kahn and Markovic [28]. Combining Theorems 1 and 2, one obtains a resolution to the long-standing Virtual Haken conjecture by Waldhausen and Virtual Fiber Conjecture by Thurston [39].

Classifying isomorphism types of subgroups in RAAGs can be quite a complicated task. For example, it is in general impossible to solve the isomorphism problem for finitely presented subgroups of a given RAAG [13]. So it is reasonable to consider a smaller class of subgroups. A question we found interesting and useful for later uses is:

Question 1. *For given finite graphs X and Y , when does $A(X)$ embed into $A(Y)$?*

On the other hand, RAAGs arise quite ubiquitously as *subgroups* of homeomorphism groups. This is due to the *universal property of RAAGs*, defined as follows. Let M be a manifold. The *support* of $f \in \text{Homeo}(M)$ is

$$\text{supp}(f) = \overline{\{x \in M : f(x) \neq x\}}.$$

Suppose $f_1, f_2, \dots, f_k \in \text{Homeo}(M)$. We let Γ be the *intersection graph* of $\{\text{supp}(f_1), \dots, \text{supp}(f_k)\}$; this means, $V(\Gamma) = \{f_1, \dots, f_k\}$ and $\{f_i, f_j\} \in E(\Gamma)$ if and only if $\text{supp}(f_i) \cap \text{supp}(f_j) \neq \emptyset$. Since the disjointness of supports imply commutativity of homeomorphisms, we have a natural group homomorphism $\phi : G(\Gamma) \rightarrow \text{Homeo}(M)$ defined as $\phi(f_i) = f_i$ for each i . In general, it is a highly non-trivial question whether or not such ϕ is injective. One can also ask the same question in different categories such as $\text{Diff}(M)$ or $\text{Diff}^k(M)$ (meaning, C^k -diffeomorphism group).

We let $\text{Mod}(S)$ denote the mapping class group of a surface S ; see [23] for fundamental facts and notations regarding this group. We denote by $\text{Symp}(S)$ the area-preserving diffeomorphism group of S , which plays an important role in fluid mechanics [3].

Question 2. *Given a surface S and a finite graph X , when does $A(X)$ embed into $\text{Mod}(S)$ or into $\text{Symp}(S)$?*

Finally, we address the question on 1-manifolds.

Question 3. *Given a one-manifold M , a finite graph X and a positive integer k , when does $A(X)$ embed into $\text{Diff}^k(M)$?*

In this paper, we will survey our results on these three questions.

§2. RAAGs in RAAGs

A motivation for Question 1 comes from the question which RAAGs contain closed hyperbolic surface subgroups [24]. It is not still known whether or not a RAAG contains a closed hyperbolic surface subgroup if and only if it contains a one-ended word-hyperbolic group; see [31, 30] for related results.

Recall that an *induced subgraph* X of a graph Y is a subgraph X of Y that satisfies $E(X) = E(Y) \cap \binom{V(X)}{2}$; we denote $X \leq Y$ in this case. It was originally suspected that $A(\Gamma)$ contains a closed hyperbolic surface subgroup if and only if Γ contains an induced 5-cycle. This suspicion was determined in the negative by showing that the RAAG on the complement graph of a cycles of length at least six contains a hyperbolic surface subgroup [32, 17, 6].

In [32] and by a different method in [6], it was proved that $G(X)$ embeds into $G(Y)$ if Y topologically contracts onto X . This can be much further generalized using the concept of *extension graph*. Let X be a finite graph. We define the *extension graph* X^e by the relations $V(X^e) = \{v^g : v \in V(\Gamma), g \in A(\Gamma)\}$ and $\{u^g, v^h\} \in E(X^e)$ iff $[u^g, v^h] = 1$ in $A(\Gamma)$. There is a natural right-action of $A(\Gamma)$ on X^e defined by $v^g.h = v^{gh}$.

The *opposite graph* X^{opp} of a graph X is defined by $V(X^{\text{opp}}) = V(X)$ and $E(X^{\text{opp}}) = \binom{V(X)}{2} \setminus E(X)$. A graph X is *anti-connected* if X^{opp} is connected. Note that every graph is a join of anti-connected graphs.

Proposition 3. [36, 35] *Let X be a finite, connected and anti-connected graph, which is not a single vertex.*

- (1) X^e is a connected, locally infinite quasi-tree of infinite diameter.
- (2) The action of $A(\Gamma)$ on X^e is acylindrical.

For the definition of *acylindricity* and related results for mapping class group actions curve complexes, see [10]. The above proposition illustrates obvious similarity between extension graphs of RAAGs and curve complexes of surfaces. Coming back to our original motivation of Question 1, let us define another concept, *clique graph*. A *clique* in a graph is a collection of vertices which are pairwise adjacent. For a possibly infinite graph Y , we define the clique graph Y_k of Y by declaring that $V(Y_k)$ is the set of cliques in Y and we join $K, L \in V(Y_k)$ if and only if $K \cup L \in V(Y_k)$.

Theorem 4. [36] *Let X and Y be finite graphs.*

- (1) If X is an induced subgraph of Y^e , then $A(X)$ embeds into $A(Y)$.
- (2) If $A(X)$ embeds into $A(Y)$, then X is an induced subgraph of $(Y^e)_k$.
- (3) If Y is triangle-free, then $X \leq Y^e$ and $A(X) \hookrightarrow A(Y)$ are equivalent.

For two groups G and H with given metrics, we say G is a *q.i. embedded subgroup* of H if there exists a group embedding $f : G \rightarrow H$ and constants $C \geq 1$ such that every $x, y \in G$ satisfies

$$\frac{1}{C}d(x, y) - C \leq d(f(x), f(y)) \leq Cd(x, y) + C.$$

We will need another type of embeddings between RAAGs. Let X be a finite graph. Consider its universal cover \tilde{X} and an arbitrary finite subtree $T \leq \tilde{X}$. We can define a map $\phi(X, T) : G(X) \rightarrow G(T)$ by

$$\phi(v) = \prod_{u \in p^{-1}(v) \cap T} u.$$

Note that $\phi(X, T)$ is well-defined, but not necessarily injective. An important observation is that it becomes injective for sufficiently large T .

Theorem 5 ([33]). *For each X , there exists a finite tree T such that $G(X)$ is a q.i. embedded subgroups of $G(T)$.*

§3. RAAGs in Mods

The embeddability of RAAGs in mapping class groups have surprising similarity with Theorem 4. For a surface S , we let $\mathcal{C}(S)$ be the curve graph of S , namely the one-skeleton of the curve complex.

Theorem 6. *Let X be a finite graph and S be a surface.*

- (1) [38] *If X is an induced subgraph of $\mathcal{C}(S)$, then $A(X)$ embeds into $\text{Mod}(S)$.*
- (2) [37] *If $A(X)$ embeds into $\text{Mod}(S)$, then X is an induced subgraph of $\mathcal{C}(S)_k$.*
- (3) [34] *If (and only if) $\mathcal{C}(S)$ is triangle-free, then $X \leq \mathcal{C}(S)$ and $A(X) \hookrightarrow \text{Mod}(S)$ are equivalent.*

In particular, if $A(X)$ embeds into $\text{Mod}(S)$, then the chromatic number of X is at most 2^N where N is the chromatic number of $\mathcal{C}(S)$. This gives a new obstruction for a RAAG from embedding into a mapping class group; compare this with the abelian rank obstruction given in [9].

Each RAAG embeds into some mapping class group. This was first seen by Crisp and Farb (unpublished), and can be deduced again from Theorem 6 (1). Regarding genus zero case, note first that we can embed $G(T)$ for each finite tree T into some planar braid group B_n [19]. Combining with Theorem 5, we have the following.

Theorem 7. *Each RAAG embeds into some braid group.*

Bridson found a RAAG which does not have a solution to the isomorphism problem for finitely presented subgroups [13]. Using the above mentioned result of Crisp and Farb, Bridson deduced that sufficiently large genus mapping class group does not have a solution to that isomorphism problem. Theorem 7 immediately implies the same for braid groups:

Corollary 8 ([33]). *The isomorphism problem for f.p. subgroups is not solvable for B_n if $n \gg 0$.*

§4. RAAGs in Diffeos

Let S be a surface with a fixed Riemannian metric. We denote by $\text{Symp}(S)$ the group of orientation- and area-preserving smooth diffeomorphisms on S . Suppose $\alpha : [0, 1] \rightarrow \text{Symp}(S)$ be a smooth isotopy on S . For $p \geq 1$, we define the L^p -length of α as the integral

$$\ell_p(\alpha) = \int_0^1 \left(\int_{x \in S} \left\| \frac{\partial \alpha_t}{\partial t} \right\|^p dx \right)^{1/p} dt.$$

This defines a right-invariant length metric d_p and a norm $\|\cdot\|_p$ on $\text{Symp}(S)$.

In [19], it was shown that if X is a finite *planar* graph then $G(X)$ is a q.i. embedded subgroup of $\text{Symp}(D^2)$ equipped with L^2 metric; see [11] for a generalization for L^p metric for $p \geq 1$. M. Kapovich proved that every RAAG embeds into $\text{Symp}(S^2)$, asking whether or not it could be q.i. embedded with respect to the L^2 metric of the latter group [29]. A remarkable inequality relating word length of a spherical braid group and the L^p metric on $\text{Symp}(S^2)$ for $p > 2$ is given in [12]. Built on these ideas and using the embedding (which happens to be quasi-isometric) found in Theorem 5 (1), we have the following results.

- Theorem 9.** (1) *Every RAAG is a q.i. embedded subgroup of $\text{Symp}(D^2)$ with the L^p metric for $p \geq 1$.*
 (2) *Every RAAG is a q.i. embedded subgroup of $\text{Symp}(S^2)$ with the L^p metric for $p > 2$.*

Let us describe an argument for (2) above. By Theorem 5, we have only to consider $G(T)$ for a tree T . Let $V(T) = \{v_1, v_2, \dots, v_k\}$. There exists a collection of simply connected subsurfaces S_1, \dots, S_k in S^2 whose intersection graph is T . Let us consider a set $P = \{m_1, m_2, \dots, m_n\}$ of marked points in S^2 , which contain at least two points from each of the regions in $S^2 \setminus \cup_i \partial S_i$.

Denote by $X_n \subseteq S^2 \times \dots \times S^2$ the configuration space of n distinct points. We let $\{D_1, D_2, \dots, D_n\}$ be a fixed collection of disjoint neighbourhoods of m_i , and put $D_0 = D_1 \times \dots \times D_n \subseteq X_n$. We define $\mathcal{P}_n \leq \text{Symp}(S^2)$ as the subgroup of diffeomorphisms that restrict to the identities on each D_i . We can choose a pseudo-Anosov diffeomorphism f_i on $S_i \setminus \cup_j D_j$ and extend f_i as the identity outside S_i and within $\cup_j D_j$. Recall we have a commutative diagram

$$\begin{array}{ccc} G(T) & \xrightarrow{\psi} & \mathcal{P}_n \\ & \searrow \phi & \downarrow f \mapsto [f] \\ & & \text{Mod}(S^2 \setminus P) \end{array}$$

where $\phi : v_i \mapsto f_i^N$. By a theorem of Clay, Leininger and Mangahas in [16], the map ϕ embeds $G(T)$ into $\text{Mod}(S \setminus P)$ as a q.i. embedded subgroup for each sufficiently large N .

It is more convenient to lift the map $\mathcal{P}_n \rightarrow \text{Mod}(S^2 \setminus P)$ with respect to the two-sheeted universal covering map $p : \text{Symp}(S^2) \rightarrow \text{Symp}(S^2)$. Namely, let us consider the quotient map from $\widetilde{\mathcal{P}}_n = p^{-1}(\mathcal{P}_n)$ to the

pure braid group $PB_n(S^2)$. For each isotopy α in $\text{Symp}(S^2)$ joining the identity to an element in \mathcal{P}_n and an n -tuple $x \in D_0$, we denote by $[\alpha(x)]$ the n -strand braid defined as the trace of x over the isotopy α . We let $\|[\alpha(x)]\|$ as the word-length in $PB_n(S^2)$. It now suffices to show the following claim.

Claim. *Let $p > 2$. There exists C such that for each isotopy $\alpha : [0, 1] \rightarrow \text{Symp}(S^2)$ with $\alpha(0) = \text{Id}$ and $\alpha(1) \in \mathcal{P}_n$, we have*

$$\|[\alpha(P)]\| \leq C\ell_p(\alpha) + C.$$

The proof of the claim is essentially given in [12], and can be summarised as follows. We may first assume that $\alpha(D_0) \subseteq \mathbb{C}^n \subseteq (S^2)^n$ possibly after disregarding a measure zero set. So we can consider the stereographic projection of $\alpha(x)$ onto \mathbb{C} for $x \in D_0$. Let $1 \leq i \neq j \leq n$. For each $x = (x_1, x_2, \dots, x_n) \in D_0$, let us define the *crossing number* $c_\alpha(x_i, x_j, \omega)$ of $\alpha(x)$ for $\omega \in S^1$ as the sum of the counts of t such that $\alpha(t)(x_i) - \alpha(t)(x_j)$ is parallel to ω . We let $c_\alpha(x_i, x_j)$ as the average of $c_\alpha(x_i, x_j, \omega)$ over $\omega \in S^1$. By the minimality of the word-length, $\sum_{i,j} c_\alpha(x_i, x_j)$ is asymptotically bounded below by $\|[\alpha(P)]\| = \|[\alpha(x)]\|$ for $x \in D_0$.

For each $x \in D_0$ and $i \neq j$, let us consider the closed curve $\gamma_\alpha(x_i, x_j; t)$ in S^1 defined by the unit vector in the direction of $\alpha(t)(x_i) - \alpha(t)(x_j)$ at each time t . By the co-area formula, $c_\alpha(x_i, x_j)$ is equal to the length of the curve $\gamma_\alpha(x_i, x_j; t)$.

The final step of the proof is a clever application of Cauchy–Schwarz [7] or Hölder inequalities [12], to show that there is C' such that for each $p > 2$

$$\int_0^1 \int_{D_i} \int_{D_j} \left| \frac{\partial \gamma_\alpha(x_i, x_j; t)}{\partial t} \right| dx_j dx_i dt \leq C' \ell_p(\alpha) + C'.$$

We finally remark that Theorem 5 is also used in the proof of the following.

Theorem 10 ([14]). *Every RAAG embeds into the area-preserving, boundary-fixing piecewise linear homeomorphism group of the I^2 .*

§5. One-dimensional manifold

The mapping class group of a pictured surface embeds into $\text{Homeo}(S^1)$ but not in $\text{Diff}^2(S^1)$ [22]. It is a famous open question whether or not mapping class groups virtually embed into $\text{Diff}^1(S^1)$ or $\text{Homeo}(\mathbb{R})$. Every RAAG A embeds into some mapping class group G ,

and once there is such an embedding, A embeds into every finite-index subgroup of G . With Baik, the authors proved that RAAGs can be given C^∞ regularity in the following sense.

Theorem 11. [4] *Every RAAG embeds into $\text{Diff}^\infty(\mathbb{R})$.*

Related to the aforementioned question on mapping class groups, we have the following result:

Theorem 12. [5] *For a closed hyperbolic surface S , the mapping class group of S does not virtually embed into $\text{Diff}^2(S^1)$.*

References

- [1] Ian Agol, *Criteria for virtual fibering*, J. Topol. **1** (2008), no. 2, 269–284. MR 2399130 (2009b:57033)
- [2] ———, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087, With an appendix by Agol, Daniel Groves, and Jason Manning. MR 3104553
- [3] Vladimir I. Arnold and Boris A. Khesin, *Topological methods in hydrodynamics*, Applied Mathematical Sciences, vol. 125, Springer-Verlag, New York, 1998. MR 1612569 (99b:58002)
- [4] Hyungryul Baik, Sang hyun Kim, and Thomas Koberda, *Right-angled Artin subgroups of the C^∞ diffeomorphism group of the real line*, Israel J. Math., to appear.
- [5] Hyungryul Baik, Sang hyun Kim, and Thomas Koberda, *Unsmoothable group actions on compact one-manifolds*, J. Eur. Math. Soc, accepted.
- [6] Robert W Bell, *Combinatorial methods for detecting surface subgroups in right-angled Artin groups*, ISRN Algebra (2011), Article ID 102029, 6 pages.
- [7] Michel Benaïm and Jean-Marc Gambaudo, *Metric properties of the group of area preserving diffeomorphisms*, Trans. Amer. Math. Soc. **353** (2001), no. 11, 4661–4672 (electronic). MR 1851187 (2002g:58010)
- [8] Nicolas Bergeron and Daniel T. Wise, *A boundary criterion for cubulation*, Amer. J. Math. **134** (2012), no. 3, 843–859. MR 2931226
- [9] J. S. Birman, A. Lubotzky, and J. McCarthy, *Abelian and solvable subgroups of the mapping class groups*, Duke Math. J. **50** (1983), no. 4, 1107–1120. MR 726319 (85k:20126)
- [10] B. H. Bowditch, *Tight geodesics in the curve complex*, Invent. Math. **171** (2008), no. 2, 281–300. MR 2367021 (2008m:57040)
- [11] Michael Brandenbursky, *Quasi-morphisms and L^p -metrics on groups of volume-preserving diffeomorphisms*, J. Topol. Anal. **4** (2012), no. 2, 255–270. MR 2949242

- [12] Michael Brandenbursky and Egor Shelukhin, *On the large-scale geometry of the l^p -metric on the symplectomorphism group of the two-sphere*, preprint, (2013).
- [13] Martin R. Bridson, *On the subgroups of right angled artin groups and mapping class groups*, To appear in *Math. Res. Lett.*
- [14] Danny Calegari and Dale Rolfsen, *Groups of PL homeomorphisms of cubes*, preprint (2014).
- [15] Ruth Charney, *An introduction to right-angled Artin groups*, *Geom. Dedicata* **125** (2007), 141–158.
- [16] Matt T. Clay, Christopher J. Leininger, and Johanna Mangahas, *The geometry of right-angled Artin subgroups of mapping class groups*, *Groups Geom. Dyn.* **6** (2012), no. 2, 249–278. MR 2914860
- [17] John Crisp, Michah Sageev, and Mark Sapir, *Surface subgroups of right-angled Artin groups*, *Internat. J. Algebra Comput.* **18** (2008), no. 3, 443–491. MR 2422070 (2009f:20054)
- [18] John Crisp and Bert Wiest, *Embeddings of graph braid and surface groups in right-angled Artin groups and braid groups*, *Algebr. Geom. Topol.* **4** (2004), 439–472. MR 2077673 (2005e:20052)
- [19] ———, *Quasi-isometrically embedded subgroups of braid and diffeomorphism groups*, *Trans. Amer. Math. Soc.* **359** (2007), no. 11, 5485–5503. MR 2327038 (2008i:20048)
- [20] Carl Droms, *Graph groups, coherence, and three-manifolds*, *J. Algebra* **106** (1987), no. 2, 484–489. MR 880971 (88e:57003)
- [21] G. Duchamp and D. Krob, *The lower central series of the free partially commutative group*, *Semigroup Forum* **45** (1992), no. 3, 385–394. MR 1179860 (93e:20047)
- [22] Benson Farb and John Franks, *Groups of homeomorphisms of one-manifolds, I: Actions of nonlinear groups*, Preprint.
- [23] Benson Farb and Dan Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, vol. 49, Princeton University Press, Princeton, NJ, 2012. MR 2850125 (2012h:57032)
- [24] C. McA. Gordon, D. D. Long, and A. W. Reid, *Surface subgroups of Coxeter and Artin groups*, *J. Pure Appl. Algebra* **189** (2004), no. 1–3, 135–148. MR 2038569 (2004k:20077)
- [25] Frédéric Haglund and Daniel T. Wise, *Special cube complexes*, *Geom. Funct. Anal.* **17** (2008), no. 5, 1551–1620. MR 2377497 (2009a:20061)
- [26] James Howie, *On locally indicable groups*, *Math. Z.* **180** (1982), no. 4, 445–461. MR 667000 (84b:20036)
- [27] Stephen P. Humphries, *On representations of Artin groups and the Tits conjecture*, *J. Algebra* **169** (1994), no. 3, 847–862. MR 1302120 (95k:20057)
- [28] J. Kahn and V. Markovic, *Immersing almost geodesic surfaces in a closed hyperbolic three manifold*, *Ann. of Math. (2)* **175** (2012), no. 3, 1127–1190. MR 2912704
- [29] Michael Kapovich, *RAAGs in Ham*, *Geom. Funct. Anal.* **22** (2012), no. 3, 733–755. MR 2972607

- [30] S. Kim and S. Oum, *Hyperbolic surface subgroups of one-ended doubles of free groups*, J. Topol. (to appear).
- [31] S. Kim and H. Wilton, *Polygonal words in free groups*, Q. J. Math. **63** (2012), no. 2, 399–421. MR 2925298
- [32] Sang-hyun Kim, *Co-contractions of graphs and right-angled Artin groups*, Algebr. Geom. Topol. **8** (2008), no. 2, 849–868. MR 2443098 (2010h:20093)
- [33] Sang-hyun Kim and Thomas Koberda, *Anti-trees and right-angled Artin subgroups of planar braid groups*, preprint.
- [34] ———, *Right-angled Artin groups and finite subgraphs of curve graphs*, preprint.
- [35] ———, *The geometry of the curve complex of a right-angled Artin group*, submitted.
- [36] ———, *Embeddability between right-angled Artin groups*, Geom. Topol. **17** (2013), no. 1, 493–530. MR 3039768
- [37] ———, *An obstruction to embedding right-angled Artin groups in mapping class groups*, Int. Math. Res. Notices (2013 (Advance Access)).
- [38] Thomas Koberda, *Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups*, Geom. Funct. Anal. **22** (2012), no. 6, 1541–1590. MR 3000498
- [39] William P. Thurston, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.) **6** (1982), no. 3, 357–381. MR 648524 (83h:57019)
- [40] Daniel T. Wise, *From riches to raags: 3-manifolds, right-angled Artin groups, and cubical geometry*, CBMS Regional Conference Series in Mathematics, vol. 117, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2012. MR 2986461

Department of Mathematical Sciences, Seoul National University, Seoul 151-747, Republic of Korea

E-mail address: `s.kim@snu.ac.kr`

Department of Mathematics, Yale University, 20 Hillhouse Ave, New Haven, CT 06520, USA

E-mail address: `thomas.koberda@gmail.com`