

As above we have

$$\| |u|^{\lambda} \|_{H^s((t_0, t_1) \times \mathbf{R}^n)} \leq C X(u)^{\lambda}.$$

Applying Theorem 9.2.2, we obtain

$$\| |u(\sigma \cdot)|^{\lambda} \|_{H^{s, 1/2}(X)} \leq C \| u(\sigma \cdot) \|_{H^{s, 1/2}(X)} \| u(\sigma \cdot) \|_{L^{\infty}(X)}^{\lambda-1}.$$

A comparison with (12.2.34) shows that

$$\Omega_0^{n/2} |u(\sigma \Omega)| \leq \frac{C}{\sigma^{n/2}} X_{s, \rho}(u),$$

$$\sigma^{n/2} \| u(\sigma \cdot) \|_{H^{s, 1/2}(X)} \leq C X_{s, \rho}(u).$$

Hence, we arrive at

$$\int_{\rho_0}^{\rho} \| |u(\sigma \cdot)|^{\lambda} \|_{H^{s, 1/2}(X)} \sigma^{n/2} d\sigma \leq C X(u)^{\lambda} \int_{\rho_0}^{\rho} \sigma^{-(\lambda-1)n/2} d\sigma.$$

Our assumption that $\lambda > 1 + 2/n$ implies that

$$\int_{\rho_0}^{\rho} \sigma^{-(\lambda-1)n/2} d\sigma < \infty$$

so we have

$$\int_{\rho_0}^{\rho} \| |u(\sigma \cdot)|^{\lambda} \|_{H^{s, 1/2}(X)} \sigma^{n/2} d\sigma \leq C X(u)^{\lambda}.$$

This observation leads to the estimate

$$\rho^{n/2} \| N(u)(\rho \cdot) \|_{H^{s, 1/2}(X)} \leq C \varepsilon + X(u)^{\lambda}.$$

This estimate and (12.2.35) give (12.2.28) and completes the proof of (12.2.5).

References

- [1] H. Bateman and A. Erdelyi, Higher transcendental functions, Vol.1 and Vol. 2, *Mc Graw-Hill Company, INC*, New York, Toronto, London, 1953 .
- [2] J. Bergh and J. Löfström, Interpolation spaces, *Springer Berlin, Heidelberg, New York*, 1976.
- [3] Ph. Brenner, $L^p - L^{p'}$ estimates for Fourier integral operators related to hyperbolic equations, *Math. Z.* **152** (1977) 273 - 286.
- [4] H. Brezis, Analyse Functionnelle - Theorie et applications, Masson Editeur, Paris, 1983.

- [5] Y.Choquet - Bruhat and D.Christodoulou, Elliptic $H_{s,\delta}$ spaces on manifolds which are Euclidean at infinity *Acta Matematica* **146** (1981) p.129 - 150.
- [6] Y. Choquet - Bruhat, C. De Witt - Morette and M. Dillard-Bleick, Analysis, Manifolds and Physics, North - Holland, Amsterdam, 1982.
- [7] R.Coifman and Y. Meyer, Au-delà des opérateurs pseudo-différentiels, *Asterisque* 57, Société de Mathématique de France 1973
- [8] D.Christodoulou, Global solutions of nonlinear hyperbolic equations for small initial data. *Comm. Pure Appl. Math.* **39** (1986), no. 2, 267–282.
- [9] H. O. CORDES, *The Technique of Pseudodifferential Operators*, London Mathematical Society Lectures Note Series, **202**, Cambridge University Press, Cambridge (1995).
- [10] I.Gelfand, G.Shilov, Generalized functions, vol I, Properties and Operations, Academic Press, 1964, New York.
- [11] I. Gelfand, M. Graev and Ya. Vilenkin, Generalized Functions, vol. V, Integral Geometry and Representation Theory, Academic Press, 1966, New York.
- [12] V.Georgiev, Une estimation avec poids pour l'équation des ondes, *C.R.Acad.Sci. Paris* **322**, Serie I (1996) 829 – 834.
- [13] V.Georgiev, Existence of global solutions to supercritical wave equations *Serdica Math. J.* **22** (1996) 125-164.
- [14] V.Georgiev, H.Lindblad and C. Sogge, Weighted Strichartz estimates and global existence for semilinear wave equations, *Amer. J. Math.* **119** (6) (1997) 1291 – 1319.
- [15] V.Georgiev and S. Lucente, Decay for nonlinear Klein – Gordon equation, NoDEA, 2004, Volume 11, Number 4, 529 - 555.
- [16] J. Ginibre and G. Velo, The global Cauchy problem for the Nonlinear Klein - Gordon Equation, *Math. Z.* **189** (1985), 487 - 505.
- [17] J.Ginibre and G.Velo, Generalized Strichartz inequalities for the wave equation, *J. Func. Anal.* **133** (1995), 50 - 68.
- [18] R. Glassey, Existence in the large for $\square u = F(u)$ in two dimensions *Math. Z.* **178** (1981) 233–261.

- [19] N.Hayashi and P. Naumkin, Asymptotics for large time of solutions to the nonlinear Schrödinger and Hartree Equations, *Amer. J.Math* **120** (1998) 369 – 391.
- [20] S. Helgason, Groups and geometrical transformations, Academic Press, New York , 1984
- [21] L. Hörmander, The analysis of linear partial differential operators, vol. I, Distribution Theory and Fourier analysis, Springer - Verlag, Berlin, New York, Tokyo 1983.
- [22] L.Hörmander, The analysis of linear partial differential operators, vol. III, Pseudodifferential Operators, Springer - Verlag, Berlin, New York, Tokyo, 1985.
- [23] L.Hörmander, On Sobolev spaces associated with some Lie algebras, Report No. 4, *Instutut Mittag Leffler*(1985)
- [24] L. Hörmander, Lectures on Nonlinear Hyperbolic Differential Equations, *Mathematiques et Applications* **26** Springer (1997).
- [25] F.John, Blow-up for quasi – linear wave equation in three space dimensions. *Comm. Pure Appl. Math.* **34** (1981), 29 – 51.
- [26] F. John, Blow-up of solutions of nonlinear wave equations in three space dimensions, *Manuscripta Math.* **28** (1979) 235–265.
- [27] T.Kato and G.Ponce, Commutator estimates and the Euler and Navier-Stokes equations, CPAM, **41** (1988) p. 891 - 907.
- [28] S. Klainerman, Global existence of small amplitude solutions to nonlinear Klein – Gordon equations in four space time dimensions, *Comm. Pure Appl. Math* **38** (1985), 631 – 641.
- [29] S. Klainerman, Remarks on the global Sobolev inequalities in the Minkowski space \mathbf{R}^{n+1} . *Comm. Pure Appl. Math* **40** (1987), 111 – 117.
- [30] H. Kubo, On the critical decay and power for semilinear wave equations in odd space dimensions *preprint*
- [31] P. Lax and R. Philips, Scattering Theory, Academic Press, New York, 1967.
- [32] H. Lindblad and C. D. Sogge, On existence and scattering with minimal regularity for semilinear wave equations *J. Funct. Anal.* **130** (1995) 357–426.

- [33] H. Lindblad and C. D. Sogge, Long - time existence for small amplitude semilinear wave equation *Amer. J. Math.* **118** (1996) 1047–1135.
- [34] H. Lindblad and C. D. Sogge, Restriction theorems and semilinear wave equations in $(1 + 3)$ -dimensions *Duke Math. J.* **85**, 1 (1996), 227-252 .
- [35] Jacques-Louis Lions, and Enrico Magenes *Problemes aux limites non homogenes et applications*. Dunod, Paris, 1968.
- [36] V. Maz'ya, Sobolev Spaces, Edition of Leningrad Univ., Leningrad, 1985.
- [37] Benjamin Muckenhoupt, *Weighted norm inequalities for the Fourier transform* *Transactions A.M.S.*, 276:729–742, 1983.
- [38] J. Peetre, Zur Interpolation von Operatorenräumen, *Arch. Math.* **21** (1970), 601 -608.
- [39] J. Peetre, On the connection between the theory of interpolation spaces and approximation theory, *Proc. Conf. Constructive Theory Func. Budapest 1969*, Akademiai Kiado, Budapest 1971, 351-363.
- [40] G.Ponce and T.Sideris, Local regularity of nonlinear wave equations in three space dimensions, *Comm. Part. Diff. Equations* **18** (1993) 169 -177.
- [41] R. Racke, Lectures on nonlinear evolution equations. Initial value problems. *Aspects of Mathematics*, Friedr. Vieweg & Sohn, Braunschweig, 1992. viii+259 pp.
- [42] M.Reed and B.Simon, Methods of Modern Mathematical Physics, vol. I, Functional Analysis, Academic Press, New York 1975.
- [43] M. Reed and B.Simon, Methods of Modern Mathematical Physics, vol. II, Fourier Analysis and Self Adjointness, 1975 Academic Press, New York, San Francisco , London .
- [44] W.Rudin, Functional analysis, McGraw-Hill Book Company, 1973, New York.
- [45] T.Runst, Mapping properties of non-linear operators in spaces of Triebel-Lizorkin and Besov type, *Anal. Math.* **12** (1986) 313-346.
- [46] T. RUNST AND W. SINCKEL, *Sobolev Spaces of Fractional Order, Nemytskij Operators and Nonlinear Partial Differential Equations*, Walter de Gruyter, Berlin (1996).

- [47] J. Schaeffer, The equation $\square u = |u|^p$ for the critical value of p , *Proc. Royal Soc. Edinburgh* **101** (1985) 31–44.
- [48] J. Shatah, M. Struwe, Geometric wave equations. *Courant Lecture Notes in Mathematics*, 2. New York University, Courant Institute of Mathematical Sciences, New York, 1998.
- [49] T. Sideris, Nonexistence of global solutions to semilinear wave equations in high dimensions *Comm. Partial Diff. Equations* **12** (1987) 378–406.
- [50] I. Sigal, Nonlinear semi – groups. *Ann. of Math.* **78** No. 2 (1963) 339 – 364.
- [51] E. Stein, Interpolation of linear operators, *Trans. Amer. Math. Soc.* **83** (1956) 482–492.
- [52] E. Stein, Singular integrals and differential properties of functions, Princeton Mathematical Series, Princeton Univ. Press, Princeton 1970.
- [53] W. Strauss, Nonlinear scattering theory, in Scattering theory in mathematical physics *Reidel Dordrect* (1979) 53– 79.
- [54] W. Strauss, Nonlinear scattering at low energy, *J. Funct. Anal.* **41** (1981) 110-133.
- [55] R. Strichartz, A priori estimates for the wave equation and some applications *J. Funct. Analysis* **5** (1970) 218–235.
- [56] R. Strichartz, Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations *Duke Math. J.* **44** (1977) 705–714.
- [57] R. Strichartz, Harmonic analysis as Spectral theory of Laplacians. *Journal of Functional Analysis* **87** (1989) 51 – 148.
- [58] R. Strichartz, Multipliers on fractional Sobolev spaces, *J. Math. and Mech.*, **16**, 9, (1967), 1031-1060.
- [59] M. Taylor, Partial Differential Equations I, Basic Theory, Springer - Verlag, 1996, New York.
- [60] M. Taylor, Pseudodifferential operators, Princeton Mathematical Series, Princeton Univ. Press, Princeton 1981.
- [61] D. Tataru, Strichartz estimates in the hyperbolic space and global existence for the semilinear wave equation, *Trans. Amer. Math. Soc.* **353** (2001), no. 2, 795–807 .

- [62] H. Triebel, Interpolation theory, functional spaces, differential operators, *VEB Deutscher Verlag der Wissenschaften*, Berlin, 1978.
- [63] W. von Wahl, L^p decay rates for homogeneous wave equation, *Math. Z.* **120** (1971) 93 – 106.
- [64] Y. Zhou, Cauchy problem for semilinear wave equations with small data in four space dimensions *J. Diff. Equations* **8** (1995) 135–144.
- [65] Eb. Zeidler, Applied Functional Analysis, Applications to Mathematical Physics, series: Applied Math. Sciences, vol. 108, (Springer - Verlag, New York) 1995.