

# The geometry of the knot concordance space

TIM D COCHRAN

SHELLY HARVEY

Most of the 50 years of study of the set of knot concordance classes,  $\mathcal{C}$ , has focused on its structure as an abelian group. Here we take a different approach, namely we study  $\mathcal{C}$  as a metric space admitting many natural geometric operators. We focus especially on the coarse geometry of satellite operators. We consider several knot concordance spaces, corresponding to different categories of concordance, and two different metrics. We establish the existence of quasi- $n$ -flats for every  $n$ , implying that  $\mathcal{C}$  admits no quasi-isometric embedding into a finite product of (Gromov) hyperbolic spaces. We show that every satellite operator is a quasihomomorphism  $P: \mathcal{C} \rightarrow \mathcal{C}$ . We show that winding number one satellite operators induce quasi-isometries with respect to the metric induced by slice genus. We prove that strong winding number one patterns induce isometric embeddings for certain metrics. By contrast, winding number zero satellite operators are bounded functions and hence quasicontractions. These results contribute to the suggestion that  $\mathcal{C}$  is a fractal space. We establish various other results about the large-scale geometry of arbitrary satellite operators.

20F65, 57M25, 57M27

## 1 Introduction

A *classical knot* is an embedded oriented  $S^1$  in  $S^3$ . We are interested in the “4-dimensional” equivalence relation on knots, called *concordance*, due to Fox and Milnor [15; 16]. Two knots,  $K_0 \hookrightarrow S^3 \times \{0\}$  and  $K_1 \hookrightarrow S^3 \times \{1\}$ , are *concordant* if there is an annulus smoothly and properly embedded in  $S^3 \times [0, 1]$  which restricts on its boundary to the given knots. Knot concordance is an important microcosm for the general study of 4-dimensional manifolds; see Casson and Freedman [6]. Our overarching goal is to discover more about the set,  $\mathcal{C}$ , of concordance classes of knots.

Most of the 50-year history of the study of  $\mathcal{C}$  has focused on its structure as an abelian group under the operation of connected sum. Here we take a different approach,

---

Note from SH: We are saddened by the loss of our coauthor, Tim D Cochran, on December 16, 2014, not long after the paper was submitted to AGT. Since he was not able to view page proofs or influence the final version, any errors in the paper should be attributed to me.

namely we study  $\mathcal{C}$  as a metric space admitting many natural geometric operators called *satellite operations*. Since the simplest example of a satellite operator is connected sum with a fixed knot, this approach can be argued to be more general than focusing on  $\mathcal{C}$  as an abelian group. In particular, we address to what extent satellite operations are isometries, quasi-isometries, self-similarities or approximate self-similarities, and whether they are quasihomomorphisms. Moreover, it was conjectured in Cochran, Harvey and Leidy [13] that  $\mathcal{C}$  is a *fractal space*. A fractal space is a metric space that admits systems of natural self-similarities, which in the weakest context (without a metric) are merely injective functions; see Bartholdi, Grigorchuk and Nekrashevych [3, Definition 3.1]. By endowing  $\mathcal{C}$  with a metric we are able to address this conjecture in a more meaningful way. The metrics we consider have certainly been considered previously. However, few have considered the metric properties of satellite operators.

We will, in fact, consider four different “concordance” equivalence relations on  $\mathcal{K}$  (corresponding to different categories), with the sets of equivalence classes being denoted by  $\mathcal{C}$ ,  $\mathcal{C}^{\text{top}}$ ,  $\mathcal{C}^{\text{ex}}$  and  $\mathcal{C}^{1/n}$ . Here  $\mathcal{C}^{\text{top}}$  is the (usual) set of *topological knot concordance* classes wherein two knots are equivalent if there exists a locally flat proper topological embedding of an annulus into  $S^3 \times [0, 1]$  which restricts on its boundary to the given knots.  $\mathcal{C}^{\text{ex}}$ , short for  $\mathcal{C}^{\text{exotic}}$ , is the set of equivalence classes of knots where two are equivalent if they cobound a properly, smoothly embedded annulus in a smooth manifold *homeomorphic* to  $S^3 \times [0, 1]$ ; that is to say they are concordant in  $S^3 \times [0, 1]$  equipped with a possibly exotic smooth structure (see Boyer [4], Sato [30, Definition 2] and Cochran, Davis and Ray [9]). If the smooth 4–dimensional Poincaré conjecture is true then  $\mathcal{C}^{\text{ex}} = \mathcal{C}$ . Finally, for a fixed nonzero integer  $n$ , let  $\mathcal{C}^{1/n}$  denote the set of equivalence classes of knots in  $S^3$  where two are equivalent if they cobound a smoothly embedded annulus in a smooth 4–manifold that is  $\mathbb{Z}[1/n]$ –homology cobordant to  $S^3 \times [0, 1]$ . For odd  $n$  it seems to be unknown whether or not  $\mathcal{C} = \mathcal{C}^{1/n}$ ! For economy we will sometimes use the notation  $\mathcal{C}^*$  to denote either  $* = \emptyset$  (ie  $\mathcal{C}$ ),  $* = \text{top}$ ,  $* = \text{ex}$  or  $* = 1/n$ . Each of these is an abelian group under connected sum, with identity the class of the trivial knot  $U$ , and where the inverse of  $K$ , denoted by  $-K$ , is the reverse of the mirror image of  $K$ , denoted by  $r\bar{K}$ . If  $K = 0 = U$  in  $\mathcal{C}$  (respectively  $\mathcal{C}^{\text{ex}}$ ,  $\mathcal{C}^{\text{top}}$ ,  $\mathcal{C}^{1/n}$ ) then  $K$  is called a (smooth) *slice knot* (respectively pseudoslice, topologically slice,  $\mathbb{Z}[1/n]$ –slice). This is equivalent to saying that  $K$  bounds a smoothly embedded disk in a manifold diffeomorphic to  $B^4$  (respectively bounds a smoothly embedded disk in an exotic  $B^4$ , bounds a locally flat, topologically embedded disk in  $B^4$ , bounds a smoothly embedded disk in a smooth manifold that is  $\mathbb{Z}[1/n]$ –homology equivalent to  $B^4$ ).

In [Section 2](#) we define a norm and a metric on a group. In [Section 3](#), we define several natural metrics on  $\mathcal{C}^*$ , each induced by a norm. To motivate these, consider two strategies to measure how far a knot is from the trivial knot, which is to say how far it is from bounding a disk in  $B^4$ . One could ask what is the least genus among all surfaces that it *does* bound in  $B^4$ . Alternatively, one could ask what is the simplest 4-manifold in which it bounds an embedded 2-disk. For us, these two types will be referred to as “slice genus norms” or “homology norms”, respectively. Specifically, on  $\mathcal{C}$  the most natural and well-studied norm is that given by the *slice genus of  $K$* ,  $\|K\|_s \equiv g_s(K)$ , which is the minimum genus among all compact oriented surfaces smoothly embedded in  $B^4$  with boundary  $K$ . The *homology norm*, denoted by  $\|-\|_H$ , is essentially the minimal  $\beta_2$  of a certain class of 4-manifolds whose boundary is the zero-framed surgery on  $K$  (see [Section 3](#) for details). The homology norm is only known to be a pseudonorm on  $\mathcal{C}$ . One of our first results is that these two really are quite different.

**Proposition 3.19** *The identity map  $i: (\mathcal{C}, d_s) \rightarrow (\mathcal{C}, d_H)$  is not a quasi-isometry.*

On  $\mathcal{C}^{\text{ex}}$  these notions yield two norms. One,  $\|-\|_{\text{ex}}$ , is given by the “slice genus” in a 4-ball with a potentially exotic smooth structure. The other is the homology norm which is a true norm on  $\mathcal{C}^{\text{ex}}$ . There are also analogues of these two metrics for the other categories  $\mathcal{C}^{\text{top}}$  and  $\mathcal{C}^{1/n}$  but we leave the precise definitions to the body of the paper.

In [Section 4](#) we show the existence of *quasi- $n$ -flats* for most of these metric spaces. Recall this means:

**Theorem 4.1** *For each  $n \geq 1$  there are subspaces of  $(\mathcal{C}, d_s)$  and  $(\mathcal{C}, d_H)$  that are quasi-isometric to  $\mathbb{R}^n$ . The same holds for  $(\mathcal{C}^{\text{ex}}, d_s^{\text{ex}})$ ,  $(\mathcal{C}^{\text{ex}}, d_H^{\text{ex}})$ ,  $(\mathcal{C}^{\text{top}}, d_s^{\text{top}})$  and  $(\mathcal{C}^{\text{top}}, d_H^{\text{top}})$ .*

It follows that  $(\mathcal{C}, d_s)$  cannot be isometrically embedded in any finite product of  $\delta$ -hyperbolic spaces. We also show that *every* infinite cyclic subgroup of  $(\mathcal{C}, d_s)$  is a quasi-1-flat if and only if the stable slice genus (defined by Livingston [\[27\]](#)) induces a norm (as opposed to a pseudonorm) on  $\mathcal{C} \otimes \mathbb{Q}$ .

The proposed self-similarities are classical *satellite operations*. Let  $P$  be a knot in a solid torus,  $K$  be a knot in  $S^3$  and  $P(K)$  be the satellite knot of  $K$  with pattern  $P$ . The algebraic intersection number of  $P$  with a cross-sectional disk of the solid torus

is called *the winding number  $w$  of  $P$* . It is known that any such operator induces a satellite operator  $P: \mathcal{C}^* \rightarrow \mathcal{C}^*$  on the various sets of concordance classes of knots. The importance of satellite operations extends beyond knot theory. Such operations have been generalized to operations on 3- and 4-manifolds, where they produce very subtle variations while fixing the homology type; see Harvey [18, Section 5.1]. In particular, winding number one satellites are closely related to Mazur 4-manifolds and Akbulut *corks*; see Akbulut and Kirby [1]. These can be used to alter the smooth structure on 4-manifolds (see for example Akbulut and Yasui [2]).

Satellite operators seem to almost never be homomorphisms, but the presence of a metric allows one to ask whether they are “close to being homomorphisms”.

**Definition 1.1** Suppose  $G$  is a group and  $A$  is an abelian group equipped with a (group) norm  $\|-\|$ . A function  $f: X \rightarrow A$  is called a *quasihomomorphism* if there exists a constant  $D_f \geq 0$ , called the *defect* of  $f$ , such that, for all  $x, y \in X$ ,

$$\|f(xy) - f(x) - f(y)\| \leq D_f.$$

**Theorem 7.1** Any satellite operator  $P: \mathcal{C} \rightarrow \mathcal{C}$  is a quasihomomorphism with respect to either the norm given by the slice genus or the pseudonorm  $d_H$ .

Moreover:

**Proposition 7.3** Suppose that  $q: \mathcal{C} \rightarrow \mathbb{R}$  is a quasihomomorphism whose absolute value gives a lower bound for some positive multiple of the slice genus. Then, for any satellite operator  $P$ , the composition  $q \circ P: \mathcal{C} \rightarrow \mathbb{R}$  is a quasihomomorphism.

There are many concordance invariants, such as Levine–Tristram signatures,  $\tau$ ,  $s$  and  $\epsilon$ , that provide such  $q$ .

There has been considerable interest in whether satellite operators are injective (especially in light of the fractal conjecture). For example, it is a famous open problem as to whether or not the Whitehead double operator is *weakly injective* (an operator is called weakly injective if  $P(K) = P(0)$  implies  $K = 0$ , where  $0$  is the class of the trivial knot); see Kirby [21, Problem 1.38] (see also Hedden and Kirk [19] for a survey). In Cochran, Harvey and Leidy [13], large classes of winding number zero operators, called “robust doubling operators” were introduced and evidence was presented for their injectivity. It was recently shown by Cochran, Davis and Ray [9, Theorem 5.1] that certain winding number  $\pm 1$  satellite operators, called *strong* winding number  $\pm 1$

operators (see [Section 5](#)), induce injective satellite operators on  $\mathcal{C}^{\text{ex}}$ ,  $\mathcal{C}^{\text{top}}$  and on  $\mathcal{C}$ , modulo the smooth 4-dimensional Poincaré conjecture; while an arbitrary (nonzero) winding number  $n$  satellite operator is injective on  $\mathcal{C}^{1/n}$ . We generalize their injectivity result to the following rather striking result with regard to the homology norm:

**Theorem 6.6** *If  $P$  is a strong winding number  $\pm 1$  pattern then*

$$P: (\mathcal{C}, d_H) \rightarrow (\mathcal{C}, d_H)$$

*preserves the pseudonorm  $d_H$  and is quasisurjective (defined just below), so that if the 4D Poincaré conjecture is true then  $P$  is a quasisurjective isometric embedding of  $\mathcal{C}$ . Moreover,*

$$P: (\mathcal{C}^*, d_H^*) \rightarrow (\mathcal{C}^*, d_H^*)$$

*is an isometric embedding for  $*$  = ex, top or  $1/n$  and is quasisurjective.*

This result, taken together with the recent result of A Levine [\[23\]](#) that some of these operators are far from surjective, means that these operators and their iterates map the concordance space isometrically onto proper subspaces. This establishes that  $(\mathcal{C}^{\text{ex}}, d_H)$  is a fractal metric space, as is  $(\mathcal{C}, d_H)$  if the 4D Poincaré conjecture is true.

What can be said using the very important slice genus norm? Recall:

**Definition 1.2** If  $(X, d)$  and  $(Y, d')$  are metric spaces then a function  $f: X \rightarrow Y$  is a *quasi-isometry* if there are constants  $A \geq 1$ ,  $B \geq 0$  and  $C \geq 0$  such that

$$(1-1) \quad \frac{1}{A}d(x, y) - B \leq d'(f(x), f(y)) \leq A d(x, y) + B$$

and, for every  $z \in Y$ , there exists some  $x \in X$  such that

$$(1-2) \quad d'(z, f(x)) \leq C.$$

A function that satisfies the second condition is called *quasisurjective*. A function that satisfies only the first condition is called a *quasi-isometric embedding of  $X$  into  $Y$* . Note the definitions make sense even if one has only pseudometrics.

**Theorem 6.5** *If  $P$  is a winding number  $\pm 1$  pattern then  $P: \mathcal{C} \rightarrow \mathcal{C}$  is a quasi-isometry with respect to the metric  $d_s$ . In fact,  $P$  is a quasi-isometry for each of the metrics we discuss on each  $\mathcal{C}^*$ .*

By contrast, we show that any winding number zero operator is an approximate contraction. This follows from the much stronger:

**Proposition 6.11** Any winding number zero satellite operator on  $(C^*, d_*)$  is a bounded function, where  $(C^*, d_*)$  is any of the metric spaces we define.

Certain of our results follow from the basic result that any winding number  $n$  satellite operator is within a bounded distance of the  $(n, 1)$ -cable operator, which leads to:

**Proposition 6.14** Suppose  $m \neq \pm n$ . Then no winding number  $m$  satellite operator is within a bounded distance of any winding number  $n$  satellite operator.

There are many other norms on  $\mathcal{C}$  that we will not consider. For example there is the minimal number of crossing changes necessary to change a knot to a slice knot (sometimes called the *slicing number*; see Livingston [25]). There is the smallest 3-genus among all knots in the concordance class of  $K$  (called the *concordance genus* of  $K$ ); see Livingston [26]. The *stable 4-genus* is an interesting pseudonorm; see Livingston [27]. In particular, the latter paper contains some significant calculations and estimations of the slice genera for certain families of knots.

## Acknowledgments

We gratefully acknowledge the helpful advice and questions of our colleague Andy Putman, the comments of Ian Agol which led to Proposition 4.3, as well as the helpful comments given by the referee.

Cochran was partially supported by the National Science Foundation DMS-1309081 and the Simons Foundation. Harvey was partially supported by the National Science Foundation DMS-0748458 (CAREER), DMS-1309070 and the Simons Foundation.

## 2 Norms and metrics on groups

**Definition 2.1** A *metric* on a set  $X$  is a function  $d: X \times X \rightarrow \mathbb{R}$  such that, for all  $x, y, z \in X$ :

- (M1) **Positivity**  $d(x, y) \geq 0$ , and  $d(x, y) = 0$  if and only if  $x = y$ .
- (M2) **Triangle inequality**  $d(x, z) \leq d(x, y) + d(y, z)$ .
- (M3) **Symmetry**  $d(x, y) = d(y, x)$ .

A function that only satisfies the first part of (M1) is called a *pseudometric*.

If the set  $X$  has a group structure then often metrics are induced from *group norms*.

**Definition 2.2** A *norm* on a group  $G$  is a function  $\|-\|: G \rightarrow \mathbb{R}$  such that, for all  $x, y \in G$ :

(GN1) **Positivity**  $\|x\| \geq 0$ , and  $\|x\| = 0$  if and only if  $x = e$ .

(GN2) **Subadditivity**  $\|xy\| \leq \|x\| + \|y\|$ .

(GN3) **Symmetry**  $\|x^{-1}\| = \|x\|$ .

A function that satisfies only the first part of (GN1) is called a *pseudonorm* or a *seminorm*.

Beware that, even for abelian groups  $G$ , we do *not* have the usual property (for normed vector spaces):  $\|nx\| = n\|x\|$  for  $n \in \mathbb{Z}$ . This is assumed only when  $n$  is a unit, as in (GN3).

The following propositions are easy to verify and thus we omit the proofs.

**Proposition 2.3** If  $\|-\|$  is a norm (respectively pseudonorm) on the group  $G$  then  $d(x, y) = \|xy^{-1}\|$  is a metric (respectively pseudometric) on the underlying set  $G$ , called the **metric induced by the norm**  $\|-\|$ . This metric is **right-invariant**, meaning that  $d(xg, yg) = d(x, y)$  for all  $g \in G$ .

**Proposition 2.4** If the metric  $d$  is induced from a group norm  $\|-\|$  on  $G$  then

$$-d(x, y) \leq \|x\| - \|y\| \leq d(x, y).$$

The reader may easily verify the following, which will not be used in the paper.

**Proposition 2.5** A metric on the underlying set of a group  $G$  is induced from a group norm on  $G$  if and only if it is right-invariant.

### 3 Norms and metrics on concordance classes of knots

In this section, we will define various pseudonorms on  $\mathcal{K}$ , the monoid of ambient isotopy classes of knots. These will induce norms (and metrics) on the four different concordance groups,  $\mathcal{C}$ ,  $\mathcal{C}^{\text{ex}}$ ,  $\mathcal{C}^{\text{top}}$  and  $\mathcal{C}^{1/n}$ . We will abuse notation and use  $K$  for both the knot and its concordance class. We then discuss what is known about how these pseudometrics compare. In particular, we establish that the slice genus norm and the homology norm are quite different.

### 3.1 The slice genus norm and the homology pseudonorm on $\mathcal{C}$

**Definition 3.1** The *slice genus* of  $K$ , denoted by  $\|K\|_s$  or  $g_s(K)$ , is the minimum  $g$  such that  $K$  is the boundary of a smoothly embedded compact oriented surface of genus  $g$  in  $B^4$ .

The following is well known.

**Proposition 3.2**  $\|-\|_s$  is a norm on the group  $\mathcal{C}$  and so  $d_s(K, J) = \|K - J\|_s$  is a metric on the set  $\mathcal{C}$ .

It is also easy to see that  $d_s$  has an alternative definition:

**Proposition 3.3**  $d_s(K, J)$  is equal to the minimal genus of a compact oriented surface properly embedded in  $S^3 \times [0, 1]$ , whose boundary is the disjoint union of  $K \hookrightarrow S^3 \times \{0\}$  and  $-J \hookrightarrow -S^3 \times \{1\}$ .

**Proof** Suppose  $\Sigma$  is such a surface in  $S^3 \times [0, 1]$ . Choose an arc on  $\Sigma$  going from  $K$  to  $J$ . Deleting a neighborhood of this arc leaves a 4-ball containing an embedded surface  $\Sigma'$  of the same genus, whose boundary is  $K \# -J$ . Hence,  $d_s(K, J) \leq \text{genus}(\Sigma)$ . This process is reversible, so the other inequality follows.  $\square$

Next we define the so-called *homology norm* on  $\mathcal{C}$ , which is only known to be a pseudonorm.

**Definition 3.4**  $\|K\|_H$ , the *homology norm* of  $K$ , has two equivalent definitions:

- (1)  $\|K\|_H$  is the minimum of  $\frac{1}{2}(\beta_2(V) + |\sigma(V)|)$ , where  $V$  ranges over all smooth, oriented, compact, simply connected 4-manifolds with  $\partial V = S^3$  in which  $K$  is slice, that is, in which  $K$  bounds a smoothly embedded disk that represents 0 in  $H_2(V, \partial V)$  (compare [31]).
- (2)  $\|K\|_H$  is the minimum of  $\frac{1}{2}(\beta_2(W) + |\sigma(W)|)$ , where  $W$  ranges over all smooth, compact, oriented 4-manifolds  $W$  whose boundary is  $M_K$  (zero-framed surgery of  $S^3$  along  $K$ ), whose  $\pi_1$  is normally generated by the meridian of  $K$ , and for which  $H_1(M_K) \rightarrow H_1(W)$  is an isomorphism.

We sketch the proof of the equivalence. Suppose that  $K = \partial \Delta$  is slice in a 4-manifold  $V$  satisfying the first part of Definition 3.4 with  $\frac{1}{2}(\beta_2(V) + |\sigma(V)|) = n$ . Let  $W$  be the exterior in  $V$  of a tubular neighborhood of  $\Delta$ . Then  $\partial W = M_K$  and one can easily check



that  $W$  has the properties of part two of [Definition 3.4](#) and  $\frac{1}{2}(\beta_2(W) + |\sigma(W)|) = n$ . For the other direction, one simply adds a 2-handle to  $W$  along the meridian of  $K$  to arrive at a suitable  $V$ .

**Remark** It might seem more natural to the reader to minimize the quantity  $\beta_2(V)$  and indeed this gives another pseudometric  $d_{H'}$ . But the inequalities

$$\frac{1}{2}\beta_2(V) \leq \frac{1}{2}(\beta_2(V) + |\sigma(V)|) \leq \beta_2(V)$$

show that these metrics are quite closely related. For example, it is easy to show that the identity map is a quasi-isometry  $(\mathcal{C}, d_H) \rightarrow (\mathcal{C}, d_{H'})$ , and that [Theorem 6.6](#) holds for  $d_{H'}$ . Our choice has the property that it gives a better approximation to  $d_s$  (see [Proposition 3.20](#) below).

**Proposition 3.5**  $\|-\|_H$  is a pseudonorm on the group  $\mathcal{C}$  and so  $d_H(K, J) = \|K - J\|_H$  is a pseudometric on  $\mathcal{C}$ . If the 4-dimensional smooth Poincaré conjecture is true then  $\|-\|_H$  is a norm on  $\mathcal{C}$  and  $d_H(K, J)$  is a metric on  $\mathcal{C}$ .

**Proof** First we establish (GN1). Clearly  $\|K\|_H \geq 0$ , with equality if and only if  $K$  is slice in a contractible manifold, which is, by the work of Freedman, a possibly exotic 4-ball. It follows that if the 4-dimensional Poincaré is true then  $K = 0$  in  $\mathcal{C}$ .

Next, suppose  $K$  and  $J$  are slice in 4-manifolds  $V_K$  and  $V_J$  satisfying the first part of [Definition 3.4](#) with  $\frac{1}{2}(\beta_2(V_K) + |\sigma(V_K)|) = \|K\|_H$  and  $\frac{1}{2}(\beta_2(V_J) + |\sigma(V_J)|) = \|J\|_H$ . Then  $K \# J$  is slice in the boundary connected sum of  $V_K$  and  $V_J$ . Hence,  $\|K \# J\|_H \leq \|K\|_H + \|J\|_H$ , establishing (GN2).

The homology norm of  $K$  is clearly the same as that of its mirror image (by reversing the orientation of  $W$ ) and that of its reverse. Thus, (GN3) holds.  $\square$

As for  $d_s$ , the metric  $d_H$  has an alternative definition, whose proof we leave to the reader:

**Proposition 3.6**  $d_H(K, J)$  is equal to the minimum of  $\frac{1}{2}(\beta_2(Z) + |\sigma(Z)|)$ , where  $Z$  ranges over all smooth, oriented, compact, simply connected 4-manifolds with  $\partial Z = S^3 \sqcup -S^3$  in which  $K$  and  $J$  are concordant, that is, in which  $K \hookrightarrow S^3 \times 0$  and  $-J \hookrightarrow -S^3 \times 1$  cobound a null-homologous smoothly embedded annulus.

We will need the following:

**Proposition 3.7** *If the zero framed surgeries  $M_J$  and  $M_K$  are homology cobordant via a cobordism whose fundamental group is normally generated by either meridian, then  $\|K\|_H = \|J\|_H$ .*

**Proof** Suppose  $C$  is the homology cobordism and  $W_K$  is a manifold with boundary  $M_K$  which realizes the minimum  $\frac{1}{2}(\beta_2(W_K) + |\sigma(W_K)|)$  as in the second version of Definition 3.4. Then let  $W_J = W_K \cup C$ . Then  $\partial(W_J) = M_J$  and  $H_*(W_K) \cong H_*(W_J)$ , so  $\|J\|_H \leq \|K\|_H$ . The result follows by symmetry.  $\square$

### 3.2 The exotic slice genus norm and the homology norm on $\mathcal{C}^{\text{ex}}$

We will discuss two norms on  $\mathcal{C}^{\text{ex}}$ , defined as above.

**Definition 3.8** The *exotic slice genus* of  $K$ , denoted by  $\|K\|_s^{\text{ex}}$  or by  $g_{\text{ex}}(K)$ , is the minimum  $n$  such that  $K$  is the boundary of a smoothly embedded, compact, oriented surface of genus  $n$  in a smooth manifold that is homeomorphic to  $B^4$ .

If the 4-dimensional smooth Poincaré conjecture is true then  $\|-\|_s^{\text{ex}} = \|-\|_s$

**Proposition 3.9**  $\|-\|_s^{\text{ex}}$  is a norm on the group  $\mathcal{C}^{\text{ex}}$  and so  $d_s^{\text{ex}}(K, J) = \|K - J\|_s^{\text{ex}}$  is a metric on the set  $\mathcal{C}^{\text{ex}}$ .

**Proof** Clearly  $\|K\|_s^{\text{ex}} \geq 0$ , with equality if and only if  $K$  is slice in a smooth 4-manifold that is homeomorphic to  $B^4$ . This is equivalent to  $K = 0$  in  $\mathcal{C}^{\text{ex}}$ . This establishes (GN1).

Suppose  $K_1$  and  $K_2$  bound embedded surfaces  $F_1$  and  $F_2$  of genera  $n_1$  and  $n_2$  in 4-manifolds  $\mathcal{B}_1$  and  $\mathcal{B}_2$ , respectively, that are homeomorphic to  $B^4$ . Then  $K_1 \# K_2$  bounds in the boundary connected sum  $\mathcal{B}_1 \natural \mathcal{B}_2$ , the boundary connected sum of  $F_1$  and  $F_2$ . The latter has genus  $n_1 + n_2$ . This establishes (GN2).

Since the exotic slice genus of  $K$  is clearly the same as that of its mirror image and that of its reverse, (GN3) holds.  $\square$

The metric  $d_s^{\text{ex}}$  has an alternative definition, whose verification is the same as that of Proposition 3.3.

**Proposition 3.10**  $d_s^{\text{ex}}(K, J)$  is equal to the minimal genus of a compact oriented surface properly embedded in a smooth 4-manifold homeomorphic to  $S^3 \times [0, 1]$ , whose boundary is the disjoint union of  $K \hookrightarrow S^3 \times \{0\}$  and  $-J \hookrightarrow S^3 \times \{1\}$ .

**Proposition 3.11** *The homology norm, as defined in Definition 3.4 induces a norm on the group  $\mathcal{C}^{\text{ex}}$  and so  $d_H(K, J) = \|K - J\|_H$  induces a metric on  $\mathcal{C}^{\text{ex}}$ .*

**Proof** First we establish (GN1). Clearly  $\|K\|_H \geq 0$ , with equality if and only if  $K$  is slice in a possibly exotic 4-ball, so  $K = 0$  in  $\mathcal{C}^{\text{ex}}$ . Otherwise the proof is identical to that of Proposition 3.5.  $\square$

The metric  $d_H$  has an alternative definition, whose proof we leave to the reader:

**Proposition 3.12**  *$d_H(K, J)$  is equal to the minimum of  $\frac{1}{2}(\beta_2(Z) + |\sigma(Z)|)$ , where  $Z$  ranges over all smooth, oriented, compact, simply connected 4-manifolds with  $\partial Z = S^3 \sqcup -S^3$  in which  $K$  and  $J$  are concordant, that is, in which  $K \hookrightarrow S^3 \times 0$  and  $-J \hookrightarrow S^3 \times 1$  cobound a null-homologous, smoothly embedded annulus.*

### 3.3 The topological slice genus norm and the homology norm on $\mathcal{C}^{\text{top}}$

Just as in the smooth category, there are norms on  $\mathcal{C}^{\text{top}}$  given by the “slice genus” and the “homology norm”. The proofs that these are indeed norms are straightforward.

**Definition 3.13**  $\|K\|_s^{\text{top}}$ , also denoted by  $g_{\text{top}}(K)$ , the *topological slice genus* of  $K$ , is the minimum  $n$  such that  $K$  is the boundary of a compact oriented surface of genus  $n$  topologically and flatly embedded in  $B^4$ . This induces the metric  $d_s^{\text{top}}$ .

**Definition 3.14**  $\|K\|_H^{\text{top}}$ , the *topological homology norm* of  $K$ , has two equivalent definitions:

- (1)  $\|K\|_H^{\text{top}}$  is the minimum of  $\frac{1}{2}(\beta_2(V) + |\sigma(V)|)$ , where  $V$  ranges over all oriented, compact, simply connected topological 4-manifolds with  $\partial V = S^3$  in which  $K$  is slice, that is, in which  $K$  bounds a topologically flatly embedded disk that represents 0 in  $H_2(V, \partial V)$ .
- (2)  $\|K\|_H^{\text{top}}$  is the minimum of  $\frac{1}{2}(\beta_2(W) + |\sigma(W)|)$ , where  $W$  ranges over all compact, oriented topological 4-manifolds  $W$  whose boundary is  $M_K$ , whose  $\pi_1$  is normally generated by the meridian of  $K$ , and for which  $H_1(M_K) \rightarrow H_1(W)$  is an isomorphism.

This induces the metric  $d_H^{\text{top}}$ .

By the same proof as in Proposition 3.7 we have:

**Proposition 3.15** *If the zero framed surgeries  $M_J$  and  $M_K$  are topologically homology cobordant via a cobordism whose fundamental group is normally generated by either meridian, then  $\|K\|_H^{\text{top}} = \|J\|_H^{\text{top}}$ .*

### 3.4 The slice genus norm and the homology norm on $\mathcal{C}^{1/n}$

The reader may note that we could have defined another version of  $\mathcal{C}^{1/n}$ , in the topological category, but we resist doing so. We will discuss two norms on  $\mathcal{C}^{1/n}$ . The proofs that these are norms are straightforward.

**Definition 3.16** The  $\mathbb{Z}[1/n]$ -slice genus of  $K$ , denoted by  $\|K\|_{1/n}$  or by  $g_{1/n}(K)$ , is a norm given by the minimum  $n$  such that  $K$  is the boundary of a smoothly embedded compact oriented surface of genus  $n$  in a smooth manifold that is  $\mathbb{Z}[1/n]$ -homology equivalent to  $B^4$ . Let  $d_s^{1/n}$  denote the induced metric.

**Definition 3.17**  $\|K\|_H^{1/n}$ , the  $\mathbb{Z}[1/n]$ -homology norm of  $K$ , has two equivalent definitions:

- (1) The minimum of  $\frac{1}{2}(\beta_2(V) + |\sigma(V)|)$ , where  $V$  ranges over all smooth, oriented, compact, 4-manifolds with  $H_1(V; \mathbb{Z}[1/n]) = 0$  and  $\partial V = S^3$  in which  $K$  is slice, that is, in which  $K$  bounds a smoothly embedded disk that represents 0 in  $H_2(V, \partial V)$ .
- (2) The minimum of  $\frac{1}{2}(\beta_2(W) + |\sigma(W)|)$ , where  $W$  ranges over all smooth, compact, oriented 4-manifolds  $W$  whose boundary is  $M_K$  and for which  $H_1(M_K; \mathbb{Z}[1/n]) \rightarrow H_1(W; \mathbb{Z}[1/n])$  is an isomorphism.

Let  $d_H^{1/n}$  denote the induced metric.

By the same proof as in [Proposition 3.7](#) we have:

**Proposition 3.18** If the zero framed surgeries  $M_J$  and  $M_K$  are smoothly  $\mathbb{Z}[1/n]$ -homology cobordant, then  $\|K\|_H^{1/n} = \|J\|_H^{1/n}$ .

### 3.5 Comparison of metrics

The homology metric is not only different from  $d_s$  and  $d_s^{\text{ex}}$ , but is *very* different, as quantified by the following result. Surprisingly, the elements of this proof have only very recently come to light.

**Proposition 3.19** Neither of the functions  $i: (C, d_s) \rightarrow (C, d_H)$  and  $j: (C^{\text{ex}}, d_s^{\text{ex}}) \rightarrow (C^{\text{ex}}, d_H)$ , induced by the identity map, is a quasi-isometry.

**Proof** It suffices to exhibit a family of knots  $\{K_i \mid i \in \mathbb{Z}_+\}$  on which both the slice genus,  $g_s$ , and the exotic slice genus,  $g_{\text{ex}}$ , are unbounded functions of  $i$ , but on which the homology norm is a bounded function. For then there can be no constants  $A$  and  $B$  satisfying

(3-1) 
$$\frac{1}{A}g_*(K_i) - B \equiv \frac{1}{A}d^*_s(K_i, U) - B \leq d_H(K_i, U) \equiv \|K_i\|_H$$

for  $* = s$  or  $* = \text{ex}$  since the right-hand side is a bounded function of  $i$  whereas the left-hand side is not.

Such a family of knots was exhibited in [29, Proposition 3.4] (building on [10]). There it was shown that for a fixed satellite operator  $P$  (in fact the mirror image of the one shown in Figure 5.1) and for  $T$  the right-handed trefoil knot, the family  $K_i = P^i(T)$  for  $i \geq 0$  has the property that the slice genus and the exotic slice genus of  $K_i$  are equal to  $i + 1$ . Therefore, both  $\|-\|_s$  and  $\|-\|_s^{\text{ex}}$  are unbounded on this family. On the other hand, since  $P$  is a strong winding number one operator with  $P(U)$  unknotted, by [9, Corollary 4.4], for any knot  $J$ , the zero framed surgeries  $M_J$  and  $M_{P(J)}$  are smoothly homology cobordant via a cobordism whose fundamental group is normally generated by either meridian. Thus, by Proposition 3.7,  $\|J\|_H = \|P(J)\|_H$  for any knot  $J$ . In particular,  $\|T\|_H = \|P(T)\|_H = \|P(P(T))\|_H$ , et cetera. Thus,  $\|K_i\|_H = \|T\|_H$  for each  $i$ .  $\square$

We now compare all of the metrics. Since each  $d_*$  defined above has a well-defined meaning for any pair of (isotopy classes of) knots, their values can be compared, even though the functions  $d_*$  only give metrics on the appropriate set of concordance classes. Below, an inequality means for every pair of knots  $J$  and  $K$ , while a strict inequality means that, in addition, there exist knots  $K$  and  $J$  for which the metrics differ.

**Proposition 3.20** *We have*

$$\begin{aligned} d_H^{\text{top}} &\leq d_H < d_s^{\text{ex}} \leq d_s, \\ d_H^{\text{top}} &\leq d_s^{\text{top}} < d_s^{\text{ex}}, \\ d_H^{1/n} &< d_s^{1/n} \leq d_s^{\text{ex}}, \\ d_H^{1/n} &\leq d_H \end{aligned}$$

**Proof** The first and third inequalities in the first row are obvious. For the second inequality in the first row, it suffices to show that  $\|K\|_H \leq \|K\|_s^{\text{ex}}$  for all  $K$ . Suppose that  $\|K\|_s^{\text{ex}} = g$ , so that  $K$  is the boundary of a smoothly embedded, compact oriented surface  $\Sigma$  of genus  $g$  in a smooth manifold  $\mathcal{B}$  that is homeomorphic to  $B^4$ . Choose

disjoint simple closed curves,  $\{\gamma_1, \dots, \gamma_g\}$ , on  $\Sigma$  representing half of a symplectic basis for  $H_1$ . There exist framings of the normal bundles of these circles whose first vector field is tangent to  $\Sigma$ . Performing surgery on these circles using these framings transforms  $\mathcal{B}$  to  $V$ , in which  $\Sigma$  can be ambiently surgered to a disk. Hence,  $K$  is smoothly slice in  $V$ . Since each  $\gamma_i$  is null-homotopic, the collection  $\{\gamma_1, \dots, \gamma_g\}$  bounds a disjoint collection of smoothly embedded disks in  $\mathcal{B}$  (by pushing intersections off the boundary) [17, Section 1.5]. Hence, surgery on these circles alters the manifold by a connected sum with either  $S^2 \times S^2$  or  $S^2 \tilde{\otimes} S^2$ . In either case  $V$  is simply connected, with  $\beta_2(V) = 2g$  and  $\sigma(V) = 0$ . Thus,  $\|K\|_H \leq \|K\|_s^{\text{ex}}$ . The second inequality in the second row is obvious. For the first inequality in the second row, repeat the argument above. The second inequality in the third row is obvious. For the first inequality in the third row, repeat the argument above. The inequality in the fourth row is obvious.

To see that  $d_H$  can be strictly less than  $d_s^{\text{ex}}$ , let  $T$  be the right-handed trefoil and let  $P$  be the mirror image of the satellite operator in Figure 5.1. It was shown in [9, Corollary 4.4] that the zero framed surgeries on  $T$  and  $P(T)$  are homology cobordant via a cobordism whose fundamental group is normally generated by either meridian. hence by Proposition 3.7,  $\|P(T)\|_H = \|T\|_H = 1$ . Thus,  $d_H(P(T), U) = 1$ . But in [10, Section 3] it is shown that  $\tau(P(T)) > \tau(T) = 1$ . This implies that  $d_s^{\text{ex}}(P(T), U) \geq 2$ . To see that  $d_s^{\text{top}}$  can be strictly less than  $d_s^{\text{ex}}$ , consider the Whitehead double of the right handed trefoil  $D(T)$ . It is well known that  $D(T)$  is topologically slice but not exotic slice [11, Theorem 2.17]. This was first shown by Selman Akbulut (unpublished). Note that the theorem in [11] states that  $D(T)$  is not smoothly slice, however the same proof shows that the knot is not exotically slice. To see that  $d_H^{1/n} < d_s^{1/n}$ , consider  $T$  and its  $(n, 1)$ -cable, as explained in [10, Theorem 5.1]. □

Additionally, it is known that for  $n$  even,  $d_s^{1/n} < d_s^{\text{ex}}$ , since the figure-eight knot is slice in a  $\mathbb{Z}[\frac{1}{2}]$ -homology ball [7, page 63; 12, Lemma 2.2] but is not even a topologically slice knot since its Arf invariant is nontrivial. This same example shows that, for  $n$  even,  $d_H^{1/n} < d_H$ .

The others could be equalities! In particular, the question of whether or not  $d_s^{1/n} = d_s$  for  $n$  an odd prime is fascinating. It ought to be true that  $d_H^{\text{top}} < d_s^{\text{top}}$ , but [10, Section 3] was unable to show that  $T$  and  $P(T)$  are not topologically concordant. As mentioned above, if the 4-dimensional smooth Poincare conjecture is true then  $d_s = d_s^{\text{ex}}$ .

4 Existence of quasiflats

If  $(X, d)$  is a metric space then a *quasi- $n$ -flat* in  $X$  is a subspace of  $X$  that is quasi-isometric to  $\mathbb{R}^n$ , using the Euclidean metric on  $\mathbb{R}^n$ . We will show that  $(\mathcal{C}, d_s)$  has quasi- $n$ -flats for each  $n$ .

**Theorem 4.1** *For each  $n \geq 1$  there are subspaces of  $(\mathcal{C}, d_s)$  that are quasi-isometric to  $\mathbb{R}^n$ . The same holds for  $(\mathcal{C}, d_H)$ ,  $(\mathcal{C}^{\text{ex}}, d_s^{\text{ex}})$ ,  $(\mathcal{C}^{\text{ex}}, d_H)$ ,  $(\mathcal{C}^{\text{top}}, d_s^{\text{top}})$  and  $(\mathcal{C}^{\text{top}}, d_H^{\text{top}})$ .*

**Proof** First consider the case of  $(\mathcal{C}, d_s)$ . It is well known (and easy to see) that the inclusion of the integer lattice  $\mathbb{Z}^n \hookrightarrow \mathbb{R}^n$  with the taxicab (or  $\ell_1$ ) metric,  $d_t$ , on  $\mathbb{Z}^n$ , is a quasi-isometry. Hence, it suffices to exhibit a quasi-isometric embedding  $(\mathbb{Z}^n, d_t) \hookrightarrow (\mathcal{C}, d_s)$ . It is well known that  $\mathcal{C}$  contains free abelian subgroups of arbitrarily large rank. Specifically, suppose that  $K_1, \dots, K_n$  are linearly independent concordance classes, each of slice genus one, that are detected by homomorphisms  $\sigma_j: \mathcal{C} \rightarrow \mathbb{Z}$  for  $1 \leq j \leq n$ , meaning that  $\sigma_j(K_i) = 2\delta_{ij}$ . These homomorphisms show that the free abelian group on  $\{K_i\}$  is a subgroup of  $\mathcal{C}$ . Assume also that these  $\sigma_j$  give lower bounds on the slice genus in the sense that

(4-1) 
$$g_s(K_i) \geq \frac{1}{2}|\sigma_j(K_i)|.$$

Such classes  $K_i$  can easily be found by taking a certain family of (genus one) twist-knots and considering certain Tristram signature functions as the  $\sigma_j$  (see [32, Theorem 2.27]). We will show that the embedding  $(\mathbb{Z}^n, d_t) \hookrightarrow (\mathcal{C}, d_s)$  given by the  $K_i$  is a quasi-isometric embedding. Suppose that  $\vec{x}, \vec{y} \in \mathbb{Z}^n$ , where  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$ . It now suffices to show that

$$\frac{1}{n}d_t(\vec{x}, \vec{y}) \leq d_s(\vec{x}, \vec{y}) \leq n d_t(\vec{x}, \vec{y}).$$

Using the definitions of the metrics, this is equivalent to

(4-2) 
$$\frac{1}{n} \sum (|x_i - y_i|) \leq g_s\Big(\sum (x_i - y_i) K_i\Big) \leq n \sum (|x_i - y_i|).$$

By the subadditivity and symmetry of Definition 2.2,

$$g_s\Big(\sum (x_i - y_i) K_i\Big) \leq \sum (|x_i - y_i|)g_s(K_i) = \sum (|x_i - y_i|) \leq n \sum (|x_i - y_i|),$$

which verifies the right-hand side of inequality (4-2). On the other hand, by (4-1), for each  $j$ ,

$$g_s\Big(\sum (x_i - y_i) K_i\Big) \geq \frac{1}{2}\Big|\sigma_j\Big(\sum (x_i - y_i) K_i\Big)\Big| = |x_j - y_j|.$$

Hence,

$$g_s\left(\sum (x_i - y_i)K_i\right) \geq \frac{1}{n} \sum |x_i - y_i|,$$

which confirms the left-hand side of inequality (4-2).

Thus, we have shown that  $(\mathcal{C}, d_s)$  admits a quasi- $n$ -flat. The same proof works for the other cases since inequality (4-1) is known to hold for all these other norms [31, Section 1]. □

There is an interesting connection between the induced metrics on  $\mathbb{Z}$  subgroups of  $\mathcal{C}$  and the stable slice genus of knots, as defined by Livingston.

**Definition 4.2** [27] The stable slice genus of  $K$  is

$$g_{\text{st}}(K) \equiv \lim_{n \rightarrow +\infty} \frac{g_s(nK)}{n}.$$

Livingston showed that this induces pseudogroup norms on  $\mathcal{C}$  and  $\mathcal{C}$  modulo torsion, and a vector space pseudonorm on  $\mathcal{C} \otimes \mathbb{Q}$ .

**Proposition 4.3** The stable slice genus induces a group norm on  $\mathcal{C}$  modulo torsion (or a vector space norm on  $\mathcal{C} \otimes \mathbb{Q}$ ) if and only if every infinite cyclic subgroup of  $(\mathcal{C}, d_s)$  is a quasiflat.

**Proof** The stable slice genus induces a norm on  $\mathcal{C}$  modulo torsion if and only if  $g_{\text{st}}(K) > 0$  for each class  $[K]$  of infinite order in  $\mathcal{C}$ . First, suppose  $g_{\text{st}}(K) > 0$ . Then there is some positive integer  $N$  such that, for all  $n \geq N$ ,

$$\frac{g_{\text{st}}(K)}{2} \leq \frac{g_s(nK)}{n}.$$

Let  $A = \max\{g_s(K), 2/g_{\text{st}}(K)\}$  and  $B = Ng_s(K)$ .

To prove that the infinite cyclic subgroup of  $(\mathcal{C}, d_s)$  generated by  $K$  is a quasiflat, we will show for all  $m, n \in \mathbb{Z}$  that

$$\frac{d_t(m, n)}{A} - B \leq d_s(mK, nK) \leq Ad_t(m, n) + B.$$

Since  $g_s(-K) = g_s(K)$ , this is equivalent to showing for all  $n \geq 1$  that

$$\frac{1}{A} - \frac{B}{n} \leq \frac{g_s(nK)}{n} \leq A + \frac{B}{n}.$$



Let  $n$  be a positive integer. To prove the rightmost inequality, note that  $A \geq g_s(K)$ ,  $B \geq 0$  and  $g_s(nK) \leq ng_s(K)$ , hence

$$\frac{g_s(nK)}{n} \leq g_s(K) \leq A \leq A + \frac{B}{n}.$$

To show the leftmost inequality, we break it into two cases. For the first case, suppose  $n \geq N$ . Since  $A \geq 2/g_{\text{st}}(K)$ , we have

$$\frac{1}{A} - \frac{B}{n} \leq \frac{1}{A} \leq \frac{g_{\text{st}}(K)}{2} \leq \frac{g_s(nK)}{n}.$$

Now, for the second case, suppose  $1 \leq n < N$ . Then  $B/n \geq g_s(K)$ . In addition, since  $\frac{1}{2}g_{\text{st}}(K)$  and  $\frac{1}{n}g_s(nK)$  are both bounded above by  $g_s(K)$ , we have that

$$\frac{g_{\text{st}}(K)}{2} - \frac{g_s(nK)}{n} \leq \left| \frac{g_{\text{st}}(K)}{2} - \frac{g_s(nK)}{n} \right| \leq g_s(K).$$

Thus,

$$\frac{1}{A} - \frac{B}{n} \leq \frac{g_{\text{st}}(K)}{2} - g_s(K) \leq \frac{g_s(nK)}{n}.$$

Hence,  $(\mathbb{Z}, d_t) \hookrightarrow (\mathcal{C}, d_s)$  given by  $n \mapsto nK$  is a quasi-isometric embedding.

On the other hand, if  $n \mapsto nK$  is a quasi-isometric embedding then, for some  $A \geq 1$  and  $B \geq 0$ ,

$$\frac{n}{A} - B \leq g_s(nK),$$

so  $g_{\text{st}}(K) \geq 1/A > 0$ . □

## 5 Satellite operators and other natural operators

In this section we review some natural operators on  $\mathcal{C}^*$  given by taking the reverse, taking the mirror image, the connected sum with a fixed class, satellite operators, and multiplication by an integer with respect to the group structure.

Let  $\text{ST} \equiv S^1 \times D^2$ , where both  $S^1$  and  $D^2$  have their usual orientations. We will always think of  $\text{ST}$  as embedded in  $S^3$  in the standard unknotted fashion. Suppose  $P \subset \text{ST}$  is an embedded oriented circle, called a *pattern knot*, that is geometrically essential (even after isotopy,  $P$  has nontrivial intersection with a meridional 2-disk). The *geometric winding number* of  $P$ , denoted by  $\text{gw}(P)$ , is the minimum number of these intersection points over all patterns isotopic to  $P$ . The *winding number* of  $P$  is the algebraic number of such intersections. We say that  $P$  has *strong winding number*  $\pm 1$  if the meridian of the solid torus  $\text{ST}$  normally generates  $\pi_1(S^3 - \tilde{P})$ ,




Figure 5.1: A strong winding number one pattern  $P$

where  $\tilde{P}$  is the knot  $P \subset \text{ST} \subset S^3$  [9, Definition 1.1]. Note that if  $\tilde{P}$  is unknotted then winding number one is the same as strong winding number one. Suppose  $K$  is an oriented knot in  $S^3$  given as the image of the embedding  $f_K: S^1 \rightarrow S^3$ . Then there is an orientation-preserving diffeomorphism  $\tilde{f}_K: S^1 \times D^2 \rightarrow N(K)$ , where  $N(K)$  is a tubular neighborhood of  $K$ , such that  $\tilde{f}_K = f$  on  $S^1 \times \{0\}$  and  $\tilde{f}_K$  takes the oriented meridian  $\eta$  of  $\text{ST}$  to the oriented meridian of  $K$ , and takes a preferred longitude of  $\text{ST}$ ,  $S^1 \times \{1\}$ , to a preferred oriented longitude of  $K$ . The (oriented) knot type of the image of  $P$  under  $\tilde{f}_K: \text{ST} \rightarrow N(K) \hookrightarrow S^3$  is called the (untwisted) *satellite* of  $K$  with *pattern knot*  $P$  [24, page 10]. In this paper this will be denoted by  $P(K)$ . Note that  $\tilde{P} = P(U)$ . In this paper,  $P$  will denote, depending on the context, either a knot in the solid torus or the corresponding induced function on a set of equivalence classes of knots in  $S^3$

$$P: \mathcal{K}/\sim \rightarrow \mathcal{K}/\sim$$

given by  $K \mapsto P(K)$ . Such functions seem rarely to be additive with respect to the monoidal structure on  $\mathcal{K}$  given by connected sum. It is well known that satellite functions descend to yield what we call *satellite operators* on  $\mathcal{K}/\sim$  for various important equivalence relations on knots. In particular, any such operator descends to  $P: \mathcal{C}^* \rightarrow \mathcal{C}^*$  on the various sets of “concordance classes of knots” as defined in Section 1. For a fixed pattern knot  $P$ , we will use the same notation for each of these satellite operators. Some examples are of particular importance.

**Definition 5.1** If  $J$  is a knot then the *connected sum operator* (corresponding to  $J$ ), denoted by  $C_J: \mathcal{C}^* \rightarrow \mathcal{C}^*$ , is the function given by  $C_J(K) = J \# K$ .

Note that if  $P$  is a pattern whose geometric winding number is  $+1$ , then the operator  $P: \mathcal{C}^* \rightarrow \mathcal{C}^*$  is  $C_{\tilde{P}}$ . In particular, a connected sum operator is a satellite operator.

**Definition 5.2** The *reverse operator*, denoted by  $r: \mathcal{C}^* \rightarrow \mathcal{C}^*$ , is the function that sends each knot  $J$  to the class represented by  $rJ$ , the reverse of the knot.

The reverse operator is a satellite operator whose pattern knot is the core of  $ST$ , oriented so that its winding number is  $-1$ .

**Definition 5.3** If  $n$  is an integer then the  $(n, 1)$ -cable operator,  $C_{n,1}: \mathcal{C}^* \rightarrow \mathcal{C}^*$ , is the satellite operator given by the pattern which is the  $(n, 1)$ -torus knot. When  $n = 0$  it is understood that  $C_{n,1}: \mathcal{C}^* \rightarrow \mathcal{C}^*$  is the *zero operator*  $Z: \mathcal{C}^* \rightarrow \mathcal{C}^*$  that sends every class to the class of the unknot.

There are other natural operators on  $\mathcal{C}^*$  that are not necessarily satellite operators, but which we will consider.

**Definition 5.4** The *mirror image operator* is the function  $J \mapsto \bar{J}$ , where  $\bar{J}$  is the mirror image.

**Definition 5.5** The *times  $m$  operator* is the function  $J \mapsto mJ$ , where  $mJ$  denotes the connected sum of  $|m|$  copies of  $J$  if  $m \geq 0$  or  $|m|$  copies of  $-J$  if  $m < 0$ .

## 6 Metric aspects of operators

The simplest operators are bijective isometries. For example:

**Proposition 6.1** Any connected sum operator  $P$  induces a bijective isometry  $P: \mathcal{C}^* \rightarrow \mathcal{C}^*$  with respect to any metric induced from a group norm. This is a quasi-isometry with respect to  $A = 1$ ,  $B = 0$  and  $C = 0$ . The same holds for the “reverse” operator  $J \mapsto rJ$  and the mirror image operator,  $J \mapsto \bar{J}$ , as long as  $\|rJ\| = \|J\|$  (which holds for all the norms we are discussing).

**Proof** Clearly any connected sum operator,  $P$ , has an inverse which is also a connected sum operator. Thus,  $P$  is surjective. Then note

$$d(P(K), P(J)) = d(K \# \tilde{P}, J \# \tilde{P}) = \|K \# \tilde{P} \# -\tilde{P} \# -J\| = \|K \# -J\| = d(K, J).$$

Thus,  $P$  is a (surjective) isometry and so  $P$  is a quasi-isometry with respect to  $A = 1$ ,  $B = 0$  and  $C = 0$ .

A similar argument works in the other cases, after noting that  $\|\bar{J}\| = \|r(-J)\| = \|-J\|$ , by assumption, and  $\|-J\| = \|J\|$  by definition of a group norm.  $\square$

**Lemma 6.2** *Suppose  $d$  is a metric on a group  $G$  and suppose that  $f: G \rightarrow G$  is a quasi-isometry with respect to constants  $A$ ,  $B$  and  $C$  in the notation of Definition 1.2. If  $g: G \rightarrow G$  is within a bounded distance  $D$  of  $f$ , then  $g$  is also a quasi-isometry, with respect to constants  $A' = A$ ,  $B' = B + 2D$  and  $C' = C + D$ .*

The proof of Lemma 6.2 is straightforward so we omit it here. The following key result says that any satellite operator is within a bounded distance of a simpler operator.

**Proposition 6.3** *Any winding number  $n$  operator  $P: (C^*, d_*) \rightarrow (C^*, d_*)$  is within a bounded distance of the  $(n, 1)$ -cable operator, with respect to any of the metrics we have introduced.*

**Proof** The operator  $P$  corresponds to a pattern knot  $P \hookrightarrow \text{ST}$ . The operator  $C_{n,1}$  corresponds to a different pattern knot  $P' \hookrightarrow \text{ST}$ , both of which have winding number  $n$ . Consider  $P \hookrightarrow \text{ST} \times \{0\}$  and  $P' \hookrightarrow \text{ST} \times \{1\}$ . Since these two oriented circles are homologous in  $\text{ST} \times [0, 1]$ , it is easily seen that they cobound a compact oriented surface  $\Sigma$  in  $\text{ST} \times [0, 1]$ . Let  $D$  be the genus of  $\Sigma$ . Note that  $D$  depends only on  $P$ . Then, for any knot  $J$  whose tubular neighborhood is given by an embedding  $f_J: \text{ST} \hookrightarrow S^3$ , consider the image of  $\Sigma$  under the map  $f_J \times \text{id}: \text{ST} \times [0, 1] \hookrightarrow S^3 \times [0, 1]$ . This surface forms a smooth cobordism of genus  $D$  from  $P(J)$  to  $P'(J)$ . By Proposition 3.3,  $d_s(P(J), P'(J)) \leq D$ . Thus, by Proposition 3.20, for each of the metrics we have defined,  $d_*(P(J), P'(J)) \leq D$  for each  $J$ .

With more attention, one can get an explicit bound for  $D$  involving only  $n$  and the geometric winding number of  $P$ , namely  $D \leq \|\tilde{P}\|_* + \frac{1}{2}(\text{gw}(P) - |n|) + (|n| - 1)$ .  $\square$

**Corollary 6.4** *Any winding number 1 satellite operator  $P$  is within a bounded  $d_*$ -distance of the identity map. Any winding number  $-1$  satellite operator  $P$  is within a bounded  $d_*$ -distance of the reverse operator.*

## 6.1 Winding number one satellite operators

By Corollary 6.4, a general winding number  $+1$  operator behaves roughly like a connected sum operator. Consequently:

**Theorem 6.5** *If  $P$  is a winding number  $\pm 1$  pattern then  $P: (\mathcal{C}^*, d_*) \rightarrow (\mathcal{C}^*, d_*)$  is a quasi-isometry.*

**Proof** By Corollary 6.4, such a  $P$  is within a bounded  $d_*$ -distance of either the identity or the reverse operator. By Proposition 6.1, both of the latter are bijective isometries, hence quasi-isometries with  $A = 1$ ,  $B = 0$  and  $C = 0$ . Hence, by Lemma 6.2,  $P$  is a quasi-isometry with respect to  $d_*$ . In fact, using the last line of the proof of Proposition 6.3, and applying Lemma 6.2 where  $f$  is an isometry, we see that  $P$  is a quasi-isometry with respect to the constants  $A = 1$ ,  $B = \text{gw}(P) - 1$  and  $C = \frac{1}{2}(\text{gw}(P) - 1)$ .

Using a slightly different approach, one can do slightly better and get  $B = \frac{1}{2}(\text{gw}(P) - 1)$ . Namely, as we will see in Figures 6.1 and 6.2 in the proof of Theorem 6.6, we have that

$$-P(K) \# P(J) = -P(K) \# P(K \# -K \# J) = R(-K \# J),$$

where  $R$  is a pattern with the same winding number, with  $\text{gw}(R) \leq \text{gw}(P)$  and for which  $\tilde{R}$  is a slice knot. Then, by Corollary 6.4,  $R$  is within a distance  $\frac{1}{2}(\text{gw}(P) - 1)$  of a surjective isometry. Hence,

$$\begin{aligned} d_*(P(K), P(J)) &\equiv d_*(-P(K) \# P(J), U) = d_*(R(-K \# J), U) \\ &\leq \| -K \# J \|_* + \frac{1}{2}(\text{gw}(P) - 1) \\ &= d_*(K, J) + \frac{1}{2}(\text{gw}(P) - 1). \end{aligned} \qquad \square$$

Strong winding number one operators have an even better behavior with respect to the homology norm.

**Theorem 6.6** *If  $P$  is a strong winding number  $\pm 1$  pattern then*

$$P: (\mathcal{C}, d_H) \rightarrow (\mathcal{C}, d_H)$$

*preserves the pseudonorm  $d_H$  and is quasisurjective, so that if the 4D Poincaré conjecture is true then  $P$  is an isometric embedding of  $\mathcal{C}$  that is quasisurjective. Moreover, for  $*$  = ex, top or  $1/n$ ,*

$$P: (\mathcal{C}^*, d_H^*) \rightarrow (\mathcal{C}^*, d_H^*)$$

*is an isometric embedding and is quasisurjective.*

**Proof** By Theorem 6.5,  $P$  is a quasi-isometry and hence is quasisurjective. Now we show that  $P$  preserves the norm (or pseudonorm). Since any isometry is an injective function, our proof should be viewed as a generalization of [9, Theorem 5.1],

where it was shown that any such operator  $P$  is injective in the cases  $\ast = \text{ex}$ ,  $\text{top}$  or  $1/n$ , and injective on  $\mathcal{C}$  if the 4D Poincaré conjecture holds. First we prove the theorem in the very special case that  $\tilde{P} = 0$  in  $\mathcal{C}^\ast$  and  $J = 0 = P(J)$ :

**Lemma 6.7** *If  $R$  is a strong winding number  $\pm 1$  pattern and  $\tilde{R} = 0$  in  $\mathcal{C}^\ast$  then the satellite operator  $R: \mathcal{C}^\ast \rightarrow \mathcal{C}^\ast$  preserves the homology norm  $\|-\|_H^\ast$ , that is,  $\|R(K)\|_H^\ast = \|K\|_H^\ast$  for each  $K$ .*

**Proof** Since  $R$  has strong winding number  $\pm 1$ , by [9, Corollary 4.4] and, in the case  $\ast = 1/n$ , by [10, Theorem 2.1], the zero framed surgeries  $M_{R(K)}$  and  $M_K$  are smoothly (respectively topologically, smoothly  $\mathbb{Z}[1/n]$ -) homology cobordant. Moreover, except in the case  $\ast = 1/n$ , we may assume the cobordism has fundamental group normally generated by each meridian. Thus, by Propositions 3.7, 3.15 and 3.18,  $\|R(K)\|_H^\ast = \|K\|_H^\ast$ .

We sketch the proof of [9, Corollary 4.4] in the case  $\ast = \text{ex}$  for the convenience of the reader. The other cases and the proof of [10, Theorem 2.1] are similar. There is a standard cobordism  $E$  whose boundary is the disjoint union of  $-M_{R(K)}$ ,  $M_R$  and  $M_K$  (see for example [10, page 2198]). This is obtained by gluing  $M_{\tilde{R}} \times [0, 1]$  to  $M_K \times [0, 1]$  by identifying the surgery solid torus in  $M_K \times \{1\}$  with the solid torus:  $\eta \times D^2 \hookrightarrow M_{\tilde{R}} \times \{1\}$ . Since  $\tilde{R} = 0$  in  $\mathcal{C}^\ast$ , it bounds a slice disk  $\Delta$  in some smooth homotopy 4-ball  $\mathcal{B}$ . Use the manifold  $\mathcal{B} - N(\Delta)$  to cap off the  $M_{\tilde{R}}$  boundary component of  $E$ , yielding a cobordism  $V$  between  $M_K$  and  $M_{R(K)}$ . This is the required homology cobordism.  $\square$

We return to the general situation in the proof of Theorem 6.6. Recall that, by definition,

$$d_H^\ast(P(J), P(K)) = \|-P(K) \# P(J)\|_H^\ast.$$

Now we mimic one of the key steps in the proof of [9, Theorem 5.1]. Since  $K \# -K$  is a slice knot in any category,  $J = K \# -K \# J$  in  $\mathcal{C}^\ast$ , so  $P(J) = P(K \# -K \# J)$  in  $\mathcal{C}^\ast$ . This last knot is pictured on the right-hand side of Figure 6.1. Hence,

$$(6-1) \quad d_H^\ast(P(J), P(K)) = \|-P(K) \# (P(K \# -K \# J))\|_H^\ast.$$

A picture of the connected sum of knots on the right-hand side of (6-1) is shown in Figure 6.1. The particular form we have pictured for the  $-[P(K)]$  summand is not important. This form will not be used. Let  $R$  be the pattern knot shown in Figure 6.2.




Figure 6.1

In terms of this operator, (6-1) becomes

(6-2) 
$$d_H^*(P(J), P(K)) = \|R(-K \# J)\|_H^*.$$

Furthermore, observe that  $\tilde{R} = -P(K) \# P(K)$  is a ribbon knot, hence a slice knot in any category. Also note that the winding number of  $R$  is the same as that of  $P$ , which is  $\pm 1$ . In fact it was shown in the proof of [9, Theorem 5.1] that  $R$  has strong winding number one since  $P$  does. We also observe that the geometric winding number of  $R$  is at most that of  $P$ . Thus, by Lemma 6.7,

$$d_H^*(P(J), P(K)) = \|R(-K \# J)\|_H^* = \|-K \# J\|_H^* \equiv d_H^*(J, K).$$

Thus, we have shown that  $P$  is an isometry, hence injective. Thus,  $P$  is a bijection to its image. □

6.2 General nonzero winding number satellite operators

**Theorem 6.8** *If  $P$  is a winding number  $n$  pattern with  $n \neq 0$ , then the satellite operator*

$$P: (\mathcal{C}^{1/n}, d_H^{1/n}) \rightarrow (\mathcal{C}^{1/n}, d_H^{1/n})$$

*is an isometric embedding. Moreover,  $P$  is quas surjective.*

**Proof** The proof is almost identical to that of Theorem 6.6 above. Note that it was already shown in [9, Theorem 5.1] that  $P$  is injective. To show that  $P$  preserves the norm, we repeat the proof of Theorem 6.6, replacing Lemma 6.7 by the following. □




Figure 6.2: The operator  $R = -[P(K)] \# P(K)$

**Lemma 6.9** *If  $R$  is a pattern of nonzero winding number  $n$  and  $\tilde{R} = 0$  in  $\mathcal{C}^{1/n}$ , then the satellite operator  $R: \mathcal{C}^{1/n} \rightarrow \mathcal{C}^{1/n}$  preserves the homology norm  $\|-\|_H^{1/n}$ .*

**Proof** By [10, Theorem 2.1], the zero framed surgeries  $M_{R(K)}$  and  $M_K$  are smoothly  $\mathbb{Z}[1/n]$ -homology cobordant. Thus, by Proposition 3.18,  $\|R(K)\|_H^{1/n} = \|K\|_H^{1/n}$ .  $\square$

6.3 Winding number zero satellite operators

We show that every winding number zero operator is a bounded function, hence is approximately a constant map and thus is an approximate contraction. Not all winding number zero operators are injective functions. It remains possible that some are injective, or nearly so.

**Definition 6.10** A function  $f: X \rightarrow Z$  is an *approximate contraction* if there is some constant  $D > 0$  such that  $d(f(x), f(y)) \leq \max\{D, d(x, y)\}$  for all  $x, y \in X$ .

Any bounded function is within a bounded distance of a constant map and hence is approximately a contraction, in that, as long as  $x$  and  $y$  are not too close to each other,  $d(f(x), f(y)) < d(x, y)$ . Thus, it follows from the  $n = 0$  case of Proposition 6.3 that:

**Proposition 6.11** *Any winding number zero satellite operator on  $(\mathcal{C}^*, d_*)$  is a bounded function, where  $(\mathcal{C}^*, d_*)$  is any of the metric spaces we have defined. Thus, any winding number zero satellite operator is an approximate contraction.*






Figure 6.3: A family of nontrivial noninjective operators  $R^k$

**Proof** By Proposition 6.3, such a  $P$  is within a bounded distance  $C$  of the constant operator that sends every class to  $\tilde{P}$ . Hence,  $d_*(P(K), P(J)) \leq 2C$ .  $\square$

We should not expect that every winding number zero operator is an injective function, because the zero operator may be viewed as a (degenerate) satellite operator whose pattern knot is an unknot which has zero geometric winding number. But it is easy to find nondegenerate patterns that yield the zero operator by choosing a pattern of nonzero geometric winding number which is concordant, inside the solid torus, to this unknot. Such a pattern will be called a *trivial pattern* since it induces the zero operator. There are various algebraic conditions on the pattern that ensure that an operator is nontrivial (see [13, Definition 7.2]) and is indeed injective on very large subsets of  $\mathcal{C}$ .

But even here the precise situation is unclear. For example, consider the family of winding number zero patterns  $R^k$  shown on the left-hand side of Figure 6.3. The solid torus is the exterior of a neighborhood of the dashed circle  $\eta$ . This pattern has an obvious genus one Seifert surface. The  $-k$  signifies the number of full twists between the two bands of that surface (without twisting the two strands of a fixed band). A knot in a box indicates that all strands passing through that box are tied into parallel copies of the indicated knot. The value of this operator on a knot  $K$  is shown on the right-hand side of the figure. Then  $R^k(U)$  is a slice knot since the core of the right-hand band has self-linking zero and has the knot type of the unknot. Similarly,  $R^k(\bar{J})$  is a slice knot since the left-hand band has zero self-linking and has the knot type of the ribbon knot  $J \# -J$ . Hence,  $R^k$  is not injective since both  $U$  and  $\bar{J}$  are sent to  $0 \in \mathcal{C}$ . Since an arbitrary pattern that is a genus one ribbon knot has two such “metabolizing curves”, this is the generic situation.




Figure 6.4

But recent examples indicate that the situation is even more complicated. In [8], the knot  $\tilde{R}$  in Figure 6.4 was shown to be a slice knot for any knot  $J$ . If we let  $\eta$  be a circle linking the right-hand band, the resulting operator,  $R$ , has winding number zero. Then  $R(U) = \tilde{R}$  is slice. But as above there are two metabolizing curves. The core of the right-hand band of the Seifert surface for  $\tilde{R}$  has self-linking zero and has the knot type of  $J_1 = J_{(2,1)} \# -J$  so  $R(-J_1)$  is also slice. There is a circle that goes over each band once that has self-linking zero which has a knot type, say  $J_2$ , so  $R(-J_2)$  is also slice. Thus, there are at least three knots (provably distinct for most  $J$ ) that are sent to 0 by the operator  $R$ .

Nonetheless there is a lot of evidence that some winding number zero operators are injective, or are nearly so [13].

### 6.4 Comparison of satellite operators and other natural operators

**Proposition 6.12** *For  $m \notin \{0, 1\}$  the function  $f_m$  given by multiplication by  $m$  on  $\mathcal{C}$  (respectively on  $\mathcal{C}^{\text{ex}}$ ) is not within a bounded distance of any satellite operator, where the metric is  $d_s$  (respectively  $d_s^{\text{ex}}$ ). The function  $f_0$  (namely the zero operator) is not within a bounded distance of any nonzero winding number satellite operator. The identity (ie  $f_1$ ) is not within a bounded distance of any satellite operator whose winding number is not  $\pm 1$ .*

**Proof** The proof will be by contradiction. Suppose that  $m \notin \{0, 1\}$  and that  $f_m$  is within  $D$  of the winding number  $n$  satellite operator  $P$ . By Proposition 6.3,  $P$  is

within some bounded distance  $D'$  of the cabling operator  $C_{(n,1)}$ . Let  $E = D + D'$ . Consequently, for every knot  $J$  we would have that

$$d(mJ, J_{(n,1)}) \leq E,$$

or, equivalently,

$$\| -mJ \# J_{(n,1)} \| \leq E.$$

But both the slice genus and the exotic slice genus are bounded below by one half of the absolute value of the classical knot signature. Hence, to reach a contradiction we need only specify a  $J$  (given  $m$ ,  $n$  and  $\tilde{P}$ ) such that

$$|\sigma(-mJ) + \sigma(J_{(n,1)})| > 2E.$$

We assume that  $m \notin \{0, 1\}$ . We can assure that

$$|\sigma(J)| > 2E,$$

by choosing  $J$  to be the connected sum of a sufficiently large number of left-handed trefoil knots. Note that for such a  $J$  the Levine–Tristram signature function takes on only two values, namely 0 and  $\sigma(J)$ . Thus,  $\sigma(J_{(n,1)})$  is either zero (if  $n = 0$ ) or is equal to the  $\exp(\pi/|n|)$ -signature of  $J$  [20], so in any case is either zero or is equal to  $\sigma(J)$ . Hence, we have

$$|\sigma(-mJ) + \sigma(J_{(n,1)})| \geq |(m-1)| |\sigma(J)| > 2E,$$

resulting in a contradiction.

Moreover, if  $m = 0$ , then, as long as  $n \neq 0$ , we can choose  $J$  such that  $|\sigma(J_{(n,1)})|$  is very large and such that

$$|\sigma(-mJ) + \sigma(J_{(n,1)})| = |\sigma(J_{(n,1)})| > 2E,$$

for a contradiction. For example, consider the twist knot  $T_k$  with  $k$  twists. The Alexander polynomial of  $T_k$  is  $-kt^2 + (2k+1)t - k$  and hence, for  $k \leq 1$ , the Alexander polynomial has exactly two roots on the unit circle, with real value  $1 - \left| \frac{1}{2k} \right|$ ; these knots also has nonzero signature. So, for  $k$  negative with  $|k|$  large,  $\sigma((T_k)_{(n,1)})$  is nonzero. Hence, one can choose  $J$  to be the connected sum of a large number of twist knots with  $k$  negative and  $|k|$  sufficiently large. It follows that the times zero operator is not within a bounded distance of any satellite operator with *nonzero* winding number.

In the case that  $m = 1$  we can find a twist knot  $T$  of signature 2 wherein the roots of its Alexander polynomial have real part less than zero. Then  $\sigma(T_{(n,1)}) = 0$  if  $n \neq \pm 1$ . Then let  $J$  be the connected sum of a large number of such  $T$ , so that

$$|\sigma(J)| > 2E.$$

and  $\sigma(J_{(n,1)}) = 0$ . Hence, the times one operator (namely the identity) is not within a bounded distance of any satellite operator whose winding number is not  $\pm 1$ . □

**Corollary 6.13** *Neither the mirror image operator nor the inverse operator on  $\mathcal{C}$  (respectively on  $\mathcal{C}^{\text{ex}}$ ) is within a bounded distance of any satellite operator, where the metric is  $d_s$  (respectively  $d_s^{\text{ex}}$ ).*

**Proof** The inverse operator is the case  $m = -1$  of [Proposition 6.12](#). Since the inverse operator is the composition of the mirror image operator followed by the reverse operator, and since the reverse operator is both an isometry and a satellite operator, if the mirror image operator were within a bounded distance of a satellite operator then the inverse operator would be within a bounded distance of a composition of satellite operators, which is a satellite operator. This is a contradiction. □

The second two statements of [Proposition 6.12](#) can easily be generalized.

**Proposition 6.14** *Suppose  $m \neq \pm n$ . Then no winding number  $m$  satellite operator is within a bounded distance of any winding number  $n$  satellite operator.*

**Proof** As above, for a proof by contradiction, it suffices to show that, given  $E > 0$ , we can find a knot  $J$  such that

$$\| -J_{(m,1)} \# J_{(n,1)} \| > E.$$

For this, as above it suffices to find  $\omega \in S^1$  such that

$$|\sigma_\omega(-J_{(m,1)}) + \sigma_\omega(J_{(n,1)})| > 2E,$$

where  $\sigma_\omega$  is the value of the (normalized) Levine–Tristram signature function at  $\omega$ . This is true because both the slice genus and the exotic slice genus are bounded below by one half of  $|\sigma_\omega|$  [\[31\]](#). Let  $J$  be the connected sum of more than  $2E$  copies of the left-handed trefoil  $-T$ , and let  $\alpha \in S^1$  be the root of the Alexander polynomial of the trefoil with positive imaginary part. Then, for any  $\beta \in S^1$  (with positive imaginary part) whose argument is greater than that of  $\alpha$ , we have  $\sigma_\beta(-T) = 2$ , so  $\sigma_\beta(J) > 2E$ ;

whereas, for any  $\beta \in S^1$  (with positive imaginary part) whose argument is less than that of  $\alpha$ , we have  $\sigma_\beta(-T) = 0$ , so  $\sigma_\beta(J) = 0$ .

Suppose  $|n| < |m|$ . Let  $r_n$  and  $r_m$  be the  $|n|^{\text{th}}$  and  $|m|^{\text{th}}$  roots of  $\alpha$  that have smallest positive argument. Then  $\arg(r_n) > \arg(r_m)$ . Let  $\omega \in S^1$  have argument between these and be close to  $r_m$ . Then

$$\sigma_\omega(J_{(n,1)}) = \pm \sigma_{\omega^{|n|}}(J) = 0$$

since the argument of  $\omega^{|n|}$  is less than that of  $\alpha$  [20]. On the other hand,

$$|\sigma_\omega(J_{(m,1)})| = |\pm \sigma_{\omega^{|m|}}(J)| > 2E$$

since the argument of  $\omega^{|m|}$  is slightly greater than that of  $\alpha$ . These choices lead to the desired contradiction.  $\square$

**Remark** With more work, the  $\pm$  in Proposition 6.14 should be able to be replaced by  $+$ . It has been shown, using Casson–Gordon invariants, that some knots are not concordant to their reverses [22; 14, Theorem 5.5.2; 28]. It is also known that Casson–Gordon invariants, in a certain sense, give lower bounds for the slice genus. Thus, it may be possible to show, with some effort beyond what is in the literature, that the reverse operator is not within a bounded distance of the identity. This would imply that no winding number  $-1$  satellite operator is within a bounded distance of any winding number  $+1$  satellite operator. For  $n > 1$ , to show that no winding number  $n$  satellite operation is within a bounded distance of any winding number  $-n$  operation would seem to require calculations (of, say, Casson–Gordon invariants) far beyond what is currently in the literature. Nonetheless, there is little doubt, in the authors’ opinion, as to its veracity.

## 7 Satellite operators as quasimorphisms

**Theorem 7.1** Any satellite operator  $P: \mathcal{C} \rightarrow \mathcal{C}$  is a quasihomomorphism with respect to norm given by the slice genus,  $\|-\|_s$ . Similarly, any satellite operator  $P: \mathcal{C}^{\text{ex}} \rightarrow \mathcal{C}^{\text{ex}}$  is a quasihomomorphism with respect to either  $d_s^{\text{ex}}$  or  $d_H$ .

**Proof** The following is well known:

**Lemma 7.2** If  $f: G \rightarrow (A, d)$  is a quasihomomorphism and  $g: G \rightarrow A$  is within a bounded distance of  $f$ , then  $g$  is a quasihomomorphism.

**Proof of Lemma 7.2** By subadditivity,  $\|a\| \leq \|a-b\| + \|b\|$ . The lemma now follows quickly from setting  $a = g(xy) - g(x) - g(y)$  and  $b = f(xy) - f(x) - f(y)$ .  $\square$

By Proposition 6.3, a satellite operator of winding number  $n$  is within a bounded distance of the operator  $C_{n,1}$ , so by Lemma 7.2, it suffices to show that latter is a quasihomomorphism. This is done geometrically. By adding  $n$  carefully placed bands, the  $(n, 1)$ -cable of the knot  $K \# J$  can be transformed to the disjoint union of the  $(n, 1)$ -cable of  $K$  with an  $n$ -component link consisting of parallel copies of  $J$ . By adding  $n-1$  more bands, the latter can be transformed to the  $(n, 1)$ -cable of  $J$ . Adding one more band transforms the disjoint union to the connected sum. Hence, there is a cobordism in  $S^3 \times [0, 1]$  from the  $(K \# J)_{(n,1)}$  to  $K_{(n,1)} \# J_{(n,1)}$  whose genus is independent of  $K$  and  $J$ . Hence,

$$\|(K \# J)_{(n,1)} - K_{(n,1)} - J_{(n,1)}\|_s \equiv d_s((K \# J)_{(n,1)}, K_{(n,1)} \# J_{(n,1)})$$

is bounded by a function of  $n$  alone. Thus, the  $(n, 1)$ -cable operator is a quasihomomorphism.  $\square$

**Proposition 7.3** Suppose that  $q: \mathcal{C} \rightarrow \mathbb{R}$  is a quasihomomorphism whose absolute value gives a lower bound for a positive multiple of the slice genus, that is, there is some  $C > 0$  such that, for all  $K$ ,  $|q(K)| \leq Cg_s(K)$ . Then, for any satellite operator  $P$ , the composition  $q \circ P: \mathcal{C} \rightarrow \mathbb{R}$  is a quasihomomorphism.

**Proof** The proof follows that of [5, Corollary 1.C], where the slightly stronger assumption was made that  $q$  is Lipschitz. Let  $E = P(K \# J) - P(K) - P(J)$ . Since  $P$  is a quasihomomorphism,  $\|E\|_s \leq D_P$ . Then

$$\begin{aligned} |q(P(K \# J)) - q(P(K)) - q(P(J))| &= |q(P(K) + P(J) + E) - q(P(K)) - q(P(J))| \\ &\leq 2D_q + |q(E)| \\ &\leq 2D_q + C\|E\|_s \\ &\leq 2D_q + CD_P. \end{aligned} \quad \square$$

## References

- [1] S Akbulut, R Kirby, *Mazur manifolds*, Michigan Math. J. 26 (1979) 259–284 [MR](#)
- [2] S Akbulut, K Yasui, *Corks, plugs and exotic structures*, J. Gökova Geom. Topol. 2 (2008) 40–82 [MR](#)

- [3] **L Bartholdi, R Grigorchuk, V Nekrashevych**, *From fractal groups to fractal sets*, from “Fractals in Graz 2001” (P Grabner, W Woess, editors), Birkhäuser, Basel (2003) 25–118 [MR](#)
- [4] **S Boyer**, *Shake-slice knots and smooth contractible 4-manifolds*, Math. Proc. Cambridge Philos. Soc. 98 (1985) 93–106 [MR](#)
- [5] **M Brandenbursky, J Kędra**, *Concordance group and stable commutator length in braid groups*, Algebr. Geom. Topol. 15 (2015) 2861–2886 [MR](#)
- [6] **A Casson, M Freedman**, *Atomic surgery problems*, from “Four-manifold theory” (C Gordon, R Kirby, editors), Contemp. Math. 35, Amer. Math. Soc., Providence, RI (1984) 181–199 [MR](#)
- [7] **J C Cha**, *The structure of the rational concordance group of knots*, Mem. Amer. Math. Soc. 885, Amer. Math. Soc., Providence, RI (2007) [MR](#)
- [8] **T D Cochran, C W Davis**, *Counterexamples to Kauffman’s conjectures on slice knots*, Adv. Math. 274 (2015) 263–284 [MR](#)
- [9] **T D Cochran, C W Davis, A Ray**, *Injectivity of satellite operators in knot concordance*, J. Topol. 7 (2014) 948–964 [MR](#)
- [10] **T D Cochran, B D Franklin, M Hedden, P D Horn**, *Knot concordance and homology cobordism*, Proc. Amer. Math. Soc. 141 (2013) 2193–2208 [MR](#)
- [11] **T D Cochran, R E Gompf**, *Applications of Donaldson’s theorems to classical knot concordance, homology 3-spheres and property P*, Topology 27 (1988) 495–512 [MR](#)
- [12] **T D Cochran, S Harvey, C Leidy**, *2-torsion in the  $n$ -solvable filtration of the knot concordance group*, Proc. Lond. Math. Soc. 102 (2011) 257–290 [MR](#)
- [13] **T D Cochran, S Harvey, C Leidy**, *Primary decomposition and the fractal nature of knot concordance*, Math. Ann. 351 (2011) 443–508 [MR](#)
- [14] **J Collins**, *On the concordance orders of knots*, PhD thesis, University of Edinburgh (2012) [arXiv](#)
- [15] **R H Fox, J W Milnor**, *Singularities of 2-spheres in 4-space and equivalence of knots*, Bull. Amer. Math. Soc. 63 (1957) 406
- [16] **R H Fox, J W Milnor**, *Singularities of 2-spheres in 4-space and cobordism of knots*, Osaka J. Math. 3 (1966) 257–267 [MR](#)
- [17] **M H Freedman, F Quinn**, *Topology of 4-manifolds*, Princeton Mathematical Series 39, Princeton Univ. Press (1990) [MR](#)
- [18] **S L Harvey**, *Homology cobordism invariants and the Cochran–Orr–Teichner filtration of the link concordance group*, Geom. Topol. 12 (2008) 387–430 [MR](#)
- [19] **M Hedden, P Kirk**, *Instantons, concordance, and Whitehead doubling*, J. Differential Geom. 91 (2012) 281–319 [MR](#)

- [20] **C Kearton**, *The Milnor signatures of compound knots*, Proc. Amer. Math. Soc. 76 (1979) 157–160 [MR](#)
- [21] **R C Kirby**, editor, *Problems in low-dimensional topology*, from “Geometric topology, II” (W H Kazez, editor), AMS/IP Stud. Adv. Math. 2, Amer. Math. Soc., Providence, RI (1997) 35–473 [MR](#)
- [22] **P Kirk**, **C Livingston**, *Twisted knot polynomials: inversion, mutation and concordance*, Topology 38 (1999) 663–671 [MR](#)
- [23] **A S Levine**, *Nonsurjective satellite operators and piecewise-linear concordance*, Forum Math. Sigma 4 (2016) art. id. e34 [MR](#)
- [24] **W B R Lickorish**, *An introduction to knot theory*, Graduate Texts in Mathematics 175, Springer (1997) [MR](#)
- [25] **C Livingston**, *The slicing number of a knot*, Algebr. Geom. Topol. 2 (2002) 1051–1060 [MR](#)
- [26] **C Livingston**, *The concordance genus of knots*, Algebr. Geom. Topol. 4 (2004) 1–22 [MR](#)
- [27] **C Livingston**, *The stable 4-genus of knots*, Algebr. Geom. Topol. 10 (2010) 2191–2202 [MR](#)
- [28] **S Naik**, *Casson–Gordon invariants of genus one knots and concordance to reverses*, J. Knot Theory Ramifications 5 (1996) 661–677 [MR](#)
- [29] **A Ray**, *Satellite operators with distinct iterates in smooth concordance*, Proc. Amer. Math. Soc. 143 (2015) 5005–5020 [MR](#)
- [30] **Y Sato**, *3–dimensional homology handles and minimal second Betti numbers of 4–manifolds*, Osaka J. Math. 35 (1998) 509–527 [MR](#)
- [31] **L R Taylor**, *On the genera of knots*, from “Topology of low-dimensional manifolds” (R A Fenn, editor), Lecture Notes in Math. 722, Springer (1979) 144–154 [MR](#)
- [32] **A G Tristram**, *Some cobordism invariants for links*, Proc. Cambridge Philos. Soc. 66 (1969) 251–264 [MR](#)

Department of Mathematics, Rice University  
Houston, TX, United States

Department of Mathematics, Rice University  
Houston, TX, United States

[shelly@rice.edu](mailto:shelly@rice.edu)

Received: 16 September 2014      Revised: 15 May 2017