

MUTUALLY CATALYTIC BRANCHING IN THE PLANE: FINITE MEASURE STATES

BY DONALD A. DAWSON,¹ ALISON M. ETHERIDGE,²
KLAUS FLEISCHMANN,³ LEONID MYTNIK,⁴
EDWIN A. PERKINS⁵ AND JIE XIONG

*Carleton University, University of Oxford, Weierstrass Institute for Applied
Analysis and Stochastics, Technion—Israel Institute of Technology,
University of British Columbia and University of Tennessee*

We study a pair of populations in \mathbb{R}^2 which undergo diffusion and branching. The system is interactive in that the branching rate of each type is proportional to the local density of the other type. For a diffusion rate sufficiently large compared with the branching rate, the model is constructed as the unique pair of finite measure-valued processes which satisfy a martingale problem involving the collision local time of the solutions. The processes are shown to have densities at fixed times which live on disjoint sets and explode as they approach the interface of the two populations. In the long-term limit, global extinction of one type is shown. The process constructed is a rescaled limit of the corresponding \mathbb{Z}^2 -lattice model studied by D. A. Dawson and E. A. Perkins [*Ann. Probab.* **26** (1998) 1088–1138] and resolves the large scale mass–time–space behavior of that model.

Contents

1. Introduction and statement of results
 - 1.1. Background and motivation
 - 1.2. A martingale problem for mutually catalytic branching
 - 1.3. Segregated densities
 - 1.4. Global extinction of one type
2. Preliminaries
 - 2.1. Green function representation
 - 2.2. First and second moments: proof of Proposition 15
 - 2.3. State spaces for \mathbf{X}
3. Dual processes for higher moments

Received September 2000; revised May 2002.

¹Supported in part by an NSERC research grant and a Max Planck award.

²Supported in part by an EPSRC advanced fellowship.

³Supported in part by the DFG.

⁴Supported in part by U.S.–Israel Binational Science Foundation Grant 2000065 and Israel Science Foundation Grant 116/01-10.0.

⁵Supported in part by an NSERC research grant.

AMS 2000 subject classifications. Primary 60K35; secondary 60G57, 60J80.

Key words and phrases. Catalytic super-Brownian motion, catalytic super-random walk, collision local time, duality, superprocesses, martingale problem, segregation of types, stochastic PDE.

- 3.1. Lattice approximation moment dual V^ε and self-duality
- 3.2. Limiting moment dual V
- 4. Construction of a solution
- 5. Long-term behavior
- 6. Existence of densities and segregation of types
- 7. Some open questions
- Appendix A. Random walk kernels
- Appendix B. Integration lemmas
- Acknowledgments
- References

1. Introduction and statement of results.

1.1. *Background and motivation.* In [14] solutions to the following system of stochastic partial differential equations (SPDEs) were studied:

$$(1) \quad \frac{\partial}{\partial t} X_t^i(x) = \frac{\sigma^2}{2} \Delta X_t^i(x) + \sqrt{\gamma X_t^1(x) X_t^2(x)} \dot{W}_t^i(x),$$

$(t, x) \in \mathbb{R}_+ \times \mathbb{R}$, $i = 1, 2$. Here Δ is the one-dimensional Laplacian, σ, γ are (strictly) positive constants (the migration and collision rate, respectively) and \dot{W}^1, \dot{W}^2 are independent standard time–space white noises on $\mathbb{R}_+ \times \mathbb{R}$. Our goal is to study the same system of equations for $x \in \mathbb{R}^2$. As we explain below, from one point of view, existence in two dimensions appears to be counterintuitive. This was one reason six different people were attracted to this question and finally combined their efforts.

Recall that

$$(2) \quad \frac{\partial}{\partial t} X_t(x) = \frac{\sigma^2}{2} \Delta X_t(x) + \sqrt{\rho_t(x) X_t(x)} \dot{W}_t(x) \quad \text{on } \mathbb{R}_+ \times \mathbb{R}$$

is the stochastic partial differential equation for the density of a one-dimensional *super-Brownian motion* (SBM) [31, 38] with branching rate at time t at x equal to $\rho_t(x)$ (bounded in t and x). As a measure-valued process it arises as the large population (N particles), small mass (N^{-1}) per particle limit of a system of critical binary branching Brownian motions with diffusion rate σ^2 which branch at rate $N\rho_t(x)$ at site x at time t . Equivalently each Brownian particle with path $s \mapsto \xi_s$ branches according to the additive functional $t \mapsto N \int_0^t ds \rho_s(\xi_s)$. Although the limit exists in higher dimensions as the unique solution of an appropriate martingale problem, the resulting process takes values in the space of singular measures and it is easy to use this fact to see that (2) has no solutions in higher dimensions (see [15], Remark 1.4). The problem is that in higher dimensions the critical branching (which tends to cluster the population on a small set) overpowers the diffusion. This situation is typical of parabolic SPDEs driven by white noise: solutions seem to only exist in one spatial dimension (see [45]).

One way to rectify this situation in the branching context is to replace $[\rho_t(x) dx, t \geq 0]$ by a collection of singular measures, that is, have the branching

only take place on singular sets. Delmas [16] showed if the branching takes place on a Lebesgue null set (the catalyst) independent of time and satisfies a mild regularity condition guaranteeing that the null set is not polar for Brownian motion (more precisely, particles branch according to an additive functional with Revuz measure supported by this null set), then the associated super-Brownian motion (reactant) has a density at all times with probability 1.

A particular time-dependent case was introduced by Dawson and Fleischmann [10] and different aspects of this model were investigated in [11, 17, 21, 26, 27]. In this model the catalyst itself is a super-Brownian motion ρ and the resulting reactant model X^ρ exists and has a nice density in three dimensions and less. In higher dimensions an intrinsic Brownian reactant particle's path will not hit the support of an independent super-Brownian catalyst and hence the reactant process degenerates into heat flow as there can be no branching. The construction of such a model poses no difficulties in principle as one first constructs the super-Brownian catalyst and then builds a super-Brownian motion (reactant) whose branching rate is governed by this catalyst.

The situation in (1) is quite different as one has a truly interacting system consisting of two types in which the branching rate of one type is given by the local density of mass of the other; that is, each type catalyzes the branching of the other. It is a natural question as to whether such an interacting system exists in \mathbb{R}^d for $d > 1$ (recall for instance that the super-Brownian reactant [10] exists nontrivially only in dimensions $d \leq 3$, and that the continuum stepping stone [41] exists only in $d = 1$).

Let $S(\mu)$ denote the closed support of a measure μ . Assume for the moment that $\mathbf{X} = (X^1, X^2)$ is a solution to (1) for $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$, where the \dot{W}^1, \dot{W}^2 are independent white noises on $\mathbb{R}_+ \times \mathbb{R}^2$. Then the singularity of ordinary (two-dimensional) SBM (or of SBM with a strictly positive branching rate) suggests that $S(X_t^1) \cap S(X_t^2)$ is Lebesgue null, and the requirement in (1) that X^i solves the heat equation away from this null set shows that X_t^i should have a density away from this null set. In fact this would suggest that $X_t^1(x)X_t^2(x) = 0$ for almost all x and so (1) degenerates into a pair of heat flows which of course do not solve (1).

To circumvent this nonexistence argument, we will work with the following martingale problem formulation of (1) in two dimensions. We write $\langle \mu, \varphi \rangle$ to denote the integral of a function φ with respect to a measure μ . For fixed constants $\sigma, \gamma > 0$, let $\mathbf{X} = (X^1, X^2)$ be a pair of continuous measure-valued processes such that, for an appropriate class of test functions φ_i ,

$$(3) \quad M_t^i(\varphi_i) := \langle X_t^i, \varphi_i \rangle - \langle \mu^i, \varphi_i \rangle - \int_0^t ds \left\langle X_s^i, \frac{\sigma^2}{2} \Delta \varphi_i \right\rangle,$$

$t \geq 0$, $i = 1, 2$, are orthogonal continuous square integrable martingales starting from 0 at time $t = 0$ and with continuous square function

$$(4) \quad \langle \langle M^i(\varphi_i) \rangle \rangle_t = \gamma \int_{[0, t] \times \mathbb{R}^2} L_{\mathbf{X}}(d(s, x)) \varphi_i^2(x).$$

Here $L_{\mathbf{X}}$ is the collision local time of X^1 and X^2 , loosely described by

$$(5) \quad L_{\mathbf{X}}(d(s, x)) = ds X_s^1(dx) \int_{\mathbb{R}^2} X_s^2(dy) \delta_x(y)$$

(a precise description is given in Definition 1 below via a smoothing procedure). It is not hard to see that *if* a solution to (1) (for two dimensions) is locally bounded (in both space and time) and has the appropriate square integrability properties then the associated measure-valued processes will satisfy (3) and (4), and so the above martingale problem is a natural generalization of (1). We will show (see Theorems 11 and 17) that, under appropriate conditions on the finite initial measures and for γ/σ^2 sufficiently small, solutions to this martingale problem exist and satisfy the intuitive description given in the paragraph prior to (3): each population X_t^i has a density denoted by the same symbol X_t^i , and $X_t^1(x)X_t^2(x) = 0$ for Lebesgue-a.a. x . Indeed we will give an explicit expression for the joint law of these densities for fixed values of t and x (see Theorem 17). Evidently these densities cannot be locally bounded since in that case we can easily show that

$$(6) \quad L_{\mathbf{X}}([0, \infty) \times \mathbb{R}^2) = \int_0^\infty ds \int_{\mathbb{R}^2} dx X_s^1(x)X_s^2(x) = 0 \quad \text{a.s.},$$

and again our solutions become a pair of solutions to the heat equation; hence $L_{\mathbf{X}}([0, \infty) \times \mathbb{R}^2) > 0$, contradicting (6). In fact, we will show that each of these densities becomes unbounded near any point in the interface of the two types given by the support of the collision local time (Corollary 19). This bad behavior of the densities near the interface is borne out by simulations of Achim Klenke which you can find on his web page <http://www.aklenke.de/~klenke>.

There are a number of reasons to study mutually catalytic, or symbiotic, branching models such as (1), (3) or their discrete counterparts. Diploid organisms require the presence of two types for reproduction. There are of course a number of features of these models which are biologically inaccurate (e.g., males and females do not, we believe, live forever if they avoid contact with the other sex). The deterministic analogue of mutually catalytic branching was introduced in the work of Eigen [19, 20] to describe the catalytic growth of self-reproducing molecules. He considered a closed chain of K equations (called a catalytic hypercycle):

$$(7) \quad \dot{y}_i = k_i y_i y_{i-1}, \quad i = 0, \dots, K - 1,$$

where the arithmetic is done mod K . The special case $K = 2$ is a deterministic growth model analogue of mutually catalytic branching. Work on the generalization of our model to K types has already been carried out by Fleischmann and Xiong [28]. Mutually catalytic branching is a fixed point under a renormalization scheme as is the case for the Fisher–Wright and continuous state branching diffusions. This suggests that it might also be the attracting element of a universality class for two component systems on the lattice. However, the question of identification and domains of attraction of universality classes of two component systems including mutually catalytic branching is considerably more involved (see [6])

for a discussion of this question). In the general context of interactive branching measure-valued diffusions the study of interactive branching mechanisms has proved to be a difficult problem. Singular interactions such as those considered here have proved to be particularly challenging and it is perhaps rather surprising that we can say so much about the mutually catalytic branching processes on the plane in light of the difficulties encountered in the study of branching diffusions with singular interactions in the spatial motion or growth rates (e.g., [23]).

The basic uniqueness questions for interactive branching models in which the branching rate depends on the current state of the system remain unresolved in general in spite of recent progress on uniqueness for a variety of interactive branching models (e.g., [1, 4, 13, 18, 36]). In the one-dimensional case (1), Mytnik [35] obtained uniqueness by an exponential self-duality argument. It will be more difficult to implement this approach here due to the bad behavior of the densities. Nevertheless, the problem of uniqueness will be resolved in a companion paper [12] under an additional integrability condition (**IntC**) involving the trajectories of \mathbf{X} , introduced in Definition 7 below. In the latter paper this condition will be verified for the solutions constructed in Theorem 11 by means of the moment calculations in Section 3 which are carried out in terms of a function-valued dual. We state the uniqueness result and associated Markov property as Theorem 11(b) as it will play an important role in our study of the longtime behavior of the solutions (Theorem 21) and the proof of segregation of the two populations [Theorem 17(b)].

The existence of our solutions will be established by means of rescaling the lattice versions of (1), constructed in [14] (in any number of dimensions). We will use the moment bounds in Sections 3 and 4 (for finite initial conditions satisfying a suitable energy condition) to establish tightness of these rescaled processes providing γ/σ^2 is small enough. This restriction on the parameters is needed to ensure that the higher (specifically fourth) moments used in the tightness arguments are finite. It is not hard to show that the approximating fourth moments blow up for γ/σ^2 large enough, but we have not tried to find the best value of this ratio here. We conjecture that solutions to (3) and (4) should exist for any positive values of γ and σ . This is because $2 + \delta$ moments should suffice and as $\delta \rightarrow 0$, this should allow any values of these parameters. The situation in higher dimensions is intriguing and unresolved.

Many of the results of this paper had been obtained independently and at the same time by two subgroups of the present authors and others were obtained after we coalesced.

The present paper is completely restricted to the finite measure-valued case. For the infinite measure case, we refer to our forthcoming paper [8].

1.2. *A martingale problem for mutually catalytic branching.* We start by formulating our martingale problem for finite measures. Let $\mathcal{M}_f = \mathcal{M}_f(\mathbb{R}^2)$ denote the space of finite measures on the Borel subsets $\mathcal{B}(\mathbb{R}^2)$ of \mathbb{R}^2 , with the topology

of weak convergence. $C_b(\mathbb{R}^2)$ is the space of bounded continuous functions on \mathbb{R}^2 with the supnorm $\|\cdot\|_\infty$ topology, and $C_b^n(\mathbb{R}^2)$ is the subspace consisting of those functions whose partial derivatives of order n or less are also in C_b (n could be a natural number or ∞). We let $\mathcal{C}_{\text{com}} = \mathcal{C}_{\text{com}}(\mathbb{R}^2)$ denote the space of continuous function on \mathbb{R}^2 with compact support. γ and σ are fixed positive constants. Write $(\xi, \Pi_x, x \in \mathbb{R}^2)$ for the Brownian motion in \mathbb{R}^2 with variance parameter σ^2 ,

$$(8) \quad p_t(x, y) := \frac{1}{2\pi\sigma^2 t} \exp\left[-\frac{|y-x|^2}{2\sigma^2 t}\right], \quad t > 0, x, y \in \mathbb{R}^2,$$

for its transition density ($|\cdot|$ denotes the Euclidean norm), and $\{S_t : t \geq 0\}$ for the corresponding semigroup. If μ is a measure on \mathbb{R}^2 , set $S_t\mu(x) := \int d\mu(y) p_t(x, y)$.

DEFINITION 1 (Collision local time). Let $\mathbf{X} = (X^1, X^2)$ denote an \mathcal{M}_f^2 -valued continuous process, where $\mathcal{M}_f^2 = \mathcal{M}_f \times \mathcal{M}_f$. The collision local time of \mathbf{X} (if it exists) is a continuous nondecreasing \mathcal{M}_f -valued stochastic process $t \mapsto L_{\mathbf{X}}(t) = L_{\mathbf{X}}(t, \cdot)$ such that

$$(9) \quad \langle L_{\mathbf{X}}^{*,\delta}(t), \varphi \rangle \rightarrow \langle L_{\mathbf{X}}(t), \varphi \rangle \quad \text{as } \delta \downarrow 0 \text{ in probability,}$$

for all $t > 0$ and $\varphi \in \mathcal{C}_{\text{com}}(\mathbb{R}^2)$, where

$$(10) \quad L_{\mathbf{X}}^{*,\delta}(t, dx) := \frac{1}{\delta} \int_0^\delta dr \int_0^t ds S_r X_s^1(x) S_r X_s^2(x) dx, \quad t \geq 0, \delta > 0.$$

The collision local time $L_{\mathbf{X}}$ will also be considered as a (locally finite) measure $L_{\mathbf{X}}(ds, dx)$ on $\mathbb{R}_+ \times \mathbb{R}^2$.

Note that we used an additional smoothing in time in defining the collision local time, compared with other sources as, for example, [2]. Clearly if it exists as in [2], it will exist in the above sense and the processes will coincide. It is also easy to see that the above definition is independent of the choice of $\sigma^2 > 0$. If $X_t^i(dx) = X_t^i(x) dx$ for some bounded densities $X_t^i(x)$, then it is easy to see that $L_{\mathbf{X}}(t)(dx) = \int_0^t X_s^1(x) X_s^2(x) ds dx$. At the end of the next section we give a simple deterministic example of a pair of unbounded densities for which this equality fails as the product of the densities vanishes but the collision local time of the corresponding measures is nonzero.

All filtrations will be assumed to be right-continuous and contain the null sets at time 0.

DEFINITION 2 [Martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$]. A continuous \mathcal{F} -adapted and $\mathcal{M}_f^2(\mathbb{R}^2)$ -valued process $\mathbf{X} = (X^1, X^2)$ on some probability space $(\Omega, \mathcal{F}, \mathcal{F}, P)$ is said to satisfy the martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$, if for all $\varphi_i \in C_b^2(\mathbb{R}^2)$, $i = 1, 2$,

$$(11) \quad M_t^i(\varphi_i) = \langle X_t^i, \varphi_i \rangle - \langle X_0^i, \varphi_i \rangle - \int_0^t ds \left\langle X_s^i, \frac{\sigma^2}{2} \Delta \varphi_i \right\rangle, \quad t \geq 0, i = 1, 2,$$

are orthogonal continuous $L^2 \mathcal{F}$ -martingales such that $M_0^i(\varphi_i) = 0$ and

$$(12) \quad \langle \langle M^i(\varphi_i) \rangle \rangle_t = \gamma \langle L_{\mathbf{X}}(t), \varphi_i^2 \rangle, \quad t \geq 0, \quad i = 1, 2.$$

Note that in this definition the initial state \mathbf{X}_0 may be random. To construct solutions to this martingale problem we will need to impose a bivariate regularity condition on the initial state.

NOTATION 3 (Energy function). Introduce the energy function

$$(13) \quad g(x_1, x_2) := 1 + \log^+ \frac{1}{|x_2 - x_1|}, \quad x_1, x_2 \in \mathbb{R}^2,$$

(recall that $|\cdot|$ denotes the Euclidean norm).

DEFINITION 4 (State space versions).

(a) *Energy condition*—Write $\mu = (\mu^1, \mu^2) \in \mathcal{M}_{f,e}$ and say μ satisfies the energy condition, iff $\mu \in \mathcal{M}_f^2(\mathbb{R}^2)$ and

$$(14) \quad \|\mu\|_g := \langle \mu^1 \times \mu^2, g \rangle < \infty.$$

(b) *Strong energy condition*—Write $\mu = (\mu^1, \mu^2) \in \mathcal{M}_{f,se}$ and say μ satisfies the strong energy condition, iff $\mu \in \mathcal{M}_f^2(\mathbb{R}^2)$ and for any $p \in (0, 1)$, there is a constant $c = c(p, \mu)$ such that

$$(15) \quad \max_{1 \leq i, j \leq 2} \langle \mu^i \times \mu^j, p_r \rangle \leq cr^{-p}, \quad r > 0.$$

REMARK 5. (a) Inequality (15) is trivially fulfilled for $r \geq 1$, and so we only need to consider $0 < r < 1$. By an elementary interpolation argument it actually suffices to consider only $r = 2^{-n}$ and so $\mathcal{M}_{f,e}$ is clearly a Borel subset of \mathcal{M}_f^2 .

(b) An elementary calculation shows that for all $T > 0$ there are constants c_T and C_T such that

$$(16) \quad c_T g \leq 1 + \int_0^T dr p_r \leq C_T g.$$

In particular, by (15),

$$(17) \quad \mathcal{M}_{f,se} \subseteq \mathcal{M}_{f,e}.$$

Next we introduce a *lattice system of approximating processes* we will use to construct solutions to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$.

Fix a deterministic $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ and $\varepsilon \in (0, 1]$. Set

$$(18) \quad X_0^{i,\varepsilon}(x) = \varepsilon^{-2} X_0^i(\varepsilon x + [0, \varepsilon)^2), \quad x = (x_1, x_2) \in \mathbb{Z}^2, \quad i = 1, 2.$$

Let $\{W^i(x) : x \in \mathbb{Z}^2, i = 1, 2\}$ be a collection of independent standard one-dimensional Brownian motions on $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$, and consider the unique (in law) solution of

$$(19) \quad X_t^{i,\varepsilon}(x) = X_0^{i,\varepsilon}(x) + \int_0^t ds \frac{\sigma^2}{2} {}^1\Delta X_s^{i,\varepsilon}(x) + \int_0^t dW_s^i(x) \sqrt{\gamma X_s^{1,\varepsilon}(x) X_s^{2,\varepsilon}(x)},$$

$i = 1, 2, t \geq 0, x \in \mathbb{Z}^2$. Here ${}^1\Delta$ is the discrete Laplacian on \mathbb{Z}^2 defined in (21) below. See ([14], Theorems 2.2 and 2.4) for the existence and uniqueness of these solutions.

Via scaling we pass to processes indexed by $\varepsilon\mathbb{Z}^2$ (instead of \mathbb{Z}^2):

$$(20) \quad {}^\varepsilon X_t^i(x) := X_{t\varepsilon^{-2}}^{i,\varepsilon}(x\varepsilon^{-1}), \quad i = 1, 2, t \geq 0, x \in \varepsilon\mathbb{Z}^2.$$

Write $x \sim^\varepsilon y$ if x and y are neighbors in $\varepsilon\mathbb{Z}^2$, and introduce the discrete Laplacian on $\varepsilon\mathbb{Z}^2$:

$$(21) \quad {}^\varepsilon\Delta\varphi(x) := \sum_{y \sim^\varepsilon x} \frac{\varphi(y) - \varphi(x)}{\varepsilon^2}, \quad x \in \varepsilon\mathbb{Z}^2.$$

If $\ell^\varepsilon := \sum_{y \in \varepsilon\mathbb{Z}^2} \varepsilon^2 \delta_y$ and $d^\varepsilon x$ denotes integration with respect to ℓ^ε , let ${}^\varepsilon\mathcal{M}_f(\mathbb{R}^2)$ denote the subspace of $\mathcal{M}_f(\mathbb{R}^2)$ of measures with densities with respect to ℓ^ε . Also denote by $t \mapsto {}^\varepsilon X_t^i$ the ${}^\varepsilon\mathcal{M}_f(\mathbb{R}^2)$ -valued process with densities ${}^\varepsilon X_t^i(x)$, that is,

$$(22) \quad \langle {}^\varepsilon X_t^i, \varphi \rangle = \int_{\varepsilon\mathbb{Z}^2} d^\varepsilon x {}^\varepsilon X_t^i(x) \varphi(x) = \sum_{x \in \varepsilon\mathbb{Z}^2} {}^\varepsilon X_t^i(x) \varphi(x) \varepsilon^2.$$

Then ${}^\varepsilon X_0^i(\{x\}) = X_0^i(x + [0, \varepsilon)^2)$ for $x \in \varepsilon\mathbb{Z}^2$ and so clearly these initial states satisfy ${}^\varepsilon X_0^i \rightarrow X_0^i$ in $\mathcal{M}_f(\mathbb{R}^2)$ as $\varepsilon \downarrow 0$. The following lemma can easily be derived.

LEMMA 6 [Martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon}$]. *The process ${}^\varepsilon\mathbf{X}$ on $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$ defined via (22), (20), (19) and (18), based on $\mathbf{X}_0 \in \mathcal{M}_{f,e}$, satisfies the following approximate martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon}$.*

For each pair of bounded functions $\varphi_i : \varepsilon\mathbb{Z}^2 \rightarrow \mathbb{R}, i = 1, 2,$

$$(23) \quad \langle {}^\varepsilon X_t^i, \varphi_i \rangle = \langle {}^\varepsilon X_0^i, \varphi_i \rangle + \int_0^t ds \left\langle {}^\varepsilon X_s^i, \frac{\sigma^2}{2} {}^\varepsilon\Delta\varphi_i \right\rangle + {}^\varepsilon M_t^i(\varphi_i),$$

where

$$(24) \quad {}^\varepsilon M_t^i(\varphi_i) = \int_{\varepsilon\mathbb{Z}^2} d^\varepsilon x \varphi_i(x) \int_0^{t\varepsilon^{-2}} dW_s^i(x\varepsilon^{-1}) \sqrt{\gamma X_s^{1,\varepsilon}(x\varepsilon^{-1}) X_s^{2,\varepsilon}(x\varepsilon^{-1})},$$

$i = 1, 2,$ are orthogonal continuous $L^2 = (\mathcal{F}_\cdot)$ -martingales such that

$$(25) \quad \langle \langle {}^\varepsilon M^i(\varphi_i) \rangle \rangle_t = \gamma \int_0^t ds \int_{\varepsilon\mathbb{Z}^2} d^\varepsilon x \varphi_i^2(x) {}^\varepsilon X_s^1(x) {}^\varepsilon X_s^2(x) =: \gamma \langle {}^\varepsilon L_{\varepsilon\mathbf{X}}(t), \varphi_i^2 \rangle,$$

$i = 1, 2.$

Existence of solutions to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ will later follow by taking a weak limit point of ${}^\varepsilon \mathbf{X}$ as $\varepsilon \downarrow 0$. Our proof of uniqueness will require an additional integrability condition:

DEFINITION 7 (Integrability conditions on path space). For $\varepsilon > 0$ and a pair $\mu = (\mu^1, \mu^2)$ of measures in $\mathcal{M}_f(\mathbb{R}^2)$, we write

$$(26) \quad H_\varepsilon(\mu) := \int_{\mathbb{R}^2} dx \int_{\mathbb{R}^2} dy \left[1 + \frac{1}{|x - y|} \right] S_\varepsilon \mu^1(x) S_\varepsilon \mu^2(x) S_\varepsilon \mu^1(y) S_\varepsilon \mu^2(y).$$

Integrability condition (IntC)—A continuous \mathcal{M}_f^2 -valued process $\mathbf{X} = (X^1, X^2)$ on a probability space $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$ is said to satisfy the integrability condition **(IntC)** if, for all $0 < \delta < T < \infty$,

$$E \left\{ \int_\delta^T ds H_\varepsilon(\mathbf{X}_s) \mid \mathcal{F}_\delta \right\} \quad \text{is bounded in probability as } \varepsilon \downarrow 0;$$

that is, for all $\eta > 0$, there is an M such that

$$(27) \quad \limsup_{\varepsilon \downarrow 0} P \left(E \left\{ \int_\delta^T ds H_\varepsilon(X_s) \mid \mathcal{F}_\delta \right\} > M \right) < \eta.$$

Strong integrability condition (SIntC)— \mathbf{X} is said to satisfy the stronger (and simpler) integrability condition **(SIntC)** if

$$(28) \quad \limsup_{\varepsilon \downarrow 0} E \int_0^T ds H_\varepsilon(\mathbf{X}_s) < \infty, \quad T > 0.$$

To describe the restriction on γ/σ^2 , let $({}^\varepsilon \xi, \Pi_x^\varepsilon, x \in \varepsilon \mathbb{Z}^2)$ denote the continuous time simple symmetric random walk on $\varepsilon \mathbb{Z}^2$ with generator $\frac{\sigma^2}{2} \varepsilon \Delta$. That is, ${}^\varepsilon \xi$ jumps to a nearest neighbor site at rate $2\varepsilon^{-2}\sigma^2$. Introduce the corresponding transition density ${}^\varepsilon p_t(x, y) = \varepsilon^{-2} \Pi_x({}^\varepsilon \xi_t = y)$, $x, y \in \varepsilon \mathbb{Z}^2$, with respect to ℓ^ε , and $\{{}^\varepsilon S_t : t \geq 0\}$ the related semigroup.

The following elementary result is proved in Appendix A for the sake of completeness.

LEMMA 8 (Random walk kernel estimates). (a) *Local central limit theorem*—For all $s > 0$, with the heat kernel \mathbf{p} from (8),

$$(29) \quad \lim_{\varepsilon \rightarrow 0} \sup_{x, y \in \varepsilon \mathbb{Z}^2} |{}^\varepsilon p_s(x, y) - \mathbf{p}_s(x, y)| = 0.$$

(b) *Uniform bound*—There is a universal constant c_{rw} (independent of σ^2) such that

$$(30) \quad \sup_{s \geq 0, x, y \in \mathbb{Z}^2} \varepsilon p_s(x, y) s \sigma^2 = \sup_{s \geq 0} \varepsilon p_s(0, 0) s \sigma^2 = c_{\text{rw}},$$

for all $\varepsilon > 0$.

REMARK 9 (Size of c_{rw}). Statement (a) is of course a standard local central limit theorem. The value of the constant c_{rw} of (b) enters in Theorem 11 below. To estimate its value, write \tilde{p} instead of εp in the case $\varepsilon = \sigma = 1$. Then

$$(31) \quad c_{\text{rw}} = \sup_{t \geq 0} t \tilde{p}_t(0, 0).$$

Now a direct calculation and exploiting Stirling’s formula (see [24], page 52) gives $c_{\text{rw}} \leq e^{1/12}/2 < 0.55$. On the other hand, $c_{\text{rw}} \geq \varepsilon p_t(0, 0) t \sigma^2$, and it follows from (a) that

$$(32) \quad c_{\text{rw}} \geq t p_t(0, 0) = (2\pi)^{-1} > 0.15.$$

Consequently, $c_{\text{rw}} \in (0.15, 0.55)$.

NOTATION 10 (Path space). Let $\Omega_\circ := C(\mathbb{R}_+, \mathcal{M}_f^2(\mathbb{R}^2))$ with the usual topology of uniform convergence on compact subsets of \mathbb{R}_+ .

Recall the spaces $\mathcal{M}_{f,e}$ and $\mathcal{M}_{f,se}$ introduced in Definition 4.

THEOREM 11 (Mutually catalytic SBM in \mathbb{R}^2). Assume

$$(33) \quad \gamma/\sigma^2 < (3\sqrt{6\pi} c_{\text{rw}})^{-1}$$

and $\mathbf{X}_0 \in \mathcal{M}_{f,e}$:

(a) *Existence*—There is a process \mathbf{X} on some $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$ satisfying the martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ and the integrability condition (\mathbf{IntC}) , and such that $\mathbf{X}_t \in \mathcal{M}_{f,e}$ for all $t \geq 0$ a.s. If moreover $\mathbf{X}_0 \in \mathcal{M}_{f,se}$, then \mathbf{X} will satisfy (\mathbf{SIntC}) .

(b) *Strong Markov and uniqueness*—There is a (time-homogeneous) Borel Markov transition kernel $\mathbf{P} = \{\mathbf{P}_t(\mu, d\nu) : t > 0, \mu \in \mathcal{M}_{f,e}\}$ on $\mathcal{M}_{f,e}$ such that any process satisfying $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ and (\mathbf{IntC}) on $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$ is (\mathcal{F}_\cdot) -strong Markov with transition kernel \mathbf{P} . In particular, the law $P_{\mathbf{X}_0}$ on Ω_0 of the solution in (a) is unique.

(c) *Lattice approximation*—Let ${}^\varepsilon \mathbf{X}$ denote the lattice system of approximating processes given by (19), (20), with initial conditions (18) and let ${}^\varepsilon L_{\varepsilon \mathbf{X}}$ be as defined in Lemma 6. As $\varepsilon \downarrow 0$,

$$(34) \quad P(({}^\varepsilon \mathbf{X}, {}^\varepsilon L_{\varepsilon \mathbf{X}}) \in \cdot) \Rightarrow P((\mathbf{X}, L_{\mathbf{X}}) \in \cdot)$$

weakly on $C(\mathbb{R}_+, \mathcal{M}_f^3(\mathbb{R}^2))$, where \mathbf{X} satisfies (\mathbf{IntC}) and is a solution to the martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ with $L_{\mathbf{X}}$ as its collision local time.

(d) *Scaling property*—Assume that \mathbf{X} satisfies $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ and (\mathbf{IntC}) , $\varepsilon, \theta > 0$, $z \in \mathbb{R}^2$ and $\widehat{X}_t^i(A) := \theta X_{\varepsilon^2 t}^i(z + \varepsilon A)$, $t \geq 0$, $A \in \mathcal{B}(\mathbb{R}^2)$, $i = 1, 2$. Then $(\widehat{X}^1, \widehat{X}^2)$ satisfies $(\mathbf{MP})_{\widehat{\mathbf{X}}_0}^{\sigma, \gamma}$ and (\mathbf{IntC}) and so has law $P_{\widehat{\mathbf{X}}_0}$.

The proof of (b) will be completed in a companion paper [12], but much of the groundwork is laid in Section 3. The verification of the integrability conditions (\mathbf{IntC}) and (\mathbf{SIntC}) is also deferred to [12] as its main use is the proof of (b) [although (\mathbf{SIntC}) is also used in our description of the long-term behavior (Theorem 21)]. The main ingredient in the proof of (\mathbf{IntC}) is a bound on its conditional fourth moments in terms of a function-valued dual (Theorem 53).

REMARK 12. (i) Part (c) remains true for a wider class of lattice approximations of the initial measure. It suffices that ${}^\varepsilon \mathbf{X}_0$ approaches \mathbf{X}_0 weakly and satisfies the conclusions of Lemmas 35 and 45(a) below.

(ii) Part (a) of Theorem 11 is valid if we only assume $\gamma/\sigma^2 < 2/\sqrt{6}$. To allow for this weaker condition, solutions may be constructed as limits as $\varepsilon \downarrow 0$ of smoothed models in \mathbb{R}^2 in which the branching rate of type i at time t at site x is $dx \int_{\mathbb{R}^2} X_t^j(dy) p_\varepsilon(x, y)$ (where $j \neq i$), instead of $X_t^j(dx)$. The proof in fact is simpler than that for our lattice approximation but the latter is in many ways more natural and is used in [8] to shed some light on the large mass–time–space behavior of the lattice systems studied in [14]. Part (b) remains valid for $\gamma/\sigma^2 < 1/\sqrt{6}$.

(iii) The space $\mathcal{M}_{f,se}$ seems to be needed to get unconditional fourth moment bounds (see, e.g., Theorem 54) and a simple second moment argument shows that $X_t \in \mathcal{M}_{f,se}$ a.s. $\forall t > 0$ [see Proposition 25(a) below]. We have not, however, been able to show $X_t \in \mathcal{M}_{f,se} \forall t > 0$ a.s. and this leads to an additional conditioning argument in our construction and the use of the larger $\mathcal{M}_{f,e}$ as our state space.

(iv) It is easy to extend all our results to populations X^1, X^2 with distinct branching rates γ_1 and γ_2 , respectively, since $(\sqrt{\gamma_2} X^1, \sqrt{\gamma_1} X^2)$ will then satisfy $(\mathbf{MP})^{\sigma, \gamma}$ with $\gamma = \sqrt{\gamma_1 \gamma_2}$. The situation for distinct diffusion rates σ_1^2 and σ_2^2 is not as straightforward. If (33) holds with $\sigma^2 = \min(\sigma_1^2, \sigma_2^2)$, then the proof of (a) and (c) given below is readily modified to show that the approximating lattice systems are tight and all limit points provide solutions to the corresponding martingale problem in (a). The proof of (b), however, is no longer valid as the underlying exponential duality breaks down. This also invalidates the proof of Theorem 17 (existence of densities and segregation of types). (See also the end of this section for more on this setting.)

We now state the key self-duality result, Proposition 1.14 from [12], both because it is used below in the proof of Theorem 17(b) in Section 6 and because its proof uses our existence results Theorem 11(a).

PROPOSITION 13. Assume (33), $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ and $\tilde{\mathbf{X}}_0 = (\tilde{x}_0^1(x), \tilde{x}_0^2(x))$, where \tilde{x}_0^i is bounded, nonnegative, integrable and continuous. Then

$$\begin{aligned} & P_{\mathbf{X}_0}(\exp\{-\langle X_t^1 + X_t^2, \tilde{x}_0^1 + \tilde{x}_0^2 \rangle + i\langle X_t^1 - X_t^2, \tilde{x}_0^1 - \tilde{x}_0^2 \rangle\}) \\ &= \lim_{\varepsilon \downarrow 0} P_{\tilde{\mathbf{X}}_0}(\exp\{-\langle X_0^1 + X_0^2, S_\varepsilon \tilde{X}_t^1 + S_\varepsilon \tilde{X}_t^2 \rangle + i\langle X_0^1 - X_0^2, S_\varepsilon \tilde{X}_t^1 - S_\varepsilon \tilde{X}_t^2 \rangle\}). \end{aligned}$$

In [12] this proposition plays a major role in the proof of uniqueness in Theorem 11(b), which is assumed implicitly in our notation. The result is therefore stated there for any solution \mathbf{X} of $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ and for a particular limit point, $\tilde{\mathbf{X}}$ from Theorem 11(c). In fact, \tilde{x}_0^i need not be integrable in that setting.

We now introduce an integrability hypothesis on a possibly random initial state. Recall the norm $\|\cdot\|_g$ introduced in (14).

DEFINITION 14 [Random energy condition **(EnC)**]. We say a possibly random initial state $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ satisfies the random energy condition **(EnC)** if

$$(35) \quad \sum_{i=1,2} E\langle X_0^i, 1 \rangle^2 + E\|\mathbf{X}_0\|_g < \infty.$$

[If $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ is deterministic, then **(EnC)** clearly holds.]

Although we will need either a dual process calculation or some explicit differential equation calculations to handle some higher moments, the covariance structure of the solutions to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ only requires some integrability conditions and **(IntC)** is more than enough.

PROPOSITION 15 (First two moments). Let \mathbf{X} satisfy $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ on some filtered space $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$ for a possibly random \mathbf{X}_0 satisfying **(EnC)**:

(a) Expectation—Let $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ be a bounded Borel map. Then

$$(36) \quad E\langle X_t^i, \varphi \rangle = E\langle X_0^i, S_t \varphi \rangle < \infty, \quad t \geq 0, i = 1, 2.$$

(b) Correlation—For bounded measurable $\psi : (\mathbb{R}^2)^2 \rightarrow \mathbb{R}_+$, $t \geq 0$ and $i, j = 1, 2$,

$$\begin{aligned} E\langle X_t^i \times X_t^j, \psi \rangle &\leq E \int_{\mathbb{R}^2} dx_1 S_t X_0^i(x_1) \int_{\mathbb{R}^2} dx_2 S_t X_0^j(x_2) \psi(x_1, x_2) \\ &\quad + \delta_{ij} \gamma E \int_0^t ds \int_{\mathbb{R}^2} dx S_s X_0^1(x) S_s X_0^2(x) \\ &\quad \times \int_{\mathbb{R}^2} dx_1 p_{t-s}(x, x_1) \int_{\mathbb{R}^2} dx_2 p_{t-s}(x, x_2) \psi(x_1, x_2), \end{aligned}$$

where all expressions are finite. Moreover, equality holds if $i \neq j$.

(c) *Expected collision local time*—For measurable $\psi : \mathbb{R}_+ \times \mathbb{R}^2 \rightarrow \mathbb{R}_+$, bounded on each $[0, T] \times \mathbb{R}^2$, $T > 0$,

$$(37) \quad E \int_{[0, T] \times \mathbb{R}^2} dL_{\mathbf{X}} \psi \leq \int_0^T ds \int_{\mathbb{R}^2} dx \psi(s, x) E S_s X_0^1(x) S_s X_0^2(x) < \infty.$$

(d) *Identities under (IntC)*—If, in addition, \mathbf{X} satisfies the integrability condition **(IntC)**, then equality holds in both (b) and (c).

Note that it follows from (a) that the solution to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ constructed in Theorem 11 is not deterministic since $\langle X_t^i, \varphi \rangle \equiv \langle X_0^i, S_t \varphi \rangle$ will not satisfy $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$. Alternatively we can see from (d) that the covariance structure of this solution is not trivial.

We will now be able to state some more interesting properties of the solutions to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$. We begin by stating the absolute continuity and segregation-of-types results mentioned in the Introduction.

1.3. *Segregated densities.*

NOTATION 16 (Brownian exit time). Consider the (planar) Brownian motion $\xi = (\xi^1, \xi^2)$ with law Π_x , $x \in \mathbb{R}_+^2$, and introduce its exit time

$$(38) \quad \tau_{\text{ex}} := \inf\{t : \xi_t^1 \xi_t^2 = 0\},$$

from the first quadrant.

Let $\ell(dx) = dx$ denote Lebesgue measure. Here and elsewhere we will identify integrable functions $X(x)$ in C_b^+ with the finite absolutely continuous measure $X(x) dx$.

Here is our segregation result.

THEOREM 17 (Segregated densities). Fix $t > 0$:

(a) *Absolute continuity*—If \mathbf{X} is a solution to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ on $(\Omega, \mathcal{F}, \mathcal{F}_\cdot, P)$ with a possibly random initial condition $\mathbf{X}_0 \in \mathcal{M}_f^2(\mathbb{R}^2)$, then $X_t^i \ll \ell$ a.s. and so $X_t^i(dx) = X_t^i(x) dx$ a.s., where

$$(39) \quad X_t^i(x) = \begin{cases} \lim_{n \rightarrow \infty} S_{2^{-n}} X_t^i(x), & \text{if it exists,} \\ 0, & \text{otherwise.} \end{cases}$$

(b) *Local segregation*—Let $\mathbf{X}_0 \in M_{f,e}$ be fixed and let $\mathbf{X}_t = (X_t^1, X_t^2)$ be the functions from (39), and set $\mathbf{S}_t \mathbf{X}_0(x) := (S_t X_0^1(x), S_t X_0^2(x))$. Then the following two statements hold:

(b1) For ℓ -a.a. x ,

$$(40) \quad P_{\mathbf{X}_0}(\mathbf{X}_t(x) \in \cdot) = \Pi_{\mathbf{S}_t \mathbf{X}_0(x)}(\xi_{\tau_{\text{ex}}} \in \cdot).$$

(b2) With $P_{\mathbf{X}_0}$ -probability 1, $X_t^1(x)X_t^2(x) = 0$ for ℓ -a.a. x , and so

$$(41) \quad \int_{\mathbb{R}^2} dx X_t^1(x) X_t^2(x) = 0, \quad P_{\mathbf{X}_0}\text{-a.s.}$$

REMARK 18 (Infinite variance). (i) Note that (b1) implies

$$E_{\mathbf{X}_0}(X_t^i(x))^2 = \infty \quad \text{for } \ell\text{-a.a. } x \in \mathbb{R}_+^2 \text{ and } i = 1, 2,$$

for any $\mathbf{X}_0 \in M_{f,e}$ with $X_0^i \neq 0$, $i = 1, 2$.

(ii) It follows from (b) that the two populations segregate at each fixed time. The “interface” between the two types, although Lebesgue null must be rather active to generate a nontrivial collision local time and we show below (Corollary 19) that the densities typically explode near it. The particular distribution arising in (b1) also gave the large time limit for the lattice system (19) starting in constant initial states. In fact, the counterpart of this latter result for solutions to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ (Theorem 21 below) plays a central role in the proof. Basically a scaling argument shows that locally the joint densities $x \mapsto \mathbf{X}_t(x)$ relax to an equilibrium state instantaneously. In fact, when both types are present, the infinitely large branching rate will immediately drive one type to local extinction. The type to die is determined by the exit distribution of planar Brownian motion from the first quadrant.

Let (39) define our canonical and jointly measurable densities

$$(42) \quad X^i : \mathbb{R}_+ \times \mathbb{R}^2 \times \Omega_o \rightarrow [0, \infty), \quad i = 1, 2.$$

Let $\|X^i\|_U$ denote the essential supremum of X^i (with respect to Lebesgue measure) on the open set $U \subseteq \mathbb{R}_+ \times \mathbb{R}^2$.

COROLLARY 19 (Explosion at the interface). *If $\mathbf{X}_0 \in M_{f,e}$, then $P_{\mathbf{X}_0}$ -a.s. for any open set $U \subseteq \mathbb{R}_+ \times \mathbb{R}^2$,*

$$(43) \quad L_{\mathbf{X}}(U) > 0 \quad \text{implies} \quad \|X^1\|_U = \infty = \|X^2\|_U.$$

This corollary is proved in Section 6.

EXAMPLE 20. Here is a simple example of two measure-valued paths, constant in time, involving measures on \mathbb{R} with unbounded densities with disjoint supports, which nonetheless have a nonzero collision local time. Let $\alpha_i \in (0, 1)$ and set

$$\begin{aligned} X_t^1(dx) &= u^1(x) dx = x^{-\alpha_1} \mathbf{1}(x > 0) dx, \\ X_t^2(dx) &= u^2(x) dx = |x|^{-\alpha_2} \mathbf{1}(x < 0) dx. \end{aligned}$$

Then clearly $u^1(x)u^2(x) \equiv 0$ but if $\alpha_1 + \alpha_2 = 1$, then for $\varphi \in \mathcal{C}_{\text{com}}(\mathbb{R}^2)$,

$$\begin{aligned} & \lim_{\varepsilon \downarrow 0} \int S_\varepsilon X_t^1(x) S_\varepsilon X_t^2(x) \varphi(x) dx \\ &= \lim_{\varepsilon \downarrow 0} \iiint \varphi(\sqrt{\varepsilon} w) p_1(w - z_1) p_1(w - z_2) z_1^{-\alpha_1} |z_2|^{-\alpha_2} \\ & \quad \times \mathbf{1}(z_2 < 0 < z_1) dz_1 dz_2 dw \\ &= \varphi(0) \iint p_2(z_1 - z_2) z_1^{-\alpha_1} |z_2|^{-\alpha_2} \mathbf{1}(z_2 < 0 < z_1) dz_1 dz_2, \end{aligned}$$

where we have used dominated convergence in the last line. Therefore the collision local time, $L_{\mathbf{X}}(t)$ of \mathbf{X} is a (nonzero) constant multiple of $t\delta_0$.

1.4. *Global extinction of one type.* The one-dimensional version of the following theorem is proved in [14], Theorem 6.6.

THEOREM 21 (Global extinction of one type). *Let $\mathbf{X}_0 \in M_{f,e}$. Then*

$$(44) \quad (\langle X_t^1, 1 \rangle, \langle X_t^2, 1 \rangle) \xrightarrow[t \uparrow \infty]{} (X_\infty^1, X_\infty^2), \quad P_{\mathbf{X}_0}\text{-a.s.},$$

where

$$(45) \quad P((X_\infty^1, X_\infty^2) \in \cdot) = \Pi_{(\langle X_0^1, 1 \rangle, \langle X_0^2, 1 \rangle)}(\xi_{\text{tex}} \in \cdot).$$

The a.s. convergence is immediate from the martingale convergence theorem, as $t \mapsto \langle X_t^i, 1 \rangle$ are nonnegative martingales by $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$. The fact that $X_\infty^1 X_\infty^2 = 0$ a.s. will require a refinement of the proof for the lattice case given in [14], Theorem 1.2(b). In particular, we need to consider the rate of convergence in that result. The proof of Theorem 21 relies on the properties established in Theorem 11(a), the strong Markov property and a third moment bound (Lemma 56) which is verified for the solutions constructed in Theorem 11. If the populations have distinct diffusion rates (and so uniqueness remains open), the argument is readily modified to establish Theorem 21 for strong Markov solutions as in Theorem 11(a) satisfying the above third moment bound.

2. Preliminaries. In this section we prove Proposition 15 and identify the natural state space for \mathbf{X} .

2.1. *Green function representation.* Assume \mathbf{X} is a solution of $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ on $(\Omega, \mathcal{F}, \mathcal{F}, P)$, where \mathbf{X}_0 is an \mathcal{F}_0 -measurable $\mathcal{M}_f^2(\mathbb{R}^2)$ -valued initial state. Let \mathbf{M}_{loc} denote the space of continuous (\mathcal{F}, \cdot) -local martingales such that $M_0 = 0$ and, for $T > 0$ fixed, let $\mathbf{M}^2[0, T]$ be the space of continuous square integrable (\mathcal{F}, \cdot) -martingales on $[0, T]$, where processes which agree off an evanescent set are

identified. Let \mathbf{M}^2 be the space of continuous square integrable (\mathcal{F}_t) -martingales on \mathbb{R}_+ .

Let \mathcal{P} denote the σ -field of (\mathcal{F}_\cdot) -predictable sets in $\mathbb{R}_+ \times \Omega$ and define

$$(46) \quad \mathcal{L}_{\text{loc}}^2 := \left\{ \psi : \mathbb{R}_+ \times \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R} : \psi \text{ is } \mathcal{P} \times \mathcal{B}(\mathbb{R}^2)\text{-measurable} \right. \\ \left. \text{and } \int_{[0,t] \times \mathbb{R}^2} L_{\mathbf{X}(\omega)}(d(s,x)) \psi^2(s,\omega,x) < \infty, \forall t > 0, \text{ a.s.} \right\}.$$

By starting with functions ψ of the form

$$(47) \quad \psi(s,\omega,x) = \sum_{m=1}^k \psi_{m-1}(\omega) \varphi_m(x) \mathbf{1}_{(t_{m-1}, t_m]}(s)$$

for some $\varphi_m \in C_b^2(\mathbb{R}^2)$, $\psi_{m-1} \in b\mathcal{F}_{t_{m-1}}$ (the space of bounded $\mathcal{F}_{t_{m-1}}$ -measurable maps) and $0 = t_0 < \dots < t_k \leq \infty$, and defining [with M^i from the martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$], for $i = 1, 2$,

$$(48) \quad M_t^i(\psi) = \int_{[0,t] \times \mathbb{R}^2} dM^i(s,x) \psi(s,x) \\ := \sum_{m=1}^k \psi_{m-1} (M_{t \wedge t_m}^i(\varphi_m) - M_{t \wedge t_{m-1}}^i(\varphi_m)),$$

we may uniquely extend M^i to linear maps $M^i : \mathcal{L}_{\text{loc}}^2 \rightarrow \mathbf{M}_{\text{loc}}$, such that

$$(49) \quad \langle \langle M^i(\psi_i), M^j(\psi_j) \rangle \rangle_t = \gamma \delta_{ij} \int_{[0,t] \times \mathbb{R}^2} L_{\mathbf{X}}(d(s,x)) \psi_i(s,x) \psi_j(s,x),$$

$t \geq 0$, a.s. for all $\psi_i \in \mathcal{L}_{\text{loc}}^2$. This may be done as in [37], Proposition II.5.4, or [45], Chapter 2. The M^i are orthogonal martingale measures. If, in addition,

$$(50) \quad \psi \in \mathcal{L}^2 := \left\{ \psi \in \mathcal{L}_{\text{loc}}^2 : E \int_{[0,t] \times \mathbb{R}^2} dL_{\mathbf{X}} \psi^2 < \infty, \forall t > 0 \right\},$$

then $M^i(\psi) \in \mathbf{M}^2$. The martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ shows that $M^i(\mathbf{1})$ belongs to \mathbf{M}^2 , hence the constant function $\mathbf{1}$ is in \mathcal{L}^2 and so

$$(51) \quad \text{every bounded and } \mathcal{P} \times \mathcal{B}(\mathbb{R}^2)\text{-measurable } \psi \text{ is in } \mathcal{L}^2 \text{ and } M^i(\psi) \in \mathbf{M}^2.$$

We need to extend $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ to time-dependent test functions.

NOTATION 22 (Time–space test functions). If $T > 0$, let \mathcal{D}_T denote the set of all bounded Borel maps $\psi : [0, T] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying the following:

(a) For any $x \in \mathbb{R}^2$, the map $t \mapsto \psi(t,x)$ is absolutely continuous and $\dot{\psi}(t,x) = \frac{\partial \psi}{\partial t}(t,x)$ is uniformly bounded in (t,x) and continuous in x for each $t \in [0, T]$.

(b) For each t in $[0, T]$, the mapping $x \mapsto \psi(t, x)$ belongs to $C_b^2(\mathbb{R}^2)$, and $\Delta\psi(t, \cdot)(x)$ is uniformly bounded in (t, x) .

LEMMA 23 [Extension of the martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$]. *If $\psi_i \in \mathcal{D}_T$, $i = 1, 2$, then*

$$(52) \quad \langle X_t^i, \psi_i(t) \rangle = \langle X_0^i, \psi_i(0) \rangle + \int_0^t ds \left\langle X_s^i, \dot{\psi}_i(s) + \frac{\sigma^2}{2} \Delta\psi_i(s) \right\rangle + M_t^i(\psi_i),$$

$t \in [0, T]$, where $M^i(\psi_i)$ belongs to \mathbf{M}^2 , and

$$(53) \quad \langle \langle M^i(\psi_i), M^j(\psi_j) \rangle \rangle_t = \delta_{ij} \gamma \int_{[0,t] \times \mathbb{R}^2} L_{\mathbf{X}}(d(s, x)) \psi_i(s, x) \psi_j(s, x).$$

PROOF. This may be done just as for ordinary superprocesses; see, for example, [37], Proposition II.5.7. The argument proceeds by approximating $\psi(s, x)$ by an appropriate sequence of step functions in t . \square

COROLLARY 24 (Green function representation). *Let $i = 1, 2$. If $\varphi_i : \mathbb{R}^2 \rightarrow \mathbb{R}$ is bounded and measurable, then, for any fixed $T > 0$,*

$$(54) \quad \langle X_t^i, S_{T-t}\varphi_i \rangle = \langle X_0^i, S_T\varphi_i \rangle + N_t^{i,T}(\varphi_i), \quad 0 \leq t \leq T, \text{ a.s.},$$

where

$$(55) \quad t \mapsto N_t^{i,T}(\varphi_i) = \int_{[0,t] \times \mathbb{R}^2} dM^i(r, x) S_{T-r}\varphi_i(x) \quad \text{belongs to } \mathbf{M}^2[0, T],$$

and

$$(56) \quad \langle \langle N^{i,T}(\varphi_i), N^{j,T}(\varphi_j) \rangle \rangle_t = \delta_{ij} \gamma \int_{[0,t] \times \mathbb{R}^2} L_{\mathbf{X}}(d(s, x)) S_{T-s}\varphi_i(x) S_{T-s}\varphi_j(x).$$

In particular,

$$(57) \quad \langle X_T^i, \varphi_i \rangle = \langle X_0^i, S_T\varphi_i \rangle + N_T^{i,T}(\varphi_i) \quad \text{a.s. } \forall T > 0.$$

PROOF. Let $\varphi \in C_b^2(\mathbb{R}^2)$ and $\psi(s, x) = S_{T-s}\varphi(x)$ for $(s, x) \in [0, T] \times \mathbb{R}^2$. Then $\psi \in \mathcal{D}_T$ because $\dot{\psi}(s, x) = (-\sigma^2/2) \Delta S_{T-s}\varphi(x) = (-\sigma^2/2) S_{T-s} \Delta\varphi(x)$. The result follows for such $\varphi \in C_b^2(\mathbb{R}^2)$ by Lemma 23. Now pass to the bounded pointwise closure to get the result for all bounded measurable ϕ . \square

2.2. *First and second moments: Proof of Proposition 15.* We proceed in several steps.

STEP 1 [Proof of (a)]. The equality in (a) is immediate upon taking expectations in Corollary 24 and using Definition 14 of (\mathbf{EnC}) for the finiteness of the mean.

STEP 2. Assume that $\psi = \varphi_1 \otimes \varphi_2$ with $\varphi_1, \varphi_2 \in b\mathcal{B}(\mathbb{R}^2)$. Corollary 24 shows that

$$(58) \quad \begin{aligned} E\langle X_t^i, \varphi_i \rangle \langle X_t^j, \varphi_j \rangle &= E\langle X_0^i, S_t \varphi_i \rangle \langle X_0^j, S_t \varphi_j \rangle \\ &\quad + \delta_{ij} \gamma E \int_{[0,t] \times \mathbb{R}^2} L_{\mathbf{X}}(d(s, x)) S_{t-s} \varphi_i(x) S_{t-s} \varphi_j(x), \end{aligned}$$

since by conditioning on \mathbf{X}_0 the cross terms vanish.

STEP 3 [Proof of (c)]. Before completing the proof of (b) we will consider (c). Assume $\psi(s, x) = \varphi(x)$ with $\varphi \in \mathcal{C}_{\text{com}}^+(\mathbb{R}^2)$. By Definition 1 and Fatou's lemma,

$$(59) \quad E\langle L_{\mathbf{X}}(T), \varphi \rangle \leq \liminf_{\delta \downarrow 0} E\langle L_{\mathbf{X}}^{*,\delta}(T), \varphi \rangle$$

$$(60) \quad = \liminf_{\delta \downarrow 0} E \frac{1}{\delta} \int_0^\delta dr \int_0^T ds \int_{\mathbb{R}^2} dx S_r X_s^1(x) S_r X_s^2(x) \varphi(x)$$

$$(61) \quad = \liminf_{\delta \downarrow 0} E \int_{\mathbb{R}^2} X_0^1(dy_1) \int_{\mathbb{R}^2} X_0^2(dy_2)$$

$$(62) \quad \times \frac{1}{\delta} \int_0^\delta dr \int_0^T ds \int_{\mathbb{R}^2} dx p_{r+s}(x, y_1) p_{r+s}(x, y_2) \varphi(x),$$

where we used (58) to continue after (60). The term in (62) is bounded by

$$(63) \quad \|\varphi\|_\infty \frac{1}{\delta} \int_0^\delta dr \int_0^T ds p_{2(r+s)}(y_1, y_2) \leq c \|\varphi\|_\infty g(y_1, y_2),$$

where in the last step we used (16). However, by **(EnC)** the bound in (63) is integrable with respect to $E X_0^1 \times X_0^2$. Hence, the limit inferior can be taken through the three integrals in (61). It is then easy to let $\delta \rightarrow 0$ in the resulting integrand as we only need to consider $y_1 \neq y_2$ by **(EnC)**. This gives

$$(64) \quad E\langle L_{\mathbf{X}}(T), \varphi \rangle \leq E \int_0^T ds \int_{\mathbb{R}^2} dx S_s X_0^1(x) S_s X_0^2(x) \varphi(x)$$

$$(65) \quad \leq c_T \|\varphi\|_\infty E \|\mathbf{X}_0\|_g < \infty.$$

By an obvious monotone class argument, claim (c) follows for bounded measurable ψ on $[0, t] \times \mathbb{R}^2$.

STEP 4 [Proof of (b)]. We may apply (c) to (58) to get the claim (b) for the special functions ψ used in Step 2. A monotone class argument then gives the desired extension.

STEP 5 [Proof of (d)]. Assume **(IntC)**. First consider again the case $\psi(s, x) = \varphi(x)$ with a function $\varphi \in \mathcal{C}_{\text{com}}^+(\mathbb{R}^2)$. Fix $0 < \varepsilon < T$. Suppose we can show

$$(66) \quad E\{\langle L_{\mathbf{X}}(T) - L_{\mathbf{X}}(\varepsilon), \varphi \rangle \mid \mathcal{F}_\varepsilon\} = \int_0^{T-\varepsilon} ds \int_{\mathbb{R}^2} dx \varphi(x) S_s X_\varepsilon^1(x) S_s X_\varepsilon^2(x).$$

Then, by (58),

$$(67) \quad E \langle L_{\mathbf{X}}(T) - L_{\mathbf{X}}(\varepsilon), \varphi \rangle = \int_{\varepsilon}^T ds \int_{\mathbb{R}^2} dx \varphi(x) E S_s X_0^1(x) S_s X_0^2(x).$$

Now let $\varepsilon \downarrow 0$. By (c), the left-hand side of (67) converges to $E \langle L_{\mathbf{X}}(T), \varphi \rangle$, whereas by monotone convergence on the right-hand side we obtain the required expression. Provided we have (66), this proves equality in (c) under **(IntC)** for the considered special ψ , hence for all required ψ by dominated convergence and (c).

By (58), we then also get the equality in (b) under **(IntC)** for functions ψ of the form $\varphi_1 \otimes \varphi_2$ with $\varphi_1, \varphi_2 \in b\mathcal{B}(\mathbb{R}^2)$, thus for all required ψ .

STEP 6. To finish the proof, it remains to show (66). First, (57) and (54) in Corollary 24 give

$$(68) \quad \langle X_s^i, \varphi \rangle - \langle X_\varepsilon^i, S_{s-\varepsilon} \varphi \rangle = N_s^{i,s}(\varphi) - N_\varepsilon^{i,s}(\varphi) \quad \text{a.s., } s \geq \varepsilon, i = 1, 2.$$

Therefore,

$$(69) \quad E \{ \langle X_s^1, \varphi \rangle \langle X_s^2, \varphi \rangle \mid \mathcal{F}_\varepsilon \} = \langle X_\varepsilon^1, S_{s-\varepsilon} \varphi \rangle \langle X_\varepsilon^2, S_{s-\varepsilon} \varphi \rangle \quad \text{a.s., } s \geq \varepsilon.$$

On the other hand, for $\delta > 0$, by the definition (Definition 1) of $L_{\mathbf{X}}^{*,\delta}$,

$$(70) \quad \langle L_{\mathbf{X}}^{*,\delta}(T) - L_{\mathbf{X}}^{*,\delta}(\varepsilon), \varphi \rangle = \frac{1}{\delta} \int_0^\delta dr \int_\varepsilon^T ds \int_{\mathbb{R}^2} dx \varphi(x) S_r X_s^1(x) S_r X_s^2(x).$$

Thus, by (69),

$$(71) \quad \begin{aligned} & E \{ \langle L_{\mathbf{X}}^{*,\delta}(T) - L_{\mathbf{X}}^{*,\delta}(\varepsilon), \varphi \rangle \mid \mathcal{F}_\varepsilon \} \\ &= \frac{1}{\delta} \int_0^\delta dr \left[\int_\varepsilon^T ds \int_{\mathbb{R}^2} dx \varphi(x) S_{r+s-\varepsilon} X_\varepsilon^1(x) S_{r+s-\varepsilon} X_\varepsilon^2(x) \right] \\ &= \frac{1}{\delta} \int_0^\delta dr \left[\int_r^{r+T-\varepsilon} ds \int_{\mathbb{R}^2} dx \varphi(x) S_s X_\varepsilon^1(x) S_s X_\varepsilon^2(x) \right]. \end{aligned}$$

Since $r \in [0, \varepsilon]$, the term in square brackets in the last line of (71) can be bounded above by

$$(72) \quad \int_0^T ds \int_{\mathbb{R}^2} dx \varphi(x) S_s X_\varepsilon^1(x) S_s X_\varepsilon^2(x).$$

However, by (58), the expectation of this can be computed and equals

$$(73) \quad \int_\varepsilon^{T+\varepsilon} ds \int_{\mathbb{R}^2} dx \varphi(x) S_s X_0^1(x) S_s X_0^2(x),$$

which is finite by (37). Hence, (72) is finite a.s. Therefore we may let $\delta \downarrow 0$ in (71) and conclude that, for any sequence $\delta_n \downarrow 0$,

$$(74) \quad \begin{aligned} & \lim_{n \uparrow \infty} E \{ \langle L_{\mathbf{X}}^{*,\delta_n}(T) - L_{\mathbf{X}}^{*,\delta_n}(\varepsilon), \varphi \rangle \mid \mathcal{F}_\varepsilon \} \\ &= \int_0^{T-\varepsilon} ds \int_{\mathbb{R}^2} dx \varphi(x) S_s X_\varepsilon^1(x) S_s X_\varepsilon^2(x) \quad \text{a.s.} \end{aligned}$$

Thus, to prove (66) it suffices to show that in probability

$$(75) \quad E\{\{L_{\mathbf{X}}^{*,\delta_n}(T) - L_{\mathbf{X}}^{*,\delta_n}(\varepsilon), \varphi\} | \mathcal{F}_\varepsilon\} \xrightarrow{n \uparrow \infty} E\{\{L_{\mathbf{X}}(T) - L_{\mathbf{X}}(\varepsilon), \varphi\} | \mathcal{F}_\varepsilon\}.$$

Note that by the definition (Definition 1) of the collision local time, there is convergence in probability of the corresponding expressions inside the conditional expectations. On the other hand, by (70) and Jensen’s inequality, we have

$$\begin{aligned} & (L_{\mathbf{X}}^{*,\delta_n}(T) - L_{\mathbf{X}}^{*,\delta_n}(\varepsilon), \varphi)^2 \\ & \leq \|\varphi\|_\infty^2 \frac{T}{\delta_n} \int_0^{\delta_n} dr \int_\varepsilon^T ds \int_{\mathbb{R}^2} dx \int_{\mathbb{R}^2} dy S_r X_s^1(x) S_r X_s^2(x) S_r X_s^1(y) S_r X_s^2(y) \\ & \leq \|\varphi\|_\infty^2 \frac{T}{\delta_n} \int_0^{\delta_n} dr \int_\varepsilon^T ds H_r(\mathbf{X}_s) \end{aligned}$$

[recall notation (26)]. Therefore,

$$(76) \quad \begin{aligned} & E\{\{L_{\mathbf{X}}^{*,\delta_n}(T) - L_{\mathbf{X}}^{*,\delta_n}(\varepsilon), \varphi\}^2 | \mathcal{F}_\varepsilon\} \\ & \leq \|\varphi\|_\infty^2 \frac{T}{\delta_n} \int_0^{\delta_n} dr \int_\varepsilon^T ds E\{H_r(\mathbf{X}_s) | \mathcal{F}_\varepsilon\}, \end{aligned}$$

which is bounded in probability as $\delta_n \downarrow 0$ by our assumption **(IntC)** (recall Definition 7). A standard uniform integrability argument for conditional expectations (Lemma 63 in Appendix B) now gives (75) and completes the proof of (d). \square

2.3. *State spaces for \mathbf{X} .* Recall the state space versions $\mathcal{M}_{f,s}$ and $\mathcal{M}_{f,se}$ from Definition 4.

PROPOSITION 25 (State spaces). *Assume \mathbf{X}_0 is a random initial state in $\mathcal{M}_{f,e}$ satisfying the random energy condition **(EnC)** from Definition 14, and \mathbf{X} satisfies **(MP)** $_{\mathbf{X}_0}^{\sigma,\gamma}$. Then the following hold:*

- (a) $X_t \in \mathcal{M}_{f,se}$ a.s. for each $t > 0$.
- (b) $X_t \in \mathcal{M}_{f,e}$ for all $t \geq 0$ a.s.

PROOF. (a) Fix $t > 0$. By Remark 5, for the verification of (15) it suffices to consider $0 < r < 1$. By Proposition 15(b),

$$\begin{aligned} E\langle X_t^i \times X_t^j, p_r \rangle & \leq E \int_{\mathbb{R}^2} dx_1 S_t X_0^i(x_1) \int_{\mathbb{R}^2} dx_2 S_t X_0^j(x_2) p_r(x_1, x_2) \\ & \quad + \delta_{ij} \gamma E \int_0^t ds \int_{\mathbb{R}^2} dx S_s X_0^1(x) S_s X_0^2(x) \\ & \quad \times \int_{\mathbb{R}^2} dx_1 p_{t-s}(x, x_1) \int_{\mathbb{R}^2} dx_2 p_{t-s}(x, x_2) p_r(x_1, x_2). \end{aligned}$$

The right-hand side of this inequality can be written as

$$(77) \quad E\langle X_0^i \times X_0^j, p_{2t+r} \rangle + \delta_{ij} \gamma E \int_0^t ds p_{r+2(t-s)}(0, 0) E\langle X_0^1 \times X_0^2, p_{2s} \rangle.$$

For the first term in (77), use $p_{2t+r}(y_1, y_2) \leq p_{2t+r}(0, 0) \leq c(t)$ to get the bound $c(t)E\langle X_0^i, 1 \rangle \langle X_0^j, 1 \rangle$. In the second term of (77), break the integral at $t/2$. For the lower part, apply $p_{r+2(t-s)}(0, 0) \leq c(t)$, whereas for the second part, use $p_{2s}(y_1, y_2) \leq c(t)$. This gives the bound

$$(78) \quad c(t) \int_0^{t/2} ds E\langle X_0^1 \times X_0^2, p_{2s} \rangle$$

$$(79) \quad + c(t) \int_{t/2}^t ds p_{r+2(t-s)}(0, 0) E\langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle$$

for the second term in (77). For (78) use (16) to bound it by $c(t) \|\mathbf{X}_0\|_g$, whereas in (79) the ds -integral can be bounded by $c(t)[1 + \log(1/r)]$. Altogether,

$$(80) \quad E\langle X_t^i \times X_t^j, p_r \rangle \leq c(t)[1 + \log(1/r)] E \left[\sum_{i=1,2} \langle X_0^i, 1 \rangle^2 + \|\mathbf{X}_0\|_g \right] \\ = c[1 + \log(1/r)],$$

where in the last step we used our assumption **(EnC)**, and the constant c is independent of r .

Next we want to apply this estimate for special values of r . In fact, if r belongs to $[2^{-n-1}, 2^{-n})$, $n \geq 0$, then $p_r \leq 2p_{2^{-n}}$, and if $p \in (0, 1)$, then from (80),

$$E \sup_{0 < r < 1} r^p \langle X_t^i \times X_t^j, p_r \rangle \leq 2c \sum_{n=0}^{\infty} 2^{-np} [1 + \log 2^n] < \infty.$$

This proves $\mathbf{X}_t \in M_{f,se}$ a.s.

(b) We will use a *Tanaka formula* approach from [2]. To prepare for this, for $\alpha, \varepsilon \geq 0$, set

$$g_{\alpha,\varepsilon}(x_1, x_2) := \frac{1}{2} e^{\varepsilon \alpha/2} \int_{\varepsilon}^{\infty} du e^{-\alpha u/2} p_u(x_1, x_2), \quad x_1, x_2 \in \mathbb{R}^2.$$

Note that

$$(81) \quad g_{\alpha,\varepsilon} \leq e^{\alpha} g_{\alpha,0}, \quad 0 \leq \varepsilon \leq 1, \alpha \geq 0,$$

and we have pointwise convergence

$$(82) \quad \lim_{\varepsilon \downarrow 0} g_{\alpha,\varepsilon} = g_{\alpha,0}, \quad \alpha \geq 0.$$

It is easy to see ([2], (5.6)) that to each $\alpha > 0$ there are positive constants c_α and C_α such that

$$(83) \quad c_\alpha g \leq 1 + g_{\alpha,0} \leq C_\alpha g$$

[with the energy function g from (13)].

Let $\mathbf{X}_t = X_t^1 \times X_t^2$. It follows from $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ and a bit of stochastic calculus, just as in the derivation of (T_ε) in Section 5 of [2], that

$$\begin{aligned}
 \langle \mathbf{X}_t, g_{\alpha, \varepsilon} \rangle &= \langle \mathbf{X}_0, g_{\alpha, \varepsilon} \rangle \\
 &+ \int_0^t \int \int g_{\alpha, \varepsilon}(x_1, x_2) [X_s^1(dx_1)M^2(ds, dx_2) \\
 &+ X_s^2(dx_2)M^1(ds, dx_1)] \\
 &+ \alpha \int_0^t \int \int g_{\alpha, \varepsilon}(x_1, x_2) X_s^1(dx_1)X_s^2(dx_2) ds - \hat{L}_t^\varepsilon(\mathbf{X}),
 \end{aligned}
 \tag{84}$$

where $\hat{L}_t^\varepsilon(\mathbf{X}) = \int_0^t \int \int p_\varepsilon(x_1 - x_2) X_s^1(dx_1)X_s^2(dx_2) ds$. As $g_{\alpha, \varepsilon}$ is bounded the above stochastic integral in (84), $I^\varepsilon(t)$, is a continuous local martingale and we may choose stopping times $T_n \uparrow \infty$ a.s. such that $\sup_{t \leq T_n} I^\varepsilon(t) \leq n$. Then (84) implies

$$\begin{aligned}
 &E(\langle \mathbf{X}_{t \wedge T_n}, g_{\alpha, \varepsilon} \rangle) \\
 &\leq E(\langle \mathbf{X}_0, g_{\alpha, \varepsilon} \rangle) + \alpha \int_0^t E(\langle \mathbf{X}_{s \wedge T_n}, g_{\alpha, \varepsilon} \rangle) ds \\
 &\leq C_\alpha E(\langle \mathbf{X}_0, g \rangle) + \alpha \int_0^t E(\langle \mathbf{X}_{s \wedge T_n}, g_{\alpha, \varepsilon} \rangle) ds \quad [\text{by (81) and (83)}].
 \end{aligned}
 \tag{85}$$

Note also that $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ implies that $\langle \mathbf{X}_t, 1 \rangle = \langle X_t^1, 1 \rangle \langle X_t^2, 1 \rangle$ is a martingale (we also use **(EnC)** here) and so

$$E(\langle \mathbf{X}_{t \wedge T_n}, g_{\alpha, \varepsilon} \rangle) \leq \|g_{\alpha, \varepsilon}\|_\infty E(\langle \mathbf{X}_{t \wedge T_n}, 1 \rangle) = \|g_{\alpha, \varepsilon}\|_\infty E(\langle \mathbf{X}_0, 1 \rangle) < \infty.$$

It therefore follows from (85) that

$$E(\langle \mathbf{X}_{t \wedge T_n}, g_{\alpha, \varepsilon} \rangle) \leq c(\alpha) E(\langle \mathbf{X}_0, g \rangle) e^{\alpha t} \quad \forall t \geq 0, n \in \mathbb{N}.
 \tag{86}$$

Note also, by Proposition 15(b),

$$\begin{aligned}
 E(\hat{L}_t^\varepsilon(\mathbf{X})) &= E\left(\int_0^t \int \int p_\varepsilon(y_1 - y_2) S_s X_0^1(y_1) S_s X_0^2(y_2) dy_1 dy_2 ds\right) \\
 &= E\left(\int_0^t \int \int p_{\varepsilon+2s}(y_1 - y_2) X_0^1(dy_1) X_0^2(dy_2) ds\right) \\
 &\leq c'(t) E(\langle \mathbf{X}_0, g \rangle).
 \end{aligned}
 \tag{87}$$

It follows from (84) and the integrability implied by (86) and (87) that

$$Y_t^n \equiv \langle \mathbf{X}_{t \wedge T_n}, g_{\alpha, \varepsilon} \rangle + \hat{L}_{t \wedge T_n}^\varepsilon(\mathbf{X})$$

is a nonnegative submartingale. Therefore by the weak maximal inequality for

any $t, K > 0$ fixed

$$\begin{aligned} P\left(\sup_{s \leq t \wedge T_n} \langle \mathbf{X}_s, g_{\alpha, \varepsilon} \rangle > K\right) &\leq P\left(\sup_{s \leq t} Y_s^n > K\right) \\ &\leq K^{-1} E(Y_t^n) \\ &\leq K^{-1} [c(\alpha)e^{\alpha t} + c'(t)] E(\langle \mathbf{X}_0, g \rangle). \end{aligned}$$

First let $n \rightarrow \infty$ and then $\varepsilon \rightarrow 0$ in the above and use Fatou’s lemma and (82) to see that

$$P\left(\sup_{s \leq t} \langle \mathbf{X}_s, g_{\alpha, 0} \rangle > K\right) \leq K^{-1} [c(\alpha)e^{\alpha t} + c'(t)] E(\langle \mathbf{X}_0, g \rangle).$$

In view of the lower bound in (83), the required result is immediate. \square

3. Dual processes for higher moments. In this section function-valued and measure-valued duals are presented which are used to compute higher moments.

3.1. *Lattice approximation moment dual \mathbf{V}^ε and self-duality.* Since it has not been explicitly mentioned in [14], we start by pointing out that our lattice approximations have finite moments of all orders:

LEMMA 26 (Moments of all orders). *Let $\varepsilon > 0$. Assume ${}^\varepsilon\mathbf{X}$ satisfies the martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon}$ of Lemma 6 with deterministic initial condition, $\mathbf{X}_0 \in \mathcal{M}_f^2(\mathbb{R}^2)$. Then for any integer $m \geq 1$ and $T > 0$ there is a constant $C = C(\varepsilon, T, m, \langle \mathbf{X}_0, 1 \rangle)$ such that*

$$(88) \quad \sum_{i=1,2} E\left(\sup_{t \leq T} \langle {}^\varepsilon X_t^i, 1 \rangle^m\right) \leq C.$$

PROOF. Clearly we may assume $m \geq 2$ and $\varepsilon = 1$, and we will suppress the index $\varepsilon = 1$ in our notation. Then, for $i \in \{1, 2\}$ fixed, $t \mapsto \langle X_t^i, 1 \rangle - \langle X_0^i, 1 \rangle = M_t^i(1)$ is a continuous L^2 -martingale such that, for $T > 0$ fixed and $t \leq T$,

$$\begin{aligned} \langle \langle M^i(1) \rangle \rangle_t^{m/2} &= \left(\gamma \int_0^t ds \sum_{x \in \mathbb{Z}^2} X_s^1(x) X_s^2(x) \right)^{m/2} \\ &\leq c \int_0^t ds \left(\sum_{x \in \mathbb{Z}^2} X_s^1(x) X_s^2(x) \right)^{m/2} \leq c \int_0^t ds \sum_{i=1,2} \langle X_s^i, 1 \rangle^m \end{aligned}$$

(where $c = c_{m, \gamma, T}$). For the moment fix $K \geq 1$, and consider the stopping time $\tau_K := T \wedge \inf\{t : \sum_{i=1}^2 \langle X_t^i, 1 \rangle \geq K\}$. Burkholder’s inequality then shows that, for

any $r \in [0, T]$,

$$(89) \quad \begin{aligned} & E\left(\sum_{i=1,2} \sup_{t \leq r \wedge \tau_K} \langle X_t^i, 1 \rangle^m\right) \\ & \leq c \sum_{i=1,2} \langle X_0^i, 1 \rangle^m + c \int_0^r ds E\left(\sum_{i=1,2} \langle X_{s \wedge \tau_K}^i, 1 \rangle^m\right), \end{aligned}$$

with the constant c independent of r (and K). Since the expectation in the integrand on the right-hand side of this inequality can further be bounded from above by $E(\sum_{i=1,2} \sup_{t \leq s \wedge \tau_K} \langle X_t^i, 1 \rangle^m)$, Gronwall's lemma implies

$$(90) \quad E\left(\sum_{i=1,2} \sup_{t \leq \tau_K} \langle X_t^i, 1 \rangle^m\right) \leq C,$$

where $C = C(T, m, \langle \mathbf{X}_0, 1 \rangle)$ is independent of K . Letting $K \uparrow \infty$ completes the proof since $\tau_K \uparrow T$. \square

Although in this paper we only use fourth order moments, we now introduce a *function-valued dual process* $\mathbf{V}^\varepsilon = \mathbf{V}^{\varepsilon, m}$ which will describe moments of arbitrary but fixed order $m \geq 1$ for solutions ${}^\varepsilon \mathbf{X}$ of $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon}$, with a fixed $\varepsilon \in (0, 1]$. The state space of the dual is $\mathcal{S}^\varepsilon = \mathcal{S}^{\varepsilon, m} := C_b^+(\varepsilon \mathbb{Z}^2)^m \times 2^{\{1, \dots, m\}}$ (with $2^{\{1, \dots, m\}}$ denoting the power set of $\{1, \dots, m\}$), and elements in \mathcal{S}^ε are denoted by (ϕ, I) . It is convenient to think of the argument of ϕ as the spatial positions of a system of m particles. Particles take two types: those corresponding to a coordinate whose index is in I are of type 1; those corresponding to indices in I^c are of type 2. These m particles have positions described by $x \in (\varepsilon \mathbb{Z}^2)^m$. We give $C_b^+(\varepsilon \mathbb{Z}^2)^m$ the topology of pointwise convergence, to make \mathcal{S}^ε a separable metric space.

Let ${}^\varepsilon S_t^{(m)}$ denote the semigroup on $C_b(\varepsilon \mathbb{Z}^2)^m$ obtained by running m independent copies of our simple random walk ${}^\varepsilon \xi$ (each with generator $\frac{\sigma^2}{2} \varepsilon \Delta$), and let $\frac{\sigma^2}{2} \varepsilon \Delta^{(m)}$ denote the associated generator.

For $1 \leq j, j' \leq m$ with $j \neq j'$, define maps $\pi_{j, j'} : (\mathbb{R}^2)^m \rightarrow (\mathbb{R}^2)^m$ and $f_{j, j'} : C_b^+(\varepsilon \mathbb{Z}^2)^m \rightarrow C_b^+(\varepsilon \mathbb{Z}^2)^m$ by

$$(91) \quad (\pi_{j, j'} x)_i := \begin{cases} x_i, & \text{if } i \neq j', \\ x_j, & \text{if } i = j', \end{cases} \quad x = (x_1, \dots, x_m) \in (\mathbb{R}^2)^m,$$

and

$$(92) \quad \begin{aligned} f_{j, j'}(\phi)(x) & := \phi(\pi_{j, j'} x) \varepsilon^{-2} \mathbf{1}(x_j = x_{j'}) \\ & = \phi(x)^\varepsilon p_0(x_j, x_{j'}). \end{aligned}$$

DEFINITION 27 (Dual process \mathbf{V}^ε). For fixed $m \geq 1$, denote by $\mathbf{V}^\varepsilon = \mathbf{V}^{\varepsilon,m} = \{\mathbf{V}_t^\varepsilon : t \geq 0\}$ the Markov process which has sample paths in the Skorohod space $D(\mathbb{R}_+, \mathcal{S}^\varepsilon)$, and evolves as follows:

(a) *Jumps*—If \mathbf{V}^ε is in the state (ϕ, I) , for each (ordered) pair (j, j') in I^2 satisfying $j \neq j'$, the process \mathbf{V}^ε jumps to $(f_{j,j'}(\phi), I \setminus \{j'\})$ with rate $\gamma/2$, and for each $(j, j') \in (I^c)^2$ with $j \neq j'$, it jumps to $(f_{j,j'}(\phi), I \cup \{j'\})$, also with rate $\gamma/2$. (In particular, a jumping particle changes its type.) In these cases we say j' switches via j .

Let $\{T_j : j \geq 1\}$ denote the successive jump times, and set $T_0 = 0$.

(b) *Between jumps*—Between jump times, the component ϕ of \mathbf{V}^ε evolves according to the semigroup ${}^\varepsilon S^{(m)}$, whereas the component I is frozen. That is

$$(93) \quad \text{if } T_n \leq t < T_{n+1}, \text{ then } \phi_t(x) = {}^\varepsilon S_{t-T_n}^{(m)} \phi_{T_n}(x), \text{ and } I_t = I_{T_n}.$$

Let $\mathcal{A}^\varepsilon = \mathcal{A}^{\varepsilon,m}$ denote the (weak) infinitesimal generator of \mathbf{V}^ε , and let $\hat{P}_{\mathbf{V}_0^\varepsilon}^\varepsilon$ denote the law of \mathbf{V}^ε if \mathbf{V}^ε starts in \mathbf{V}_0^ε (deterministic).

Define a duality function $F : \mathcal{S}^\varepsilon \times \mathcal{M}_f^2(\varepsilon\mathbb{Z}^2) \rightarrow \mathbb{R}_+$ by

$$(94) \quad F(\phi, I, \mu^1, \mu^2) := \prod_{i \in I} \int_{\varepsilon\mathbb{Z}^2} \mu^1(dx_i) \prod_{j \in I^c} \int_{\varepsilon\mathbb{Z}^2} \mu^2(dx_j) \phi(x).$$

Then, for $(\phi, I, \mu) \in \mathcal{S}^\varepsilon \times \mathcal{M}_f^2(\varepsilon\mathbb{Z}^2)$,

$$(95) \quad \begin{aligned} \mathcal{A}^\varepsilon F(\cdot, \cdot, \mu)(\phi, I) &= F\left(\frac{\sigma^2}{2} {}^\varepsilon \Delta^{(m)} \phi, I, \mu\right) \\ &+ \frac{\gamma}{2} \sum_{\substack{(j,j') \in I^2 \\ j \neq j'}} (F(f_{j,j'}^\varepsilon(\phi), I \setminus \{j'\}, \mu) - F(\phi, I, \mu)) \\ &+ \frac{\gamma}{2} \sum_{\substack{(j,j') \in (I^c)^2 \\ j \neq j'}} (F(f_{j,j'}^\varepsilon(\phi), I \cup \{j'\}, \mu) - F(\phi, I, \mu)). \end{aligned}$$

Hence, for $\mu \in \mathcal{M}_f^2(\varepsilon\mathbb{Z}^2)$,

$$(96) \quad F(\mathbf{V}_t^\varepsilon, \mu) - F(\mathbf{V}_0^\varepsilon, \mu) - \int_0^t ds \mathcal{A}^\varepsilon F(\mathbf{V}_s^\varepsilon, \mu)$$

is a $\hat{P}_{\mathbf{V}_0^\varepsilon}^\varepsilon$ -martingale. [See (98) below for the integrability of $F(\mathbf{V}_t^\varepsilon, \mu)$ with respect to $\hat{P}_{\mathbf{V}_0^\varepsilon}^\varepsilon$.]

Let ${}^\varepsilon \mathbf{X}$ be our solution to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon}$ from Lemma 6 and denote the underlying probability by $P_{\mathbf{X}_0}^\varepsilon$. As usual \mathbf{X}_0 is a fixed element in $\mathcal{M}_{f,e}$. If $(\phi, I) \in \mathcal{S}^\varepsilon$, then

Itô's lemma and the system of stochastic equations (19) defining the process ${}^\varepsilon\mathbf{X}$ show that

$$\begin{aligned}
 F(\phi, I, {}^\varepsilon\mathbf{X}_t) &= F(\phi, I, {}^\varepsilon\mathbf{X}_0) \\
 &+ \int_0^t ds \left[\mathcal{A}^\varepsilon F(\phi, I, {}^\varepsilon\mathbf{X}_s) + \gamma \left\{ \binom{|I|}{2} + \binom{|I^c|}{2} \right\} F(\phi, I, {}^\varepsilon\mathbf{X}_s) \right] \\
 &+ M_t^{\phi, I},
 \end{aligned}$$

where $M^{\phi, I}$ is a continuous L^2 -martingale which can be explicitly written in terms of the Brownian motions arising in (19). [Note that the integrals in the duality function (94) are actually sums.]

On the other hand, if A^ε is the weak generator of ${}^\varepsilon\mathbf{X}$, then we have

$$(97) \quad A^\varepsilon F(\phi, I, \cdot)(\mu) = \mathcal{A}^\varepsilon F(\cdot, \cdot, \mu)(\phi, I) + \gamma \left\{ \binom{|I|}{2} + \binom{|I^c|}{2} \right\} F(\phi, I, \mu),$$

$$(\phi, I, \mu) \in \mathcal{S}^\varepsilon \times \mathcal{M}_f^2(\mathbb{R}^2).$$

PROPOSITION 28 (Moment duality for \mathbf{X}^ε). *For any $\mathbf{V}_0^\varepsilon \in \mathcal{S}^\varepsilon$, $\mathbf{X}_0 \in \mathcal{M}_{f,e}(\mathbb{R}^2)$, $\varepsilon \in (0, 1]$ and $t > 0$,*

$$E_{\mathbf{X}_0}^\varepsilon F(\mathbf{V}_0^\varepsilon, {}^\varepsilon\mathbf{X}_t) = \hat{E}_{\mathbf{V}_0^\varepsilon}^\varepsilon \left(F(\mathbf{V}_t^\varepsilon, {}^\varepsilon\mathbf{X}_0) \exp \left[\gamma \int_0^t ds \left\{ \binom{|I_s|}{2} + \binom{|I_s^c|}{2} \right\} \right] \right) < \infty.$$

PROOF. In view of (97) we only need to check the hypotheses (4.50) and (4.51) of [22], Theorem 4.11, with $\alpha = 0$ and $\beta(\phi, I) = \binom{|I|}{2} + \binom{|I^c|}{2}$. Note that $\beta(\phi, I) \leq 2 \binom{m}{2}$, so that (4.51) is obvious. Let N_s be the number of jumps of \mathbf{V}^ε up to time s . Note that

$$\begin{aligned}
 (98) \quad &\hat{E}_{\mathbf{V}_0^\varepsilon}^\varepsilon \times E_{\mathbf{X}_0}^\varepsilon \left(\sup_{0 \leq s, t \leq T} F(\phi_s, I_s, {}^\varepsilon\mathbf{X}_t) \right) \\
 &\leq c \hat{E}_{\mathbf{V}_0^\varepsilon}^\varepsilon (\varepsilon^{-2N_T} \|\phi_0\|_\infty) E_{\mathbf{X}_0}^\varepsilon \left(\sup_{t \leq T} \langle {}^\varepsilon X_t^1, 1 \rangle^m + \sup_{t \leq T} \langle {}^\varepsilon X_t^2, 1 \rangle^m \right) < \infty,
 \end{aligned}$$

by Lemma 26. Then (4.50) in Theorem 4.11 of [22] is a simple consequence of this. \square

It is not hard to see that the above moments grow too quickly for the moment problem to be well posed and hence do not characterize the law of ${}^\varepsilon\mathbf{X}$. An exponential self-duality [35] is required for this.

3.2. *Limiting moment dual V.* To let $\varepsilon \downarrow 0$ in Proposition 28 we specialize to $m = 4$ and introduce the natural candidate for a limiting dual process V . To define the state space we introduce some notation.

NOTATION 29. If φ is a (real-valued) function on \mathbb{R}^d , put

$$|\varphi|_\lambda := \sup_{x \in \mathbb{R}^d} |\varphi(x)|/\phi_\lambda(x), \quad \lambda \in \mathbb{R},$$

where

$$\phi_\lambda(x) := e^{-\lambda|x|}, \quad x \in \mathbb{R}^d.$$

For $\lambda \in \mathbb{R}$, let \mathcal{C}_λ denote the set of all continuous functions such that $|\varphi|_\lambda$ is finite. Introduce the space

$$\mathcal{C}_{\text{exp}} = \mathcal{C}_{\text{exp}}(\mathbb{R}^d) := \bigcup_{\lambda > 0} \mathcal{C}_\lambda$$

of exponentially decreasing continuous functions. Let $\mathcal{M}_{\text{tem}} = \mathcal{M}_{\text{tem}}(\mathbb{R}^d)$ denote the subset of all measures μ on \mathbb{R}^d such that $\langle \mu, \phi_\lambda \rangle < \infty$ for all $\lambda > 0$. We topologize the set of tempered measures \mathcal{M}_{tem} by the metric

$$d_{\text{tem}}(\mu, \nu) := d_0(\mu, \nu) + \sum_{n=1}^{\infty} 2^{-n} (|\mu - \nu|_{1/n} \wedge 1), \quad \mu, \nu \in \mathcal{M}_{\text{tem}}.$$

Here d_0 is a complete metric on the space of Radon measures on \mathbb{R}^d inducing the vague topology, and $|\mu - \nu|_\lambda$ is an abbreviation for $|\langle \mu, \phi_\lambda \rangle - \langle \nu, \phi_\lambda \rangle|$. Note that $(\mathcal{M}_{\text{tem}}, d_{\text{tem}})$ is a Polish space and that $\mu_n \rightarrow \mu$ in \mathcal{M}_{tem} if and only if $\langle \mu_n, \varphi \rangle \rightarrow \langle \mu, \varphi \rangle$ for all $\varphi \in \mathcal{C}_{\text{exp}}$.

The state space for this dual V will be $\mathcal{S} = \mathcal{M}_{\text{tem}}((\mathbb{R}^2)^4) \times 2^{\{1, \dots, 4\}}$, although our starting point V_0 will be in

$$(99) \quad C_b^+((\mathbb{R}^2)^4) \times 2^{\{1, \dots, 4\}} =: \mathcal{S}_0.$$

As before, we will identify functions ϕ_0 in C_b^+ with the measure $\phi_0(x) dx$ in \mathcal{M}_{tem} . We abuse our earlier notation slightly and define $F : \mathcal{S} \times \mathcal{M}_f^2(\mathbb{R}^2) \mapsto \mathbb{R}_+$ by

$$F(\phi, I, \mu) = \begin{cases} \int \phi(x_1, \dots, x_4) \prod_{i \in I} \mu^1(dx_i) \prod_{j \notin I} \mu^2(dx_j), & \text{if } (\phi, I) \in \mathcal{S}_0, \\ 0, & \text{otherwise,} \end{cases}$$

and define $\pi_{j, j'} : (\mathbb{R}^2)^4 \mapsto (\mathbb{R}^2)^4$ for $1 \leq j, j' \leq 4$ as before. If $1 \leq j, j' \leq 4$, then $f_{j, j'} : C_b^+((\mathbb{R}^2)^4) \mapsto \mathcal{M}_{\text{tem}}((\mathbb{R}^2)^4)$ is given by

$$(100) \quad f_{j, j'}(\phi)(A) = \int_A dx_1 \cdots dx_4 \phi(\pi_{j, j'}x) \delta_0(x_j - x_{j'}).$$

It is easy to check this measure is in \mathcal{M}_{tem} .

DEFINITION 30 (Dual process \mathbf{V}). Let \mathbf{S}_t be the eight-dimensional Brownian semigroup with variance parameter σ^2 , let $\frac{\sigma^2 \Delta}{2}$ denote its generator and $\mathbf{p}_t(x, y)$ the associated transition function. The dynamics of the dual process $\mathbf{V} = (\phi, I) \in D(\mathbb{R}_+, \mathcal{I})$ are as follows:

(a) For each $(j, j') \in I_t^2$, $j \neq j'$, with rate $\gamma/2$, (ϕ_{t-}, I_{t-}) jumps to $(f_{j,j'}(\phi_{t-}), I_{t-} \setminus \{j'\})$, and for each $(j, j') \in (I_t^c)^2$, $j \neq j'$, with rate $\gamma/2$, (ϕ_{t-}, I_{t-}) jumps to $(f_{j,j'}(\phi_{t-}), I_{t-} \cup \{j'\})$.

Let $0 = T_0 < T_1 < T_2 < \dots$ be the successive jump times.

(b) For $T_n \leq t < T_{n+1}$, $\mathbf{V}_t = (\mathbf{S}_{t-T_n} \phi_{T_n}, I_{T_n})$.

REMARK 31. To ensure that this does define a process V_t we need to check that $\phi_{T_n-} \in C_b^+(\mathbb{R}^2)^4$ for all $n \geq 1$ a.s. so that $f_{j,j'}(\phi_{T_n-})$ is well defined. For this we will use induction to show if $T_n < T_{n+1}$ for all $n \geq 0$. Then

$$(101) \quad \begin{aligned} &\text{on } [T_n, T_{n+1}), \phi_t \text{ is a continuous } \mathcal{M}_{\text{tem}}\text{-valued process taking values in} \\ &C_b^+(\mathbb{R}^8) \text{ for } t \in (T_n, T_{n+1}), \text{ and } \phi_{T_{n+1}-} = \mathbf{S}_{T_{n+1}-T_n} \phi_{T_n} \in C_b^+(\mathbb{R}^8). \end{aligned}$$

For $n = 0$ this is clear as $\phi_0 \in C_b^+$. Assume (101) for $n - 1$ and consider n . Then $\phi_{T_n} = f_{j,j'}(\phi_{T_n-}) \in \mathcal{M}_{\text{tem}}$ and, for $t \in [T_n, T_{n+1})$,

$$\phi_t(x) = \mathbf{S}_{t-T_n} \phi_{T_n}(x) = \int \mathbf{p}_{t-T_n}(x, y) \phi_{T_n}(dy).$$

It is easy to see that if $f \in \mathcal{C}_{\text{exp}}$, then $\langle \mathbf{S}_{t-T_n} \phi_{T_n}, f \rangle = \langle \phi_{T_n}, \mathbf{S}_{t-T_n} f \rangle$ is continuous in t (e.g., use dominated convergence and Lemma 6.2(ii) of [42]) and so ϕ_t is continuous on $[T_n, T_{n+1})$ and $\phi_{T_{n+1}-} = \mathbf{S}_{T_{n+1}-T_n} \phi_{T_n}$. For $t > T_n$, use the bound $\mathbf{p}_{t-T_n}(x, y) \leq c e^{\lambda|x|} e^{-\lambda|y|}$ [c, λ may depend on (t, T_n)] and dominated convergence to conclude that $\phi_t(\cdot)$ is continuous for all $t \in (T_n, T_{n+1})$ and the same is true for $\phi_{T_{n+1}-}(\cdot)$. For boundedness use the induction hypothesis to see that

$$\phi_{T_n} \leq \|\phi_{T_n-}\|_\infty \delta_{x_j-x_{j'}} dx$$

and so (take $j = 1$ and $j' = 2$ for definiteness)

$$\begin{aligned} \phi_t(x) &\leq \|\phi_{T_n-}\|_\infty \int \mathbf{p}_{t-T_n}(x, y_1, y_1, y_3, y_4) dy_1 dy_3 dy_4 \\ &\leq \|\phi_{T_n-}\|_\infty p_{2(t-T_n)}(x_1, x_2) \\ &\leq c(t - T_n)^{-1} \|\phi_{T_n-}\|_\infty < \infty. \end{aligned}$$

The same reasoning shows that $\phi_{T_{n+1}-}$ is bounded. This completes the inductive proof of (101).

It is clear from (101) that V_t has sample paths in $D(\mathbb{R}_+, \mathcal{I})$ a.s. Let $\hat{P}_{\mathbf{V}_0}$ denote the law of \mathbf{V} on $D(\mathbb{R}_+, \mathcal{I})$.

THEOREM 32 (Limiting moment dual process **V**). *Assume $\gamma/\sigma^2 < (c_{rw}\pi\sqrt{6})^{-1}$, $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ where c_{rw} is given by (31) and ${}^\varepsilon\mathbf{X}$ is the solution to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma,\varepsilon}$ of Lemma 6. Let $\psi : \mathcal{M}_f^2(\mathbb{R}^2) \rightarrow \mathbb{R}_+$ be a bounded continuous map and let $\{\varepsilon_m\}_{m \geq 1}$ be a sequence of positive numbers with $\varepsilon_m \downarrow 0$. Assume either of the following holds:*

- (a) $0 = \delta < t$ and $\mathbf{X}_0 \in \mathcal{M}_{f,se}$;
- (b) $0 < \delta < t$, $\{\psi \neq 0\} \subseteq \{(\mu^1, \mu^2) : \mu^1(\mathbb{R}^2) + \mu^2(\mathbb{R}^2) \leq K\}$ for some K , and the law of ${}^{\varepsilon_m}\mathbf{X}_\delta$ converges weakly in $\mathcal{M}_f^2(\mathbb{R}^2)$ as $m \rightarrow \infty$ to a law $P_{\mathbf{X}_0}(\mathbf{X}_\delta \in \cdot)$.

Then, for any $\phi_0 \in C_b^+(\mathbb{R}^8)$, $I_0 \subseteq \{1, \dots, 4\}$,

$$\begin{aligned} & \lim_{m \rightarrow \infty} E_{\varepsilon_m \mathbf{X}_0} F(\mathbf{V}_0, {}^{\varepsilon_m}\mathbf{X}_t) \psi({}^{\varepsilon_m}\mathbf{X}_\delta) \\ &= \hat{E}_{\mathbf{V}_0} \times E_{\mathbf{X}_0} \left(F(\mathbf{V}_{t-\delta}, \mathbf{X}_\delta) \psi(\mathbf{X}_\delta) \right. \\ & \quad \left. \times \exp \left[2\gamma \int_0^{t-\delta} ds \left\{ \binom{|I_s|}{2} + \binom{|I_s^c|}{2} \right\} \right] \right) < \infty. \end{aligned}$$

REMARK 33. The proof (given below) is independent of the uniqueness results in Theorem 11 and will in fact be used in the derivation of uniqueness in [12]. By (101), $\phi_{t-\delta} \in C_b^+(R^8)$ a.s. and so, on the right-hand side of the above,

$$\begin{aligned} & F(\phi_{t-\delta}, I_{t-\delta}, \mathbf{X}_\delta) \\ (102) \quad &= \int \phi_{t-\delta}(x_1, \dots, x_4) \prod_{i \in I_{t-\delta}} X_\delta^1(dx_i) \prod_{j \notin I_{t-\delta}} X_\delta^2(dx_j) \quad \text{a.s.} \end{aligned}$$

The proof requires the following bound on ${}^\varepsilon p$ which is proved in Appendix A.

LEMMA 34. *If $r \in (0, 1)$, then*

$$(103) \quad \sup\{{}^\varepsilon p_s(x, y) : 0 < s, 0 < \varepsilon, |y - x| > s^{r/2} + \varepsilon^r\} =: c_{34} < \infty.$$

If $p, \varepsilon > 0$, define

$$(104) \quad \mathcal{E}_{\varepsilon,p}({}^\varepsilon\mathbf{X}_t) := \sup_{s>0} s^p \langle {}^\varepsilon X_t^1 \times {}^\varepsilon X_t^2, {}^\varepsilon p_s \rangle$$

and

$$(105) \quad \mathcal{E}_p(\mathbf{X}_t) := \sup_{s>0} s^p \langle X_t^1 \times X_t^2, p_s \rangle + \langle X_t^1, 1 \rangle \langle X_t^2, 1 \rangle.$$

The proof of case (a) also uses the following result, which is a simple consequence of the previous lemma.

LEMMA 35. *If $\mathbf{X}_0 \in \mathcal{M}_{f,se}$, then for any $0 < p' < p < 1$ there is a $c_{35} = c_{35}(p, p', \sigma)$ so that*

$$(106) \quad \sup_{0 < \varepsilon} \mathfrak{E}_{\varepsilon, p}(\varepsilon \mathbf{X}_0) \leq c_{35} \mathfrak{E}_{p'}(\mathbf{X}_0) < \infty.$$

PROOF. By Lemma 8 and the definition of $\varepsilon p_s(x)$ we have

$$(107) \quad \varepsilon p_s(x) \leq c_0(\sigma)(s^{-1} \wedge \varepsilon^{-2}).$$

If $\varepsilon > 0$ and $p' \in (0, 1)$, then

$$(108) \quad \begin{aligned} & \int_{\mathbb{R}^2} X_0^1(dx_1) \int_{\mathbb{R}^2} X_0^2(dx_2) \mathbf{1}(|x_1 - x_2| \leq \varepsilon) \\ & \leq c_1(\sigma) \varepsilon^2 \langle X_0^1 \times X_0^2, p_{\varepsilon^2} \rangle \leq c_1 \varepsilon^{2(1-p')} \mathfrak{E}_{p'}(\mathbf{X}_0). \end{aligned}$$

If $p, r \in (0, 1)$, then (107) and (108) show that

$$\begin{aligned} & s^p \iint \mathbf{1}(|x_1 - x_2| \leq s^{r/2} + \varepsilon^r) \varepsilon p_s(x_1, x_2) \varepsilon X_0^1(dx_1) \varepsilon X_0^2(dx_2) \\ & \leq s^p c_0(\sigma)(s^{-1} \wedge \varepsilon^{-2}) \iint \mathbf{1}(|x_1 - x_2| \leq 4(s^{r/2} + \varepsilon^r)) X_0^1(dx_1) X_0^2(dx_2) \\ & \leq s^p c_0(\sigma)(s^{-1} \wedge \varepsilon^{-2}) \iint \mathbf{1}(|x_1 - x_2| \leq 8(s^{r/2} \vee \varepsilon^r)) X_0^1(dx_1) X_0^2(dx_2) \\ & \leq c_2 \mathfrak{E}_{p'}(\mathbf{X}_0) s^p (s \vee \varepsilon^2)^{r(1-p')-1} \\ & \leq 2c_2 \mathfrak{E}_{p'}(\mathbf{X}_0) (s \vee \varepsilon^2)^{p-1+r(1-p')}. \end{aligned}$$

Let $0 < p' < p < 1$ and choose $r = r(p', p)$ sufficiently close to 1 so that the exponent of s in the above is positive. Use the above to bound $s \leq 1$ and Lemma 8(b) to handle $s > 1$ and conclude that

$$(109) \quad \begin{aligned} & \sup_{0 < \varepsilon, s} s^p \iint \mathbf{1}(|x_1 - x_2| \leq s^{r/2} + \varepsilon^r) \varepsilon p_s(x_1, x_2) \varepsilon X_0^1(dx_1) \varepsilon X_0^2(dx_2) \\ & \leq 2c_2 \mathfrak{E}_{p'}(\mathbf{X}_0) + c_{rw} \sigma^{-2} \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle. \end{aligned}$$

Combine this with Lemma 34 and (108) to see that

$$(110) \quad \sup_{0 < \varepsilon} \mathfrak{E}_{\varepsilon, p}(\varepsilon \mathbf{X}_0) \leq 2c_2 \mathfrak{E}_{p'}(\mathbf{X}_0) + (c_{34} + c_{rw} \sigma^{-2}) \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle.$$

The result follows. \square

The proof of case (b) of Theorem 32 will use the following lemma.

LEMMA 36. Let $0 < p < 1$ and $\delta > 0$:

(a) There is a $c_{36} = c_{36}(\sigma, p)$ so that for any $\varepsilon > 0, \eta \in (0, 1]$ there is a random variable $Z(\varepsilon, \eta, p, \delta)$ satisfying

$$\mathcal{E}_{\varepsilon,p}(\varepsilon \mathbf{X}_\delta) \leq c_{36} \eta^{p-1} \langle \varepsilon X_\delta^1, 1 \rangle \langle \varepsilon X_\delta^2, 1 \rangle + Z(\varepsilon, \eta, p, \delta),$$

and $E(Z(\varepsilon, \eta, p, \delta)) \leq c_{36} \delta^{-1} \eta^{p/2} \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle$.

(b) $\sup_{0 < \varepsilon} E(\mathcal{E}_{\varepsilon,p}(\varepsilon \mathbf{X}_\delta)) \leq c_{36} (1 + \delta^{-1}) \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle$.

PROOF. (a) Lemma 8 implies that $\varepsilon p_s \leq c_1 (s^{-1} \wedge \varepsilon^{-2})$. This, together with Lemma 34, implies, for $\varepsilon > 0$ and $r = 1 - \frac{p}{2}$,

$$\begin{aligned} & \mathcal{E}_{\varepsilon,p}(\varepsilon \mathbf{X}_\delta) \\ & \leq c_1 \langle \varepsilon X_\delta^1, 1 \rangle \langle \varepsilon X_\delta^2, 1 \rangle + \sup_{s \leq 1} s^p \int \mathbf{p}_s(x_1, x_2) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2) \\ (111) \quad & \leq (c_1 + c_{34}) \langle \varepsilon X_\delta^1, 1 \rangle \langle \varepsilon X_\delta^2, 1 \rangle \\ & \quad + \sup_{s \leq 1} c_1 s^p (s^{-1} \wedge \varepsilon^{-2}) \int \mathbf{1}(|x_1 - x_2| \leq (s^{r/2} + \varepsilon^r)) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2). \end{aligned}$$

The second term on the far right-hand side is bounded by

$$\begin{aligned} & \sup_{s \leq 1} c_1 (s \vee \varepsilon^2)^p (s \vee \varepsilon^2)^{-1} \int \mathbf{1}(|x_1 - x_2| \leq 2(s \vee \varepsilon^2)^{r/2}) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2) \\ (112) \quad & \leq c_1 \eta^{p-1} \langle \varepsilon X_\delta^1, 1 \rangle \langle \varepsilon X_\delta^2, 1 \rangle \\ & \quad + c_1 \sup_{\varepsilon^2 \leq s \leq \eta} s^{p-1} \int \mathbf{1}(|x_1 - x_2| \leq 2s^{r/2}) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2), \end{aligned}$$

where the second term is defined to be 0 if $\varepsilon^2 > \eta$. If $s \in [2^{-k-1}, 2^{-k}]$, then

$$\begin{aligned} & s^{p-1} \int \mathbf{1}(|x_1 - x_2| \leq 2s^{r/2}) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2) \\ & \leq 2^{1-p} 2^{-k(p-1)} \int \mathbf{1}(|x_1 - x_2| \leq 2^{1-rk/2}) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2). \end{aligned}$$

Use this in (112) and then (111) to see that

$$(113) \quad \mathcal{E}_{\varepsilon,p}(\varepsilon \mathbf{X}_\delta) \leq c_2(\sigma, p) \eta^{p-1} \langle \varepsilon X_\delta^1, 1 \rangle \langle \varepsilon X_\delta^2, 1 \rangle + Z(\varepsilon, \eta, p, \delta),$$

where

$$Z(\varepsilon, \eta, p, \delta) = c_2 \sum_{2^{-k} \leq 2\eta} 2^{k(1-p)} \int \mathbf{1}(|x_1 - x_2| \leq 2^{1-rk/2}) \varepsilon X_\delta^1(dx_1) \varepsilon X_\delta^2(dx_2).$$

Proposition 15(b) shows that

$$\begin{aligned}
 & E(Z(\varepsilon, \eta, p, \delta)) \\
 &= c_2 \sum_{2^{-k} \leq 2\eta} 2^{k(1-p)} \iiint \mathbf{1}(|x_1 - x_2| \leq 2^{1-rk/2}) \\
 &\quad \times \varepsilon p_\delta(x_1, y_1) \varepsilon p_\delta(x_2, y_2) d^\varepsilon x_1 d^\varepsilon x_2 \varepsilon X_0^1(dy_1) \varepsilon X_0^2(dy_2) \\
 &\leq c_2 c_{\text{rw}} \sigma^{-2} \delta^{-1} \sum_{2^{-k} \leq 2\eta} 2^{k(1-p)} \pi 2^{2-rk} \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle \\
 &\leq c_3(p, \sigma) \delta^{-1} \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle \sum_{2^{-k} \leq 2\eta} 2^{-kp/2} \\
 &\leq c_4(p, \sigma) \delta^{-1} \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle \eta^{p/2}.
 \end{aligned}$$

Therefore, (113) implies (a). To derive (b), take $\eta = 1$ in (a) and note that $E(\langle \varepsilon X_\delta^1, 1 \rangle \langle \varepsilon X_\delta^2, 1 \rangle) = \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle$ by Proposition 15(b). \square

NOTATION 37. Let $c_{37}(\sigma^2) = c_{\text{rw}} \sigma^{-2}$. Then Lemma 8 implies

$$(114) \quad \varepsilon p_t(x) \leq c_{37} t^{-1} \quad \forall \varepsilon > 0, t > 0, x \in \varepsilon \mathbb{Z}^2.$$

Let $U_n = T_n - T_{n-1}$ ($n \geq 1$) be the interjump times for the dual process \mathbf{V}_t^ε .

The next result is the key technical bound on our lattice dual process. It will be used in Corollary 40 and Lemma 43 below to get a uniform (in ε and x) bound on $\phi_\varepsilon(x)$ and will provide the uniform integrability required in the proof of Theorem 32. Finally it will play a critical role in the derivation of the L^2 bounds on the increments of the approximate collision local times of the rescaled lattice processes which underly the tightness needed in Theorem 11(c) (see the proof of Proposition 46 at the end of Section 4).

LEMMA 38. Let $\phi_0 \in C_b^+(\mathbb{R}^8)$, $I_0 \subset \{1, 2, 3, 4\}$ and $n_0 \in \mathbb{Z}_+$. Assume there are distinct random indices $\{i_1, i_2\} \subset \{1, 2, 3, 4\}$ and a measurable map $f: \mathbb{R}_+ \Omega \rightarrow \mathbb{R}_+$ such that $t \rightarrow f(t, \omega)$ is continuous $\hat{P}_{\phi_0, I_0}^\varepsilon$ -a.s. and

$$(115) \quad \begin{aligned} \phi_t^\varepsilon(y_1, y_2, y_3, y_4) &\leq f(t, \omega) \varepsilon p_{2(t-T_{n_0})}(y_{i_1} - y_{i_2}), \\ &i_1 \in I_t, i_2 \notin I_t, \text{ for } T_{n_0} \leq t < T_{n_0+1}, \hat{P}_{\phi_0, I_0}^\varepsilon\text{-a.s.} \end{aligned}$$

Let

$$\rho_{n_0}^f(s) = \begin{cases} f(T_{n_0+1}) \left(\prod_{k=n_0+2}^n \left(\frac{c_{37}}{U_{k-1} + U_k} \right) \right) \frac{c_{37}}{U_n + s - T_n}, & \text{if } T_n \leq s < T_{n+1}, n > n_0, \\ f(s), & \text{if } T_{n_0} \leq s < T_{n_0+1}. \end{cases}$$

Then there are random indices $\{i_1^n, i_2^n : n \geq n_0\} \subset \{1, 2, 3, 4\}$ such that

$$\begin{aligned} \phi_s^\varepsilon(y) &\leq \rho_{n_0}^f(s)^\varepsilon p_{2(s-T_n)}(y_{i_1^n} - y_{i_2^n}), \\ i_1^n \in I_s, i_2^n \in I_s^c, T_n \leq s < T_{n+1}, \forall n \geq n_0, \hat{P}_{\phi_0, I_0}^\varepsilon\text{-a.s.} \end{aligned}$$

PROOF. We proceed by induction on $n \geq n_0$. If $n = n_0$, the required result is our hypothesis (115). Assume the result holds for $n - 1$ ($n - 1 \geq n_0$) and consider n . Then

$$\phi_{T_n-}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_1^{n-1}} - y_{i_2^{n-1}}), \quad i_1^{n-1} \in I_{T_n-}, i_2^{n-1} \notin I_{T_n-}.$$

We consider several cases in analyzing the jump at T_n . We will write (i_1, i_2) for (i_1^{n-1}, i_2^{n-1}) and use i_3, i_4 to denote the distinct indices in $\{1, 2, 3, 4\} - \{i_1, i_2\}$.

Case 1. i_1 switches via $i_3 \in I_{T_n-}$.

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_3} - y_{i_2})^\varepsilon p_0(y_{i_1} - y_{i_3}), \quad I_{T_n} \supset \{i_3\}, I_{T_n}^c \supset \{i_1, i_2\}.$$

Case 2. i_2 switches via $i_3 \in I_{T_n-}^c$.

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_1} - y_{i_3})^\varepsilon p_0(y_{i_2} - y_{i_3}), \quad I_{T_n} \supset \{i_1, i_2\}, I_{T_n}^c \supset \{i_3\}.$$

Case 3. $i_3 \in I_{T_n-}$ switches via i_1 .

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_1} - y_{i_2})^\varepsilon p_0(y_{i_3} - y_{i_1}), \quad I_{T_n} \supset \{i_1\}, I_{T_n}^c \supset \{i_2, i_3\}.$$

Case 4. $i_3 \in I_{T_n-}^c$ switches via i_2 .

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_1} - y_{i_2})^\varepsilon p_0(y_{i_3} - y_{i_2}), \quad I_{T_n} \supset \{i_1, i_3\}, I_{T_n}^c \supset \{i_2\}.$$

Case 5. $i_3 \in I_{T_n-}$ switches via $i_4 \in I_{T_n-}$.

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_1} - y_{i_2})^\varepsilon p_0(y_{i_3} - y_{i_4}), \quad I_{T_n} = \{i_1, i_4\}, I_{T_n}^c = \{i_2, i_3\}.$$

Case 6. $i_3 \in I_{T_n-}^c$ switches via $i_4 \in I_{T_n-}^c$.

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{i_1} - y_{i_2})^\varepsilon p_0(y_{i_3} - y_{i_4}), \quad I_{T_n} = \{i_1, i_3\}, I_{T_n}^c = \{i_2, i_4\}.$$

We can now introduce new random indices $\{\hat{i}_j : j \leq 4\} = \{1, 2, 3, 4\}$ and reduce these six cases to essentially two cases.

Case A.

$$\begin{aligned} \phi_{T_n}^\varepsilon(y) &\leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{\hat{i}_1} - y_{\hat{i}_3})^\varepsilon p_0(y_{\hat{i}_1} - y_{\hat{i}_2}), \\ \hat{i}_1 \in I_{T_n}, \{\hat{i}_2, \hat{i}_3\} &\subset I_{T_n}^c \text{ or } \hat{i}_1 \in I_{T_n}^c, \{\hat{i}_2, \hat{i}_3\} \subset I_{T_n}. \end{aligned}$$

Case B.

$$\phi_{T_n}^\varepsilon(y) \leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n}(y_{\hat{i}_3} - y_{\hat{i}_4})^\varepsilon p_0(y_{\hat{i}_1} - y_{\hat{i}_2}), \quad I_{T_n} = \{\hat{i}_1, \hat{i}_3\}, I_{T_n}^c = \{\hat{i}_2, \hat{i}_4\}.$$

For Case A use (114) to see that, for $T_n \leq t < T_{n+1}$,

$$\begin{aligned} \phi_t^\varepsilon(y) &\leq \rho_{n_0}^f(T_n-) \\ &\quad \times \int^\varepsilon p_{2U_n+t-T_n}(z_{\hat{i}_1} - y_{\hat{i}_3})^\varepsilon p_{t-T_n}(z_{\hat{i}_1} - y_{\hat{i}_2})^\varepsilon p_{t-T_n}(z_{\hat{i}_1} - y_{\hat{i}_1})^\varepsilon d^\varepsilon z_{\hat{i}_1} \\ &\leq \rho_{n_0}^f(T_n-) c_{37} (2U_n + t - T_n)^{-1} \varepsilon p_{2(t-T_n)}(y_{\hat{i}_2} - y_{\hat{i}_1}) \\ &\leq \rho_{n_0}^f(t) \varepsilon p_{2(t-T_n)}(y_{\hat{i}_2} - y_{\hat{i}_1}), \end{aligned}$$

where $\hat{i}_1 \in I_t$, $\hat{i}_2 \in I_t^c$ or conversely.

For Case B we use (114) to see that, for $T_n \leq t < T_{n+1}$,

$$\begin{aligned} \phi_t^\varepsilon &\leq \rho_{n_0}^f(T_n-)^\varepsilon p_{2U_n+t-T_n}(y_{\hat{i}_3} - y_{\hat{i}_4})^\varepsilon p_{2(t-T_n)}(y_{\hat{i}_2} - y_{\hat{i}_1}) \\ &\leq \rho_{n_0}^f(T_n-) c_{37} (U_n + t - T_n)^{-1} \varepsilon p_{2(t-T_n)}(y_{\hat{i}_2} - y_{\hat{i}_1}) \\ &\leq \rho_{n_0}^f(t) \varepsilon p_{2(t-T_n)}(y_{\hat{i}_2} - y_{\hat{i}_1}), \end{aligned}$$

where $\hat{i}_1 \in I_t$ and $\hat{i}_2 \in I_t^c$.

In either case it is clear how to define i^n so that the required result holds on $T_n \leq t < T_{n+1}$, $\hat{P}_{\phi_0, T_0}^\varepsilon$ -a.s. This completes the inductive proof. \square

NOTATION 39. Write $\rho_{n_0}(s)$ for $\rho_{n_0}^f(s)$ when $f \equiv 1$.

COROLLARY 40. Let $\phi_0 \in C_b^+((\mathbb{R}^2)^4)$ and $I_0 \subset \{1, 2, 3, 4\}$. There are random indices $\{i_1^n, i_2^n : n \geq 1\} \subset \{1, 2, 3, 4\}$ such that, $\hat{P}_{\phi_0, I_0}^\varepsilon$ -a.s. $\forall n \geq 1$,

$$i_1^n \in I_s, \quad i_2^n \in I_s^c \quad \text{if } T_n \leq s < T_{n+1}$$

and

$$(116) \quad \begin{aligned} \phi_s^\varepsilon(y) &\leq \|\phi_0\|_\infty \left[\mathbf{1}(s < T_1) \right. \\ &\quad \left. + \sum_{n=1}^\infty \mathbf{1}(T_n \leq s < T_{n+1}) \rho_1(s)^\varepsilon p_{2(s-T_n)}(y_{i_1^n} - y_{i_2^n}) \right] \end{aligned}$$

$$(117) \quad \begin{aligned} &\leq \|\phi_0\|_\infty \left[\mathbf{1}(s < T_1) + \sum_{n=1}^\infty \mathbf{1}(T_n \leq s < T_{n+1}) \rho_1(s) c_{37} (s - T_n)^{-1} \right] \\ &\equiv \bar{\phi}(s). \end{aligned}$$

PROOF. We check (115) of the previous lemma for $n_0 = 1$ and $f \equiv \|\phi_0\|_\infty$. Clearly $\phi_{T_1}^\varepsilon = \lim_{t \uparrow T_1} \varepsilon S_t^{(4)} \phi_0 \leq \|\phi_0\|_\infty$. Therefore the definition of ϕ_t^ε shows that

$\phi_{T_1}^\varepsilon(y) \leq \|\phi_0\|_\infty \varepsilon p_0(y_{i_1} - y_{i_2})$ for some $i_1 \in I_{T_1}, i_2 \notin I_{T_1}$. It follows that, for $T_1 \leq t < T_2, i_1 \in I_t, i_2 \notin I_t$, and

$$\begin{aligned} \phi_t^\varepsilon(y) &\leq \|\phi_0\|_\infty \int \varepsilon p_{s-T_1}(z - y_{i_1}) \varepsilon p_{s-T_1}(z - y_{i_2}) d^\varepsilon z \\ &= \|\phi_0\|_\infty \varepsilon p_{2(s-T_1)}(y_{i_1} - y_{i_2}). \end{aligned}$$

This verifies (115), and (116) follows from Lemma 38, as this inequality is trivial for $s < T_1$. The second inequality is then clear by (114). \square

LEMMA 41. *Let ${}^\varepsilon S_t$ denote the semigroup of the nearest neighbor continuous time random walk ξ_t^ε on $\varepsilon\mathbb{Z}^d$ which jumps to a nearest neighbor at rate $d\varepsilon^{-2}\sigma^2$ and let S_t denote the semigroup of the d -dimensional Brownian motion with variance parameter σ^2 . Let $f^\varepsilon : \varepsilon\mathbb{Z}^d \rightarrow \mathbb{R}$, let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy $\sup_{\varepsilon>0} \|f^\varepsilon\|_\infty < \infty$ and let $\lim_{\varepsilon\downarrow 0} f^\varepsilon(x_\varepsilon) = f(x)$ whenever $\lim_{\varepsilon\downarrow 0} x_\varepsilon = x (x_\varepsilon \in \varepsilon\mathbb{Z}^d)$. Then $\|f\|_\infty < \infty$ and*

$$\lim_{\varepsilon\downarrow 0} {}^\varepsilon S_t f^\varepsilon(x_\varepsilon) = S_t f(x).$$

PROOF. The first assertion is obvious. Let $\varepsilon_n \downarrow 0$. By Skorohod's theorem we may assume $\xi_t^{\varepsilon_n} \rightarrow B_t$ a.s. where $\xi_0^{\varepsilon_n} = B_0 = 0$. Then $x_{\varepsilon_n} + \xi_t^{\varepsilon_n} \rightarrow x + B_t$ a.s. and the result follows by dominated convergence. \square

NOTATION 42. If $x \in \mathbb{R}$, let $[x]_\varepsilon = [\varepsilon^{-1}x]\varepsilon$ for each $\varepsilon > 0$. If $x = (x_1, \dots, x_d)$, let $[x]_\varepsilon = ([x_1]_\varepsilon, \dots, [x_d]_\varepsilon) \in (\varepsilon\mathbb{Z})^d$.

LEMMA 43. *If $\phi_0 \in C_b^+(\mathbb{R}^8)$ and $I_0 \subset \{1, 2, 3, 4\}$, then, for each $t > 0$, the following hold:*

- (a) $\sup_{\varepsilon>0} \sup_{x \in \varepsilon\mathbb{Z}^8} \phi_t^\varepsilon(x) < \infty, \hat{P}_{\phi_0, I_0}$ -a.s.
- (b) $\lim_{\varepsilon\downarrow 0} \sup_{x \in \varepsilon\mathbb{Z}^8, |x| \leq K} |\phi_t^\varepsilon(x) - \phi_t(x)| = 0, \forall K > 0, \hat{P}_{\phi_0, I_0}$ -a.s.

PROOF. Statement (a) follows from Corollary 40 since $\bar{\phi}(t) < \infty$ for $t \notin \{T_n : n \geq 1\}$ which holds \hat{P}_{ϕ_0, I_0} -a.s.

For (b), it suffices to show that, for a fixed sequence $\varepsilon_k \downarrow 0$,

$$(118) \quad \begin{aligned} \lim_{k \rightarrow \infty} \phi_t^{\varepsilon_k}(x_k) &= \phi_t(x) \quad \hat{P}_{\phi_0, I_0}\text{-a.s. whenever } x_k \in \varepsilon_k\mathbb{Z}^8, x \in \mathbb{R}^8 \text{ are} \\ &\text{random points satisfying } \lim_{k \rightarrow \infty} x_k = x \hat{P}_{\phi_0, I_0}\text{-a.s.} \end{aligned}$$

This in turn will follow by establishing

$$(118_n) \text{ (a) } (118) \text{ holds a.s. on } \{\omega : T_n < t < T_{n+1}\} \text{ for } \{x_k\}, x \text{ as above, and}$$

(119)

$$(118_n) \text{ (b) } \lim_{k \rightarrow \infty} \phi_{T_{n+1}-}^{\varepsilon_k}(x_k) = \phi_{T_{n+1}-}(x) \text{ for } \{x_k\}, x \text{ as above, for all } n \in \mathbb{Z}_+.$$

Clearly (118_n)(a) $\forall n \geq 0$ suffices but (b) helps in our inductive proof. On $\{t < T_1\}$,

$$\phi_t^{\varepsilon_k}(x_k) = \mathbf{S}_t^{\varepsilon_k} \phi_0(x_k), \quad \phi_t(x) = \mathbf{S}_t \phi_0(x)$$

and so Lemma 41 implies (118₀)(a). Since $\phi_{T_1-}^{\varepsilon_k}(x_k) = \varepsilon_k \mathbf{S}_{T_1} \phi_0(x_k)$ and $\phi_{T_1-}(x_k) = \mathbf{S}_{T_1}(x_k)$, the same result also gives (118₀)(b). Assume (118_m) for $m < n$. Consider

$$\{T_n < t < T_{n+1}\} \cap \{j \text{ switches via } j' \text{ at } T_n\}.$$

On this event

$$(120) \quad \begin{aligned} \phi_t^{\varepsilon_k}(x_k) &= \int \phi_{T_n-}^{\varepsilon_k}([\pi_{j,j'}y]_{\varepsilon_k})^{\varepsilon_k} p_{t-T_n}([y_j]_{\varepsilon_k} - (x_k)_{j'}) \\ &\quad \times \prod_{i \neq j'}^{\varepsilon_k} p_{t-T_n}([y_i]_{\varepsilon_k} - (x_k)_i) d\hat{y}^{j'} \end{aligned}$$

and

$$(121) \quad \phi_t(x) = \int \phi_{T_n-}(\pi_{j,j'}y) p_{t-T_n}(y_j - x_{j'}) \prod_{i \neq j'} p_{t-T_n}(y_i - x_i) d\hat{y}^{j'},$$

where $d\hat{y}^{j'}$ is the three-dimensional Lebesgue integral with the $y_{j'}$ variable omitted and we write $(x_k)_j$ for the j th component of $x_k \in (\varepsilon_k \mathbb{Z}^2)^4$. By (118_{n-1})(b) and Lemma 8 if $(y_i^k)_{i \neq j'} \rightarrow (y_i)_{i \neq j'}$ as $k \rightarrow \infty$ when $(y_i^k)_{i \neq j'} \in (\varepsilon_k \mathbb{Z}^2)^3$ and $(y_i)_{i \neq j'} \in (\mathbb{R}^2)^3$, then

$$\lim_{k \rightarrow \infty} \phi_{T_n-}^{\varepsilon_k}([\pi_{j,j'}y^k]_{\varepsilon_k})^{\varepsilon_k} p_{t-T_n}([y_j^k]_{\varepsilon_k} - (x_k)_{j'}) = \phi_{T_n-}(\pi_{j,j'}y) p_{t-T_n}(y_j - x_{j'}).$$

Moreover (a) and Lemma 8(a) show that

$$\sup_{\varepsilon_k, y} \phi_{T_n-}^{\varepsilon_k}([\pi_{j,j'}y]_{\varepsilon_k})^{\varepsilon_k} p_{t-T_n}([y_j] - (x_k)_{j'}) < \infty, \quad \hat{P}_{\phi_0, I_0}\text{-a.s.}$$

Now apply Lemma 41 to the six-dimensional random walks with transition function

$$\prod_{i \neq j'}^{\varepsilon_k} p_{t-T_n}([y_i]_{\varepsilon_k} - (x_k)_i)$$

to see that $\lim_{k \rightarrow \infty} \phi_t^{\varepsilon_k}(x_k) = \phi_t(x)$ on $\{T_n < t < T_{n+1}\} \cap \{j \text{ switches via } j' \text{ at } T_n\}$. The same reasoning also proves (118_n)(b). This completes the induction and hence the proof of (b). \square

PROOF OF THEOREM 32. Use the Markov property of $\varepsilon_m \mathbf{X}$ at $t = \delta \geq 0$ and Proposition 28 to see that it suffices to prove

$$(122) \quad \begin{aligned} &\lim_{m \rightarrow \infty} \hat{E}_{\phi_0, I_0}^{\varepsilon_m} \times E_{\varepsilon_m \mathbf{X}_0}^{\varepsilon_m} (F(\phi_{t-\delta}^{\varepsilon_m}, I_{t-\delta}, \varepsilon_m \mathbf{X}_\delta) \psi(\varepsilon_m \mathbf{X}_\delta) \mathcal{E}_{t-\delta}) \\ &= \hat{E}_{\phi_0, I_0} \times E_{\mathbf{X}_0} (F(\phi_{t-\delta}, I_{t-\delta}, \mathbf{X}_\delta) \psi(\mathbf{X}_\delta) \mathcal{E}_{t-\delta}), \end{aligned}$$

where

$$\varepsilon_{t-\delta} = \exp\left\{\gamma \int_0^{t-\delta} \left(\frac{|I_s|}{2}\right) + \left(\frac{|I_s^c|}{2}\right) ds\right\}.$$

By Skorohod’s theorem we may assume $\{\varepsilon_m \mathbf{X}_\delta\}$ and \mathbf{X}_δ are defined on a common $(\Omega, \mathcal{F}, \mathcal{P})$ such that $\varepsilon_m \mathbf{X}_\delta \rightarrow \mathbf{X}_\delta$, P -a.s. and replace the expectations $E_{\varepsilon_m \mathbf{X}_0}$ and $E_{\mathbf{X}_0}$ in (122) with E . We now claim that

$$(123) \quad \begin{aligned} &\lim_{m \rightarrow \infty} F(\phi_{t-\delta}^{\varepsilon_m}, I_{t-\delta}, \varepsilon_m \mathbf{X}_\delta) \psi(\varepsilon_m \mathbf{X}_\delta) \\ &= F(\phi_{t-\delta}, I_{t-\delta}, \mathbf{X}_\delta) \psi(\mathbf{X}_\delta) \quad \hat{P}_{\phi_0, I_0} \times P\text{-a.s.} \end{aligned}$$

As ψ is continuous we only need focus on the “ F terms.” Since $\varepsilon_m \mathbf{X}_\delta \rightarrow \mathbf{X}_\delta$ in $\mathcal{M}_f^2(\mathbb{R})^2$ a.s., $\{\varepsilon_m \mathbf{X}_\delta : m \in \mathbb{N}\}$ are a.s. tight. This together with Lemma 43(b), the fact that $\phi_{t-\delta} \in C_b^+(\mathbb{R}^8)$ \hat{P}_{ϕ_0, I_0} -a.s. [recall (101)] and $\varepsilon_m \mathbf{X}_\delta \rightarrow \mathbf{X}_\delta$ a.s. allow one to prove (123) by an elementary weak convergence argument.

To prove (122) it now suffices to show

$$(124) \quad \begin{aligned} &\{F(\phi_{t-\delta}^{\varepsilon_m}, I_{t-\delta}, \varepsilon_m \mathbf{X}_\delta) \psi(\varepsilon_m \mathbf{X}_\delta) : m \in \mathbb{N}\} \text{ is uniformly integrable} \\ &\text{with respect to } \hat{E}_{\phi_0, I_0} \times E. \end{aligned}$$

Bound ϕ_0 by $\|\phi\|_\infty$ and hence verify (115) with $n_0 = 1$ and $f = \|\phi_0\|_\infty$ through a short calculation. Lemma 38 shows (recall $\rho_{n_0} = \rho_{n_0}^1$) that if $M = M^{m, \delta}(\omega) = \varepsilon_m X_\delta^1(\mathbb{R}^2) + \varepsilon_m X_\delta^2(\mathbb{R}^2)$ and $p \in (0, \frac{1}{2})$, then, $\hat{P}_{\phi_0, I_0} \times P$ -a.s.,

$$(125) \quad \begin{aligned} &F(\phi_{t-\delta}^{\varepsilon_m}, I_{t-\delta}, \varepsilon_m \mathbf{X}_\delta) \\ &\leq \|\phi_0\|_\infty \left[\mathbf{1}(t - \delta < T_1) \left((\varepsilon_m X_\delta^1(\mathbb{R}^2))^4 + (\varepsilon_m X_\delta^2(\mathbb{R}^2))^4 \right) \right. \\ &\quad \left. + \sum_{n=1}^\infty \mathbf{1}(T_n \leq t - \delta < T_{n+1}) \rho_1(t - \delta) (2(t - \delta - T_n))^{-p} \right. \\ &\quad \left. \times \mathcal{E}_{\varepsilon_m, p}(\varepsilon_m \mathbf{X}_\delta) \left((\varepsilon_m X_\delta^1(\mathbb{R}^2))^2 + (\varepsilon_m X_\delta^2(\mathbb{R}^2))^2 \right) \right] \\ &\leq \|\phi_0\|_\infty (M^4 + M^2) \\ &\quad \times \left[\mathbf{1}(t - \delta < T_1) + \left(\mathbf{1}(T_1 \leq t - \delta < T_2) (t - \delta - T_1)^{-p} \right. \right. \\ &\quad \left. \left. + \sum_{n=2}^\infty \mathbf{1}(T_n \leq t - \delta < T_{n+1}) c_{37}^{n-1} \left(\prod_{k=3}^n (U_{k-1} + U_k)^{-1} \right) \right. \right. \\ &\quad \left. \left. \times (U_n + t - \delta - T_n)^{-1} (t - \delta - T_n)^{-p} \right) \mathcal{E}_{\varepsilon_m, p}(\varepsilon_m \mathbf{X}_\delta) \right]. \end{aligned}$$

Now for $n \geq 2$, either U_n is exponential with rate 2γ and U_{n+1} is exponential with rate 3γ or conversely. Therefore if α_n is the rate of U_n we have

$$\begin{aligned}
 & \hat{E}_{\phi_0, I_0} \left(\mathbf{1}(T_n \leq s < T_{n+1}) \left(\prod_{k=3}^n (U_{k-1} + U_k)^{-1} \right) (U_n + s - T_n)^{-1} (s - T_n)^{-p} \right) \\
 &= \int_0^\infty du_1 \cdots \int_0^\infty du_n \mathbf{1} \left(\sum_1^n u_i \leq s \right) e^{-\alpha_{n+1}(s - \sum_1^n u_j)} e^{-\sum_1^n \alpha_j u_j} \\
 (126) \quad & \times \prod_1^n \alpha_j \left(\prod_{k=3}^n (u_{k-1} + u_k)^{-1} \right) \left(s - \sum_1^{n-1} u_i \right)^{-1} \left(s - \sum_1^n u_i \right)^{-p} \\
 & \leq \alpha_1 6^{n/2} \gamma^{n-1} \int_0^\infty du_1 \cdots \int_0^\infty du_n \mathbf{1} \left(\sum_1^n u_i \leq s \right) \\
 & \quad \times \left(\prod_{k=2}^{n-1} (u_{k+1} + u_k)^{-1} \right) \left(s - \sum_1^{n-1} u_i \right)^{-1} \left(s - \sum_1^n u_i \right)^{-p}.
 \end{aligned}$$

Now change variables and set $v_k = u_{k+1}$, $1 \leq k \leq n-1$, $v_n = \sum_1^n u_i$. Note also $\alpha_1 \leq 6\gamma$ as the largest jump rate occurs when $I_0 = \emptyset$ or $I_0^c = \emptyset$. If $J_n(s, T)$ is as in Corollary 61 in Appendix B, then the far right-hand side of (126) is at most

$$\begin{aligned}
 & \gamma^n 6^{1+n/2} \int_0^s (s - v_n)^{-p} \\
 & \quad \times \left[\int_{\mathbb{R}_+^{n-1}} \mathbf{1} \left(\sum_1^{n-1} v_i \leq v_n \right) \right. \\
 & \quad \quad \left. \times \prod_{i=1}^{n-2} (v_i + v_{i+1})^{-1} (s - v_n + v_{n-1})^{-1} dv_1 \cdots dv_{n-1} \right] dv_n \\
 & \leq \gamma^n 6^{1+n/2} \int_0^s (s - v_n)^{-p} J_{n-1}(s - v_n, v_n) dv_n \\
 & \leq c_{61} \gamma^n 6^{1+n/2} \pi^{n-2} \int_0^s (s - v_n)^{-p} (v_n)^{1/2} (s - v_n)^{-1/2} dv_n,
 \end{aligned}$$

where we have used Corollary 61 with $p = \frac{1}{2}$ in the last line. A simple change of variables shows that if we use the above to bound (126) we arrive at

$$\begin{aligned}
 & \hat{E}_{\phi_0, I_0} \left(\mathbf{1}(T_n \leq s < T_{n+1}) \left(\prod_{k=3}^n (U_{k-1} + U_k)^{-1} \right) (U_n + s - T_n)^{-1} (s - T_n)^{-p} \right) \\
 (127) \quad & \leq c_{61} (1/2) \gamma^n 6^{1+n/2} \pi^{n-2} \int_0^1 w^{1/2} (1-w)^{-1/2-p} dw s^{1-p} \\
 & \leq c_{127}(p) (\gamma \sqrt{6} \pi)^n s^{1-p}.
 \end{aligned}$$

We first establish (124) in case (a). As $\delta = 0$, $M = X_0^1(\mathbb{R}^2) + X_0^2(\mathbb{R}^2)$ is a constant. Lemma 35 and (125) imply that if $p' \in (0, p)$, and

$$\begin{aligned} W(s) &= 1 + \mathbf{1}(T_1 \leq s < T_2)(s - T_1)^{-p} \\ &\quad + \sum_{n=2}^{\infty} \mathbf{1}(T_n \leq s < T_{n+1})c_{37}^{n-1} \\ &\quad \times \prod_{k=3}^n (U_{k-1} + U_k)^{-1} \times (U_n + s - T_n)^{-1}(s - T_n)^{-p}, \end{aligned}$$

then

$$(128) \quad F(\phi_t^{\varepsilon_m}, I_t, \varepsilon_m \mathbf{X}_0) \leq \|\phi_0\|_{\infty}(M^4 + M^2)(1 + c_{35}\mathcal{E}_{p'}(\mathbf{X}_0))W(t).$$

Our assumption on $\gamma\sigma^{-2}$ implies $c_{37}\gamma\sqrt{6\pi} < 1$ and (127) easily implies

$$(129) \quad \hat{E}_{\phi_0 I_0}(W(s)) < \infty \quad \forall s > 0.$$

As the upper bound in (128) is \hat{P}_{ϕ_0, I_0} -integrable and independent of m , and ψ is bounded, the required uniform integrability in (124) follows and the proof is complete in case (a).

Consider the case (b) and write $(\hat{\omega}, \omega)$ for our sample points under $\hat{P}_{\phi_0, I_0} \times P$. Note that $W(t - \delta) \equiv W(t - \delta, \hat{\omega})$. Our hypothesis on ψ and (125) imply, for some $0 < c(\psi) < \infty$,

$$(130) \quad \psi(\varepsilon_m \mathbf{X}_\delta)F(\phi_{t-\delta}^{\varepsilon_m} I_{t-\delta}, \varepsilon_m \mathbf{X}_\delta) \leq c(\psi)W(t - \delta, \hat{\omega})(1 + \mathcal{E}_{\varepsilon_m, p}(\varepsilon_m \mathbf{X}_\delta(\omega))).$$

Fix $\eta > 0$. By Lemma 36 there are random variables $Z(\varepsilon_m, \eta, p, \delta) \equiv Z_m(\omega)$ such that

$$(131) \quad \begin{aligned} \mathcal{E}_{\varepsilon_m, p}(\varepsilon_m \mathbf{X}_\delta(\omega)) &\leq c_{36}\eta^{p-1}X_\delta^1(\mathbb{R}^2)X_\delta^2(\mathbb{R}^2)(\omega) + Z_m(\omega), \\ E(Z_m) &\leq c_{36}\delta^{-1}\eta^{p/2}X_0^1(\mathbb{R}^2)X_0^2(\mathbb{R}^2). \end{aligned}$$

By (129) and Proposition 15(b) we may choose $\varepsilon > 0$ so that $\hat{P}_{\phi_0, I_0} \times P(A) < \varepsilon$ implies $\hat{E}_{\phi_0, I_0} \times E(\mathbf{1}_A W(t - \delta)(1 + X_\delta^1(\mathbb{R}^2)X_\delta^2(\mathbb{R}^2))) < \eta(1 + c_{36}\eta^{p-1})^{-1}$. Then (130) and (131) imply, for A as above,

$$\begin{aligned} &\hat{E}_{\phi_0, I_0} \times E(\mathbf{1}_A F(\phi_{t-\delta}^{\varepsilon_m} I_{t-\delta}, \varepsilon_m \mathbf{X}_\delta)\psi(\varepsilon_m \mathbf{X}_\delta)) \\ &\leq c(\psi)(1 + c_{36}\eta^{p-1})\hat{E}_{\phi_0, I_0} \times E(\mathbf{1}_A W(t - \delta)(1 + X_\delta^1(\mathbb{R}^2)X_\delta^2(\mathbb{R}^2))) \\ &\quad + c(\psi)\hat{E}_{\phi_0, I_0}(W(t - \delta))E(Z_m) \\ &\leq c(\psi)\eta + c(\psi)\hat{E}_{\phi_0, I_0}(W(t - \delta))c_{36}\delta^{-1}X_0^1(\mathbb{R}^2)X_0^2(\mathbb{R}^2)\eta^{p/2}. \end{aligned}$$

This goes to zero as $\eta \downarrow 0$, independently of m , and so (124) holds and the proof is complete in case (b). \square

4. Construction of a solution. In this section we prove Theorem 11(a), (c). Recall that $[x]_\varepsilon = (y_1, y_2) \in \varepsilon\mathbb{Z}^2$ iff $x \in \prod_{i=1}^2 [y_i, y_i + \varepsilon) \equiv C_\varepsilon((y_1, y_2))$. In Section 1.2 we fixed $\mathbf{X}_0 \in \mathcal{M}_{f,\varepsilon}$ and constructed a solution ${}^\varepsilon\mathbf{X}$ to the approximate martingale problem $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma,\varepsilon}$ starting at

$${}^\varepsilon X_0^i(\{x\}) = X_0^i(C_\varepsilon(x)), \quad x \in \varepsilon\mathbb{Z}^2.$$

We assume (33) throughout this section. We use this stronger condition in the proof of a key L^2 estimate in Proposition 46. The following elementary bound is proved in Appendix A.

LEMMA 44. *There is a constant $c_{44} = c_{44}(\sigma^2)$ such that*

$$\int_0^\delta {}^\varepsilon p_s(x) ds \leq c_{44} \left[\left(\frac{\sqrt{\delta}}{\|x\|} \right) \wedge 1 + \log^+ \left(\frac{\sqrt{\delta}}{\|x\|} \right) \right] \quad \forall x \in \varepsilon\mathbb{Z}^2, \forall \delta, \varepsilon > 0.$$

LEMMA 45.

$$(a) \limsup_{\delta \downarrow 0} \sup_{\varepsilon > 0} \int \left[\int_0^\delta {}^\varepsilon p_s(x_1 - x_2) ds \right] {}^\varepsilon X_0^1(dx_1) {}^\varepsilon X_0^2(dx_2) = 0.$$

$$(b) \sup_{\varepsilon > 0} \int \left[\int_0^T {}^\varepsilon p_s(x_1 - x_2) ds \right] {}^\varepsilon X_0^1(dx_1) {}^\varepsilon X_0^2(dx_2) = c_{45}(T) < \infty$$

$\forall T > 0$.

PROOF. Define

$$G_\varepsilon(\mathbf{X}_0) = \int \mathbf{1}(\|x_1 - x_2\| < \sqrt{2}\varepsilon) g(x_1 - x_2) X_0^1(dx_1) X_0^2(dx_2).$$

Note that if $[x_1]_\varepsilon \neq [x_2]_\varepsilon$, then $\|[x_1]_\varepsilon - [x_2]_\varepsilon\| \geq \varepsilon$ and so

$$(132) \quad \frac{\|[x_1]_\varepsilon - [x_2]_\varepsilon\|}{\|x_1 - x_2\|} \geq \frac{\|[x_1]_\varepsilon - [x_2]_\varepsilon\|}{\|[x_1]_\varepsilon - [x_2]_\varepsilon\| + 2\sqrt{2}\varepsilon} \geq (1 + 2\sqrt{2})^{-1} \equiv c_0.$$

We have ${}^\varepsilon p_s(0) \leq c_1(s^{-1} \wedge \varepsilon^{-2})$ by Lemma 8, and so, by Lemma 44,

$$(133) \quad \begin{aligned} & \iint \left[\int_0^\delta {}^\varepsilon p_s(x_1 - x_2) ds \right] {}^\varepsilon X_0^1(dx_1) {}^\varepsilon X_0^2(dx_2) \\ &= \iint \left[\int_0^\delta {}^\varepsilon p_s([x_1]_\varepsilon - [x_2]_\varepsilon) ds \right] X_0^1(dx_1) X_0^2(dx_2) \\ &\leq \iint \left[\int_0^\delta c_1(s^{-1} \wedge \varepsilon^{-2}) ds \right] \mathbf{1}([x_1]_\varepsilon = [x_2]_\varepsilon) X_0^1(dx_1) X_0^2(dx_2) \\ &+ \iint c_{44} \left[1 \wedge \left(\frac{\sqrt{\delta}}{\|[x_1]_\varepsilon - [x_2]_\varepsilon\|} \right) + \log^+ \left(\frac{\sqrt{\delta}}{\|[x_1]_\varepsilon - [x_2]_\varepsilon\|} \right) \right] \\ &\quad \times \mathbf{1}([x_1]_\varepsilon \neq [x_2]_\varepsilon) X_0^1(dx_1) X_0^2(dx_2) \end{aligned}$$

$$\begin{aligned} &\leq c_1 \left(\frac{\varepsilon^2 \wedge \delta}{\varepsilon^2} + \log^+ \frac{\delta}{\varepsilon^2} \right) \left(1 + \log^+ \frac{1}{\sqrt{2\varepsilon}} \right)^{-1} G_\varepsilon(\mathbf{X}_0) \\ &\quad + c_{44} \iint \left[1 \wedge \left(\frac{\sqrt{\delta}}{c_0 \|x_1 - x_2\|} \right) + \log^+ \left(\frac{\sqrt{\delta}}{c_0 \|x_1 - x_2\|} \right) \right] dX_0^1(x_1) dX_0^2(x_2). \end{aligned}$$

We have used (132) in the last line. The second term approaches 0 as $\delta \downarrow 0$ by dominated convergence since $\mathbf{X}_0 \in \mathcal{M}_{f,e}$. This also implies $\lim_{\varepsilon \downarrow 0} G_\varepsilon(\mathbf{X}_0) = 0$ and so the first term in (133) clearly approaches 0 uniformly in $\delta \in (0, 1]$ as $\varepsilon \downarrow 0$. As $G_\varepsilon(\mathbf{X}_0)$ is uniformly bounded in ε , it then follows easily that the first term in (133) approaches 0 uniformly in $\varepsilon > 0$ as $\delta \downarrow 0$. This proves (a). Relation (b) is immediate from (a). \square

Tightness of ${}^\varepsilon \mathbf{X}$ will be proved using bounds on its moments. First and second moments for $\varepsilon = 1$ are easy to derive from (19) and were given in Theorem 2.2(b)(iii) of [14]. Using our definition of ${}^\varepsilon X_t^i$, we then easily get, for $\phi_i : \varepsilon \mathbb{Z}^2 \rightarrow \mathbb{R}_+, i = 1, 2$,

$$(134) \quad \begin{aligned} (i) \quad &E(\langle {}^\varepsilon X_t^i, \phi_i \rangle) = \langle {}^\varepsilon X_0^i, {}^\varepsilon S_t \phi_i \rangle, \\ (ii) \quad &E(\langle {}^\varepsilon X_t^1, \phi_1 \rangle \langle {}^\varepsilon X_t^2, \phi_2 \rangle) = \langle {}^\varepsilon X_0^1, {}^\varepsilon S_t \phi_1 \rangle \langle {}^\varepsilon X_0^2, {}^\varepsilon S_t \phi_2 \rangle. \end{aligned}$$

Our key L^2 -bound is on the increments of

$$\langle L^\varepsilon(t), \phi \rangle \equiv \langle L_{\varepsilon \mathbf{X}}^\varepsilon(t), \phi \rangle = \int_0^t \int \phi(x) {}^\varepsilon X_s^1(x) {}^\varepsilon X_s^2(x) d^\varepsilon x ds.$$

Recall that $d^\varepsilon x$ denotes integration with respect to $\ell^\varepsilon = \sum_{y \in \varepsilon \mathbb{Z}^2} \varepsilon^2 \delta_y$. Recall the notation $\mathcal{E}_{\varepsilon,p}({}^\varepsilon \mathbf{X}_0)$ from Lemma 34 and let

$$\overline{\mathcal{E}}_{\varepsilon,p}({}^\varepsilon \mathbf{X}_0) = \mathcal{E}_{\varepsilon,p}({}^\varepsilon \mathbf{X}_0) [({}^\varepsilon X_0^1(\mathbb{R}^2))^2 + ({}^\varepsilon X_0^2(\mathbb{R}^2))^2].$$

PROPOSITION 46. *There is an $\varepsilon_0 = \varepsilon_0(\gamma, \sigma^2) > 0$, and for any $T > 0$ there is a $c_{46} = c_{46}(T, \gamma, \sigma^2) > 0$ such that for any bounded Borel $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ and any $0 < \varepsilon < \varepsilon_0$,*

$$\begin{aligned} E(\langle L^\varepsilon(t_2), \phi \rangle - \langle L^\varepsilon(t_1), \phi \rangle)^2 &\leq c_{46} \overline{\mathcal{E}}_{\varepsilon,1/2}({}^\varepsilon \mathbf{X}_0) |t_2 - t_1|^{3/2} t_2^{-1} \|\phi\|_\infty^2 \\ &\text{for all } 0 \leq t_1 < t_2 \leq T. \end{aligned}$$

REMARK 47. The power 3/2 is by no means sharp and can easily be improved to $2 - \delta$ for any $\delta > 0$, at the cost of a stronger assumption on $\gamma \sigma^{-2}$. The factor t_2^{-1} will not pose any problems as t_2 is the greater of the two times.

The proof will be given at the end of this section and uses the following bound on a family of iterated integrals for $p = \frac{1}{2}$. We include the more general case here because it will be used in [12] to verify **(IntC)**.

NOTATION 48. If $n \in \mathbb{N} \geq 2$, $p \in (0, 1)$ and $s_0 > s_1 > 0$, let

$$K_n^{(p)}(s_0, s_1) = \int_0^{s_1} ds_2 \cdots \int_0^{s_{n-1}} ds_n \prod_{k=2}^n (s_{k-2} - s_k)^{-1} s_{n-1}^{-1} s_n^{-p} (1 + [(s_{n-1} - s_n)/s_n]^{-p}).$$

LEMMA 49. Let $p \in (0, 1)$ and $c_{49}(p) = 3\pi / \sin(\pi(1 - p))$:

(a) If $\phi_p(x) = x(1 + (x - 1)^{-p})^{-1} \int_0^1 (x - w)^{-1} (w^{-p} + (1 - w)^{-p}) dw$, $x > 1$, then $\sup_{x>1} \phi_p(x) \leq c_{49}(p)$.

(b) $K_n^{(p)}(s_0, s_1) \leq c_{49}(p)^{n-1} s_1^{-p} s_0^{-1} (1 + (s_0/s_1 - 1)^{-p})$, $\forall n \in \mathbb{N}^{\geq 2}$, $s_0 > s_1 > 0$.

See Appendix B for the proof.

LEMMA 50. If $\phi \in C_b(\mathbb{R}^2)$, then, $\forall T > 0$,

$$(135) \quad \lim_{\varepsilon \downarrow 0} E(\langle L^\varepsilon(T), \phi \rangle) = \int_0^T \int S_s X_0^1(x) S_s X_0^2(x) \phi(x) dx ds \in \mathbb{R}.$$

PROOF. By (134),

$$(136) \quad E({}^\varepsilon X_s^1(x) {}^\varepsilon X_s^2(x)) = {}^\varepsilon S_s {}^\varepsilon X_0^1(x) {}^\varepsilon S_s {}^\varepsilon X_0^2(x),$$

and therefore

$$(137) \quad E(\langle L^\varepsilon(T), \phi \rangle) = \int_0^T \int {}^\varepsilon S_s {}^\varepsilon X_0^1(x) {}^\varepsilon S_s {}^\varepsilon X_0^2(x) \phi(x) d^\varepsilon x ds.$$

Lemma 45(a) shows that

$$(138) \quad \begin{aligned} & \limsup_{\delta \downarrow 0} \sup_{\varepsilon > 0} \int_0^\delta \int {}^\varepsilon S_s {}^\varepsilon X_0^1(x) {}^\varepsilon S_s {}^\varepsilon X_0^2(x) |\phi(x)| d^\varepsilon x ds \\ & \leq \|\phi\|_\infty \limsup_{\delta \downarrow 0} \sup_{\varepsilon > 0} \int \left[\int_0^\delta {}^\varepsilon p_{2s}(y_1 - y_2) ds \right] {}^\varepsilon X_0^1(dy_1) {}^\varepsilon X_0^2(dy_2) = 0. \end{aligned}$$

If $\delta > 0$, then

$$(139) \quad \begin{aligned} & \lim_{\varepsilon \downarrow 0} \int_\delta^T \int {}^\varepsilon S_s {}^\varepsilon X_0^1(x) {}^\varepsilon S_s {}^\varepsilon X_0^2(x) \phi(x) d^\varepsilon x ds \\ & = \lim_{\varepsilon \downarrow 0} \int_\delta^T \int \left[\int {}^\varepsilon p_s([y_1]_\varepsilon - [x]_\varepsilon) {}^\varepsilon p_s([y_2]_\varepsilon - [x]_\varepsilon) \phi([x]_\varepsilon) dx \right] \\ & \quad \times X_0^1(dy_1) X_0^2(dy_2) ds \\ & = \int_\delta^T \int \left[\int p_s(y_1 - x) p_s(y_2 - x) \phi(x) dx \right] X_0^1(dy_1) X_0^2(dy_2) ds \\ & = \int_\delta^T \int S_s X_0^1(x) S_s X_0^2(x) \phi(x) dx ds, \end{aligned}$$

where in the next to last line we used Lemma 8 and dominated convergence. Note that the finiteness of the right-hand side of (135) is clear since $\mathbf{X}_0 \in \mathcal{M}_{f,e}$. Relations (137), (138) and (139) now easily give (135). \square

PROPOSITION 51. *If $\varepsilon_n \downarrow 0$, then $\{(\varepsilon_n X^1, \varepsilon_n X^2, L_{\varepsilon_n \mathbf{X}}^{\varepsilon_n}) : n \in \mathbb{N}\}$ is a tight sequence in $C(\mathbb{R}_+, \mathcal{M}_f(\mathbb{R}^2)^3)$.*

PROOF. Write $({}^n \mathbf{X}, L^n)$ for $(\varepsilon_n \mathbf{X}, L_{\varepsilon_n \mathbf{X}}^{\varepsilon_n})$. It suffices to show tightness of each of the three coordinates separately ([29], page 317) and to this end we specialize a result of Jakubowski [30] (see [37], Theorem II.4.1). To show a sequence of processes $\{Y^n\}$, with sample paths in $C(\mathbb{R}_+, \mathcal{M}_f(\mathbb{R}^2))$, is tight, it suffices to show the following:

(i) $\forall \varepsilon, T > 0$, there is a compact set, $K_{T,\varepsilon}$, in \mathbb{R}^2 such that

$$\sup_n P\left(\sup_{t \leq T} Y_t^n(K_{T,\varepsilon}^c) > \varepsilon\right) < \varepsilon.$$

(ii) $\forall \phi \in C_b^2(\mathbb{R}^2)$, $\{\langle Y^n, \phi \rangle : n \in \mathbb{N}\}$ is tight in $C(\mathbb{R}_+, \mathbb{R})$.

We start by proving (i) for $Y^n = L^n$. Fix $\psi : \mathbb{R}^2 \rightarrow [0, 1]$ in $C_b^2(\mathbb{R}^2)$ such that $[-1, 1]^2 \subset \{\psi = 0\} \subset \{\psi < 1\} \subset [-2, 2]^2$ and define $\psi_k(x) = \psi(xk^{-1})$. Lemma 50 implies

$$(140) \quad \lim_{n \rightarrow \infty} E(\langle L^n(T), \psi_k \rangle) = \int_0^T \int S_s X_0^1(x) S_s X_0^2(x) \psi_k(x) dx ds < \infty.$$

The right-hand side of (140) approaches 0 as $k \rightarrow \infty$ and so it follows from (140) that for any $\eta > 0$ there is a k_0 such that

$$(141) \quad \sup_n E(\langle L^n(T), \psi_{k_0} \rangle) < \eta.$$

This proves (i) for $Y^n = L^n$ by the monotonicity of $L^n(t)$ in t .

Statement (ii) for $Y^n = L^n$ would be a simple consequence of Proposition 46 and Lemma 35 if $\mathbf{X}_0 \in \mathcal{M}_{f,se}$. To handle $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ we will condition on $\mathcal{F}_\delta^\varepsilon \equiv \sigma(\varepsilon \mathbf{X}_s : s \leq \delta)$ and use the elementary equivalence between (ii) and the following:

- (ii)_a $\forall \delta > 0, \forall \phi \in C_b^2(\mathbb{R}^2)$, $\{\langle Y^n, \phi \rangle : n \in \mathbb{N}\}$ is tight in $C([\delta, \infty], \mathbb{R})$;
- (ii)_b $\forall \phi \in C_b^2(\mathbb{R}^2)$, $\{\langle Y_0^n, \phi \rangle : n \in \mathbb{N}\}$ is tight in \mathbb{R} and, $\forall \eta > 0$,

$$\limsup_{\delta \downarrow 0} \sup_n P\left(\sup_{t \leq \delta} |\langle Y_t^n, \phi \rangle - \langle Y_0^n, \phi \rangle| > \eta\right) = 0.$$

To verify (ii)_a we may choose $\delta = k_0 2^{-m_0}$ for some $k_0, m_0 \in \mathbb{N}$. For $\phi \in C_b^2(\mathbb{R}^2)$ and $m \geq m_0$, use the Markov property of ${}^n \mathbf{X}$ and Proposition 46 to see that

$$\begin{aligned}
 &P\left(\max_{k_0 2^{m-m_0} < k \leq N 2^m} \langle L^n(k 2^{-m}) - L^n((k-1) 2^{-m}), \phi \rangle > 2^{-m/8} \mid \mathcal{F}_\delta^{\varepsilon_n}\right) \\
 &\leq \sum_{k=k_0 2^{m-m_0} + 1}^{N 2^m} 2^{m/4} c_{46}(N) \bar{\mathcal{E}}_{\varepsilon_n, 1/2}({}^n \mathbf{X}_\delta) 2^{-m 3/2} (k 2^{-m} - k_0 2^{-m_0})^{-1} \|\phi\|_\infty^2 \\
 &\leq c(N, \|\phi\|_\infty) \bar{\mathcal{E}}_{\varepsilon_n, 1/2}({}^n \mathbf{X}_\delta) 2^{-m/4} \sum_{k=1}^{N 2^m} k^{-1} \\
 &\leq c(N, \|\phi\|_\infty) \bar{\mathcal{E}}_{\varepsilon_n, 1/2}({}^n \mathbf{X}_\delta) 2^{-m/4} m,
 \end{aligned}$$

which is summable over m . The standard binary expansion argument of Lévy shows that, for some $c_1 > 0$ and any $\eta, M > 0$,

$$\begin{aligned}
 &P\left(\sup_{\substack{\delta \leq s < t \leq N \\ t-s < \eta}} |\langle L^n(t), \phi \rangle - \langle L^n(s), \phi \rangle| |t-s|^{-1/8} > c_1\right) \\
 &\leq P(\mathcal{E}_{\varepsilon_n, 1/2}({}^n \mathbf{X}_\delta) > M) + P({}^n X_\delta^1(\mathbb{R}^2) + {}^n X_\delta^2(\mathbb{R}^2) > M) \\
 &\quad + c'(N, \|\phi\|_\infty) M^3 \delta_1(\eta),
 \end{aligned}$$

where $\lim_{\eta \downarrow 0} \delta_1(\eta) = 0$. Lemma 36(b) and (134)(i) allow us to choose M so that the first two terms are small, uniformly in n . Then choose η small enough to make the last term small. This proves (ii)_a; (ii)_b is immediate from (137) and (138), which imply

$$(142) \quad \limsup_{\delta \downarrow 0} \sup_n E(\langle L^n(\delta), |\phi| \rangle) = 0.$$

This proves the tightness of $\{L^n(\cdot) : n \in \mathbb{N}\}$ in $C(\mathbb{R}_+, \mathcal{M}_f(\mathbb{R}^2))$.

Next consider (i) for $Y^n = {}^n X^i$. If ψ_k is as above, then a second-order Taylor expansion shows that

$$(143) \quad |\varepsilon \Delta \psi_k(x)| \leq c_\psi k^{-2}.$$

Let $\eta > 0$. The definition of ${}^n X_0^i$ shows we may choose k_0 so that

$$(144) \quad \sup_n \langle {}^n X_0^i, \psi_k \rangle = < \eta \quad \forall k \geq k_0,$$

and (141) holds but with η^3 in place of η . Let $k \geq k_0$. Then $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon_n}$ implies

$$\sup_{t \leq T} {}^n X_t^i(\psi_k) \leq \eta + (c_\psi \sigma^2 / 2) k^{-2} \int_0^T {}^n X_s^i(\mathbb{R}^2) ds + \sup_{t \leq T} |\varepsilon^n M_t^i(\psi_k)|.$$

Therefore, for $k \geq k_0$,

$$\begin{aligned} P\left(\sup_{t \leq T} {}^n X_t^i(\psi_k) > 3\eta\right) &\leq \frac{c_\psi \sigma^2}{2\eta k^2} \int_0^T E({}^n X_s^i(\mathbb{R}^2)) ds + \eta^{-2} E\left(\sup_{t \leq T} |{}^{\varepsilon_n} M_t^i(\psi_k)|^2\right) \\ &\leq \frac{c_\psi \sigma^2 T X_0^i(\mathbb{R}^2)}{2\eta k^2} + c\gamma\eta, \end{aligned}$$

where in the last line we have used (134)(i), Burkholder’s inequality and (141) (with η^3 in place of η). Take k larger still to ensure the above bound is at most $c'\eta$, thus verifying (i) for $Y^n = {}^n X^i$.

Let $\phi \in C_b^2(\mathbb{R}^2)$ and consider (ii)_a for $Y^n = {}^n X^i$. A second order Taylor approximation shows that

$$(145) \quad \left| \frac{\sigma^{2\varepsilon_n} \Delta \phi(x)}{2} \right| \leq c\phi \quad \text{all } x, n.$$

Use the Markov property of ${}^{\varepsilon_n} \mathbf{X}$ together with $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon_n}$ and Burkholder’s inequality to see that for $\delta \leq s \leq T$,

$$\begin{aligned} (146) \quad &E(({}^n X_s^i(\mathbb{R}^2))^4 | \mathcal{F}_\delta^{\varepsilon_n})(\omega) \\ &= E(({}^n X_s^i(\mathbb{R}^2))^4 | {}^n X_0^i = {}^n X_\delta^i(\omega)) \\ &\leq c[{}^n X_\delta^i(\mathbb{R}^2)(\omega)^4 + \gamma^2 E(\langle L^n(s - \delta), 1 \rangle^2 | {}^n X_0^i = {}^n X_\delta^i(\omega))] \\ &\leq c[{}^n X_\delta^i(\mathbb{R}^2)(\omega)^4 + \gamma^2 c_{46}(T) \bar{\mathcal{E}}_{\varepsilon_n, 1/2}({}^n \mathbf{X}_\delta(\omega)) T^{1/2}], \end{aligned}$$

where Proposition 46 is used in the last line. Now use (145) and (146) in $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon_n}$ to conclude that, for $0 < \delta \leq t_1 < t_2 \leq T$,

$$\begin{aligned} &E(\langle ({}^n X_{t_2}^i, \phi) - \langle {}^n X_{t_1}^i, \phi \rangle \rangle^4 | \mathcal{F}_\delta^{\varepsilon_n}) \\ &\leq c \left[E\left(\left[\int_{t_1}^{t_2} \left\langle {}^n X_s^i, \frac{\sigma^{2\varepsilon_n} \Delta \phi}{2} \right\rangle ds \right]^4 \middle| \mathcal{F}_\delta^{\varepsilon_n}\right) \right. \\ &\quad \left. + E(\langle [{}^{\varepsilon_n} M_{t_2}^i(\phi) - {}^{\varepsilon_n} M_{t_1}^i(\phi)] \rangle^4 | \mathcal{F}_\delta^{\varepsilon_n}) \right] \\ &\leq c \left[c_\phi^4 (t_2 - t_1)^3 \int_{t_1}^{t_2} E({}^n X_s^i(\mathbb{R}^2)^4 | \mathcal{F}_\delta^{\varepsilon_n}) ds \right. \\ &\quad \left. + c\gamma^2 E(\langle (L^n(t_2), \phi) - \langle L^n(t_1), \phi \rangle \rangle^2 | \mathcal{F}_\delta^{\varepsilon_n}) \right] \\ &\leq c(T, \phi, \gamma, \sigma^2) [{}^n X_\delta^i(\mathbb{R}^2)^4 (t_2 - t_1)^3 + \bar{\mathcal{E}}_{\varepsilon_n, 1/2}({}^n \mathbf{X}_\delta)(t_2 - t_1)^{3/2} (t_2 - \delta)^{-1}], \end{aligned}$$

by Proposition 46. Lemma 36(b) and the fact that $E({}^n X_\delta^i(\mathbb{R}^2)) = X_0^i(\mathbb{R}^2)$ [from (134)(i)] show that ${}^n X_\delta^i(\mathbb{R}^2)^4 + \bar{\mathcal{E}}_{\varepsilon_n, 1/2}({}^n X_\delta^i)$ remains bounded in probability

as $n \rightarrow \infty$. We can therefore argue just as for L^n , using the above conditional L^4 bound, to see that (ii)_a holds for $Y^n = {}^nX^i$.

To check (ii)_b, let $\delta \in (0, 1]$ and use $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon_n}$ to see

$$\begin{aligned} & E\left(\sup_{t \leq \delta} (\langle {}^nX_t^i, \phi \rangle - \langle {}^nX_0^i, \phi \rangle)^2\right) \\ & \leq 2E\left(\left(\int_0^\delta \left\langle {}^nX_s^i, \frac{\sigma^2 \varepsilon_n \Delta \phi}{2} \right\rangle ds\right)^2\right) + 2E\left(\sup_{s \leq \delta} (\varepsilon_n M_s^i(\phi))^2\right) \\ & \leq 2c_\phi^2 \delta \int_0^\delta E(\langle {}^nX_s^i, 1 \rangle^2) ds + c\gamma \|\phi\|_\infty^2 E(\langle L^n(\delta), 1 \rangle) \\ & \leq 2c_\phi^2 \delta [2\langle X_0^i, 1 \rangle^2 + 2\delta E(\langle L^n(\delta), 1 \rangle)] + c\gamma \|\phi\|_\infty^2 E(\langle L^n(\delta), 1 \rangle). \end{aligned}$$

The above bound converges to zero uniformly in n as $\delta \downarrow 0$ by (142). Since $\langle {}^nX_0^i, \phi \rangle \rightarrow \langle X_0^i, \phi \rangle$ as $n \rightarrow \infty$, (ii)_b follows for $Y^n = {}^nX^i$ and we are done. \square

To show the limit points obtained from Proposition 51 solve $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ we first introduce some notation:

$$\begin{aligned} \Omega_{X,L} &= C(\mathbb{R}_+, \mathcal{M}_f(\mathbb{R}^2)^3) \text{ with its Borel sigma-field } \mathcal{F}_{X,L} \\ & \text{and canonical right-continuous filtration } (\mathcal{F}_t^{X,L}). \end{aligned}$$

Let $(\mathbf{X}, L) = (X^1, X^2, L)$ denote the coordinate maps on $\Omega_{X,L}$.

PROPOSITION 52. *Let P be a weak limit point of the laws of $\{(\varepsilon_n \mathbf{X}, L_{\varepsilon_n \mathbf{X}}) : n \in \mathbb{N}\}$ on $(\Omega_{X,L}, \mathcal{F}_{X,L})$, as $\varepsilon_n \downarrow 0$. Let \mathcal{F} and \mathcal{F}_t be the P -completions of $\mathcal{F}_{X,L}$ and $\mathcal{F}_t^{X,L}$, respectively. Then \mathbf{X} solves $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma^2, \gamma}$ on $(\Omega_{X,L}, \mathcal{F}, \mathcal{F}_t, P)$ and $L = L_{\mathbf{X}}$ is the collision local time of \mathbf{X} P -a.s. Moreover,*

$$(147) \quad \langle L_{\mathbf{X}}^{*, \delta}(t), \phi \rangle \rightarrow \langle L_{\mathbf{X}}(t), \phi \rangle \quad \text{in } L^1(P) \text{ as } \delta \downarrow 0, \forall \phi \in C_b(\mathbb{R}^2),$$

where $L_{\mathbf{X}}^{*, \delta}$ was defined in (10).

PROOF. By Skorohod's theorem we may assume that, on some $(\Omega', \mathcal{F}', P')$,

$$(148) \quad ({}^n \mathbf{X}, L^n) \equiv (\varepsilon_n \mathbf{X}, L_{\varepsilon_n \mathbf{X}}) \xrightarrow{\text{a.s.}} (\mathbf{X}, L) \quad \text{in } \Omega_{X,L}, \varepsilon_n \downarrow 0.$$

Let \mathcal{F}'_t (respectively, $\mathcal{F}'_t{}^n$) be the right-continuous P' -complete filtration generated by (\mathbf{X}, L) (respectively, ${}^n \mathbf{X}$) and let $\phi \in C_b^2(\mathbb{R}^2)$. An elementary argument shows that

$$(149) \quad \varepsilon_n \Delta \phi([x]_{\varepsilon_n}) \rightarrow \Delta \phi(x) \quad \text{boundedly and uniformly on compacts.}$$

From $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma, \varepsilon_n}$ we have

$$\begin{aligned} \langle {}^n X_t^i, \phi \rangle &= \langle {}^n X_0^i, \phi \rangle + \int_0^t \left\langle {}^n X_s^i, \frac{\sigma^2 \varepsilon_n \Delta \phi}{2} \right\rangle ds + \varepsilon_n M_t^i(\phi), \quad t \geq 0, \\ (150) \quad \varepsilon_n M_t^i(\phi) &\text{ is a continuous } L^2(\mathcal{F}_t^n)\text{-martingale, } i = 1, 2, \text{ and} \\ \langle \varepsilon_n M^i(\phi), \varepsilon_n M^j(\phi) \rangle_t &= \delta_{ij} \gamma \langle L^n(t), \phi^2 \rangle. \end{aligned}$$

As each of the first three terms in (150) converges a.s. in $C(\mathbb{R}_+, \mathbb{R})$, we see that $\varepsilon_n M_t^i(\phi) \rightarrow M_t^i(\phi)$ a.s. in $C(\mathbb{R}_+, \mathbb{R})$ for some \mathcal{F}_t' -adapted continuous process $M_t^i(\phi)$. Lemma 50, (150) and Burkholder's inequality imply that $\{\sup_{t \leq T} |\varepsilon_n M_t^i(\phi)| : n \in \mathbb{N}\}$ is L^2 -bounded for each $T > 0$. It follows easily that $M_t^i(\phi)$ is a continuous $L^2(\mathcal{F}_t')$ -martingale. Theorem VI.6.1(b) of [29] implies that $\langle M^i(\phi), M^j(\phi) \rangle_t = \gamma \delta_{ij} \langle L(t), \phi^2 \rangle \forall t \geq 0$ a.s. We may now let $n \rightarrow \infty$ in (150) to see that, for $\phi \in C_b^2(\mathbb{R}^2)$,

$$\begin{aligned} \langle X_t^i, \phi \rangle &= \langle X_0^i, \phi \rangle + \int_0^t \left\langle X_s^i, \frac{\sigma^2 \Delta \phi}{2} \right\rangle ds + M_t^i(\phi), \quad i = 1, 2, \\ (151) \quad M_t^i(\phi) &\text{ is a continuous } L^2(\mathcal{F}_t')\text{-martingale such that} \\ \langle M^i(\phi), M^j(\phi) \rangle_t &= \delta_{ij} \gamma \langle L(t), \phi^2 \rangle \quad \forall t \geq 0 \text{ a.s.} \end{aligned}$$

By polarization the last equality implies $\langle M^1(\phi_1), M^2(\phi_2) \rangle = 0$ a.s. for all $\phi_1, \phi_2 \in C_b^2(\mathbb{R}^2)$.

To show \mathbf{X} satisfies $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ it remains to prove $L = L_{\mathbf{X}}$, P' -a.s. The Markov property of ${}^n \mathbf{X}$ and (137) imply, for $0 \leq s < t$ and $\phi \in C_b(\mathbb{R}^2)$,

$$\begin{aligned} (152) \quad E(\langle L^n(t) - L^n(s), \phi \rangle \mid \mathcal{F}_s^n) &= \int_0^{t-s} \left[\int \varepsilon_n S_r {}^n X_s^1(x) \varepsilon_n S_r {}^n X_s^2(x) \phi(x) d^{\varepsilon_n} x \right] dr \quad \text{a.s.} \end{aligned}$$

For each $r > 0$, it is straightforward to use Lemma 8, (148) and a dominated convergence argument to see that

$$\begin{aligned} (153) \quad \lim_{n \rightarrow \infty} \int \varepsilon_n S_r {}^n X_s^1(x) \varepsilon_n S_r {}^n X_s^2(x) \phi(x) d^{\varepsilon_n} x &= \int S_r X_s^1(x) S_r X_s^2(x) \phi(x) dx \\ &< \infty \quad \text{a.s. } \forall r > 0. \end{aligned}$$

To take an L^1 limit on the right-hand side of (152) we first show

$$\begin{aligned} (154) \quad f_n(r) &= \int \varepsilon_n S_r {}^n X_s^1(x) \varepsilon_n S_r {}^n X_s^2(x) d^{\varepsilon_n} x \\ &\text{ is a uniformly integrable sequence on } ([0, t-s] \times \Omega', dr \times P'). \end{aligned}$$

If $s = 0$, this is an easy consequence of Lemma 45(a), so assume $s > 0$. We see from (153) that

$$(155) \quad \lim_{n \rightarrow \infty} f_n(r) = \int S_r X_s^1(x) S_r X_s^2(x) dx \equiv f(r) \quad \text{a.s. } \forall r > 0.$$

Now use (151) just as in the proof of Proposition 15(b) [more specifically, (58)] to conclude that

$$(156) \quad E'(S_r X_s^1(x) S_r X_s^2(x)) = S_{r+s} X_0^1(x) S_{r+s} X_0^2(x) \quad \forall r > 0, x \in \mathbb{R}^2.$$

From (134) we have

$$\begin{aligned} \lim_{n \rightarrow \infty} E' \left(\int_0^{t-s} f_n(r) dr \right) &= \lim_{n \rightarrow \infty} \int_0^{t-s} \int \varepsilon_n S_{r+s} X_0^1(x) \varepsilon_n S_{r+s} X_0^2(x) d^{\varepsilon_n} x dr \\ &= \int_0^{t-s} \int S_{r+s} X_0^1(x) S_{r+s} X_0^2(x) dx dr \end{aligned}$$

by dominated convergence and $s > 0$, as for (139). This with (155) and (156) shows that $\lim_{n \rightarrow \infty} E'(\int_0^{t-s} f_n(r) dr) = E'(\int_0^{t-s} f(r) dr)$, which, together with (155), gives (154). Therefore the same uniform integrability holds for

$$\left\{ \int \varepsilon_n S_r X_s^1(x) \varepsilon_n S_r X_s^2(x) \phi(x) d^{\varepsilon_n} x : n \in \mathbb{N} \right\}.$$

Use this and (153) to let $n \rightarrow \infty$ in the right-hand side of (152) and conclude

$$(157) \quad \begin{aligned} &E'(\langle L^n(t) - L^n(s), \phi \rangle | \mathcal{F}_s^n) \\ &\xrightarrow{L^1} \int_0^{t-s} \int S_r X_s^1(x) S_r X_s^2(x) \phi(x) dx dr \quad (\text{as } n \rightarrow \infty) \\ &\equiv \Phi(\mathbf{X}_s). \end{aligned}$$

Now let $\psi : \Omega_{X,L} \rightarrow \mathbb{R}$ be bounded continuous and $\mathcal{F}_s^{X,L}$ -measurable and satisfy $\psi(\mathbf{X}, L) = 0$ if $X_s^1(\mathbb{R}^2) + X_s^2(\mathbb{R}^2) > K$ for some $K > 0$. If $J > 0$, then (157) implies

$$(158) \quad \begin{aligned} &E'(\Phi(\mathbf{X}_s) \psi(\mathbf{X}, L)) \\ &= \lim_{n \rightarrow \infty} E'(E'(\langle L^n(t) - L^n(s), \phi \rangle \\ &\quad \times \mathbf{1}(|\langle L^n(t) - L^n(s), \phi \rangle| > J) | \mathcal{F}_s^n) \psi(\mathbf{X}, L^n)) \\ &\quad + E'(E'(\langle L^n(t) - L^n(s), \phi \rangle \mathbf{1}(|\langle L^n(t) - L^n(s), \phi \rangle| \leq J) | \mathcal{F}_s^n) \psi(\mathbf{X}, L^n)) \\ &= \lim_{n \rightarrow \infty} T_n^{(1)} + T_n^{(2)}. \end{aligned}$$

Choose J so that $P'(|\langle L^n(t) - L^n(s), \phi \rangle| = J) = 0$. By Proposition 46, the Markov property of ${}^n\mathbf{X}$ and our assumption on the support of ψ , if $s > 0$, then

$$\begin{aligned} |T_n^{(1)}| &\leq J^{-1} E'(E'(\langle L^n(t) - L^n(s), \phi \rangle^2 | \mathcal{F}_s^n) \psi({}^n\mathbf{X}, L^n)) \\ (159) \quad &\leq J^{-1} c_{46}(t) t^{1/2} \|\phi\|_\infty^2 K^2 E'(\mathcal{E}_{\varepsilon_n, 1/2}({}^n\mathbf{X}_s)) \|\psi\|_\infty \\ &\leq c(t, \phi, \psi) c_{36}(1 + s^{-1}) K^4 J^{-1}, \end{aligned}$$

the last by Lemma 36(b). Next use (148), our choice of J and dominated convergence to see that

$$\begin{aligned} (160) \quad \lim_{n \rightarrow \infty} T_n^{(2)} &= E'(\langle L(t) - L(s), \phi \rangle \mathbf{1}(|\langle L(t) - L(s), \phi \rangle| \leq J) \psi(\mathbf{X}, L)) \\ &\rightarrow E'(\langle L(t) - L(s), \phi \rangle \psi(\mathbf{X}, L)) \quad \text{as } J \rightarrow \infty. \end{aligned}$$

The last line is clear from $E'(\langle L(t), 1 \rangle) < \infty$ [by (151)]. Use (159) and (160) in (158) and then let $J \rightarrow \infty$ to conclude

$$E'(\Phi(\mathbf{X}_s) \psi(\mathbf{X}, L)) = E'(\langle L(t) - L(s), \phi \rangle \psi(\mathbf{X}, L)), \quad t > s > 0.$$

It follows that, for $\phi \in C_b(\mathbb{R}^2)$,

$$E'(\langle L(t) - L(s), \phi \rangle | \mathcal{F}_s') = \int_0^{t-s} \int S_r X_s^1(x) S_r X_s^2(x) \phi(x) dx dr \quad \text{a.s. } \forall s > 0,$$

and therefore, by the definition of $L_{\mathbf{X}}^{*,\delta}(t)$, that

$$(161) \quad \langle L_{\mathbf{X}}^{*,\delta}(t), \phi \rangle = \int_0^t \frac{1}{\delta} E'(\langle L(s + \delta) - L(s), \phi \rangle | \mathcal{F}_s') ds, \quad P'\text{-a.s.}$$

Theorem 37 on page 126 of [32] and the continuity and integrability of $\langle L(t), \phi \rangle$ yield that the right-hand side of (161) converges in $L^1(P')$ to $\langle L(t), \phi \rangle$ as $\delta \downarrow 0$ for each $t \geq 0$ and $\phi \in C_b(\mathbb{R}^2)$. Therefore $L_{\mathbf{X}}$ exists and equals L a.s., and (147) holds on $(\Omega', \mathcal{F}', P')$. It is now trivial to transfer these results over to the canonical space in Proposition 52. \square

PROOF OF THEOREM 11. Parts (a) and (c) are immediate from Propositions 51, 52 and 25(b), except for the verification of **(IntC)** and **(SIntC)**, the latter for $\mathbf{X}_0 \in \mathcal{M}_{f,se}$. These are derived in [12] using the moment dual process from Section 3 and, more specifically, Theorems 53, 54 and Remark 55 below.

Part (b) is proved in [12].

The first assertion of (d) follows by a direct change of variables calculation in $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma,\gamma}$ and the second assertion then follows from (b). \square

Let $P_{\mathbf{X}_0}$ denote the law on $\Omega_o = C(\mathbb{R}_+, \mathcal{M}_f^2(\mathbb{R}^2))$ of the process \mathbf{X} constructed in Proposition 52.

THEOREM 53. *Assume (33) and $\mathbf{X}_0 \in \mathcal{M}_{f,e}$. For any $0 < \delta < t$, any bounded continuous $\phi_0: (\mathbb{R}^2)^4 \rightarrow \mathbb{R}_+$, $I_0 \subset \{1, \dots, 4\}$, and any Borel map $\psi: \mathcal{M}_f^2(\mathbb{R}^2) \rightarrow \mathbb{R}_+$,*

$$\begin{aligned}
 & E_{\mathbf{X}_0} \left(\int \phi_0(x_1, \dots, x_4) \prod_{i \in I_0} X_t^1(dx_i) \prod_{j \notin I_0} X_t^2(dx_j) \psi(\mathbf{X}_\delta) \right) \\
 (162) \quad & \leq \hat{E}_{\phi_0, I_0} \times E_{\mathbf{X}_0} \left(\int \phi_{t-\delta}(x_1, \dots, x_4) \prod_{i \in I_{t-\delta}} X_\delta^1(dx_i) \prod_{j \notin I_{t-\delta}} X_\delta^2(dx_j) \psi(\mathbf{X}_\delta) \right. \\
 & \quad \left. \times \exp \left\{ \gamma \int_0^{t-\delta} \binom{|I_s|}{2} + \binom{|I_s^c|}{2} ds \right\} \right).
 \end{aligned}$$

If, in addition, ψ is bounded and

$$(163) \quad \{\psi \neq 0\} \subset \{(\mu^1, \mu^2) : \mu^1(\mathbb{R}^2) + \mu^2(\mathbb{R}^2) \leq K\} \quad \text{for some } K > 0,$$

then the above expressions are both finite.

PROOF. If $\delta > 0$, ψ is bounded, continuous, and satisfies (163), then both the above results are immediate from Theorems 32(b) and 11(c) and Fatou’s lemma. By taking bounded pointwise limits in ψ , these results extend to bounded nonnegative Borel ψ satisfying (163). Next, use monotone convergence to get the first inequality for all nonnegative Borel ψ and $\delta > 0$. \square

THEOREM 54. *Assume (33) and $\mathbf{X}_0 \in \mathcal{M}_{f,se}$. For any $t > 0$, any bounded continuous $\phi_0: (\mathbb{R}^2)^4 \rightarrow \mathbb{R}_+$, any $I_0 \subset \{1, \dots, 4\}$ and any Borel map $\psi: \mathcal{M}_f(\mathbb{R}^2)^2 \rightarrow \mathbb{R}_+$,*

$$\begin{aligned}
 & E_{\mathbf{X}_0} \left(\int \phi_0(x_1, \dots, x_4) \prod_{i \in I_0} X_t^1(dx_i) \prod_{j \notin I_0} X_t^2(dx_j) \psi(\mathbf{X}_0) \right) \\
 & \leq \hat{E}_{\phi_0, I_0} \left(\int \phi_t(x_1, \dots, x_4) \prod_{i \in I_t} X_0^1(dx_i) \prod_{j \notin I_t} X_0^2(dx_j) \psi(\mathbf{X}_0) \right. \\
 & \quad \left. \times \exp \left\{ \gamma \int_0^t \binom{|I_s|}{2} + \binom{|I_s^c|}{2} ds \right\} \right) < \infty.
 \end{aligned}$$

In particular,

$$E_{\mathbf{X}_0} \left(\sup_{t \leq T} X_t^1(\mathbb{R}^2)^4 + X_t^2(\mathbb{R}^2)^4 \right) < \infty \quad \text{for all } T > 0.$$

PROOF. The first two inequalities are proved as in Theorem 53 but using Theorem 32(a) instead of Theorem 32(b) [the proof is simpler as the $\psi(\mathbf{X}_0)$ term is deterministic and hence trivial to include]. The last result is obtained by taking $\phi_0 = 1$, $I_0 = \emptyset$ or $I_0^c = \emptyset$, and using the L^4 maximal inequality for martingales. \square

REMARK 55. We will use Theorem 53 in [12] to show the solution constructed in Theorem 11(a),(c) satisfies (IntC). Note that, without any uniqueness result, the above proof and Propositions 51 and 52 show that any weak limit point of $\{\varepsilon_n \mathbf{X}\}$ satisfies $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ and the conclusions of Theorems 53 and 54.

We complete this section with the proof of our key L^2 estimate.

PROOF OF PROPOSITION 46. Clearly it suffices to consider the case $\phi \equiv 1$. Let $(\mathcal{F}_t^\varepsilon)$ denote the right continuous filtration generated by ${}^\varepsilon \mathbf{X}$. Use the Markov property of ${}^\varepsilon \mathbf{X}$ and (134)(ii) to see that, for T fixed and $0 \leq t_1 < t_2 \leq T$,

$$\begin{aligned} & E(\langle L^\varepsilon(t_2) - L^\varepsilon(t_1), 1 \rangle^2) \\ &= 2 \int_{t_1}^{t_2} dr \int_{t_1}^r dt \int d^\varepsilon y \int d^\varepsilon x E(E({}^\varepsilon X_r^1(x) {}^\varepsilon X_r^2(x) | \mathcal{F}_t^\varepsilon) {}^\varepsilon X_t^1(y) {}^\varepsilon X_t^2(y)) \\ &= 2 \int_{t_1}^{t_2} dr \int_{t_1}^r dt \int d^\varepsilon y \int d^\varepsilon x E({}^\varepsilon S_{r-t} {}^\varepsilon X_t^1(x) {}^\varepsilon S_{r-t} {}^\varepsilon X_t^2(x) {}^\varepsilon X_t^1(y) {}^\varepsilon X_t^2(y)) \\ &= 2 \int_{t_1}^{t_2} dr \int_{t_1}^r dt E\left(\int {}^\varepsilon p_{2(r-t)}(y_1 - y_2) {}^\varepsilon p_0(y_3 - y_4) \right. \\ &\quad \left. \times {}^\varepsilon X_t^1(dy_1) {}^\varepsilon X_t^2(dy_2) {}^\varepsilon X_t^1(dy_3) {}^\varepsilon X_t^2(dy_4)\right). \end{aligned}$$

Let $\phi_0^{\varepsilon, v}(y) = {}^\varepsilon p_v(y_1 - y_2) {}^\varepsilon p_0(y_3 - y_4)$ ($v \geq 0$) and let $(\phi_t^{\varepsilon, v}, I_t)$ denote the moment dual process in Proposition 28 starting at $(\phi_0^{\varepsilon, v}, I_0 = \{1, 3\})$. Then a simple change of variables in the above, together with Proposition 28, implies

$$\begin{aligned} & E(\langle L^\varepsilon(t_2) - L^\varepsilon(t_1), 1 \rangle^2) \\ (164) \quad &= \int_{t_1}^{t_2} dt \int_0^{2(t_2-t)} dv E(F(\phi_0^{\varepsilon, v}, I_0, {}^\varepsilon \mathbf{X}_t)) \\ &\leq e^{3\gamma T} \int_{t_1}^{t_2} dt \int_0^{2(t_2-t)} dv \hat{E}_{\phi_0^{\varepsilon, v}, I_0}^\varepsilon(F(\phi_t^{\varepsilon, v}, I_t, {}^\varepsilon \mathbf{X}_0)). \end{aligned}$$

To bound the expectation on the right-hand side of (164) we will use Lemma 38. Note first that

$$\begin{aligned} & \phi_t^{\varepsilon, v}(y) = {}^\varepsilon p_{v+2t}(y_1 - y_2) {}^\varepsilon p_{2t}(y_3 - y_4) \quad \text{for } T_0 \leq t < T_1, \\ (165) \quad & \phi_{T_1-}^{\varepsilon, v}(y) = {}^\varepsilon p_{v+2T_1}(y_1 - y_2) {}^\varepsilon p_{2T_1}(y_3 - y_4). \end{aligned}$$

We now will verify (115) for $n_0 = 1$. Suppose i_1 switches via i_2 , where $\{i_1, i_2\} = \{2, 4\}$ are distinct random indices. Then $I_{T_1} = \{1, 3, i_1\}$ and

$$\phi_{T_1}^{\varepsilon, v}(y) = \varepsilon p_{v+2T_1}(y_1 - y_{i_2}) \varepsilon p_{2T_1}(y_3 - y_{i_2}) \varepsilon p_0(y_{i_1} - y_{i_2}).$$

It follows that, for $T_1 \leq t < T_2$,

$$\begin{aligned} \phi_t^{\varepsilon, v}(y) &= \int \varepsilon p_{v+2T_1+t-T_1}(y_1 - z_{i_2}) \varepsilon p_{2T_1+t-T_1}(y_3 - z_{i_2}) \\ &\quad \times \varepsilon p_{t-T_1}(z_{i_2} - y_{i_1}) \varepsilon p_{t-T_1}(z_{i_2} - y_{i_2}) d^\varepsilon z_{i_2} \\ &\leq \left(\frac{c_{37}}{v + T_1 + t} \right) \left(\frac{c_{37}}{T_1 + t} \right) \varepsilon p_{2(t-T_1)}(y_{i_1} - y_{i_2}) \end{aligned}$$

and $i_1 \in I_t, i_2 \notin I_t$. A similar result holds if 1 switches via 3, or conversely, at T_{n_0} . This establishes (115) with

$$(166) \quad f(t, \omega) = c_{37}^2 (v + t)^{-1} t^{-1}.$$

Then, according to the definition in Lemma 38, after some algebra

$$\begin{aligned} \rho(s) \equiv \rho_1^f(s) &= \begin{cases} c_{37}^{n+1} \prod_{k=1}^{n-1} (U_k + U_{k+1})^{-1} (v + T_2)^{-1} (s - T_{n-1})^{-1}, & T_n \leq s < T_{n+1}, n \geq 2, \\ c_{37}^2 (v + s)^{-1} s^{-1}, & T_1 \leq s < T_2, \end{cases} \\ &= c_{37}^{n+1} \prod_{k=1}^{n-1} (U_k + U_{k+1})^{-1} (v + T_2 \wedge s)^{-1} (s - T_{n-1})^{-1}, \\ &\quad T_n \leq s < T_{n+1}, n \geq 1. \end{aligned}$$

Extend $\rho(s)$ to $[0, T_1]$ by defining

$$\rho(s) = c_{37} (v + s)^{-1} \quad \text{if } 0 \leq s < T_1.$$

Then Lemma 38, (114) and (165) imply there are random indices $\{i_1^n, i_2^n\} \subset \{1, 2, 3, 4\}$, such that

$$\begin{aligned} \phi_s^{\varepsilon, v}(y) &\leq \rho(s) \varepsilon p_{2(s-T_n)}(y_{i_1^n} - y_{i_2^n}), \\ i_1^n &\in I_s, i_2^n \in I_s^c, T_n \leq s < T_{n+1}, n \geq 0, \hat{P}_{\phi_0^{\varepsilon, v}, I_0} \text{-a.s.} \end{aligned}$$

Therefore, if \hat{E}^ε denotes $\hat{E}_{\phi_0^{\varepsilon, v}, I_0}^\varepsilon$ and $N(t) = n$ iff $T_n \leq t < T_{n+1}$, then

$$\begin{aligned} &\hat{E}^\varepsilon(F(\phi_t^{\varepsilon, v}, I_t, \varepsilon \mathbf{X}_0)) \\ (167) \quad &\leq \hat{E}^\varepsilon \left(\rho(t) \int \varepsilon p_{2(t-T_{N(t)})}(y_1 - y_2) \varepsilon X_0^1(dy_1) \varepsilon X_0^2(dy_2) \varepsilon X_0^1(\mathbb{R}^2) \varepsilon X_0^2(\mathbb{R}^2) \right) \\ &\leq \bar{\varepsilon}_{\varepsilon, 1/2}(\varepsilon \mathbf{X}_0) \hat{E}^\varepsilon(\rho(t)(t - T_{N(t)})^{-1/2}). \end{aligned}$$

Let α_n be the rate of the exponential time U_n . Then $\alpha_{2n} = 3\gamma, \alpha_{2n+1} = 2\gamma$. The definition of $\rho(t)$ gives

$$\begin{aligned} & \hat{E}^\varepsilon(\rho(t)(t - T_{N(t)})^{-1/2}) \\ &= \hat{E}^\varepsilon(1(t \in [0, T_1])c_{37}(v + t)^{-1}t^{-1/2}) \\ & \quad + \sum_{n=1}^\infty \hat{E}^\varepsilon\left(\mathbf{1}(T_n \leq t < T_{n+1})c_{37}^{n+1} \right. \\ & \quad \quad \times \prod_{k=1}^{n-1} (U_k + U_{k+1})^{-1}(v + T_2 \wedge t)^{-1}(t - T_{n-1})^{-1}(t - T_n)^{-1/2}\left. \right) \\ & \leq c_{37}(v + t)^{-1}t^{-1/2} \\ (168) \quad & + \sum_{n=1}^\infty c_{37}^{n+1} \prod_{i=1}^n \alpha_i \int_{\mathbb{R}_+^n} \mathbf{1}\left(\sum_1^n u_i \leq t\right) e^{-\alpha_{n+1}(t - \sum_1^n u_i)} e^{-\sum_1^n \alpha_i u_i} \\ & \quad \times \prod_{i=1}^{n-1} (u_i + u_{i+1})^{-1}(v + (u_1 + u_2) \wedge t)^{-1} \\ & \quad \times \left(t - \sum_1^{n-1} u_i\right)^{-1} \left(t - \sum_1^n u_i\right)^{-1/2} du_1 \cdots du_n \\ & \leq c_{37}(v + t)^{-1}t^{-1/2} + c_{37}e^{3\gamma t} \sum_{n=1}^\infty (c_{37}\sqrt{6\gamma})^n I_n(t, v), \end{aligned}$$

where

$$\begin{aligned} I_n(t, v) &= \int_{\mathbb{R}_+^n} \mathbf{1}\left(\sum_1^n u_i \leq t\right) \prod_{i=1}^{n-1} (u_i + u_{i+1})^{-1}(v + (u_1 + u_2) \wedge t)^{-1} \\ & \quad \times \left(t - \sum_1^{n-1} u_i\right)^{-1} \left(t - \sum_1^n u_i\right)^{-1/2} du_1 \cdots du_n. \end{aligned}$$

Let $s_j = t - \sum_{i=1}^j u_i, j = 0, 1, \dots, n$, and recall the notation $K_n^{(p)}(s_0, s)$ in Lemma 49. Then, for $n \geq 1$,

$$\begin{aligned} I_n(t, v) &= \int_{\mathbb{R}_+^n} \mathbf{1}(s_n \leq s_{n-1} \leq \cdots \leq s_1 \leq s_0 = t) \\ & \quad \times \prod_{i=1}^n (s_{i-2} - s_i)^{-1}(v + (s_0 - s_1))^{-1} s_{n-1}^{-1} s_n^{-1/2} ds_1 \cdots ds_n \end{aligned}$$

$$\begin{aligned} &\leq K_{n+1}^{(1/2)}(v + s_0, s_0) \\ &\leq (3\pi)^n t^{-1/2}(v + t)^{-1}(1 + (v/t)^{-1/2}), \end{aligned}$$

the last by Lemma 49(b). Our hypothesis (33) on $\gamma\sigma^{-2}$ implies $c_{37}3\pi\sqrt{6}\gamma < 1$ so we may use the above bound in (168) to conclude that, for $t \leq T$,

$$\begin{aligned} &\hat{E}^\varepsilon(\rho(t)(t - T_{N(t)})^{-1/2}) \\ &\leq c_{37}(v + t)^{-1}t^{-1/2} \\ &\quad + c_{37}e^{3\gamma t}(1 - c_{37}3\pi\sqrt{6}\gamma)^{-1}(v + t)^{-1}t^{-1/2}(1 + (v/t)^{-1/2}) \\ &\leq c_1(\gamma, \sigma^2, T)(v + t)^{-1}t^{-1/2}(1 + (v/t)^{-1/2}). \end{aligned}$$

Employing this bound in (167) and (164), we get (for $0 \leq t_1 < t_2 \leq T$)

$$\begin{aligned} &E((L^\varepsilon(t_2) - L^\varepsilon(t_1), 1)^2) \\ (169) \quad &\leq c_2(\gamma, \sigma^2, T)\bar{\mathcal{E}}_{\varepsilon,1/2}(\varepsilon \mathbf{X}_0) \int_{t_1}^{t_2} dt \int_0^{2(t_2-t)} dv (v + t)^{-1}(t^{-1/2} + v^{-1/2}) \\ &\equiv c_2\bar{\mathcal{E}}_{\varepsilon,1/2}(\varepsilon \mathbf{X}_0)I(t_1, t_2). \end{aligned}$$

Substitute $u = v/t$ for v to see that

$$\begin{aligned} I(t_1, t_2) &= \int_{t_1}^{t_2} dt t^{-1/2} \int_0^{2(t_2/t-1)} du \left(\frac{1 + u^{-1/2}}{1 + u} \right) \\ &\leq c_3 \int_{t_1}^{t_2} t^{-1/2} \left(\left(\frac{t_2}{t} - 1 \right) \wedge 1 \right)^{1/2} dt \\ &= c_3 \int_{t_1 \vee (t_2/2)}^{t_2} t^{-1}(t_2 - t)^{1/2} dt + c_3 \int_{t_1}^{t_1 \vee (t_2/2)} t^{-1/2} dt \\ &\leq \frac{2c_3}{t_2} \int_{t_1}^{t_2} (t_2 - t)^{1/2} dt + c_3 \mathbf{1}(t_2 > 2t_1)2(t_2/2)^{1/2} \\ &\leq \frac{c_4}{t_2}((t_2 - t_1)^{3/2} + \mathbf{1}(t_2 > 2t_1)t_2^{3/2}) \\ &\leq \frac{c_5}{t_2}(t_2 - t_1)^{3/2}, \end{aligned}$$

where we use $t_2 - t_1 > t_2/2$ if $t_2 > 2t_1$ in the last line. Use this in (169) to complete the proof. \square

5. Long-term behavior. In this section we prove Theorem 21. Recall this gives the limiting law of $(X_t^1(\mathbb{R}^2), X_t^2(\mathbb{R}^2))$ as $t \rightarrow \infty$. We will adapt the proof of the corresponding result for the lattice case (Theorem 1.2(b) from [14]). Assume X_0 is a fixed initial state in $\mathcal{M}_{f,e}$ and (33) holds throughout this section. The

following third moment bound is simpler than the fourth moment bounds in Section 4 but we include a proof for completeness.

Recall the notation $\mathcal{E}_p(\mathbf{X}_0)$ introduced prior to Lemma 35. We set

$$\bar{\mathcal{E}}_p(\mathbf{X}_0) = \mathcal{E}_p(\mathbf{X}_0)[\langle X_0^1, 1 \rangle + \langle X_0^2, 1 \rangle].$$

For those keeping track, in this particular argument (33) could be weakened to $\gamma/\sigma^2 < (c_{rw}\pi)^{-1}$.

LEMMA 56. Assume $X_0 \in \mathcal{M}_{f,se}$. For any $p' \in (0, 1/2)$ there is a $c_{56} = c_{56}(\gamma, \sigma, p')$ so that the law $P_{\mathbf{X}_0}$ in Theorem 11 satisfies

$$E_{X_0} \int_0^T \int \mathbb{P}_s(x_1, x_2) X_r^1(dx_1) L_{\mathbf{X}}(d[r, x_2]) \leq c_{56} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) s^{-1/2} < \infty \quad \forall T > 0.$$

PROOF. Fix $s > 0$. Let ${}^\varepsilon X^i$ and $L^\varepsilon = {}^\varepsilon L_{\varepsilon \mathbf{X}}$ denote our usual rescalings of the process and its collision local time on $\varepsilon \mathbb{Z}^2$. An application of Fatou's lemma, Theorem 11(c), Skorohod's a.s. representation and Lemma 8 show that it suffices to prove that, for all $\varepsilon > 0$ sufficiently small,

$$E \int_0^T \int \varepsilon \mathbb{P}_s(x_1, x_2) {}^\varepsilon X_r^1(dx_1) L^\varepsilon[d(r, x_2)] \leq c_{56} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) s^{-1/2}.$$

We calculate the left-hand side using the moment dual process in Proposition 28 with $p = 3$.

Let $T_n = U_1 + \dots + U_n$ ($T_0 = 0$) be the jump times of the moment dual process $(\phi_t(x_1, x_2, x_3), I_t)$ for third order moments with

$$\phi_0(x_1, x_2, x_3) = \varepsilon \mathbb{P}_s(x_1, x_2) \varepsilon \mathbb{P}_0(x_2, x_3) \quad \text{and} \quad I_0 = \{1, 2\}.$$

Then $\{U_i\}$ are i.i.d. exponential with rate γ and Proposition 28 gives

$$\begin{aligned} (170) \quad & E \left(\int \phi_0(x_1, x_2, x_3) {}^\varepsilon X_r^1(dx_1) {}^\varepsilon X_r^1(dx_2) {}^\varepsilon X_r^2(dx_3) \right) \\ & = \hat{E}_{\mathbf{V}_0^\varepsilon}^\varepsilon \left(e^{\gamma r} \int \phi_r(x_1, x_2, x_3) \prod_{i \in I_r} {}^\varepsilon X_0^1(dx_i) \prod_{j \in I_r^c} {}^\varepsilon X_0^2(dx_j) \right). \end{aligned}$$

Recall from Lemma 8 that $\varepsilon \mathbb{P}_r \leq c_{rw} \sigma^{-2} r^{-1} \equiv c_1 r^{-1}$. We claim that, setting $U_0 \equiv s$ for all $n \in \mathbb{Z}_+$,

(171) $T_n \leq r < T_{n+1}$ implies

$$\begin{aligned} (170_n) \quad & \phi_r(x_1, x_2, x_3) \\ & \leq (c_1)^{n+1} \prod_{\ell=0}^{n-1} (U_\ell + U_{\ell+1})^{-1} (U_n + r - T_n)^{-1} \varepsilon \mathbb{P}_{2(r-T_n)}(x_i, x_j) \quad \text{and} \\ & I_r = \{i, k\} \quad \text{or} \quad I_r = \{i\} \quad (i, j, k \text{ distinct random indices}). \end{aligned}$$

Assume (170_n) with, say, $I_{T_{n+1}-} = \{i, k\}$ (a similar argument goes through if $I_{T_{n+1}-} = \{i\}$). Then, if k changes type at T_{n+1} ,

$$\begin{aligned} & \phi_{T_{n+1}}(x_1, x_2, x_3) \\ & \leq c_1^{n+1} \prod_{\ell=0}^n (U_\ell + U_{\ell+1})^{-1} \varepsilon p_{2U_{n+1}}(x_i, x_j) \varepsilon p_0(x_i, x_k), \quad I_{T_{n+1}} = \{i\}. \end{aligned}$$

Therefore if $T_{n+1} \leq r < T_{n+2}$,

$$\begin{aligned} & \phi_r(x_1, x_2, x_3) \\ & \leq c_1^{n+1} \prod_{\ell=0}^n (U_\ell + U_{\ell+1})^{-1} \\ & \quad \times \int \varepsilon p_{2U_{n+1}+(r-T_{n+1})}(x_j, y_i) \varepsilon p_{r-T_{n+1}}(x_k, y_i) \varepsilon p_{r-T_{n+1}}(x_i, y_i) d^\varepsilon y_i \\ & \leq c_1^{n+2} \prod_{\ell=0}^n (U_\ell + U_{\ell+1})^{-1} (U_{n+1} + (r - T_{n+1}))^{-1} \varepsilon p_{2(r-T_{n+1})}(x_i, x_k). \end{aligned}$$

If i changes type at T_{n+1} , then

$$\begin{aligned} & \phi_{T_{n+1}}(x_1, x_2, x_3) \\ & \leq c_1^{n+1} \prod_{\ell=0}^n (U_\ell + U_{\ell+1})^{-1} \varepsilon p_{2U_{n+1}}(x_k, x_j) \varepsilon p_0(x_i, x_k), \quad I_{T_{n+1}} = \{k\}, \end{aligned}$$

and so if $T_{n+1} \leq r < T_{n+2}$,

$$\begin{aligned} & \phi_r(x_1, x_2, x_3) \\ & \leq c_1^{n+1} \prod_{\ell=0}^n (U_\ell + U_{\ell+1})^{-1} \\ & \quad \times \int \varepsilon p_{2U_{n+1}+(r-T_{n+1})}(x_j, y_k) \varepsilon p_{r-T_{n+1}}(x_i, y_k) \varepsilon p_{r-T_{n+1}}(x_k, y_k) d^\varepsilon y_k \\ & \leq c_1^{n+2} \prod_{\ell=0}^n (U_\ell + U_{\ell+1})^{-1} (U_{n+1} + (r - T_{n+1}))^{-1} \varepsilon p_{2(r-T_{n+1})}(x_i, x_k), \end{aligned}$$

which gives (170_{n+1}). Finally if $T_0 \leq r < T_1$, $I_r = \{1, 2\}$ and

$$\begin{aligned} \phi_r(x_1, x_2, x_3) &= \int \varepsilon p_{s+r}(x_1, y_2) \varepsilon p_r(y_2, x_3) \varepsilon p_r(x_2, y_2) d^\varepsilon y_2 \\ &\leq c_1(s+r)^{-1} \varepsilon p_{2r}(x_2, x_3) \end{aligned}$$

and so (170₀) holds. This completes the inductive proof of (170_n) for $n \in \mathbb{Z}_+$.

It follows from (170) and (170_n) that

$$\begin{aligned}
 & E^{\varepsilon_{\mathbf{X}_0}} \int \phi_0 \varepsilon X_r^1(dx_1) \varepsilon X_r^1(dx_2) \varepsilon X_r^2(dx_3) \\
 & \leq \sum_{n=0}^{\infty} E \left(\mathbf{1}(T_n \leq r < T_{n+1}) e^{\gamma r} c_1^{n+1} \prod_{\ell=0}^{n-1} (U_\ell + U_{\ell+1})^{-1} \right. \\
 & \quad \left. \times (U_n + r - T_n)^{-1} \int \varepsilon p_{2(r-T_n)}(x_1, x_2) \varepsilon X_0^1(dx_1) \varepsilon X_0^2(dx_2) \right) \\
 & \quad \times (\langle X_0^1, 1 \rangle + \langle X_0^2, 1 \rangle) \\
 & \leq \sum_{n=0}^{\infty} (c_1 \gamma)^{n+1} \int_{\mathbb{R}_+^n} \mathbf{1} \left(\sum_1^n u_i \leq r \right) e^{\gamma r} \exp \left(-\gamma \left(r - \sum_1^n u_i \right) \right) \\
 & \quad \times \exp \left(-\gamma \sum_1^n u_i \right) (s + u_1)^{-1} \prod_{\ell=1}^{n-1} (u_\ell + u_{\ell+1})^{-1} \\
 & \quad \times \left(u_n + r - \sum_1^n u_i \right)^{-1} c_{35} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) \left(r - \sum_1^n u_i \right)^{-1/2} d\mathbf{u},
 \end{aligned}$$

where $0 < p' < 1/2$ and we have used Lemma 35 in the last inequality.

Therefore

$$\begin{aligned}
 & E \int_0^T \int \varepsilon p_s(x_1, x_2) \varepsilon X_r^1(dx_1) L^\varepsilon(d[r, x_2]) \\
 & = E \int_0^T dr \int \varepsilon p_s(x_1, x_2) \varepsilon p_0(x_2, x_3) \varepsilon X_r^1(dx_1) \varepsilon X_r^1(dx_2) \varepsilon X_r^2(dx_3) \\
 & \leq \sum_{n=0}^{\infty} (c_1 \gamma)^{n+1} \int_0^T dr \int_{\mathbb{R}_+^n} \mathbf{1} \left(\sum_1^n u_i \leq r \right) (s + u_1)^{-1} \\
 & \quad \times \prod_{\ell=1}^{n-1} (u_\ell + u_{\ell+1})^{-1} \left(u_n + r - \sum_1^n u_i \right)^{-1} \\
 & \quad \times \left(r - \sum_1^n u_i \right)^{-1/2} d\mathbf{u} \cdot c_{35} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) \\
 & \leq c_{35} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) \sum_{n=0}^{\infty} (c_1 \gamma)^{n+1} \\
 & \quad \times \int_{\mathbb{R}_+^{n+1}} \mathbf{1} \left(\sum_1^{n+1} u_i \leq T \right) (s + u_1)^{-1} \prod_{\ell=1}^n (u_\ell + u_{\ell+1})^{-1} u_{n+1}^{-1/2} d\mathbf{u} \\
 & \leq c_{35} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) \sum_{n=0}^{\infty} (c_1 \gamma \pi)^{n+1} s^{-1/2},
 \end{aligned}$$

the last by Lemma 60 in Appendix B below. Our choice of γ/σ^2 in (33) ensures the series is summable and so the above expected value is bounded by the required quantity. \square

PROOF OF THEOREM 21. The a.s. convergence of $\mathbf{X}_t(\mathbb{R}^2)$ is immediate from the martingale convergence theorem as $X_t^i(\mathbb{R}^2)$ is a nonnegative (hence L^1 -bounded) martingale. Since $\mathbf{X}_t(\mathbb{R}^2)$ is a conformal martingale [$X_t^i(\mathbb{R}^2)$ are orthogonal martingales with the same square function] $\mathbf{X}_t(\mathbb{R}^2) = B(A_t)$ for some planar Brownian motion B starting at $\mathbf{X}_0(\mathbb{R}^2)$, where $A_t = L_X(t)(\mathbb{R}^2)$. Clearly $\mathbf{X}_\infty(\mathbb{R}^2) = B(A_\infty)$, where $A_\infty \leq \tau_{\text{ex}}$ because $\mathbf{X}(\mathbb{R}^2)$ stays in the first quadrant. To complete the proof we need only prove

$$(172) \quad X_\infty^1(1)X_\infty^2(1) = 0 \quad \text{a.s.,}$$

as this clearly implies $A_\infty = \tau_{\text{ex}}$ a.s.

To prove (172) we may assume $\mathbf{X}_0 \in \mathcal{M}_{f,\text{se}}$ by applying the Markov property at a fixed time $\delta > 0$ and using Proposition 25(a). Let \mathbf{S}_t denote the four-dimensional Brownian semigroup, let M^i denote the martingale measures associated with X^i ($i = 1, 2$) and let $\Pi_t = X_t^1 \times X_t^2$ denote the product measure on \mathbb{R}^4 . We claim that if ϕ is bounded and Borel measurable on \mathbb{R}^4 , then for each $s > 0$, with probability 1,

$$(173) \quad \begin{aligned} \Pi_s(\phi) = \langle \Pi_0, \mathbf{S}_s \phi \rangle + \int_0^s \int \mathbf{S}_{s-r} \phi(x_1, x_2) [X_r^1(dx_1)M^2(dr, dx_2) \\ + X_r^2(dx_2)M^1(dr, dx_1)]. \end{aligned}$$

If $\phi(x_1, x_2) = \phi_1(x_1)\phi_2(x_2)$ for bounded measurable ϕ_i , then this is immediate from Corollary 24 and an integration by parts. The general result follows by passing to the bounded pointwise closure of the linear span of this class. Let $\varepsilon > 0$ and define

$$M_s = \int_0^s \int p_{\varepsilon+2(s-r)}(x_1, x_2) [X_r^1(dx_1)M^2(dr, dx_2) + X_r^2(dx_2)M^1(dr, dx_1)].$$

Now let $\phi(x_1, x_2) = p_\varepsilon(x_1, x_2)$ in (173) to get

$$(174) \quad \int p_\varepsilon(x_1, x_2)\Pi_s(dx_1, dx_2) = \int p_{\varepsilon+2s}(x_1, x_2)\Pi_0(dx_1, dx_2) + M_s.$$

Integrate s over $[0, T]$ and use a stochastic Fubini theorem ([45], Theorem 2.6) to conclude

$$(175) \quad \begin{aligned} & \int_0^T \int p_\varepsilon(x_1, x_2)\Pi_s(dx_1, dx_2) ds \\ & = \int_0^T \int p_{\varepsilon+2s}(x_1, x_2)\Pi_0(dx_1, dx_2) ds + \int_0^T M_s ds \end{aligned}$$

$$\begin{aligned}
 &= \int \left[\frac{1}{2} \int_{\varepsilon}^{\varepsilon+2T} p_r(x_1, x_2) dr \right] \Pi_0(dx_1, dx_2) \\
 &\quad + \int_0^T \int \left[\frac{1}{2} \int_{\varepsilon}^{\varepsilon+2(T-r)} p_u(x_1, x_2) du \right] \\
 &\quad \times [X_r^1(dx_1)M^2(dr, dx_2) + X_r^2(dx_2)M^1(dr, dx_1)].
 \end{aligned}$$

To check the integrability condition required for the stochastic Fubini theorem, note first that the expression on the left-hand side of (174) is L^2 -bounded in s (by Theorem 54 and our assumption that the initial measure is in $\mathcal{M}_{f,se}$) and the first term on the right-hand side of (174) is bounded. This shows that M_s is also L^2 -bounded in s and so $E(\int_0^T \langle M \rangle_s ds) < \infty$, which is the required condition in [45].

Let $h_{\delta,T} : \mathbb{R}_+ \rightarrow [0, 1]$ be the piecewise linear function satisfying $h_{\delta,T}(0) = h_{\delta,T}(x) = 0$ for all $x \geq 2T + \delta$ and $h_{\delta,T}(r) = 1$ for $r \in [\delta, 2T]$. Let $q_{\delta}(x, y) = \delta^{-1} \int_0^{\delta} p_r(x, y) dr$. The left-hand side of (175) equals

$$(176) \quad \int_0^T \int S_{\varepsilon/2} X_s^1(x) S_{\varepsilon/2} X_s^2(x) dx ds$$

by Chapman–Kolmogorov. By Theorem 11(a) (**SIntC**) holds, and this [we do not require the factor $|x - y|^{-1}$ in the definition of H_{ε} in this application of (**SIntC**)], together with the Cauchy–Schwarz inequality, shows that (176), and so the left-hand side of (175), remains L^2 -bounded as $\varepsilon \downarrow 0$. The first term on the right-hand side of (175) approaches

$$\frac{1}{2} \int G_{2T}(x_1, x_2) \Pi_0(dx_1, dx_2) < \infty,$$

where $G_{2T}(x, y) = \int_0^{2T} p_r(x, y) dr$ and the above is finite since $\mathbf{X}_0 \in \mathcal{M}_{f,se}$. This means the stochastic integral on the far right-hand side of (175) is also L^2 -bounded as $\varepsilon \downarrow 0$. This allows us to integrate (175) with respect to $\varepsilon \in (0, \delta]$ and again use the stochastic Fubini theorem to see that

$$\begin{aligned}
 &\int_0^T \int q_{\delta}(x_1, x_2) \Pi_s(dx_1, dx_2) ds \\
 &= \int \frac{1}{2} \int_0^{\delta+2T} p_r(x_1, x_2) h_{\delta,T}(r) dr \Pi_0(dx_1, dx_2) \\
 (177) \quad &+ \int_0^T \int \left[\frac{1}{2} \int_0^{\delta+2(T-r)} p_u(x_1, x_2) h_{\delta,T-r}(u) du \right] \\
 &\quad \times [X_r^1(dx_1)M^2(dr, dx_2) + X_r^2(dx_2)M^1(dr, dx_1)].
 \end{aligned}$$

As $\delta \downarrow 0$, the first term on the right-hand side approaches

$$\frac{1}{2} \int G_{2T}(x_1, y_2) \Pi_0(dx_1, dx_2) < \infty$$

by dominated convergence (recall that $\mathbf{X}_0 \in M_{f,se}$). The left-hand side converges in L^1 to $L_{\mathbf{X}}(T)(\mathbb{R}^2)$ by (147), and is L^2 -bounded as $\delta \downarrow 0$, by **(SIntC)** and the Cauchy–Schwarz inequality (as above), respectively. It follows that the square function of the stochastic integral remains L^1 -bounded as $\delta \downarrow 0$ and so by Fatou’s lemma

$$E\left(\int_0^T \int \left[\int G_{2(T-r)}(x_1, x_2) X_r^1(dx_1) \right]^2 L_{\mathbf{X}}(dr, dx_2) + \int_0^T \int \left[\int G_{2(T-r)}(x_1, x_2) X_r^2(dx_2) \right]^2 L_{\mathbf{X}}(dr, dx_1) \right) < \infty.$$

This and the above L^2 -boundedness readily allow us to see that the above integrals are still finite if $G_{2(T-r)}(x_1, x_2)$ is replaced with

$$\int_0^{\delta+2(T-r)} p_u(x_1, x_2) \tilde{h}_{\delta, T-r}(u) du,$$

where $\tilde{h}_{\delta, T-r}(u) = 1$ for $0 \leq u \leq 2(T - r)$ and agrees with $h_{\delta, T-r}$ elsewhere. Therefore we may apply dominated convergence to see that the stochastic integral in (177) converges in L^2 and conclude that

$$\begin{aligned} A_T &\equiv L_{\mathbf{X}}(T)(\mathbb{R}^2) \\ &= \frac{1}{2} \int G_{2T}(x_1, x_2) \Pi_0(dx_1, dx_2) \\ (178) \quad &+ \int_0^T \int \left[\frac{1}{2} G_{2(T-r)}(x_1, x_2) \right] \\ &\quad \times [X_r^1(dx_1)M^2(dr, dx_2) + X_r^2(dx_2)M^1(dr, dx_1)] \\ &\equiv A_T^1 + N_T, \end{aligned}$$

where N_T is in L^2 .

Choose $M = M(X_0) \in \mathbb{N}$ so that $X_0^i(B(0, M/2)) \geq \frac{1}{2} X_0^i(\mathbb{R}^2)$. If $T \geq M^4$ and $G_{2T}(M) = \inf_{|x_1-x_2|<M} G_{2T}(x_1, x_2)$, then

$$(179) \quad A_T^1 \geq \frac{1}{8} G_{2T}(M) X_0^1(1) X_0^2(1) \geq c_1(\log T) X_0^1(1) X_0^2(1)$$

for some universal constant $c_1(M)$ by an elementary calculation. Let $p > 1$ and set

$$\begin{aligned} \eta_T^i &= \int_0^T \int \left\{ \int \left[\int_0^{T-p} p_s(x_1, x_2) ds \right] X_r^i(dx_1) \right\}^2 L_{\mathbf{X}}(dr, dx_2), \\ \delta_T^i &= \int_0^{T-p} \log(T^{-p}/s) \left[\int_0^T \int p_s(x_1, x_2) X_r^i(dx_1) L_{\mathbf{X}}(dr, dx_2) \right] ds. \end{aligned}$$

If $X_T^{i*}(1) = \sup_{r \leq T} X_r^i(1)$, then (for $T \geq 2$)

$$\begin{aligned}
 \langle N \rangle_T &= \frac{\gamma}{4} \sum_{i=1,2} \int_0^T \int \left[\int G_{2(T-r)}(x_1, x_2) X_r^i(dx_1) \right]^2 L_{\mathbf{X}}(dr, dx_2) \\
 &\leq \frac{\gamma}{2} \sum_{i=1,2} \left[\int_0^T \int \left[\iint_{T-p}^{2T} p_s(x_1, x_2) ds X_r^i(dx_1) \right]^2 L_{\mathbf{X}}(dr, dx_2) + \eta_T^i \right] \\
 &\leq c(p, \gamma, \sigma) (\log T)^2 \left(\sum_{i=1,2} X_T^{i*}(1)^2 \right) L_{\mathbf{X}}(T)(\mathbb{R}^2) \\
 (180) \quad &+ \sum_{i=1,2} c(\gamma, \sigma) \int_0^T \iint \left[\int_0^{T-p} p_{s_1}(x_1, x_2) \int_{s_1}^{T-p} \frac{ds}{s} ds_1 \right] \\
 &\quad \times X_r^i(dx'_1) X_r^i(dx_1) L_{\mathbf{X}}(dr, dx_2) \\
 &\leq c_2 (\log T)^2 \left[\sum_{l=1,2} X_T^{l*}(1)^2 \right] A_T + c_2 \sum_{i=1,2} X_T^{i*}(1) \delta_T^i,
 \end{aligned}$$

recalling $A_T \equiv L_{\mathbf{X}}(T)(\mathbb{R}^2)$.

Lemma 56 shows that if $0 < p' < 1/2$,

$$\begin{aligned}
 E(\delta_T^i) &\leq c_{56} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) \int_0^{T-p} (\log(T^{-p}/s)) s^{-1/2} ds \\
 (181) \quad &= c_{56} \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) \left[\int_0^1 (\log 1/u) u^{-1/2} du \right] T^{-p/2} \\
 &\leq c_3 \bar{\mathcal{E}}_{p'}(\mathbf{X}_0) T^{-p/2}.
 \end{aligned}$$

Assume $X_0^1(1)X_0^2(1) > 0$ and $T_1(\mathbf{X}_0)$ is chosen large enough so that, for $T \geq T_1(\mathbf{X}_0)$,

$$(182) \quad c_1 (\log T) X_0^1(1) X_0^2(1) \geq 2 \quad \text{and} \quad T \geq 2 \vee M^4.$$

Then (178) and (179) imply that, for $T \geq T_1(\mathbf{X}_0)$,

$$\begin{aligned}
 &P(A_T \leq 1) \\
 &\leq P(c_1 (\log T) X_0^1(1) X_0^2(1) + N_T \leq 1, A_T \leq 1) \\
 &\leq P\left(N_T \leq -\frac{c_1}{2} (\log T) X_0^1(1) X_0^2(1), A_T \leq 1, \right. \\
 &\quad \left. X_T^{1*}(1) \vee X_T^{2*}(1) \leq R, \delta_T^1 \vee \delta_T^2 \leq R(\log T)^2 \right) \\
 (183) \quad &+ \sum_{i=1,2} P(X_T^{i*}(1) > R, A_T \leq 1) + P(\delta_T^i > R(\log T)^2) \quad [\text{by (182)}]
 \end{aligned}$$

$$\begin{aligned} &\leq P\left(N_T \leq -\frac{c_1}{2}(\log T)X_0^1(1)X_0^2(1), \langle N \rangle_T \leq 4c_2(\log T)^2R^2\right) \\ &\quad + \left(\sum_{i=1,2} P_{X_0^i(1)}\left(\sup_{s \leq 1} B_s \geq R\right)\right) + 2R^{-1}(\log T)^{-2}c_3\bar{\varepsilon}_{p'}(\mathbf{X}_0)T^{-p/2}, \end{aligned}$$

where in the last line B is a linear Brownian motion starting at x under P_x , and we have used (180), (181) and the Dubins–Schwarz theorem to write $X_t^i(\mathbb{R}^2)$ as $B(A_t)$. Assume

$$R \geq \max(2(\langle X_0^1, 1 \rangle + \langle X_0^2, 1 \rangle), \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle).$$

Then an elementary calculation with Brownian motion, again using Dubins–Schwarz (see [14], (3.12) and (3.13)) shows that the first term on the right-hand side of (183) is at most

$$1 - c_4 \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle R^{-1}$$

($c_4 > 0$ universal) and the second term is at most

$$\frac{8}{R} \exp\left(-\frac{R^2}{8}\right).$$

Now set

$$\begin{aligned} R &= R(\langle X_0^1, 1 \rangle, \langle X_0^2, 1 \rangle) \\ &= \max\left(2(\langle X_0^1, 1 \rangle + \langle X_0^2, 1 \rangle), \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle, \right. \\ &\quad \left. [8|\log[32(c_4 \langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle)^{-1}]|]^{1/2}\right) \end{aligned}$$

and then assume $T \in \mathbb{N}$, in addition to (182), also satisfies $T \geq T_2(\mathbf{X}_0)$ to ensure the last term on the right-hand side of (183) is at most $\frac{c_4}{4}X_0^1(1)X_0^2(1)R^{-1}$. In fact, define $T(\mathbf{X}_0)$ to be the smallest such T in \mathbb{N} . Set $T \equiv \infty$ if $X_0 \notin M_{f,se}$ or $X_0^1(1)X_0^2(1) = 0$. Combining the above bounds and using them in (183), we get

$$\begin{aligned} P(A_T > 1) &\geq c_4 X_0^1(\mathbb{R}^2) X_0^2(\mathbb{R}^2) R^{-1} - \frac{8}{R} e^{-R^2/8} - \frac{c_4}{4} X_0^1(\mathbb{R}^2) X_0^2(\mathbb{R}^2) R^{-1} \\ (184) \quad &\geq \frac{c_4}{4} X_0^1(\mathbb{R}^2) X_0^2(\mathbb{R}^2) R(X_0^1(\mathbb{R}^2), X_0^2(\mathbb{R}^2))^{-1} \quad (\text{by the choice of } R) \\ &\equiv q(X_0^1(\mathbb{R}^2), X_0^2(\mathbb{R}^2)). \end{aligned}$$

Set $q(0, x) = q(x, 0) = 0$ so that (184) remains valid if $\langle X_0^1, 1 \rangle \langle X_0^2, 1 \rangle = 0$. Note that

$$(185) \quad \inf\{q(u, v) : u \geq \delta, v \geq \delta\} = \varepsilon(\delta) > 0 \quad \forall \delta > 0.$$

Inductively define \mathbb{N} -valued stopping times by $T_{n+1} = T(X_{T_n}^1, X_{T_n}^2) + T_n \leq \infty$. By the Markov property for X if $\mathcal{F}_t^X = \sigma(\mathbf{X}_r : r \leq t)$,

$$\begin{aligned} P_{\mathbf{X}_0}(A_{T_{n+1}} - A_{T_n} \geq 1 \mid \mathcal{F}_{T_n}^X) \\ = P_{\mathbf{X}_{T_n}}(A(T_1(\mathbf{X}_0)) \geq 1) \mathbf{1}(T_n < \infty) \geq q(\mathbf{X}_{T_n}(1)) \mathbf{1}(T_n < \infty) \quad [\text{by (184)}]. \end{aligned}$$

Now use the conditional version of the Borel–Cantelli lemma and the fact that $\lim_{t \rightarrow \infty} A_t = A_\infty < \infty$ a.s. [because $X_t^i(\mathbb{R}^2) \xrightarrow{\text{a.s.}} X_\infty^i(\mathbb{R}^2) < \infty$] as in (3.18) of [14] to conclude that

$$(186) \quad \sum_{n=1}^{\infty} q(\mathbf{X}_{T_n}(\mathbb{R}^2)) \mathbf{1}(T_n < \infty) < \infty \quad \text{a.s.}$$

If $T_n < \infty$ for all n , then (185) and (186) imply $\lim_{n \rightarrow \infty} X_{T_n}^1(\mathbb{R}^2) X_{T_n}^2(\mathbb{R}^2) = 0$ a.s. and so $\lim_{t \rightarrow \infty} X_t^1(\mathbb{R}^2) X_t^2(\mathbb{R}^2) = 0$ a.s. by martingale convergence. If $T_n = \infty$ for some n , then let n_0 be the first such n . Since $X_k \in M_{f,se}$ for all $k \in \mathbb{Z}_+$ a.s. by Proposition 25(a), this implies $X_{T_{n_0-1}}^1(\mathbb{R}^2) X_{T_{n_0-1}}^2(\mathbb{R}^2) = 0$ and therefore, $X_t^1(\mathbb{R}^2) X_t^2(\mathbb{R}^2) = 0$ for all $t \geq T_{n_0-1}$. The required result is established in either case. \square

6. Existence of densities and segregation of types. We start with a general result giving the existence of densities for a class of measure-valued martingale problems based on a conformal martingale argument. Write $\mathcal{M} = \mathcal{M}(\mathbb{R}^d)$ for the space of all Radon measures on \mathbb{R}^d equipped with the topology of vague convergence and let $\mathcal{C}_{\text{com}}^\infty(\mathbb{R}^d)$ be the space of infinitely differentiable functions on \mathbb{R}^d with compact support.

THEOREM 57. *Let Q_t denote a Feller semigroup on \mathbb{R}^d , let $T > 0$ and assume $\mathbf{X}_t = (X_t^1, X_t^2)$, $0 \leq t \leq T$, is an adapted continuous \mathcal{M}^2 -valued process on $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$. Suppose that for some $c > 0$, for all nonnegative $\varphi^j \in \mathcal{C}_{\text{com}}^\infty(\mathbb{R}^d)$, $j = 1, 2$,*

$$(187) \quad N_t^j(\varphi^j) = \langle X_t^j, Q_{T-t}\varphi^j \rangle, \quad t \leq T, j = 1, 2,$$

are orthogonal \mathcal{F}_t -martingales whose predictable square functions satisfy

$$\langle \langle N^1(\varphi^1) \rangle \rangle_t = c \langle \langle N^2(\varphi^1) \rangle \rangle_t, \quad 0 \leq t \leq T.$$

Then $X_T^j \ll \ell$, P -a.s. for $j = 1, 2$ if and only if $Q_T X_0^j \ll \ell$, P -a.s. for $j = 1, 2$.

PROOF. By working with the regular conditional probability for \mathbf{X} given \mathbf{X}_0 we may assume that \mathbf{X}_0 is deterministic (it suffices to assume the above for a countable supnorm dense set of φ^j 's).

STEP 1. First we assume that $c = 1$. Fix a nonnegative $\varphi \in \mathcal{C}_{\text{com}}^\infty(\mathbb{R}^d)$. Set $X := X^1 + iX^2$ and $N(\varphi) := N^1(\varphi) + iN^2(\varphi)$. Then $N(\varphi)$ is a conformal martingale (see, e.g., [39], Section V.2), and Itô's lemma shows that the bounded process $t \mapsto e^{-N_t(\varphi)}$ is a continuous \mathcal{F} -martingale. We therefore have

$$(188) \quad E\langle X_T, \varphi \rangle = EN_T(\varphi) = N_0(\varphi) = \langle X_0, Q_T\varphi \rangle$$

and

$$(189) \quad Ee^{-\langle X_T, \varphi \rangle} = Ee^{-N_T(\varphi)} = e^{-N_0(\varphi)} = e^{-\langle X_0, Q_T\varphi \rangle}.$$

Let $\{\varphi_n : n \geq 1\}$ denote a (nonnegative) radially symmetric approximate identity (which is approximating the δ_0 -function) in $\mathcal{C}_{\text{com}}^\infty(\mathbb{R}^d)$. Set $\varphi_n^x(y) := \varphi_n(y - x)$, $x, y \in \mathbb{R}^d$. Since $B \mapsto \langle X_0, Q_T(\mathbf{1}_B\varphi) \rangle =: \mu(B)$ is a finite complex measure, we may apply standard differentiation theory of measures (see, e.g., [40], Theorem 8.6). From the identity (188) we conclude that

$$(190) \quad \begin{aligned} EN_T(\varphi_n^x\varphi) &= \langle X_0, Q_T(\varphi_n^x\varphi) \rangle \xrightarrow[n \uparrow \infty]{} f(x) \\ &=: f^1(x) + if^2(x) \quad \text{for } \ell\text{-a.a. } x, \end{aligned}$$

where f is the density of the absolutely continuous part of μ :

$$(191) \quad \mu(\cdot) = \int_{(\cdot)} \ell(dx)f(x) + \nu(\cdot), \quad \nu \perp \ell.$$

Note that $f^j \geq 0$, $j = 1, 2$ and

$$(192) \quad \int \ell(dx)f^j(x) \leq \langle X_0^j, Q_T\varphi \rangle = N_0^j(\varphi) < \infty$$

hence $f^j(x) < \infty$ for ℓ -almost all x . Applying the same argument to the random finite complex measure $B \mapsto \langle X_T, \mathbf{1}_B\varphi \rangle$, we see that

$$(193) \quad \langle X_T, \varphi_n^x\varphi \rangle \xrightarrow[n \uparrow \infty]{} \eta(x) =: \eta^1(x) + i\eta^2(x) \quad \text{for } \ell \times P\text{-a.a. } (x, \omega),$$

where η is the density of the absolutely continuous part of $\langle X_T, \mathbf{1}_{(\cdot)}\varphi \rangle$. Fatou's lemma gives

$$(194) \quad \begin{aligned} E\eta^j(x) &\leq \liminf_{n \uparrow \infty} E\langle X_T^j, \varphi_n^x\varphi \rangle = \liminf_{n \uparrow \infty} \langle X_0^j, Q_T(\varphi_n^x\varphi) \rangle \\ &= f^j(x) < \infty \quad \text{for } \ell\text{-a.a. } x. \end{aligned}$$

Now (190) shows that, for ℓ -a.a. x and for $\theta \geq 0$,

$$e^{-\theta f(x)} = \lim_{n \uparrow \infty} \exp[-\theta \langle X_0, Q_T(\varphi_n^x\varphi) \rangle],$$

which by (189), (193) and bounded convergence, equals

$$\lim_{n \uparrow \infty} E \exp[-\theta \langle X_T, \varphi_n^x\varphi \rangle] = Ee^{-\theta \eta(x)}.$$

We use the finiteness in (194) to differentiate $Pe^{-\theta\eta(x)}$ with respect to θ at $\theta = 0+$ and conclude

$$(195) \quad E\eta(x) = f(x) < \infty \quad \text{for } \ell\text{-a.a. } x.$$

STEP 2. Assume now that $\mathbf{X}_0 * \mathbf{Q}_T \ll (\ell, \ell)$. Then, by (188) and since $\nu = 0$ in the decomposition (191),

$$E\langle X_T^j, \varphi \rangle = \langle X_0^j, \mathbf{Q}_T \varphi \rangle = \int \ell(dx) f^j(x) = E \int \ell(dx) \eta^j(x),$$

where in the last step we used (195). This shows the singular part of $B \mapsto \langle X_T^j, \mathbf{1}_B \varphi \rangle$ is a.s. 0 and as φ is an arbitrary smooth nonnegative function with compact support, we may conclude that $X_T^j \ll \ell$ P -a.s.

STEP 3. Conversely, assume that $X_T^j \ll \ell$, P -a.s., $j = 1, 2$. Then, if B is a Lebesgue null set in \mathbb{R}^2 , we get $X_T^j(B) = 0$, P -a.s., and so

$$\langle X_0^j, \mathbf{Q}_T \mathbf{1}_B \rangle = E\langle X_T^j, \mathbf{1}_B \rangle = 0.$$

In fact, in the first equality we have extended (188) from $\varphi \in \mathcal{C}_{\text{com}}^\infty$ to bounded measurable φ by a standard monotone class argument.

STEP 4. The result for general c now follows by applying the above to $(c^{-1/2}X^1, X^2)$. \square

Although the above result may appear to be fairly general, a bit of thought will convince the reader that these hypotheses are not readily satisfied. Of course we have just worked rather hard to find at least one case where they are satisfied.

PROOF OF THEOREM 17(a). Corollary 24 shows that the hypothesis of Theorem 57 holds with $Q_t = S_t$, the Brownian semigroup, and $c = 1$. The absolute continuity of the Brownian semigroup and Theorem 57 completes the proof. \square

REMARK 58. Note that the proof of Theorem 17(a) only relied on a result (Corollary 24) which was established for any solution of $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \nu}$ independently of our uniqueness results, and on the general Theorem 57, which is independent of the other results in this paper. This will allow us to use the above existence of densities in the derivation of uniqueness in law and the strong Markov property in [12].

The proof of the segregation-of-types result, Theorem 17(b), will be an adaptation of the method of Cox, Klenke and Perkins [7], which was designed to prove convergence to equilibria from more general initial conditions once it is established from uniform initial measures, and will be used for precisely this purpose in [8]. Given the close links between the local and longtime behaviors (cf. [8]), this connection is not surprising.

As $\varepsilon \downarrow 0$ and $k \rightarrow \infty$, the integrand converges pointwise to 0, and for $r \in [0, k(t - \eta))$ is uniformly bounded by c/η . Therefore

$$(200) \quad \mathbf{1}(r < kt) S_{kt-r} \Delta_{k,\varepsilon}^i \rightarrow 0 \text{ pointwise as } \varepsilon \rightarrow 0, k \rightarrow \infty \text{ and} \\ \sup\{S_{kt-r} \Delta_{k,\varepsilon}^i(z) : r \leq k(t - \eta), z \in \mathbb{R}^2, k \in \mathbb{N}, \varepsilon > 0\} \leq \frac{c}{\eta} X_0^i(\mathbb{R}^2).$$

By dominated convergence, the first term on the right-hand side of (198) approaches 0 as $\varepsilon \downarrow 0$ and $k \rightarrow \infty$. Turning to the second term, let

$$N_{k,\varepsilon}^{i,j}(s) = \int_0^s \int_{\mathbb{R}^2} S_{kt-r} \Delta_{k,\varepsilon}^i(y) d\tilde{M}^j(r, y), \quad s \leq kt.$$

Then

$$\langle N_{k,\varepsilon}^{i,j}(kt - \eta) \rangle = \gamma \int_0^{k(t-\eta)} \int_{\mathbb{R}^2} (S_{kt-r} \Delta_{k,\varepsilon}^i(y))^2 L_{\tilde{\mathbf{X}}}(dr, dy),$$

which approaches 0 a.s. as $\varepsilon \downarrow 0$ and $k \rightarrow \infty$ by (200), dominated convergence, and the fact that $L_{\tilde{\mathbf{X}}}(t, \mathbb{R}^2) \rightarrow L_{\tilde{\mathbf{X}}}(\infty, \mathbb{R}^2) < \infty$ a.s. as $t \rightarrow \infty$. The latter is true because $\gamma L_{\tilde{\mathbf{X}}}(t, \mathbb{R}^2)$ is the square function of the nonnegative martingale $\tilde{X}_t^i(\mathbb{R}^2)$ which therefore must converge a.s. Now use Proposition 15(c) to see that

$$E_{a_1\mu_0, a_2\mu_0}(\langle N_{k,\varepsilon}^{i,j}(kt) \rangle - \langle N_{k,\varepsilon}^{i,j}(kt - \eta) \rangle) \\ \leq \gamma \int_{k(t-\eta)}^{kt} \int_{\mathbb{R}^2} (S_{kt-r} \Delta_{k,\varepsilon}^i(y))^2 a_1 a_2 S_r \mu_0(y)^2 dy dr \\ \leq c(t, \mathbf{X}_0) \\ \times \left[\int_{k(t-\eta)}^{kt} \int_{\mathbb{R}^2} \left[\int p_{t+\varepsilon-rk^{-1}}(w - x - yk^{-1/2}) X_0^i(dw) \right]^2 p_{r+1}(y)^2 dy dr \right. \\ \left. + \int_{k(t-\eta)}^{kt} \int_{\mathbb{R}^2} p_{r+1}(y)^2 dy dr \right],$$

where we have used (199) in the last line. This in turn is bounded by

$$c(t, \mathbf{X}_0) \left[\int_{k(t-\eta)}^{kt} \int p_{t+\varepsilon-rk^{-1}}(w_1 - x - yk^{-1/2}) \right. \\ \times p_{t+\varepsilon-rk^{-1}}(w_2 - x - yk^{-1/2}) dy (r + 1)^{-2} \\ \times X_0^i(dw_1) X_0^i(dw_2) dr + \int_{k(t-\eta)}^{kt} p_{2(r+1)}(0) dr \left. \right] \\ \leq c(t, \mathbf{X}_0) \left[\int_{k(t-\eta)}^{kt} \int k p_{2(t+\varepsilon-rk^{-1})}(w_1 - w_2) X_0^i(dw_1) X_0^i(dw_2) (r + 1)^{-2} dr \right. \\ \left. + \log\left(\frac{t}{t - \eta}\right) \right]$$

$$\begin{aligned} &\leq c(t, \mathbf{X}_0) \left[\int_{k(t-\eta)}^{kt} (t + \varepsilon - rk^{-1})^{-1/2} (r + 1)^{-2} k \, dr + \log\left(\frac{t}{t-\eta}\right) \right] \\ &\hspace{20em} (\text{since } \mathbf{X}_0 \in M_{f,se}) \\ &\leq c'(t, \mathbf{X}_0)[\eta^{1/2} + \eta] \rightarrow 0 \quad \text{as } \eta \downarrow 0. \end{aligned}$$

It follows from the above results that

$$\langle N_{k,\varepsilon}^{i,j} \rangle(kt) \xrightarrow{P_{a_1\mu_0, a_2\mu_0}} 0 \quad \text{a.s., } \varepsilon \downarrow 0 \text{ and } k \rightarrow \infty,$$

and so by a standard martingale inequality, the second term on the right-hand side of (198) [i.e., $N_{k,\varepsilon}^{i,j}(kt)$] also converges to 0 in $P_{a_1\mu_0, a_2\mu_0}$ -probability as $\varepsilon \downarrow 0$ and $k \rightarrow \infty$. We have proved $\langle \tilde{X}_{kt}^j, \Delta_{k,\varepsilon}^i \rangle \xrightarrow{P_{a_1\mu_0, a_2\mu_0}} 0$ as $\varepsilon \downarrow 0$ and $k \rightarrow \infty$ and so (197) now gives

$$\begin{aligned} &E_{\mathbf{X}_0}(e^{-aX_t(x)+ibY_t(x)}) \\ (201) \quad &= \lim_{k \rightarrow \infty} E_{a_1\mu_0, a_2\mu_0}(\{-\langle \tilde{X}_{kt}, 1 \rangle S_t X_0(x) + i\langle \tilde{Y}_{kt}, 1 \rangle S_t Y_0(x)\}) \\ &= E_{a_1, a_2}^0(e^{-S_t X_0(x)(B_{\tau_{\text{ex}}}^1 + B_{\tau_{\text{ex}}}^2) + iS_t Y_0(x)(B_{\tau_{\text{ex}}}^1 - B_{\tau_{\text{ex}}}^2)}) \quad (\text{by Theorem 21}) \\ &= E_{S_t X_0^1(x), S_t X_0^2(x)}^0(e^{-a(B_{\tau_{\text{ex}}}^1 + B_{\tau_{\text{ex}}}^2) + ib(B_{\tau_{\text{ex}}}^1 - B_{\tau_{\text{ex}}}^2)}). \end{aligned}$$

The last equality is an easy exercise on harmonic functions which may be found in the proof of Theorem 1.5 in [14]. An easy application of the Stone–Weierstrass theorem, as in the proof of Lemma 2.3(b) in [14], shows that the above joint Laplace–Fourier transforms for $a_1, a_2 \geq 0$ uniquely determine the law of $(x_t(x), y_t(x))$ and the result follows for $\mathbf{X}_0 \in \mathcal{M}_{f,se}$.

Assume now that $\mathbf{X}_0 \in \mathcal{M}_{f,e}$. Let $\delta_n \in (0, t)$ decrease to 0. By Proposition 25(a), $\mathbf{X}_{\delta_n} \in \mathcal{M}_{f,se}$ a.s. and so the Markov property and (201) imply

$$\begin{aligned} E_{\mathbf{X}_0}(e^{-aX_t(x)+ibY_t(x)}) &= E_{\mathbf{X}_0}(E_{\mathbf{X}_{\delta_n}}(e^{-ax_{t-\delta_n}(x)+iby_{t-\delta_n}(x)})) \\ &= E_{\mathbf{X}_0}(E_{a_1, a_2}^0(e^{-S_{t-\delta_n} X_{\delta_n}(x)(B_{\tau_{\text{ex}}}^1 + B_{\tau_{\text{ex}}}^2) + iS_{t-\delta_n} Y_{\delta_n}(x)(B_{\tau_{\text{ex}}}^1 - B_{\tau_{\text{ex}}}^2)})) \\ &\rightarrow E_{a_1, a_2}^0(e^{-S_t X_0(x)(B_{\tau_{\text{ex}}}^1 + B_{\tau_{\text{ex}}}^2) + iS_t Y_0(x)(B_{\tau_{\text{ex}}}^1 - B_{\tau_{\text{ex}}}^2)}) \quad \text{as } n \rightarrow \infty. \end{aligned}$$

In the last line we have used dominated convergence, the a.s. continuity of \mathbf{X}_t and the uniform convergence of $p_{t-\delta_n}(\cdot)$ to $p_t(\cdot)$. This establishes (201) for $\mathbf{X}_0 \in \mathcal{M}_{f,e}$ and the proof may be completed just as in the previous case. \square

PROOF OF COROLLARY 19. Let $\{B_k : k \in \mathbb{N}\}$ be the set of open balls in \mathbb{R}^2 with rational centers and radii. Choose nonnegative $\{\phi_k\} \subset \mathcal{C}_{\text{com}}(\mathbb{R}^2)$ such that $\{\phi_k > 0\} = B_k$. We may fix $\varepsilon_n \downarrow 0$ such that

$$(202) \quad \langle L_{\mathbf{X}}^{*, \varepsilon_n}(t), \phi_k \rangle \rightarrow \langle L_{\mathbf{X}}(t), \phi_k \rangle \quad \forall t \in \mathbb{Q}_+, \forall k, \text{ a.s.}$$

By Theorem 17 we may fix ω outside a null set such that (202) holds,

$$(203) \quad X_s^i(dx) = X_s^i(x) dx \quad \text{for Lebesgue a.a. } s > 0$$

and

$$(204) \quad \int_0^\infty \int_{\mathbb{R}^2} X_s^1(x) X_s^2(x) dx ds = 0.$$

It clearly suffices to show that, for this fixed choice of ω , the desired conclusion holds for $U = (r_1, r_2) \times B_k$ for a fixed k and fixed rationals $0 \leq r_1 < r_2$. Assume

$$L_{\mathbf{X}}(r_2)(B_k) - L_{\mathbf{X}}(r_1)(B_k) > 0$$

and, say, $\|x^1\|_U < \infty$. Clearly $\exists B_{k'} \subset \overline{B_{k'}} \subset B_k$ such that $\langle L_{\mathbf{X}}(r_2), \phi_{k'} \rangle - \langle L_{\mathbf{X}}(r_1), \phi_{k'} \rangle > 0$. Then by (202)

$$\begin{aligned} & \lim_{n \rightarrow \infty} \varepsilon_n^{-1} \int_0^{\varepsilon_n} dr \int_{r_1}^{r_2} ds \left[\int \phi_{k'}(y) S_r X_s^1(y) S_r X_s^2(y) dy \right] \\ & = \langle L_{\mathbf{X}}(r_2), \phi_{k'} \rangle - \langle L_{\mathbf{X}}(r_1), \phi_{k'} \rangle > 0, \end{aligned}$$

which by Fubini implies

$$(205) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{r_1}^{r_2} \int_{\mathbb{R}^2} \left[\frac{1}{\varepsilon_n} \int_0^{\varepsilon_n} S_r X_s^1(y) S_r X_s^2(y) dr \right] \phi_{k'}(y) dy ds \\ & = \langle L_{\mathbf{X}}(r_2), \phi_{k'} \rangle - \langle L_{\mathbf{X}}(r_1), \phi_{k'} \rangle > 0. \end{aligned}$$

On the other hand (203) implies

$$(206) \quad S_r X_s^i(y) \rightarrow X_s^i(y) \quad \text{for Lebesgue a.a. } (s, y) \text{ a.s., } r \downarrow 0.$$

Let $d(B_{k'}, B_k^c) = \eta_k > 0$. Recall $\|\cdot\|_U$ denotes the essential supremum with respect to Lebesgue measure on the space-time open set U . We abuse this notation slightly and let $\|\cdot\|_{B_k}$ denote the essential sup with respect to Lebesgue measure on B_k . We may fix $s \in (r_1, r_2)$ outside a Lebesgue null set so that (203) holds and $\|x_s^1\|_{B_k} \leq \|x^1\|_U < \infty$. If $y \in B_{k'}$, then

$$\begin{aligned} S_r X_s^1(y) & \leq \int_{B_k} p_r(z-y) x_s^1(z) dz + \int_{B_k^c} p_r(z-y) X_s^1(dz) \\ & \leq \|x_s^1\|_{B_k} + p_r(\eta_k) X_s^1(\mathbb{R}^2) \\ & \leq \|x^1\|_U + X_s^1(\mathbb{R}^2) \end{aligned}$$

providing $r < r(k)$, where $r(k) > 0$. This implies that, for $\varepsilon_n < r(k)$, $y \in B_{k'}$ and Lebesgue a.a. $s \in (r_1, r_2)$, we have

$$(207) \quad \frac{1}{\varepsilon_n} \int_0^{\varepsilon_n} S_r X_s^1(y) S_r X_s^2(y) dr \leq \left[\|x^1\|_U + \sup_{s \leq r_2} X_s^1(\mathbb{R}^2) \right] \frac{1}{\varepsilon_n} \int_0^{\varepsilon_n} S_r X_s^2(y) dr.$$

Assume for the moment that

$$(208) \quad H_n(s, y) = \frac{1}{\varepsilon_n} \int_0^{\varepsilon_n} S_r X_s^2(y) dr \quad (n \in \mathbb{N})$$

is a uniformly integrable family on $((r_1, r_2) \times \mathbb{R}^2, \phi_{k'}(y) ds dy)$.

Then (207) allows us to take the limit in (205) through the first two integrals and conclude that the limit on the left-hand side of (205) equals

$$\begin{aligned} & \int_{r_1}^{r_2} \int_{\mathbb{R}^2} \left[\lim_{n \rightarrow \infty} \frac{1}{\varepsilon_n} \int_0^{\varepsilon_n} S_r X_s^1(y) S_r X_s^2(y) dr \right] \phi_{k'}(y) dy ds \\ &= \int_{r_1}^{r_2} \int_{\mathbb{R}^2} X_s^1(y) X_s^2(y) \phi_{k'}(y) dy ds \quad [\text{by (206)}] \\ &= 0 \quad [\text{by (204)}]. \end{aligned}$$

This contradicts (205) and so shows that for ω as above $L_{\mathbf{X}}(U) > 0$ implies $\|x^1\|_U = \infty$. By symmetry the proof is complete except for the verification of (208). To this end note that, by (206), $\lim_{n \rightarrow \infty} H_n(s, y) = x_s^2(y)$ for Lebesgue a.a. (s, y) and

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_{r_1}^{r_2} \int H_n(s, y) \phi_{k'}(y) dy ds \\ &= \lim_{n \rightarrow \infty} \int_{r_1}^{r_2} \int \frac{1}{\varepsilon_n} \int_0^{\varepsilon_n} S_r \phi_{k'}(y) dr x_s^2(y) dy ds \\ &= \int_{r_1}^{r_2} \int \phi_{k'}(y) x_s^2(y) dy ds \quad (\text{by dominated convergence}). \end{aligned}$$

Since $H_n \geq 0$ (208) follows, and the proof is complete. \square

7. Some open questions. An intriguing feature of this process is the volatile nature of its densities. There are a number of interesting open problems about the qualitative nature of the densities but, after spending three papers just to get existence, uniqueness and the basic features of the process straight, we will leave these for another day and perhaps another bunch of authors. Throughout this section (X_t^1, X_t^2) denotes the unique solution of $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ starting at $\mathbf{X}_0 \in \mathcal{M}_{f,e}$.

We know from Theorem 17 that at a fixed time the densities segregate and the measures are mutually singular. This does not, however, say anything about their closed supports. Let $S(X_t^i)$ denote the closed support of X_t^i and let

$$G(X^i) = \text{cl}\{(t, x) : x \in S(X_t^i), t > 0\}$$

(cl denotes closure in $\mathbb{R}_+ \times \mathbb{R}^2$) be the closed graph of X^i for $i = 1, 2$.

CONJECTURE 1. The interface $I = G(X^1) \cap G(X^2)$ is a.s. Lebesgue null in $\mathbb{R}_+ \times \mathbb{R}^2$ and there are versions of the densities $x^i(\cdot, \cdot)$ which are smooth on I^c and satisfy $\frac{\partial x^i}{\partial t} = \frac{\sigma^2 \Delta x^i}{2}$ on I^c .

CONJECTURE 2. For $t > 0$, the fixed time interface $S(X_t^1) \cap S(X_t^2)$ is a.s. Lebesgue null.

Assuming the second conjecture is correct we have the following question.

QUESTION 3. What is the Hausdorff dimension of $S(X_t^1) \cap S(X_t^2)$?

Uniform-in-time behavior leads to an even more difficult set of problems. Even the simplest kind of uniform-in- t nonsingularity (membership in $\mathcal{M}_{f,e}$ for all $t \geq 0$ a.s.) led to some nontrivial arguments in Proposition 25(b) and we were never able to decide if in fact $X_t^i \in \mathcal{M}_{f,se}$ for all $t > 0$ a.s. The fact that the existence of the densities at a fixed time is rather delicate means the existence for all t is uncertain.

QUESTION 4. Is $X_t^i(dx) \ll dx$ for all $t > 0$ a.s.? Is $S(X_t^1) \cap S(X_t^2)$ Lebesgue null for all $t > 0$ a.s.?

We showed in Corollary 19 that the densities blow up at typical points in the interface.

QUESTION 5. Can one find a canonical rate of explosion of $x^i(t, x)$ as x approaches x_0 for L_X a.a. (t, x_0) ?

As mentioned near the end of Section 1.1 we feel that the results of this paper should hold for any (γ, σ^2) .

PROBLEM 6. Prove this.

Having done this, the reader may then want to turn to higher dimensions. Recall for super-Brownian motion branching in a super-Brownian medium, the process will only exist in three or fewer dimensions as it is critical that a typical Brownian path collides with the time-dependent catalyst [10]. The situation for mutually catalytic branching is less clear and, depending on the time of day, you may be able to convince yourself that it should exist in any dimension, or only for $d \leq 3$, or only for $d \leq 2$.

PROBLEM 7. Construct a solution to $(\mathbf{MP})_{\mathbf{X}_0}^{\sigma, \gamma}$ in higher dimensions or prove they cannot exist for sufficiently high dimensions.

APPENDIX A: RANDOM WALK KERNELS

In this Appendix we gather together the results we need for the transition kernel of our continuous time random walk ${}^\varepsilon \xi$ on $\varepsilon \mathbb{Z}^2$ which jumps to a randomly chosen nearest neighbor with rate $2\varepsilon^{-2}\sigma^2$. One would have thought that references

containing proofs of Lemma 8 are commonplace but we could not locate one. Recall that

$${}^\varepsilon p_t(x) = \varepsilon^{-2} \Pi({}^\varepsilon \xi_t = x), \quad x \in \varepsilon \mathbb{Z}^2$$

and

$$p_t(x) = (2\pi t \sigma^2)^{-1} e^{-|x|^2/2\sigma^2 t}.$$

Let ${}^\varepsilon q_t(x)$ and $q_t(x)$ be the one-dimensional counterparts of ${}^\varepsilon p_t(x)$ and $p_t(x)$, respectively, so that ${}^\varepsilon p_t(x_1, x_2) = {}^\varepsilon q_t(x_1) {}^\varepsilon q_t(x_2)$ and $p_t(x_1, x_2) = q_t(x_1) q_t(x_2)$. Lemma 8 then is immediate from its one-dimensional version which we now prove.

LEMMA 59. (a) $\forall s > 0, \lim_{\varepsilon \rightarrow 0} \sup_{x \in \varepsilon \mathbb{Z}} |{}^\varepsilon q_s(x) - q_s(x)| = 0$.
 (b) *There is a universal constant $c_{A.1} < \infty$ such that, for all $\varepsilon > 0$,*

$$\sup_{s \geq 0, x \in \varepsilon \mathbb{Z}} {}^\varepsilon q_s(x) \sqrt{s} \sigma = \sup_{s \geq 0} {}^\varepsilon q_s(0) \sqrt{s} \sigma = c_{A.1}.$$

PROOF. The characteristic function of ${}^\varepsilon q_s(x)$ is given by

$$\begin{aligned} {}^\varepsilon \widehat{q}_s(\theta) &= e^{-\sigma^2 s / \varepsilon^2} \sum_n \frac{(\sigma^2 s / \varepsilon^2)^n}{n!} (\cos \theta \varepsilon)^n \\ &= \exp\left(-\sigma^2 s \left[\frac{1 - \cos \theta \varepsilon}{\varepsilon^2}\right]\right). \end{aligned}$$

Then by Fourier inversion (see [25], page 511) we have

$$(209) \quad {}^\varepsilon q_s(x) = (2\pi)^{-1} \int_{-\pi/\varepsilon}^{\pi/\varepsilon} \cos x\theta \exp\left(-\sigma^2 s \left[\frac{1 - \cos \theta \varepsilon}{\varepsilon^2}\right]\right) d\theta$$

and

$$(210) \quad q_s(x) = (2\pi)^{-1} \int_{-\infty}^{\infty} \cos x\theta \exp(-\sigma^2 s \theta^2 / 2) d\theta.$$

Let $K > 1$ and assume $\varepsilon < \frac{\pi}{K} s^{1/2}$. Then

$$\begin{aligned} &|{}^\varepsilon q_s(x) - q_s(x)| \\ &\leq \pi^{-1} \int_0^{Ks^{-1/2}} \left| \exp\left(\frac{-\sigma^2 s \theta^2}{2}\right) - \exp\left(-\sigma^2 s \left[\frac{1 - \cos(\theta \varepsilon)}{\varepsilon^2}\right]\right) \right| d\theta \\ &\quad + \pi^{-1} \left[\int_{Ks^{-1/2}}^{\infty} \exp\left(\frac{-\sigma^2 s \theta^2}{2}\right) d\theta \right. \\ &\quad \left. + \int_{Ks^{-1/2}}^{\pi/\varepsilon} \exp\left(-\sigma^2 s \left[\frac{1 - \cos(\theta \varepsilon)}{\varepsilon^2}\right]\right) d\theta \right] \\ &\equiv I_1 + I_2. \end{aligned}$$

Let $\eta > 0$ and define $c_0 = \inf_{|x| \leq \pi} (1 - \cos x)x^{-2} \in (0, \frac{1}{2}]$. Then

$$\begin{aligned}
 I_2 &\leq 2\pi^{-1} \int_{Ks^{-1/2}}^{\infty} \exp(-\sigma^2 s c_0 \theta^2) d\theta \\
 &\leq 2\pi^{-1} \int_{Ks^{1/2}}^{\infty} \frac{\theta}{K} s^{-1/2} e^{-\sigma^2 s c_0 \theta^2} d\theta \\
 (211) \quad &= (\pi \sigma^2 c_0)^{-1} s^{-1/2} e^{-\sigma^2 c_0 K^2} \\
 &\leq \eta s^{-1/2},
 \end{aligned}$$

where the last line is valid provided $K \geq K_0(\sigma, \eta)$. For I_1 use a second order Taylor expansion to write

$$\frac{1 - \cos \theta \varepsilon}{\varepsilon^2} = \frac{\theta^2}{2} \cos X_\theta \quad \text{for some } X_\theta \in (0, \theta \varepsilon),$$

and conclude that

$$\begin{aligned}
 I_1 &= \pi^{-1} \int_0^{Ks^{-1/2}} \left| \exp\left(\frac{-\sigma^2 s \theta^2}{2}\right) \left[1 - \exp\left(\frac{-\sigma^2 s \theta^2}{2} (\cos X_\theta - 1)\right) \right] \right| d\theta \\
 &\leq \pi^{-1} \int_0^{Ks^{-1/2}} \exp\left(\frac{-\sigma^2 s \theta^2}{2}\right) \left| \exp\left(\frac{\sigma^2 s \theta^4 \varepsilon^2}{4}\right) - 1 \right| d\theta.
 \end{aligned}$$

The elementary inequality $1 - \cos x \leq x^2/2$ is used in the last line.

For $0 \leq \theta \leq Ks^{-1/2}$, our assumed bound on ε gives

$$\sigma^2 s \theta^4 \varepsilon^2 \leq \sigma^2 s^{-1} K^4 \varepsilon^2 < \pi^2 \sigma^2 K^2,$$

and so

$$\exp(\sigma^2 s \theta^4 \varepsilon^2 / 4) - 1 \leq \exp(\pi^2 \sigma^2 K^2 / 4) \sigma^2 s \theta^4 \varepsilon^2 / 4.$$

This gives

$$\begin{aligned}
 I_1 &\leq \pi^{-1} \exp\left(\frac{\pi^2 \sigma^2 K^2}{4}\right) \int_0^{Ks^{-1/2}} \exp\left(\frac{-\sigma^2 s \theta^2}{2}\right) \sigma^2 s \theta^4 \varepsilon^2 / 4 d\theta \\
 &\leq \pi^{-1} \exp\left(\frac{\pi^2 \sigma^2 K^2}{4}\right) \frac{\varepsilon^2}{\sqrt{2} \sigma^2 s^{3/2}} \int_0^{\infty} e^{-u} u^{3/2} du \\
 &\leq c(K, \sigma) \varepsilon^2 s^{-3/2}.
 \end{aligned}$$

Combine this with (211) and set $K = K_0(\sigma, \eta)$ to see that

$$(212) \quad \sup_{x \in \mathbb{Z}} |{}^\varepsilon q_s(x) - q_s(x)| \leq \eta s^{-1/2} + c(K_0, \sigma) \varepsilon^2 s^{-3/2} \quad \text{for } \varepsilon < \frac{\pi}{K_0} s^{1/2}.$$

Statement (a) is immediate from the above.

The first equality in (b) is clear from (209). For the second, note that by (209)

$$\begin{aligned} \varepsilon q_s(0)\sqrt{s}\sigma &= \frac{\sqrt{s}\sigma}{2\pi} \int_{-\pi/\varepsilon}^{\pi/\varepsilon} \exp\left(-\sigma^2 s \left[\frac{1-\cos\theta\varepsilon}{\varepsilon^2}\right]\right) d\theta \\ &\leq \frac{\sqrt{s}\sigma}{2\pi} \int_{-\infty}^{\infty} \exp(-\sigma^2 s c_0 \theta^2) d\theta + \frac{\sqrt{s}\sigma}{\pi} \int_{\pi/2\varepsilon}^{\pi/\varepsilon} \exp\left(-\frac{\sigma^2 s}{\varepsilon^2}\right) d\theta \\ &\leq \frac{1}{\sqrt{\pi c_0}} + \frac{1}{2} \sup_{\sqrt{s}\sigma/\varepsilon > 0} \frac{\sqrt{s}\sigma}{\varepsilon} \exp\left(-\frac{\sigma^2 s}{\varepsilon^2}\right) \\ &= c_{A.1}, \end{aligned}$$

where $c_0 = \inf_{0 \leq u \leq \pi/2} \frac{1-\cos u}{u^2}$ and $c_{A.1}$ is a constant independent of σ and ε . \square

PROOF OF LEMMA 44. We may consider $x = (x_1, x_2) \in \varepsilon\mathbb{Z}^2$ such that $x_1 = |x_1| \geq |x_2|$. By scaling, Lemma 59(b) and (209),

$$\begin{aligned} &\int_0^\delta \varepsilon p_s(x_1, x_2) ds \\ &= \int_0^{\delta\varepsilon^{-2}} {}^1p_u(x_1\varepsilon^{-1}, x_2\varepsilon^{-1}) du \\ (213) \quad &\leq c_1(\sigma) \int_0^{\delta\varepsilon^{-2}} {}^1q_u(x_1\varepsilon^{-1})u^{-1/2} du \\ &= c_1(\sigma)\pi^{-1} \int_0^{\delta\varepsilon^{-2}} \int_0^\pi \cos(x_1\theta/\varepsilon) \exp(-\sigma^2(1-\cos\theta)u) d\theta u^{-1/2} du \\ &= c_2(\sigma) \int_0^\pi \cos(x_1\theta/\varepsilon) g_{\delta,\varepsilon}(\theta) d\theta, \end{aligned}$$

where $g_{\delta,\varepsilon}(\theta) = \int_0^{\delta\varepsilon^{-2}} \exp(-\sigma^2(1-\cos\theta)u)u^{-1/2} du$. Note that $g_{\delta,\varepsilon}$ is a decreasing function on $[0, \pi]$ and if $c(\theta) = \sigma\sqrt{1-\cos\theta}$, then

$$(214) \quad g_{\delta,\varepsilon}(\theta) = \int_0^{c(\theta)^2\delta\varepsilon^{-2}} e^{-v}v^{-1/2} dv c(\theta)^{-1} \leq c_3 \left[c(\theta)^{-1} \wedge \frac{\sqrt{\delta}}{\varepsilon} \right].$$

This implies

$$\begin{aligned} &\int_0^{(\varepsilon/x_1)\wedge\pi} \left| \cos\left(\frac{x_1\theta}{\varepsilon}\right) g_{\delta,\varepsilon}(\theta) \right| d\theta \\ (215) \quad &\leq c_3 \int_0^{(\varepsilon/x_1)\wedge(\varepsilon/\sqrt{\delta})} \frac{\sqrt{\delta}}{\varepsilon} d\theta + c_3 \int_{(\varepsilon/\sqrt{\delta})\wedge(\varepsilon/x_1)}^{\varepsilon/x_1} c(\theta)^{-1} \mathbf{1}(\theta \leq \pi) d\theta \\ &\leq c_4(\sigma) \left[1 \wedge \frac{\sqrt{\delta}}{x_1} + \int_{(\varepsilon/\sqrt{\delta})\wedge(\varepsilon/x_1)}^{\varepsilon/x_1} \theta^{-1} d\theta \right] \\ &\leq c_4(\sigma) \left[1 \wedge \frac{\sqrt{\delta}}{x_1} + \log^+\left(\frac{\sqrt{\delta}}{x_1}\right) \right]. \end{aligned}$$

An integration by parts shows that if $x_1 > \varepsilon/\pi$, then

$$\begin{aligned}
 & \int_{\varepsilon/x_1}^{\pi} \cos\left(\frac{x_1\theta}{\varepsilon}\right) g_{\delta,\varepsilon}(\theta) d\theta \\
 (216) \quad &= \frac{\varepsilon}{x_1} \sin\left(\frac{x_1\theta}{\varepsilon}\right) g_{\delta,\varepsilon}(\theta) \Big|_{\varepsilon/x_1}^{\pi} - \int_{\varepsilon/x_1}^{\pi} \frac{\varepsilon}{x_1} \sin\left(\frac{x_1\theta}{\varepsilon}\right) g'_{\delta,\varepsilon}(\theta) d\theta \\
 &\leq \frac{\varepsilon}{x_1} g_{\delta,\varepsilon}(\pi) + \frac{\varepsilon}{x_1} \left(g_{\delta,\varepsilon}\left(\frac{\varepsilon}{x_1}\right) - g_{\delta,\varepsilon}(\pi) \right) = \frac{\varepsilon}{x_1} g_{\delta,\varepsilon}\left(\frac{\varepsilon}{x_1}\right).
 \end{aligned}$$

In the last line we bounded the integrand in absolute value by $\frac{\varepsilon}{x_1}(-g'_{\delta,\varepsilon}(\theta))$. Now use (214) in (216) to conclude that, for $x_1 > \varepsilon/\pi$,

$$\begin{aligned}
 \int_{\varepsilon/x_1}^{\pi} \cos\left(\frac{x_1\theta}{\varepsilon}\right) g_{\delta,\varepsilon}(\theta) d\theta &\leq \frac{\varepsilon}{x_1} g_{\delta,\varepsilon}\left(\frac{\varepsilon}{x_1}\right) \\
 &\leq \frac{\varepsilon}{x_1} c_3 \left[c\left(\frac{\varepsilon}{x_1}\right)^{-1} \wedge \frac{\sqrt{\delta}}{\varepsilon} \right] \\
 &\leq c_5(\sigma) \frac{\varepsilon}{x_1} \frac{x_1 \wedge \sqrt{\delta}}{\varepsilon} = c_5(\sigma) \left(1 \wedge \frac{\sqrt{\delta}}{x_1} \right).
 \end{aligned}$$

Combine this with (215) in (213) to see that

$$\int_0^{\delta} \varepsilon p_s(x_1, x_2) ds \leq c_6(\sigma) \left[1 \wedge \frac{\sqrt{\delta}}{x_1} + \log^+ \frac{\sqrt{\delta}}{x_1} \right].$$

Recalling our assumption that $x_1 = |x_1| \geq |x_2|$, we see that the result follows. \square

PROOF OF LEMMA 34. By Lemma 59(b) and the fact that $|(x_1, x_2)| > s^{r/2} + \varepsilon^r$ implies $|x_i| > \frac{s^{r/2} + \varepsilon^r}{2}$ for $i = 1$ or 2 , the result follows from

$$(217) \quad \sup \left\{ s^{-1/2} \varepsilon q_s(x) : 0 < s, \varepsilon, |x| > \frac{s^{r/2} + \varepsilon^r}{2}, x \in \varepsilon\mathbb{Z} \right\} < \infty.$$

Another application of Lemma 59(b) shows that we need only consider $s \leq 1$. If τ_1 is the first jump time of the one-dimensional random walk ${}^\varepsilon\xi$ then, for $x \neq 0$,

$$\varepsilon q_s(x) \leq \varepsilon^{-1} P(\tau_1 < s) = \varepsilon^{-1} (1 - \exp(-\sigma^2 s \varepsilon^{-2})) \leq \sigma^2 s \varepsilon^{-3}$$

and so

$$\sup \{ s^{-1/2} \varepsilon q_s(x) : s^{1/6} \leq \varepsilon, x \neq 0 \} \leq \sigma^2.$$

These observations show that it now suffices to prove

$$(218) \quad \sup \left\{ s^{-1/2} \varepsilon q_s(x) : |x| > \frac{s^{r/2} + \varepsilon^r}{2}, 0 < \varepsilon \leq s^{1/6} \leq 1 \right\} < \infty.$$

To get bounds for larger values of s we first use some exponential bounds. Let S_n be a simple symmetric random walk on $\varepsilon\mathbb{Z}$ and let $N^\varepsilon(s)$ be a Poisson process with rate $\sigma^2\varepsilon^{-2}$ which is independent of $\{S_n\}$. Then we may take ${}^\varepsilon\xi(s) = S_{N^\varepsilon(s)}$ and a simple calculation shows that if $0 < \lambda \leq \varepsilon^{-1}$, then

$$E(e^{\lambda {}^\varepsilon\xi(s)}) = \exp(\sigma^2\varepsilon^{-2}s(\cosh \lambda\varepsilon - 1)) \leq e^{c_1s\lambda^2}$$

for some $c_1 = c_1(\sigma^2) > 0$. If $\lambda = \varepsilon^{-1} \wedge s^{-1/2}$ and $x \geq (\varepsilon^r + s^{r/2})/2$, then

$$\begin{aligned} {}^\varepsilon q_s(x) &\leq \varepsilon^{-1} P(e^{\lambda {}^\varepsilon\xi(s)} \geq e^{\lambda x}) \leq \varepsilon^{-1} \exp(-\lambda x + c_1s\lambda^2) \\ &\leq \varepsilon^{-1} \exp(-(\varepsilon^{-1} \wedge s^{-1/2})(\varepsilon^r + s^{r/2})/2 + c_1) \\ &\leq \varepsilon^{-1} \exp(-(\varepsilon \vee s^{1/2})^{r-1}/2 + c_1). \end{aligned}$$

By symmetry in x this gives

$$\begin{aligned} &\sup\left\{s^{-1/2} {}^\varepsilon q_s(x) : |x| \geq \frac{\varepsilon^r + s^{r/2}}{2}, 0 < s^9 \leq \varepsilon \leq s^{1/6} \leq 1\right\} \\ &\leq \sup\{s^{-1/2}\varepsilon^{-1} \exp(-(\varepsilon \vee s^{1/2})^{r-1}/2 + c_1) : 0 < s^9 \leq \varepsilon \leq s^{1/6} \leq 1\} \\ &\leq \sup\{s^{-19/2} \exp(-s^{(r-1)/6}/2 + c_1) : 0 < s \leq 1\} = c_2 < \infty. \end{aligned}$$

To obtain (218) it therefore now suffices to show

$$(219) \quad \sup\left\{s^{-1/2} {}^\varepsilon q_s(x) : |x| \geq \frac{\varepsilon^r + s^{r/2}}{2}, 0 < \varepsilon < s^9 \leq 1\right\} < \infty.$$

For this use (209) to see that

$$\begin{aligned} (220) \quad {}^\varepsilon q_s(x) &= \pi^{-1} \int_0^{\pi/\varepsilon} \cos x\theta \exp\left(-\sigma^2s \left[\frac{1 - \cos \theta\varepsilon}{\varepsilon^2}\right]\right) d\theta \\ &= \pi^{-1} \int_0^{s^{-1}} \cos x\theta \left[\exp\left(-\sigma^2s \left[\frac{1 - \cos \theta\varepsilon}{\varepsilon^2}\right]\right) - \exp\left(\frac{-\sigma^2s\theta^2}{2}\right) \right] d\theta \\ &\quad + \pi^{-1} \int_{s^{-1}}^{\pi/\varepsilon} \exp(-\varepsilon^{-2}\sigma^2s[1 - \cos \theta\varepsilon]) d\theta - \pi^{-1} \int_{s^{-1}}^\infty \exp\left(\frac{-\sigma^2s\theta^2}{2}\right) d\theta \\ &\quad + \pi^{-1} \int_0^\infty \cos \theta x \exp\left(\frac{-\sigma^2s\theta^2}{2}\right) d\theta \\ &\leq \pi^{-1} \left| \int_0^{s^{-1}} \cos x\theta \left[\exp\left(-\sigma^2s \left[\frac{1 - \cos \theta\varepsilon}{\varepsilon^2}\right]\right) - \exp\left(\frac{-\sigma^2s\theta^2}{2}\right) \right] d\theta \right| \\ &\quad + \pi^{-1} \int_{s^{-1}}^{\pi/\varepsilon} \exp(-\varepsilon^{-2}\sigma^2s[1 - \cos \theta\varepsilon]) d\theta + \pi^{-1} \int_{s^{-1}}^\infty \exp\left(\frac{-\sigma^2s\theta^2}{2}\right) d\theta \\ &\quad + \pi^{-1} \left| \int_0^\infty \cos \theta x \exp\left(\frac{-\sigma^2s\theta^2}{2}\right) d\theta \right| \\ &\equiv I + II + III + IV. \end{aligned}$$

By Fourier inversion we see that, for $|x| \geq s^{r/2}$ and $s \in (0, 1]$,

$$(221) \quad IV = p_s(x) \leq c(\sigma^2)s^{-1/2} \exp(-s^{r-1}/2\sigma^2) \leq c(\sigma^2)s^{1/2}.$$

Use the fact that $(1 - \cos \theta \varepsilon)\varepsilon^{-2} \geq c_2\theta^2$ for all $|\theta| \leq \pi/\varepsilon$ and some $c_2 \in (0, 1/2]$, to see that

$$(222) \quad \begin{aligned} II + III &\leq \int_{s^{-1}}^{\infty} \exp(-\sigma^2 c_2 s \theta^2) d\theta \\ &\leq \int_{s^{-1}}^{\infty} \exp(-\sigma^2 c_2 s \theta^2) \theta s d\theta \\ &= (2\sigma^2 c_2)^{-1} \exp(-\sigma^2 c_2 s^{-1}) \leq c(\sigma^2)s^{1/2}. \end{aligned}$$

To bound I , use Taylor's formula to write $1 - \cos \theta \varepsilon = \frac{\theta^2 \varepsilon^2}{2} - \frac{(\cos X)\theta^4 \varepsilon^4}{4!}$ for some $|X| < \theta \varepsilon$ and note that, for $0 \leq \theta \leq s^{-1}$ and $\varepsilon \leq s^9 \leq 1$,

$$(223) \quad \sigma^2 s \theta^4 \varepsilon^2 \leq \sigma^2.$$

Therefore, for $\varepsilon < s^9 \leq 1$,

$$(224) \quad \begin{aligned} I &\leq \int_0^{s^{-1}} \exp(-\sigma^2 s \theta^2/2) |\exp((\cos X)\theta^4 \sigma^2 s \varepsilon^2/24) - 1| d\theta \\ &\leq \int_0^{s^{-1}} c(\sigma^2) s \varepsilon^2 \theta^4 d\theta \quad [\text{by (223)}] \\ &\leq c(\sigma^2) s^{1/2}. \end{aligned}$$

We use the condition on ε and s in the last line. Now use (221), (222) and (224) in (220) to derive (219) and complete the proof. \square

APPENDIX B: INTEGRATION LEMMAS

LEMMA 60. *Let $p \in (0, 1)$ and*

$$I_{n,p}(s) = \int_{\mathbb{R}_+^n} \frac{1}{s + u_1} \prod_{i=1}^{n-1} \frac{1}{u_i + u_{i+1}} \frac{1}{u_n^p} du_1 \cdots du_n.$$

Then

$$I_{n,p}(s) = \left(\frac{\pi}{\sin((1-p)\pi)} \right)^n s^{-p} \quad \text{for all } n \in \mathbb{N}.$$

PROOF. Let $z = (u_n/u_{n-1})^{1-p}$ to see that $(u_n = u_{n-1}z^{1/(1-p)})$

$$\begin{aligned} \int_0^\infty \frac{1}{u_{n-1} + u_n} \frac{1}{u_n^p} du_n &= \int_0^\infty \frac{1}{u_{n-1}(1 + z^{1/(1-p)})} \frac{u_{n-1}z^{p/(1-p)}}{u_{n-1}^p z^{p/(1-p)}(1-p)} dz \\ &= \frac{u_{n-1}^{-p}}{1-p} \int_0^\infty \frac{dz}{1 + z^{1/(1-p)}} = \frac{\pi}{\sin(\pi(1-p))} u_{n-1}^{-p} \end{aligned}$$

by a standard residue calculation. This shows

$$\begin{aligned}
 I_{n,p}(s) &= \int_{\mathbb{R}_+^{n-1}} \frac{1}{s + u_1} \prod_1^{n-2} \frac{1}{u_i + u_{i+1}} \left[\int_0^\infty \frac{1}{u_{n-1} + u_n} \frac{1}{u_n^p} du_n \right] du_1 \cdots du_{n-1} \\
 &= \frac{\pi}{\sin(\pi(1-p))} I_{n-1,p}(s)
 \end{aligned}$$

and $I_{1,p}(s) = \frac{\pi}{\sin(\pi(1-p))} s^{-p}$. The obvious induction completes the proof. \square

COROLLARY 61. *Let $0 < p < 1$ and, for $s, T > 0$, define*

$$J_n(s, T) = \int_{\mathbb{R}_+^n} \frac{1}{s + u_1} \mathbf{1}(u_n \leq T) \prod_{i=1}^{n-1} \frac{1}{u_i + u_{i+1}} du_1 \cdots du_n.$$

Then there is a constant $c_{61}(p)$ such that

$$J_n(s, T) \leq c_{61}(p) \left(\frac{\pi}{\sin(\pi(1-p))} \right)^{n-1} \left(\frac{T}{s} \right)^p \quad \text{for all } n \in \mathbb{N}.$$

PROOF.

$$\int_0^T \frac{1}{u_{n-1} + u_n} du_n = \log \left(1 + \frac{T}{u_{n-1}} \right) \leq c_{61}(p) \left(\frac{T}{u_{n-1}} \right)^p$$

because $\log(1+x) \leq c_{61}x^p$ for all $x \geq 0$. Therefore, by Lemma 60,

$$\begin{aligned}
 J_n(s, T) &\leq c_{61} \int_{\mathbb{R}_+^{n-1}} \frac{1}{s + u_1} \prod_{i=1}^{n-2} \frac{1}{u_i + u_{i+1}} \left(\frac{T}{u_{n-1}} \right)^p du_1 \cdots du_{n-1} \\
 &= c_{61} T^p I_{n-1}(s) \leq c_{61} \left(\frac{\pi}{\sin((1-p)\pi)} \right)^{n-1} \left(\frac{T}{s} \right)^p. \quad \square
 \end{aligned}$$

COROLLARY 62. *Assume $h : (0, \infty) \rightarrow [0, \infty)$ satisfies $h(u) \leq c(1 + u^{-p})$ for $u \in [0, T]$ and some $p \in (0, 1)$. Then*

$$\begin{aligned}
 J_n(s, h) &\equiv \int_{\mathbb{R}_+^n} \frac{\mathbf{1}(u_n \leq T)}{s + u_1} \left(\prod_1^{n-1} \frac{1}{u_i + u_{i+1}} \right) h(u_n) du_1 \cdots du_n \\
 &\leq c c_{62}(p) \left(\frac{\pi}{\sin((1-p)\pi)} \right)^n s^{-p} (T^p + 1).
 \end{aligned}$$

PROOF. Immediate from the previous two results. \square

PROOF OF LEMMA 49. (a) Let $u = (1 - w)/(x - 1)$ in the integral defining ϕ_p to see that

$$\begin{aligned}
 \phi_p(x) &= \frac{x}{1 + (x - 1)^{-p}} \left[\int_0^{1/(x-1)} \frac{x - 1}{(x - 1)(1 + u)} (1 - (x - 1)u)^{-p} du \right. \\
 &\quad \left. + \int_0^{1/(x-1)} \frac{x - 1}{(x - 1)(1 + u)} (x - 1)^{-p} u^{-p} du \right] \\
 (225) \quad &= \frac{x}{(x - 1)^p + 1} \left[\int_0^{1/(x-1)} \frac{((x - 1)^{-1} - u)^{-p}}{1 + u} du \right. \\
 &\quad \left. + \int_0^{1/(x-1)} (1 + u)^{-1} u^{-p} du \right].
 \end{aligned}$$

If $x \geq 2$, then

$$\begin{aligned}
 \phi_p(x) &\leq \frac{2x}{(x - 1)^p + 1} \int_0^{1/(x-1)} u^{-p} du \\
 (226) \quad &= \left(\frac{2}{1 - p} \right) \frac{x}{x - 1 + (x - 1)^{1-p}} \leq \frac{2}{1 - p}.
 \end{aligned}$$

Assume now that $x \in (1, 2)$. If $R = (x - 1)^{-1} \geq 1$ and we set $w = R - u$, then

$$\begin{aligned}
 &\int_0^R (R - u)^{-p} (1 + u)^{-1} du \\
 &\leq \int_0^{R/2} u^{-p} (1 + u)^{-1} du + \int_{R/2}^R (R - u)^{-p} (1 + R - u)^{-1} du \\
 &= \int_0^{R/2} u^{-p} (1 + u)^{-1} du + \int_0^{R/2} w^{-p} (1 + w)^{-1} dw \\
 &\leq 2 \int_0^\infty u^{-p} (1 + u)^{-1} du.
 \end{aligned}$$

Use this together with the fact $(x - 1)^p + 1 \geq x - 1 + 1 = x$ for $x < 2$, to see that (225) implies

$$(227) \quad \phi_p(x) \leq 3 \int_0^\infty u^{-p} (1 + u)^{-1} du = \frac{3\pi}{\sin(1 - p)\pi},$$

the last by a standard contour integration. As $\frac{3\pi}{\sin(1-p)\pi} > \frac{2}{1-p}$, the result follows from (226) and (227).

(b) If $w = \frac{s_2}{s_1}$, then

$$\begin{aligned}
 K_2^{(p)}(s_0, s_1) &= \int_0^{s_1} (s_0 - s_2)^{-1} s_1^{-1} s_2^{-p} (1 + ((s_1 - s_2)/s_2)^{-p}) ds_2 \\
 &= s_1^{-1-p} \int_0^1 ((s_0/s_1) - w)^{-1} (w^{-p} + (1 - w)^{-p}) dw \\
 &\leq c_{49} s_1^{-p} s_0^{-1} (1 + ((s_0/s_1) - 1)^{-p}).
 \end{aligned}$$

In the last line we used (a). This gives the result for $n = 2$. Assume the result for $n \geq 2$. Then

$$\begin{aligned} K_{n+1}^{(p)}(s_0, s_1) &= \int_0^{s_1} K_n(s_1, s_2)(s_0 - s_2)^{-1} ds_2 \\ &\leq c_{49}^{n-1} s_1^{-1} \int_0^{s_1} (s_0 - s_2)^{-1} s_2^{-p} (1 + ((s_1 - s_2)/s_2)^{-p}) ds_2 \\ &= c_{49}^{n-1} K_2^{(p)}(s_0, s_1). \end{aligned}$$

Use the result derived for $n = 2$ to obtain the required bound for $n + 1$ and hence complete the induction. \square

LEMMA 63. Let $\{X_n\}$ be a sequence of nonnegative random variables on (Ω, \mathcal{F}, P) and let \mathcal{G} be a sub- σ -field of \mathcal{F} . Assume, for some $p > 1$, $\{E(X_n^p | \mathcal{G}) : n \in \mathbb{N}\}$ is bounded in probability and X_n converges in probability to X_∞ . Then

$$E(X_n | \mathcal{G}) \text{ converges in probability to } E(X_\infty | \mathcal{G}) < \infty \text{ a.s.}$$

PROOF. This may be shown by making the obvious changes in the standard proof of the unconditional version of this result. \square

Acknowledgments. Thanks go to Carl Mueller who suggested studying this model several years ago, and to an anonymous referee for a number of suggestions for improving the exposition.

REFERENCES

- [1] ATHREYA, S. and TRIBE, R. (2000). Uniqueness for a class of one-dimensional stochastic PDEs using moment duality. *Ann. Probab.* **28** 1711–1734.
- [2] BARLOW, M. T., EVANS, S. N. and PERKINS, E. A. (1991). Collision local times and measure-valued processes. *Canad. J. Math.* **43** 897–938.
- [3] BARLOW, M. T. and PERKINS, E. A. (1994). On the filtration of historical Brownian motion. *Ann. Probab.* **22** 1273–1294.
- [4] BASS, R. F. and PERKINS, E. A. (2003). Degenerate stochastic differential equations with Hölder continuous coefficients and super-Markov chains. *Trans. Amer. Math. Soc.* **335** 373–405.
- [5] BILLINGSLEY, P. (1986). *Probability and Measure*, 2nd ed. Wiley, New York.
- [6] COX, J. T., DAWSON, D. A. and GREVEN, A. (2002). Mutually catalytic super branching random walks: Large finite systems and renormalization analysis. Preprint.
- [7] COX, T., KLENKE, A. and PERKINS, E. A. (2000). Convergence to equilibrium and linear system duality. *CMS Conf. Proc.* **26** 41–66.
- [8] DAWSON, D. A., ETHERIDGE, A. M., FLEISCHMANN, K., MYTNIK, L., PERKINS, E. A. and XIONG, J. (2002). Mutually catalytic branching in the plane: Infinite measure states. *Electron. J. Probab.* **7**.
- [9] DAWSON, D. A. and FLEISCHMANN, K. (1995). Super-Brownian motions in higher dimensions with absolutely continuous measure states. *J. Theoret. Probab.* **8** 179–206.

- [10] DAWSON, D. A. and FLEISCHMANN, K. (1997). A continuous super-Brownian motion in a super-Brownian medium. *J. Theoret. Probab.* **10** 213–276.
- [11] DAWSON, D. A. and FLEISCHMANN, K. (1997). Longtime behavior of a branching process controlled by branching catalysts. *Stochastic Process. Appl.* **71** 241–257.
- [12] DAWSON, D. A., FLEISCHMANN, K., MYTNIK, L. T., PERKINS, E. A. and XIONG, J. (2002). Mutually catalytic branching in the plane: Uniqueness. *Ann. Inst. H. Poincaré Probab. Statist.* To appear.
- [13] DAWSON, D. A. and MARCH, P. (1995). Resolvent estimates for Fleming–Viot operators and uniqueness of solutions to related martingale problems. *J. Funct. Anal.* **132** 417–472.
- [14] DAWSON, D. A. and PERKINS, E. A. (1998). Long-time behaviour and coexistence in a mutually catalytic branching model. *Ann. Probab.* **26** 1088–1138.
- [15] DAWSON, D. A. and PERKINS, E. A. (1999). Measure-valued processes and renormalization of branching particle systems. In *Stochastic Partial Differential Equations: Six Perspectives* (R. Carmona and B. Rozovskii, eds.) 45–106. Amer. Math. Soc., Providence, RI.
- [16] DELMAS, J.-F. (1996). Super-mouvement brownien avec catalyse. *Stochastics Stochastics Rep.* **58** 303–347.
- [17] DELMAS, J.-F. and FLEISCHMANN, K. (2001). On the hot spots of a catalytic super-Brownian motion. *Probab. Theory Related Fields* **121** 389–421.
- [18] DONNELLY, P. and KURTZ, T. G. (1999). Particle representations for measure-valued population models. *Ann. Probab.* **27** 166–205.
- [19] EIGEN, M. (1971). Selforganization of matter and the evolution of biological macromolecules. *Die Naturwissenschaften* **58** 465–523.
- [20] EIGEN, M. (1982). *The Hypercycle: Principle of Natural Selforganization*. Springer, Berlin.
- [21] ETHERIDGE, A. M. and FLEISCHMANN, K. (1998). Persistence of a two-dimensional super-Brownian motion in a catalytic medium. *Probab. Theory Related Fields* **110** 1–12.
- [22] ETHIER, S. N. and KURTZ, T. G. (1986). *Markov Processes: Characterization and Convergence*. Wiley, New York.
- [23] EVANS, S. N. and PERKINS, E. A. (1994). Measure-valued branching diffusions with singular interactions. *Canad. J. Math.* **46** 120–168.
- [24] FELLER, W. (1968). *An Introduction to Probability Theory and Its Applications* **1**, 3rd ed. Wiley, New York.
- [25] FELLER, W. (1971). *An Introduction to Probability Theory and Its Applications* **2**, 2nd ed. Wiley, New York.
- [26] FLEISCHMANN, K. and KLENKE, A. (1999). Smooth density field of catalytic super-Brownian motion. *Ann. Appl. Probab.* **9** 298–318.
- [27] FLEISCHMANN, K. and KLENKE, A. (2000). The biodiversity of catalytic super-Brownian motion. *Ann. Appl. Probab.* **10** 1121–1136.
- [28] FLEISCHMANN, K. and XIONG, J. (2001). A cyclically catalytic super-Brownian motion. *Ann. Probab.* **29** 820–861.
- [29] JACOD, J. and SHIRYAEV, A. N. (1987). *Limit Theorems for Stochastic Processes*. Springer, Berlin.
- [30] JAKUBOWSKI, A. (1986). On the Skorohod topology. *Ann. Inst. H. Poincaré Ser. B* **22** 263–285.
- [31] KONNO, N. and SHIGA, T. (1988). Stochastic differential equations for some measure-valued diffusions. *Probab. Theory Related Fields* **79** 201–225.
- [32] MEYER, P.-A. (1966). *Probability and Potentials*. Blaisdell, Toronto.
- [33] MITOMA, I. (1985). An ∞ -dimensional inhomogeneous Langevin equation. *J. Funct. Anal.* **61** 342–359.
- [34] MYTNIK, L. (1996). Superprocesses in random environments. *Ann. Probab.* **24** 1953–1978.
- [35] MYTNIK, L. (1998). Uniqueness for a mutually catalytic branching model. *Probab. Theory Related Fields* **112** 245–253.

- [36] PERKINS, E. A. (1995). On the martingale problem for interactive measure-valued branching diffusions. *Mem. Amer. Math. Soc.* **115** 1–89.
- [37] PERKINS, E. A. (2000). Dawson–Watanabe superprocesses and measure-valued diffusions. *École d'Été de Probabilités de Saint-Flour 1999. Lecture Notes in Math.* **1781** 125–329. Springer, New York.
- [38] REIMERS, M. (1989). One-dimensional stochastic partial differential equations and the branching measure diffusion. *Probab. Theory Related Fields* **81** 319–340.
- [39] REVUZ, D. and YOR, M. (1991). *Continuous Martingales and Brownian Motion*. Springer, Berlin.
- [40] RUDIN, W. (1974). *Real and Complex Analysis*, 2nd ed. McGraw-Hill, New York.
- [41] SHIGA, T. (1988). Stepping stone models in population genetics and population dynamics. In *Stochastic Processes in Physics and Engineering* (S. Albeverio et al., eds.) 345–355. Reidel, Dordrecht.
- [42] SHIGA, T. (1994). Two contrasting properties of solutions for one-dimensional stochastic pde's. *Canad. J. Math.* **46** 415–437.
- [43] SHIGA, T. and SHIMIZU, A. (1980). Infinite-dimensional stochastic differential equations and their applications. *J. Math. Kyoto Univ.* **20** 395–416.
- [44] SPITZER, F. (1964). *Principles of Random Walk*. Van Nostrand, Princeton, NJ.
- [45] WALSH, J. B. (1986). An introduction to stochastic partial differential equations. *École d'Été de Probabilités de Saint-Flour XIV. Lecture Notes in Math.* **1180** 266–439. Springer, Berlin.

D. A. DAWSON
 SCHOOL OF MATHEMATICS
 AND STATISTICS
 CARLETON UNIVERSITY
 OTTAWA
 CANADA K1S 5B6
 E-MAIL: ddawson@math.carleton.ca

K. FLEISCHMANN
 WEIERSTRASS INSTITUTE FOR
 APPLIED ANALYSIS AND STOCHASTICS
 MOHRENSTR. 39
 D-10117 BERLIN
 GERMANY
 E-MAIL: fleischmann@wias-berlin.de

E. A. PERKINS
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF BRITISH COLUMBIA
 1984 MATHEMATICS ROAD
 VANCOUVER
 BRITISH COLUMBIA
 CANADA V6T 1Z2
 E-MAIL: perkins@math.ubc.ca

A. M. ETHERIDGE
 DEPARTMENT OF STATISTICS
 UNIVERSITY OF OXFORD
 1 SOUTH PARKS ROAD
 OXFORD OX1 3TG
 UNITED KINGDOM
 E-MAIL: etheridg@stats.ox.ac.uk

L. MYTNIK
 TECHNION—ISRAEL INSTITUTE
 OF TECHNOLOGY
 HAIFA 32000
 ISRAEL
 E-MAIL: leonid@ie.technion.ac.il

J. XIONG
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF TENNESSEE
 KNOXVILLE, TENNESSEE 37996–1300
 E-MAIL: jxiong@math.utk.edu