

# ANALYSIS & PDE

Volume 9

No. 3

2016

YAKINE BAHRI

**ASYMPTOTIC STABILITY IN ENERGY SPACE FOR DARK  
SOLITONS  
OF THE LANDAU-LIFSHITZ EQUATION**

# ASYMPTOTIC STABILITY IN ENERGY SPACE FOR DARK SOLITONS OF THE LANDAU–LIFSHITZ EQUATION

YAKINE BAHRI

We prove the asymptotic stability in energy space of nonzero speed solitons for the one-dimensional Landau–Lifshitz equation with an easy-plane anisotropy

$$\partial_t m + m \times (\partial_{xx} m - m_3 e_3) = 0$$

for a map  $m = (m_1, m_2, m_3) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{S}^2$ , where  $e_3 = (0, 0, 1)$ . More precisely, we show that any solution corresponding to an initial datum close to a soliton with nonzero speed is weakly convergent in energy space as time goes to infinity to a soliton with a possible different nonzero speed, up to the invariances of the equation. Our analysis relies on the ideas developed by Martel and Merle for the generalized Korteweg–de Vries equations. We use the Madelung transform to study the problem in the hydrodynamical framework. In this framework, we rely on the orbital stability of the solitons and the weak continuity of the flow in order to construct a limit profile. We next derive a monotonicity formula for the momentum, which gives the localization of the limit profile. Its smoothness and exponential decay then follow from a smoothing result for the localized solutions of the Schrödinger equations. Finally, we prove a Liouville type theorem, which shows that only the solitons enjoy these properties in their neighbourhoods.

## 1. Introduction

We consider the one-dimensional Landau–Lifshitz equation

$$\partial_t m + m \times (\partial_{xx} m + \lambda m_3 e_3) = 0 \tag{LL}$$

for a map  $m = (m_1, m_2, m_3) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{S}^2$ , where  $e_3 = (0, 0, 1)$  and  $\lambda \in \mathbb{R}$ . This equation was introduced by Landau and Lifshitz [1935]. It describes the dynamics of magnetization in a one-dimensional ferromagnetic material, for example in CsNiF<sub>3</sub> or TMNC (see, e.g., [Kosevich et al. 1990; Hubert and Schäfer 1998] and the references therein). The parameter  $\lambda$  accounts for the anisotropy of the material. The choices  $\lambda > 0$  and  $\lambda < 0$  correspond respectively to an easy-axis and an easy-plane anisotropy. In the isotropic case  $\lambda = 0$ , the equation is exactly the one-dimensional Schrödinger map equation, which has been intensively studied (see, e.g., [Guo and Ding 2008; Jerrard and Smets 2012]). In this paper, we study the Landau–Lifshitz equation with an easy-plane anisotropy ( $\lambda < 0$ ). Performing, if necessary, a suitable scaling argument on the map  $m$ , we assume from now on that  $\lambda = -1$ . Our main goal is to prove the asymptotic stability for the solitons of this equation (see [Theorem 1.1](#) below).

*MSC2010:* 35Q51, 35Q60, 37K40.

*Keywords:* asymptotic stability, solitons, Landau–Lifshitz equation, travelling waves.

The Landau–Lifshitz equation is Hamiltonian. Its Hamiltonian, the so-called Landau–Lifshitz energy, is given by

$$E(m) := \frac{1}{2} \int_{\mathbb{R}} (|\partial_x m|^2 + m_3^2).$$

In the sequel, we restrict our attention to the Hamiltonian framework in which the solutions  $m$  to (LL) have finite Landau–Lifshitz energy, i.e., belong to energy space

$$\mathcal{E}(\mathbb{R}) := \{v : \mathbb{R} \rightarrow \mathbb{S}^2 \mid v' \in L^2(\mathbb{R}) \text{ and } v_3 \in L^2(\mathbb{R})\}.$$

A soliton with speed  $c$  is a travelling-wave solution of (LL) having the form

$$m(x, t) := u(x - ct).$$

Its profile  $u$  is a solution to the ordinary differential equation

$$u'' + |u'|^2 u + u_3^2 u - u_3 e_3 + cu \times u' = 0. \tag{TWE}$$

The solutions of this equation are explicit. When  $|c| \geq 1$ , the only solutions with finite Landau–Lifshitz energy are the constant vectors in  $\mathbb{S}^1 \times \{0\}$ . In contrast, when  $|c| < 1$ , there exist nonconstant solutions  $u_c$  to (TWE), which are given by the formulae

$$[u_c]_1(x) = \frac{c}{\cosh((1-c^2)^{1/2}x)}, \quad [u_c]_2(x) = \tanh((1-c^2)^{1/2}x), \quad [u_c]_3(x) = \frac{(1-c^2)^{1/2}}{\cosh((1-c^2)^{1/2}x)},$$

up to the invariances of the problem, i.e., translations, rotations around the axis  $x_3$  and orthogonal symmetries with respect to the plane  $x_3 = 0$  (see [de Laire 2014] for more details).

Our goal is to study the asymptotic behaviour for solutions of (LL) which are initially close to a soliton in energy space. We endow  $\mathcal{E}(\mathbb{R})$  with the metric structure corresponding to the distance introduced by de Laire and Gravejat [2015],

$$d_{\mathcal{E}}(f, g) := |\check{f}(0) - \check{g}(0)| + \|f' - g'\|_{L^2(\mathbb{R})} + \|f_3 - g_3\|_{L^2(\mathbb{R})},$$

where  $f = (f_1, f_2, f_3)$  and  $\check{f} = f_1 + if_2$  (and similarly for  $g$ ). The Cauchy problem and the orbital stability of the travelling waves have been solved by de Laire and Gravejat [2015]. We are concerned with the asymptotic stability of travelling waves. The following theorem is our main result.

**Theorem 1.1.** *Let  $c \in (-1, 1) \setminus \{0\}$ . There exists a positive number  $\delta_c$ , depending only on  $c$ , such that, if*

$$d_{\mathcal{E}}(m^0, u_c) \leq \delta_c,$$

*then there exist a number  $c^* \in (-1, 1) \setminus \{0\}$ , and two functions  $b \in C^1(\mathbb{R}, \mathbb{R})$  and  $\theta \in C^1(\mathbb{R}, \mathbb{R})$  such that*

$$b'(t) \rightarrow c^* \quad \text{and} \quad \theta'(t) \rightarrow 0$$

*as  $t \rightarrow +\infty$ , and for which the map*

$$m_{\theta} := (\cos(\theta)m_1 - \sin(\theta)m_2, \sin(\theta)m_1 + \cos(\theta)m_2, m_3),$$

satisfies the convergences

$$\begin{aligned}\partial_x m_{\theta(t)}(\cdot + b(t), t) &\rightharpoonup \partial_x u_{c^*} && \text{in } L^2(\mathbb{R}), \\ m_{\theta(t)}(\cdot + b(t), t) &\rightarrow u_{c^*} && \text{in } L^\infty_{\text{loc}}(\mathbb{R}), \\ m_3(\cdot + b(t), t) &\rightharpoonup [u_{c^*}]_3 && \text{in } L^2(\mathbb{R})\end{aligned}$$

as  $t \rightarrow +\infty$ .

**Remarks.** (i) Note that the case  $c = 0$ —that is, black solitons—is excluded from the statement of [Theorem 1.1](#). In this case, the map  $\check{u}_0$  vanishes and we cannot apply the Madelung transform and the subsequent arguments. Orbital and asymptotic stability remain open problems for this case. Note that, to our knowledge, there is currently no available proof of the local well-posedness of [\(LL\)](#) in energy space, when  $u_0$  vanishes and so the hydrodynamical framework can no longer be used.

(ii) Here, we state a weak convergence result and not a local strong convergence one, like the results given by Martel and Merle [\[2008a; 2008b\]](#) for the Korteweg–de Vries equation. In their situation, they can use two monotonicity formulae for the  $L^2$  norm and the energy. This heuristically originates in the property that dispersion has negative speed in the context of the Korteweg–de Vries equation. In contrast, the possible group velocities for the dispersion of the Landau–Lifshitz equation are given by

$$v_g(k) = \pm \frac{1+2k^2}{\sqrt{1+k^2}},$$

where  $k$  is the wave number. Dispersion has both negative and positive speeds. A monotonicity formula remains for the momentum due to the existence of a gap in the possible group velocities, which satisfy the condition  $|v_g(k)| \geq 1$ . However, there is no evidence that one can establish a monotonicity formula for the energy.

Similar results were stated by Soffer and Weinstein [\[1989; 1990; 1992\]](#). They provided the asymptotic stability of ground states for the nonlinear Schrödinger equation with a potential in a regime for which the nonlinear ground-state is a close continuation of the linear one. They rely on dispersive estimates for the linearized equation around the ground state in suitable weighted spaces, and they apply a fixed point argument. This strategy was successfully extended in particular by Buslaev, Perelman, C. Sulem and Cuccagna to the nonlinear Schrödinger equations without potential (see, e.g., [\[Buslaev and Perelman 1993; 1995; Buslaev and Sulem 2003; Cuccagna 2001\]](#)) and with a potential (see, e.g., [\[Gang and Sigal 2007\]](#)). We refer to the detailed historical survey by Cuccagna [\[2003\]](#) for more details. Later, Cuccagna [\[2011\]](#) proved a stronger result for the ground state satisfying the sufficient conditions for orbital stability of M. Weinstein, for seemingly generic nonlinear Schrödinger equation which has a smooth short range nonlinearity with the presence of a very short range and smooth linear potential. In addition, asymptotic stability in spaces of exponentially localized perturbations was studied by Pego and Weinstein [\[1994\]](#) (see also [\[Mizumachi 2001\]](#) for perturbations with algebraic decay).

Our strategy for establishing the asymptotic stability result in [Theorem 1.1](#) is reminiscent of ideas developed by Martel and Merle [[2006](#); [2008a](#); [2008b](#)] for the Korteweg–de Vries equation, and successfully adapted by Béthuel, Gravejat and Smets in [[Béthuel et al. 2014](#)] for the Gross–Pitaevskii equation.

The main steps of the proof are similar to the ones for the Gross–Pitaevskii equation in [[Béthuel et al. 2015](#)]. Indeed, the solitons of the Landau–Lifshitz equation share many properties with the solitons of the Gross–Pitaevskii equation. In fact, the stereographic variable  $\psi$  defined by

$$\psi = \frac{u_1 + iu_2}{1 + u_3}$$

satisfies the equation

$$\partial_{xx}\psi + \frac{1 - |\psi|^2}{1 + |\psi|^2}\psi - ic\partial_x\psi = \frac{2\bar{\psi}}{1 + |\psi|^2}(\partial_x\psi)^2,$$

which can be seen as a perturbation of the equation for the travelling waves of the Gross–Pitaevskii equation, namely

$$\partial_{xx}\Psi + (1 - |\Psi|^2)\Psi - ic\partial_x\Psi = 0.$$

However, the analysis of the Landau–Lifshitz equation is much more difficult. Indeed, we rely on a Hasimoto like transform in order to relate the Landau–Lifshitz equation with a nonlinear Schrödinger equation. Doing so, we lose some regularity. We have to deal with a nonlinear equation at the  $L^2$ -level and not at the  $H^1$ -level as in the case of the Gross–Pitaevskii equation. This leads to important technical difficulties.

Returning to the proof of [Theorem 1.1](#), we first translate the problem into the hydrodynamical formulation. Then, we prove the asymptotic stability in that framework. In fact, we begin by refining the orbital stability. Next, we construct a limit profile, which is smooth and localized. For the proof of the exponential decay of the limit profile, we cannot rely on the Sobolev embedding  $H^1$  into  $L^\infty$  as was done in [[Béthuel et al. 2015](#)]. We use instead the results of Kenig, Ponce and Vega in [[Kenig et al. 2003](#)], and the Gagliardo–Nirenberg inequality (see the proof of [Proposition 2.9](#) for more details). We also have to deal with the weak continuity of the flow in order to construct the limit profile. For the Gross–Pitaevskii equation, this property relies on the uniqueness in a weaker space (see [[Béthuel et al. 2015](#)]). There is no similar result at the  $L^2$ -level. Instead, we use the Kato smoothing effect. The asymptotic stability in the hydrodynamical variables then follows from a Liouville type theorem. It shows that the only smooth and localized solutions in the neighbourhood of the solitons are the solitons. Finally, we deduce the asymptotic stability in the original setting from the result in the hydrodynamical framework.

In [Section 2](#) below, we explain the main tools and different steps for the proof. First, we introduce the hydrodynamical framework. Then, we state the orbital stability of the solitons under a new orthogonality condition. Next, we sketch the proof of the asymptotic stability for the hydrodynamical system and we state the main propositions. We finally complete the proof of [Theorem 1.1](#).

In [Sections 3 to 5](#), we give the proofs of the results stated in [Section 2](#). In [Section 3](#), we deal with the orbital stability in the hydrodynamical framework. In [Section 4](#), we prove the localization and the smoothness of the limit profile. In the last section, we prove a Liouville type theorem. In a separate appendix, we show some facts used in the proofs, in particular the weak continuity of the (HLL) flow.

## 2. Main steps for the proof of Theorem 1.1

**The hydrodynamical framework.** We introduce the map  $\check{m} := m_1 + im_2$ . Since  $m_3$  belongs to  $H^1(\mathbb{R})$ , it follows from the Sobolev embedding theorem that

$$|\check{m}(x)| = (1 - m_3^2(x))^{1/2} \rightarrow 1$$

as  $x \rightarrow \pm\infty$ . As a consequence, the Landau–Lifshitz equation shares many properties with the Gross–Pitaevskii equation (see, e.g., [Béthuel et al. 2008]). One of these properties is the existence of a hydrodynamical framework for the Landau–Lifshitz equation. In terms of the maps  $\check{m}$  and  $m_3$ , this equation may be written as

$$\begin{cases} i\partial_t \check{m} - m_3 \partial_{xx} \check{m} + \check{m} \partial_{xx} m_3 - \check{m} m_3 = 0, \\ \partial_t m_3 + \partial_x \langle i\check{m}, \partial_x \check{m} \rangle_{\mathbb{C}} = 0. \end{cases}$$

When the map  $\check{m}$  does not vanish, one can write it as  $\check{m} = (1 - m_3^2)^{1/2} \exp i\varphi$ . The hydrodynamical variables  $v := m_3$  and  $w := \partial_x \varphi$  satisfy the system

$$\begin{cases} \partial_t v = \partial_x ((v^2 - 1)w), \\ \partial_t w = \partial_x \left( \frac{\partial_{xx} v}{1 - v^2} + v \frac{(\partial_x v)^2}{(1 - v^2)^2} + v(w^2 - 1) \right). \end{cases} \quad (\text{HLL})$$

This system is similar to the hydrodynamical Gross–Pitaevskii equation (see, e.g., [Béthuel et al. 2015]).<sup>1</sup> We first study the asymptotic stability in the hydrodynamical framework.

In this framework, the Landau–Lifshitz energy is expressed as

$$E(\mathfrak{v}) := \int_{\mathbb{R}} e(\mathfrak{v}) := \frac{1}{2} \int_{\mathbb{R}} \left( \frac{(v')^2}{1 - v^2} + (1 - v^2)w^2 + v^2 \right), \quad (2-1)$$

where  $\mathfrak{v} := (v, w)$  denotes the hydrodynamical pair. The momentum  $P$ , defined by

$$P(\mathfrak{v}) := \int_{\mathbb{R}} vw, \quad (2-2)$$

is also conserved by the Landau–Lifshitz flow. The momentum  $P$  and the Landau–Lifshitz energy  $E$  play an important role in the study of the asymptotic stability of the solitons. When  $c \neq 0$ , the function  $\check{v}_c$  does not vanish. The hydrodynamical pair  $Q_c := (v_c, w_c)$  is given by

$$v_c(x) = \frac{(1 - c^2)^{1/2}}{\cosh((1 - c^2)^{1/2}x)} \quad \text{and} \quad w_c(x) = \frac{cv_c(x)}{1 - v_c(x)^2} = \frac{c(1 - c^2)^{1/2} \cosh((1 - c^2)^{1/2}x)}{\sinh((1 - c^2)^{1/2}x)^2 + c^2}. \quad (2-3)$$

The only invariances of (HLL) are translations and the opposite map  $(v, w) \mapsto (-v, -w)$ . We restrict our attention to the translation invariances. All the analysis developed below applies when the opposite map is also taken into account. For  $a \in \mathbb{R}$ , we define

$$Q_{c,a}(x) := Q_c(x - a) := (v_c(x - a), w_c(x - a)),$$

<sup>1</sup>The hydrodynamical terminology originates in the fact that the hydrodynamical Gross–Pitaevskii equation is similar to the Euler equation for an irrotational fluid (see, e.g., [Béthuel et al. 2014]).

a nonconstant soliton with speed  $c$ . We also set

$$\mathcal{NV}(\mathbb{R}) := \left\{ \mathbf{v} = (v, w) \in H^1(\mathbb{R}) \times L^2(\mathbb{R}) \mid \max_{\mathbb{R}} |v| < 1 \right\}.$$

This nonvanishing space is endowed in the sequel with the metric structure provided by the norm

$$\|\mathbf{v}\|_{H^1 \times L^2} := \left( \|v\|_{H^1}^2 + \|w\|_{L^2}^2 \right)^{1/2}.$$

**Orbital stability.** A perturbation of a soliton is provided by another soliton with a slightly different speed. This property follows from the existence of a continuum of solitons with different speeds. A solution corresponding to such a perturbation at initial time diverges from the soliton due to the different speeds of propagation, so that the standard notion of stability does not apply to solitons. The notion of orbital stability is tailored to deal with such situations. The orbital stability theorem below shows that a perturbation of a soliton at initial time remains a perturbation of the soliton, up to translations, for all time.

The following theorem is a variant of the result by de Laire and Gravejat [2015] concerning sums of solitons. It is useful for the proof of the asymptotic stability.

**Theorem 2.1.** *Let  $c \in (-1, 1) \setminus \{0\}$ . There exists a positive number  $\alpha_c$ , depending only on  $c$ , with the following properties. Given any  $(v_0, w_0) \in X(\mathbb{R}) := H^1(\mathbb{R}) \times L^2(\mathbb{R})$  such that*

$$\alpha_0 := \|(v_0, w_0) - Q_{c,a}\|_{X(\mathbb{R})} \leq \alpha_c \tag{2-4}$$

*for some  $a \in \mathbb{R}$ , there exist a unique global solution  $(v, w) \in C^0(\mathbb{R}, \mathcal{NV}(\mathbb{R}))$  to (HLL) with initial datum  $(v_0, w_0)$ , and two maps  $c \in C^1(\mathbb{R}, (-1, 1) \setminus \{0\})$  and  $a \in C^1(\mathbb{R}, \mathbb{R})$  such that the function  $\varepsilon$  defined by*

$$\varepsilon(\cdot, t) := \left( v(\cdot + a(t), t), w(\cdot + a(t), t) \right) - Q_{c(t)} \tag{2-5}$$

*satisfies the orthogonality conditions*

$$\langle \varepsilon(\cdot, t), \partial_x Q_{c(t)} \rangle_{L^2(\mathbb{R})^2} = \langle \varepsilon(\cdot, t), \chi_{c(t)} \rangle_{L^2(\mathbb{R})^2} = 0 \tag{2-6}$$

*for any  $t \in \mathbb{R}$ . Moreover, there exist two positive numbers  $\sigma_c$  and  $A_c$ , depending only and continuously on  $c$ , such that*

$$\max_{x \in \mathbb{R}} v(x, t) \leq 1 - \sigma_c, \tag{2-7}$$

$$\|\varepsilon(\cdot, t)\|_{X(\mathbb{R})} + |c(t) - c| \leq A_c \alpha^0, \tag{2-8}$$

$$|c'(t)| + |a'(t) - c(t)| \leq A_c \|\varepsilon(\cdot, t)\|_{X(\mathbb{R})}, \tag{2-9}$$

*for any  $t \in \mathbb{R}$ .*

**Remark.** In this statement, the function  $\chi_c$  is a normalized eigenfunction associated to the unique negative eigenvalue of the linear operator

$$\mathcal{H}_c := E''(Q_c) + cP''(Q_c).$$

The operator  $\mathcal{H}_c$  is self-adjoint on  $L^2(\mathbb{R}) \times L^2(\mathbb{R})$ , with domain  $\text{Dom}(\mathcal{H}_c) := H^2(\mathbb{R}) \times L^2(\mathbb{R})$  (see (A-42) for its explicit formula). It has a unique negative simple eigenvalue  $-\tilde{\lambda}_c$ , and its kernel is given by

$$\text{Ker}(\mathcal{H}_c) = \text{Span}(\partial_x Q_c). \tag{2-10}$$

Our statement of orbital stability relies on a different decomposition from that proposed by Grillakis, Shatah and Strauss in [Grillakis et al. 1987]. This modification is related to the proof of asymptotic stability. A key ingredient in the proof is the coercivity of the quadratic form  $G_c$ , which is defined in (2-46), under a suitable orthogonality condition. In case we use the orthogonality conditions in [Grillakis et al. 1987], the corresponding orthogonality condition for  $G_c$  is provided by the function  $v_c^{-1} S \partial_c Q_c$  (see (2-40) for the definition of  $S$ ), which does not belong to  $L^2(\mathbb{R})$ . In order to bypass this difficulty, we use the second orthogonality condition in (2-6) for which the corresponding orthogonality condition for  $G_c$  is given by the function  $v_c^{-1} S \chi_c$ , which does belong to  $L^2(\mathbb{R})$  (see the appendix for more details). This alternative decomposition is inspired by the one used by Martel and Merle [2008a].

Concerning the proof of Theorem 2.1, we first establish an orbital stability theorem with the classical decomposition of Grillakis, Shatah and Strauss [Grillakis et al. 1987]. This appears as a particular case of the orbital stability theorem in [de Laire and Gravejat 2015] for sums of solitons. We next show that if we have orbital stability for some decomposition and orthogonality conditions, then we also have it for different decomposition and orthogonality conditions (see Section 2 for the detailed proof of Theorem 2.1).

**Asymptotic stability for the hydrodynamical variables.** The following theorem shows the asymptotic stability result in the hydrodynamical framework.

**Theorem 2.2.** *Let  $c \in (-1, 1) \setminus \{0\}$ . There exists a positive constant  $\beta_c \leq \alpha_c$ , depending only on  $c$ , with the following properties. Given any  $(v_0, w_0) \in X(\mathbb{R})$  such that*

$$\|(v_0, w_0) - Q_{c,a}\|_{X(\mathbb{R})} \leq \beta_c,$$

*for some  $a \in \mathbb{R}$ , there exist a number  $c^* \in (-1, 1) \setminus \{0\}$  and a map  $b \in C^1(\mathbb{R}, \mathbb{R})$  such that the unique global solution  $(v, w) \in C^0(\mathbb{R}, \mathcal{NV}(\mathbb{R}))$  to (HLL) with initial datum  $(v_0, w_0)$  satisfies*

$$(v(\cdot + b(t), t), w(\cdot + b(t), t)) \rightharpoonup Q_{c^*} \quad \text{in } X(\mathbb{R}), \tag{2-11}$$

*and*

$$b'(t) \rightarrow c^*$$

*as  $t \rightarrow +\infty$ .*

Theorem 2.2 establishes a convergence to some orbit of the soliton. This result is stronger than the one given by Theorem 2.1 which only shows that the solution stays close to that orbit.

In the next subsections, we explain the main ideas of the proof, which follows the strategy developed by Martel and Merle [2008a; 2008b] for the Korteweg–de Vries equation.

*Construction of a limit profile.* Let  $\mathfrak{c} \in (-1, 1) \setminus \{0\}$ , and let  $(v_0, w_0) \in X(\mathbb{R})$  be any pair satisfying the assumptions of [Theorem 2.2](#). Since  $\beta_{\mathfrak{c}} \leq \alpha_{\mathfrak{c}}$  in the assumptions of [Theorem 2.2](#), we deduce from [Theorem 2.1](#) that the unique solution  $(v, w)$  to [\(HLL\)](#) with initial datum  $(v_0, w_0)$  is global.

We take an arbitrary sequence of times  $(t_n)_{n \in \mathbb{N}}$  tending to  $+\infty$ . In view of [\(2-8\)](#) and [\(2-9\)](#), we may assume, up to a subsequence, that there exist a limit perturbation  $\varepsilon_0^* \in X(\mathbb{R})$  and a limit speed  $c_0^* \in [-1, 1]$  such that

$$\varepsilon(\cdot, t_n) = (v(\cdot + a(t_n), t_n), w(\cdot + a(t_n), t_n)) - Q_{c(t_n)}) \rightharpoonup \varepsilon_0^* \quad \text{in } X(\mathbb{R}), \tag{2-12}$$

and

$$c(t_n) \rightarrow c_0^* \tag{2-13}$$

as  $n \rightarrow +\infty$ . Our main goal is to show that

$$\varepsilon_0^* \equiv 0$$

(see [Corollary 2.15](#)). For that, we establish smoothness and rigidity properties for the solution of [\(HLL\)](#) with the initial datum  $Q_{c_0^*} + \varepsilon_0^*$ .

First, we require the constant  $\beta_{\mathfrak{c}}$  to be sufficiently small so that, when the number  $\alpha^0$  which appears in [Theorem 2.1](#) satisfies  $\alpha^0 \leq \beta_{\mathfrak{c}}$ , then we infer from [\(2-8\)](#) and [\(2-9\)](#) that

$$\min\{c(t)^2, a'(t)^2\} \geq \frac{\mathfrak{c}^2}{2}, \quad \max\{c(t)^2, a'(t)^2\} \leq 1 + \frac{\mathfrak{c}^2}{2}, \tag{2-14}$$

and

$$\|v_{\mathfrak{c}}(\cdot) - v(\cdot + a(t), t)\|_{L^\infty(\mathbb{R})} \leq \min\left\{\frac{\mathfrak{c}^2}{4}, \frac{1 - \mathfrak{c}^2}{16}\right\} \tag{2-15}$$

for any  $t \in \mathbb{R}$ . This yields, in particular, that  $c_0^* \in (-1, 1) \setminus \{0\}$ , and then, that  $Q_{c_0^*}$  is well-defined and different from the black soliton.

By [\(2-8\)](#), we also have

$$|c_0^* - \mathfrak{c}| \leq A_{\mathfrak{c}}\beta_{\mathfrak{c}}, \tag{2-16}$$

and, applying again [\(2-8\)](#) well as [\(2-12\)](#) and the weak lower semicontinuity of the norm, we also know that the function

$$(v_0^*, w_0^*) := Q_{c_0^*} + \varepsilon_0^*$$

satisfies

$$\|(v_0^*, w_0^*) - Q_{\mathfrak{c}}\|_{X(\mathbb{R})} \leq A_{\mathfrak{c}}\beta_{\mathfrak{c}} + \|Q_{\mathfrak{c}} - Q_{c_0^*}\|_{X(\mathbb{R})}. \tag{2-17}$$

We next impose a supplementary smallness assumption on  $\beta_{\mathfrak{c}}$  so that

$$\|(v_0^*, w_0^*) - Q_{\mathfrak{c}}\|_{X(\mathbb{R})} \leq \alpha_{\mathfrak{c}}. \tag{2-18}$$

By [Theorem 2.1](#), there exists a unique global solution  $(v^*, w^*) \in \mathcal{C}^0(\mathbb{R}, \mathcal{NV}(\mathbb{R}))$  to [\(HLL\)](#) with initial datum  $(v_0^*, w_0^*)$ , and two maps  $c^* \in \mathcal{C}^1(\mathbb{R}, (-1, 1) \setminus \{0\})$  and  $a^* \in \mathcal{C}^1(\mathbb{R}, \mathbb{R})$  such that the function  $\varepsilon^*$  defined by

$$\varepsilon^*(\cdot, t) := (v^*(\cdot + a^*(t), t), w^*(\cdot + a^*(t), t)) - Q_{c^*(t)} \tag{2-19}$$

satisfies the orthogonality conditions

$$\langle \varepsilon^*(\cdot, t), \partial_x Q_{c^*(t)} \rangle_{L^2(\mathbb{R})^2} = \langle \varepsilon^*(\cdot, t), \chi_{c^*(t)} \rangle_{L^2(\mathbb{R})^2} = 0, \quad (2-20)$$

as well as the estimates

$$\|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})} + |c^*(t) - c| + |a^{*'}(t) - c^*(t)| \leq A_c \|v_0^*, w_0^* - Q_c\|_{X(\mathbb{R})}, \quad (2-21)$$

for any  $t \in \mathbb{R}$ .

We may take  $\beta_c$  small enough such that, combining (2-16) with (2-17) and (2-21), we obtain

$$\min\{c^*(t)^2, (a^{*}')^2(t)\} \geq \frac{c^2}{2}, \quad \max\{c^*(t)^2, (a^{*}')^2(t)\} \leq 1 + \frac{c^2}{2}, \quad (2-22)$$

and

$$\|v_c(\cdot) - v^*(\cdot + a^*(t), t)\|_{L^\infty(\mathbb{R})} \leq \min\left\{\frac{c^2}{4}, \frac{1-c^2}{16}\right\}, \quad (2-23)$$

for any  $t \in \mathbb{R}$ .

Finally, we use the weak continuity of the flow map for the Landau–Lifshitz equation. The proof relies on Proposition A.1 and follows the lines of the proof of Proposition 1 in [Béthuel et al. 2015].

**Proposition 2.3.** *Let  $t \in \mathbb{R}$  be fixed. Then*

$$(v(\cdot + a(t_n), t_n + t), w(\cdot + a(t_n), t_n + t)) \rightharpoonup (v^*(\cdot, t), w^*(\cdot, t)) \quad \text{in } X(\mathbb{R}), \quad (2-24)$$

while

$$a(t_n + t) - a(t_n) \rightarrow a^*(t) \quad \text{and} \quad c(t_n + t) \rightarrow c^*(t) \quad (2-25)$$

as  $n \rightarrow +\infty$ . In particular, we have

$$\varepsilon(\cdot, t_n + t) \rightharpoonup \varepsilon^*(\cdot, t) \quad \text{in } X(\mathbb{R}) \quad (2-26)$$

as  $n \rightarrow +\infty$ .

*Localization and smoothness of the limit profile.* Our proof of the localization of the limit profile is based on a monotonicity formula.

Consider a pair  $(v, w)$  which satisfies the conclusions of Theorem 2.1 and suppose that (2-14) and (2-15) are true. Let  $R$  and  $t$  be two real numbers, and set

$$I_R(t) \equiv I_R^{(v,w)}(t) := \frac{1}{2} \int_{\mathbb{R}} [vw](x + a(t), t) \Phi(x - R) dx,$$

where  $\Phi$  is the function defined on  $\mathbb{R}$  by

$$\Phi(x) := \frac{1}{2}(1 + \text{th}(v_c x)), \quad (2-27)$$

with  $v_c := \sqrt{1 - c^2}/8$ .

**Proposition 2.4.** *Let  $R \in \mathbb{R}$ ,  $t \in \mathbb{R}$  and  $\sigma \in [-\sigma_c, \sigma_c]$ , with  $\sigma_c := \sqrt{1 - \mathfrak{c}^2}/4$ . Under the above assumptions, there exists a positive number  $B_c$ , depending only on  $\mathfrak{c}$ , such that*

$$\frac{d}{dt}[I_{R+\sigma t}(t)] \geq \frac{1-\mathfrak{c}^2}{8} \int_{\mathbb{R}} [(\partial_x v)^2 + v^2 + w^2](x + a(t), t) \Phi'(x - R - \sigma t) dx - B_c e^{-2v_c |R+\sigma t|}. \tag{2-28}$$

In particular, we have

$$I_R(t_1) \geq I_R(t_0) - B_c e^{-2v_c |R|} \tag{2-29}$$

for any real numbers  $t_0 \leq t_1$ .

For the limit profile  $(v^*, w^*)$ , we set  $I_R^*(t) := I_R^{(v^*, w^*)}(t)$  for any  $R \in \mathbb{R}$  and any  $t \in \mathbb{R}$ .

**Proposition 2.5** [Béthuel et al. 2015]. *Given any positive number  $\delta$ , there exists a positive number  $R_\delta$ , depending only on  $\delta$ , such that we have*

$$\begin{aligned} |I_R^*(t)| &\leq \delta \quad \forall R \geq R_\delta, \\ |I_R^*(t) - P(v^*, w^*)| &\leq \delta \quad \forall R \leq -R_\delta \end{aligned}$$

for any  $t \in \mathbb{R}$ .

The proof of Proposition 2.5 is the same as that of Proposition 3 in [Béthuel et al. 2015].

From Propositions 2.4 and 2.5, as in [Béthuel et al. 2015] we derive:

**Proposition 2.6** [Béthuel et al. 2015]. *Let  $t \in \mathbb{R}$ . There exists a positive constant  $\mathcal{A}_c$  such that*

$$\int_t^{t+1} \int_{\mathbb{R}} [(\partial_x v^*)^2 + (v^*)^2 + (w^*)^2](x + a^*(s), s) e^{2v_c |x|} dx ds \leq \mathcal{A}_c.$$

We next consider the following map, which was introduced by de Laire and Gravejat [2015]:

$$\Psi := \frac{1}{2} \left( \frac{\partial_x v}{(1-v^2)^{1/2}} + i(1-v^2)^{1/2} w \right) \exp i\theta, \tag{2-30}$$

where

$$\theta(x, t) := - \int_{-\infty}^x v(y, t) w(y, t) dy. \tag{2-31}$$

The map  $\Psi$  solves the nonlinear Schrödinger equation

$$i \partial_t \Psi + \partial_{xx} \Psi + 2|\Psi|^2 \Psi + \frac{1}{2} v^2 \Psi - \text{Re}(\Psi(1 - 2F(v, \bar{\Psi}))) (1 - 2F(v, \Psi)) = 0, \tag{2-32}$$

with

$$F(v, \Psi)(x, t) := \int_{-\infty}^x v(y, t) \Psi(y, t) dy, \tag{2-33}$$

while the function  $v$  satisfies the two equations

$$\begin{cases} \partial_t v = 2\partial_x \text{Im}(\Psi(2F(v, \bar{\Psi}) - 1)), \\ \partial_x v = 2\text{Re}(\Psi(1 - 2F(v, \bar{\Psi}))). \end{cases} \tag{2-34}$$

The local Cauchy problem for (2-32)–(2-34) was analyzed by de Laire and Gravejat [2015]. We recall the following proposition which shows the continuous dependence with respect to the initial datum of the solutions to the system of equations (2-32)–(2-34) (see [de Laire and Gravejat 2015] for the proof).

**Proposition 2.7** [de Laire and Gravejat 2015]. *Suppose that the pairs  $(v^0, \Psi^0) \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$  and  $(\tilde{v}^0, \tilde{\Psi}^0) \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$  are such that*

$$\partial_x v^0 = 2 \operatorname{Re}(\Psi^0(1 - 2F(v^0, \overline{\Psi^0}))) \quad \text{and} \quad \partial_x \tilde{v}^0 = 2 \operatorname{Re}(\tilde{\Psi}^0(1 - 2F(\tilde{v}^0, \overline{\tilde{\Psi}^0}))).$$

*Given two solutions  $(v, \Psi)$  and  $(\tilde{v}, \tilde{\Psi})$  in  $C^0([0, T_*], H^1(\mathbb{R}) \times L^2(\mathbb{R}))$ , with  $(\Psi, \tilde{\Psi}) \in L^4([0, T_*], L^\infty(\mathbb{R}))^2$ , to (2-32)–(2-34) with initial data  $(v^0, \Psi^0)$  and  $(\tilde{v}^0, \tilde{\Psi}^0)$  respectively, for some positive time  $T_*$ , there exist a positive number  $\tau$ , depending only on  $\|v^0\|_{L^2}$ ,  $\|\tilde{v}^0\|_{L^2}$ ,  $\|\Psi^0\|_{L^2}$  and  $\|\tilde{\Psi}^0\|_{L^2}$ , and a universal constant  $A$  such that we have*

$$\|v - \tilde{v}\|_{C^0([0, T], L^2)} + \|\Psi - \tilde{\Psi}\|_{C^0([0, T], L^2)} + \|\Psi - \tilde{\Psi}\|_{L^4([0, T], L^\infty)} \leq A(\|v^0 - \tilde{v}^0\|_{L^2} + \|\Psi^0 - \tilde{\Psi}^0\|_{L^2}) \quad (2-35)$$

*for any  $T \in [0, \min\{\tau, T_*\}]$ . In addition, there exists a positive number  $B$ , depending only on  $\|v^0\|_{L^2}$ ,  $\|\tilde{v}^0\|_{L^2}$ ,  $\|\Psi^0\|_{L^2}$  and  $\|\tilde{\Psi}^0\|_{L^2}$ , such that*

$$\|\partial_x v - \partial_x \tilde{v}\|_{C^0([0, T], L^2)} \leq B(\|v^0 - \tilde{v}^0\|_{L^2} + \|\Psi^0 - \tilde{\Psi}^0\|_{L^2}) \quad (2-36)$$

*for any  $T \in [0, \min\{\tau, T_*\}]$ .*

This proposition plays an important role in the proof of not only the smoothing of the limit profile, but also the weak continuity of the hydrodynamical Landau–Lifshitz flow.

In order to prove the smoothness of the limit profile, we rely on the following smoothing type estimate for localized solutions of the linear Schrödinger equation (see [Béthuel et al. 2015; Escauriaza et al. 2008] for the proof of Proposition 2.8).

**Proposition 2.8** [Béthuel et al. 2015; Escauriaza et al. 2008]. *Let  $\lambda \in \mathbb{R}$ , and consider a solution  $u \in C^0(\mathbb{R}, L^2(\mathbb{R}))$  to the linear Schrödinger equation*

$$i \partial_t u + \partial_{xx} u = F, \quad (\text{LS})$$

*with  $F \in L^2(\mathbb{R}, L^2(\mathbb{R}))$ . Then there exists a positive constant  $K_\lambda$ , depending only on  $\lambda$ , such that*

$$\lambda^2 \int_{-T}^T \int_{\mathbb{R}} |\partial_x u(x, t)|^2 e^{\lambda x} dx dt \leq K_\lambda \int_{-T-1}^{T+1} \int_{\mathbb{R}} (|u(x, t)|^2 + |F(x, t)|^2) e^{\lambda x} dx dt \quad (2-37)$$

*for any positive number  $T$ .*

We apply Proposition 2.8 to  $\Psi^*$  as well as all its derivatives, where  $\Psi^*$  is the solution to (2-32) associated to the solution  $(v^*, w^*)$  of (HLL), and then express the result in terms of  $(v^*, w^*)$  to obtain:

**Proposition 2.9.** *The pair  $(v^*, w^*)$  is indefinitely smooth and exponentially decaying on  $\mathbb{R} \times \mathbb{R}$ . Moreover, given any  $k \in \mathbb{N}$ , there exists a positive constant  $A_{k, c}$ , depending only on  $k$  and  $c$ , such that*

$$\int_{\mathbb{R}} [(\partial_x^{k+1} v^*)^2 + (\partial_x^k v^*)^2 + (\partial_x^k w^*)^2](x + a^*(t), t) e^{\nu_c |x|} dx \leq A_{k, c} \quad (2-38)$$

*for any  $t \in \mathbb{R}$ .*

*The Liouville type theorem.* We next establish a Liouville type theorem, which guarantees that the limit profile constructed above is exactly a soliton. In particular, we will show that  $\varepsilon_0^* \equiv 0$ .

The pair  $\varepsilon^*$  satisfies the equation

$$\partial_t \varepsilon^* = J\mathcal{H}_{c^*(t)}(\varepsilon^*) + J\mathcal{R}_{c^*(t)}\varepsilon^* + (a^{*\prime}(t) - c^*(t))(\partial_x Q_{c^*(t)} + \partial_x \varepsilon^*) - c^{*\prime}(t)\partial_c Q_{c^*(t)}, \tag{2-39}$$

where  $J$  is the symplectic operator

$$J = -2S\partial_x := \begin{pmatrix} 0 & -2\partial_x \\ -2\partial_x & 0 \end{pmatrix}, \tag{2-40}$$

and the remainder term  $\mathcal{R}_{c^*(t)}\varepsilon^*$  is given by

$$\mathcal{R}_{c^*(t)}\varepsilon^* := E'(Q_{c^*(t)} + \varepsilon^*) - E'(Q_{c^*(t)}) - E''(Q_{c^*(t)})(\varepsilon^*).$$

We rely on the strategy developed by Martel and Merle [2008a] (see also [Martel 2006]), and then applied by B ethuel, Gravejat and Smets in [B ethuel et al. 2015] to the Gross–Pitaevskii equation. We define the pair

$$u^*(\cdot, t) := S\mathcal{H}_{c^*(t)}(\varepsilon^*(\cdot, t)). \tag{2-41}$$

Since  $S\mathcal{H}_{c^*(t)}(\partial_x Q_{c^*(t)}) = 0$ , we deduce from (2-39) that

$$\begin{aligned} \partial_t u^* &= \mathcal{H}_{c^*(t)}(J S u^*) + S\mathcal{H}_{c^*(t)}(J\mathcal{R}_{c^*(t)}\varepsilon^*) - (c^*)'(t)S\mathcal{H}_{c^*(t)}(\partial_c Q_{c^*(t)}) \\ &\quad + (c^*)'(t)S\partial_c \mathcal{H}_{c^*(t)}(\varepsilon^*) + ((a^*)'(t) - c^*(t))S\mathcal{H}_{c^*(t)}(\partial_x \varepsilon^*). \end{aligned} \tag{2-42}$$

Decreasing further the value of  $\beta_c$  if necessary, we have:

**Proposition 2.10.** *There exist two positive numbers  $A_*$  and  $R_*$ , depending only on  $c$ , such that we have<sup>2</sup>*

$$\frac{d}{dt} \left( \int_{\mathbb{R}} x u_1^*(x, t) u_2^*(x, t) dx \right) \geq \frac{1-c^2}{16} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 - A_* \|u^*(\cdot, t)\|_{X(B(0, R_*))}^2 \tag{2-43}$$

for any  $t \in \mathbb{R}$ .

We give a second monotonicity type formula to dispose of the nonpositive local term  $\|u^*(\cdot, t)\|_{X(B(0, R_*))}^2$  on the right-hand side of (2-43). If  $M$  is a smooth, bounded, two-by-two symmetric matrix-valued function, then

$$\frac{d}{dt} \langle M u^*, u^* \rangle_{L^2(\mathbb{R})^2} = 2 \langle S M u^*, \mathcal{H}_{c^*}(-2\partial_x u^*) \rangle_{L^2(\mathbb{R})^2} + \text{“superquadratic terms”}, \tag{2-44}$$

where  $S$  is the matrix

$$S := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

<sup>2</sup>In (2-43), we use the notation

$$\|(f, g)\|_{X(\Omega)}^2 := \int_{\Omega} ((\partial_x f)^2 + f^2 + g^2),$$

in which  $\Omega$  denotes a measurable subset of  $\mathbb{R}$ .

For  $c \in (-1, 1) \setminus \{0\}$ , let  $M_c$  be given by

$$M_c := \begin{pmatrix} -\frac{2cv_c \partial_x v_c}{(1-v_c)^2} & -\frac{\partial_x v_c}{v_c} \\ -\frac{\partial_x v_c}{v_c} & 0 \end{pmatrix}. \tag{2-45}$$

We have the following lemma.

**Lemma 2.11.** *Let  $c \in (-1, 1) \setminus \{0\}$  and  $u \in X^3(\mathbb{R})$ . Then*

$$\begin{aligned} G_c(u) &:= 2\langle SM_c u, \mathcal{H}_c(-2\partial_x u) \rangle_{L^2(\mathbb{R})^2} \\ &= 2 \int_{\mathbb{R}} \mu_c \left( u_2 - \frac{cv_c^2}{\mu_c} u_1 - \frac{2cv_c \partial_x v_c}{\mu_c(1-v_c^2)} \partial_x u_1 \right)^2 + 3 \int_{\mathbb{R}} \frac{v_c^4}{\mu_c} \left( \partial_x u_1 - \frac{\partial_x v_c}{v_c} u_1 \right)^2, \end{aligned} \tag{2-46}$$

where

$$\mu_c = 2(\partial_x v_c)^2 + v_c^2(1 - v_c^2) > 0. \tag{2-47}$$

The functional  $G_c$  is a nonnegative quadratic form, and

$$\text{Ker}(G_c) = \text{Span}(Q_c). \tag{2-48}$$

We have indeed chosen the matrix  $M_c$  such that  $M_c Q_c = \partial_x Q_c$  to obtain (2-48). Since  $Q_c$  does not vanish, we deduce from standard Sturm–Liouville theory that  $G_c$  is nonnegative, which is confirmed by the computation in Lemma 2.11.

By the second orthogonality condition in (2-20) and the fact that  $\mathcal{H}_{c^*}(\chi_{c^*}) = -\tilde{\lambda}_{c^*} \chi_{c^*}$ , we have

$$0 = \langle \mathcal{H}_{c^*}(\chi_{c^*}), \varepsilon^* \rangle_{L^2(\mathbb{R})^2} = \langle \mathcal{H}_{c^*}(\varepsilon^*), \chi_{c^*} \rangle_{L^2(\mathbb{R})^2} = \langle u^*, S\chi_{c^*} \rangle_{L^2(\mathbb{R})^2}. \tag{2-49}$$

On the other hand, we know that

$$\langle Q_{c^*}, S\chi_{c^*} \rangle = P'(Q_{c^*})(\chi_{c^*}) \neq 0, \tag{2-50}$$

so that the pair  $u^*$  is not proportional to  $Q_{c^*}$  under the orthogonality condition in (2-49). We claim the following coercivity property of  $G_c$  under this orthogonality condition.

**Proposition 2.12.** *Let  $c \in (-1, 1) \setminus \{0\}$ . There exists a positive number  $\Lambda_c$ , depending only and continuously on  $c$ , such that*

$$G_c(u) \geq \Lambda_c \int_{\mathbb{R}} [(\partial_x u_1)^2 + (u_1)^2 + (u_2)^2](x) e^{-2|x|} dx, \tag{2-51}$$

for any pair  $u \in X(\mathbb{R})$  verifying

$$\langle u, S\chi_c \rangle_{L^2(\mathbb{R})^2} = 0. \tag{2-52}$$

Coming back to (2-44), we can prove the next proposition.

**Proposition 2.13.** *There exists a positive number  $B_*$ , depending only on  $\mathfrak{c}$ , such that*

$$\frac{d}{dt} \langle (M_{c^*(t)} u^*(\cdot, t), u^*(\cdot, t))_{L^2(\mathbb{R}^2)} \rangle \geq \frac{1}{B_*} \int_{\mathbb{R}} [(\partial_x u_1^*)^2 + (u_1^*)^2 + (u_2^*)^2](x, t) e^{-2|x|} dx - B_* \|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 \quad (2-53)$$

for any  $t \in \mathbb{R}$ .

Propositions 2.10 and 2.13 have the following corollary.

**Corollary 2.14.** *Set*

$$N(t) := \frac{1}{2} \begin{pmatrix} 0 & x \\ x & 0 \end{pmatrix} + A_* B_* e^{2R_*} M_{c^*(t)}.$$

*There exists a positive constant  $\mathcal{A}_c$  such that we have*

$$\frac{d}{dt} \langle (N(t)u^*(\cdot, t), u^*(\cdot, t))_{L^2(\mathbb{R}^2)} \rangle \geq \mathcal{A}_c \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 \quad (2-54)$$

for any  $t \in \mathbb{R}$ . Since

$$\int_{-\infty}^{+\infty} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 dt < +\infty, \quad (2-55)$$

there exists a sequence  $(t_k^*)_{k \in \mathbb{N}}$  such that

$$\lim_{k \rightarrow +\infty} \|u^*(\cdot, t_k^*)\|_{X(\mathbb{R})}^2 = 0. \quad (2-56)$$

In view of (2-20), (2-41) and the bound for  $\mathcal{H}_{c^*}$  in (A-43), we have

$$\|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})} \leq A_c \|u^*(\cdot, t)\|_{X(\mathbb{R})}, \quad (2-57)$$

Hence, we can apply (2-56) and (2-57) in order to obtain

$$\lim_{k \rightarrow +\infty} \|\varepsilon^*(\cdot, t_k^*)\|_{X(\mathbb{R})}^2 = 0. \quad (2-58)$$

By (2-58) and the orbital stability in Theorem 2.1, this yields:

**Corollary 2.15.**  $\varepsilon_0^* \equiv 0$ .

At this stage we obtain (2-11) for some subsequence. We should extend this result for any sequence. The proof is exactly the same as the one done by Béthuel, Gravejat and Smets in [Béthuel et al. 2015] (see Subsection 1.3.4 in [Béthuel et al. 2015] for the details).

*Proof of Theorem 1.1.* We choose a positive number  $\delta_c$  such that  $\|(v_0, w_0) - Q_c\|_{X(\mathbb{R})} \leq \beta_c$ , whenever  $d_{\mathcal{E}}(m^0, u_c) \leq \delta_c$ . We next apply Theorem 2.2 to the solution  $(v, w) \in \mathcal{C}^0(\mathbb{R}, \mathcal{N}\mathcal{V}(\mathbb{R}))$  to (HLL) corresponding to the solution  $m$  to (LL). This yields the existence of a speed  $\mathfrak{c}^*$  and a position function  $b$  such that the convergences in Theorem 2.2 hold. In particular, since the weak convergence for  $m_3$  is satisfied by Theorem 2.2, it is sufficient to show the existence of a phase function  $\theta$  such that  $\exp(i\theta(t))\partial_x \check{m}(\cdot + b(t), t)$  is weakly convergent to  $\partial_x \check{u}_{\mathfrak{c}^*}$  in  $L^2(\mathbb{R})$  as  $t \rightarrow \infty$ . The locally uniform

convergence of  $\exp(i\theta(t))\check{m}(\cdot + b(t), t)$  towards  $\check{u}_{c^*}$  then follows from the Sobolev embedding theorem. We begin by constructing this phase function.

We fix a nonzero function  $\chi \in C_c^\infty(\mathbb{R}, [0, 1])$  such that  $\chi$  is even. Using the explicit formula of  $\check{u}_{c^*}$ , we have

$$\int_{\mathbb{R}} \check{u}_{c^*}(x)\chi(x) dx = 2c^* \int_{\mathbb{R}} \frac{\chi(x)}{\cosh(\sqrt{1 - (c^*)^2}x)} dx \neq 0. \quad (2-59)$$

Decreasing the value of  $\beta_c$  if needed, we deduce from the orbital stability in [de Laire and Gravejat 2015] that

$$\left| \int_{\mathbb{R}} \check{m}(x + b(t), t)\chi(x) dx \right| \geq |c^*| \int_{\mathbb{R}} \frac{\chi(x)}{\cosh(\sqrt{1 - (c^*)^2}x)} dx \neq 0 \quad (2-60)$$

for any  $t \in \mathbb{R}$ .

Let  $\Upsilon : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the  $C^1$  function defined by

$$\Upsilon(t, \theta) := \operatorname{Im} \left( e^{-i\theta} \int_{\mathbb{R}} \check{m}(x + b(t), t)\chi(x) dx \right).$$

From (2-60) we can find a number  $\theta_0$  such that  $\Upsilon(0, \theta_0) = 0$  and  $\partial_\theta \Upsilon(0, \theta_0) > 0$ . Then, using the implicit function theorem, there exists a  $C^1$  function  $\theta : \mathbb{R} \rightarrow \mathbb{R}$  such that  $\Upsilon(t, \theta(t)) = 0$ . In addition, using (2-60) another time, we can fix the choice of  $\theta$  so that there exists a positive constant  $A_{c^*}$  such that

$$\partial_\theta \Upsilon(t, \theta(t)) = \operatorname{Re} \left( e^{-i\theta(t)} \int_{\mathbb{R}} \check{m}(x + b(t), t)\chi(x) dx \right) \geq A_{c^*} > 0. \quad (2-61)$$

Differentiating the identity  $\Upsilon(t, \theta(t)) = 0$  with respect to  $t$ , this implies that

$$|\theta'(t)| = \left| \frac{\partial_t \Upsilon(t, \theta(t))}{\partial_\theta \Upsilon(t, \theta(t))} \right| \leq \frac{1}{A_{c^*}} |\partial_t \Upsilon(t, \theta(t))| \quad (2-62)$$

for all  $t \in \mathbb{R}$ . Now, we differentiate the function  $\Upsilon$  with respect to  $t$ , and we use the equation of  $\check{m}$  to obtain

$$\begin{aligned} \partial_t \Upsilon(t, \theta(t)) = \operatorname{Im} \left( e^{-i\theta} \int_{\mathbb{R}} \chi(x) (\partial_x \check{m}(x + b(t), t)b'(t) - im_3(x + b(t), t)\partial_{xx} \check{m}(x + b(t), t) \right. \\ \left. + i\check{m}(x + b(t), t)\partial_{xx} m_3(x + b(t), t) - im_3(x + b(t), t)\check{m}(x + b(t), t)) dx \right). \end{aligned} \quad (2-63)$$

Since  $b \in C_b^1(\mathbb{R}, \mathbb{R})$ , and since both  $\partial_x \check{m}$  and  $\partial_t \check{m}$  belong to  $C_b^0(\mathbb{R}, H^{-1}(\mathbb{R}))$ , it follows that the derivative  $\theta'$  is bounded on  $\mathbb{R}$ .

We denote by  $\varphi$  the phase function defined by

$$\varphi(x + b(t), t) := \varphi(b(t), t) + \int_0^x w(y + b(t), t) dy,$$

with  $\varphi(b(t), t) \in [0, 2\pi]$ , which is associated to the function  $\check{m}(x + b(t), t)$  for any  $(x, t) \in \mathbb{R}^2$  in the way that

$$\check{m}(x + b(t), t) = (1 - m_3^2(x + b(t), t))^{1/2} \exp(i\varphi(x + b(t), t)).$$

It is sufficient to prove that

$$\exp(i(\varphi(b(t), t) - \theta(t))) \rightarrow 1 \tag{2-64}$$

as  $t \rightarrow \infty$  to obtain

$$\exp(i(\varphi(\cdot + b(t), t) - \theta(t))) \rightarrow \exp(i\varphi_{c^*}(\cdot)) := \exp\left(i \int_0^\cdot w_{c^*}(y) dy\right) \quad \text{in } L^\infty_{\text{loc}}(\mathbb{R})$$

as  $t \rightarrow \infty$ . This implies, using [Theorem 2.2](#) once again as well as the Sobolev embedding theorem, that

$$\begin{aligned} e^{-i\theta(t)} \partial_x \check{m}(\cdot + b(t), t) &\rightarrow \partial_x \check{u}_{c^*} && \text{in } L^2(\mathbb{R}), \\ e^{-i\theta(t)} \check{m}(\cdot + b(t), t) &\rightarrow \check{u}_{c^*} && \text{in } L^\infty_{\text{loc}}(\mathbb{R}) \end{aligned} \tag{2-65}$$

as  $t \rightarrow \infty$ . Now let us prove [\(2-64\)](#). We have

$$\begin{aligned} e^{-i\theta(t)} \int_{\mathbb{R}} \check{m}(x + b(t), t) \chi(x) dx \\ = \exp(i[\varphi(b(t), t) - \theta(t)]) \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx. \end{aligned}$$

We use the fact that  $\Upsilon(t, \theta(t)) = 0$  to obtain

$$\begin{aligned} \cos(\varphi(b(t), t) - \theta(t)) \operatorname{Im} \left( \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx \right) \\ + \sin(\varphi(b(t), t) - \theta(t)) \operatorname{Re} \left( \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx \right) = 0. \end{aligned}$$

On the other hand, by [\(2-61\)](#), we have

$$\begin{aligned} \cos(\varphi(b(t), t) - \theta(t)) \operatorname{Re} \left( \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx \right) \\ - \sin(\varphi(b(t), t) - \theta(t)) \operatorname{Im} \left( \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx \right) > 0. \end{aligned}$$

We derive from [Theorem 2.2](#) and [\(2-59\)](#) that

$$\operatorname{Im} \left( \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx \right) \rightarrow \operatorname{Im} \left( \int_{\mathbb{R}} \check{u}_{c^*}(x) \chi(x) dx \right) = 0,$$

and

$$\operatorname{Re} \left( \int_{\mathbb{R}} (1 - m_3^2(x + b(t), t))^{1/2} \exp\left(i \int_0^x w(y + b(t), t) dy\right) \chi(x) dx \right) \rightarrow \operatorname{Re} \left( \int_{\mathbb{R}} \check{u}_{c^*}(x) \chi(x) dx \right) > 0.$$

This is enough to derive [\(2-64\)](#).

Finally, we claim that  $\theta'(t) \rightarrow 0$  as  $t \rightarrow \infty$ . Indeed, we can introduce [\(2-65\)](#) into [\(2-63\)](#), and we then obtain, using the equation satisfied by  $\check{u}_{c^*}$ , that

$$\partial_t \Upsilon(t, \theta(t)) \rightarrow 0$$

as  $t \rightarrow \infty$ . By [\(2-62\)](#), this yields  $\theta'(t) \rightarrow 0$  as  $t \rightarrow \infty$ , which finishes the proof of [Theorem 1.1](#). □

### 3. Proof of the orbital stability

First, we recall the orbital stability theorem, which was established in [de Laire and Gravejat 2015] (see Corollary 2 and Propositions 2 and 4 in [de Laire and Gravejat 2015]).

**Theorem 3.1.** *Let  $c \in (-1, 1) \setminus \{0\}$  and  $(v_0, w_0) \in X(\mathbb{R})$  satisfying (2-4). There exist a unique global solution  $(v, w) \in \mathcal{C}^0(\mathbb{R}, \mathcal{NV}(\mathbb{R}))$  to (HLL) with initial datum  $(v_0, w_0)$ , and two maps  $c_1 \in \mathcal{C}^1(\mathbb{R}, (-1, 1) \setminus \{0\})$  and  $a_1 \in \mathcal{C}^1(\mathbb{R}, \mathbb{R})$  such that the function  $\varepsilon_1$ , defined by (2-5), satisfies the orthogonality conditions*

$$\langle \varepsilon_1(\cdot, t), \partial_x Q_{c_1(t)} \rangle_{L^2(\mathbb{R}^2)} = P'(Q_{c_1(t)})(\varepsilon_1(\cdot, t)) = 0 \tag{3-1}$$

for any  $t \in \mathbb{R}$ . Moreover,  $\varepsilon_1(\cdot, t)$ ,  $c_1(t)$  and  $a_1(t)$  satisfy (2-7), (2-8) and (2-9) for any  $t \in \mathbb{R}$ .

With Theorem 3.1 at hand, we can provide the proof of Theorem 2.1.

*Proof of Theorem 2.1.* We consider the map

$$\Xi((v, w), \sigma, \mathbf{b}) := (\langle \partial_x Q_{\sigma, \mathbf{b}}, \varepsilon \rangle_{L^2 \times L^2}, \langle \chi_{\sigma, \mathbf{b}}, \varepsilon \rangle_{L^2 \times L^2}),$$

where we have set  $\varepsilon = (v, w) - Q_{\sigma, \mathbf{b}}$ , and  $\chi_{\sigma, \mathbf{b}} = \chi_\sigma(\cdot - \mathbf{b})$  (we recall that  $\chi_\sigma$  is the eigenfunction associated to the unique negative eigenvalue  $-\tilde{\lambda}_\sigma$  of the operator  $\mathcal{H}_\sigma$ ). The map  $\Xi$  is well-defined for, and depends smoothly on,  $(v, w) \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$ ,  $\sigma \in (-1, 1) \setminus \{0\}$  and  $\mathbf{b} \in \mathbb{R}$ .

We fix  $t \in \mathbb{R}$ . In order to simplify the notation, we substitute  $(c_1(t), a_1(t))$  by  $(c_1, a_1)$ . We check that

$$\Xi(Q_{c_1, a_1}, c_1, a_1) = 0,$$

and we compute

$$\begin{cases} \partial_\sigma \Xi_1(Q_{c_1, a_1}, c_1, a_1) = 0, \\ \partial_\sigma \Xi_2(Q_{c_1, a_1}, c_1, a_1) = -\langle \chi_{c_1, a_1}, \partial_\sigma Q_{c_1, a_1} \rangle_{L^2 \times L^2}. \end{cases}$$

Let  $c \in (-1, 1) \setminus \{0\}$  and suppose towards a contradiction that

$$\langle \chi_c, \partial_c Q_c \rangle_{L^2 \times L^2} = 0.$$

Using the fact that  $\mathcal{H}_c(\partial_c Q_c) = P'(Q_c)$ , this gives

$$0 = \langle \chi_c, \partial_c Q_c \rangle_{L^2 \times L^2} = -\frac{1}{\tilde{\lambda}_c} \langle \chi_c, \mathcal{H}_c(\partial_c Q_c) \rangle_{L^2 \times L^2} = -\frac{1}{\tilde{\lambda}_c} \langle \chi_c, P'(Q_c) \rangle_{L^2 \times L^2}.$$

Since  $\mathcal{H}_c$  is self-adjoint, we also have

$$\langle \chi_c, \partial_x Q_c \rangle_{L^2 \times L^2} = 0.$$

By Proposition 1 in [de Laire and Gravejat 2015], we infer that

$$0 > -\tilde{\lambda}_c \|\chi_c\|_{L^2 \times L^2}^2 = \langle \chi_c, \mathcal{H}_c(\chi_c) \rangle_{L^2 \times L^2} \geq \Lambda_c \|\chi_c\|_{L^2 \times L^2}^2 > 0,$$

which provides the contradiction and shows that

$$\langle \chi_c, \partial_c Q_c \rangle_{L^2 \times L^2} \neq 0 \tag{3-2}$$

for all  $c \in (-1, 1) \setminus \{0\}$ . In addition, we have

$$\begin{cases} \partial_b \Xi_1(Q_{c_1, a_1}, c_1, a_1) = \|\partial_x Q_{c_1}\|_{L^2}^2 = 2(1 - c_1^2)^{1/2} > 0, \\ \partial_b \Xi_2(Q_{c_1, a_1}, c_1, a_1) = 0. \end{cases}$$

Therefore, the matrix

$$d_{\sigma, b} \Xi(Q_{c_1, a_1}, c_1, a_1) = \begin{pmatrix} 0 & \langle \chi_{c_1, a_1}, \partial_\sigma Q_{c_1, a_1} \rangle_{L^2 \times L^2} \\ 2(1 - c_1^2)^{1/2} & 0 \end{pmatrix}$$

is an isomorphism from  $\mathbb{R}^2$  to  $\mathbb{R}^2$ .

Then, we can apply the version of the implicit function theorem in [Béthuel et al. 2014] in order to find a neighbourhood  $\mathcal{V}$  of  $Q_{c_1, a_1}$ , a neighbourhood  $\mathcal{U}$  of  $(c_1, a_1)$ , and a map  $\gamma_{c_1, a_1} : \mathcal{U} \rightarrow \mathcal{V}$  such that

$$\Xi((v, w), \sigma, \mathbf{b}) = 0 \Leftrightarrow (c(v, w), a(v, w)) := (\sigma, \mathbf{b}) = \gamma_{c, a}(v, w) \quad \forall (v, w) \in \mathcal{V}, \forall (\sigma, \mathbf{b}) \in \mathcal{U}.$$

In addition, there exists a positive constant  $\Lambda$ , depending only on  $c_1$ , such that

$$\|\varepsilon(t)\|_X + |c(t) - c_1(t)| + |a(t) - a_1(t)| \leq \Lambda \|\varepsilon_1(t)\|_X \leq \Lambda_{c_1} A_c \alpha_0, \tag{3-3}$$

where  $c(t) := c(v(t), w(t))$ ,  $a(t) := a(v(t), w(t))$  and  $\varepsilon(t) := (v(t), w(t)) - Q_{c(t), a(t)}$ , for any fixed  $t \in \mathbb{R}$ . Using the fact that  $(v(t), w(t))$  stays in a neighbourhood of  $Q_{c_1(t), a_1(t)}$  for all  $t \in \mathbb{R}$  by Theorem 3.1, and also the fact that  $c_1$  satisfies (2-8), we are led to the following lemma.

**Lemma 3.2.** *Under the assumptions of Theorem 3.1, there exists a unique pair  $(a, c)$  of functions in  $C^0(\mathbb{R}, \mathbb{R}^2)$  such that*

$$\varepsilon(t) := (v(t), w(t)) - Q_{c(t), a(t)}$$

*satisfies the orthogonality conditions*

$$\langle \varepsilon(t), \partial_x Q_{c(t), a(t)} \rangle_{L^2 \times L^2} = \langle \chi_{c(t), a(t)}, \varepsilon(t) \rangle_{L^2 \times L^2} = 0. \tag{3-4}$$

Moreover, we have (2-8).

This completes the proof of orbital stability. Now, let us prove the continuous differentiability of the functions  $a$  and  $c$ , as well as the inequality

$$|c'(t)| + |a'(t) - c(t)| \leq A_c \|\varepsilon(\cdot, t)\|_{X(\mathbb{R})}, \tag{3-5}$$

for all  $t \in \mathbb{R}$ . The  $C^1$  nature of  $a$  and  $c$  can be derived from a standard density argument as in [de Laire and Gravejat 2015]. Concerning (3-5), we can write the equations satisfied by  $\varepsilon$ , namely

$$\partial_t \varepsilon_v = ((a'(t) - c(t)) \partial_x v_{c, a} - c'(t) \partial_c v_{c, a}) + \partial_x ((v_{c, a} + \varepsilon_v)^2 - 1)(v_{c, a} + \varepsilon_w) - (v_{c, a}^2 - 1) w_{c, a} \tag{3-6}$$

and

$$\begin{aligned} \partial_t \varepsilon_w &= (a'(t) - c(t)) \partial_x w_{c,a} - c'(t) \partial_c w_{c,a} \\ &+ \partial_x \left( \frac{\partial_{xx} v_{c,a} + \partial_{xx} \varepsilon_v}{1 - (v_{c,a} + \varepsilon_v)^2} + (v_{c,a} + \varepsilon_v) \frac{(\partial_x v_{c,a} + \partial_x \varepsilon_v)^2}{(1 - (v_{c,a} + \varepsilon_v)^2)^2} - \frac{\partial_{xx} v_{c,a}}{1 - v_{c,a}^2} - v_{c,a} \frac{(\partial_x v_{c,a})^2}{(1 - v_{c,a}^2)^2} \right) \\ &\quad + \partial_x \left( (v_{c,a} + \varepsilon_v) ((w_{c,a} + \varepsilon_w)^2 - 1) - v_{c,a} (w_{c,a}^2 - 1) \right). \end{aligned} \quad (3-7)$$

We differentiate with respect to time the orthogonality conditions in (2-6) and we invoke equations (3-6) and (3-7) to write the identity

$$M \begin{pmatrix} c' \\ a' - c \end{pmatrix} = \begin{pmatrix} Y \\ Z \end{pmatrix}. \quad (3-8)$$

Here,  $M$  refers to the matrix of size 2 given by

$$\begin{aligned} M_{1,1} &= \langle \partial_c Q_c, \chi_c \rangle_{L^2 \times L^2} + \langle \partial_c \chi_{c,a}, \varepsilon \rangle_{L^2 \times L^2}, \\ M_{1,2} &= \langle \chi_c, \partial_x Q_c \rangle_{L^2 \times L^2} - \langle \partial_x \chi_{c,a}, \varepsilon \rangle_{L^2 \times L^2}, \\ M_{2,1} &= -\langle \partial_x Q_c, \partial_c Q_c \rangle_{L^2 \times L^2} + \langle \partial_c \partial_x Q_{c,a}, \varepsilon \rangle_{L^2 \times L^2}, \\ M_{2,2} &= \|\partial_x Q_c\|_{L^2 \times L^2}^2 - \langle \partial_{xx} Q_{c,a}, \varepsilon \rangle_{L^2 \times L^2}. \end{aligned}$$

The vectors  $Y$  and  $Z$  are defined by

$$\begin{aligned} Y &= \langle \partial_x w_{c,a}, ((v_{c,a} + \varepsilon_v)^2 - 1)(w_{c,a} + \varepsilon_w) - (v_{c,a}^2 - 1)w_{c,a} \rangle_{L^2} \\ &\quad + \langle \partial_x v_{c,a}, ((w_{c,a} + \varepsilon_w)^2 - 1)(v_{c,a} + \varepsilon_v) - (w_{c,a}^2 - 1)v_{c,a} \rangle_{L^2} \\ &\quad - \left\langle \partial_{xx} v_{c,a}, \frac{\partial_{xx} v_{c,a} + \partial_{xx} \varepsilon_v}{1 - (v_{c,a} + \varepsilon_v)^2} - \frac{\partial_{xx} v_{c,a}}{1 - v_{c,a}^2} \right\rangle_{L^2} + c \langle \partial_x \chi_{c,a}, \varepsilon \rangle_{L^2 \times L^2} \end{aligned}$$

and

$$\begin{aligned} Z &= \langle \partial_{xx} v_{c,a}, ((v_{c,a} + \varepsilon_v)^2 - 1)(w_{c,a} + \varepsilon_w) - (v_{c,a}^2 - 1)w_{c,a} \rangle_{L^2} \\ &\quad + \langle \partial_{xx} w_{c,a}, ((w_{c,a} + \varepsilon_w)^2 - 1)(v_{c,a} + \varepsilon_v) - (w_{c,a}^2 - 1)v_{c,a} \rangle_{L^2} \\ &\quad - \left\langle \partial_{xxx} w_{c,a}, \frac{\partial_{xx} v_{c,a} + \partial_{xx} \varepsilon_v}{1 - (v_{c,a} + \varepsilon_v)^2} - \frac{\partial_{xx} v_{c,a}}{1 - v_{c,a}^2} \right\rangle_{L^2} + c \langle \partial_{xx} Q_{c,a}, \varepsilon \rangle_{L^2 \times L^2}. \end{aligned}$$

We next decompose the matrix  $M$  as  $M = D + H$ , where  $D$  is the diagonal matrix of size 2 with diagonal coefficients

$$D_{1,1} = \langle \partial_c Q_c, \chi_c \rangle_{L^2 \times L^2} \neq 0,$$

by (3-2), and

$$D_{2,2} = \|\partial_x Q_{c(t)}\|_{L^2}^2 = 2(1 - c(t)^2)^{1/2},$$

so that  $D$  is invertible. Concerning the matrix  $H$ , we check that

$$\langle P'(Q_c), \partial_x Q_c \rangle_{L^2 \times L^2} = \langle \partial_x Q_c, \partial_c Q_c \rangle_{L^2 \times L^2} = 0.$$

Then,

$$H = \begin{pmatrix} \langle \partial_c \chi_{c,a}, \varepsilon \rangle_{L^2 \times L^2} & -\langle \partial_x \chi_{c,a}, \varepsilon \rangle_{L^2 \times L^2} \\ \langle \partial_c \partial_x Q_{c,a}, \varepsilon \rangle_{L^2 \times L^2} & -\langle \partial_{xx} Q_{c,a}, \varepsilon \rangle_{L^2 \times L^2} \end{pmatrix}.$$

It follows from the exponential decay of  $Q_{c,a}$  and its derivatives that

$$|H| \leq A_c \|\varepsilon\|_{L^2 \times L^2}.$$

We can make a further choice of the positive number  $\alpha_c$ , such that the operator norm of the matrix  $D^{-1}H$  is less than  $1/2$ . In this case, the matrix  $M$  is invertible and the operator norm of its inverse is uniformly bounded with respect to  $t$ . Coming back to (3-8), we are led to the estimate

$$|c'(t)| + |a'(t) - c(t)| \leq A_c (|Y(t)| + |Z(t)|). \tag{3-9}$$

It remains to estimate the quantities  $Y$  and  $Z$ . We write

$$\begin{aligned} & \left| \langle \partial_x w_{c,a}, ((v_{c,a} + \varepsilon_v)^2 - 1)(w_{c,a} + \varepsilon_w) - (v_{c,a}^2 - 1)w_{c,a} \rangle_{L^2} \right| \\ &= \left| \langle \partial_x w_{c,a}, (\varepsilon_v^2 + 2v_{c,a}\varepsilon_v)w_{c,a} + \varepsilon_w((\varepsilon_v + v_{c,a})^2 - 1) \rangle_{L^2} \right| \\ &\leq A_c \|\varepsilon\|_{L^2 \times L^2}. \end{aligned}$$

Arguing in the same way for the other terms in  $Y$  and  $Z$ , we obtain

$$|Y| + |Z| = \mathcal{O}(\|\varepsilon\|_{L^2 \times L^2}),$$

which is enough to deduce (3-5) from (3-9).

To complete the proof, we show (2-7). Using the Sobolev embedding theorem of  $H^1(\mathbb{R})$  into  $C^0(\mathbb{R})$ , we can write

$$\max_{x \in \mathbb{R}} v(x, t) \leq \|v_{c(t)}\|_{L^\infty(\mathbb{R})} + \|v(\cdot, t) - v_{c(t),a(t)}\|_{L^\infty(\mathbb{R})} \leq \|v_{c(t)}\|_{L^\infty(\mathbb{R})} + \|\varepsilon(t)\|_{X(\mathbb{R})}.$$

By (2-3),  $\|v_c\|_{L^\infty(\mathbb{R})} < 1$ , so that (2-8) implies that there exists a small positive number  $\gamma_c$  such that  $\|v_{c(t)}\|_{L^\infty(\mathbb{R})} \leq 1 - \gamma_c$ . We obtain

$$\max_{x \in \mathbb{R}} v(x, t) \leq 1 - \gamma_c + \|\varepsilon(t)\|_{X(\mathbb{R})} \leq 1 - \gamma_c + \alpha_c.$$

For  $\alpha_c$  small enough, the estimate (2-7) follows, with  $\sigma_c := -\alpha_c + \gamma_c$ . □

#### 4. Proofs of localization and smoothness of the limit profile

*Proof of Proposition 2.4.* The proof relies on the conservation law for the density of momentum  $vw$ . Let  $R$  and  $t$  be two real numbers, and recall that

$$I_R(t) \equiv I_R^{(v,w)}(t) := \frac{1}{2} \int_{\mathbb{R}} [vw](x + a(t), t) \Phi(x - R) dx,$$

where  $\Phi$  is the function defined on  $\mathbb{R}$  by

$$\Phi(x) := \frac{1}{2}(1 + \text{th}(v_c x)),$$

with  $v_c := \sqrt{1 - c^2}/8$ . First, we deduce from the conservation law for  $vw$  (see Lemma 3.1 in [de Laire and Gravejat 2015] for more details) the identity

$$\begin{aligned} \frac{d}{dt}[I_{R+\sigma t}(t)] &= -(a'(t) + \sigma) \int_{\mathbb{R}} [vw](x + a(t), t) \Phi'(x - R - \sigma t) dx \\ &+ \int_{\mathbb{R}} \left[ v^2 + w^2 - 3v^2 w^2 + \frac{3-v^2}{(1-v^2)^2} (\partial_x v)^2 \right] (x + a(t), t) \Phi'(x - R - \sigma t) dx \\ &+ \int_{\mathbb{R}} [\ln(1 - v^2)](x + a(t), t) \Phi'''(x - R - \sigma t) dx. \end{aligned} \quad (4-1)$$

Our goal is to provide a lower bound for the integrands on the right-hand side of (4-1).

Notice that the function  $\Phi$  satisfies the inequality

$$|\Phi'''| \leq 4v_c^2 \Phi'. \quad (4-2)$$

In view of the bound (2-14) on  $a'(t)$  and the definition of  $\sigma_c$ , we obtain that

$$|a'(t) + \sigma|^2 \leq \frac{9+7c^2}{8}. \quad (4-3)$$

Hence, we deduce

$$\begin{aligned} \frac{d}{dt}[I_{R+\sigma t}(t)] &\geq \int_{\mathbb{R}} \left[ 4v_c^2 \ln(1 - v^2) + v^2 + w^2 - 3v^2 w^2 \right. \\ &\left. + (\partial_x v)^2 - \sqrt{\frac{9+7c^2}{8}} |vw| \right] (x + a(t), t) \Phi'(x - R - \sigma t) dx =: J_1 + J_2. \end{aligned} \quad (4-4)$$

At this step, we decompose the real line into two domains,  $[-R_0, R_0]$  and its complement, where  $R_0$  is to be defined below, and we denote by  $J_1$  and  $J_2$  the value of the integral on the right-hand side of (4-4) on each region. On  $\mathbb{R} \setminus [-R_0, R_0]$ , we bound the integrand pointwise from below by a positive quadratic form in  $(v, w)$ . Exponentially small error terms arise from integration on  $[-R_0, R_0]$ .

For  $|x| \geq R_0$ , using Theorem 2.1 and the Sobolev embedding theorem, and choosing  $\alpha_0$  small enough and  $R_0$  large enough, we obtain

$$|v(x + a(t), t)| \leq |\varepsilon_v(x, t)| + |v_{c(t)}(x)| \leq A_c(\alpha_0 + \exp(-\sqrt{1 - c^2} R_0)) \leq \frac{1}{12} \quad (4-5)$$

for any  $t \in \mathbb{R}$ . Using the fact that  $\ln(1 - s) \geq -2s$  for all  $s \in [0, \frac{1}{2}]$  and introducing (4-5) in (4-4), we obtain

$$J_1 \geq \frac{1-c^2}{8} \int_{|x| \geq R_0} [v^2 + w^2 + (\partial_x v)^2] (x + a(t), t) \Phi'(x - R - \sigma t) dx. \quad (4-6)$$

We next consider the case  $x \in [-R_0, R_0]$ . In that region, we have

$$|x - R - \sigma t| \geq -R_0 + |R + \sigma t|.$$

Hence,

$$\Phi'(x - R - \sigma t) \leq 2v_c e^{2v_c R_0} e^{-2v_c |R + \sigma t|}. \quad (4-7)$$

Since the function  $|\ln(\cdot)|$  is decreasing on  $(0, 1]$ , in view of (2-7) and (4-4),

$$|J_2| \leq A_c \int_{|x| \leq R_0} [v^2 + w^2 + (\partial_x v)^2](x + a(t), t) \Phi'(x - R - \sigma t) dx.$$

Then, by (4-7) and the control on the norm of  $(v, w)$  in  $X(\mathbb{R})$  provided by the conservation of the energy, we obtain

$$|J_2| \leq B_c e^{-2\nu_c |R + \sigma t|}.$$

This finishes the proof of (2-28). It remains to prove (2-29). For that, we distinguish two cases. If  $R \geq 0$ , we integrate (2-28) from  $t = t_0$  to  $t = (t_0 + t_1)/2$ , choosing  $\sigma = \sigma_c$  and  $R = R - \sigma_c t_0$ , and then from  $t = (t_0 + t_1)/2$  to  $t = t_1$  choosing  $\sigma = -\sigma_c$  and  $R = R + \sigma_c t_1$ . If  $R \leq 0$ , we use the same arguments for the reverse choices  $\sigma = -\sigma_c$  and  $\sigma = \sigma_c$ . This implies (2-29), and finishes the proof of Proposition 2.4.  $\square$

*Proof of Proposition 2.9.* Let  $\Psi^*$  and  $v^*$  be the solutions of (2-32)–(2-34) expressed in terms of the hydrodynamical variables  $(v^*, w^*)$  as in (2-30). We split the proof into five steps.

**Step 1.** *There exists a positive number  $A_c$ , depending only on  $c$ , such that*

$$\int_t^{t+1} \int_{\mathbb{R}} |\partial_x \Psi^*(x + a^*(t), s)|^2 e^{\nu_c |x|} dx ds \leq A_c \tag{4-8}$$

for any  $t \in \mathbb{R}$ .

By (2-23) and (2-30),

$$|\Psi^*| \leq A_c (|\partial_x v^*| + |w^*|). \tag{4-9}$$

In view of Proposition 2.6 and the fact that  $|a^*(t) - a^*(s)|$  is uniformly bounded for  $s \in [t - 1, t + 2]$  by (2-22), this yields

$$\int_{t-1}^{t+2} \int_{\mathbb{R}} |\Psi^*(x + a^*(t), s)|^2 e^{2\nu_c |x|} dx ds \leq A_c. \tag{4-10}$$

We define

$$F^* := -\frac{1}{2}(v^*)^2 \Psi^* + \text{Re}(\Psi^*(1 - 2F(v^*, \bar{\Psi}^*))(1 - 2F(v^*, \Psi^*))).$$

We recall that  $\|v^*\|_{L^\infty(\mathbb{R} \times \mathbb{R})} < 1 - \sigma_c$  by (2-23). Using the Cauchy–Schwarz inequality, the Sobolev embedding theorem and the control of the norm in  $X(\mathbb{R})$  provided by the conservation of energy, we have  $F(v^*, \Psi^*) \in L^\infty(\mathbb{R} \times \mathbb{R})$ . Hence,

$$|F^*| \leq A_c |\Psi^*|, \tag{4-11}$$

where  $A_c$  is a positive number depending only on  $c$ . Then, by (4-10),

$$\int_{t-1}^{t+2} \int_{\mathbb{R}} |F^*(x + a^*(t), s)|^2 e^{2\nu_c |x|} dx ds \leq A_c \tag{4-12}$$

for any  $t \in \mathbb{R}$ . Next, by Proposition 2.7, we have

$$\|\Psi^*\|_{L^4([t-1, t+2], L^\infty)} \leq A_c. \tag{4-13}$$

Indeed, we fix  $t \in \mathbb{R}$  and we denote by

$$\begin{aligned}(\Psi_1^0, v_1^0) &:= (\Psi^*(\cdot + a^*(t-1), t-1), v^*(\cdot + a^*(t-1), t-1)), \\ (\Psi_1(s), v_1(s)) &:= (\Psi^*(\cdot + a^*(t-1), t-1+s), v^*(\cdot + a^*(t-1), t-1+s))\end{aligned}$$

the corresponding solution to (2-32)–(2-34). Denote also by

$$\begin{aligned}(\Psi_2^0, v_2^0) &:= (\Psi_{c^*(t-1)}, v_{c^*(t-1)}), \\ (\Psi_2(s), v_2(s)) &:= (\Psi_{c^*(t-1)}(x - c^*(t-1)s), v_{c^*(t-1)}(x - c^*(t-1)s))\end{aligned}$$

the corresponding solution to (2-32)–(2-34), where  $\Psi_{c^*(t)}$  is the solution to (2-32) associated to the soliton  $Q_{c^*(t)}$ . We have, by (2-35),

$$\|\Psi_1(s) - \Psi_2(s)\|_{L^4([0, \tau_c], L^\infty)} \leq A(\|v_1^0 - v_2^0\|_{L^2} + \|\Psi_1^0 - \Psi_2^0\|_{L^2}).$$

Using (2-21), we obtain

$$\|\Psi_1(s) - \Psi_2(s)\|_{L^4([0, \tau_c], L^\infty)} \leq A_c,$$

where  $\tau_c = \tau_c(\|v_1^0\|_{L^2}, \|v_2^0\|_{L^2}, \|\Psi_1^0\|_{L^2}, \|\Psi_2^0\|_{L^2})$  depend only on  $c$ . Since  $[0, 3] \subseteq \bigcup_{0 \leq k \leq 3/\tau_c} [k\tau_c, (k+1)\tau_c]$ , we can infer (4-13) inductively.

In addition, by (4-9), we have

$$\|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^\infty([t-1, t+2], L^2)} \leq A_c. \quad (4-14)$$

Hence, applying the Cauchy–Schwarz inequality to the integral with respect to the time variable, (4-10), (4-13) and (4-14),

$$\begin{aligned}& \int_{t-1}^{t+2} \int_{\mathbb{R}} |\Psi^*(x + a^*(t), s)|^4 e^{\nu_c |x|} dx ds \\ & \leq \int_{t-1}^{t+2} \int_{\mathbb{R}} |\Psi^*(x + a^*(t), s)|^2 e^{\nu_c |x|} dx \|\Psi^*(s)\|_{L^\infty(\mathbb{R})}^2 ds \\ & \leq \|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^4([t-1, t+2], L^2(\mathbb{R}))}^2 \|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^4([t-1, t+2], L^\infty(\mathbb{R}))}^2 \\ & \leq \|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^2([t-1, t+2], L^2(\mathbb{R}))}^2 \|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^\infty([t-1, t+2], L^2(\mathbb{R}))}^2 \\ & \quad \|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^4([t-1, t+2], L^\infty(\mathbb{R}))}^2 \\ & \leq A_c.\end{aligned} \quad (4-15)$$

In order to use Proposition 2.8 on  $\Psi^*$ , it is sufficient to verify

$$\sup_{s \in [t-1, t+2]} \int_{\mathbb{R}} |\Psi^*(x + a^*(t), s)|^2 e^{2\nu_c |x|} dx ds \leq A_c. \quad (4-16)$$

Indeed, using (4-16) and (4-13), we can write

$$\begin{aligned} & \int_{t-1}^{t+2} \int_{\mathbb{R}} |\Psi^*(x + a^*(t), s)|^6 e^{2\nu_c|x|} dx ds \\ & \leq \|\Psi^*(\cdot + a^*(t), \cdot) e^{\nu_c|\cdot|}\|_{L^\infty([t-1, t+2], L^2(\mathbb{R}))}^2 \|\Psi^*(\cdot + a^*(t), \cdot)\|_{L^4([t-1, t+2], L^\infty(\mathbb{R}))}^4 \\ & \leq A_c, \end{aligned} \tag{4-17}$$

which proves that  $\Psi^*$  satisfies the assumptions of Proposition 2.8. Then, we apply Proposition 2.8 with  $u := \Psi^*(\cdot + a^*(t), \cdot + (t + 1/2))$ ,  $T := 1/2$ ,  $F := |u|^2u + F^*(\cdot, t + 1/2)$  and successively  $\lambda := \pm\nu_c$ , and we use (4-10) and (4-12) to obtain (4-8).

Now let us prove (4-16). First, we recall the next lemma stated by Kenig, Ponce and Vega [Kenig et al. 2003].

**Lemma 4.1.** *Let  $a \in [-2, -1]$  and  $b \in [2, 3]$ . Assume that  $u \in C^0([a, b] : L^2(\mathbb{R}))$  is a solution of the inhomogeneous Schrödinger equation*

$$i \partial_t u + \partial_{xx} u = H, \tag{4-18}$$

with  $H \in L^1([a, b] : L^2(e^{\beta x} dx))$ , for some  $\beta \in \mathbb{R}$ , and

$$u_a \equiv u(\cdot, a), u_b \equiv u(\cdot, b) \in L^2(e^{\beta x} dx). \tag{4-19}$$

Then there exists a positive number  $K$  such that

$$\sup_{a \leq t \leq b} \|u(\cdot, t)\|_{L^2(e^{\beta x} dx)} \leq K (\|u_a\|_{L^2(e^{\beta x} dx)} + \|u_b\|_{L^2(e^{\beta x} dx)} + \|H\|_{L^1([a, b], L^2(e^{\beta x} dx))}). \tag{4-20}$$

In order to apply the lemma, we need to verify the existence of numbers  $a$  and  $b$  such that (4-19) holds for  $u := \Psi^*(\cdot + a^*(t), \cdot + t)$  and such that  $H := |u|^2u + F^*(\cdot, \cdot + t) \in L^1([a, b], L^2(e^{\beta x} dx))$  for  $\beta = \pm\nu_c$  respectively and any  $t \in \mathbb{R}$ . Our first claim is a consequence of (4-10) and the Markov inequality. Indeed, there exist  $s_0 \in [-2, -1]$  and  $s_1 \in [2, 3]$  such that

$$\int_{\mathbb{R}} |\Psi^*(x + a^*(t), s_j + t)|^2 e^{2\nu_c|x|} dx \leq A_c \quad \text{for } j = 0, 1.$$

For the second claim, due to (4-12) and the Cauchy–Schwarz estimate, it is sufficient to show that  $|u|^2u \in L^1([-2, 3], L^2(e^{\nu_c|x|} dx))$ . To prove this we use the Cauchy–Schwarz inequality for the time variable, (4-10) and (4-13),

$$\begin{aligned} & \int_{-2}^3 \left( \int_{\mathbb{R}} |\Psi^*(x + a^*(t), s + t)|^6 e^{2\nu_c|x|} dx \right)^{1/2} ds \\ & \leq \|\Psi^*(\cdot + a^*(t), \cdot + t) e^{\nu_c|\cdot|}\|_{L^2([-2, 3], L^2)} \|\Psi^*(\cdot + a^*(t), \cdot + t)\|_{L^4([-2, 3], L^\infty)}^2 \\ & \leq A_c. \end{aligned}$$

Now we may apply Lemma 4.1 with  $a = s_0$  and  $b = s_1$  to deduce (4-16). This finishes the proof of the first step.

In the next step, we prove that (4-8) remains true for all the derivatives of  $\Psi^*$  and  $v^*$ .

**Step 2.** Let  $k \geq 1$ . There exists a positive number  $A_{k,c}$ , depending only on  $k$  and  $c$ , such that

$$\int_t^{t+1} \int_{\mathbb{R}} |\partial_x^k \Psi^*(x + a^*(t), s)|^2 e^{\nu_c |x|} dx ds \leq A_{k,c} \quad (4-21)$$

and

$$\int_t^{t+1} \int_{\mathbb{R}} |\partial_x^k v^*(x + a^*(t), s)|^2 e^{\nu_c |x|} dx \leq A_{k,c} \quad (4-22)$$

for any  $t \in \mathbb{R}$ .

The proof of Step 2 is by induction on  $k \geq 1$ . We are going to differentiate (2-32)  $k$  times with respect to the space variable and write the resulting equation as

$$i \partial_t (\partial_x^k \Psi^*) + \partial_{xx} (\partial_x^k \Psi^*) = R_k(v^*, \Psi^*), \quad (4-23)$$

where  $R_k(v^*, \Psi^*) = \partial_x^k (|\Psi^*|^2 \Psi^*) + \partial_x^k F^*$ . We are going to prove by induction that (4-21), (4-22) and

$$\int_t^{t+1} \int_{\mathbb{R}} |R_k(v^*, \Psi^*)(x + a^*(t), s)|^2 e^{\nu_c |x|} dx ds \leq A_{k,c} \quad (4-24)$$

hold simultaneously for any  $t \in \mathbb{R}$ . Notice that (4-21) implies that  $\partial_x^k \Psi^* \in L^2_{\text{loc}}(\mathbb{R}, L^2(\mathbb{R}))$ , while (4-24) implies that  $R_k(v^*, \Psi^*) \in L^2_{\text{loc}}(\mathbb{R}, L^2(\mathbb{R}))$ . Therefore, if (4-21), (4-22) and (4-24) are established for some  $k \geq 1$ , then applying Proposition 2.8 to  $\partial_x^k \Psi^*$  can be justified by a standard approximation procedure.

For  $k = 1$ , (4-21) is exactly (4-8). Equation (4-22) holds from Proposition 2.6 and the fact that  $|a^*(t) - a^*(s)|$  is uniformly bounded for  $s \in [t - 1, t + 2]$ . Next, we write

$$\begin{aligned} R_1(v^*, \Psi^*) &= -v^* \partial_x v^* \Psi^* - \frac{1}{2} (v^*)^2 \partial_x \Psi^* + \text{Re}(\partial_x \Psi^* (1 - 2F(v^*, \bar{\Psi}^*))) (1 - 2F(v^*, \Psi^*)) \\ &\quad - 2v^* |\Psi^*|^2 (1 - 2F(v^*, \Psi^*)) - 2v^* \Psi^* \text{Re}(\Psi^* (1 - 2F(v^*, \bar{\Psi}^*)) - 2\partial_x (\Psi^* |\Psi^*|^2)). \end{aligned}$$

We will show that

$$\Psi^* \in L^\infty([t - 1, t + 2], L^\infty(\mathbb{R})) \quad (4-25)$$

in order to control the derivative of the cubic nonlinearity by  $|\partial_x \Psi^*|$ , and then we will use the fact that  $F(v^*, \Psi^*) \in L^\infty(\mathbb{R} \times \mathbb{R})$ ,  $\|v^*\|_{L^\infty(\mathbb{R} \times \mathbb{R})} < 1$  and the second equation in (2-34) to get

$$R_1(v^*, \Psi^*) \leq K (|\partial_x \Psi^*| + |\partial_x v^*| |\Psi^*| + |\Psi^*|^2). \quad (4-26)$$

Let us prove (4-25). We define the function  $H$  on  $\mathbb{R}$  by

$$H(s) := \frac{1}{2} \int_{\mathbb{R}} (|\partial_x \Psi^*(x, s)|^2 - |\Psi^*(x, s)|^4) dx.$$

We differentiate it with respect to  $s$ , integrate by parts and use (2-32) to obtain

$$\begin{aligned} H'(s) &= -\operatorname{Re}\left(\int_{\mathbb{R}} \partial_s \Psi^*(x, s) \left[\overline{\partial_{xx} \Psi^* + 2\Psi^*|\Psi^*|^2}\right](x, s) dx\right) \\ &= \operatorname{Re}\left(\int_{\mathbb{R}} \partial_s \Psi^*(x, s) F^*(x, s) dx\right) \\ &\leq \|\partial_s \Psi^*(s)\|_{H^{-1}(\mathbb{R})} \|F^*(s)\|_{H^1(\mathbb{R})}. \end{aligned} \tag{4-27}$$

We have

$$|\partial_x F^*| \leq K(|\partial_x \Psi^*| + |\partial_x v^*| |\Psi^*| + |\Psi^*|^2),$$

using the fact that  $F(v^*, \Psi^*) \in L^\infty(\mathbb{R} \times \mathbb{R})$ ,  $\|v^*\|_{L^\infty(\mathbb{R} \times \mathbb{R})} < 1$  and the second equation in (2-34).

Hence, by (4-8), (4-10), (4-15) and the fact that  $|\partial_x v^*| \leq |\Psi^*|$  on  $\mathbb{R} \times \mathbb{R}$ , we obtain

$$\|\partial_x F^*\|_{L^2([t-1, t+2], L^2(\mathbb{R}))} \leq A_c. \tag{4-28}$$

On the other hand, we infer

$$\|\partial_s \Psi^*\|_{L^2([t-1, t+2], H^{-1}(\mathbb{R}))} \leq A_c \tag{4-29}$$

from (2-32), (4-8), (4-12) and the fact that  $\Psi^* \in L^4([t-1, t+2], L^\infty(\mathbb{R})) \cap L^8([t-1, t+2], L^4(\mathbb{R}))$ .

Next, we integrate (4-27) between  $t-1$  and  $t+2$  and apply the Cauchy–Schwarz inequality to obtain  $H \in W^{1,1}([t-1, t+2])$  for all  $t \in \mathbb{R}$  using (4-28) and (4-29). Notice that all these computations can be justified by a standard approximation procedure. This yields, by the Sobolev embedding theorem, that  $H \in L^\infty([t-1, t+2])$ . We conclude that the derivative  $\partial_x \Psi^* \in L^\infty([t-1, t+2], L^2(\mathbb{R}))$ . Indeed, we can use the Gagliardo–Nirenberg inequality and the fact that  $\Psi^*$  is uniformly bounded in  $L^2(\mathbb{R})$  by a positive number to write

$$\begin{aligned} H(s) &\geq \frac{1}{2} \int_{\mathbb{R}} |\partial_x \Psi^*(x, s)|^2 dx - A \|\Psi^*(s)\|_{L^2(\mathbb{R})}^3 \|\partial_x \Psi^*(\cdot)\|_{L^2(\mathbb{R})} \\ &\geq \frac{1}{2} \int_{\mathbb{R}} |\partial_x \Psi^*(x, s)|^2 dx - AK^3 \|\partial_x \Psi^*(\cdot)\|_{L^2(\mathbb{R})}. \end{aligned}$$

The function  $x \mapsto \frac{1}{2}x^2 - AM^3x$  diverges to  $+\infty$  when  $x$  goes to  $+\infty$ . Since  $H$  is bounded, we infer that  $\|\partial_x \Psi^*(\cdot)\|_{L^2(\mathbb{R})}$  is uniformly bounded on  $[t-1, t+2]$  for all  $t \in \mathbb{R}$ . This finishes the proof of (4-25) by the Sobolev embedding theorem. Then, by (4-26), (4-24) for  $k = 1$  is a consequence of (4-8), (4-15) and the fact that  $|\partial_x v^*| \leq |\Psi^*|$  on  $\mathbb{R} \times \mathbb{R}$ .

Assume now that (4-21), (4-22) and (4-24) are satisfied for any integer  $1 \leq k \leq k_0$  and any  $t \in \mathbb{R}$ . Let us prove these three estimates for  $k = k_0 + 1$ . We apply Proposition 2.8 with  $u := \partial_x^{k_0} \Psi^*(\cdot + a^*(t), \cdot + (t+1/2))$ ,  $T := 1/2$  and successively  $\lambda := \pm v_c$ . In view of (4-21), (4-23), (4-24) and the fact that  $|a^*(t) - a^*(s)|$  is uniformly bounded for  $s \in [t-1, t+2]$ , this yields

$$\int_t^{t+1} \int_{\mathbb{R}} |\partial_x^{k_0+1} \Psi^*(x + a^*(t), s)|^2 e^{\nu_c |x|} dx ds \leq A_c, \tag{4-30}$$

so that (4-21) is satisfied for  $k = k_0 + 1$ .

Let  $k \in \{1, \dots, k_0\}$ . We use the induction hypothesis and (4-30) to infer that

$$\partial_x^{k-1} \Psi^* \in L^2([t, t + 1], H^2(\mathbb{R})).$$

Also, we have

$$\partial_x^{k-1} \Psi^* \in H^1([t, t + 1], L^2(\mathbb{R}))$$

using (4-23) and (4-24). This yields, by interpolation,

$$\partial_x^{k-1} \Psi^* \in H^{2/3}([t, t + 1], H^{2/3}(\mathbb{R})).$$

Hence, using the Sobolev embedding theorem, we obtain

$$\partial_x^{k-1} \Psi^* \in L^\infty([t, t + 1], L^\infty(\mathbb{R})) \quad \text{for all } t \in \mathbb{R}. \tag{4-31}$$

On the other hand, since  $|\partial_x v^*| \leq |\Psi^*|$ , we have, by (4-25), that  $\partial_x v^* \in L^\infty([t, t + 1], L^\infty(\mathbb{R}))$ . For  $k \in \{2, \dots, k_0\}$ , we differentiate the second equation in (2-34)  $k$  times and we use (4-31) to obtain

$$|\partial_x^k v^*| \leq K \left( \sum_{j=1}^{k-1} |\partial_x^j \Psi^*| + \sum_{j=0}^{k-2} |\partial_x^j v^*| \right), \tag{4-32}$$

where  $K$  is a positive constant. By induction we infer from (4-31) that

$$\partial_x^k v^* \in L^\infty([t, t + 1], L^\infty(\mathbb{R})) \quad \text{for all } t \in \mathbb{R}, \tag{4-33}$$

for all  $k \in \{2, \dots, k_0\}$ . Then, we just compute explicitly  $R_{k_0+1}(v^*, \Psi^*)$  and we use (4-31) and (4-33) to obtain

$$|R_{k_0+1}(v^*, \Psi^*)| \leq A_{k_0+1,c,K} \left( \sum_{j=0}^{k_0+1} |\partial_x^j \Psi^*| + \sum_{j=1}^{k_0} |\partial_x^j v^*| \right).$$

Hence, by (4-21) for all  $k \leq k_0$ , (4-22) and (4-30), we obtain (4-24) for  $k = k_0 + 1$ . Finally, we introduce (4-21) for all  $k \leq k_0 + 1$  and (4-22) for all  $k \leq k_0$  into (4-32) to deduce (4-22) for  $k = k_0 + 1$ . This finishes the proof of this step.

In order to finish the proof of Proposition 2.9, we now turn these  $L^2_{\text{loc}}$  in time estimates into  $L^\infty$  in time estimates, and then into uniform estimates.

**Step 3.** *Let  $k \geq 0$ . There exists a positive number  $A_{k,c}$ , depending only on  $k$  and  $c$ , such that*

$$\int_{\mathbb{R}} |\partial_x^k \Psi^*(x + a^*(t), t)|^2 e^{v_c|x|} dx \leq A_{k,c} \tag{4-34}$$

for any  $t \in \mathbb{R}$ . In particular, we have

$$\left\| \partial_x^k \Psi^*(\cdot + a^*(t), t) e^{(v_c/2)|\cdot|} \right\|_{L^\infty(\mathbb{R})} \leq A_{k,c} \tag{4-35}$$

for any  $t \in \mathbb{R}$ , and for a possibly different choice of the positive constant  $A_{k,c}$ .

Here, we use the Sobolev embedding theorem in time and (4-23) for the proof. By the Sobolev embedding theorem, we have

$$\begin{aligned} \|\partial_x^k \Psi^*(\cdot + a^*(t), t)e^{(v_c/2)|\cdot|}\|_{L^2(\mathbb{R})}^2 &\leq K \left( \|\partial_s(\partial_x^k \Psi^*(\cdot + a^*(t), s)e^{(v_c/2)|\cdot|})\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \right. \\ &\quad \left. + \|\partial_x^k \Psi^*(\cdot + a^*(t), s)e^{(v_c/2)|\cdot|}\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \right), \end{aligned}$$

while, by (4-23),

$$\begin{aligned} \|\partial_s(\partial_x^k \Psi^*(\cdot + a^*(t), s)e^{(v_c/2)|\cdot|})\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 &\leq 2 \left( \|\partial_x^{k+2} \Psi^*(\cdot + a^*(t), s)e^{(v_c/2)|\cdot|}\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \right. \\ &\quad \left. + \|\mathcal{R}_k(\Psi^*)(\cdot + a^*(t), s)e^{(v_c/2)|\cdot|}\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \right), \end{aligned}$$

so that we finally deduce (4-34) from (4-21) and (4-24). The estimate (4-35) follows from applying the Sobolev embedding theorem to (4-34).

The function  $v^*$  satisfies a similar inequality:

**Step 4.** Let  $k \in \mathbb{N}$ . There exists a positive number  $A_{k,c}$ , depending only on  $k$  and  $c$ , such that

$$\int_{\mathbb{R}} (\partial_x^k v^*(x + a^*(t), t))^2 e^{v_c|x|} dx \leq A_{k,c} \tag{4-36}$$

and

$$\|\partial_x^k v^*(\cdot + a^*(t), t)e^{(v_c/2)|\cdot|}\|_{L^\infty(\mathbb{R})} \leq A_{k,c} \tag{4-37}$$

for any  $t \in \mathbb{R}$ .

The proof is similar to the proof of Step 3 using the first equation in (2-34) instead of (2-32). We use the Sobolev embedding theorem to write

$$\begin{aligned} \|\partial_x^k v^*(\cdot + a^*(t), t)e^{v_c|\cdot|}\|_{L^2(\mathbb{R})}^2 &\leq K \left( \|\partial_s(\partial_x^k v^*(\cdot + a^*(t), s)e^{v_c|\cdot|})\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \right. \\ &\quad \left. + \|\partial_x^k v^*(\cdot + a^*(t), s)e^{v_c|\cdot|}\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \right). \end{aligned}$$

By the first equation in (2-34), (4-21), (4-23) and (4-33), we have

$$\|\partial_s(\partial_x^k v^*(\cdot + a^*(t), s)e^{v_c|\cdot|})\|_{L^2([t-1, t+1], L^2(\mathbb{R}))}^2 \leq A_c.$$

This leads to (4-36). The uniform bound in (4-37) is then a consequence of the Sobolev embedding theorem.

Finally, we provide the estimates for the function  $w^*$ .

**Step 5.** Let  $k \in \mathbb{N}$ . There exists a positive number  $A_{k,c}$ , depending only on  $k$  and  $c$ , such that

$$\int_{\mathbb{R}} |\partial_x^k w^*(x + a^*(t), t)|^2 e^{v_c|x|} dx \leq A_{k,c} \tag{4-38}$$

and

$$\|\partial_x^k w^*(\cdot + a^*(t), t)e^{(v_c/2)|\cdot|}\|_{L^\infty(\mathbb{R})} \leq A_{k,c} \tag{4-39}$$

for any  $t \in \mathbb{R}$ .

The proof relies on the last two steps. First, we write

$$v^* \Psi^* = -\frac{1}{2} \partial_x ((1 - (v^*)^2)^{1/2} \exp i\theta^*).$$

Since  $(1 - v^*(x, t)^2)^{1/2} \exp i\theta^*(x, t) \rightarrow 1$  as  $x \rightarrow -\infty$  for any  $t \in \mathbb{R}$ , we obtain the formula

$$2F(v^*, \Psi^*) = 1 - (1 - (v^*)^2)^{1/2} \exp i\theta^*. \quad (4-40)$$

Hence, using (2-30), we have

$$w^* = 2 \operatorname{Im} \left( \frac{\Psi^* (1 - 2F(v^*, \Psi^*))}{1 - (v^*)^2} \right). \quad (4-41)$$

Combining (2-7) and (4-40), we recall that

$$\frac{|1 - 2F(v^*, \Psi^*)|}{1 - (v^*)^2} \leq A_c. \quad (4-42)$$

Hence, we obtain

$$|w^*| \leq A_c |\Psi^*|.$$

Then, (4-38) and (4-39) follow from (4-34) and (4-35) for  $k = 0$ . For  $k \geq 1$ , we differentiate (4-41)  $k$  times with respect to the space variable, and using (4-35), (4-37) and (4-42), we are led to

$$|\partial_x^k w^*| \leq A_{k,c} \left( \sum_{j=0}^k |\partial_x^j \Psi^*| + \sum_{j=1}^{k-1} |\partial_x^j v^*| \right).$$

We finish the proof of this step using Steps 3 and 4. This completes the proof of Proposition 2.9.  $\square$

## 5. Proof of the Liouville theorem

*Proof of Proposition 2.10.* First, by (2-38) and the explicit formula for  $v_c$  and  $w_c$  in (2-3), there exists a positive number  $A_{k,c}$  such that

$$\int_{\mathbb{R}} ((\partial_x^k \varepsilon_v^*(x, t))^2 + (\partial_x^k \varepsilon_w^*(x, t))^2) e^{\nu_c |x|} dx \leq A_{k,c}, \quad (5-1)$$

for any  $k \in \mathbb{N}$  and any  $t \in \mathbb{R}$ . In view of the formulae of  $\mathcal{H}_c$  in (A-42) and for  $u^*$  in (2-41), a similar estimate holds for  $u^*$ , for a possibly different choice of the constant  $A_{k,c}$ . As a consequence, we are allowed to differentiate with respect to the time variable the quantity

$$\mathcal{I}^*(t) := \int_{\mathbb{R}} x u_1^*(x, t) u_2^*(x, t) dx$$

on the left-hand side of (2-43). Moreover, we can compute

$$\begin{aligned} \frac{d}{dt}(\mathcal{I}^*) &= -2 \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(\partial_x u^*), u^* \rangle_{\mathbb{R}^2} + \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(J\mathcal{R}_{c^*}\varepsilon^*), u^* \rangle_{\mathbb{R}^2} \\ &\quad - (c^*)' \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(\partial_c Q_{c^*}), u^* \rangle_{\mathbb{R}^2} + (c^*)' \int_{\mathbb{R}} \mu \langle \partial_c \mathcal{H}_{c^*}(\varepsilon^*), u^* \rangle_{\mathbb{R}^2} \\ &\quad + ((a^*)' - c^*) \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(\partial_x \varepsilon^*), u^* \rangle_{\mathbb{R}^2}, \end{aligned} \tag{5-2}$$

where we have set  $\mu(x) = x$  for any  $x \in \mathbb{R}$ .

At this stage, we split the proof into five steps. The proof of these steps is similar to the proof of Proposition 7 in [B ethuel et al. 2015].

**Step 1.** *There exist two positive numbers  $A_1$  and  $R_1$ , depending only on  $c$ , such that*

$$\mathcal{I}_1^*(t) := -2 \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(\partial_x u^*), u^* \rangle_{\mathbb{R}^2} \geq \frac{1-c^2}{8} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 - A_1 \|u^*(\cdot, t)\|_{X(B(0, R_1))}^2 \tag{5-3}$$

for any  $t \in \mathbb{R}$ .

We introduce the explicit formula of the operator  $\mathcal{H}_{c^*}$  in the definition of  $\mathcal{I}_1^*(t)$  to obtain

$$\begin{aligned} \mathcal{I}_1^*(t) &= 2 \int_{\mathbb{R}} \mu \partial_x \left( \frac{\partial_{xx} u_1^*}{1-v_{c^*}^2} \right) u_1^* - 2 \int_{\mathbb{R}} \mu (1-(c^*)^2 - (5+(c^*)^2)v_{c^*}^2 + 2v_{c^*}^4) \frac{\partial_x u_1^*}{(1-v_{c^*}^2)^2} u_1^* \\ &\quad + 2 \int_{\mathbb{R}} \mu c^* \frac{1+v_{c^*}^2}{1-v_{c^*}^2} (\partial_x u_2^*) u_1^* - 2 \int_{\mathbb{R}} \mu (c^*)^2 \frac{(1+v_{c^*}^2)^2}{(1-v_{c^*}^2)^3} (\partial_x u_1^*) u_1^* \\ &\quad + 2 \int_{\mathbb{R}} \mu c^* \frac{1+v_{c^*}^2}{1-v_{c^*}^2} (\partial_x u_1^*) u_2^* - 2 \int_{\mathbb{R}} \mu (1-v_{c^*}^2) (\partial_x u_2^*) u_2^*. \end{aligned}$$

Integrating each term by parts, we obtain

$$\mathcal{I}_1^*(t) = \int_{\mathbb{R}} \iota_1^*(x, t) dx,$$

with

$$\begin{aligned} \iota_1^* &= \left( \frac{2}{1-v_{c^*}^2} + 2x \frac{\partial_x v_{c^*} v_{c^*}}{1-v_{c^*}^2} \right) (\partial_x u_1^*)^2 - 2c^* \left( \frac{1+v_{c^*}^2}{1-v_{c^*}^2} + \frac{4x \partial_x v_{c^*} v_{c^*}}{(1-v_{c^*}^2)^2} \right) u_2^* u_1^* \\ &\quad + (1-v_{c^*}^2 - 2x \partial_x v_{c^*} v_{c^*}) (u_2^*)^2 + \frac{1+2((c^*)^2-3)v_{c^*}^2 + (2(c^*)^2-3)v_{c^*}^4 - 2v_{c^*}^6}{(1-v_{c^*}^2)^3} (u_1^*)^2 \\ &\quad + 4x \partial_x v_{c^*} v_{c^*} \frac{((c^*)^2-3) + (2(c^*)^2-3)v_{c^*}^2 - 3v_{c^*}^4}{(1-v_{c^*}^2)^4} (u_1^*)^2. \end{aligned}$$

Let  $\delta$  be a small positive number. We next use the exponential decay of the function  $v_c$  and its derivatives to guarantee the existence of a radius  $R$ , depending only on  $c$  and  $\delta$  (in view of the bound on  $c^* - c$  in

(2-21)), such that

$$v_1^*(x, t) \geq (2 - \delta)(\partial_x u_1^*)^2(x, t) + \left(\frac{1 - c^2}{4} - \delta\right)((u_1^*)^2(x, t) + (u_2^*)^2(x, t))$$

when  $|x| \geq R$ .

Then, we choose  $\delta$  small enough and fix the number  $R_1$  according to the value of the corresponding  $R$ , to obtain

$$\int_{|x| \geq R_1} v_1^*(x, t) dx \geq \frac{1 - c^2}{8} \int_{|x| \geq R_1} ((\partial_x u_1^*(x, t))^2 + u_1^*(x, t)^2 + u_2^*(x, t)^2) dx. \quad (5-4)$$

On the other hand, it follows from (2-3), and again (2-8), that

$$\int_{|x| \leq R_1} v_1^*(x, t) dx \geq \left(\frac{1 - c^2}{8} - A_1\right) \int_{|x| \leq R_1} ((\partial_x u_1^*(x, t))^2 + u_1^*(x, t)^2 + u_2^*(x, t)^2) dx$$

for a positive number  $A_1$  depending only on  $c$ . Combining with (5-4), we obtain (5-3).

**Step 2.** *There exist two positive numbers  $A_2$  and  $R_2$ , depending only on  $c$ , such that*

$$|\mathcal{I}_2^*(t)| := \left| \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(J\mathcal{R}_{c^*}\varepsilon^*), u^* \rangle_{\mathbb{R}^2} \right| \leq \frac{1 - c^2}{64} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 + A_2 \|u^*(\cdot, t)\|_{X(B(0, R_2))}^2 \quad (5-5)$$

for any  $t \in \mathbb{R}$ .

We refer to the proof of Step 2 in the proof of Proposition 7 in [Béthuel et al. 2015] for more details.

We infer the next step from (2-9), (2-57), the explicit formula of  $\mathcal{H}_{c^*}$  in (A-42) and the exponential decay of the function  $\partial_c Q_{c^*}$  and its derivatives.

**Step 3.** *There exist two positive numbers  $A_3$  and  $R_3$ , depending only on  $c$ , such that*

$$|\mathcal{I}_4^*(t)| := \left| (c^*)' \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(\partial_c Q_{c^*}), u^* \rangle_{\mathbb{R}^2} \right| \leq \frac{1 - c^2}{64} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 + A_3 \|u^*(\cdot, t)\|_{X(B(0, R_3))}^2 \quad (5-6)$$

for any  $t \in \mathbb{R}$ .

We decompose the real line into two regions,  $[-R, R]$  and its complement, for any  $R > 0$ . We use the fact that  $|x| \leq e^{\nu_c|x|/4}$  for all  $|x| \geq R$ , to write

$$\begin{aligned} |\mathcal{I}_4^*(t)| \leq R |(c^*)'(t)| \int_{|x| \leq R} |\mathcal{H}_{c^*(t)}(\partial_c Q_{c^*(t)})(x)| |u^*(x, t)| dx \\ + \delta |(c^*)'(t)| \int_{|x| \geq R} |\mathcal{H}_{c^*(t)}(\partial_c Q_{c^*(t)})(x)| |u^*(x, t)| e^{\nu_c|x|/4} dx \end{aligned}$$

for any  $t \in \mathbb{R}$ . We deduce from (2-9), the explicit formula of  $\mathcal{H}_{c^*}$  in (A-42) and the exponential decay of the function  $\partial_c Q_{c^*}$  and its derivatives that

$$|\mathcal{I}_4^*(t)| \leq A_c \left( R \|u^*(\cdot, t)\|_{X(B(0, R))} + \delta \|u^*(\cdot, t)\|_{X(\mathbb{R})} \right) \|\varepsilon^*(\cdot, t)\|_{L^2(\mathbb{R})^2}$$

for any  $t \in \mathbb{R}$ . Hence, by (2-57),

$$|\mathcal{I}_4^*(t)| \leq A_c \left( \frac{R^2}{\delta} \|u^*(\cdot, t)\|_{X(B(0, R))}^2 + 2\delta \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 \right).$$

We choose  $\delta$  so that  $2A_c\delta \leq (1 - c^2)/64$ , and we denote by  $R_4$  the corresponding number  $R$ , to obtain (5-6), with  $A_4 = A_c R_4^2/\delta$ .

Similarly, we use (2-9), (2-21) and (2-57) to obtain:

**Step 4.** *There exists two positive numbers  $A_4$  and  $R_4$ , depending only on  $c$ , such that*

$$|\mathcal{I}_3^*(t)| := \left| (c^*)' \int_{\mathbb{R}} \mu \langle \partial_c \mathcal{H}_{c^*}(\varepsilon^*), u^* \rangle_{\mathbb{R}^2} \right| \leq \frac{1 - c^2}{64} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 + A_4 \|u^*(\cdot, t)\|_{X(B(0, R_4))}^2 \tag{5-7}$$

for any  $t \in \mathbb{R}$ .

The last step follows from an argument as in Step 3.

**Step 5.** *There exist two positive numbers  $A_5$  and  $R_5$ , depending only on  $c$ , such that*

$$\begin{aligned} |\mathcal{I}_5^*(t)| &:= \left| ((a^*)' - c^*) \int_{\mathbb{R}} \mu \langle \mathcal{H}_{c^*}(\partial_x \varepsilon^*), u^* \rangle_{\mathbb{R}^2} \right| \\ &\leq \frac{1 - c^2}{64} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 + A_5 \|u^*(\cdot, t)\|_{X(B(0, R_5))}^2 \end{aligned} \tag{5-8}$$

for any  $t \in \mathbb{R}$ .

Finally, combining the estimates in Steps 1 to 5 with the identity (5-2), we obtain

$$\frac{d}{dt}(\mathcal{I}^*(t)) \geq \frac{1 - c^2}{16} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2 - (A_1 + A_2 + A_3 + A_4 + A_5) \|u^*(\cdot, t)\|_{X(B(0, R_*))}^2,$$

allowing us to conclude the proof of (2-43) with

$$\begin{aligned} R_* &= \max\{R_1, R_2, R_3, R_4, R_5\}, \\ A_* &= A_1 + A_2 + A_3 + A_4 + A_5. \end{aligned} \quad \square$$

*Proof of Lemma 2.11.* When  $u \in H^3(\mathbb{R}) \times H^1(\mathbb{R})$ , the function  $\partial_x u$  is in the space  $H^2(\mathbb{R}) \times L^2(\mathbb{R})$  which is the domain of  $\mathcal{H}_c$ . The scalar product on the right-hand side of (2-46) is well-defined in view of (2-45). Next, we use the formula for  $\mathcal{H}_c$  in (A-42) to express  $G_c(u)$  as

$$\begin{aligned} &\langle SM_c u, \mathcal{H}_c(-2\partial_x u) \rangle_{L^2(\mathbb{R})^2} \\ &= 2 \int_{\mathbb{R}} \frac{\partial_x v_c}{v_c} \left( \frac{1 - c^2 - (5 + c^2)v_c^2 + 2v_c^4}{(1 - v_c^2)^2} + c^2 \frac{(1 + v_c^2)^2}{(1 - v_c^2)^3} - 2c^2 \frac{v_c^2(1 + v_c^2)}{(1 - v_c^2)^3} \right) u_1 \partial_x u_1 \\ &\quad - 2 \int_{\mathbb{R}} \frac{\partial_x v_c}{v_c} \partial_x \left( \frac{\partial_{xx} u_1}{1 - v_c^2} \right) + 2 \int_{\mathbb{R}} \frac{\partial_x v_c (1 - v_c^2)}{v_c} u_2 \partial_x u_2 \\ &\quad + 2c \int_{\mathbb{R}} \left( 2 \frac{v_c \partial_x v_c}{1 - v_c^2} u_1 \partial_x u_2 - \frac{\partial_x v_c (1 + v_c^2)}{v_c (1 - v_c^2)} \partial_x (u_1 u_2) \right). \end{aligned} \tag{5-9}$$

We recall that  $v_c$  solves the equation

$$\partial_{xx} v_c = (1 - c^2 - 2v_c^2)v_c, \tag{5-10}$$

which leads to

$$(\partial_x v_c)^2 = (1 - c^2 - v_c^2)v_c^2 \quad \text{and} \quad \partial_x \left( \frac{\partial_x v_c}{v_c} \right) = -v_c^2. \tag{5-11}$$

Then, the third integral on the right-hand side of (5-9) can be written as

$$2 \int_{\mathbb{R}} \frac{\partial_x v_c (1 - v_c^2)}{v_c} u_2 \partial_x u_2 = \int_{\mathbb{R}} \mu_c u_2^2, \quad (5-12)$$

with  $\mu_c := 2(\partial_x v_c)^2 + (1 - v_c^2)v_c^2$ . Similarly, the last integral is given by

$$\int_{\mathbb{R}} \left( 2 \frac{v_c \partial_x v_c}{1 - v_c^2} u_1 \partial_x u_2 - \frac{\partial_x v_c (1 + v_c^2)}{v_c (1 - v_c^2)} \partial_x (u_1 u_2) \right) = - \int_{\mathbb{R}} \left( v_c^2 u_1 u_2 + 2 \frac{v_c \partial_x v_c}{1 - v_c^2} u_2 \partial_x u_1 \right). \quad (5-13)$$

Combining (5-12) and (5-13) with (5-9), we obtain the identity

$$\langle SM_c u, \mathcal{H}_c(-2\partial_x u) \rangle_{L^2(\mathbb{R})^2} = I + \int_{\mathbb{R}} \mu_c \left( u_2 - \frac{c v_c^2}{\mu_c} u_1 - \frac{2c v_c \partial_x v_c}{\mu_c (1 - v_c^2)} \partial_x u_1 \right)^2,$$

where

$$\begin{aligned} I = \int_{\mathbb{R}} 2 \left( \frac{\partial_x v_c}{v_c} \left( \frac{1 - c^2 - (5 + c^2)v_c^2 + 2v_c^4}{(1 - v_c^2)^2} + c^2 \frac{1 + v_c^2}{(1 - v_c^2)^2} \right) - 2c^2 \frac{v_c^3 \partial_x v_c}{\mu_c (1 - v_c^2)} \right) u_1 \partial_x u_1 \\ - \int_{\mathbb{R}} \frac{\partial_x v_c}{v_c} u_1 \partial_x \left( \frac{\partial_{xx} u_1}{1 - v_c^2} \right) - c^2 \int_{\mathbb{R}} \frac{v_c^4}{\mu_c} u_1^2 - 4c^2 \int_{\mathbb{R}} \frac{(\partial_x v_c)^2 v_c^2}{\mu_c (1 - v_c^2)^2} (\partial_x u_1)^2. \end{aligned}$$

Using (5-10) and (5-11), we finally deduce that

$$I = \frac{3}{2} \int_{\mathbb{R}} \frac{v_c^4}{\mu_c} \left( \partial_x u_1 - \frac{\partial_x v_c}{v_c} u_1 \right)^2,$$

which finishes the proof of (2-46).  $\square$

*Proof of Proposition 2.12.* We first rely on (2-3) and (2-46) to check that the quadratic form  $G_c$  is well-defined and continuous on  $X(\mathbb{R})$ . Next, setting

$$v = (v_c u_1, v_c u_2), \quad (5-14)$$

and using (5-10), we can express it as

$$G_c(u) = K_c(v) := \int_{\mathbb{R}} \frac{v_c^2}{\mu_c} \left( \partial_x v_1 - \frac{2\partial_x v_c}{v_c} v_1 \right)^2 + \int_{\mathbb{R}} \frac{\mu_c}{v_c^2} \left( v_2 + \frac{c\lambda_c}{\mu_c (1 - v_c^2)} v_1 - 2 \frac{c v_c \partial_x v_c}{\mu_c (1 - v_c^2)} \partial_x v_1 \right)^2, \quad (5-15)$$

where we have set  $\lambda_c := -\mu_c + 4(\partial_x v_c)^2$ . From (2-48) and (5-14) we deduce that

$$\text{Ker}(K_c) = \text{Span}(v_c Q_c). \quad (5-16)$$

Let  $w$  be the pair defined in the following way

$$w = \left( v_1, v_2 - 2 \frac{c v_c \partial_x v_c}{\mu_c (1 - v_c^2)} \partial_x v_1 \right).$$

We compute

$$K_c(v) = \langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2}, \quad (5-17)$$

with

$$\mathcal{T}_c(w) = \begin{pmatrix} -3\partial_x \left( \frac{v_c^2}{\mu_c} \partial_x w_1 \right) + \left( \frac{8v_c^4(\partial_x v_c)^2 - 2v_c^6(1-v_c^2)}{\mu_c^2} + \frac{4(\partial_x v_c)^2}{\mu_c} + \frac{c^2(2c^2-1+v_c^2)v_c^2}{\mu_c(1-v_c^2)} \right) w_1 - \frac{c(2c^2-1+v_c^2)}{(1-v_c^2)} w_2 \\ - \frac{c(2c^2-1+v_c^2)}{(1-v_c^2)} w_1 + \frac{\mu_c}{v_c^2} w_2 \end{pmatrix}. \tag{5-18}$$

The operator  $\mathcal{T}_c$  in (5-18) is self-adjoint on  $L^2(\mathbb{R})^2$ , with domain  $\text{Dom}(\mathcal{T}_c) = H^2(\mathbb{R}) \times L^2(\mathbb{R})$ . In addition, combining (5-15) with (5-17) we deduce that  $\mathcal{T}_c$  is nonnegative, with a kernel equal to

$$\text{Ker}(\mathcal{T}_c) = \text{Span} \left\{ \left( v_c^2, \frac{2cv_c^2(\partial_x v_c)^2}{\mu_c(1-v_c^2)} \right) \right\}.$$

At this stage, we divide the proof into three steps.

**Step 1.** *Let  $c \in (-1, 1) \setminus \{0\}$ . There exists a positive number  $\Lambda_1$ , depending continuously on  $c$ , such that*

$$\langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2} \geq \Lambda_1 \int_{\mathbb{R}} (w_1^2 + w_2^2), \tag{5-19}$$

for any pair  $w \in X^1(\mathbb{R})$  such that

$$\left\langle w, \left( v_c^2, \frac{2cv_c^2(\partial_x v_c)^2}{\mu_c(1-v_c^2)} \right) \right\rangle_{L^2(\mathbb{R})^2} = 0. \tag{5-20}$$

We claim that the essential spectrum of  $\mathcal{T}_c$  is given by

$$\sigma_{\text{ess}}(\mathcal{T}_c) = [\tau_c, +\infty), \tag{5-21}$$

with

$$\tau_c = \tau_{1,c} - \frac{1}{2} \tau_{2,c}^{1/2} > 0. \tag{5-22}$$

Here, we have set

$$\tau_{1,c} = \frac{4(1-c^2) + c^2(2c^2-1)^2}{2(3-2c^2)} + \frac{3-2c^2}{2}$$

and

$$\tau_{2,c} = \left( \frac{4(1-c^2) + c^2(2c^2-1)^2}{3-2c^2} - (3-2c^2) \right)^2 + 4c^2(2c^2-1)^2.$$

In particular, 0 is an isolated eigenvalue in the spectrum of  $\mathcal{T}_c$ . The inequality (5-19) follows with  $\Lambda_1$  either equal to  $\tau_c$ , or to the smallest positive eigenvalue of  $\mathcal{T}_c$ . In view of the analytic dependence on  $c$  of the operator  $\mathcal{T}_c$ ,  $\Lambda_1$  depends continuously on  $c$ .

Now, let us prove (5-21). We rely on the Weyl criterion. It follows from (2-47) and (5-10) that

$$\frac{\mu_c(x)}{v_c^2(x)} \rightarrow 3 - 2c^2 \quad \text{and} \quad \frac{(\partial_x v_c)^2(x)}{\mu_c(x)} \rightarrow \frac{1-c^2}{3-2c^2}$$

as  $x \rightarrow \pm\infty$ . Coming back to (5-18), we introduce the operator  $\mathcal{T}_\infty$  given by

$$\mathcal{T}_\infty(w) = \begin{pmatrix} -\frac{3}{3-2c^2} \partial_{xx} w_1 + \frac{4(1-c^2)+c^2(2c^2-1)^2}{3-2c^2} w_1 - c(2c^2-1)w_2 \\ -c(2c^2-1)w_1 + (3-2c^2)w_2 \end{pmatrix}.$$

By the Weyl criterion, the essential spectrum of  $\mathcal{T}_c$  is equal to the spectrum of  $\mathcal{T}_\infty$ .

We next apply again the Weyl criterion to establish that a real number  $\lambda$  belongs to the spectrum of  $\mathcal{T}_\infty$  if and only if there exists a complex number  $\xi$  such that

$$\lambda^2 - \left( \frac{3}{3-2c^2} |\xi|^2 + \frac{4(1-c^2)+c^2(2c^2-1)^2}{3-2c^2} + 3-2c^2 \right) \lambda + 3|\xi|^2 + 4(1-c^2) = 0.$$

This is the case if and only if

$$\begin{aligned} \lambda = & \frac{4(1-c^2)+c^2(2c^2-1)^2+3|\xi|^2}{2(3-2c^2)} + \frac{3-2c^2}{2} \\ & \pm \frac{1}{2} \left( \left( \frac{4(1-c^2)+c^2(2c^2-1)^2+3|\xi|^2}{3-2c^2} - (3-2c^2) \right)^2 + 4c^2(2c^2-1)^2 \right)^{1/2}. \end{aligned}$$

This leads to  $\sigma_{\text{ess}}(\mathcal{T}_c) = \sigma(\mathcal{T}_\infty) = [\tau_c, +\infty)$ , with  $\tau_c$  as in (5-22). This completes the proof of Step 1.

**Step 2.** *There exists a positive number  $\Lambda_2$ , depending continuously on  $c$ , such that*

$$K_c(v) \geq \Lambda_2 \int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + v_2^2), \tag{5-23}$$

for any pair  $v \in X^1(\mathbb{R})$  such that

$$\langle v, v_c^{-1} S \chi_c \rangle_{L^2(\mathbb{R})^2} = 0. \tag{5-24}$$

We start by improving the estimate in (5-19). Given a pair  $w \in X^1(\mathbb{R})$ , we observe that

$$\left| \langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2} - 3 \int_{\mathbb{R}} \frac{v_c^2}{\mu_c} (\partial_x w_1)^2 \right| \leq A_c \int_{\mathbb{R}} (w_1^2 + w_2^2).$$

Here and in the sequel,  $A_c$  refers to a positive number depending continuously on  $c$ . For  $0 < \tau < 1$ , we have

$$\langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2} \geq (1-\tau) \langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2} + 3\tau \int_{\mathbb{R}} \frac{v_c^2}{\mu_c} (\partial_x w_1)^2 - A_c \tau \int_{\mathbb{R}} (w_1^2 + w_2^2).$$

Since  $v_c^2/\mu_c \geq 1/(3-2c^2)$ , this yields

$$\langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2} \geq ((1-\tau)\Lambda_1 - A_c \tau) \int_{\mathbb{R}} (w_1^2 + w_2^2) + \frac{3\tau}{3-2c^2} \int_{\mathbb{R}} (\partial_x w_1)^2$$

under condition (5-20). For  $\tau$  small enough, this leads to

$$\langle \mathcal{T}_c(w), w \rangle_{L^2(\mathbb{R})^2} \geq A_c \int_{\mathbb{R}} ((\partial_x w_1)^2 + w_1^2 + w_2^2) \tag{5-25}$$

when  $w$  satisfies condition (5-20).

Since the pair  $w$  depends on the pair  $v$ , we can write (5-25) in terms of  $v$ . By (5-17),  $K_c(v)$  is equal to the left-hand side of (5-25). We deduce that (5-25) may be expressed as

$$K_c(v) \geq A_c \int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2) + A_c \int_{\mathbb{R}} \left( v_2 - \frac{2cv_c(\partial_x v_c)}{\mu_c(1-v_c^2)} \partial_x v_1 \right)^2.$$

We recall that, given two vectors  $a$  and  $b$  in a Hilbert space  $H$ , we have

$$\|a - b\|_H^2 \geq \tau \|a\|_H^2 - \frac{\tau}{1-\tau} \|b\|_H^2$$

for any  $0 < \tau < 1$ . Then, we deduce that

$$K_c(v) \geq A_c \int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + \tau v_2^2) - \frac{\tau A_c}{1-\tau} \int_{\mathbb{R}} \left( \frac{v_c(\partial_x v_c)}{\mu_c(1-v_c^2)} \partial_x v_1 \right)^2.$$

We choose  $\tau$  small enough so that we can infer from (2-3) that

$$K_c(v) \geq A_c \int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + v_2^2) \tag{5-26}$$

when  $w$  satisfies condition (5-20), i.e., when  $v$  is orthogonal to the pair

$$\mathbf{v}_c = \left( v_c^2 - \partial_x \left( \frac{2cv_c^2(\partial_x v_c)^2}{\mu_c(1-v_c^2)} \right), \frac{2cv_c^2(\partial_x v_c)^2}{\mu_c(1-v_c^2)} \right). \tag{5-27}$$

Next, we verify that (5-26) remains true, decreasing possibly the value of  $A_c$ , when we replace this orthogonality condition by

$$\langle v, v_c Q_c \rangle_{L^2(\mathbb{R})^2} = 0. \tag{5-28}$$

We remark that

$$\langle \mathbf{v}_c, v_c Q_c \rangle_{L^2(\mathbb{R})^2} \neq 0.$$

Indeed, we would deduce from (5-26) that

$$0 = K_c(v_c Q_c) \geq A_c \int_{\mathbb{R}} ((\partial_x v_c^2)^2 + v_c^4 + (v_c w_c)^2) > 0,$$

which is impossible. In addition, the number  $\langle \mathbf{v}_c, v_c Q_c \rangle_{L^2(\mathbb{R})^2}$  depends continuously on  $c$  in view of (5-27). Given a pair  $\tilde{v}$  satisfying (5-28), we denote by  $\lambda$  the real number such that  $\mathbf{v} = \lambda v_c Q_c + \tilde{v}$  is orthogonal to  $v_c$ . Since  $v_c Q_c$  belongs to the kernel of  $K_c$ , using (5-26) we obtain

$$K_c(\tilde{v}) = K_c(\mathbf{v}) \geq A_c \int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + v_2^2). \tag{5-29}$$

On the other hand, since  $\tilde{v}$  satisfies (5-28), we have

$$\lambda = \frac{\langle \mathbf{v}, v_c Q_c \rangle_{L^2(\mathbb{R})^2}}{\|v_c Q_c\|_{L^2(\mathbb{R})^2}^2}.$$

Using the Cauchy–Schwarz inequality, this yields

$$\lambda^2 \leq A_c \left( \int_{\mathbb{R}} (v_c^4 + (v_c w_c)^2) \right) \left( \int_{\mathbb{R}} (v_1^2 + v_2^2) \right).$$

Hence, by (2-3) and (5-29),

$$\lambda^2 \leq A_c K_c(\mathbf{v}) = A_c K_c(\tilde{\mathbf{v}}).$$

Using (5-29), this leads to

$$\int_{\mathbb{R}} ((\partial_x \tilde{v}_1)^2 + \tilde{v}_1^2 + \tilde{v}_2^2) \leq 2 \left( \lambda^2 \int_{\mathbb{R}} v_c^2 ((\partial_x v_c)^2 + v_c^2 + w_c^2) + \int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + v_2^2) \right) \leq A_c K_c(\tilde{\mathbf{v}}).$$

We finish the proof of this step applying again the same argument. We write  $v = \lambda v_c S Q_c + \tilde{v}$ , with  $\langle \tilde{v}, v_c Q_c \rangle_{L^2(\mathbb{R})^2} = 0$ . Since  $v_c Q_c$  belongs to the kernel of  $K_c$ , we infer from the same argument that

$$K_c(v) = K_c(\tilde{\mathbf{v}}) \geq \Lambda_2 \int_{\mathbb{R}} (\partial_x \tilde{v}_1)^2 + \tilde{v}_1^2 + \tilde{v}_2^2. \quad (5-30)$$

Using the orthogonality condition in (5-24), we obtain

$$\lambda = - \frac{\langle \tilde{\mathbf{v}}, v_c^{-1} S \chi_c \rangle_{L^2(\mathbb{R})^2}}{\langle Q_c, S \chi_c \rangle_{L^2(\mathbb{R})^2}}.$$

By the Cauchy–Schwarz inequality, we are led to

$$\lambda^2 \leq A_c \|v_c^{-1} S \chi_c\|_{L^2 \times L^2}^2 \int_{\mathbb{R}} (\tilde{v}_1^2 + \tilde{v}_2^2).$$

Invoking the exponential decay of  $\chi_c$  in (A-46), we deduce

$$\|v_c^{-1} S \chi_c\|_{L^2 \times L^2}^2 \leq A_c.$$

As a consequence, we can derive from (5-30) that

$$\lambda^2 \leq A_c K_c(\tilde{\mathbf{v}}) = A_c K_c(v).$$

Combining again with (5-30), we are led to

$$\int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + v_2^2) \leq 2 \left( \lambda^2 \int_{\mathbb{R}} v_c^2 ((\partial_x v_c)^2 + v_c^2 + w_c^2) + \int_{\mathbb{R}} ((\partial_x \tilde{v}_1)^2 + \tilde{v}_1^2 + \tilde{v}_2^2) \right) \leq A_c K_c(v).$$

which completes the proof of Step 2.

**Step 3.** *End of the proof.*

Since the pair  $v$  depends on the pair  $u$  as in (5-14), we can write (5-23) in terms of  $u$ . The left-hand side of (5-23) is equal to  $G_c(u)$  by (5-15). Moreover, for the right-hand side, we have

$$\int_{\mathbb{R}} ((\partial_x v_1)^2 + v_1^2 + v_2^2) = \int_{\mathbb{R}} v_c^2 ((\partial_x u_1)^2 + (2v_c^2 + c^2)u_1^2 + u_2^2).$$

We deduce that (5-23) may be written as

$$G_c(u) \geq A_c \int_{\mathbb{R}} v_c^2 ((\partial_x u_1)^2 + u_1^2 + u_2^2), \quad (5-31)$$

when  $v_c u$  verifies the orthogonality condition (5-24), which means that  $u$  verifies the orthogonality condition (2-52). We recall that

$$v_c(x) \geq A_c e^{-|x|}$$

by (2-3), which is sufficient to obtain (2-51). This completes the proof of Proposition 2.12. □

*Proof of Proposition 2.13.* First we check that we are allowed to differentiate the quantity

$$\mathcal{J}^*(t) := \langle M_{c^*(t)} u^*(\cdot, t), u^*(\cdot, t) \rangle_{L^2(\mathbb{R})^2}.$$

Indeed, by (2-41), (5-1) and (A-42), there exists a positive number  $A_{k,c}$  such that

$$\int_{\mathbb{R}} ((\partial_x^k u_1^*(x, t))^2 + (\partial_x^k u_2^*(x, t))^2) e^{v_c |x|} dx \leq A_{k,c}. \tag{5-32}$$

Next, using (2-42) and (2-45), we obtain

$$\begin{aligned} \frac{d}{dt}(\mathcal{J}^*) &= 2\langle SM_{c^*} u^*, \mathcal{H}_{c^*}(JSu^*) \rangle_{L^2(\mathbb{R})^2} + 2\langle SM_{c^*} u^*, \mathcal{H}_{c^*}(J\mathcal{R}_{c^*} \varepsilon^*) \rangle_{L^2(\mathbb{R})^2} \\ &\quad + 2((a^*)' - c^*) \langle SM_{c^*} u^*, \mathcal{H}_{c^*}(\partial_x \varepsilon^*) \rangle_{L^2(\mathbb{R})^2} - 2(c^*)' \langle SM_{c^*} u^*, \mathcal{H}_{c^*}(\partial_c Q_{c^*}) \rangle_{L^2(\mathbb{R})^2} \\ &\quad + (c^*)' \langle \partial_c M_{c^*} u^*, u^* \rangle_{L^2(\mathbb{R})^2} + 2(c^*)' \langle M_{c^*} u^*, S\partial_c \mathcal{H}_{c^*}(\varepsilon^*) \rangle_{L^2(\mathbb{R})^2}. \end{aligned} \tag{5-33}$$

The proof of (2-53) is the same as in [B ethuel et al. 2015]. We will give only the main ideas of the proof. We will estimate all the terms on the right-hand side of (5-33) except the fourth term, which vanishes.

For the first one, we infer from Proposition 2.12 the following estimate.

**Step 1.** *There exists a positive number  $B_1$ , depending only on  $c$ , such that*

$$\mathcal{J}_1^*(t) := 2\langle SM_{c^*} u^*, \mathcal{H}_{c^*}(JSu^*) \rangle_{L^2(\mathbb{R})^2} \geq B_1 \int_{\mathbb{R}} [(\partial_x u_1^*)^2 + (u_1^*)^2 + (u_2^*)^2](x, t) e^{-2|x|} dx$$

for any  $t \in \mathbb{R}$ .

From (2-21), (2-57) and (5-1), we get an estimate for the second term.

**Step 2.** *There exists a positive number  $B_2$ , depending only on  $c$ , such that*

$$|\mathcal{J}_2^*(t)| := 2|\langle SM_{c^*} u^*, \mathcal{H}_{c^*}(J\mathcal{R}_{c^*} \varepsilon^*) \rangle_{L^2(\mathbb{R})^2}| \leq B_2 \|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2$$

for any  $t \in \mathbb{R}$ .

For the third one, we use (2-21) to obtain:

**Step 3.** *There exists a positive number  $B_3$ , depending only on  $c$ , such that*

$$|\mathcal{J}_3^*(t)| := 2|(a^*)' - c^*| |\langle SM_{c^*} u^*, \mathcal{H}_{c^*}(\partial_x \varepsilon^*) \rangle_{L^2(\mathbb{R})^2}| \leq B_3 \|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2$$

for any  $t \in \mathbb{R}$ .

We now prove the following statement for the fourth term.

**Step 4.** We have

$$\mathcal{J}_4^*(t) := 2(c^*)' \langle SM_{c^*} u^*, \mathcal{H}_{c^*}(\partial_c Q_{c^*}) \rangle_{L^2(\mathbb{R})^2} = 0$$

for any  $t \in \mathbb{R}$ .

Since  $\mathcal{H}_{c^*}(\partial_c Q_{c^*}) = P'(Q_{c^*}) = S Q_{c^*}$  and  $M_{c^*} Q_{c^*} = S \partial_x Q_{c^*}$ , we have

$$\begin{aligned} \langle SM_{c^*} u^*, \mathcal{H}_{c^*}(\partial_c Q_{c^*}) \rangle_{L^2(\mathbb{R})^2} &= \langle M_{c^*} u^*, Q_{c^*} \rangle_{L^2(\mathbb{R})^2} = \langle u^*, S \partial_x Q_{c^*} \rangle_{L^2(\mathbb{R})^2} \\ &= \langle \varepsilon^*, \mathcal{H}_{c^*}(\partial_x Q_{c^*}) \rangle_{L^2(\mathbb{R})^2} = 0. \end{aligned}$$

This is the reason why we do not need to establish a quadratic dependence of  $(c^*)'(t)$  on  $\varepsilon^*$ .

Next, we use (2-3), (2-9), (2-21) and (2-45) to bound the fifth term.

**Step 5.** There exists a positive number  $B_5$ , depending only on  $c$ , such that

$$|\mathcal{J}_5^*(t)| := |(c^*)'| \langle \partial_c M_{c^*} u^*, u^* \rangle_{L^2(\mathbb{R})^2} \leq B_5 \|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2$$

for any  $t \in \mathbb{R}$ .

Finally, we acquire a bound on the sixth term in the same way.

**Step 6.** There exists a positive number  $B_6$ , depending only on  $c$ , such that

$$|\mathcal{J}_6^*(t)| := |(c^*)'| \langle M_{c^*} u^*, S \partial_c \mathcal{H}_{c^*}(\varepsilon^*) \rangle_{L^2(\mathbb{R})^2} \leq B_6 \|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2$$

for any  $t \in \mathbb{R}$ .

We conclude the proof of Proposition 2.13 by combining the six previous steps to obtain (2-53), with  $B_* := \max\{1/B_1, B_2 + B_3 + B_5 + B_6\}$ .  $\square$

*Proof of Corollary 2.14.* Corollary 2.14 is a consequence of Propositions 2.10 and 2.13. We combine the two estimates (2-43) and (2-53) with the definition of  $N(t)$  to obtain

$$\frac{d}{dt} \langle N(t) u^*(\cdot, t), u^*(\cdot, t) \rangle_{L^2(\mathbb{R})^2} \geq \left( \frac{1-c^2}{16} - A_* B_*^2 e^{2R_*} \|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \right) \|u^*(\cdot, t)\|_{X(\mathbb{R})}^2$$

for any  $t \in \mathbb{R}$ . In view of (2-21), we fix the parameter  $\beta_c$  such that

$$\|\varepsilon^*(\cdot, t)\|_{X(\mathbb{R})}^{1/2} \leq \frac{1-c^2}{32A_* B_*^2 e^{2R_*}}$$

for any  $t \in \mathbb{R}$ , to obtain (2-54). In view of (2-3), (2-21) and (2-45), we notice that there exists a positive number  $A_c$ , depending only on  $c$ , such that

$$\|M_{c^*}(t)\|_{L^\infty(\mathbb{R})} \leq A_c \tag{5-34}$$

for any  $t \in \mathbb{R}$ . Moreover, since the map  $t \mapsto \langle N(t) u^*(\cdot, t), u^*(\cdot, t) \rangle_{L^2(\mathbb{R})^2}$  is uniformly bounded by (5-32) and (5-34), the estimate (2-55) follows by integrating (2-54) from  $t = -\infty$  to  $t = +\infty$ . Finally, statement (2-56) is a direct consequence of (2-55).  $\square$

### Appendix A. Appendix

**Weak continuity of the hydrodynamical flow.** In this section, we prove the weak continuity of the hydrodynamical flow, which is stated in the following proposition.

**Proposition A.1.** *We consider a sequence  $(v_{n,0}, w_{n,0})_{n \in \mathbb{N}} \in \mathcal{NV}(\mathbb{R})^{\mathbb{N}}$ , and a pair  $(v_0, w_0) \in \mathcal{NV}(\mathbb{R})$  such that*

$$v_{n,0} \rightharpoonup v_0 \quad \text{in } H^1(\mathbb{R}) \quad \text{and} \quad w_{n,0} \rightharpoonup w_0 \quad \text{in } L^2(\mathbb{R}) \tag{A-1}$$

as  $n \rightarrow +\infty$ . We denote by  $(v_n, w_n)$  the unique solution to (HLL) with initial datum  $(v_{n,0}, w_{n,0})$  and we assume that there exists a positive number  $T_n$  such that the solutions  $(v_n, w_n)$  are defined on  $(-T_n, T_n)$ , and satisfy the condition

$$\sup_{n \in \mathbb{N}} \sup_{t \in (-T_n, T_n)} \max_{x \in \mathbb{R}} v_n(x, t) \leq 1 - \sigma \tag{A-2}$$

for a given positive number  $\sigma$ . Then, the unique solution  $(v, w)$  to (HLL) with initial datum  $(v_0, w_0)$  is defined on  $(-T_{\max}, T_{\max})$ , with<sup>3</sup>

$$T_{\max} = \liminf_{n \rightarrow +\infty} T_n,$$

and for any  $t \in (-T_{\max}, T_{\max})$ , we have

$$v_n(t) \rightharpoonup v(t) \quad \text{in } H^1(\mathbb{R}) \quad \text{and} \quad w_n(t) \rightharpoonup w(t) \quad \text{in } L^2(\mathbb{R}) \tag{A-3}$$

as  $n \rightarrow +\infty$ .

First we prove a weak continuity property of the flow of equations (2-32)–(2-34). Next, we deduce the weak convergence of  $w_n$  from (4-41).

More precisely, we consider now a sequence of initial conditions  $(\Psi_{n,0}, v_{n,0}) \in L^2(\mathbb{R}) \times H^1(\mathbb{R})$ , such that the norms  $\|\Psi_{n,0}\|_{L^2}$  and  $\|v_{n,0}\|_{L^2}$  are uniformly bounded with respect to  $n$ , and we assume that

$$\sup_{n \in \mathbb{N}} \|v_{n,0}\|_{L^\infty(\mathbb{R})} < 1. \tag{A-4}$$

Then, there exist two functions  $\Psi_0 \in L^2(\mathbb{R})$  and  $v_0 \in H^1(\mathbb{R})$  such that, going possibly to a subsequence,

$$\Psi_{n,0} \rightharpoonup \Psi_0 \quad \text{in } L^2(\mathbb{R}), \tag{A-5}$$

$$v_{n,0} \rightharpoonup v_0 \quad \text{in } H^1(\mathbb{R}), \tag{A-6}$$

and, for any compact subset  $K$  of  $\mathbb{R}$ ,

$$v_{n,0} \rightarrow v_0 \quad \text{in } L^\infty(K) \tag{A-7}$$

as  $n \rightarrow +\infty$ . We claim that this convergence is conserved along the flow corresponding to equations (2-32)–(2-34).<sup>4</sup>

<sup>3</sup>See Theorem 1 in [de Laire and Gravejat 2015] for more details.

<sup>4</sup>We only consider here positive time but the proof remains valid for negative time.

**Proposition A.2.** *We consider two sequences  $(\Psi_{n,0})_{n \in \mathbb{N}} \in L^2(\mathbb{R})^{\mathbb{N}}$  and  $(v_{n,0})_{n \in \mathbb{N}} \in H^1(\mathbb{R})^{\mathbb{N}}$ , and two functions  $\Psi_0 \in L^2(\mathbb{R})$  and  $v_0 \in H^1(\mathbb{R})$ , such that assumptions (A-4)–(A-7) are satisfied, and we denote by  $(\Psi_n, v_n)$  and  $(\Psi, v)$ , respectively, the unique global solutions to (2-32)–(2-34) with initial data  $(\Psi_{n,0}, v_{n,0})$  and  $(\Psi_0, v_0)$ , which we assume to be defined on  $[0, T]$  for a positive number  $T$ . For any fixed  $t \in [0, T]$ , we have*

$$\Psi_n(\cdot, t) \rightharpoonup \Psi(\cdot, t) \quad \text{in } L^2(\mathbb{R}), \quad (\text{A-8})$$

$$v_n(\cdot, t) \rightharpoonup v(\cdot, t) \quad \text{in } H^1(\mathbb{R}) \quad (\text{A-9})$$

when  $n \rightarrow +\infty$ .

*Proof.* We split the proof into four steps.

**Step 1.** *There exist three functions  $\Phi \in L^2([0, T], L^2(\mathbb{R}))$  and  $\mathfrak{v} \in L^2([0, T], H^1(\mathbb{R}))$  such that, up to a further subsequence,*

$$\Psi_n(t) \rightharpoonup \Phi(t) \quad \text{in } L^2(\mathbb{R}), \quad (\text{A-10})$$

$$v_n(\cdot, t) \rightharpoonup \mathfrak{v}(\cdot, t) \quad \text{in } H^1(\mathbb{R}), \quad (\text{A-11})$$

$$v_n(\cdot, t) \rightarrow \mathfrak{v}(\cdot, t) \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}) \quad (\text{A-12})$$

for all  $t \in [0, T]$ , and

$$|\Psi_n|^2 \Psi_n \rightharpoonup |\Phi|^2 \Phi \quad \text{in } L^2([0, T], L^2(\mathbb{R})), \quad (\text{A-13})$$

when  $n \rightarrow +\infty$ .

*Proof.* We recall that there exists a constant  $M$  such that

$$\|\Psi_{n,0}\|_{L^2} \leq M \quad \text{and} \quad \|v_{n,0}\|_{H^1} \leq M$$

uniformly on  $n$ . Applying [Proposition 2.7](#) to the pairs  $(\Psi_n, v_n)$  and  $(0, 0)$ , we obtain

$$\|\Psi_n\|_{C_T^0 L_x^2} + \|v_n\|_{C_T^0 H_x^1} + \|\Psi_n\|_{L_T^4 L_x^\infty} \leq A(\|\Psi_{n,0}\|_{L^2} + \|v_{n,0}\|_{H^1}).$$

This leads to

$$\|\Psi_n\|_{L_T^4 L_x^\infty} \leq 2AM, \quad \|\Psi_n\|_{L_T^\infty L_x^2} \leq 2AM \quad \text{and} \quad \|v_n\|_{L_T^\infty H_x^1} \leq 2AM. \quad (\text{A-14})$$

Hence, there exist two functions  $\Phi \in L^\infty([0, T], L^2(\mathbb{R})) \cap L^4([0, T], L^\infty(\mathbb{R}))$  and  $\mathfrak{v} \in L^\infty([0, T], H^1(\mathbb{R}))$  such that

$$\Psi_n \xrightarrow{*} \Phi \quad \text{in } L^\infty([0, T], L^2(\mathbb{R})),$$

$$v_n \xrightarrow{*} \mathfrak{v} \quad \text{in } L^\infty([0, T], H^1(\mathbb{R})).$$

Let us prove (A-10) and (A-11). We argue as in [\[Béthuel et al. 2015\]](#) and we introduce a cutoff function  $\chi \in C_c^\infty(\mathbb{R})$  such that  $\chi \equiv 1$  on  $[-1, 1]$  and  $\chi \equiv 0$  on  $(-\infty, 2] \cup [2, +\infty)$ . Set  $\chi_p(\cdot) := \chi(\cdot/p)$  for any integer  $p \in \mathbb{N}^*$ . By (A-14), the sequences  $(\chi_p \Psi_n)_{n \in \mathbb{N}}$  and  $(\chi_p v_n)_{n \in \mathbb{N}}$  are bounded in  $C^0([0, T], L^2(\mathbb{R}))$  and

$C^0([0, T], H^1(\mathbb{R}))$ , respectively. In view of the Rellich–Kondrachov theorem, the sets  $\{\chi_p \Psi_n(\cdot, t) \mid n \in \mathbb{N}\}$  and  $\{\chi_p v_n(\cdot, t) \mid n \in \mathbb{N}\}$  are relatively compact in  $H^{-2}(\mathbb{R})$  and  $H^{-1}(\mathbb{R})$ , respectively, for any fixed  $t \in [0, T]$ . In addition, since the pair  $(\Psi_n, v_n)$  is a solution to (2-32)–(2-34), we have that  $(\partial_t \Psi_n, \partial_t v_n)$  belongs to  $C^0([0, T], H^{-2}(\mathbb{R}) \times H^{-1}(\mathbb{R}))$  and satisfies

$$\|\partial_t \Psi_n(\cdot, t)\|_{H^{-2}(\mathbb{R})} \leq K_M \quad \text{and} \quad \|\partial_t v_n(\cdot, t)\|_{H^{-1}(\mathbb{R})} \leq K_M.$$

This leads to the fact that the pair  $(\chi_p \Psi_n, \chi_p v_n)$  is equicontinuous in  $C^0([0, T], H^{-2}(\mathbb{R}) \times H^{-1}(\mathbb{R}))$ . Then, we apply the Arzelà–Ascoli theorem and the Cantor diagonal argument to find a further subsequence (independent of  $p$ ), such that, for each  $p \in \mathbb{N}^*$ ,

$$\chi_p \Psi_n \rightarrow \chi_p \Phi \quad \text{in } C^0([0, T], H^{-2}(\mathbb{R})), \tag{A-15}$$

$$\chi_p v_n \rightarrow \chi_p \mathfrak{v} \quad \text{in } C^0([0, T], H^{-1}(\mathbb{R})) \tag{A-16}$$

as  $n \rightarrow +\infty$ . Combining this with (A-14) we infer that (A-10) and (A-11) hold. By the Sobolev embedding theorem, (A-12) is a consequence of (A-11).

Now, let us prove (A-13). Using the Hölder inequality, we infer that

$$\int_0^T \int_{\mathbb{R}} |\Psi_n(x, t)|^6 dx dt \leq \|\Psi_n\|_{L^\infty L_x^2}^2 \|\Psi_n\|_{L_T^4 L_x^\infty}^4.$$

By (A-14), we conclude that

$$\| |\Psi_n|^2 \Psi_n \|_{L_T^2 L_x^2} \leq M. \tag{A-17}$$

So, there exists a function  $\Phi_1 \in L^2(\mathbb{R} \times [0, T])$  such that up to a further subsequence,

$$|\Psi_n|^2 \Psi_n \rightharpoonup \Phi_1 \quad \text{in } L^2(\mathbb{R} \times [0, T]).$$

Let us prove that  $\Phi_1 \equiv |\Phi|^2 \Phi$ . To obtain this it is sufficient to prove that, up to a subsequence,

$$\Psi_n \rightarrow \Phi \quad \text{in } L^2([0, T], L^2([-R, R])) \tag{A-18}$$

for any  $R > 0$ , i.e., the sequence  $(\Psi_n)$  is relatively compact in  $L^2([-R, R] \times [0, T])$ . Indeed, using the Hölder inequality, we obtain

$$\begin{aligned} \| |\Psi_n|^2 \Psi_n - |\Phi|^2 \Phi \|_{L_{T,R}^{6/5}} &= \| (\Psi_n - \Phi)(|\Psi_n|^2 + |\Phi|^2) + \Psi_n \Phi(\bar{\Psi}_n - \bar{\Phi}) \|_{L_{T,R}^{6/5}} \\ &\leq 2 \| |\Psi_n - \Phi| (|\Psi_n|^2 + |\Phi|^2) \|_{L_{T,R}^{6/5}} \\ &\leq 2 \| \Psi_n - \Phi \|_{L_{T,R}^2} (\| \Psi_n \|_{L_{T,R}^6}^2 + \| \Phi \|_{L_{T,R}^6}^2) \end{aligned} \tag{A-19}$$

for any  $R > 0$ . By (A-17),  $(\Psi_n)$  is uniformly bounded in  $L^6(\mathbb{R} \times [0, T])$  and  $\Phi \in L^6(\mathbb{R} \times [0, T])$ . Then

$$|\Psi_n|^2 \Psi_n \rightarrow |\Phi|^2 \Phi \quad \text{in } L^{6/5}([-R, R] \times [0, T]),$$

so that  $\Phi_1 \equiv |\Phi|^2 \Phi$ . Now, let us prove that the sequence  $(\Psi_n)$  is relatively compact in  $L^2([-R, R] \times [0, T])$ . The main point of the proof is the following claim.

**Claim 1.** Let  $\Psi$  be a solution of (2-32) in

$$C^0([0, T], L^2(\mathbb{R})) \cap L^4([0, T], L^\infty(\mathbb{R})).$$

Then  $\Psi \in L^2([0, T], H_{\text{loc}}^{1/2}(\mathbb{R}))$ .

*Proof.* The proof relies on the Kato smoothing effect for the linear Schrödinger group (see [Linares and Ponce 2009]). Let  $S(t) = e^{it\partial_{xx}}$ , and

$$\mathcal{F}(\Psi, v) := \frac{1}{2}v^2\Psi - \text{Re}(\Psi(1 - 2F(v, \bar{\Psi}))(1 - 2F(v, \Psi))). \tag{A-20}$$

We recall that there exists a positive constant  $M$  such that

$$\sup_{x \in \mathbb{R}} \int_{-\infty}^{+\infty} |D_x^{1/2} S(t) f(x)|^2 dt \leq M \|f\|_{L^2}^2 \tag{A-21}$$

and

$$\left\| \int_{\mathbb{R}} S(-t') D_x^{1/2} h(\cdot, t') dt' \right\|_{L^2} \leq M \|h\|_{L_x^1 L_t^2} \tag{A-22}$$

when  $f \in L^2(\mathbb{R})$  and  $h \in L^1(\mathbb{R}, L^2(\mathbb{R}))$  (see [Linares and Ponce 2009] for more details). We prove that there exists a positive constant  $M$  such that

$$\|D_x^{1/2} \Psi\|_{L_x^\infty L_T^2} \leq M \|\Psi_0\|_{L^2} + M \|\Psi\|_{L_{T,x}^2} (\|\Psi\|_{L_{T,x}^6}^2 + T^{1/2} (\|v\|_{L_{T,x}^\infty}^2 + \|1 - 2F(v, \Psi)\|_{L_{T,x}^\infty}^2)). \tag{A-23}$$

The claim is a consequence of this estimate, so that it is sufficient to prove (A-23).

We write

$$\Psi(x, t) = S(t)\Psi_0(x) + i \int_0^t S(t-t') (2(|\Psi|^2\Psi)(x, t') + \mathcal{F}(\Psi, v)(x, t')) dt'$$

for all  $(x, t) \in \mathbb{R}$ . First, using (A-21), we obtain

$$\sup_{x \in \mathbb{R}} \int_{-\infty}^{+\infty} |D_x^{1/2} S(t)\Psi_0(x)|^2 dt \leq M \|\Psi_0\|_{L^2}^2.$$

For the nonlinear term, we can argue as in [Goubet and Molinet 2009] to prove that

$$\left\| \int_0^t S(t-t') D_x^{1/2} g(\cdot, t') dt' \right\|_{L_x^\infty L_T^2} \leq M \|g\|_{L_T^1 L_x^2}. \tag{A-24}$$

Using a duality argument, it is equivalent to prove that for any smooth function  $h$  that satisfies  $\|h\|_{L_x^1 L_t^2} \leq 1$ , we have

$$\left| \int_{\mathbb{R} \times [0, T]^2} S(t-t') D_x^{1/2} g(x, t') \bar{h}(x, t) dt' dx dt \right| \leq M \|g\|_{L_T^1 L_x^2}. \tag{A-25}$$

Using the Cauchy–Schwarz and Strichartz estimates and (A-22), the left-hand side can be written as

$$\begin{aligned} & \left| \int_{\mathbb{R}} \left( \int_0^T S(-t') D_x^{1/2} g(x, t') dt' \right) \left( \int_0^T \overline{S(-t) h(x, t)} dt \right) dx \right| \\ &= \left| \int_{\mathbb{R}} \left( \int_0^T S(-t') g(x, t') dt' \right) \left( \int_0^T \overline{S(-t) D_x^{1/2} h(x, t)} dt \right) dx \right| \\ &\leq M \left\| \int_0^T S(-t') g(x, t') dt' \right\|_{L^2} \leq M \|g\|_{L_T^1 L_x^2}. \end{aligned}$$

This achieves the proof of (A-24). Similarly, we have

$$\left\| \int_0^t S(t-t') D_x^{1/2} g(\cdot, t') dt' \right\|_{L_x^\infty L_T^2} \leq M \|g\|_{L_{T,x}^{5/6}}. \tag{A-26}$$

We next apply (A-24) and (A-26) on the nonlinear terms to obtain, using the Cauchy–Schwarz and Hölder estimates,

$$\left\| \int_0^t D_x^{1/2} S(t-t') (|\Psi|^2 \Psi)(\cdot, t') dt' \right\|_{L_x^\infty L_T^2} \leq M \|\Psi^3\|_{L_{T,x}^{6/5}} \leq M \|\Psi\|_{L_{T,x}^2} \|\Psi\|_{L_{T,x}^6}^2$$

and

$$\begin{aligned} \left\| \int_0^t D_x^{1/2} S(t-t') \mathcal{F}(\Psi, v)(\cdot, t') dt' \right\|_{L_x^\infty L_T^2} &\leq M \|\mathcal{F}(\Psi, v)\|_{L_T^1 L_x^2} \\ &\leq M \|\Psi\|_{L_T^1 L_x^2} (\|v\|_{L_{T,x}^\infty}^2 + \|1 - 2F(v, \Psi)\|_{L_{T,x}^\infty}^2) \\ &\leq MT^{1/2} \|\Psi\|_{L_{T,x}^2} (\|v\|_{L_{T,x}^\infty}^2 + \|1 - 2F(v, \Psi)\|_{L_{T,x}^\infty}^2). \end{aligned}$$

Since  $v \in L^\infty([0, T], H^1(\mathbb{R}))$  and  $\Psi \in L^\infty([0, T], L^2(\mathbb{R}))$ , we know that  $\Psi \in L^\infty([0, T], L^2(\mathbb{R}))$  and  $F(\Psi, v) \in L^\infty(\mathbb{R} \times [0, T])$ . Using the fact that  $\Psi \in L^6(\mathbb{R} \times [0, T])$ , we finish the proof of this claim.  $\square$

Applying this claim to the sequence  $(\Psi_n)$  yields that  $(\Psi_n)$  is uniformly bounded in  $L^2([0, T], H_{\text{loc}}^{1/2}(\mathbb{R}))$ . On the other hand, we have  $\mathcal{F}(\Psi_n, v_n) \in L^\infty([0, T], L^2(\mathbb{R}))$ , since

$$v_n \in L^\infty([0, T], H^1(\mathbb{R})), \quad \Psi_n \in L^\infty([0, T], L^2(\mathbb{R})) \quad \text{and} \quad F(\Psi_n, v_n) \in L^\infty(\mathbb{R} \times [0, T]).$$

Then, using (2-32) and (A-17), we obtain that  $(\Psi_n)$  is uniformly bounded in  $H^1([0, T], H^{-2}(\mathbb{R}))$ . Hence, by interpolation,  $(\Psi_n) \in H^{1/10}([0, T], H_{\text{loc}}^{1/4}(\mathbb{R}))$ , so that it converges in  $L^2([-R, R] \times [0, T])$  for any  $R > 0$ . This finishes the proofs of (A-18) and of Step 1.  $\square$

**Step 2.** *We have*

$$\mathcal{F}(\Psi_n, v_n) \rightharpoonup \mathcal{F}(\Phi, \mathbf{v}) \quad \text{in } L^2(\mathbb{R}), \tag{A-27}$$

for any  $t \in [0, T]$ , and

$$\mathcal{F}(\Psi_n, v_n) \rightarrow \mathcal{F}(\Phi, \mathbf{v}) \quad \text{in } L^1([0, T], L_{\text{loc}}^2(\mathbb{R})). \tag{A-28}$$

*Proof.* Let  $\phi \in L^2(\mathbb{R})$ . We compute

$$\begin{aligned} & \int_{\mathbb{R}} (v_n^2(x, t)\Psi_n(x, t) - v^2(x, t)\Phi(x, t))\phi(x) dx \\ &= \int_{\mathbb{R}} (v_n^2(x, t) - v^2(x, t))\Psi_n(x, t)\phi(x) dx + \int_{\mathbb{R}} (\Psi_n(x, t) - \Phi(x, t))v^2(x, t)\phi(x) dx. \end{aligned} \tag{A-29}$$

The second term on the right-hand side goes to 0 when  $n$  goes to  $+\infty$ , since  $v^2(t)\phi \in L^2(\mathbb{R})$  for all  $t$  on the one hand and using (A-10) on the other hand. For the first term on the right-hand side, we consider a cutoff function  $\chi$  with support in  $[-1, 1]$  and let  $\chi_R(x) = \chi(x/R)$  for all  $(x, R) \in \mathbb{R} \times (0, +\infty)$ . We set

$$\begin{aligned} I_n(t) &:= \int_{\mathbb{R}} (v_n^2(x, t) - v^2(x, t))\Psi_n(x, t)\phi(x) dx, \\ I_n^{(1)}(t) &:= \int_{\mathbb{R}} (v_n^2(x, t) - v^2(x, t))\Psi_n(x, t)\chi_R(x)\phi(x) dx, \\ I_n^{(2)}(t) &:= \int_{\mathbb{R}} (v_n^2(x, t) - v^2(x, t))\Psi_n(x, t)(1 - \chi_R(x))\phi(x) dx, \end{aligned}$$

so that  $I_n(t) = I_n^{(1)}(t) + I_n^{(2)}(t)$ . By the Cauchy-Schwarz inequality, we have

$$|I_n^{(1)}(t)| \leq \|\Psi_n(t)\|_{L^2(\mathbb{R})} \|\phi\|_{L^2(\mathbb{R})} \|v_n^2(t) - v^2(t)\|_{L^\infty([-R, R])}. \tag{A-30}$$

Using (A-12) and (A-14), we infer that

$$I_n^{(1)}(t) \rightarrow 0 \quad \text{for any } t \in [0, T], \tag{A-31}$$

as  $n \rightarrow +\infty$ . Next, we write

$$|I_n^{(2)}(t)| \leq (\|v_n(t)\|_{L^\infty(\mathbb{R})}^2 + \|v(t)\|_{L^\infty(\mathbb{R})}^2) \|\Psi_n(t)\|_{L^2(\mathbb{R})} \|(1 - \chi_R)\phi\|_{L^2(\mathbb{R})}.$$

Since  $\phi \in L^2(\mathbb{R})$ , we have

$$\lim_{R \rightarrow \infty} \|(1 - \chi_R)\phi\|_{L^2(\mathbb{R})} = 0.$$

In view of (A-14), this is sufficient to prove that

$$I_n(t) \rightarrow 0 \tag{A-32}$$

as  $n \rightarrow +\infty$ , for all  $t \in [0, T]$ . This yields

$$(v_n^2\Psi_n)(t) \rightharpoonup (v^2\Phi)(t) \quad \text{in } L^2(\mathbb{R}) \tag{A-33}$$

for any  $t \in [0, T]$ . Now, we prove

$$v_n^2\Psi_n \rightarrow v^2\Phi \quad \text{in } L^1([0, T], L^2_{\text{loc}}(\mathbb{R})). \tag{A-34}$$

As in (A-29), we write

$$\|v_n^2\Psi_n - v^2\Phi\|_{L^1_T L^2_R} \leq \|(v_n^2 - v^2)\Psi_n\|_{L^1_T L^2_R} + \|(\Psi_n - \Phi)v^2\|_{L^1_T L^2_R}.$$

For the first term on the right-hand side, we infer from the Cauchy–Schwarz inequality that

$$\begin{aligned} \|(v_n^2 - \mathfrak{v}^2)\Psi_n\|_{L_T^1 L_R^2} &\leq \|v_n^2 - \mathfrak{v}^2\|_{L_T^2 L_R^2} \|\Psi_n\|_{L_T^2 L_R^\infty} \\ &\leq \|v_n - \mathfrak{v}\|_{L_T^4 L_R^4} (\|v_n\|_{L_T^4 L_R^4} + \|\mathfrak{v}\|_{L_T^4 L_R^4}) T^{1/2} \|\Psi_n\|_{L_T^4 L_R^\infty}. \end{aligned}$$

On the other hand, by (A-14),  $v_n$  is uniformly bounded on  $L^2([0, T], H^1(\mathbb{R}))$ . By the first equation of (2-34) and (A-14),  $v_n$  is uniformly bounded in  $H^1([0, T], H^{-1}(\mathbb{R}))$ . We deduce that  $v_n$  is uniformly bounded in  $H^{1/3}([0, T], H^{1/3}(\mathbb{R}))$  and so that  $v_n$  converges to  $\mathfrak{v}$  in  $L^4([0, T], L^4([-R, R]))$  as  $n \rightarrow +\infty$ . Hence, using (A-14) once again, we obtain

$$\|(v_n^2 - \mathfrak{v}^2)\Psi_n\|_{L_T^1 L_R^2} \rightarrow 0$$

as  $n \rightarrow +\infty$ . For the second term we have, by the Cauchy–Schwarz inequality and the Sobolev embedding theorem,

$$\|(\Psi_n - \Phi)\mathfrak{v}^2\|_{L_T^1 L_R^2} \leq \|\Psi_n - \Phi\|_{L_T^2 L_R^2} \|\mathfrak{v}^2\|_{L_T^2 L_R^\infty} \leq M^2 T^{1/2} \|\Psi_n - \Phi\|_{L_T^2 L_R^2}.$$

This yields, using (A-18),

$$\|(\Psi_n - \Phi)\mathfrak{v}^2\|_{L_T^1 L_R^2} \rightarrow 0$$

as  $n \rightarrow +\infty$ , which proves (A-34). Next, we set

$$\mathcal{G}(v_n, \Psi_n) = \Psi_n(1 - F(v_n, \bar{\Psi}_n))(1 - F(v_n, \Psi_n)).$$

We have, by (2-33),

$$\partial_x F(v_n, \Psi_n) = v_n \Psi_n \quad \text{and} \quad \partial_x F(\mathfrak{v}, \Phi) = \mathfrak{v} \Phi.$$

Using the same arguments as in the proof of (A-32), we obtain

$$\partial_x F(v_n, \Psi_n) \rightharpoonup \partial_x F(\mathfrak{v}, \Phi) \quad \text{in } L^2(\mathbb{R})$$

for any  $t \in [0, T]$ . Hence,

$$F(v_n, \Psi_n) \rightarrow F(\mathfrak{v}, \Phi) \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}) \tag{A-35}$$

for any  $t \in [0, T]$ . Using (A-10), (A-35) and the same arguments as in the proof of (A-33), we conclude that

$$\mathcal{G}(v_n, \Psi_n) \rightharpoonup \mathcal{G}(\mathfrak{v}, \Phi) \quad \text{in } L^2(\mathbb{R}) \tag{A-36}$$

for any  $t \in [0, T]$ . Next, we use (A-18) and (A-35) to prove that

$$\mathcal{G}(v_n, \Psi_n) \rightarrow \mathcal{G}(\mathfrak{v}, \Phi) \quad \text{in } L^1([0, T], L_{\text{loc}}^2(\mathbb{R})). \tag{A-37}$$

This finishes the proof of this step. □

**Step 3.**  $(\Phi, \mathfrak{v})$  is a weak solution of (2-32)–(2-34).

*Proof.* By (A-18), we have

$$i \partial_t \Psi_n \rightarrow i \partial_t \Phi \quad \text{in } \mathcal{D}'(\mathbb{R} \times [0, T]) \quad \text{and} \quad \partial_{xx}^2 \Psi_n \rightarrow \partial_{xx}^2 \Phi \quad \text{in } \mathcal{D}'(\mathbb{R} \times [0, T])$$

as  $n \rightarrow +\infty$ . It remains to invoke (A-13) and (A-35) and to take the limit  $n \rightarrow +\infty$  in the expression

$$\int_0^T \int_{\mathbb{R}} (i\partial_t \Psi_n + \partial_{xx}^2 \Psi_n + 2|\Psi_n|^2 \Psi_n + \frac{1}{2}v_n^2 \Psi_n - \operatorname{Re}(\Psi(1 - 2F(v_n, \overline{\Psi_n}))(1 - 2F(v_n, \Psi_n)))\bar{h} = 0,$$

where  $h \in C_c^\infty(\mathbb{R} \times [0, T])$ , in order to establish that  $(\Phi, v)$  is a solution to (2-32) in the sense of distributions. In addition, using the same arguments as above and (A-35), we prove that  $(\Phi, v)$  is a solution to (2-34) in the sense of distributions. Moreover, we infer from (A-5) that  $\Phi(\cdot, 0) = \Psi_0$  and from (A-6) that  $v(\cdot, 0) = v_0$ .  $\square$

In order to prove that the function  $(\Phi, v)$  coincides with the solution  $(\Psi, v)$  in Proposition A.2, it is sufficient, in view of the uniqueness result given by Proposition 2.7, to establish the following.

**Step 4.**  $\Phi \in \mathcal{C}([0, T], L^2(\mathbb{R}))$  and  $v \in \mathcal{C}([0, T], H^1(\mathbb{R}))$ .

*Proof.* First, we prove that  $\Phi \in \mathcal{C}([0, T], L^2(\mathbb{R}))$ . This is a direct consequence of the identity

$$\Phi(x, t) = S(t)\Phi_0 + \int_0^t S(t-t')(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt'. \tag{A-38}$$

Indeed, let us define

$$G(\Phi, v)(t) = \int_0^t S(t-t')(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt'.$$

Since  $S(t)\Phi_0 \in \mathcal{C}([0, T], L^2(\mathbb{R}))$ , it suffices to show  $G(\Phi, v) \in \mathcal{C}([0, T], L^2(\mathbb{R}))$ . We take  $(t_1, t_2) \in [0, T]^2$  and write

$$\begin{aligned} G(\Phi, v)(t_1) - G(\Phi, v)(t_2) &= \int_0^{t_1} (S(t_1-t') - S(t_2-t'))(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt' \\ &\quad - \int_{t_1}^{t_2} S(t-t')(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt'. \end{aligned}$$

For the second term on the right-hand side, we use the Strichartz and Cauchy–Schwarz inequalities to obtain

$$\begin{aligned} &\left\| \int_{t_1}^{t_2} S(t-t')(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt' \right\|_{L^2} \\ &\leq M \left\| 2|\Phi|^2\Phi + \mathcal{F}(\Phi, v) \right\|_{L^1([t_1, t_2], L^2(\mathbb{R}))} \\ &\leq M|t_1 - t_2|^{1/2} \|\Phi\|_{L_{T,x}^2}^2 + M|t_1 - t_2| \|\mathcal{F}(\Phi, v)\|_{L_T^\infty L_x^2}. \tag{A-39} \end{aligned}$$

For the first term, we write

$$S(t_1 - t') - S(t_2 - t') = S(t_1 - t')(1 - S(t_2 - t_1)).$$

Hence,

$$\begin{aligned} &\left\| \int_0^{t_1} (S(t_1-t') - S(t_2-t'))(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt' \right\|_{L^2} \\ &= \|(1 - S(t_2 - t_1))G(\Phi, v)(t_1)\|_{L^2}. \tag{A-40} \end{aligned}$$

Taking the limit  $t_2 \rightarrow t_1$  in (A-39) and (A-40), we obtain that  $\Phi \in C([0, T], L^2(\mathbb{R}))$ .

Now, let us prove (A-38). Denote by  $\tilde{\Phi}$  the function given by the right-hand side of (A-38). We will prove that

$$\Psi_n(t) \rightharpoonup \tilde{\Phi}(t) \quad \text{in } L^2(\mathbb{R}) \tag{A-41}$$

for all  $t \in \mathbb{R}$ . This yields  $\Phi \equiv \tilde{\Phi}$  by uniqueness of the weak limit. Let  $R > 0$  and denote by  $\chi_R$  the function defined in Step 2. Set

$$\begin{aligned} G_n^{(1)}(\cdot, t) &= \int_0^t S(t-t')\chi_R(2(|\Psi_n|^2\Psi_n)(\cdot, t') + \mathcal{F}(\Psi_n, v_n)(\cdot, t')) dt', \\ G_n^{(2)}(\cdot, t) &= \int_0^t S(t-t')(1-\chi_R)(2(|\Psi_n|^2\Psi_n)(\cdot, t') + \mathcal{F}(\Psi_n, v_n)(\cdot, t')) dt', \\ G^{(1)}(\cdot, t) &= \int_0^t S(t-t')\chi_R(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt', \\ G^{(2)}(\cdot, t) &= \int_0^t S(t-t')(1-\chi_R)(2(|\Phi|^2\Phi)(\cdot, t') + \mathcal{F}(\Phi, v)(\cdot, t')) dt', \end{aligned}$$

for all  $t \in \mathbb{R}$ , so that  $G(\Phi, v) = G^{(1)} + G^{(2)}$  and  $G(\Psi_n, v_n) = G_n^{(1)} + G_n^{(2)}$ . Since  $S(t)\Psi_{n,0} \rightharpoonup S(t)\Phi_0$  in  $L^2(\mathbb{R})$  as  $n \rightarrow +\infty$  for all  $t \in \mathbb{R}$ , it is sufficient to show that

$$G(\Psi_n, v_n)(t) \rightharpoonup G(\Phi, v)(t) \quad \text{in } L^2(\mathbb{R})$$

as  $n \rightarrow +\infty$  for all  $t \in \mathbb{R}$ . Let  $\varphi \in L^2(\mathbb{R})$ . We write

$$\begin{aligned} &(G(\Psi_n, v_n)(t) - G(\Phi, v)(t), \varphi)_{L^2} \\ &= \int_{-\infty}^{+\infty} [G_n^{(1)}(x, t) - G^{(1)}(x, t)]\overline{\varphi(x)} dx + \int_{-\infty}^{+\infty} [G_n^{(2)}(x, t) - G^{(2)}(x, t)]\overline{\varphi(x)} dx \\ &= I_n^R(t) + J_n^R(t). \end{aligned}$$

For the first integral, using the Cauchy–Schwartz inequality, the Strichartz estimates for the admissible pairs (6, 6) and ( $\infty$ , 2), the Hölder inequality and (A-19), there exists a positive constant  $M$  such that for all  $t \in [0, T]$  we have

$$\begin{aligned} |I_n^R(t)| &\leq \|G_n^{(1)}(t) - G^{(1)}(t)\|_{L^2} \|\varphi\|_{L^2} \\ &\leq M \|\varphi\|_{L^2} \left( \| |\Psi_n|^2\Psi_n - |\Phi|^2\Phi \|_{L_{T,R}^{6/5}} + \|\mathcal{F}(\Psi_n, v_n) - \mathcal{F}(\Phi, v)\|_{L_T^1 L_R^2} \right) \\ &\leq M \|\varphi\|_{L^2} \left( \|\mathcal{F}(\Psi_n, v_n) - \mathcal{F}(\Phi, v)\|_{L_T^1 L_R^2} + \|\Psi_n - \Phi\|_{L_{T,R}^2} \left( \|\Psi_n\|_{L_{T,R}^6}^2 + \|\Phi\|_{L_{T,R}^6}^2 \right) \right). \end{aligned}$$

Then, using (A-18) and (A-28), we obtain for all  $t \in \mathbb{R}$

$$|I_n^R(t)| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Next, using the Hölder inequality we have

$$|J_n^R(t)| \leq 2 \left( \int_0^T \int_{-\infty}^{\infty} \left| |\Psi_n|^2 \Psi_n(x, t') - |\Phi|^2 \Phi(x, t') \right|^{6/5} dx dt' \right)^{5/6} \left( \int_0^T \int_{|x| \geq R} |S(t-t')\varphi|^6 dx dt' \right)^{1/6} \\ + \int_0^T \left( \int_{-\infty}^{\infty} |\mathcal{F}(\Psi_n, v_n)(x, t') - \mathcal{F}(\Phi, v)(x, t')|^2 dx \right)^{1/2} dt' \sup_{t' \in [0, T]} \left( \int_{|x| \geq R} |S(t-t')\varphi(x)|^2 dx \right)^{1/2}.$$

The terms on the right-hand side are bounded by a constant independent of  $n$ . Besides, since (6, 6) and  $(\infty, 2)$  are admissible pairs, we have

$$\|S(t)\varphi\|_{L^6_{T,x}} \leq M \|\varphi\|_{L^2(\mathbb{R})}, \\ \|S(t)\varphi\|_{L^\infty_T L^2(\mathbb{R})} \leq M \|\varphi\|_{L^2(\mathbb{R})},$$

so that, by the dominated convergence theorem and the fact that  $t \mapsto S(t)$  is uniformly continuous from  $[0, T]$  to  $L^2(\mathbb{R})$ , we obtain

$$\lim_{R \rightarrow \infty} \int_0^T \int_{|x| \geq R} |S(t)\varphi|^6 dx dt = \lim_{R \rightarrow \infty} \sup_{t \in [0, T]} \left( \int_{|x| \geq R} |S(t)\varphi(x)|^2 dx \right)^{1/2} = 0.$$

Hence,

$$\lim_{R \rightarrow \infty} |J_n^R(t)| = 0 \quad \text{uniformly with respect to } n \in \mathbb{N}$$

for any  $t \in [0, T]$ . This completes the proof of (A-41) and then of (A-38). This leads to the fact that  $\Phi \in C^0([0, T], L^2(\mathbb{R}))$ .

Now, let us prove that  $v \in C^0([0, T], H^1(\mathbb{R}))$ . Since  $(\Phi, v)$  satisfies the first equation in (2-34),  $\Phi \in L^\infty([0, T], L^2(\mathbb{R}))$  and  $F(\Psi, v) \in L^\infty([0, T], L^\infty(\mathbb{R}))$ , we have  $v \in H^1([0, T], H^{-1}(\mathbb{R}))$ . This yields, using the Sobolev embedding theorem,  $v \in C^0([0, T], H^{-1}(\mathbb{R}))$ . Let  $(t_1, t_2) \in [0, T]^2$ . We can write that

$$\int_{\mathbb{R}} |v(t_1, x) - v(t_2, x)|^2 dx = \langle v(t_1, x) - v(t_2, x), v(t_1, x) - v(t_2, x) \rangle_{H^{-1}, H^1} \\ \leq \|v(t_1, x) - v(t_2, x)\|_{H^{-1}} \|v(t_1, x) - v(t_2, x)\|_{H^1}.$$

Since  $v \in C^0([0, T], H^{-1}(\mathbb{R})) \cap L^\infty([0, T], H^1(\mathbb{R}))$ , we obtain  $v \in C^0([0, T], L^2(\mathbb{R}))$ . Next, we write

$$\|F(v, \Phi)(t_1) - F(v, \Phi)(t_2)\|_{L^\infty(\mathbb{R})} \leq \|v(t_1) - v(t_2)\|_{L^2} \|\Phi(t_1)\|_{L^2} + \|\Phi(t_2) - \Phi(t_1)\|_{L^2} \|v(t_2)\|_{L^2}.$$

Using the fact that  $\Phi, v \in C^0([0, T], L^2(\mathbb{R}))$ , we infer that  $F(v, \Phi) \in C^0([0, T], L^\infty(\mathbb{R}))$ . Then, by the second equation in (2-34),  $v \in C^0([0, T], H^1(\mathbb{R}))$ . This finishes the proof of this step, and of Proposition A.2. □

Finally, we give the proof of Proposition A.1.

*Proof of Proposition A.1.* In view of Proposition A.2, it is sufficient to prove the convergence of  $w_n$ . The proof follows the arguments in the proof of (A-27). Let  $\phi \in L^2(\mathbb{R})$ . We rely on (4-41) to write

$$\begin{aligned} & \int_{\mathbb{R}} [w^*(t, x) - w_n(t, x)]\phi(x) dx \\ &= 2 \int_{\mathbb{R}} \operatorname{Im} \left( \frac{\Psi^*(t, x)(1-2F(v^*, \Psi^*)(t, x))}{1-(v^*)^2(t, x)} - \frac{\Psi_n(t, x)(1-2F(v_n, \Psi_n)(t, x))}{1-(v_n)^2(t, x)} \right) \phi(x) dx \\ &= 2 \int_{\mathbb{R}} \operatorname{Im} \left( \frac{\Psi^*(t, x)}{1-(v^*)^2(t, x)} - \frac{\Psi_n(t, x)}{1-(v_n)^2(t, x)} \right) \phi(x) dx \\ &\quad - 4 \int_{\mathbb{R}} \operatorname{Im} \left( \frac{\Psi^*(t, x)F(v^*, \Psi^*)(t, x)}{1-(v^*)^2(t, x)} - \frac{\Psi_n(t, x)F(v_n, \Psi_n)(t, x)}{1-(v_n)^2(t, x)} \right) \phi(x) dx \end{aligned}$$

for all  $t \in [0, T]$ . Then, we use the same arguments as in the proof of (A-27) to show that the two last terms on the right-hand side go to 0 when  $n$  goes to  $+\infty$ . This finishes the proof of the proposition.  $\square$

**Exponential decay of  $\chi_c$ .** In this subsection, we recall the explicit formula and some useful properties of the operator  $\mathcal{H}_c$ , and then study its negative eigenfunction  $\chi_c$ . For  $c \in (-1, 1) \setminus \{0\}$ , the operator  $\mathcal{H}_c$  is given in explicit terms by

$$\mathcal{H}_c(\varepsilon) = \begin{pmatrix} \mathcal{L}_c(\varepsilon_v) + c^2 \frac{(1+v_c^2)^2}{(1-v_c^2)^3} \varepsilon_v - c \frac{1+v_c^2}{1-v_c^2} \varepsilon_w \\ -c \frac{1+v_c^2}{1-v_c^2} \varepsilon_v + (1-v_c^2) \varepsilon_w \end{pmatrix}, \tag{A-42}$$

where  $\varepsilon = (\varepsilon_v, \varepsilon_w)$  and

$$\mathcal{L}_c(\varepsilon_v) = -\partial_x \left( \frac{\partial_x \varepsilon_v}{1-v_c^2} \right) + (1-c^2 - (5+c^2)v_c^2 + 2v_c^4) \frac{\varepsilon_v}{(1-v_c^2)^2}.$$

In view of (A-42), the operator  $\mathcal{H}_c$  is an isomorphism from  $H^2(\mathbb{R}) \times L^2(\mathbb{R}) \cap \operatorname{Span}(\partial_x Q_c)^\perp$  onto  $\operatorname{Span}(\partial_x Q_c)^\perp$ . In addition, there exists a positive number  $A_c$ , depending continuously on  $c$ , such that

$$\|\mathcal{H}_c^{-1}(f, g)\|_{H^{k+2}(\mathbb{R}) \times H^k(\mathbb{R})} \leq A_c \|(f, g)\|_{H^k(\mathbb{R})^2} \tag{A-43}$$

for any  $(f, g) \in H^k(\mathbb{R})^2 \cap \operatorname{Span}(\partial_x Q_c)^\perp$  and any  $k \in \mathbb{N}$ .

The following proposition establishes the coercivity of the quadratic form  $H_c$  under suitable orthogonality conditions.

**Proposition A.3.** *Let  $c \in (-1, 1) \setminus \{0\}$ . There exists a positive number  $\Lambda_c$ , depending only on  $c$ , such that*

$$H_c(\varepsilon) \geq \Lambda_c \|\varepsilon\|_{H^1 \times L^2}^2 \tag{A-44}$$

for any pair  $\varepsilon \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$  satisfying the two orthogonality conditions

$$\langle \partial_x Q_c, \varepsilon \rangle_{L^2 \times L^2} = \langle \chi_c, \varepsilon \rangle_{L^2 \times L^2} = 0. \tag{A-45}$$

Moreover, the map  $c \mapsto \Lambda_c$  is uniformly bounded from below on any compact subset of  $(-1, 1) \setminus \{0\}$ .

The proof relies on standard Sturm–Liouville theory (see, e.g., the proof of Proposition 1 in [de Laire and Gravejat 2015] for more details).

Now, we turn to the analysis of the pair  $\chi_c$ .

**Lemma A.4.** *The pair  $\chi_c$  belongs to  $C^\infty(\mathbb{R}) \times C^\infty(\mathbb{R})$ . In addition, there exist two positive numbers  $A_c$  and  $a_c$ , depending continuously on  $c$ , such that  $a_c > \sqrt{1-c^2}$  and*

$$|\partial_x^k \chi_c| \leq A_c e^{-a_c |x|} \quad \text{on } \mathbb{R} \text{ for } k \in \{0, 1, 2\}. \quad (\text{A-46})$$

*Proof.* We set  $\chi_c := (\zeta_c, \xi_c)$ . Since  $\mathcal{H}_c(\chi_c) = -\tilde{\lambda}_c \chi_c$ , we have the following system

$$-\partial_x \left( \frac{\partial_x \zeta_c}{1-v_c^2} \right) + (1-c^2 - (5+c^2)v_c^2 + 2v_c^4) \frac{\zeta_c}{(1-v_c^2)^2} + c^2 \frac{(1+v_c^2)^2}{(1-v_c^2)^3} \zeta_c - c \frac{1+v_c^2}{1-v_c^2} \xi_c = -\tilde{\lambda}_c \zeta_c, \quad (\text{A-47})$$

$$c \frac{1+v_c^2}{1-v_c^2} \zeta_c = (1-v_c^2 + \tilde{\lambda}_c) \xi_c. \quad (\text{A-48})$$

It follows from standard elliptic theory that  $\chi_c \in H^2(\mathbb{R}) \times L^2(\mathbb{R})$ . Since the coefficients in (A-48) are smooth and bounded from above and below, we infer from a standard bootstrap argument that  $\chi_c \in C^\infty(\mathbb{R}) \times C^\infty(\mathbb{R})$ . Notice in particular that, by the Sobolev embedding theorem,  $\chi_c$  and  $\partial_x \chi_c$  are bounded on  $\mathbb{R}$ . Then, we deduce from the first statement in (5-11) that<sup>5</sup>

$$-\partial_{xx} \zeta_c + (1 + \tilde{\lambda}_c) \zeta_c - c \xi_c = \mathcal{O}(v_c^2), \quad (\text{A-49})$$

$$\zeta_c = \frac{1 + \tilde{\lambda}_c}{c} \xi_c + \mathcal{O}(v_c^2). \quad (\text{A-50})$$

Note that we have

$$B_c \exp(-\sqrt{1-c^2}|x|) \leq v_c(x) \leq A_c \exp(-\sqrt{1-c^2}|x|) \quad \text{for all } x \in \mathbb{R}, \quad (\text{A-51})$$

where  $B_c$  and  $A_c$  are two positive numbers.

In order to prove (A-46), we now introduce (A-50) into (A-49) to obtain

$$-\partial_{xx} \zeta_c + b_c^2 \zeta_c = \mathcal{O}(\exp(-2\sqrt{1-c^2}|x|)), \quad (\text{A-52})$$

$$\xi_c = \frac{c}{1 + \tilde{\lambda}_c} \zeta_c + \mathcal{O}(\exp(-2\sqrt{1-c^2}|x|)), \quad (\text{A-53})$$

with  $b_c^2 = \frac{1-c^2 + 2\tilde{\lambda}_c + (\tilde{\lambda}_c)^2}{1 + \tilde{\lambda}_c} > 1-c^2$ . Next, we set

$$g_c := -\partial_{xx} \zeta_c + b_c^2 \zeta_c, \quad (\text{A-54})$$

so that  $g_c(x) = \mathcal{O}(\exp(-2\sqrt{1-c^2}|x|))$  for all  $x \in \mathbb{R}$ . Using the variation of constants method, we obtain, for all  $x \in \mathbb{R}$ ,

$$\zeta_c(x) = A(x)e^{b_c x} + A_c e^{b_c x} + B(x)e^{-b_c x} + B_c e^{-b_c x},$$

<sup>5</sup>The notation  $\mathcal{O}(v_c^2)$  refers to a quantity bounded by  $A_c v_c^2$  (pointwise), where the positive number  $A_c$  depends only on  $c$ .

with

$$A(x) = \frac{-1}{2b_c} \int_0^x e^{-b_c t} g_c(t) dt$$

and

$$B(x) = \frac{-1}{2b_c} \int_0^x e^{b_c t} g_c(t) dt.$$

Since  $\zeta_c \in L^2(\mathbb{R})$ , this leads to

$$\zeta_c(x) = \mathcal{O}(\exp(-2\sqrt{1-c^2}|x|) + \exp(-b_c|x|)).$$

Hence, we can take  $a_c = \min\{2\sqrt{1-c^2}, b_c\}$  and invoke (A-50) to obtain (A-46) for  $k = 0$ . Using (5-10), (5-11), (A-47), (A-48) and (A-51), we extend (A-46) to  $k \in 1, 2$ .  $\square$

### Acknowledgments

I would like to thank both my supervisors R. Côte and P. Gravejat for their support throughout the time I spent to finish this manuscript, offering invaluable advice. I am also grateful to Y. Martel for interesting and helpful discussions. I am also indebted to the anonymous referee for valuable comments and remarks. This work is supported by a Ph.D. grant from “Région Ile-de-France” and is partially sponsored by the project “Schrödinger equations and applications” (ANR-12-JS01-0005-01) of the Agence Nationale de la Recherche.

### References

- [Béthuel et al. 2008] F. Béthuel, P. Gravejat, and J.-C. Saut, “Existence and properties of travelling waves for the Gross–Pitaevskii equation”, pp. 55–103 in *Stationary and time dependent Gross–Pitaevskii equations*, edited by A. Farina and J.-C. Saut, *Contemp. Math.* **473**, American Mathematical Society, Providence, RI, 2008. [MR 2522014](#) [Zbl 1216.35132](#)
- [Béthuel et al. 2014] F. Béthuel, P. Gravejat, and D. Smets, “Stability in the energy space for chains of solitons of the one-dimensional Gross–Pitaevskii equation”, *Ann. Inst. Fourier (Grenoble)* **64**:1 (2014), 19–70. [MR 3330540](#) [Zbl 06387265](#)
- [Béthuel et al. 2015] F. Béthuel, P. Gravejat, and D. Smets, “Asymptotic stability in the energy space for dark solitons of the Gross–Pitaevskii equation”, *Ann. Sci. Éc. Norm. Supér.* (4) **48**:6 (2015), 1327–1381.
- [Buslaev and Perelman 1993] V. S. Buslaev and G. S. Perelman, “Scattering for the nonlinear Schrödinger equation: states that are close to a soliton”, *St. Petersburg Math. J.* **4**:6 (1993), 1111–1142. [MR 1199635](#) [Zbl 0853.35112](#)
- [Buslaev and Perelman 1995] V. S. Buslaev and G. S. Perelman, “On the stability of solitary waves for nonlinear Schrödinger equations”, pp. 75–98 in *Nonlinear evolution equations*, edited by N. N. Uraltseva, *Amer. Math. Soc. Transl. Ser. 2* **164**, American Mathematical Society, Providence, RI, 1995. [MR 1334139](#) [Zbl 0841.35108](#)
- [Buslaev and Sulem 2003] V. S. Buslaev and C. Sulem, “On asymptotic stability of solitary waves for nonlinear Schrödinger equations”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **20**:3 (2003), 419–475. [MR 1972870](#) [Zbl 1028.35139](#)
- [Cuccagna 2001] S. Cuccagna, “Stabilization of solutions to nonlinear Schrödinger equations”, *Comm. Pure Appl. Math.* **54**:9 (2001), 1110–1145. [MR 1835384](#) [Zbl 1031.35129](#)
- [Cuccagna 2003] S. Cuccagna, “On asymptotic stability of ground states of NLS”, *Rev. Math. Phys.* **15**:8 (2003), 877–903. [MR 2027616](#) [Zbl 1084.35089](#)
- [Cuccagna 2011] S. Cuccagna, “The Hamiltonian structure of the nonlinear Schrödinger equation and the asymptotic stability of its ground states”, *Comm. Math. Phys.* **305**:2 (2011), 279–331. [MR 2805462](#) [Zbl 1222.35183](#)
- [Escarriaza et al. 2008] L. Escarriaza, C. E. Kenig, G. Ponce, and L. Vega, “Hardy’s uncertainty principle, convexity and Schrödinger evolutions”, *J. Eur. Math. Soc. (JEMS)* **10**:4 (2008), 883–907. [MR 2443923](#) [Zbl 1158.35018](#)

- [Gang and Sigal 2007] Z. Gang and I. M. Sigal, “Relaxation of solitons in nonlinear Schrödinger equations with potential”, *Adv. Math.* **216**:2 (2007), 443–490. [MR 2351368](#) [Zbl 1126.35065](#)
- [Goubet and Molinet 2009] O. Goubet and L. Molinet, “Global attractor for weakly damped nonlinear Schrödinger equations in  $L^2(\mathbb{R})$ ”, *Nonlinear Anal.* **71**:1–2 (2009), 317–320. [MR 2518038](#) [Zbl 1170.35534](#)
- [Grillakis et al. 1987] M. Grillakis, J. Shatah, and W. Strauss, “Stability theory of solitary waves in the presence of symmetry. I”, *J. Funct. Anal.* **74**:1 (1987), 160–197. [MR 901236](#) [Zbl 0656.35122](#)
- [Guo and Ding 2008] B. Guo and S. Ding, *Landau–Lifshitz equations*, Frontiers of Research with the Chinese Academy of Sciences **1**, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2008. [MR 2432099](#) [Zbl 1158.35096](#)
- [Hubert and Schäfer 1998] A. Hubert and R. Schäfer, *Magnetic domains: the analysis of magnetic microstructures*, Springer, New York, 1998.
- [Jerrard and Smets 2012] R. L. Jerrard and D. Smets, “On Schrödinger maps from  $T^1$  to  $S^2$ ”, *Ann. Sci. Éc. Norm. Supér.* (4) **45**:4 (2012), 637–680. [MR 3059243](#) [Zbl 1308.58023](#)
- [Kenig et al. 2003] C. E. Kenig, G. Ponce, and L. Vega, “On unique continuation for nonlinear Schrödinger equations”, *Comm. Pure Appl. Math.* **56**:9 (2003), 1247–1262. [MR 1980854](#) [Zbl 1041.35072](#)
- [Kosevich et al. 1990] A. M. Kosevich, B. A. Ivanov, and A. S. Kovalev, “Magnetic Solitons”, *Phys. Rep.* **194**:3–4 (1990), 117–238.
- [de Laire 2014] A. de Laire, “Minimal energy for the traveling waves of the Landau–Lifshitz equation”, *SIAM J. Math. Anal.* **46**:1 (2014), 96–132. [MR 3148081](#) [Zbl 1305.35123](#)
- [de Laire and Gravejat 2015] A. de Laire and P. Gravejat, “Stability in the energy space for chains of solitons of the Landau–Lifshitz equation”, *J. Differential Equations* **258**:1 (2015), 1–80. [MR 3271297](#) [Zbl 1301.35173](#)
- [Landau and Lifshitz 1935] L. D. Landau and E. M. Lifshitz, “On the theory of the dispersion of magnetic permeability in ferromagnetic bodies”, *Phys. Z. Sowjetunion* **8** (1935), 153–169. [Zbl 0012.28501](#)
- [Linares and Ponce 2009] F. Linares and G. Ponce, *Introduction to nonlinear dispersive equations*, Springer, New York, 2009. [MR 2492151](#) [Zbl 1178.35004](#)
- [Martel 2006] Y. Martel, “Linear problems related to asymptotic stability of solitons of the generalized KdV equations”, *SIAM J. Math. Anal.* **38**:3 (2006), 759–781. [MR 2262941](#) [Zbl 1126.35055](#)
- [Martel and Merle 2008a] Y. Martel and F. Merle, “Asymptotic stability of solitons of the gKdV equations with general nonlinearity”, *Math. Ann.* **341**:2 (2008), 391–427. [MR 2385662](#) [Zbl 1153.35068](#)
- [Martel and Merle 2008b] Y. Martel and F. Merle, “Refined asymptotics around solitons for gKdV equations”, *Discrete Contin. Dyn. Syst.* **20**:2 (2008), 177–218. [MR 2358258](#) [Zbl 1137.35062](#)
- [Mizumachi 2001] T. Mizumachi, “Large time asymptotics of solutions around solitary waves to the generalized Korteweg-de Vries equations”, *SIAM J. Math. Anal.* **32**:5 (2001), 1050–1080. [MR 1828318](#) [Zbl 0981.35066](#)
- [Pego and Weinstein 1994] R. L. Pego and M. I. Weinstein, “Asymptotic stability of solitary waves”, *Comm. Math. Phys.* **164**:2 (1994), 305–349. [MR 1289328](#) [Zbl 0805.35117](#)
- [Soffer and Weinstein 1989] A. Soffer and M. I. Weinstein, “Multichannel nonlinear scattering theory for nonintegrable equations”, pp. 312–327 in *Integrable systems and applications* (Île d’Oléron, 1988), edited by M. Balabane et al., Lecture Notes in Phys. **342**, Springer, Berlin, 1989. [MR 1034551](#) [Zbl 0712.35074](#)
- [Soffer and Weinstein 1990] A. Soffer and M. I. Weinstein, “Multichannel nonlinear scattering for nonintegrable equations”, *Comm. Math. Phys.* **133**:1 (1990), 119–146. [MR 1071238](#) [Zbl 0721.35082](#)
- [Soffer and Weinstein 1992] A. Soffer and M. I. Weinstein, “Multichannel nonlinear scattering for nonintegrable equations, II. The case of anisotropic potentials and data”, *J. Differential Equations* **98**:2 (1992), 376–390. [MR 1170476](#) [Zbl 0795.35073](#)

Received 16 Jul 2015. Revised 25 Nov 2015. Accepted 30 Jan 2016.

YAKINE BAHRI: [yakine.bahri@polytechnique.edu](mailto:yakine.bahri@polytechnique.edu)

Centre de Mathématiques Laurent Schwartz, École Polytechnique, 91128 Palaiseau Cedex, France

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

[patrick.gerard@math.u-psud.fr](mailto:patrick.gerard@math.u-psud.fr)

Université Paris Sud XI

Orsay, France

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>	Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy <a href="mailto:berti@sissa.it">berti@sissa.it</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>	Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Vaughan Jones	U.C. Berkeley & Vanderbilt University <a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a>	Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>
Vadim Kaloshin	University of Maryland, USA <a href="mailto:vadim.kaloshin@gmail.com">vadim.kaloshin@gmail.com</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Richard B. Melrose	Massachusetts Inst. of Tech., USA <a href="mailto:rbb@math.mit.edu">rbb@math.mit.edu</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>
William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>	Maciej Zworski	University of California, Berkeley, USA <a href="mailto:zvorski@math.berkeley.edu">zvorski@math.berkeley.edu</a>
Clément Mouhot	Cambridge University, UK <a href="mailto:c.mouhot@dpms.cam.ac.uk">c.mouhot@dpms.cam.ac.uk</a>		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2016 is US \$/year for the electronic version, and \$/year (+\$, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 9 No. 3 2016

---

Local analytic regularity in the linearized Calderón problem	515
JOHANNES SJÖSTRAND and GUNTHER UHLMANN	
Dispersive estimates for the Schrödinger operator on step-2 stratified Lie groups	545
HAJER BAHOURI, CLOTILDE FERMANIAN-KAMMERER and ISABELLE GALLAGHER	
Obstructions to the existence of limiting Carleman weights	575
PABLO ANGULO-ARDOY, DANIEL FARACO, LUIS GUIJARRO and ALBERTO RUIZ	
Finite chains inside thin subsets of $\mathbb{R}^d$	597
MICHAEL BENNETT, ALEXANDER IOSEVICH and KRYSTAL TAYLOR	
Advection-diffusion equations with density constraints	615
ALPÁR RICHÁRD MÉSZÁROS and FILIPPO SANTAMBROGIO	
Asymptotic stability in energy space for dark solitons of the Landau–Lifshitz equation	645
YAKINE BAHRI	
On the well-posedness of the generalized Korteweg–de Vries equation in scale-critical $\hat{L}^r$ -space	699
SATOSHI MASAKI and JUN-ICHI SEGATA	
Regularity for parabolic integro-differential equations with very irregular kernels	727
RUSSELL W. SCHWAB and LUIS SILVESTRE	