

ANALYSIS & PDE

Volume 10

No. 7

2017

JOSÉ A. CAÑIZO, AMIT EINAV AND BERTRAND LODS

**TREND TO EQUILIBRIUM FOR THE BECKER-DÖRING
EQUATIONS:
AN ANALOGUE OF CERCIGNANI'S CONJECTURE**

TREND TO EQUILIBRIUM FOR THE BECKER–DÖRING EQUATIONS: AN ANALOGUE OF CERCIGNANI’S CONJECTURE

JOSÉ A. CAÑIZO, AMIT EINAV AND BERTRAND LODS

We investigate the rate of convergence to equilibrium for subcritical solutions to the Becker–Döring equations with physically relevant coagulation and fragmentation coefficients and mild assumptions on the given initial data. Using a discrete version of the log-Sobolev inequality with weights, we show that in the case where the coagulation coefficient grows linearly and the detailed balance coefficients are of typical form, one can obtain a linear functional inequality for the dissipation of the relative free energy. This results in showing Cercignani’s conjecture for the Becker–Döring equations and consequently in an exponential rate of convergence to equilibrium. We also show that for all other typical cases, one can obtain an “almost” Cercignani’s conjecture, which results in an algebraic rate of convergence to equilibrium.

1. Introduction	1663
2. A discrete weighted logarithmic Sobolev inequality	1674
3. Energy dissipation inequalities	1683
4. Rate of convergence to equilibrium	1696
5. Consequences for general coagulation and fragmentation models	1697
Appendix A. Additional computations for discrete log-Sobolev with weights	1701
Appendix B. Additional useful computations	1705
References	1707

1. Introduction

The Becker–Döring equations. The Becker–Döring equations are a fundamental set of equations which describe the kinetics of a first-order phase transition. Amongst the phenomena to which they are relevant one can find crystallisation [Kelton et al. 1983], nucleation of polymers [Capasso 2003], vapour condensation, aggregation of lipids [Neu et al. 2002] and phase separation in alloys [Xiao and Haasen 1991]. For more general reviews of nucleation theory, see, for instance, [Schmelzer 2005; Oxtoby 1992].

Cañizo was supported by the Marie-Curie CIG grant KineticCF and the Spanish Ministerio de Economía y Competitividad/European Regional Development Fund (ERDF/FEDER), project MTM2014-52056-P. Einav was supported by EPSRC grant EP/L002302/1. Lods acknowledges support of the “de Castro” Statistics Initiative, Collegio C. Alberto, Moncalieri, Italy.

MSC2010: 34D05, 35Q82, 82C05, 82C40.

Keywords: Becker–Döring, nucleation, exponential convergence, entropy method.

The Becker–Döring equations give the time evolution of the size distribution of clusters of a certain substance. Denoting by $\{c_i(t)\}_{i \in \mathbb{N}}$ the density of clusters of size i at time $t \geq 0$ (i.e., the density of clusters that are composed of i particles), the equations read


$$\frac{d}{dt}c_i(t) = W_{i-1}(t) - W_i(t), \quad i \in \mathbb{N} \setminus \{1\}, \tag{1-1a}$$

$$\frac{d}{dt}c_1(t) = -W_1(t) - \sum_{k=1}^{\infty} W_k(t), \tag{1-1b}$$

where

$$W_i(t) := a_i c_1(t)c_i(t) - b_{i+1} c_{i+1}(t), \quad i \in \mathbb{N}, \tag{1-2}$$

and a_i, b_i , assumed to be strictly positive, are the *coagulation and fragmentation coefficients*. They determine, respectively, the rate at which clusters of size i combine with clusters of size 1 to create clusters of size $i + 1$, and the rate at which clusters of size $i + 1$ split into clusters of sizes i and 1. This corresponds to the basic assumption of the underlying model: if we represent symbolically by $\{i\}$ the chemical species of clusters of size i , then the only (relevant) chemical reactions that take place are



The quantity $W_i(t)$ defined in (1-2) represents the *net rate* of the reaction $\{i\} + \{1\} \rightleftharpoons \{i + 1\}$, and under the above set of equations it is easy to see that the *density*, or *mass*, of the solution, defined by

$$\varrho := \sum_{i=1}^{\infty} i c_i(0) = \sum_{i=1}^{\infty} i c_i(t), \tag{1-3}$$

is formally conserved under time evolution. The original equations proposed by Becker and Döring [1935] were similar to (1-1), with the slight change that the density of one particle c_1 , usually called the monomer density, was assumed to be constant. The current version, motivated by the conservation of total density, was first discussed in [Burton 1977] and [Penrose and Lebowitz 1979] and is widely used in classical nucleation theory.

Much like in other kinetic equations, the study of a state of equilibrium and the convergence to it is a fundamental question in the study of the Becker–Döring equations. Defining the *detailed balance coefficients* Q_i recursively by

$$Q_1 = 1, \quad Q_{i+1} = \frac{a_i}{b_{i+1}} Q_i, \quad i \in \mathbb{N}, \tag{1-4}$$

one can see that for a given $z \geq 0$ the sequence

$$c_i = Q_i z^i \tag{1-5}$$

is formally an equilibrium of (1-1). However, depending on the coagulation and fragmentation coefficients a_i and b_i , many of these formal equilibria do not have a finite mass. The largest $z_s \geq 0$, possibly $z_s = +\infty$, for which

$$\sum_{i=1}^{\infty} i Q_i z^i < +\infty \quad \text{for all } 0 \leq z < z_s$$

is called the *critical monomer density*, or sometimes the monomer saturation density. The *critical mass* (or, again, saturation mass) is then defined by

$$\varrho_s := \sum_{i=1}^{\infty} i Q_i z_s^i \in [0, +\infty]. \quad (1-6)$$

It is important to note that both z_s and ϱ_s are uniquely determined by a_i and b_i and that $\{Q_i z^i\}_{i \in \mathbb{N}}$ is a finite-mass equilibrium only for $0 \leq z < z_s$, with the possibility for the equality $z = z_s$ only when $\varrho_s < +\infty$. Additionally, it is easy to see that for a given finite mass $\varrho \leq \varrho_s$ there exists a unique $\bar{z} \geq 0$ such that

$$\varrho = \sum_{i=1}^{\infty} i Q_i \bar{z}^i,$$

giving us a candidate for the asymptotic equilibrium state of (1-1) under a given initial data. These are in fact the only finite-mass equilibria (see [Ball et al. 1986]), and \bar{z} defined above is called the *equilibrium monomer density* for a given mass ϱ .

A finite mass solution is called *subcritical* when its mass ϱ is strictly less than ϱ_s . It is called *critical* if $\varrho = \varrho_s$ and *supercritical* if $\varrho > \varrho_s$, assuming $\varrho_s < +\infty$. In this paper we will only concern ourselves with subcritical solutions. Thus, to avoid triviality we always assume that $z_s > 0$.

The critical density ϱ_s , if finite, marks a change in the behaviour of equilibrium states: if $\varrho < \varrho_s$ then a unique equilibrium state with mass ϱ exists, while if $\varrho > \varrho_s$, no such equilibrium can occur and a phase transition phenomenon takes place — reflected in the fact that the excess density $\varrho - \varrho_s$ is concentrated in larger and larger clusters as time progresses.

Previous results. Let us briefly review existing results on the mathematical theory of the Becker–Döring equations, which has advanced much since the first rigorous works on the topic [Ball and Carr 1988; Ball et al. 1986]. In [Ball et al. 1986] the authors showed (among other things) existence and uniqueness of a global solution to (1-1) when

$$a_i \leq C_1 i, \quad b_i \leq C_2 i, \quad \sum_{i=1}^{\infty} i^{1+\varepsilon} c_i(0) < +\infty \quad (1-7)$$

for some constants $C_1, C_2, \varepsilon > 0$. As expected, under the above assumptions the unique solution conserves mass (that is, (1-3) holds rigorously). This basic existence theory is applicable to all solutions we consider in this work.

The asymptotic behaviour of solutions to (1-1) is one of the most interesting aspects of the equation. Supercritical behaviour, while not dealt with in this work, has a particularly interesting link to late-stage coarsening and has been studied extensively in [Penrose 1997; Velázquez 1998; Collet et al. 2002; Niethammer 2003], with several questions still open. Asymptotic approximations of such solutions have been developed in [Farjoun and Neu 2008; 2011; Neu et al. 2005].

Regarding the subcritical regime, it was proved in [Ball and Carr 1988; Ball et al. 1986] that solutions with subcritical mass ϱ approach the unique equilibrium with this mass, determined by (1-3). A

fundamental quantity in understanding this approach is the *free energy*, $H(\mathbf{c})$, defined for any nonnegative sequence $\mathbf{c} = \{c_i\}_{i \in \mathbb{N}}$ by

$$H(\mathbf{c}) := \sum_{i=1}^{\infty} c_i \left(\log \frac{c_i}{Q_i} - 1 \right) \tag{1-8}$$

whenever the sum converges. It can be shown that $H(\mathbf{c}(t))$ decreases along solutions $\mathbf{c} = \mathbf{c}(t)$ to the Becker–Döring equations; in fact, for a (strictly positive, suitably decaying for large i) solution $\mathbf{c}(t) = \{c_i(t)\}_{i \in \mathbb{N}}$ of (1-1) we have

$$\frac{d}{dt} H(\mathbf{c}(t)) = -D(\mathbf{c}(t)) := - \sum_{i=1}^{\infty} a_i Q_i \left(\frac{c_1 c_i}{Q_i} - \frac{c_{i+1}}{Q_{i+1}} \right) \left(\log \frac{c_1 c_i}{Q_i} - \log \frac{c_{i+1}}{Q_{i+1}} \right) \leq 0. \tag{1-9}$$

This free energy is motivated by physical considerations and constitutes a Lyapunov functional for our equation. Since it does not have a definite sign, we define a more natural candidate to measure the distance of $\mathbf{c}(t) = \{c_i(t)\}_{i \in \mathbb{N}}$ to the equilibrium. Using the notation

$$(Q_z)_i = Q_i z^i$$

and denoting $Q_{\bar{z}}$ by \mathcal{Q} , we can define the *relative free energy* as

$$H(\mathbf{c}|\mathcal{Q}) := \sum_{i=1}^{\infty} c_i \left(\log \frac{c_i}{\bar{z}^i Q_i} - 1 \right) + \sum_{i=1}^{\infty} \bar{z}^i Q_i = H(\mathbf{c}) - \log \bar{z} \sum_{i=1}^{\infty} i c_i + \sum_{i=1}^{\infty} \bar{z}^i Q_i. \tag{1-10}$$

The relative free energy has the same time derivative as the free energy, and thus the same monotonicity property

$$\frac{d}{dt} H(\mathbf{c}(t)|\mathcal{Q}) = -D(\mathbf{c}(t)) \quad \forall t \geq 0,$$

where the *free energy dissipation* D is defined in (1-9). The relative free energy also satisfies

- $H(\mathbf{c}|\mathcal{Q}) \geq 0$, as can be seen by writing

$$H(\mathbf{c}|\mathcal{Q}) = \sum_{i=1}^{\infty} Q_i \varphi \left(\frac{c_i}{Q_i} \right), \quad \text{with } \varphi(r) := r \log r - r + 1 \geq 0, \tag{1-11}$$

- $H(\mathbf{c}|\mathcal{Q}) = 0$ if and only if $c_i = Q_i = Q_i \bar{z}^i$ for any $i \in \mathbb{N}$, which is readily seen from (1-11).

This hints that $H(\mathbf{c}|\mathcal{Q})$ is the right “distance” to investigate. Indeed, while $H(\mathbf{c}|\mathcal{Q})$ is not a distance strictly speaking, it does control the ℓ^1 distance between \mathbf{c} and \mathcal{Q} by the celebrated Csiszár–Kullback inequality,¹ which in our case translates to

$$\|\mathbf{c} - \mathcal{Q}\|_{\ell^1(\mathbb{N})} = \sum_{i=1}^{\infty} |c_i - Q_i| \leq \sqrt{2\varrho H(\mathbf{c}|\mathcal{Q})}. \tag{1-12}$$

(See also [Jabin and Niethammer 2003, Corollary 2.2] for a version involving the ℓ^1 distance with weight i .) The issue of estimating the rate of convergence to equilibrium of subcritical solutions is the main concern

¹Sometimes called the Pinsker or Kullback–Pinsker inequality.

of this paper. The first result in this direction was [Jabin and Niethammer 2003], where they investigated the possibility of applying the so-called *entropy method* to the Becker–Döring equation. This consists roughly in looking for functional inequalities between a suitable Lyapunov functional of the equation (generally called the entropy; it corresponds to the relative free energy in our case) and its dissipation, so that one obtains a differential inequality that estimates the rate of convergence to equilibrium. In the case of the Becker–Döring equation, it was proved in [Jabin and Niethammer 2003] that there exists a constant $C > 0$, depending only on the fixed parameters of the problem and the initial data, such that

$$D(\mathbf{c}) \geq C \frac{H(\mathbf{c}|\mathcal{Q})}{(\log H(\mathbf{c}|\mathcal{Q}))^2} \quad (1-13)$$

for all nonnegative sequences \mathbf{c} with subcritical mass ϱ , satisfying $\varepsilon \leq c_1 \leq z_s - \varepsilon$ for some $\varepsilon > 0$ and

$$\sum_{i=1}^{\infty} e^{\mu i} c_i =: M^{\text{exp}} < +\infty. \quad (1-14)$$

The constant C depends on ε and M^{exp} . This result applies under reasonable conditions on the coefficients a_i and b_i ; in particular, it applies to the coefficients (1-23) and (1-25), which we give as examples below. If we consider now a solution $\mathbf{c} = \mathbf{c}(t)$ to (1-1), we may apply the inequality (1-13) to $\mathbf{c}(t)$ as long as $\mathbf{c}(t)$ satisfies the appropriate conditions, obtaining

$$\frac{d}{dt} H(\mathbf{c}(t)|\mathcal{Q}) = -D(\mathbf{c}(t)) \leq -C \frac{H(\mathbf{c}(t)|\mathcal{Q})}{(\log H(\mathbf{c}(t)|\mathcal{Q}))^2}.$$

Adding to this some additional considerations for the times t for which the inequality (1-13) is not applicable to $\mathbf{c}(t)$, one can deduce that $H(\mathbf{c}(t)|\mathcal{Q})$ is (roughly) bounded above by the solution of the above differential inequality, namely that

$$H(\mathbf{c}(t)|\mathcal{Q}) \leq H(\mathbf{c}(0)|\mathcal{Q}) e^{-Kt^{1/3}}$$

for some $K > 0$. Using inequality (1-12), this gives an almost-exponential rate of convergence to equilibrium for subcritical solutions in the $\ell^1(\mathbb{N})$ norm.

The question remained open of whether the convergence is in fact exponential or not. Recently this has been answered positively in [Cañizo and Lods 2013] by two of the authors of the present paper, through a different approach involving a detailed study of the spectrum of the linearisation of equation (1-1) around a subcritical equilibrium. This is an approach with a strong analogy to results in the theory of the Boltzmann equation; we refer to [Cañizo and Lods 2013; Villani 2003; Desvillettes et al. 2011] for more details on this parallel. The idea of the argument is to use the inequality (1-13) when one is far from equilibrium. Then, once we have reached a region which is close enough to equilibrium, the linearised regime is dominant and one can use the spectral study of the linearised operator in order to show that the convergence is in fact exponential. The outcome of this strategy is the following: for many interesting coefficients (including (1-23) and (1-25)), subcritical solutions $\mathbf{c} = \mathbf{c}(t)$ to (1-1) with

$$\sum_{i=1}^{\infty} e^{\mu i} c_i(0) =: M^{\text{exp}} < +\infty \quad \text{for some } \mu > 0$$

satisfy

$$\sum_{i=1}^{\infty} e^{\mu' i} |c_i(t) - Q_i| \leq C e^{-\lambda t} \quad \text{for } t \geq 0$$

for some $0 < \mu' < \mu$, $C > 0$ and $\lambda > 0$ which depend on the parameters of the problem and on M^{exp} . In fact, μ' and C only depend on the initial data $c(0)$ through its mass and the value of M^{exp} ; λ depends only on the coefficients and the initial mass and can be estimated explicitly. The value of λ is bounded above by (and can be taken very close to) the size of the spectral gap of the linearised operator. Recently Murray and Pego [2015] have used this spectral gap and developed the local estimates of the linearised operator in order to obtain convergence to equilibrium at a polynomial rate with milder conditions on the decay of the initial data. These results, like those in [Cañizo and Lods 2013], are local in nature and require the use of some global estimate such as (1-13) in order to provide global rates of convergence to equilibrium.

Main results. Our main goal in this work is to complete the picture of convergence to equilibrium by investigating modified and improved versions of the inequality (1-13). We show optimal inequalities and settle the question of whether full exponential convergence can be obtained through a *linear* inequality of the form

$$D(c) \geq KH(c|Q)$$

for some constant $K > 0$. In analogy to the Boltzmann equation, we refer to the question of whether such K exists along solutions to (1-1) as *Cercignani’s conjecture for the Becker–Döring equations*. In fact, we show that under relatively mild conditions on the initial data, typical coagulation and fragmentation coefficients (covering the physically relevant situations; see the next subsection) admit an “almost” Cercignani conjecture for the energy dissipation, i.e., an inequality bounding $D(c)$ below by a power of $H(c|Q)$, yielding an explicit rate of convergence to equilibrium. Surprisingly, we also find a relevant case ($a_i \sim i$ for all i) for which the conjecture is actually valid.

We will often require the following assumptions on the coagulation and fragmentation coefficients. Some of these are similar to those in [Jabin and Niethammer 2003], and always include physically relevant coefficients, such as those described in the next subsection. We recall that we always assume $a_i, b_i > 0$ for all $i \in \mathbb{N}$, and that the detailed balance coefficients Q_i were defined in (1-4) — given a_i one can determine b_i through Q_i , and vice versa.

Hypothesis 1. $0 < z_s < +\infty$.

Hypothesis 2. For all $i \in \mathbb{N}$, we have $Q_i = z_s^{1-i} \alpha_i$, where $\{\alpha_i\}_{i \in \mathbb{N}}$ is a nonincreasing positive sequence with $\alpha_1 = 1$ and $\lim_{i \rightarrow \infty} \alpha_{i+1}/\alpha_i = 1$.

Hypothesis 3. There exist $C_1, C_2 > 0$ such that

$$C_1 i^\nu \leq a_i \leq C_2 i^\nu \quad \text{for all } i \in \mathbb{N}.$$

Hypothesis 2 on the form of Q_i is given as a compromise that allows us to give simple quantitative estimates of the constants in our theorems while allowing for the most commonly used types of coefficients. As one can see from the proofs, this assumption may be relaxed at the price of obtaining more involved estimates for our constants, particularly the logarithmic Sobolev constant in [Proposition 3.4](#).

In most of our estimates a crucial role will be played by the *lower free energy dissipation*, $\bar{D}(\mathbf{c})$, defined for a given nonnegative sequence \mathbf{c} by

$$\bar{D}(\mathbf{c}) = \sum_{i=1}^{\infty} a_i Q_i \left(\sqrt{\frac{c_1 c_i}{Q_i}} - \sqrt{\frac{c_{i+1}}{Q_{i+1}}} \right)^2. \quad (1-15)$$

At this point one notices that the elementary inequality $(x - y)(\log x - \log y) \geq 4(\sqrt{x} - \sqrt{y})^2$ when $x, y > 0$ implies

$$D(\mathbf{c}) \geq 4\bar{D}(\mathbf{c})$$

for any nonnegative sequence \mathbf{c} . Thus, any lower bound that is obtained for $\bar{D}(\mathbf{c})$ will transfer immediately to $D(\mathbf{c})$.

We now state our main result on general functional inequalities for the free energy dissipation, from which later we conclude a quantitative rate of convergence to equilibrium. It can be divided into two parts: functional inequalities when c_1 is not too small and not too far from \bar{z} , and inequalities in the case where c_1 escapes the above region.

Theorem 1.1. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ satisfy Hypotheses 1–3 and let $\mathbf{c} = \{c_i\}_{i \in \mathbb{N}}$ be an arbitrary positive sequence with finite total density $0 < \varrho < \varrho_s$.*

(i) **Estimate for $a_i \sim i$.** *Assume that $\gamma = 1$ and that there exist $\delta > 0$ such that*

$$\delta < c_1 < z_s - \delta. \quad (1-16)$$

Then there exists $K > 0$ depending only on δ, ϱ and the coefficients $\{a_i\}_{i \geq 1}, \{b_i\}_{i \geq 2}$, such that

$$\bar{D}(\mathbf{c}) \geq KH(\mathbf{c}|\mathcal{Q}). \quad (1-17)$$

(ii) **Estimate for $a_i \sim i^\gamma$ with $\gamma < 1$.** *Assume that $0 \leq \gamma < 1$ and that c_1 satisfies (1-16) for some $\delta > 0$. If, in addition, there exists $\beta > 1$ with*

$$M_\beta(\mathbf{c}) = \sum_{i=1}^{\infty} i^\beta c_i < +\infty \quad (1-18)$$

then there exists $K > 0$ depending only on $\delta, \varrho, M_\beta(\mathbf{c})$ and the coefficients $\{a_i\}_{i \geq 1}, \{b_i\}_{i \geq 2}$ such that

$$\bar{D}(\mathbf{c}) \geq KH(\mathbf{c}|\mathcal{Q})^{\frac{\beta-\gamma}{\beta-1}}. \quad (1-19)$$

(iii) **Estimate for c_1 far from equilibrium.** *Assume that $\gamma = 1$, or that $0 \leq \gamma < 1$ and (1-18) holds for some $\beta > 1$. Assume also that for some $\delta > 0$*

$$c_1 \leq \delta$$

or that

$$c_1 \geq z_s - \delta;$$

i.e., c_1 is outside of the range given in (1-16). Then if $\delta > 0$ is small enough (depending only on ϱ and $\{Q_i\}_{i \geq 1}$), there exists $\varepsilon > 0$ depending only on δ, ϱ and the coefficients $\{a_i\}_{i \geq 1}, \{b_i\}_{i \geq 2}$ if

$\gamma = 1$ (and additionally on $M_\beta(\mathbf{c})$ if $\gamma < 1$) such that

$$\bar{D}(\mathbf{c}) \geq \varepsilon. \tag{1-20}$$

The constants K and ε can be estimated explicitly in all cases.

We emphasise that all constants in the above theorem depend only on ϱ , the coefficients $\{a_i\}_{i \in \mathbb{N}}$, $\{b_i\}_{i \in \mathbb{N}}$, and the additional bounds δ or M_β (notice that ϱ_s is determined by the coefficients alone). Case (ii) of Theorem 1.1 is optimal in the following sense:

Theorem 1.2. Call X_ϱ the set of nonnegative sequences $\mathbf{c} = \{c_i\}_{i \in \mathbb{N}}$ with mass ϱ , i.e., such that $\sum_{i=1}^\infty i c_i = \varrho$. Then, there exist $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ that satisfy Hypotheses 1–3 with $\gamma < 1$ such that

$$\inf_{X_\varrho} \frac{D(\mathbf{c})}{H(\mathbf{c}|Q)} = 0$$

for any $\varrho < \varrho_s$.

In other words, this shows that a linear inequality as that of Theorem 1.1(i) cannot hold if $a_i \sim i^\gamma$ with $\gamma < 1$.

The idea behind the proof of Theorem 1.1 is to use a discrete logarithmic Sobolev inequality with weights, motivated by works of Bobkov and Götze [1999] and Barthe and Roberto [2003], to show part (i). As the conditions for the validity of the log-Sobolev inequality are *not* satisfied under the conditions of part (ii), a simple interpolation is used to show the desired result in that case. Part (iii) is proved by two estimates: The case where c_1 is too large follows an idea essentially already stated in [Jabin and Niethammer 2003], while the case where c_1 is too small seems to be a new result which we provide.

From Theorem 1.1 one can conclude in a straightforward way the following theorem, our main result on the rate of convergence to equilibrium:

Theorem 1.3. Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ satisfy Hypotheses 1–3 with $0 \leq \gamma \leq 1$, and let $\mathbf{c} = \mathbf{c}(t) = \{c_i(t)\}_{i \in \mathbb{N}}$ be a solution to the Becker–Döring equations with mass $\varrho \in (0, \varrho_s)$.

(i) **Rate for $a_i \sim i$.** If $\gamma = 1$ then there exists a constant $K > 0$ depending only on δ , ϱ and the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 2}$, and a constant $C > 0$ depending only on $H(\mathbf{c}(0)|Q)$, ϱ and the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 2}$ such that

$$H(\mathbf{c}(t)|Q) \leq C e^{-Kt} \quad \text{for } t \geq 0.$$

(ii) **Rate for $a_i \sim i^\gamma$, $\gamma < 1$.** If $\gamma < 1$ and $M_\beta(\mathbf{c}(0)) < +\infty$ for some $\beta \geq \max\{2 - \gamma, 1 + \gamma\}$ then there exists a constant $K > 0$ depending only on M_β , δ , ϱ and the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 2}$, and a constant $C > 0$ depending only on $H(\mathbf{c}(0)|Q)$, M_β , δ , ϱ and the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 2}$ such that

$$H(\mathbf{c}(t)|Q) \leq \frac{1}{\left(C + \frac{1-\gamma}{\beta-1} Kt\right)^{\frac{\beta-1}{1-\gamma}}} \quad \text{for } t \geq 0.$$

The constants K and C can be estimated explicitly.

In order to deduce [Theorem 1.3](#) we use the inequalities in [Theorem 1.1](#) when they are applicable. Of course, the assumption that $c_1(t)$ is in the “good” region given by (1-16) becomes eventually true, since $c_1(t)$ is known to converge to \bar{z} . More explicitly, one can apply the Csiszár–Kullback inequality (1-12) to obtain that for any $t > t_0$ we have

$$\bar{z} - H(c(t_0)|\mathcal{Q}) \leq c_1(t) \leq \bar{z} + H(c(t_0)|\mathcal{Q}), \quad t \geq t_0.$$

If $H(c(t_0)|\mathcal{Q})$ is small enough, this implies (1-16). For times t such that $c_1(t)$ is outside this “good” region, we use the inequality in [Theorem 1.1\(iii\)](#); details are given in [Section 4](#).

There are several improvements in these theorems with respect to the existing theory. One of them is that they apply to more general initial conditions, removing the need for a finite exponential moment present in [[Cañizo and Lods 2013](#); [Jabin and Niethammer 2003](#)]. Another one is that they answer the question of whether one can obtain a linear inequality such as (1-17) (i.e., whether the equivalent of Cercignani’s conjecture holds), making clear the link to discrete logarithmic Sobolev inequalities. It does hold in the case $a_i \sim i$, which is physically relevant, for example, in modelling polymer chains [[Farjoun and Neu 2011](#); [Neu et al. 2002](#)]. As a result, the statement for $a_i \sim i$ is quite strong: it gives full exponential convergence, with explicit constants in terms of the parameters, with no restriction on the initial data except that of subcritical mass. Point (ii) in [Theorem 1.3](#) also relaxes the requirements on the initial data, at the price of obtaining a slower convergence than that of [[Cañizo and Lods 2013](#)]; we do not know whether this rate is optimal for initial conditions with polynomially decaying tails (so that $M_\beta < \infty$ for some $\beta > 1$, but $M_{\beta'} = +\infty$ for some $\beta' > \beta$). Recently, Murray and Pego [[2015](#)] investigated this rate of convergence, concluding an algebraic rate of decay as well. It would be interesting to verify the optimality of this result by determining whether the corresponding linearised operator admits a spectral gap in ℓ^1 spaces with polynomial weights (in ℓ^1 spaces with exponential weights, the answer is positive and an estimate of the spectral gap can be found in [[Cañizo and Lods 2013](#)]). We believe that no such spectral gap exists for $0 \leq \gamma < 1$, i.e., that the algebraic rate of convergence is optimal even for close to equilibrium initial data.

One may wonder if the method presented here can be used to reach an inequality like Jabin and Niethammer’s (1-13) under the additional condition of an exponential moment. The answer is indeed positive:

Theorem 1.4. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ satisfy Hypotheses 1–3 with $0 \leq \gamma < 1$.*

- (i) **Functional inequality.** *Let $c = \{c_i\}_{i \in \mathbb{N}}$ be an arbitrary positive sequence with mass $\varrho \in (0, \varrho_s)$ for which there exists $\mu > 0$ such that*

$$M_\mu^{\text{exp}}(c) := \sum_{i=1}^{\infty} e^{\mu i} c_i < +\infty. \tag{1-21}$$

Then there exist $K_1, K_2, \varepsilon > 0$ depending only on $M_\mu^{\text{exp}}(c), \delta, \varrho$ and the coefficients $\{a_i\}_{i \geq 1}, \{b_i\}_{i \geq 2}$ such that

$$\bar{D}(c) \geq \min \left(\frac{K_1 H(c|\mathcal{Q})}{|\log(K_2 H(c|\mathcal{Q}))|^{1-\gamma}}, \varepsilon \right). \tag{1-22}$$

Moreover, K_1, K_2 and ε can be given explicitly.

(ii) **Rate of convergence.** If $\mathbf{c}(t) = \{c_i(t)\}_{i \in \mathbb{N}}$ is a solution to the Becker–Döring equations with mass $0 < \varrho < \varrho_s$ such that there exists $\mu > 0$ with

$$M_\mu^{\text{exp}}(\mathbf{c}(0)) := \sum_{i=1}^{\infty} e^{\mu i} c_i(0) < +\infty,$$

then there exists a constant $K > 0$ depending only on $M_\mu^{\text{exp}}(\mathbf{c}(0))$, δ , ϱ and the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 2}$, and a constant $C > 0$ depending only on $H(\mathbf{c}(0)|\mathcal{Q})$, $M_\mu^{\text{exp}}(\mathbf{c}(0))$, δ , ϱ and the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 2}$ such that

$$H(\mathbf{c}(t)|\mathcal{Q}) \leq C e^{-Kt^{1/(2-\gamma)}}.$$

Moreover, K and C can be given explicitly.

Typical coefficients. The above results are valid for coagulation and fragmentation coefficients satisfying Hypotheses 1 – 3. To motivate our choice of assumptions, we briefly recall here some physically motivated coagulation and fragmentation coefficients found in the literature.

Common model coefficients appearing in the theory of density-conserving phase transitions (see [Niethammer 2003; Penrose 1989]) are given by

$$a_i = i^\gamma, \quad b_i = a_i \left(z_s + \frac{q}{i^{1-\mu}} \right) \quad \text{for all } i \geq 1 \tag{1-23}$$

for some $0 < \gamma \leq 1$, $z_s > 0$, $q > 0$ and $0 < \mu < 1$. These coefficients may be derived from simple assumptions on the mechanism of the reactions taking place; we take particular values from [Niethammer 2003]:

$$\begin{aligned} \gamma = \frac{1}{3}, \quad \mu = \frac{2}{3} & \quad (\text{diffusion-limited kinetics in 3-D}), \\ \gamma = 0, \quad \mu = \frac{1}{2} & \quad (\text{diffusion-limited kinetics in 2-D}), \\ \gamma = \frac{2}{3}, \quad \mu = \frac{2}{3} & \quad (\text{interface-reaction-limited kinetics in 3-D}), \\ \gamma = \frac{1}{2}, \quad \mu = \frac{1}{2} & \quad (\text{interface-reaction-limited kinetics in 2-D}). \end{aligned} \tag{1-24}$$

The case $\gamma = 1$ appears, for example, in modelling polymer chains, where the binding energy increases by a constant each time a monomer is added.

A different kind of reasoning, based on a statistical mechanics argument involving the binding energy of clusters, results in the coefficients

$$a_i = i^\gamma, \quad b_i = z_s(i - 1)^\gamma \exp(\sigma i^\mu - \sigma(i - 1)^\mu), \quad i \in \mathbb{N}, \tag{1-25}$$

for appropriate constants γ, μ and where $\sigma > 0$ is related to the surface tension of the aggregates. The values of μ and γ for various situations are still those in (1-24).

As already mentioned, the choice $\gamma = 1$ corresponds to the physically relevant example in modelling polymer chains, for instance, for proteins aggregating in a cubic phase of lipid bilayers [Farjoun and Neu 2011; Neu et al. 2002].

The behaviour of (1-23) and (1-25) is similar: observe that for large i we have $i^\mu - (i-1)^\mu \sim \mu i^{\mu-1}$, so the fragmentation coefficients become roughly

$$b_i \sim z_s a_i \exp(\sigma \mu i^{\mu-1}) \sim a_i \left(z_s + \frac{z_s \sigma \mu}{i^{1-\mu}} \right),$$

which is like (1-23) with $q = z_s \sigma \mu$. Moreover, for both classes of coefficients, we can write (by the definition of Q_i)

$$Q_i = \frac{a_1 a_2 \cdots a_{i-1}}{b_2 b_3 \cdots b_i} = z_s^{1-i} \alpha_i, \quad (1-26)$$

where $\{\alpha_i\}_{i \in \mathbb{N}}$ is nonincreasing and satisfies

$$\lim_{i \rightarrow \infty} \frac{\alpha_{i+1}}{\alpha_i} = 1.$$

In other words, Hypotheses 1–3 hold true for both models.

Application to general coagulation and fragmentation models. The Becker–Döring equations are the simplest form of a coagulation and fragmentation process, assuming that the only relevant reactions are governed by monomers. Other models take into account the fact that clusters of size i and size j , for $i, j \in \mathbb{N}$, may interact. A discrete model — similar to the Becker–Döring equations (1-1) — can be formulated, now with coagulation and fragmentation coefficients of the form $a_{i,j}, b_{i,j}$ (see the subsection on page 1698). Together with an assumption of detailed balance, one can once again find equilibria to the process and inquire about the rate of convergence to them. Our study of the Becker–Döring equations allows us to give a quantitative answer (though not optimal) for this question. We leave the detailed description of the model we have in mind for the subsection on page 1698. For such a model, using the same notion of free relative energy we will show that:

Theorem 1.5 (asymptotic behaviour of the coagulation-fragmentation system). *Let $\{a_{i,j}\}_{i,j \in \mathbb{N}}$ and $\{b_{i,j}\}_{i,j \in \mathbb{N}}$ be the coagulation and fragmentation coefficients for equation (5-1), and assume that the detailed balance condition (5-6) holds. Assume that*

$$a_{i,j} = i^\gamma + j^\gamma \quad (1-27)$$

for some $0 \leq \gamma < 1$ and that $\{Q_i\}_{i \in \mathbb{N}}$ satisfies Hypothesis 2. Assume in addition that $M_k(c(0)) < +\infty$ for some $k \in \mathbb{N}$, $k > 1$. Then

$$H(c(t)|\mathcal{Q}) \leq \frac{1}{(C_1 + C_2 \log t)^{\frac{k-1}{1-\gamma}}}, \quad t \geq 0, \quad (1-28)$$

where $C_1, C_2 > 0$ are constants depending only on $H(c(0)|\mathcal{Q})$, z_s , ϱ , $\{\alpha_i\}_{i \in \mathbb{N}}$, k , γ and $M_k(c(0))$.

Organisation of the paper. The structure of the paper is as follows: In Section 2 we will present our main technical tool, a discrete version of the log-Sobolev inequality with weights. Section 3 contains the proof of Theorem 1.1 and uses Section 2 to show the first part of the theorem. We also show in this section that this method is optimal and that Cercignani’s conjecture cannot hold when $\gamma < 1$, proving Theorem 1.2,

and explore the additional inequality that appears under the assumption of a finite exponential moment. Section 4 deals with the consequences of our functional inequalities for the solutions to the Becker–Döring equation and contains the proof of Theorem 1.3 and part (ii) of Theorem 1.4. In Section 5 we provide the proof of Theorem 1.5 and remark on the difficulties of obtaining stronger results in this general setting. Lastly, we give two appendices where proofs to some technical lemmas can be found.

2. A discrete weighted logarithmic Sobolev inequality

One of the key ingredients in proving Cercignani’s conjecture for the Becker–Döring equations in terms of Theorem 1.1 is a discrete log-Sobolev inequality with weights. The theory presented here follows closely the work of Bobkov and Götze [1999], and that of Barthe and Roberto [2003], and can be seen as a discrete version of the aforementioned papers. It is worth noting that a discrete version is explicitly mentioned in [Barthe and Roberto 2003, Section 4], with a remark that the arguments in that paper can be adapted to prove it. Indeed, our proof is essentially an adaptation of the one in [Bobkov and Götze 1999], and we give it in this section for the sake of completeness (and since we have not been able to find an explicit proof in the discrete case). Some further technical details are postponed to Appendix A.

The main log-Sobolev inequality. We start with some basic definitions:

Definition 2.1. We say that $\mu \in P(\mathbb{N})$ if $\mu = \{\mu_i\}_{i \in \mathbb{N}}$ is a nonnegative sequence such that

$$\sum_{i=1}^{\infty} \mu_i = 1.$$

For any nonnegative sequence $g = \{g_i\}_{i \in \mathbb{N}}$ with $\sum_{i=1}^{\infty} \mu_i g_i < +\infty$, we define its *entropy* with respect to μ as

$$\text{Ent}_{\mu}(g) = \sum_{i=1}^{\infty} \mu_i g_i \log \frac{g_i}{\sum_{i=1}^{\infty} \mu_i g_i}. \tag{2-1}$$

Definition 2.2. Given $\mu \in P(\mathbb{N})$ and positive sequence $\nu = \{\nu_i\}_{i \in \mathbb{N}}$ (not necessarily normalised) we say that ν admits a log-Sobolev inequality with respect to μ with constant $0 < C_{\text{LS}} < +\infty$ if, for any sequence $f = \{f_i\}_{i \in \mathbb{N}}$,

$$\text{Ent}_{\mu}(f^2) \leq C_{\text{LS}} \sum_{i=1}^{\infty} \nu_i (f_{i+1} - f_i)^2, \tag{2-2}$$

where $f^2 = \{f_i^2\}_{i \in \mathbb{N}}$.

In what follows we will always assume that $\mu \in P(\mathbb{N})$. Setting

$$\Psi(x) = |x| \log(1 + |x|),$$

the main theorem, and its simplified corollary, that we will prove in this section are:

Theorem 2.3. *The following two conditions are equivalent:*

- (i) ν admits a log-Sobolev inequality with respect to μ with constant C_{LS} .

(ii) For any $m \in \mathbb{N}$ such that

$$\max\left(\sum_{i=1}^{m-1} \mu_i, \sum_{i=m+1}^{\infty} \mu_i\right) < \frac{2}{3},$$

we have

$$B_1 = \sup_{k \geq m} \frac{\sum_{i=1}^k 1/v_i}{\Psi^{-1}(1/\sum_{i=k+1}^{\infty} \mu_i)} < +\infty. \tag{2-3}$$

Moreover, if (ii) is valid then one can choose

$$C_{LS} = 40(B_2 + 4B_1), \quad \text{where } B_2 = \frac{\sum_{i=1}^{m-1} 1/v_i}{\Psi^{-1}(1/\sum_{i=1}^{m-1} \mu_i)}. \tag{2-4}$$

A somehow more tractable consequence is the following.

Corollary 2.4. *The following two conditions are equivalent:*

- (i) ν admits a log-Sobolev inequality with respect to μ with constant C_{LS} .
- (ii) For any $m \in \mathbb{N}$ such that

$$\max\left(\sum_{i=1}^{m-1} \mu_i, \sum_{i=m+1}^{\infty} \mu_i\right) < \frac{2}{3},$$

we have

$$D_1 = \sup_{k \geq m} \left(-\sum_{i=k+1}^{\infty} \mu_i \log\left(\sum_{i=k+1}^{\infty} \mu_i\right)\right) \left(\sum_{i=1}^k \frac{1}{v_i}\right) < \infty. \tag{2-5}$$

Moreover, if (ii) is valid then one can choose

$$C_{LS} = 120(D_2 + 4D_1), \tag{2-6}$$

where $D_2 = (-\sum_{i=1}^{m-1} \mu_i \log(\sum_{i=1}^{m-1} \mu_i))(\sum_{i=1}^{m-1} 1/v_i)$.

Remark 2.5. One can clearly see that if

$$\sup_{k \geq 1} \left(-\sum_{i=k+1}^{\infty} \mu_i \log\left(\sum_{i=k+1}^{\infty} \mu_i\right)\right) \left(\sum_{i=1}^k \frac{1}{v_i}\right) < \infty$$

then one has a log-Sobolev inequality of ν with respect to μ . However, the introduction of the “approximate median” m allows us to have an explicit estimation on the log-Sobolev constant C_{LS} .

The rest of the Section is dedicated to the proof of the above results and will be divided in various steps — each one corresponding to a subsection.

A reformulation as a Poincaré inequality in Orlicz spaces. As in the work of Bobkov and Götze [1999], a key argument in the proofs of Theorem 2.3 and Corollary 2.4 is to recast the log-Sobolev inequality as a Poincaré inequality in the Orlicz space associated to Ψ . We start with the definition:

Definition 2.6. Given $\mu \in P(\mathbb{N})$ and a Young function, $\Sigma : [0, +\infty) \rightarrow [0, +\infty)$, i.e., a convex function such that

$$\frac{\Sigma(x)}{x} \xrightarrow{x \rightarrow +\infty} +\infty, \quad \frac{\Sigma(x)}{x} \xrightarrow{x \rightarrow 0} 0,$$

we define the Orlicz space $L_{\Sigma}^{(\mu)}$ as the space of all sequences f such that there exists $k > 0$ with

$$\sum_{i=1}^{\infty} \mu_i \Sigma\left(\frac{|f_i|}{k}\right) < \infty.$$

In that case we define

$$\|f\|_{L_{\Sigma}^{(\mu)}} = \inf_{k>0} \left\{ \sum_{i=1}^{\infty} \mu_i \Sigma\left(\frac{|f_i|}{k}\right) \leq 1 \right\}.$$

In what follows we will drop the superscript μ from the Orlicz space of Ψ and its norm. Additionally we set $\Phi(x) = \Psi(x^2)$ and notice that

$$\|f^2\|_{L_{\Psi}} = \inf_{k>0} \left\{ \sum_{i=1}^{\infty} \mu_i \Psi\left(\frac{f^2}{k}\right) \leq 1 \right\} = \left(\inf_{\sqrt{k}>0} \left\{ \sum_{i=1}^{\infty} \mu_i \Phi\left(\frac{|f|}{\sqrt{k}}\right) \leq 1 \right\} \right)^2 = \|f\|_{L_{\Phi}}^2. \tag{2-7}$$

We have then the following version of Rothaus’s lemma:

Lemma 2.7. Given $\mu \in P(\mathbb{N})$ and a sequence $f = \{f_i\}_{i \in \mathbb{N}}$, we set

$$\mathcal{L}(f) = \sup_{\alpha \in \mathbb{R}} \text{Ent}_{\mu}((f + \alpha)^2), \tag{2-8}$$

where $f + \alpha = \{f_i + \alpha\}_{i \in \mathbb{N}}$. Then,

$$\text{Ent}_{\mu}(f^2) \leq \mathcal{L}(f) \leq \text{Ent}_{\mu}(f^2) + 2 \sum_{i=1}^{\infty} \mu_i f_i^2. \tag{2-9}$$

Remark 2.8. This lemma is an adaptation of the appropriate lemma in [Rothaus 1985, Lemma 9]. We leave the proof of it to Appendix A.

We have then the following equivalent formulation of the log-Sobolev inequality:

Proposition 2.9. The following conditions are equivalent:

- (i) ν admits a log-Sobolev inequality with respect to μ with constant C_{LS} .
- (ii) For any sequence f ,

$$\mathcal{L}(f) \leq C_{\text{LS}} \sum_{i=1}^{\infty} \nu_i (f_{i+1} - f_i)^2. \tag{2-10}$$

(iii) For any sequence \mathbf{f} ,

$$\|\mathbf{f} - \langle \mathbf{f} \rangle\|_{L_\Phi}^2 \leq \lambda \sum_{i=1}^{\infty} \nu_i (f_{i+1} - f_i)^2, \quad (2-11)$$

where $\langle \mathbf{f} \rangle = \sum_{i=1}^{\infty} \mu_i f_i$.

Moreover, if (i) or (ii) are valid, one can choose $\lambda = \frac{3}{2}C_{\text{LS}}$. If (iii) is valid one can choose $C_{\text{LS}} = 5\lambda$.

The proof of the proposition relies on the following lemma:

Lemma 2.10. For any sequence \mathbf{f} , one has

$$\frac{2}{3}\|\mathbf{f} - \langle \mathbf{f} \rangle\|_{L_\Phi}^2 \leq \mathcal{L}(\mathbf{f}) \leq 5\|\mathbf{f} - \langle \mathbf{f} \rangle\|_{L_\Phi}^2. \quad (2-12)$$

Proof. We start by noticing that we may assume $\langle \mathbf{f} \rangle = 0$, as well as $\|\mathbf{f} - \langle \mathbf{f} \rangle\|_{L_\Phi} = 1$. This is true as \mathcal{L} is invariant under translations and

$$\text{Ent}_\mu(\alpha \mathbf{f}) = \alpha \text{Ent}_\mu(\mathbf{f}).$$

Using Lemma 2.7, we find that

$$\begin{aligned} \mathcal{L}(\mathbf{f}) &\leq \text{Ent}_\mu(\mathbf{f}^2) + 2 \sum_{i=1}^{\infty} \mu_i f_i^2 \\ &= \sum_{i=1}^{\infty} \mu_i f_i^2 \log(f_i^2) + 2 \sum_{i=1}^{\infty} \mu_i f_i^2 - \left(\sum_{i=1}^{\infty} \mu_i f_i^2 \right) \log \left(\sum_{i=1}^{\infty} \mu_i f_i^2 \right) \\ &\leq \sum_{i=1}^{\infty} \mu_i \Phi(f_i) + h \left(\sum_{i=1}^{\infty} \mu_i f_i^2 \right), \end{aligned}$$

where $h(x) = 2x - x \log x$ for $x > 0$. As h is an increasing function on $[0, e]$ and

$$\|\mathbf{f}\|_{L_\mu^1} \leq \|\mathbf{f}\|_{L_\mu^2} \leq \sqrt{\frac{3}{2}}\|\mathbf{f}\|_{L_\Phi},$$

(see Lemma A.2 in Appendix A) we have

$$\|\mathbf{f}\|_{L_\mu^2}^2 \leq 2.$$

Thus, as

$$\sum_{i=1}^{\infty} \mu_i \Phi(f_i) = \sum_{i=1}^{\infty} \mu_i \Phi \left(\frac{f_i}{\|\mathbf{f}\|_{L_\Phi}} \right) \leq 1,$$

we find that

$$\mathcal{L}(\mathbf{f}) \leq 1 + h(2) \leq 5,$$

proving the right-hand side inequality of (2-12). To show the left-hand side of the inequality we assume that $\mathcal{L}(\mathbf{f}) = 2$. By the definition of \mathcal{L} and the fact that

$$\|\mathbf{f} - \langle \mathbf{f} \rangle\|_{L_\mu^2}^2 = \frac{1}{2} \lim_{|a| \rightarrow \infty} \text{Ent}_\mu((\mathbf{f} + a)^2)$$

(see Lemma A.3 in Appendix A), we know that

$$\|f\|_{L^2_\mu}^2 \leq \frac{1}{2} \mathcal{L}(f) = 1.$$

This implies

$$\sum_{i=1}^\infty \mu_i \Phi(f_i) \leq 1 + \sum_{i=1}^\infty \mu_i f_i^2 \log f_i^2 = 1 + \text{Ent}_\mu(f^2) + \|f\|_{L^2_\mu}^2 \log(\|f\|_{L^2_\mu}^2) \leq 1 + \mathcal{L}(f) = 3,$$

where we have used the fact that $x \log(1+x) \leq 1+x \log x$ when $x > 0$.

Since, for any $a \geq 1$,

$$\Phi\left(\frac{x}{\sqrt{a}}\right) = \frac{x^2}{a^2} \log\left(1 + \frac{x^2}{a^2}\right) \leq \frac{1}{a^2} \Phi(x),$$

the above implies

$$\sum_{i=1}^\infty \mu_i \Phi\left(\frac{f_i}{\sqrt{3}}\right) \leq 1$$

and as such, by the definition of $\|\cdot\|_{L_\Phi}$, we conclude that

$$\|f\|_{L_\Phi}^2 \leq 3 = \frac{3}{2} \mathcal{L}(f),$$

and the proof is complete. □

Proof of Proposition 2.9. The equivalence of (ii) and (iii) is immediate following Lemma 2.10, which also proves the desired connection between C_{LS} and λ . To show that (i) implies (ii) we notice that as the right-hand side of (2-2) is invariant under translation, taking the supremum over all possible translations results in (ii). The fact that (ii) implies (i) is immediate as $\text{Ent}_\mu(f^2) \leq \mathcal{L}(f)$. □

Discrete Hardy inequalities. The above observation that the log-Sobolev inequality with weights is actually a form of a Poincaré inequality brings to mind another inequality with weights that is closely connected to the Poincaré inequality — the Hardy inequality. In its discrete form, we have:

Lemma 2.11. *Let μ and ν be two sequences of positive numbers and let $m \in \mathbb{N}$. Then, the following two conditions are equivalent:*

(i) *There exists a finite constant $A_{1,m} \geq 0$ such that*

$$\sum_{i=m}^\infty \mu_i \left(\sum_{j=m}^i f_j\right)^2 \leq A_{1,m} \sum_{i=m}^\infty \nu_i f_i^2$$

for any sequence f .

(ii) *We have*

$$B_{1,m} = \sup_{k \geq m} \left(\sum_{i=k}^\infty \mu_i\right) \left(\sum_{i=m}^k \frac{1}{\nu_i}\right) < \infty.$$

Moreover, if any of the conditions holds then $B_{1,m} \leq A_{1,m} \leq 4B_{1,m}$.

The proof for the case $m = 1$ can be found in [Cañizo and Lods 2013], and the general case follows by the same method of proof.

Corollary 2.12. *Let*

$$B_m^{(1)} = \sup_{k \geq m} \left(\sum_{i=k+1}^{\infty} \mu_i \right) \left(\sum_{i=m}^k \frac{1}{v_i} \right).$$

Then for any sequence f such that $f_m = 0$, we have

$$\sum_{i=m}^{\infty} \mu_i f_i^2 \leq A_m^{(1)} \sum_{i=m}^{\infty} v_i (f_{i+1} - f_i)^2 \quad (2-13)$$

if and only if $B_m^{(1)} < \infty$. In that case $B_m^{(1)} \leq A_m^{(1)} \leq 4B_m^{(1)}$. Additionally,

$$B_{1,m} \leq B_m^{(1)} \leq B_{1,m+1}.$$

Proof. This follows immediately from Lemma 2.11 applied to the sequence $g_i = f_{i+1} - f_i$ and a simple translation argument. \square

Besides the above, we will also need to have a Hardy-type inequality for sums up to a fixed integer m .

Lemma 2.13. *Let μ and v be two sequences of positive numbers and let $m \in \mathbb{N}$. Then, for any sequence f such that $f_m = 0$, we have that if there exists $A > 0$ such that*

$$\sum_{i=1}^{m-1} \mu_i f_i^2 \leq A \sum_{i=1}^{m-1} v_i (f_{i+1} - f_i)^2, \quad (2-14)$$

then $b_{2,m} \leq A$, where

$$b_{2,m} = \sup_{k \leq m-1} \sum_{i=1}^k \mu_i \left(\sum_{j=k}^{m-1} \frac{1}{v_j} \right).$$

Moreover, one can always choose

$$A = B_{2,m} = \sum_{i=1}^{m-1} \mu_i \left(\sum_{j=i}^{m-1} \frac{1}{v_j} \right).$$

Proof. We start by noticing that for any $1 \leq i \leq m-1$, we have

$$f_i^2 = \left(\sum_{j=i}^{m-1} (f_{j+1} - f_j) \right)^2 \leq \left(\sum_{j=i}^{m-1} \frac{1}{v_j} \right) \left(\sum_{j=i}^{m-1} v_j (f_{j+1} - f_j)^2 \right) \leq \left(\sum_{j=i}^{m-1} \frac{1}{v_j} \right) \left(\sum_{j=1}^{m-1} v_j (f_{j+1} - f_j)^2 \right).$$

Thus

$$\sum_{i=1}^{m-1} \mu_i f_i^2 \leq \left(\sum_{i=1}^{m-1} \mu_i \left(\sum_{j=i}^{m-1} \frac{1}{v_j} \right) \right) \left(\sum_{j=1}^{m-1} v_j (f_{j+1} - f_j)^2 \right) = B_{2,m} \sum_{j=1}^{m-1} v_j (f_{j+1} - f_j)^2,$$

completing the second statement. Next, for any $j \leq m - 1$ we set

$$\sigma_j = \sum_{i=j}^{m-1} \frac{1}{v_i}.$$

Fix $k \leq m - 1$ and define $f^{(k)}$ to be such that $f_i^{(k)} = \sigma_k$ when $i \leq k$ and $f_i^{(k)} = \sigma_i$ when $i > k$. We have

$$\sum_{i=1}^{m-1} v_i (f_{i+1}^{(k)} - f_i^{(k)})^2 = \sum_{i=k}^{m-1} v_i (f_{i+1}^{(k)} - f_i^{(k)})^2 = \sum_{i=k}^{m-1} \frac{1}{v_i} = \sigma_k.$$

On the other hand,

$$\sum_{i=1}^{m-1} \mu_i (f_i^{(k)})^2 \geq \sum_{i=1}^k \mu_i (f_i^{(k)})^2 = \sigma_k^2 \left(\sum_{i=1}^k \mu_i \right).$$

As (2-14) is valid, we see that

$$A \geq \left(\sum_{i=k}^{m-1} \frac{1}{v_i} \right) \left(\sum_{i=1}^k \mu_i \right)$$

for all k . □

Proof of the main inequality. The last ingredient we need in order to prove [Theorem 2.3](#) is the following lemma:

Lemma 2.14. *The following conditions are equivalent:*

- (i) \mathbf{v} admits a log-Sobolev inequality with respect to $\boldsymbol{\mu}$ with constant C_{LS} .
- (ii) There exists $\eta > 0$ such that, for any sequence $\mathbf{f} = \{f_i\}$ such that $f_m = 0$ with $m \in \mathbb{N}$ satisfying

$$\max \left(\sum_{i=1}^{m-1} \mu_i, \sum_{i=m+1}^{\infty} \mu_i \right) < \frac{2}{3},$$

we have

$$\|(\mathbf{f}^{(0)})^2\|_{L_\Psi} + \|(\mathbf{f}^{(1)})^2\|_{L_\Psi} \leq \eta \sum_{i=1}^{\infty} v_i (f_{i+1} - f_i)^2,$$

where $\mathbf{f}^{(0)} = \mathbf{f} \mathbb{1}_{i < m}$ and $\mathbf{f}^{(1)} = \mathbf{f} \mathbb{1}_{i > m}$.

Moreover, if condition (ii) is valid, one can choose $C_{LS} = 40\eta$.

Proof. We notice that it is enough for us to show the equivalence of condition (ii) of our lemma and [Proposition 2.9\(ii\)](#).

Assume, to begin with, that [Proposition 2.9\(ii\)](#) is valid. As was shown in that proposition, this implies

$$\|\mathbf{f} - \langle \mathbf{f} \rangle\|_{L_\Phi}^2 \leq \frac{3C_{LS}}{2} \sum_{i=1}^{\infty} v_i (f_{i+1} - f_i)^2. \tag{2-15}$$

Due to the conditions on f and the definitions of $f^{(0)}$ and $f^{(1)}$, one has that

$$\begin{aligned}\|\langle f^{(0)} \rangle\|_{L_\Phi} &\leq |\langle f^{(0)} \rangle| \leq \|f^{(0)}\|_{L_\mu^2} \left(\sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}}, \\ \|\langle f^{(1)} \rangle\|_{L_\Phi} &\leq |\langle f^{(1)} \rangle| \leq \|f^{(1)}\|_{L_\mu^2} \left(\sum_{i=m+1}^{\infty} \mu_i \right)^{\frac{1}{2}}\end{aligned}$$

(see Lemma A.4 in Appendix A). Thus

$$\|f^{(0)}\|_{L_\Phi} \leq \|f^{(0)} - \langle f^{(0)} \rangle\|_{L_\Phi} + \|\langle f^{(0)} \rangle\|_{L_\Phi} \leq \|f^{(0)} - \langle f^{(0)} \rangle\|_{L_\Phi} + \left(\frac{3}{2} \sum_{i=1}^{m-1} \mu_i \right) \|f^{(0)}\|_{L_\Phi},$$

implying

$$\|f^{(0)}\|_{L_\Phi} \leq \frac{1}{1 - \sqrt{\frac{3}{2} \sum_{i=1}^{m-1} \mu_i}} \|f^{(0)} - \langle f^{(0)} \rangle\|_{L_\Phi},$$

and similarly

$$\|f^{(1)}\|_{L_\Phi} \leq \frac{1}{1 - \sqrt{\frac{3}{2} \sum_{i=m+1}^{\infty} \mu_i}} \|f^{(1)} - \langle f^{(1)} \rangle\|_{L_\Phi}.$$

We can conclude, by applying (2-15) to $f^{(0)}$ and $f^{(1)}$, that

$$\begin{aligned}\|f^{(0)}\|_{L_\Phi}^2 &\leq \frac{3C_{LS}}{2(1 - \sqrt{\frac{3}{2} \sum_{i=1}^{m-1} \mu_i})^2} \sum_{i=1}^{m-1} v_i (f_{i+1} - f_i)^2, \\ \|f^{(1)}\|_{L_\Phi}^2 &\leq \frac{3C_{LS}}{2(1 - \sqrt{\frac{3}{2} \sum_{i=m+1}^{\infty} \mu_i})^2} \sum_{i=m}^{\infty} v_i (f_{i+1} - f_i)^2.\end{aligned}$$

The result now follows from (2-7).

To show the converse, we use the translation invariance of Proposition 2.9(ii) to assume that $f_m = 0$. As such we have $f = f^{(0)} + f^{(1)}$. Moreover,

$$\begin{aligned}\|f - \langle f \rangle\|_{L_\Phi}^2 &\leq (\|f^{(0)} - \langle f^{(0)} \rangle\|_{L_\Phi} + \|f^{(1)} - \langle f^{(1)} \rangle\|_{L_\Phi})^2 \\ &\leq \left(\left(1 + \left(\frac{3}{2} \sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}} \right) \|f^{(0)}\|_{L_\Phi} + \left(1 + \left(\frac{3}{2} \sum_{i=m+1}^{\infty} \mu_i \right)^{\frac{1}{2}} \right) \|f^{(1)}\|_{L_\Phi} \right)^2 \\ &\leq 2 \left(1 + \left(\frac{3}{2} \sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}} \right)^2 \|f^{(0)}\|_{L_\Phi}^2 + 2 \left(1 + \left(\frac{3}{2} \sum_{i=m+1}^{\infty} \mu_i \right)^{\frac{1}{2}} \right)^2 \|f^{(1)}\|_{L_\Phi}^2 \\ &\leq 2\eta \max \left(\left(1 + \left(\frac{3}{2} \sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}} \right)^2, \left(1 + \left(\frac{3}{2} \sum_{i=m+1}^{\infty} \mu_i \right)^{\frac{1}{2}} \right)^2 \right) \sum_{i=1}^{\infty} v_i (f_{i+1} - f_i)^2,\end{aligned}$$

where we again used (2-7). This shows the desired result due to Proposition 2.9. \square

Proof of Theorem 2.3. Our main tool will be [Lemma 2.14](#). It is known that

$$\|f^2\|_{L_\Psi} = \sup \left\{ \sum_{i=1}^\infty \mu_i f_i^2 g_i : \sum_{i=1}^\infty \mu_i \Xi(g_i) \leq 1 \right\},$$

where Ξ is the Young complement of Ψ . Using [Corollary 2.12](#), we know that if $f_m = 0$ then

$$\sum_{i=m}^\infty \mu_i f_i^2 g_i \leq C_{LS} \sum_{i=m}^\infty v_i (f_{i+1} - f_i)^2$$

if and only if

$$B = \sup_{k \geq m} \left(\sum_{i=k+1}^\infty g_i \mu_i \right) \left(\sum_{i=1}^k \frac{1}{v_i} \right) < \infty.$$

Taking supremum over all appropriate $g = \{g_i\}$, we find that

$$\|f^2 \mathbb{1}_{i>m}\|_{L_\Psi} \leq C_{LS} \sum_{i=m}^\infty v_i (f_{i+1} - f_i)^2 \tag{2-16}$$

if and only if

$$B = \sup_{k \geq m} \|\mathbb{1}_{[k+1, \infty)}\|_{L_\Psi} \sum_{i=1}^k \frac{1}{v_i} < \infty.$$

As

$$\begin{aligned} \|\mathbb{1}_{[k+1, \infty)}\|_{L_\Psi} &= \inf_{\alpha > 0} \left\{ \sum_{i=k+1}^\infty \mu_i \Psi\left(\frac{1}{\alpha}\right) \leq 1 \right\} = \inf_{\alpha > 0} \left\{ \Psi\left(\frac{1}{\alpha}\right) \leq \frac{1}{\sum_{i=k+1}^\infty \mu_i} \right\} \\ &= \frac{1}{\Psi^{-1}\left(1/\sum_{i=k+1}^\infty \mu_i\right)}, \end{aligned}$$

we find that (2-16) is equivalent to $B_1 < \infty$, showing that (i) implies (ii).

Conversely, using [Lemma 2.13](#) we find that if $f_m = 0$ then

$$\begin{aligned} \sum_{i=1}^{m-1} \mu_i f_i^2 g_i &\leq \left[\sum_{i=1}^{m-1} \mu_i g_i \left(\sum_{j=i}^{m-1} \frac{1}{v_j} \right) \right] \sum_{i=1}^{m-1} v_i (f_{i+1} - f_i)^2 \\ &\leq \left[\left(\sum_{i=1}^{m-1} \mu_i g_i \right) \left(\sum_{j=1}^{m-1} \frac{1}{v_j} \right) \right] \sum_{i=1}^{m-1} v_i (f_{i+1} - f_i)^2 \end{aligned}$$

and again, by taking the supremum over the appropriate g , we find that

$$\|f^2 \mathbb{1}_{i<m}\|_{L_\Psi} \leq B_2 \sum_{i=1}^{m-1} v_i (f_{i+1} - f_i)^2. \tag{2-17}$$

Thus, if $\mathbf{f} = \{f_i\}$ is a sequence such that $f_m = 0$, and if in addition $B_1 < \infty$, we have

$$\begin{aligned} \|(\mathbf{f}^{(0)})^2\|_{L_\Psi} + \|(\mathbf{f}^{(1)})^2\|_{L_\Psi} &\leq B_2 \sum_{i=1}^{m-1} v_i (f_{i+1} - f_i)^2 + 4B_1 \sum_{i=m}^{\infty} v_i (f_{i+1} - f_i)^2 \\ &\leq (B_2 + 4B_1) \sum_{i=1}^{\infty} v_i (f_{i+1} - f_i)^2, \end{aligned}$$

where we have used [Corollary 2.12](#). We conclude, using [Lemma 2.14](#), that if $B_1 < \infty$, then ν admits a log-Sobolev inequality with respect to μ with constant C_{LS} that can be chosen to be $C_{\text{LS}} = 40(B_1 + 4B_2)$. \square

We are only left with the proof of [Corollary 2.4](#). The proof relies on the following technical lemma, whose proof is left to [Appendix A](#):

Lemma 2.15. *For any $t \geq \frac{3}{2}$, one has*

$$\frac{1}{3} \frac{t}{\log t} \leq \Psi^{-1}(t) \leq 2 \frac{t}{\log t}.$$

Proof of [Corollary 2.4](#). Due to the choice of m and [Lemma 2.15](#), we know that $\Psi^{-1}(t)$ and $t/\log t$ are equivalent for our choice of

$$t = \frac{1}{\sum_{i=m+1}^{\infty} \mu_i}.$$

This shows the desired equivalence using [Theorem 2.3](#). As for the last estimation, it follows immediately from the fact that

$$B_i \leq 3D_i$$

for $i = 1, 2$. \square

Now that we have achieved a necessary and sufficient condition for the validity of a discrete log-Sobolev inequality with weight, we will proceed to see how it can be used to prove [Theorem 1.1](#).

3. Energy dissipation inequalities

The log-Sobolev inequality and the Becker–Döring equations. Motivated by our previous section, the first step in trying to show the validity of Cercignani’s conjecture would be to relate the energy dissipation, $D(\mathbf{c})$, and a term that resembles the right-hand side of (2-2). Recall that, for any nonnegative sequence $\mathbf{c} = \{c_i\}$ we defined

$$D(\mathbf{c}) = \sum_{i=1}^{\infty} a_i Q_i \Theta\left(\frac{c_1 c_i}{Q_i}, \frac{c_{i+1}}{Q_{i+1}}\right)$$

with $\Theta(x, y) := (x - y)(\log x - \log y)$, and

$$\bar{D}(\mathbf{c}) = \sum_{i=1}^{\infty} a_i Q_i \left(\sqrt{\frac{c_1 c_i}{Q_i}} - \sqrt{\frac{c_{i+1}}{Q_{i+1}}} \right)^2.$$

We have the following properties:

Lemma 3.1. *For any nonnegative sequence \mathbf{c} , the following holds:*

(i) *We have*

$$4\bar{D}(\mathbf{c}) \leq D(\mathbf{c}). \tag{3-1}$$

(ii) *For any $z > 0$, we can rewrite $D(\mathbf{c})$ as*

$$D(\mathbf{c}) = \sum_{i=1}^{\infty} a_i Q_i z^{i+1} \Theta\left(\frac{c_1 c_i}{Q_i z^{i+1}}, \frac{c_{i+1}}{Q_{i+1} z^{i+1}}\right), \tag{3-2}$$

recalling $\Theta(x, y) := (x - y)(\log x - \log y)$, and

$$\bar{D}(\mathbf{c}) = \sum_{i=1}^{\infty} a_i Q_i z^{i+1} \left(\sqrt{\frac{c_1 c_i}{Q_i z^{i+1}}} - \sqrt{\frac{c_{i+1}}{Q_{i+1} z^{i+1}}}\right)^2. \tag{3-3}$$

Proof. Part (i) is an immediate consequence of the inequality

$$\Theta(x, y) = (x - y)(\log x - \log y) \geq 4(\sqrt{x} - \sqrt{y})^2$$

and (ii) is immediate from the homogeneity of the expressions involved. □

Property (ii) of the above lemma gives an indication of how we may be able to find a connection between $\bar{D}(\mathbf{c})$ and the relative entropy between \mathbf{c} and *some* equilibrium, by appropriately choosing z . Similar to the work of Jabin and Niethammer [2003], another equilibrium state that will play an important role in what is to follow is

$$\tilde{Q} = Q_{c_1} = \{Q_i c_1^i\}_{i \geq 1}.$$

Indeed, it is the *only* possible equilibrium under which the right-hand side of (3-3) attains a form that is suitable for the log-Sobolev theory developed in the previous section. From (3-3) we find, after cancelling c_1 , that

$$\bar{D}(\mathbf{c}) = \sum_{i=1}^{\infty} a_i \tilde{Q}_i \tilde{Q}_1 \left(\sqrt{\frac{c_i}{\tilde{Q}_i}} - \sqrt{\frac{c_{i+1}}{\tilde{Q}_{i+1}}}\right)^2. \tag{3-4}$$

This enables us to finally link $\bar{D}(\mathbf{c})$ to $H(\mathbf{c}|\mathcal{Q})$:

Proposition 3.2. *For given coagulation and detailed balance coefficients $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ and a given positive sequence \mathbf{c} with finite mass ϱ and such that*

$$\sum_{i=1}^{\infty} \tilde{Q}_i < +\infty, \quad \sum_{i=1}^{\infty} a_i \tilde{Q}_i < +\infty$$

(recall $Q_i := Q_i c_1^i$ for $i \geq 1$), we define the measures

$$\mu_i = \frac{\tilde{Q}_i}{\sum_{i=1}^{\infty} \tilde{Q}_i}, \quad \nu_i := \frac{a_i \tilde{Q}_i}{\sum_{j=1}^{\infty} a_j \tilde{Q}_j}, \quad i \in \mathbb{N}. \tag{3-5}$$

Then, if \mathbf{v} admits a log-Sobolev inequality with respect to μ with constant C_{LS} , we have

$$\bar{D}(\mathbf{c}) \geq \frac{c_1^3 (\sum_{i=1}^\infty a_i \tilde{Q}_i)}{C_{LS} (\sum_{i=1}^\infty \tilde{Q}_i) (c_1^2 + 2(\sum_{i=1}^\infty c_i) (\sum_{i=1}^\infty \tilde{Q}_i))} H(\mathbf{c}|\mathcal{Q}). \tag{3-6}$$

Proof. Let $f_i = \sqrt{c_i/\tilde{Q}_i}$. Since \mathbf{v} admits a log-Sobolev inequality with respect to μ with constant C_{LS} , we have

$$\bar{D}(\mathbf{c}) = \left(\sum_{i=1}^\infty a_i \tilde{Q}_i \tilde{Q}_1 \right) \sum_{i=1}^\infty v_i (f_{i+1} - f_i)^2 \geq \frac{c_1 (\sum_{i=1}^\infty a_i \tilde{Q}_i)}{C_{LS}} \text{Ent}_\mu(\mathbf{f}^2). \tag{3-7}$$

Next, we notice that

$$\begin{aligned} \left(\sum_{i=1}^\infty \tilde{Q}_i \right) \text{Ent}_\mu(\mathbf{f}^2) &= \sum_{i=1}^\infty c_i \log \frac{c_i}{\tilde{Q}_i} - \left(\sum_{i=1}^\infty c_i \right) \left(\log \sum_{i=1}^\infty c_i - \log \sum_{i=1}^\infty \tilde{Q}_i \right) \\ &= H(\mathbf{c}|\tilde{\mathcal{Q}}) + \sum_{i=1}^\infty c_i - \sum_{i=1}^\infty \tilde{Q}_i - \left(\sum_{i=1}^\infty c_i \right) \left(\log \sum_{i=1}^\infty c_i - \log \sum_{i=1}^\infty \tilde{Q}_i \right) \\ &= H(\mathbf{c}|\tilde{\mathcal{Q}}) - \left(\sum_{i=1}^\infty \tilde{Q}_i \right) \Lambda \left(\frac{\sum_{i=1}^\infty c_i}{\sum_{i=1}^\infty \tilde{Q}_i} \right), \end{aligned} \tag{3-8}$$

where $\Lambda(x) = x \log x - x + 1$. We now use the fact that \mathcal{Q} minimises the relative entropy to the set of equilibria to bound the first term,

$$H(\mathbf{c}|\tilde{\mathcal{Q}}) \geq H(\mathbf{c}|\mathcal{Q}) \tag{3-9}$$

(see Lemma B.1 in Appendix B). The only remaining bound is to show that the term with the negative sign at the end of (3-8) is in fact bounded by $\text{Ent}_\mu(\mathbf{f}^2)$. For this we will use the following Csiszár–Kullback inequality:

$$\text{Ent}_\mu(\mathbf{f}^2) \geq \frac{1}{2\langle \mathbf{f}^2 \rangle} \left(\sum_{i=1}^\infty |f_i^2 - \langle \mathbf{f}^2 \rangle| \mu_i \right)^2, \tag{3-10}$$

where

$$\langle \mathbf{f}^2 \rangle := \sum_{i=1}^\infty f_i^2 \mu_i.$$

With (3-10), we find that in our particular setting

$$\text{Ent}_\mu(\mathbf{f}^2) \geq \frac{\sum_{i=1}^\infty \tilde{Q}_i}{2 \sum_{i=1}^\infty c_i} \left(\sum_{i=1}^\infty \left| \frac{c_i}{\sum_{i=1}^\infty \tilde{Q}_i} - \frac{\tilde{Q}_i (\sum_{i=1}^\infty c_i)}{(\sum_{i=1}^\infty \tilde{Q}_i)^2} \right| \right)^2 = \frac{\sum_{i=1}^\infty c_i}{2 \sum_{i=1}^\infty \tilde{Q}_i} \left(\sum_{i=1}^\infty \left| \frac{c_i}{\sum_{i=1}^\infty c_i} - \frac{\tilde{Q}_i}{\sum_{i=1}^\infty \tilde{Q}_i} \right| \right)^2,$$

and keeping only the first term in the last sum we get

$$\text{Ent}_\mu(\mathbf{f}^2) \geq \frac{\sum_{i=1}^\infty c_i}{2 \sum_{i=1}^\infty \tilde{Q}_i} \left| \frac{c_1}{\sum_{i=1}^\infty c_i} - \frac{\tilde{Q}_1}{\sum_{i=1}^\infty \tilde{Q}_i} \right|^2 = \frac{c_1^2}{2 \sum_{i=1}^\infty c_i \sum_{i=1}^\infty \tilde{Q}_i} \left(1 - \frac{\sum_{i=1}^\infty c_i}{\sum_{i=1}^\infty \tilde{Q}_i} \right)^2.$$

Continuing from (3-8) and using (3-9), the above inequality and the fact that

$$\Lambda(x) \leq (x - 1)^2$$

show that

$$\begin{aligned} \left(\sum_{i=1}^{\infty} \tilde{Q}_i\right) \text{Ent}_{\mu}(f^2) &\geq H(c|\mathcal{Q}) - \left(\sum_{i=1}^{\infty} \tilde{Q}_i\right) \left(\frac{\sum_{i=1}^{\infty} c_i}{\sum_{i=1}^{\infty} \tilde{Q}_i} - 1\right)^2 \\ &\geq H(c|\mathcal{Q}) - \frac{2}{c_1^2} \left(\sum_{i=1}^{\infty} \tilde{Q}_i\right)^2 \left(\sum_{i=1}^{\infty} c_i\right) \text{Ent}_{\mu}(f^2). \end{aligned}$$

Thus,

$$H(c|\mathcal{Q}) \leq \left(\sum_{i=1}^{\infty} \tilde{Q}_i\right) \left(1 + \frac{2}{c_1^2} \left(\sum_{i=1}^{\infty} \tilde{Q}_i\right) \left(\sum_{i=1}^{\infty} c_i\right)\right) \text{Ent}_{\mu}(f^2).$$

Combining the above with (3-7) completes the proof. □

Main inequality for c_1 “close” to equilibrium. On the basis of Proposition 3.2, one obtains the following.

Proposition 3.3. *Assume the conditions of Proposition 3.2 and the additional condition that $c_1 < z_*$ for some $0 < z_* < z_s$. Setting*

$$\varrho_* := \sum_{i=1}^{\infty} i Q_i z_*^i < \infty,$$

we have

$$\bar{D}(c) \geq \frac{a_1 z_*^2 c_1^2}{C_{LS}(z_* + \varrho_*)(z_*^2 + 2\varrho(z_* + \varrho_*))} H(c|\mathcal{Q}). \tag{3-11}$$

In particular, if $0 < \delta < c_1 < z_s - \delta$ for some $\delta > 0$,

$$\bar{D}(c) \geq \lambda H(c|\mathcal{Q})$$

for some constant $\lambda > 0$ which depends only on δ, ρ, a_1 and $\{Q_i\}_{i \geq 1}$.

Proof. This follows immediately from (3-6) and the estimates

$$\begin{aligned} \sum_{i=1}^{\infty} \tilde{Q}_i &= \sum_{i=1}^{\infty} Q_i c_1^i \leq c_1 \left(1 + \frac{1}{z_*} \sum_{i=2}^{\infty} Q_i z_*^i\right) < c_1 \left(1 + \frac{\varrho_*}{z_*}\right), \\ \sum_{i=1}^{\infty} c_i &\leq \sum_{i=1}^{\infty} i c_i = \varrho, \end{aligned}$$

together with $\sum_{i=1}^{\infty} a_i \tilde{Q}_i \geq a_1 c_1$. □

Proposition 3.3 shows us that as long as c_1 is bounded away from 0 and z_s , Cercignani’s conjecture will follow immediately from a log-Sobolev inequality for ν with respect to μ (which were defined in Proposition 3.2). Our next result shows that this is indeed true for subcritical masses, under reasonable conditions on the coefficients:

Proposition 3.4. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ satisfy Hypotheses 1–3 with $\gamma = 1$ and let $\mathbf{c} = \{c_i\}_{i \in \mathbb{N}}$ be an arbitrary positive sequence with finite total density $\varrho < \varrho_s < +\infty$. Assume that there exists $\delta > 0$ such that*

$$c_1 \leq z_s - \delta.$$

Then, the measure ν admits a log-Sobolev inequality with respect to the measure μ with constant

$$C_{\text{LS}} = \frac{60z_s^3}{\delta^3} C \left(\frac{z_s - \delta}{z_s} \right) \left(4 + 2e \sup_k |\log(\alpha_{k+1}^{\frac{1}{k+1}})| + e \log \frac{z_s}{\delta} \right), \quad (3-12)$$

where μ and ν were defined in Proposition 3.2 and

$$C(\eta) = 1 + \sup_{k \geq 3} (k(1 + \log(\frac{1}{2}k))\eta^{\frac{k}{2}}) + \frac{2\eta}{1-\eta}$$

for $\eta < 1$.

Proof. We just need to estimate the constant given in Corollary 2.4. As mentioned in the Introduction, we can assume without loss of generality that $a_i = i$. We define

$$\eta = \frac{c_1}{z_s} \leq \frac{z_s - \delta}{z_s} =: \eta_1 < 1.$$

As

$$\tilde{Q}_i = \alpha_i z_s^{1-i} c_1^i \leq z_s \alpha_i \eta^i,$$

we find that due to the monotonicity of $\{\alpha_i\}_{i \in \mathbb{N}}$,

$$z_s \alpha_{k+1} \eta^{k+1} = \tilde{Q}_{k+1} \leq \sum_{i=k+1}^{\infty} \tilde{Q}_i \leq z_s \eta^{k+1} \sum_{i=1}^{\infty} \alpha_{i+k} \eta^{i-1} \leq \frac{z_s \alpha_{k+1} \eta^{k+1}}{1-\eta}.$$

As such

$$\alpha_{k+1} (1-\eta) \eta^k \leq \sum_{i=k+1}^{\infty} \mu_i \leq \alpha_{k+1} \frac{\eta^k}{1-\eta},$$

implying

$$- \sum_{i=k+1}^{\infty} \mu_i \log \left(\sum_{i=k+1}^{\infty} \mu_i \right) \leq \frac{\alpha_{k+1} \eta^k}{1-\eta} \left(k \log \left(\frac{1}{\eta} \right) - \log(\alpha_{k+1} (1-\eta)) \right). \quad (3-13)$$

Next, we notice that as

$$\sum_{i=1}^{\infty} i y^i = \frac{y}{(1-y)^2},$$

one has

$$z_s \eta \leq \sum_{i=1}^{\infty} i \alpha_i z_s \eta^i = \sum_{i=1}^{\infty} a_i \tilde{Q}_i \leq z_s \frac{\eta}{(1-\eta)^2},$$

from which we find that

$$i \alpha_i (1-\eta)^2 \eta^{i-1} \leq v_i \leq i \alpha_i \eta^{i-1}.$$

We notice that for $k \geq 3$, the monotonicity of $\{\alpha_i\}_{i \in \mathbb{N}}$ implies

$$\begin{aligned} k\alpha_k \eta^k \sum_{i=1}^k \frac{1}{i\alpha_i} \left(\frac{1}{\eta}\right)^i &= 1 + \sum_{i=1}^{k-1} \frac{k\alpha_k}{i\alpha_i} \eta^{k-i} \\ &\leq 1 + \sum_{i=1}^{k-1} \frac{k}{i} \eta^{k-i} = 1 + \sum_{i=1}^{\lfloor \frac{k}{2} \rfloor} \frac{k}{i} \eta^{k-i} + \sum_{i=\lfloor \frac{k}{2} \rfloor + 1}^{k-1} \frac{k}{i} \eta^{k-i} \\ &\leq 1 + k\eta_1^{\frac{k}{2}} \sum_{i=1}^{\lfloor \frac{k}{2} \rfloor} \frac{1}{i} + \frac{k}{\lfloor \frac{k}{2} \rfloor + 1} \sum_{j=1}^{\infty} \eta_1^j \\ &\leq 1 + k(1 + \log(\frac{1}{2}k))\eta_1^{\frac{k}{2}} + \frac{2\eta_1}{1 - \eta_1}. \end{aligned}$$

Using the definition of $C(\eta)$ and the fact that $C(\eta) > 1 + \eta$, we find that for all $k \in \mathbb{N}$,

$$k\alpha_k \eta^k \sum_{i=1}^k \frac{1}{i\alpha_i} \left(\frac{1}{\eta}\right)^i \leq C(\eta_1),$$

and as such

$$\sum_{i=1}^k \frac{1}{v_i} \leq C(\eta_1) \frac{\eta}{(1 - \eta)^2} \frac{1}{k\alpha_k} \left(\frac{1}{\eta}\right)^k. \tag{3-14}$$

Combining the above with (3-13) yields the bound

$$\left(-\sum_{i=k+1}^{\infty} \mu_i \log\left(\sum_{i=k+1}^{\infty} \mu_i\right)\right) \left(\sum_{i=1}^k \frac{1}{v_i}\right) \leq C(\eta_1) \frac{\alpha_{k+1}}{\alpha_k} \frac{\eta}{(1 - \eta)^3} \left(\log\left(\frac{1}{\eta}\right) - \frac{1}{k} \log(\alpha_{k+1}(1 - \eta))\right).$$

Thus, with the notation of Corollary 2.4,

$$\begin{aligned} D_1 &\leq \frac{C(\eta_1)}{(1 - \eta_1)^3} \left(\sup_{0 \leq x \leq 1} (-\eta \log(\eta)) + \eta_1 \sup_k \frac{k+1}{k} |\log(\alpha_{k+1}^{\frac{1}{k+1}})| + \eta_1 \log\left(\frac{1}{1 - \eta_1}\right) \right) \\ &\leq \frac{C(\eta_1)}{(1 - \eta_1)^3} \left(\frac{1}{e} + 2\eta_1 \sup_k |\log(\alpha_{k+1}^{\frac{1}{k+1}})| + \eta_1 \log\left(\frac{1}{1 - \eta_1}\right) \right). \end{aligned}$$

As m , defined in Corollary 2.4, is always finite, we conclude using the same corollary that ν admits a log-Sobolev inequality with respect to μ . However, in order to estimate the constant C_{LS} , we still need to estimate the constant D_2 in the case where $m > 1$ (otherwise, $D_2 = 0$).

Since

$$\sum_{i=m}^{\infty} \mu_i \leq \frac{\alpha_m}{1 - \eta} \eta^{m-1},$$

the requirement that $\sum_{i=1}^{m-1} \mu_i < \frac{2}{3}$ implies

$$\frac{1}{\alpha_{m-1} \eta^{m-1}} \leq \frac{\alpha_m}{\alpha_{m-1}} \frac{3}{(1 - \eta)} \leq \frac{3}{(1 - \eta)}.$$

Using the above along with the fact that $m > 1$ and inequality (3-14) shows that

$$\sum_{i=1}^{m-1} \frac{1}{v_i} \leq 3C(\eta_1) \frac{\eta_1}{(1-\eta_1)^3} \frac{1}{m-1} \leq 3C(\eta_1) \frac{\eta_1}{(1-\eta_1)^3}.$$

We can conclude that

$$\left(-\sum_{i=m-1}^{\infty} \mu_i \log\left(\sum_{i=m-1}^{\infty} \mu_i\right)\right) \left(\sum_{i=1}^{m-1} \frac{1}{v_i}\right) \leq 3 \sup_{0 \leq x \leq 1} (-x \log x) C(\eta_1) \frac{\eta_1}{(1-\eta_1)^3}, \tag{3-15}$$

from which we conclude that

$$D_2 \leq \frac{3}{e} C(\eta_1) \frac{\eta_1}{(1-\eta_1)^3},$$

which completes the proof, as the result follows directly from Corollary 2.4. □

We finally have all the tools to prove part (i) of Theorem 1.1:

Proof of part (i) of Theorem 1.1. The result follows immediately from Proposition 3.3, Proposition 3.4 and condition (1-16). □

The last part of this section will be devoted to the proof of part (ii) of Theorem 1.1. For that we will need the following lemma:

Lemma 3.5. *For any $\beta \geq 0$, any nonnegative sequence c and positive sequence $\{Q_i\}_{i \geq 1}$, it holds that*

$$\sum_{i=1}^{\infty} i^\beta Q_i \left(\sqrt{\frac{c_1 c_i}{Q_i}} - \sqrt{\frac{c_{i+1}}{Q_{i+1}}}\right)^2 \leq 2 \left(c_1 + \sup_j \frac{Q_j}{Q_{j+1}}\right) \sum_{i=1}^{\infty} i^\beta c_i. \tag{3-16}$$

Proof. The proof is a direct consequence of the inequality $(a + b)^2 \leq 2(a^2 + b^2)$:

$$\begin{aligned} \sum_{i=1}^{\infty} i^\beta Q_i \left(\sqrt{\frac{c_1 c_i}{Q_i}} - \sqrt{\frac{c_{i+1}}{Q_{i+1}}}\right)^2 &\leq 2c_1 \sum_{i=1}^{\infty} i^\beta c_i + 2 \sum_{i=1}^{\infty} i^\beta \frac{Q_i}{Q_{i+1}} c_{i+1} \\ &\leq 2 \left(c_1 + \sup_j \frac{Q_j}{Q_{j+1}}\right) \sum_{i=1}^{\infty} i^\beta c_i. \end{aligned} \tag{3-17}$$
□

Proof of part (ii) of Theorem 1.1. We denote by $\bar{D}_\gamma(c)$ the lower free energy dissipation of c associated to the coagulation coefficient $a_i = i^\gamma$. According to part (i) of Theorem 1.1, there exists $K > 0$ that depends only on δ, z_s, Q and $\{\alpha_i\}_{i \in \mathbb{N}}$ such that

$$\bar{D}_1(c) \geq KH(c|Q).$$

Using interpolation between γ and β , we find that

$$\bar{D}_1(c) \leq \bar{D}_\gamma^{\frac{\beta-1}{\beta-\gamma}}(c) \bar{D}_\beta^{\frac{1-\gamma}{\beta-\gamma}}(c) \leq 2^{\frac{1-\gamma}{\beta-\gamma}} \bar{D}_\gamma^{\frac{\beta-1}{\beta-\gamma}}(c) \left(z_s + \frac{1}{z_s} \sup_j \frac{\alpha_j}{\alpha_{j+1}}\right)^{\frac{1-\gamma}{\beta-\gamma}} M_\beta^{\frac{1-\gamma}{\beta-\gamma}}, \tag{3-17}$$

where we have used [Lemma 3.5](#), the upper bound on c_1 and [Hypothesis 2](#). Therefore

$$D(\mathbf{c}) \geq \bar{D}_\gamma(\mathbf{c}) \geq \left(\frac{z_s K^{\frac{\beta-\gamma}{1-\gamma}}}{2(z_s^2 + \sup_j \alpha_j / \alpha_{j+1}) M_\beta} \right)^{\frac{1-\gamma}{\beta-1}} H(\mathbf{c} | \mathcal{Q})^{\frac{\beta-\gamma}{\beta-1}} \tag{3-18}$$

and the proof is now complete. □

This concludes the part of the proof of [Theorem 1.1](#) that relied on the log-Sobolev inequality. In the next subsection we will address the question of what happens when c_1 escapes the “good region” delimited by [\(1-16\)](#).

Energy dissipation estimate when c_1 is “far” from equilibrium. The goal of this subsection is to show that when c_1 is far from equilibrium, in the aforementioned sense, while we may lose our desired inequality between $\bar{D}(\mathbf{c})$ and $H(\mathbf{c} | \mathcal{Q})$, the energy dissipation becomes *uniformly large* — forcing the free energy to decrease (and as a consequence, the distance between c_1 and \bar{z} decreases as well).

The next proposition, dealing with the case when c_1 is “too large”, is an adaptation of a theorem from [\[Jabin and Niethammer 2003\]](#).

Proposition 3.6. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ be the coagulation and detailed balance coefficients for the Becker–Döring equations. Assume that $\inf_i a_i > 0$ and*

$$\lim_{i \rightarrow \infty} \frac{Q_{i+1}}{Q_i} = \frac{1}{z_s}.$$

Let $\mathbf{c} = \{c_i\}$ be a nonnegative sequence with finite total density $\varrho < \varrho_s$. Then, if

$$c_1 > \bar{z} + \delta$$

for any $\delta > 0$, we have

$$\bar{D}(\mathbf{c}) > \varepsilon_1$$

for a fixed constant ε_1 that depends only on $\{Q_i\}_{i \in \mathbb{N}}$, \bar{z} , z_s and δ .

Proof. Without loss of generality we may assume that $\bar{z} + \delta < z_s$. Defining $u_i = c_i / Q_i$ we notice that

$$\bar{D}(\mathbf{c}) = \sum_{i=1}^{\infty} a_i Q_i (\sqrt{c_1 u_i} - \sqrt{u_{i+1}})^2.$$

Let $\lambda < 1$ be such that $\lambda c_1 = \bar{z} + \frac{1}{2}\delta$ and let $i_0 \in \mathbb{N}$ be the first index such that

$$u_{i+1} < \lambda c_1 u_i.$$

This index exists, else, for any $i \in \mathbb{N}$ we have

$$u_{i+1} \geq \lambda c_1 u_i \geq (\lambda c_1)^i c_1, \tag{3-19}$$

and thus

$$\varrho = \sum_{i=1}^{\infty} i c_i \geq c_1 + c_1 \sum_{i=2}^{\infty} i Q_i (\lambda c_1)^{i-1} \geq \sum_{i=1}^{\infty} i Q_i (\bar{z} + \frac{1}{2}\delta)^i,$$

which is a contradiction.

Due to the positivity of each term in the sum consisting of the lower free energy dissipation, we conclude that

$$\bar{D}(\mathbf{c}) \geq a_{i_0} Q_{i_0} (1 - \sqrt{\lambda})^2 c_1 u_{i_0} \geq a_{i_0} Q_{i_0} \lambda^{i_0-1} c_1^{i_0+1} (1 - \sqrt{\lambda})^2, \tag{3-20}$$

where we have used the fact that up to $i_0 - 1$ we have inequality (3-19).

As we know that there exists $C > 0$, depending only on $\{Q_i\}_{i \in \mathbb{N}}$, \bar{z} , z_s and δ such that

$$\sum_{i=i_0+1}^{\infty} i c_1 (\lambda c_1)^{i-1} Q_i \leq C Q_{i_0} (\lambda c_1)^{i_0} c_1$$

(see Lemma B.2 in Appendix B), we conclude that, using (3-19) again,

$$C Q_{i_0} (\lambda c_1)^{i_0} c_1 \geq \tilde{\varrho} - \sum_{i=1}^{i_0} i Q_i (\lambda c_1)^{i-1} c_1 \geq \tilde{\varrho} - \sum_{i=1}^{i_0} i c_i \geq \tilde{\varrho} - \varrho,$$

where $\tilde{\varrho} = \sum_{i=1}^{\infty} i Q_i (\lambda c_1)^{i-1} c_1$. We can estimate the difference $\varrho - \tilde{\varrho}$ as

$$\tilde{\varrho} - \varrho \geq \sum_{i=1}^{\infty} i Q_i \left((\bar{z} + \frac{1}{2}\delta)^i - \bar{z}^i \right) \geq \left(\sum_{i=1}^{\infty} i^2 Q_i \bar{z}^{i-1} \right) \frac{1}{2}\delta.$$

In conclusion, there exists a universal constant $C_1 > 0$, depending only on $\{Q_i\}_{i \in \mathbb{N}}$, \bar{z} , z_s and δ , and not on i_0 , c_1 or λ , such that

$$Q_{i_0} (\lambda c_1)^{i_0} c_1 > C_1.$$

Recalling (3-20) and using the fact that

$$\lambda = \frac{\bar{z} + \frac{1}{2}\delta}{c_1} < \frac{\bar{z} + \frac{1}{2}\delta}{\bar{z} + \delta}$$

we find that

$$\bar{D}(\mathbf{c}) \geq C_1 a_{i_0} \frac{(1 - \sqrt{\lambda})^2}{\lambda} \geq C_1 \inf_{i \geq 1} a_i \frac{(\sqrt{\bar{z} + \delta} - \sqrt{\bar{z} + \frac{1}{2}\delta})^2}{\bar{z} + \frac{1}{2}\delta},$$

completing the proof. □

Next, we present a new lower bound estimate for the energy dissipation in the case where c_1 is “too small”.

Lemma 3.7. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ be the coagulation and detailed balance coefficients for the Becker–Döring equations. Assume that*

$$\begin{aligned} \bar{Q} &= \sup_i \frac{Q_i}{Q_{i+1}} < +\infty, & \underline{Q} &= \inf_i \frac{Q_i}{Q_{i+1}} < +\infty, \\ \bar{a} &= \sup_i \frac{a_i}{a_{i+1}} < +\infty, & \underline{a} &= \inf_i \frac{a_i}{a_{i+1}} < +\infty, \end{aligned}$$

and let \mathbf{c} be a nonnegative sequence such that

$$c_1 < \delta$$

for some $\delta > 0$. Then,

$$\bar{D}(\mathbf{c}) \geq \underline{Q}a \left(\sum_{i=1}^{\infty} a_i c_i - a_1 \delta \right) - 2\sqrt{\delta} \sqrt{\bar{Q}a} \left(\sum_{i=1}^{\infty} a_i c_i \right).$$

Proof. Expanding the square, one has

$$\bar{D}(\mathbf{c}) = c_1 \sum_{i=1}^{\infty} a_i c_i + \sum_{i=1}^{\infty} a_i \frac{Q_i}{Q_{i+1}} c_{i+1} - 2\sqrt{c_1} \sum_{i=1}^{\infty} a_i \sqrt{\frac{Q_i}{Q_{i+1}}} \sqrt{c_i c_{i+1}}$$

so that

$$\begin{aligned} \bar{D}(\mathbf{c}) &\geq \underline{Q}a \left(\sum_{i=2}^{\infty} a_i c_i \right) - 2\sqrt{c_1} \sqrt{\bar{Q}a} \left(\sum_{i=2}^{\infty} a_i c_i \right)^{\frac{1}{2}} \left(\sum_{i=1}^{\infty} a_i c_i \right)^{\frac{1}{2}} \\ &\geq \underline{Q}a \left(\sum_{i=1}^{\infty} a_i c_i - a_1 \delta \right) - 2\sqrt{\delta} \sqrt{\bar{Q}a} \left(\sum_{i=1}^{\infty} a_i c_i \right), \end{aligned}$$

which is the desired result. □

Proposition 3.8. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ be the coagulation and detailed balance coefficients for the Becker–Döring equations. Assume that*

$$\bar{Q} = \sup_i \frac{Q_i}{Q_{i+1}} < +\infty, \quad \underline{Q} = \inf_i \frac{Q_i}{Q_{i+1}} < +\infty.$$

Let \mathbf{c} be a nonnegative sequence with finite total density ϱ . Then:

(i) If $a_i = i$ then there exists a $\delta_1 > 0$, depending only on \bar{Q} , \underline{Q} and ϱ such that if $c_1 < \delta_1$ then

$$\bar{D}(\mathbf{c}) \geq \frac{\underline{Q}\varrho}{4}.$$

(ii) If $a_i = i^\gamma$ for $\gamma < 1$ and there exists $\beta > 1$ such that $M_\beta < +\infty$, then there exists $\delta_1 > 0$, depending only on \bar{Q} , \underline{Q} , ϱ and M_β such that if $c_1 < \delta_1$ then

$$\bar{D}(\mathbf{c}) \geq \frac{\underline{Q}\varrho^{\frac{\beta-\gamma}{\beta-1}}}{4M_\beta^{\frac{1-\gamma}{\beta-1}}}.$$

Proof. Both (i) and (ii) will follow immediately from Lemma 3.7 and a suitable choice of δ_1 . Indeed, for (i) we notice that

$$\underline{Q}a \left(\sum_{i=1}^{\infty} a_i c_i - a_1 \delta \right) - 2\sqrt{\delta} \sqrt{\bar{Q}a} \left(\sum_{i=1}^{\infty} a_i c_i \right) = \frac{\underline{Q}}{2}(\varrho - \delta) - 2\sqrt{\delta} \sqrt{\bar{Q}a},$$

where we have used the notations of Lemma 3.7. As the above is less than $\frac{1}{2}\underline{Q}\varrho$ and converges to it as δ goes to zero, we can find δ_1 that satisfies the desired result.

For (ii) we notice that the interpolation estimate

$$\varrho = \sum_{i=1}^{\infty} i c_i \leq \left(\sum_{i=1}^{\infty} i^\gamma c_i \right)^{\frac{\beta-1}{\beta-\gamma}} (M_\beta)^{\frac{1-\gamma}{\beta-\gamma}}$$

along with the fact that $\sum_{i=1}^{\infty} i^\gamma c_i \leq \varrho$ implies

$$\underline{Q}a \left(\sum_{i=1}^{\infty} a_i c_i - a_1 \delta \right) - 2\sqrt{\delta} \sqrt{\underline{Q}a} \left(\sum_{i=1}^{\infty} a_i c_i \right) \geq \frac{\underline{Q}}{2} \left(\frac{Q^{\frac{\beta-\gamma}{\beta-1}}}{M_\beta^{\frac{1-\gamma}{\beta-1}}} - \delta \right) - 2\sqrt{\delta} \sqrt{\underline{Q}} \varrho,$$

from which the result follows. □

We are finally ready to complete the proof of [Theorem 1.1](#):

Proof of part (iii) of Theorem 1.1. This follows immediately from [Propositions 3.6](#) and [3.8](#). □

Now that we have our general functional inequality at hand, one may wonder about the sharpness of this method of using the log-Sobolev inequality. Perhaps we were too coarse in our estimation, and Cercignani’s conjecture *is* valid in the case $a_i = i^\gamma$ with $\gamma < 1$ under the restrictions of [Theorem 1.1](#). The answer, surprisingly, is that the result is optimal, as we shall see in the next subsection.

Optimality of the results. This subsection is devoted to showing that unlike the case $a_i = i$, the case $a_i = i^\gamma$ when $\gamma < 1$ does not satisfy Cercignani’s conjecture, even if c_1 is bounded appropriately. This is stated in [Theorem 1.2](#).

Proof of Theorem 1.2. We start by choosing $a_i = i^\gamma$, $\gamma < 1$ and $Q_i = e^{-\lambda(i-1)}$ ($i \geq 1$) for some $\lambda \geq 0$. We will show the desired result by constructing a family of nonnegative sequences $\{c^{(\varepsilon)}\}_{\varepsilon>0}$ with a fixed mass ϱ such that

$$\lim_{\varepsilon \rightarrow 0} \frac{D(c^{(\varepsilon)})}{H(c^{(\varepsilon)}|Q)} = 0.$$

Let $\xi > 0$ be such that

$$\frac{\varrho}{2} = \sum_{i=1}^{\infty} i e^{\lambda} e^{-\xi i} = \frac{e^{\lambda-\xi}}{(1-e^{-\xi})^2}.$$

Consider the sequence $c^{(\varepsilon)} = \{c_i^{(\varepsilon)}\}$ given by

$$c_i^{(\varepsilon)} = e^{\lambda} e^{-\xi i} + A_\varepsilon e^{-\varepsilon i}, \quad i \in \mathbb{N},$$

where $0 < \varepsilon$ is small and A_ε is chosen such that the mass of the sequence $c^{(\varepsilon)}$ is ϱ , i.e., $A_\varepsilon = \frac{1}{2} \varrho e^\varepsilon (1 - e^{-\varepsilon})^2$. Next, as $Q_i/Q_{i+1} = e^\lambda$ for any $i \geq 1$, we see that

$$\begin{aligned} \frac{Q_i}{Q_{i+1}} c_{i+1}^{(\varepsilon)} - c_1^{(\varepsilon)} c_i^{(\varepsilon)} &= e^{2\lambda} e^{-\xi(i+1)} + A_\varepsilon e^{\lambda} e^{-\varepsilon(i+1)} - e^{2\lambda} e^{-\xi(i+1)} - A_\varepsilon e^{\lambda} (e^{-\xi i - \varepsilon} + e^{-\varepsilon i - \xi}) - A_\varepsilon^2 e^{-\varepsilon(i+1)} \\ &= A_\varepsilon e^{\lambda} e^{-\varepsilon(i+1)} (1 - e^{-(\xi-\varepsilon)} - e^{-(\xi-\varepsilon)i} - A_\varepsilon e^{-\lambda}) > 0 \end{aligned}$$

for ε small enough depending only on λ, ξ and Q but not on i . Additionally, one can easily verify that

$$\frac{Q_i c_{i+1}^{(\varepsilon)}}{Q_{i+1} c_1^{(\varepsilon)} c_i^{(\varepsilon)}} \leq e^\lambda \left(1 + \frac{1}{A_\varepsilon}\right).$$

As such, setting $B_{z,\gamma} = \sum_{i=1}^\infty i^\gamma e^{-zi}$ for any $z > 0$, we find that

$$\begin{aligned} D(\mathbf{c}^{(\varepsilon)}) &= \sum_{i=1}^\infty i^\gamma \left(\frac{Q_i}{Q_{i+1}} c_{i+1}^{(\varepsilon)} - c_i^{(\varepsilon)} \right) \log \left(\frac{Q_i c_{i+1}^{(\varepsilon)}}{Q_{i+1} c_1^{(\varepsilon)} c_i^{(\varepsilon)}} \right) \\ &\leq A_\varepsilon e^\lambda B_{\varepsilon,\gamma} \log \left(e^\lambda \left(1 + \frac{1}{A_\varepsilon}\right) \right) \left((1 - A_\varepsilon e^{-\lambda}) e^{-\varepsilon} - e^{-\xi} \right) - A_\varepsilon e^\lambda B_{\xi,\gamma} \log \left(e^{\lambda-\varepsilon} \left(1 + \frac{1}{A_\varepsilon}\right) \right). \end{aligned} \tag{3-21}$$

As $A_\varepsilon \approx \frac{1}{2} Q \varepsilon^2$ when ε approaches zero, and $B_{\varepsilon,\gamma}$ is of order $\varepsilon^{-(1+\gamma)}$ (see Lemma B.3 in Appendix B) we conclude that

$$\lim_{\varepsilon \rightarrow 0} D(\mathbf{c}^{(\varepsilon)}) = 0.$$

Lastly, we turn our attention to the relative free energy. We start by denoting by $\bar{\xi} > 0$ the unique parameter for which

$$Q = e^\lambda \sum_{i=1}^\infty i e^{-\bar{\xi}i}.$$

Clearly, $\bar{\xi} < \xi$ and the associated equilibrium with mass Q is $Q_i = e^\lambda e^{-\bar{\xi}i}$. Since, for any fixed $i \geq 1$, it holds that

$$\lim_{\varepsilon \rightarrow 0} c_i^{(\varepsilon)} = c_i^{(0)} = e^\lambda e^{-\xi i},$$

using Fatou’s lemma we can conclude that

$$\liminf_{\varepsilon \rightarrow 0} H(\mathbf{c}^{(\varepsilon)} | Q) \geq H(\mathbf{c}^{(0)} | Q) > 0,$$

as $\mathbf{c}^{(0)} \neq Q$. □

Remark 3.9. We notice the following:

- In the example we provided, $z_s = e^\lambda < +\infty$ but $\varrho_s = +\infty$. This, however, is not a great obstacle as all our proofs rely on some *positive* distance from z_s and ϱ_s , and can be reformulated accordingly.
- The constructed sequence $\mathbf{c}^{(\varepsilon)}$ satisfies

$$\sup_\varepsilon \sum_{i=1}^\infty i^\beta c_i^{(\varepsilon)} = +\infty$$

for any $\beta > 1$. Thus, the conclusion of part (ii) of Theorem 1.1 does not apply to it. Actually, one can easily check that

$$\lim_{\varepsilon \rightarrow 0} \frac{D(\mathbf{c}^{(\varepsilon)})}{(H(\mathbf{c}^{(\varepsilon)} | Q))^s} = 0$$

for any $s > 0$.

Inequalities with exponential moments. Up to now, we have avoided using exponential moments in any of our functional inequalities. In this section we will show that when $0 \leq \gamma < 1$, under the additional assumption of a bounded exponential moment, one can obtain an improved functional inequality between $\bar{D}(c)$ and $H(c|\mathcal{Q})$, extending the result given by Jabin and Niethammer [2003]. The key idea in this section is to avoid using the interpolation inequality (3-17) and replace it with one that involves an exponential weight.

Proposition 3.10. *Let f be a nonnegative sequence and let $0 \leq \gamma < 1$. Assume that there exists $\mu \in (0, 4 \log 2)$ such that*

$$\sum_{i=1}^{\infty} e^{\mu i} f_i = M_{\mu}^{\text{exp}}(f) < +\infty.$$

Then,

$$M_{\gamma}(f) \geq \frac{M_1(f)}{2} \left(\frac{2}{\mu} \log \left(\frac{4M_{\mu}^{\text{exp}}(f)}{\mu e M_1(f)} \right) \right)^{-(1-\gamma)}, \quad (3-22)$$

where $M_{\alpha}(f)$ denotes the α -moment of f and $M_{\mu}^{\text{exp}}(f)$ is the exponential moment defined in (1-14).

Proof. For simplicity, we will use the notation of M_1 and M_{μ}^{exp} instead of $M_1(f)$ and $M_{\mu}^{\text{exp}}(f)$. We start with the simple inequality

$$\begin{aligned} M_1 &= \sum_{i=1}^{\infty} i f_i = \sum_{i=1}^N i^{1-\gamma} i^{\gamma} f_i + \sum_{i=N+1}^{\infty} i e^{-\frac{\mu i}{2}} e^{-\frac{\mu i}{2}} e^{\mu i} f_i \\ &\leq N^{1-\gamma} M_{\gamma} + \frac{2e^{-\frac{\mu(N+1)}{2}}}{\mu e} M_{\mu}^{\text{exp}} \quad \forall N \in \mathbb{N}, \end{aligned} \quad (3-23)$$

where we used the fact that $\sup_{x \geq 0} x e^{-\lambda x} = 1/(\lambda e)$ for any $\lambda > 0$. Our goal will be to choose a particular N to plug in the inequality above to conclude the desired result. Again, using the supremum of $g(x) = x e^{-\lambda x}$, we conclude that

$$M_1 \leq \frac{1}{\mu e} M_{\mu}^{\text{exp}}.$$

As $\mu < 4 \log 2$, we find that

$$M_1 < \frac{4M_{\mu}^{\text{exp}}}{\mu e^{1+\frac{\mu}{2}}},$$

from which we conclude that

$$N = \left\lceil \frac{2}{\mu} \log \left(\frac{4M_{\mu}^{\text{exp}}}{\mu e M_1} \right) \right\rceil \geq 1.$$

Plugging this N into (3-23) we see that

$$e^{-\frac{\mu(N+1)}{2}} \leq \frac{\mu e M_1}{4M_{\mu}^{\text{exp}}},$$

and as such

$$M_{\gamma} \geq N^{\gamma-1} \frac{M_1}{2}$$

and the result follows. \square

With this proposition at hand, we are prepared to show part (i) of [Theorem 1.4](#).

Proof of part (i) of Theorem 1.4. Without loss of generality we may assume that $\mu \in (0, 4 \log 2)$. Introduce the sequence $f = \{f_i\}$, where

$$f_i = Q_i \left(\sqrt{\frac{c_1 c_i}{Q_i}} - \sqrt{\frac{c_{i+1}}{Q_{i+1}}} \right)^2, \quad i \geq 1.$$

Following the same proof as presented in [Lemma 3.5](#), we find that

$$M_\mu^{\text{exp}}(f) \leq 2 \left(c_1 + z_s \sup_j \frac{\alpha_j}{\alpha_{j+1}} \right) M_\mu^{\text{exp}}(c).$$

Thus, using the simple fact that $M_\alpha(f) = \bar{D}_\alpha(c)$, for any $\alpha > 0$, together with [Proposition 3.10](#) and parts (i) and (iii) of [Theorem 1.1](#), yields the desired functional inequality. \square

4. Rate of convergence to equilibrium

In this section we will use all the information we gathered so far to prove [Theorem 1.3](#) and part (ii) of [Theorem 1.4](#), giving an explicit rate of convergence to equilibrium for the Becker–Döring equations.

The convergence result in [Theorem 1.3](#) is a consequence of [Theorem 1.1](#). To use the functional inequality established there, we need first to invoke uniform (and explicit) upper bounds on moments $M_\beta(c(t))$; see (1-18). This is provided by the following (see [[Cañizo et al. 2017](#)]):

Proposition 4.1. *Let $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$ satisfy Hypotheses 1–3 with $0 \leq \gamma \leq 1$, and let $c(t) = \{c_i(t)\}_{i \in \mathbb{N}}$ be a solution to the Becker–Döring equations with mass $\varrho \in (0, \varrho_s)$. Let $\beta \geq \max\{2 - \gamma, 1 + \gamma\}$ be such that*

$$M_\beta(c(0)) = \sum_{i=1}^{\infty} i^\beta c_i(0) < \infty.$$

There exists a constant $C > 0$ depending only on β , $M_\beta(0)$, the initial relative free energy $H(c(0)|\mathcal{Q})$, the coefficients $\{a_i\}_{i \geq 1}$, $\{b_i\}_{i \geq 1}$ and the mass ϱ such that

$$M_\beta(c(t)) = \sum_{i=1}^{\infty} i^\beta c_i(t) \leq C \quad \text{for all } t \geq 0.$$

Using such an estimate, the proof is easily derived from [Theorem 1.1](#) and part (i) of [Theorem 1.4](#), yet we provide a proof here for the sake of completeness and to show that we can find all the constants explicitly.

Proof of Theorem 1.3. Combining [Theorem 1.1](#) and [Proposition 4.1](#), we conclude the differential inequality

$$\frac{d}{dt} H(c(t)|\mathcal{Q}) \leq \begin{cases} -\min(KH(c(t)|\mathcal{Q}), \varepsilon), & \gamma = 1, \\ -\min(KH(c(t)|\mathcal{Q})^{\frac{\beta-\gamma}{\beta-1}}, \varepsilon), & 0 \leq \gamma < 1 \end{cases} \tag{4-1}$$

for appropriate K and ε . We claim that there exists $t_0 \geq 0$ such that for all $t \geq t_0$,

$$H(c(t)|\mathcal{Q}) \leq \begin{cases} \frac{\varepsilon}{K}, & \gamma = 1, \\ \left(\frac{\varepsilon}{K}\right)^{\frac{\beta-1}{\beta-\gamma}}, & 0 \leq \gamma < 1. \end{cases} \tag{4-2}$$

Indeed, if $H(c(t)|\mathcal{Q})$ is larger than the appropriate constants in $[0, t]$ then

$$\frac{d}{ds} H(c(s)|\mathcal{Q}) \leq -\varepsilon \quad \forall s \in (0, t),$$

implying that

$$H(c(t)|\mathcal{Q}) \leq H(c(0)|\mathcal{Q}) - \varepsilon t.$$

We define

$$t_0 = \begin{cases} \min(0, (H(c(0)|\mathcal{Q}) - \frac{\varepsilon}{K})/\varepsilon), & \gamma = 1, \\ \min(0, (H(c(0)|\mathcal{Q}) - (\frac{\varepsilon}{K})^{\frac{\beta-1}{\beta-\gamma}})/\varepsilon), & 0 \leq \gamma < 1, \end{cases}$$

and find that $H(c(t_0)|\mathcal{Q})$ satisfies the appropriate inequality in (4-2). As $H(c(t)|\mathcal{Q})$ is decreasing, we conclude that (4-2) is valid for any $t \geq t_0$.

With this in hand, along with (4-1), we have, for all $t \geq t_0$,

$$H(c(t)|\mathcal{Q}) \leq \begin{cases} H(c(t_0)|\mathcal{Q})e^{-K(t-t_0)}, & \gamma = 1, \\ (H(c(t_0)|\mathcal{Q})^{\gamma-1/\beta-1} + \frac{1-\gamma}{\beta-1}K(t-t_0))^{-\frac{\beta-1}{1-\gamma}}, & 0 \leq \gamma < 1. \end{cases}$$

As

$$H(c(t_0)|\mathcal{Q}) = \begin{cases} \min(H(c(0)|\mathcal{Q}), \frac{\varepsilon}{K}), & \gamma = 1, \\ \min(H(c(0)|\mathcal{Q}), (\frac{\varepsilon}{K})^{\frac{\beta-1}{\beta-\gamma}}), & 0 \leq \gamma < 1, \end{cases}$$

and t_0 is given explicitly, we conclude that

$$C(H(c(0)|\mathcal{Q})) = \begin{cases} H(c(0)|\mathcal{Q}), & \gamma = 1, t_0 = 0, \\ \frac{\varepsilon}{K}e^{K\frac{1}{\varepsilon}(H(c(0)|\mathcal{Q}) - \frac{\varepsilon}{K})}, & \gamma = 1, t_0 > 0, \\ H(c(0)|\mathcal{Q}), & 0 \leq \gamma < 1, t_0 = 0, \\ \left(\frac{\varepsilon}{K}\right)^{\frac{\gamma-1}{\beta-\gamma}} - \frac{1-\gamma}{\beta-1}K\frac{1}{\varepsilon}\left(H(c(0)|\mathcal{Q}) - \left(\frac{\varepsilon}{K}\right)^{\frac{\beta-1}{\beta-\gamma}}\right), & 0 \leq \gamma < 1, t_0 > 0, \end{cases}$$

completing the proof. □

Proof of part (ii) of Theorem 1.4. This follows from part (i) of Theorem 1.4 by the same methods used in the above proof and the fact that


$$\sup_{t \geq 0} M_{\mu'}^{\text{exp}}(c(t)) < +\infty$$

for some $0 < \mu' < \mu$ (a known result from [Jabin and Niethammer 2003]). □

5. Consequences for general coagulation and fragmentation models

In this final section we illustrate how the functional inequalities investigated in Section 3 provide new insights on the behaviour of solutions to general discrete coagulation-fragmentation models.

General discrete coagulation-fragmentation equation. The Becker–Döring equations (1-1) are derived under the assumption that the only relevant reactions taking place are those between monomers and clusters of any size. One can obtain a more general model by taking into account reactions between clusters of any size. Keeping the notation of the Introduction, this means that we consider reactions of the type



for any positive integer sizes i and j . We assume their coagulation rate (i.e., the reaction from left to right) is determined by a coefficient we call $a_{i,j}$, and their fragmentation rate (the reaction from right to left) by a coefficient called $b_{i,j}$. These coefficients are always assumed to be nonnegative (as before) and symmetric in i, j (that is, $a_{i,j} = a_{j,i}$ and $b_{i,j} = b_{j,i}$ for all i, j). The equation corresponding to (1-1) is then

$$\frac{d}{dt} c_i(t) = \frac{1}{2} \sum_{j=1}^{i-1} W_{j,i-j}(t) - \sum_{j=1}^{\infty} W_{i,j}(t), \quad i \in \mathbb{N}, \tag{5-1}$$

where

$$W_{i,j}(t) := a_{i,j} c_i(t)c_j(t) - b_{i,j} c_{i+j}(t), \quad i \in \mathbb{N}. \tag{5-2}$$

The system (1-1) is then a particular case of (5-1) obtained by choosing $a_{i,j}, b_{i,j}$ as

$$a_{i,j} = b_{i,j} = 0 \quad \text{when } \min\{i, j\} \geq 2, \tag{5-3}$$

$$a_{i,1} := 2a_1, \quad a_{i,1} = a_{1,i} = a_i \quad \text{for } i \geq 2, \tag{5-4}$$

$$b_{1,1} := 2b_2, \quad b_{i,1} = b_{1,i} = b_{i+1} \quad \text{for } i \geq 2. \tag{5-5}$$

The mathematical theory of this full system is much less complete than that of (1-1). Well-posedness of mass-conserving solutions has been studied in [Ball and Carr 1990], and there are a number of works on asymptotic behaviour, for instance [Cañizo 2005; 2007; Carr and da Costa 1994; Carr 1992], but it is still not fully understood. To start with, it is unclear whether equilibria of (5-1) are unique or not (when they exist). A common physical condition imposed on the coefficients $a_{i,j}, b_{i,j}$ which avoids this problem is that of *detailed balance*: we say it holds when there exists a sequence $\{Q_i\}_{i \geq 1}$ of strictly positive numbers such that

$$a_{i,j} Q_i Q_j = b_{i,j} Q_{i+j} \quad \text{for any } i, j, \tag{5-6}$$

where we always further assume without loss of generality that $Q_1 = 1$. This is the analogue of (1-4), but in this case it needs to be imposed as a condition since numbers Q_i satisfying (5-6) cannot always be found (unlike in the Becker–Döring case). If we assume (5-6) then equilibria (5-1) exist and have the same form (1-5) as in the Becker–Döring case, and a similar phase transition in the long-time behaviour has been rigorously proved in some cases (see [Cañizo 2005; 2007; Carr and da Costa 1994; Carr 1992] for more details). However, even with detailed balance, the long-time behaviour is in general not understood except in particular cases. If clusters larger than a given size N do not react among themselves (that is, if $a_{i,j} = b_{i,j} = 0$ whenever $\min\{i, j\} > N$) the system is known as the *generalised Becker–Döring system*, and has been studied in [Cañizo 2005; da Costa 1998]. For coefficients $a_{i,j}$ given by

$$a_{i,j} = i^\nu j^\eta + i^\eta j^\nu \quad \text{for any } i, j, \tag{5-7}$$

with $\eta \leq 0 \leq \gamma$ and $\gamma + \eta \leq 1$, the asymptotic behaviour was identified in [Cañizo 2007] and a constructive (though probably far from optimal) rate of convergence to equilibrium was given. Very little is known about the asymptotic behaviour for coefficients of the type (5-7) with $\gamma, \eta > 0$ and $\gamma + \eta \leq 1$. In this case the size of $a_{i,j}$ is larger than that of $a_{i,1}$ and the system (5-1) may behave quite differently from (1-1).

A natural question is whether any of the functional inequalities investigated in this paper can shed new light on the behaviour of solutions to (5-1). Assuming the detailed balance condition (5-6), along a solution $\mathbf{c}(t) = \{c_i(t)\}_{i \geq 1}$ to (5-1) we have

$$\begin{aligned} \frac{d}{dt} H(\mathbf{c}(t)) &= -D_{\text{CF}}(\mathbf{c}(t)) \\ &:= -\frac{1}{2} \sum_{i,j=1}^{\infty} a_{i,j} Q_i Q_j \left(\frac{c_i c_j}{Q_i Q_j} - \frac{c_{i+j}}{Q_{i+j}} \right) \left(\log \frac{c_i c_j}{Q_i Q_j} - \log \frac{c_{i+j}}{Q_{i+j}} \right) \\ &\leq -\sum_{i=1}^{\infty} a_i Q_i \left(\frac{c_i c_1}{Q_i} - \frac{c_{i+1}}{Q_{i+1}} \right) \left(\log \frac{c_i c_1}{Q_i} - \log \frac{c_{i+1}}{Q_{i+1}} \right) = D(\mathbf{c}(t)) \leq 0 \end{aligned} \quad (5-8)$$

(see [Cañizo 2007] for a rigorous proof), where the a_i are defined by (5-3) for any $i \geq 1$. Hence the free energy is also a Lyapunov functional for (5-1), and it dissipates at a *faster* rate than for the Becker–Döring equations (since more types of reactions are allowed). As such, it is reasonable to think that the inequalities from Section 3 can be useful also in this case. This turns out to be true, and some improvements can be made on existing results. However, it also turns out that our results are not able to extend the range of possible coefficients for which convergence to a particular subcritical equilibrium can be proved; we cannot give any new results for coefficients such as (5-7) with $\gamma, \eta > 0$ and $\gamma + \eta \leq 1$.

Proof of Theorem 1.5. We now give the proof of our main result concerning the above model (5-1). One of the main obstacles in applying directly our results to equation (5-1) is that, unlike for the Becker–Döring equations, the moments of solutions to the general coagulation and fragmentation system are not known to be bounded; i.e., Proposition 4.1 is not available for (5-1). One can for example say the following about integer moments (this result can easily be extended to noninteger powers by interpolation, and was known from the early works in the topic [Carr and da Costa 1994; Carr 1992]). From this point onward we will assume that

$$a_{i,j} = i^\gamma j^\eta + i^\eta j^\gamma \quad \text{for } i, j \in \mathbb{N}, \quad (5-9)$$

with $\eta \leq \gamma$ and $0 \leq \lambda := \gamma + \eta \leq 1$.

Lemma 5.1. *Let $k \in \mathbb{N}$ and let $\mathbf{c} = \mathbf{c}(t) = \{c_i(t)\}_{i \in \mathbb{N}}$ be a solution with mass ϱ to the coagulation and fragmentation system (5-1) with coefficients satisfying (5-9). Then*

$$M_k(\mathbf{c}(t)) \leq \begin{cases} (M_k(\mathbf{c}(0)) + \frac{1-\lambda}{k-1} (2^k - 2) \varrho^{\frac{1-\gamma}{k-1}} t)^{\frac{k-1}{1-\lambda}} & \text{if } 0 < \lambda < 1, \\ M_k(\mathbf{c}(0)) \exp(2(2^k - 2) \varrho t) & \text{if } \lambda = 1, \end{cases} \quad (5-10)$$

where $M_p(\mathbf{c}(t)) := \sum_{i=1}^{\infty} i^p c_i(t)$ for any $p \geq 0$, $t \geq 0$.

Proof. We give a formal proof for completeness; a rigorous one can be obtained by standard approximation methods, and can be found in [Ball and Carr 1990]. To simplify the notation and since $c(t)$ is fixed, we define $M_j(t) = M_j(c(t))$ for any $j \geq 1, t \geq 0$. One can check the following weak formula for the integral of the right-hand side of (5-1) against a test sequence $\{\phi(i)\}_i$:

$$\sum_{i=1}^{\infty} \phi(i) \left(\frac{1}{2} \sum_{j=1}^{i-1} W_{i-j,j} - \sum_{j=1}^{\infty} W_{i,j} \right) = \frac{1}{2} \sum_{i,j} (\phi(i+j) - \phi(i) - \phi(j)) W_{i,j}.$$

Applying this to $\phi(i) := i^k$, neglecting the negative contribution of the fragmentation terms and using the binomial formula, one obtains

$$\frac{d}{dt} M_k(t) \leq \sum_{l=1}^{k-1} \binom{k}{l} M_{l+\gamma}(t) M_{k-l+\eta}(t) \quad \forall t \geq 0.$$

Next, we use the interpolation

$$M_{\delta}(t) \leq M_1^{\frac{k-\delta}{k-1}}(t) M_k^{\frac{\delta-1}{k-1}}(t),$$

where $1 < \delta < k$, to find that

$$M_{l+\gamma}(t) M_{k-l+\eta}(t) \leq M_1(t)^{\frac{k-\lambda}{k-1}} M_k(t)^{\frac{k+\lambda-2}{k-1}}.$$

Thus,

$$\frac{d}{dt} M_k(t) \leq (2^k - 2) \rho^{\frac{k-\lambda}{k-1}} M_k^{\frac{k+\lambda-2}{k-1}}(t) \quad \forall t \geq 0$$

and the result follows from this differential inequality. □

With the above at hand, we are now able to prove our main result about the rate of convergence to equilibrium in the general setting of coagulation and fragmentation equations:

Proof of Theorem 1.5. Assume for the moment that $a_{i,j}$ is of the form (5-7), in order to see why the proof only works for coefficients of the form (1-27).

Fix $\delta > 0$ such that $0 < \delta < \bar{z} < z_s - \delta$. We use the observation (5-8) that $D_{CF}(c(t)) \geq D(c(t))$ at all times $t \geq 0$, defining $\{a_i\}_{i \in \mathbb{N}}$ by (5-3). Using Theorem 1.1 (actually, its more detailed forms in equation (3-18) and Proposition 3.8), we obtain

$$\begin{aligned} \frac{d}{dt} H(c(t)|\mathcal{Q}) &= -D_{CF}(c(t)) \leq -D(c(t)) \\ &\leq \begin{cases} -CM_k(c(t))^{\frac{\gamma-1}{k-1}} H(c(t)|\mathcal{Q})^{\frac{k-\gamma}{k-1}} & \text{if } \delta < c_1(t) < z_s - \delta, \\ -CM_k(c(t))^{\frac{\gamma-1}{k-1}} & \text{if } c_1(t) < \delta \text{ or } c_1(t) \geq z_s - \delta \end{cases} \\ &\leq -C_0 M_k(c(t))^{\frac{\gamma-1}{k-1}} H(c(t)|\mathcal{Q})^{\frac{k-\gamma}{k-1}} \end{aligned}$$

for some constant $C_0 > 0$ that depends also on $H(c(0)|\mathcal{Q})$. Using Lemma 5.1 this implies

$$\frac{d}{dt} H(c(t)|\mathcal{Q}) \leq -\frac{C_0}{(M_k(c(0)) + \frac{1-\lambda}{k-1}(2^k - 2)\rho^{\frac{k-\lambda}{k-1}}t)^{\frac{1-\gamma}{1-\lambda}}} H(c(t)|\mathcal{Q})^{\frac{k-\gamma}{k-1}}, \quad t \geq 0.$$

This implies decay of $H(\mathbf{c}(t))$ only when $\lambda = \gamma$, that is, when $\eta = 0$ (since $\lambda = \gamma + \eta$). Solving the differential inequality yields the result. \square

Remark 5.2. The same decay rate was obtained in [Cañizo 2007] by means of the particular case of inequality (1-19) for $\beta = 2 - \gamma$. Here we obtain slightly different decay rates by assuming higher moments of the initial data $\mathbf{c}(0)$ are finite, but the method does not seem to give a better decay than a power of $\log t$ in any case.

Remark 5.3. It seems to the authors that the inequality we use in the proof of Theorem 1.5 is not optimal, and could be improved to deal with the case

$$a_{i,j} = i^\gamma j^\eta + i^\eta j^\gamma,$$

with a resulting convergence rate that would depend on $\lambda = \gamma + \eta$.

Appendix A: Additional computations for the theory of the discrete log-Sobolev inequality with weights

We have collected here technical lemmas from Section 2 that we felt would have encumbered it.

Lemma A.1. *For any sequence \mathbf{f} , we have*

$$\text{Ent}_\mu(\mathbf{f}^2) \leq \mathcal{L}(\mathbf{f}) \leq \text{Ent}_\mu(\mathbf{f}^2) + 2 \sum_{i=1}^{\infty} \mu_i f_i^2.$$

Proof. From the definition of \mathcal{L} , the inequality

$$\text{Ent}_\mu(\mathbf{f}^2) \leq \mathcal{L}(\mathbf{f})$$

is trivial. We thus consider the right-hand side inequality. For a given sequence \mathbf{f} and any $\alpha \in \mathbb{R}$ we define

$$\begin{aligned} G_\alpha(t) &= \sum_{i=1}^{\infty} \mu_i (t f_i + \alpha)^2 \log \left(\frac{(t f_i + \alpha)^2}{\sum_{i=1}^{\infty} \mu_i (t f_i + \alpha)^2} \right) \\ &= 2 \sum_{i=1}^{\infty} \mu_i (t f_i + \alpha)^2 \log |t f_i + \alpha| - \left(\sum_{i=1}^{\infty} \mu_i (t f_i + \alpha)^2 \right) \log \left(\sum_{i=1}^{\infty} \mu_i (t f_i + \alpha)^2 \right), \end{aligned}$$

and notice that

$$G_0(t) = t^2 \text{Ent}_\mu(\mathbf{f}^2).$$

Next, we define $g(t) = G_0(t) + 2t^2 \sum_{i=1}^{\infty} \mu_i f_i^2$ and notice that the inequality we want to prove is equivalent to

$$G_\alpha(1) \leq g(1)$$

for any $\alpha \in \mathbb{R}$. Clearly $G_\alpha(t) \leq g(t)$ when $t = 0$. Differentiating G we find that

$$\begin{aligned} G'_\alpha(t) &= 4 \sum_{i=1}^\infty \mu_i f_i |tf_i + \alpha| \log(tf_i + \alpha) + 2 \sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha) \\ &\quad - 2 \left(\sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha) \right) \log \left(\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2 \right) - 2 \sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha) \\ &= 4 \sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha) \log|tf_i + \alpha| - 2 \left(\sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha) \right) \log \left(\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2 \right), \end{aligned}$$

which satisfies $G'_\alpha(0) = 0$ for any f and α , implying that $G'_\alpha(0) = g'(0) = 0$. As G is defined for any $t \in [0, 1]$ we see that it is enough to show that when defined,

$$G''_\alpha(t) \leq g''(t)$$

for any α . Indeed,

$$\begin{aligned} G''_\alpha(t) &= 4 \sum_{i=1}^\infty \mu_i f_i^2 \log|tf_i + \alpha| + 4 \sum_{i=1}^\infty \mu_i f_i^2 - 2 \sum_{i=1}^\infty \mu_i f_i^2 \log \left(\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2 \right) \\ &\quad - 4 \frac{(\sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha))^2}{\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2} \\ &= 2 \sum_{i=1}^\infty \mu_i f_i^2 \log \left(\frac{(tf_i + \alpha)^2}{\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2} \right) + 4 \sum_{i=1}^\infty \mu_i f_i^2 - 4 \frac{(\sum_{i=1}^\infty \mu_i f_i (tf_i + \alpha))^2}{\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2}. \end{aligned}$$

As

$$\text{Ent}_\mu(f^2) = \sup \left\{ \sum_{i=1}^\infty \mu_i f_i^2 \log h_i : \sum_{i=1}^\infty \mu_i h_i = 1 \right\},$$

we see that by choosing

$$h_i = \frac{(tf_i + \alpha)^2}{\sum_{i=1}^\infty \mu_i (tf_i + \alpha)^2}$$

we get

$$G''_\alpha(t) \leq 2 \text{Ent}_\mu(f^2) + 4 \sum_{i=1}^\infty \mu_i f_i^2 = g''(t). \quad \square$$

Lemma A.2. For all $f \in L_\Phi$, we have

$$\|f\|_{L^1_\mu} \leq \|f\|_{L^2_\mu} \leq \sqrt{\frac{3}{2}} \|f\|_{L_\Phi}. \tag{A-1}$$

Proof. The inequality

$$\|f\|_{L^1_\mu} \leq \|f\|_{L^2_\mu}$$

is immediate as μ is a probability measure. To show the last inequality we may assume that $\|f\|_{L_\Phi} = 1$. Due to Fatou's lemma we know that if $k_n \xrightarrow{n \rightarrow \infty} k > 0$ then

$$\sum_{i=1}^{\infty} \mu_i \Phi\left(\frac{|f_i|}{k}\right) \leq \liminf_{n \rightarrow \infty} \sum_{i=1}^{\infty} \mu_i \Phi\left(\frac{|f_i|}{k_n}\right),$$

implying that if $\|f\|_{L_\Phi} > 0$ then

$$\sum_{i=1}^{\infty} \mu_i \Phi\left(\frac{|f_i|}{\|f\|_{L_\Phi}}\right) \leq 1.$$

In our case, since $\Psi(x)$ is convex we find that

$$1 \geq \sum_{i=1}^{\infty} \mu_i \Phi(f_i) = \sum_{i=1}^{\infty} \mu_i \Psi(f_i^2) \geq \Psi\left(\sum_{i=1}^{\infty} \mu_i f_i^2\right) = \Psi(\|f\|_{L_\mu}^2).$$

As Ψ is increasing and $\Psi(\frac{3}{2}) > 1$ we conclude that

$$\|f\|_{L_\mu}^2 < \frac{3}{2},$$

yielding the desired result. □

Lemma A.3. *Let $f \in L_\Phi$. Then*

$$\|f - \langle f \rangle\|_{L_\mu}^2 = \frac{1}{2} \lim_{|a| \rightarrow \infty} \text{Ent}_\mu((f + a)^2). \quad (\text{A-2})$$

Proof. We start by noticing that

$$\text{Ent}_\mu((f + a)^2) = \sum_{i=1}^{\infty} \mu_i (f_i^2 + 2af_i + a^2) \log\left(\frac{(1 + f_i/a)^2}{\sum_{i=1}^{\infty} \mu_i (1 + f_i/a)^2}\right),$$

and continue by assuming that f_i is uniformly bounded, from which the result will follow with an application of an appropriate convergence theorem. There exists a_0 such that if $|a| > |a_0|$ we have that $|f_i/a| < \frac{1}{2}$ uniformly in i . As on $[-\frac{1}{2}, \frac{1}{2}]$, we have that there exists $C > 0$ such that

$$\left|\log(1 + x) - x + \frac{1}{2}x^2\right| \leq Cx^3.$$

We conclude that

$$\log\left(1 + 2\frac{f_i}{a} + \frac{f_i^2}{a^2}\right) = \left(2\frac{f_i}{a} + \frac{f_i^2}{a^2}\right) - 2\frac{f_i^2}{a^2} + \frac{E_{1,i}}{a^3} = 2\frac{f_i}{a} - \frac{f_i^2}{a^2} + \frac{E_{1,i}}{a^3}$$

and

$$\log\left(1 + 2\frac{\langle f \rangle}{a} + \frac{\|f\|_{L_\mu}^2}{a^2}\right) = 2\frac{\langle f \rangle}{a} + \frac{\|f\|_{L_\mu}^2}{a^2} - 2\frac{\langle f \rangle^2}{a^2} + \frac{E_{2,i}}{a^3},$$

where $E_{1,i}, E_{2,i}$ are uniformly bounded in i . This implies

$$\begin{aligned} \text{Ent}_\mu((f + a)^2) &= \sum_{i=1}^\infty \mu_i (f_i^2 + 2af_i + a^2) \left(2\frac{f_i}{a} - 2\frac{\langle f \rangle}{a} - \frac{f_i^2}{a^2} - \frac{\|f\|_{L_\mu^2}^2}{a^2} + 2\frac{\langle f \rangle^2}{a^2} \right) \\ &\quad + \frac{1}{a} \sum_{i=1}^\infty \mu_i \left(1 + 2\frac{f_i}{a} + \frac{f_i^2}{a^2} \right) (E_{1,i} - E_{2,i}). \end{aligned}$$

The last term clearly goes to zero as $|a|$ goes to infinity, so we are only left to deal with the first expression.

$$\begin{aligned} \sum_{i=1}^\infty \mu_i (f_i^2 + 2af_i + a^2) &\left(2\frac{f_i}{a} - 2\frac{\langle f \rangle}{a} - \frac{f_i^2}{a^2} - \frac{\|f\|_{L_\mu^2}^2}{a^2} + 2\frac{\langle f \rangle^2}{a^2} \right) \\ &= 4\|f\|_{L_\mu^2}^2 - 4\langle f \rangle^2 + 2a\langle f \rangle - 2a\langle f \rangle - \|f\|_{L_\mu^2}^2 - \|f\|_{L_\mu^2}^2 + 2\langle f \rangle^2 + \frac{E_3}{a} \\ &= 2(\|f\|_{L_\mu^2}^2 - \langle f \rangle^2) + \frac{E_3}{a}. \end{aligned}$$

This completes the proof as $\|f - \langle f \rangle\|_{L_\mu^2}^2 = \|f\|_{L_\mu^2}^2 - \langle f \rangle^2$. □

Lemma A.4. *Let f be a sequence such that $f_m = 0$ for some $m \in \mathbb{N}$. Set by $f^{(0)} = f \mathbb{1}_{i < m}$ and $f^{(1)} = f \mathbb{1}_{i > m}$. Then*

$$\begin{aligned} \|\langle f^{(0)} \rangle\|_{L_\Phi} &\leq |\langle f^{(0)} \rangle| \leq \|f^{(0)}\|_{L_\mu^2} \left(\sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}}, \\ \|\langle f^{(1)} \rangle\|_{L_\Phi} &\leq |\langle f^{(1)} \rangle| \leq \|f^{(1)}\|_{L_\mu^2} \left(\sum_{i=m+1}^\infty \mu_i \right)^{\frac{1}{2}}. \end{aligned} \tag{A-3}$$

Proof. We start by noticing that for any constant sequence $f = \alpha$ one has

$$\|\alpha\|_{L_\Phi} = \inf_{k>0} \left\{ \sum_{i=1}^\infty \mu_i \Phi\left(\frac{|\alpha|}{k}\right) \leq 1 \right\} = \inf_{k>0} \left\{ \Phi\left(\frac{|\alpha|}{k}\right) \leq 1 \right\} = \frac{|\alpha|}{\Phi^{-1}(1)} \leq |\alpha|,$$

as long as $\Phi(1) < 1$, which is valid in our case. Next we notice that

$$|\langle f^{(0)} \rangle| \leq \sum_{i=1}^{m-1} \mu_i |f_i| \leq \left(\sum_{i=1}^{m-1} \mu_i f_i^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}} = \|f^{(0)}\|_{L_\mu^2} \left(\sum_{i=1}^{m-1} \mu_i \right)^{\frac{1}{2}}.$$

This yields the first inequality and similar arguments yield the second inequality. □

Remark A.5. As was shown in the proof of Lemma A.4, one can actually improve the bounds in (A-3) by a factor of $\Psi^{-1}(1)$.

Lemma A.6. *For any $t \geq \frac{3}{2}$ one has that*

$$\frac{1}{3} \frac{t}{\log t} \leq \Psi^{-1}(t) \leq 2 \frac{t}{\log t}. \tag{A-4}$$

Proof. We start by noticing that

$$\Psi\left(\frac{1}{3} \frac{t}{\log t}\right) = \frac{1}{3} \frac{t}{\log t} \log\left(1 + \frac{1}{3} \frac{t}{\log t}\right) \leq \frac{1}{3} \frac{t}{\log t} \log\left(1 + \frac{t}{\log\left(\frac{27}{8}\right)}\right) \leq \frac{1}{3} \frac{t}{\log t} \log(1+t).$$

Thus, one notices that if

$$1+t \leq t^3$$

when $t \geq \frac{3}{2}$, we have

$$\Psi\left(\frac{1}{3} \frac{t}{\log t}\right) \leq t,$$

yielding the left-hand side of (A-4). This is indeed the case as $g(t) = t^3 - t - 1$ is increasing on $[1/\sqrt{3}, \infty)$ and $g(\frac{3}{2}) > 0$.

For the converse we notice that

$$\Psi\left(2 \frac{t}{\log t}\right) = 2 \frac{t}{\log t} \log\left(1 + 2 \frac{t}{\log t}\right) \geq t$$

if and only if

$$1 + 2 \frac{t}{\log t} \geq \sqrt{t}.$$

Considering the function $g(x) = x/\log x$ for $x > 1$, we see that it obtains a minimum at $x = e$. Thus, for any $x > 1$, we have $g(x) \geq e > 1$. We conclude that for $t > \frac{3}{2}$,

$$2 \frac{t}{\log t} = \sqrt{t} g(\sqrt{t}) \geq \sqrt{t},$$

showing the desired result. □

Appendix B: Additional useful computations

Lemma B.1. *For given coagulation and detailed balance coefficients $\{a_i\}_{i \in \mathbb{N}}$ and $\{Q_i\}_{i \in \mathbb{N}}$, and a given positive sequence \mathbf{c} with finite mass ϱ , we have, for any $z > 0$,*

$$H(\mathbf{c}|\mathcal{Q}) \leq H(\mathbf{c}|\mathcal{Q}_z),$$

where $\mathcal{Q} = \mathcal{Q}_{\bar{z}}$.

Proof. We have

$$H(\mathbf{c}|\mathcal{Q}_z) = \sum_{i=1}^{\infty} c_i \left(\log\left(\frac{c_1}{Q_i z^i}\right) - 1 \right) + \sum_{i=1}^{\infty} Q_i z^i$$

implying

$$H(\mathbf{c}|\mathcal{Q}_{z_1}) - H(\mathbf{c}|\mathcal{Q}_{z_2}) = \sum_{i=1}^{\infty} i c_i \log\left(\frac{z_2}{z_1}\right) + \sum_{i=1}^{\infty} Q_i (z_1^i - z_2^i).$$

In particular, if $z_2 = \bar{z}$ we have, for any $z > 0$,

$$H(\mathbf{c}|\mathcal{Q}_z) = H(\mathbf{c}|\mathcal{Q}) + \varrho \log\left(\frac{\bar{z}}{z}\right) + \sum_{i=1}^{\infty} Q_i (z^i - \bar{z}^i)$$

$$\begin{aligned}
 &= H(\mathbf{c}|\mathcal{Q}) + \sum_{i=1}^{\infty} i Q_i \bar{z}^i \log\left(\frac{\bar{z}}{z}\right) + \sum_{i=1}^{\infty} Q_i z^i \left(1 - \left(\frac{\bar{z}}{z}\right)^i\right) \\
 &= H(\mathbf{c}|\mathcal{Q}) + \sum_{i=1}^{\infty} Q_i z^i \left(\left(\frac{\bar{z}}{z}\right)^i \log\left(\left(\frac{\bar{z}}{z}\right)^i\right) - \left(\frac{\bar{z}}{z}\right)^i + 1\right) \\
 &= H(\mathbf{c}|\mathcal{Q}) + \sum_{i=1}^{\infty} Q_i z^i \Lambda\left(\frac{(Q_z)i}{Q_i}\right),
 \end{aligned}$$

where $\Lambda(x) = x \log x - x + 1 > 0$ when $x > 0$. □

Lemma B.2. *Let $\{Q_i\}_{i \in \mathbb{N}}$ be a nonnegative sequence such that $\lim_{i \rightarrow \infty} Q_{i+1}/Q_i = 1/r$ for some $r > 0$. Assume that $0 < x < r_1 < r$. Then*

$$\sum_{i=i_0+1}^{\infty} i Q_i x^{i-1} \leq C Q_{i_0} x^{i_0},$$

where C is a constant depending only on $\{Q_i\}_{i \in \mathbb{N}}$ and r_1 .

Proof. Define $\beta_i = Q_{i+1}/Q_i$. We have that $\lim_{i \rightarrow \infty} \beta_i = 1/r$, and as such we can find $l \in \mathbb{N}$ such that for all $i > l$

$$\Lambda_1 = \sup_{i>l} \beta_i < \frac{1}{r_1}.$$

Let $\Lambda_2 = \sup_{i \leq l} \beta_i$. Since for any $i > i_0$

$$Q_i = \left(\prod_{j=i_0}^{i-1} \beta_j\right) Q_{i_0},$$

we see that

$$\begin{aligned}
 \sum_{i=i_0+1}^{\infty} i Q_i x^{i-1} &= Q_{i_0} x^{i_0} \sum_{i=i_0+1}^{\infty} i \left(\prod_{j=i_0}^{i-1} \beta_j\right) x^{i-i_0-1} \\
 &\leq Q_{i_0} x^{i_0} \left(\Lambda_2 \sum_{j=0}^{l-i_0} i (\Lambda_2 r_1)^j + \Lambda_1 \sum_{j=l+1-i_0}^{\infty} i (\Lambda_1 r_1)^j\right) \\
 &\leq Q_{i_0} x^{i_0} \left(\Lambda_2 \sum_{j=0}^l j (\Lambda_2 r_1)^j + \Lambda_1 \sum_{j=0}^{\infty} j (\Lambda_1 r_1)^j\right),
 \end{aligned}$$

completing the proof as l, Λ_1 and Λ_2 depend solely on $\{Q_i\}_{i \in \mathbb{N}}$. □

Lemma B.3. *Let $\varepsilon > 0$ and $\gamma > 0$. Define*

$$B_{\varepsilon,\gamma} = \sum_{i=1}^{\infty} i^\gamma e^{-\varepsilon i}.$$

Then $\varepsilon^{1+\gamma} B_{\varepsilon,\gamma}$ is of order 1 when ε goes to zero.

Proof. We start by noticing that the function $g_{\varepsilon,\gamma}(x) = x^\gamma e^{-\varepsilon x}$ is increasing in $[0, \frac{\gamma}{\varepsilon}]$ and decreasing in $[\frac{\gamma}{\varepsilon}, \infty)$. As such

$$B_{\varepsilon,\gamma} \geq \sum_{i=[\frac{\gamma}{\varepsilon}]+1}^{\infty} i^\gamma e^{-\varepsilon i} \geq \int_{[\frac{\gamma}{\varepsilon}]+1}^{\infty} x^\gamma e^{-\varepsilon x} dx = \varepsilon^{-(1+\gamma)} \int_{\varepsilon([\frac{\gamma}{\varepsilon}]+1)}^{\infty} y^\gamma e^{-y} dy \geq \varepsilon^{-(1+\gamma)} \int_{\varepsilon}^{\infty} y^\gamma e^{-y} dy,$$

showing the lower bound. For the upper bound we notice that

$$B_{\varepsilon,\gamma} \leq \sup_{x \geq 0} g_{\varepsilon,\gamma}(x) \sum_{i=1}^{\infty} e^{-\frac{\varepsilon}{2}i} = \left(\frac{2\gamma}{\varepsilon}\right)^\gamma e^{-\gamma} \frac{e^{-\frac{\varepsilon}{2}}}{1 - e^{-\frac{\varepsilon}{2}}},$$

which completes the proof since

$$\sup_{\varepsilon > 0} \frac{\varepsilon e^{-\frac{\varepsilon}{2}}}{1 - e^{-\frac{\varepsilon}{2}}} < +\infty. \quad \square$$

References

- [Ball and Carr 1988] J. M. Ball and J. Carr, “Asymptotic behaviour of solutions to the Becker–Döring equations for arbitrary initial data”, *Proc. Roy. Soc. Edinburgh Sect. A* **108**:1-2 (1988), 109–116. [MR](#) [Zbl](#)
- [Ball and Carr 1990] J. M. Ball and J. Carr, “The discrete coagulation-fragmentation equations: existence, uniqueness, and density conservation”, *J. Statist. Phys.* **61**:1-2 (1990), 203–234. [MR](#) [Zbl](#)
- [Ball et al. 1986] J. M. Ball, J. Carr, and O. Penrose, “The Becker–Döring cluster equations: basic properties and asymptotic behaviour of solutions”, *Comm. Math. Phys.* **104**:4 (1986), 657–692. [MR](#) [Zbl](#)
- [Barthe and Roberto 2003] F. Barthe and C. Roberto, “Sobolev inequalities for probability measures on the real line”, *Studia Math.* **159**:3 (2003), 481–497. [MR](#) [Zbl](#)
- [Becker and Döring 1935] R. Becker and W. Döring, “Kinetische Behandlung der Keimbildung in übersättigten Dämpfen”, *Ann. Phys.* **416**:8 (1935), 719–752. [Zbl](#)
- [Bobkov and Götze 1999] S. G. Bobkov and F. Götze, “Exponential integrability and transportation cost related to logarithmic Sobolev inequalities”, *J. Funct. Anal.* **163**:1 (1999), 1–28. [MR](#) [Zbl](#)
- [Burton 1977] J. J. Burton, “Nucleation theory”, pp. 195–234 in *Statistical mechanics, Part A: Equilibrium techniques*, edited by B. J. Berne, Modern Theoretical Chemistry **5**, Springer, 1977.
- [Cañizo 2005] J. A. Cañizo, “Asymptotic behaviour of solutions to the generalized Becker–Döring equations for general initial data”, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **461**:2064 (2005), 3731–3745. [MR](#) [Zbl](#)
- [Cañizo 2007] J. A. Cañizo, “Convergence to equilibrium for the discrete coagulation-fragmentation equations with detailed balance”, *J. Stat. Phys.* **129**:1 (2007), 1–26. [MR](#) [Zbl](#)
- [Cañizo and Lods 2013] J. A. Cañizo and B. Lods, “Exponential convergence to equilibrium for subcritical solutions of the Becker–Döring equations”, *J. Differential Equations* **255**:5 (2013), 905–950. [MR](#) [Zbl](#)
- [Cañizo et al. 2017] J. A. Cañizo, A. Einav, and B. Lods, “Uniform moment propagation for the Becker–Döring equation”, preprint, 2017. [arXiv](#)
- [Capasso 2003] V. Capasso (editor), *Mathematical modelling for polymer processing: polymerization, crystallization, manufacturing*, Mathematics in Industry **2**, Springer, 2003. [MR](#) [Zbl](#)
- [Carr 1992] J. Carr, “Asymptotic behaviour of solutions to the coagulation-fragmentation equations, I: The strong fragmentation case”, *Proc. Roy. Soc. Edinburgh Sect. A* **121**:3-4 (1992), 231–244. [MR](#) [Zbl](#)
- [Carr and da Costa 1994] J. Carr and F. P. da Costa, “Asymptotic behavior of solutions to the coagulation-fragmentation equations, II: Weak fragmentation”, *J. Statist. Phys.* **77**:1-2 (1994), 89–123. [MR](#) [Zbl](#)
- [Collet et al. 2002] J.-F. Collet, T. Goudon, F. Poupaud, and A. Vasseur, “The Becker–Döring system and its Lifshitz–Slyozov limit”, *SIAM J. Appl. Math.* **62**:5 (2002), 1488–1500. [MR](#) [Zbl](#)

- [da Costa 1998] F. P. da Costa, “Asymptotic behaviour of low density solutions to the generalized Becker–Döring equations”, *NoDEA Nonlinear Differential Equations Appl.* **5**:1 (1998), 23–37. [MR](#) [Zbl](#)
- [Desvillettes et al. 2011] L. Desvillettes, C. Mouhot, and C. Villani, “Celebrating Cercignani’s conjecture for the Boltzmann equation”, *Kinet. Relat. Models* **4**:1 (2011), 277–294. [MR](#) [Zbl](#)
- [Farjoun and Neu 2008] Y. Farjoun and J. C. Neu, “Exhaustion of nucleation in a closed system”, *Phys. Rev. E* **78**:5 (2008), art. id. 051402.
- [Farjoun and Neu 2011] Y. Farjoun and J. C. Neu, “Aggregation according to classical kinetics: from nucleation to coarsening”, *Phys. Rev. E* **83**:5 (2011), art. id. 051607.
- [Jabin and Niethammer 2003] P.-E. Jabin and B. Niethammer, “On the rate of convergence to equilibrium in the Becker–Döring equations”, *J. Differential Equations* **191**:2 (2003), 518–543. [MR](#) [Zbl](#)
- [Kelton et al. 1983] K. F. Kelton, A. L. Greer, and C. V. Thompson, “Transient nucleation in condensed systems”, *J. Chem. Phys.* **79**:12 (1983), 6261–6276.
- [Murray and Pego 2015] R. W. Murray and R. L. Pego, “Polynomial decay to equilibrium for the Becker–Döring equations”, preprint, 2015. [arXiv](#)
- [Neu et al. 2002] J. C. Neu, J. A. Cañizo, and L. L. Bonilla, “Three eras of micellization”, *Phys. Rev. E* **66**:6 (2002), art. id. 061406.
- [Neu et al. 2005] J. C. Neu, L. L. Bonilla, and A. Carpio, “Igniting homogeneous nucleation”, *Phys. Rev. E* (3) **71**:2 (2005), art. id. 021601. [MR](#)
- [Niethammer 2003] B. Niethammer, “On the evolution of large clusters in the Becker–Döring model”, *J. Nonlinear Sci.* **13**:1 (2003), 115–155. [MR](#) [Zbl](#)
- [Oxtoby 1992] D. W. Oxtoby, “Homogeneous nucleation: theory and experiment”, *J. Phys. Cond. Matter* **4**:38 (1992), 7620–7650.
- [Penrose 1989] O. Penrose, “Metastable states for the Becker–Döring cluster equations”, *Comm. Math. Phys.* **124**:4 (1989), 515–541. [MR](#) [Zbl](#)
- [Penrose 1997] O. Penrose, “The Becker–Döring equations at large times and their connection with the LSW theory of coarsening”, *J. Statist. Phys.* **89**:1-2 (1997), 305–320. [MR](#) [Zbl](#)
- [Penrose and Lebowitz 1979] O. Penrose and J. L. Lebowitz, “Towards a rigorous molecular theory of metastability”, pp. 293–340 in *Fluctuation phenomena*, edited by E. Montroll and J. L. Lebowitz, Studies in Statistical Mechanics 7, North-Holland, New York, 1979.
- [Rothaus 1985] O. S. Rothaus, “Analytic inequalities, isoperimetric inequalities and logarithmic Sobolev inequalities”, *J. Funct. Anal.* **64**:2 (1985), 296–313. [MR](#) [Zbl](#)
- [Schmelzer 2005] J. Schmelzer (editor), *Nucleation theory and applications*, Wiley-VCH, Weinheim, 2005.
- [Velázquez 1998] J. J. L. Velázquez, “The Becker–Döring equations and the Lifshitz–Slyozov theory of coarsening”, *J. Statist. Phys.* **92**:1-2 (1998), 195–236. [MR](#) [Zbl](#)
- [Villani 2003] C. Villani, “Cercignani’s conjecture is sometimes true and always almost true”, *Comm. Math. Phys.* **234**:3 (2003), 455–490. [MR](#) [Zbl](#)
- [Xiao and Haasen 1991] S. Q. Xiao and P. Haasen, “HREM investigation of homogeneous decomposition in a ni-12 at. % al alloy”, *Acta. Metall. Mater.* **39**:4 (1991), 651–659.

Received 24 Dec 2016. Revised 28 Mar 2017. Accepted 29 May 2017.

JOSÉ A. CAÑIZO: canizo@ugr.es

Departamento de Matemática Aplicada, Universidad de Granada, Av. Fuentenueva S/N, 18071 Granada, Spain

AMIT EINAV: a.einav@dpms.cam.ac.uk

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Cambridge, CB3 0WB, United Kingdom

BERTRAND LODS: bertrand.lods@unito.it

Department of Economics and Statistics & Collegio Carlo Alberto, Università degli Studi di Torino,

Corso Unione Sovietica, 218/bis, 10134 Torino, Italy

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2017 is US \$265/year for the electronic version, and \$470/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 10 No. 7 2017

A vector field method for relativistic transport equations with applications DAVID FAJMAN, JÉRÉMIE JOUDIOUX and JACQUES SMULEVICI	1539
Analytic hypoellipticity for sums of squares and the Treves conjecture, II ANTONIO BOVE and MARCO MUGHETTI	1613
Pliability, or the Whitney extension theorem for curves in Carnot groups NICOLAS JUILLET and MARIO SIGALOTTI	1637
Trend to equilibrium for the Becker–Döring equations: an analogue of Cercignani’s conjecture JOSÉ A. CAÑIZO, AMIT EINAV and BERTRAND LODS	1663
The A_∞ -property of the Kolmogorov measure KAJ NYSTRÖM	1709
L^2 -Betti numbers of rigid C^* -tensor categories and discrete quantum groups DAVID KYED, SVEN RAUM, STEFAAN VAES and MATTHIAS VALVEKENS	1757