

# ANALYSIS & PDE

Volume 11

No. 8

2018

FABRICIO MACIÀ AND GABRIEL RIVIÈRE

**TWO-MICROLOCAL REGULARITY OF QUASIMODES ON THE  
TORUS**



# TWO-MICROLOCAL REGULARITY OF QUASIMODES ON THE TORUS

FABRICIO MACIÀ AND GABRIEL RIVIÈRE

We study the regularity of stationary and time-dependent solutions to strong perturbations of the free Schrödinger equation on two-dimensional flat tori. This is achieved by performing a second microlocalization related to the size of the perturbation and by analyzing concentration and nonconcentration properties at this new scale. In particular, we show that sufficiently accurate quasimodes can only concentrate on the set of critical points of the average of the potential along closed geodesics.

## 1. Introduction

The high-frequency analysis of eigenfunctions of elliptic operators on a compact Riemannian manifold has been the subject of intensive study in the past fifty years. To this day, many questions remain open, even in the simplest cases. Here we focus on eigenfunctions of Schrödinger operators on  $\mathbb{T}^d := \mathbb{R}^d / \mathbb{Z}^d$ , the standard torus endowed with its canonical metric. Eigenfunctions of a Schrödinger operator on  $\mathbb{T}^d$  are the solutions to the equation

$$-\Delta u_\lambda(x) + V(x) u_\lambda(x) = \lambda^2 u_\lambda(x), \quad x \in \mathbb{T}^d, \quad \|u_\lambda\|_{L^2(\mathbb{T}^d)} = 1, \quad (1)$$

where the potential  $V$  is real-valued and essentially bounded. In the free case  $V = 0$ , a straightforward computation shows that eigenfunctions of eigenvalue  $\lambda^2$  are linear combinations of complex exponentials  $e^{2i\pi k \cdot x}$  with frequencies  $k \in \mathbb{Z}^d$  lying on a circle of radius  $\lambda/(2\pi) > 0$  centered at the origin. However, extracting from this exact representation formula an asymptotic description of eigenfunctions in the high-frequency limit  $\lambda \rightarrow +\infty$  is a hard problem, due to the fact that multiplicities of large eigenvalues can also be very big. Instead, one can try to describe particular features of high-frequency eigenfunctions, such as formation of (asymptotic) singularities.

A natural way to quantify these singularities is through the scale of  $L^p$  spaces. This has been a classical topic in harmonic analysis, that originates with the seminal result of [Zygmund 1974] showing that, for  $d = 2$  and in the free case, there exists some universal constant  $C$  such that any solution  $u_\lambda$  of (1) satisfies  $\|u_\lambda\|_{L^4(\mathbb{T}^2)} \leq C$ . Later on, Bourgain [1993] conjectured that, again for the free case and when  $d \geq 3$ , one must have  $\|u_\lambda\|_{L^{2d/(d-2)}(\mathbb{T}^d)} \leq C_\delta \lambda^\delta$  for every  $\delta > 0$ . We refer the reader to [Bourgain 2013; Bourgain and Demeter 2015] for recent progress towards this conjecture. Note that the problem of showing the existence of an index  $p > 2$  such that  $\|u_\lambda\|_{L^p(\mathbb{T}^d)}$  is uniformly bounded remains open for  $d \geq 3$ .

---

Macià takes part in the visiting faculty program of ICMAT and is partially supported by ERC Starting Grant 277778 and the grants MTM2013-41780-P and MTM2017-85934-C3-3-P (MECD). Rivière is partially supported by the Agence Nationale de la Recherche through the Labex CEMPI (ANR-11-LABX-0007-01) and the ANR project GeRaSic (ANR-13-BS01-0007-01).

*MSC2010:* 58J51, 35P20, 35Q41, 58J50.

*Keywords:* quasimodes, Schrödinger operator, semiclassical measures, time-dependent Schrödinger equation.

There are alternative ways to describe the asymptotic structure of the solutions of (1). For instance, notice that a direct corollary of Zygmund's result is that, in the free case, any accumulation point of the sequence of probability measures,

$$\nu_\lambda(dx) = |u_\lambda(x)|^2 dx,$$

is a probability measure which is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{T}^2$  (it has in fact an  $L^2$  density). This result was refined by Jakobson [1997] who showed that the density has to be a trigonometric polynomial whose frequencies enjoy certain geometric constraints. It is natural to try to understand what happens when  $d \geq 3$ , where no analogue to Zygmund's result is known to hold, or when the Laplacian is perturbed by a lower-order term, such as a potential. Note that the problem of identifying accumulation points of sequences of moduli squares of eigenfunctions has a long history and it is connected to fundamental questions in quantum mechanics.

In dimension  $d \geq 3$  and for  $V = 0$ , Bourgain proved that any accumulation point has to be absolutely continuous even if we do not know a priori that the  $L^p$  norms of eigenfunctions are uniformly bounded for small  $p > 2$ ; this result was reported in [Jakobson 1997]. In the same reference, Jakobson obtained partial results on the structure of the densities of accumulation points. These results are based on harmonic analysis techniques and arguments on the geometry of lattice points. Absolute continuity of accumulation points also holds in the case of a nonzero potential  $V \in L^\infty(\mathbb{T}^d)$ , as was proved by Anantharaman and the first author [Anantharaman and Macià 2014]. The proof of that result is based on methods from semiclassical analysis for the time-dependent Schrödinger equation that were introduced for the particular case  $d = 2$  in [Macià 2010]. In fact, the results in [Anantharaman and Macià 2014] apply to the more general problem

$$\widehat{P}_\epsilon(\hbar)u_\hbar = \frac{1}{2}u_\hbar + o(\hbar\epsilon_\hbar), \quad \|u_\hbar\|_{L^2(\mathbb{T}^d)} = 1, \quad (2)$$

where  $\hbar \rightarrow 0^+$  is some semiclassical parameter, and where

$$\widehat{P}_\epsilon(\hbar) := -\frac{1}{2}\hbar^2\Delta + \epsilon_\hbar^2V, \quad (3)$$

with  $0 \leq \epsilon_\hbar \leq \hbar$  for  $\hbar$  small enough.<sup>1</sup> Among the main ingredients used in this approach are the two-microlocal techniques developed in [Nier 1996; Miller 1996; Fermanian-Kammerer 2000; 2005; Fermanian-Kammerer and Gérard 2002] in a different context. The results in [Anantharaman and Macià 2014] were further extended to treat the case of more general completely integrable systems in [Anantharaman et al. 2015]. This approach can also be used in order to analyze the Schrödinger equation on the planar disk [Anantharaman et al. 2016a; 2016b]. Note that studying the regularity of the solutions to (2) is also related to problems arising in control theory, as was shown by Burq and Zworski [2004]. We refer the reader to [Anantharaman and Léautaud 2014; Anantharaman et al. 2016b; Anantharaman and Macià 2014; Bourgain et al. 2013; Burq and Zworski 2004; 2012; Macià 2011] for perspectives from the point of view of control theory.

A different but related approach consists in studying the wavefront set  $\text{WF}_\hbar(u_\hbar)$  of solutions to (2). This was done in a series of works by Wunsch [2008; 2012] and Vasy and Wunsch [2009] dealing

<sup>1</sup>Note that, when  $\hbar = \epsilon_\hbar = \lambda^{-1}$ , equation (2) is essentially equation (1).

with completely integrable systems in dimension  $d = 2$ . In these articles, the authors investigated the properties of the semiclassical wavefront set  $\text{WF}_h(u_h)$  of solutions to (2) when  $0 \leq \epsilon_h \leq \hbar^{1+\delta}$  with  $\delta > 0$ . By proving some propagation of second microlocal wavefront sets, they showed that  $\text{WF}_h(u_h)$  cannot be reduced to a single geodesic and has to fill a Lagrangian torus—see for instance [Wunsch 2008, Theorem B; 2012, Theorem 3]. Note that, as in [Anantharaman et al. 2015], the results of Vasy and Wunsch hold for general classes of nondegenerate completely integrable systems. Under the assumption that  $\hbar^{1-\delta} \ll \epsilon_h \ll 1$ , Wunsch also exhibited examples of quasimodes of order  $\mathcal{O}(\hbar^\infty)$  for the operator  $\widehat{P}_\epsilon(\hbar)$  which concentrate on closed geodesics. This result was reported in [Anantharaman et al. 2015, Section 5.3], and it shows that  $\epsilon_h = \hbar$  is the critical size for which one can expect to have singular concentration phenomena for perturbations of the free semiclassical Schrödinger operator  $-\frac{1}{2}\hbar^2\Delta$ . In particular, for stronger perturbation  $\epsilon_h \gg \hbar$ , one cannot expect to have uniform bounds for  $L^p$  norms even for a small range of  $p$ . A notable feature of Wunsch’s construction is that the singularity is located on critical points of the potential  $V$  restricted to certain closed geodesics. In some sense, this type of singularity is similar to the ones that may occur in the case of Zoll manifolds [Macià and Rivièrè 2016; 2017]. Motivated by this observation, we will combine the ideas from [Anantharaman and Macià 2014; Macià and Rivièrè 2016] in order to derive some properties on the regularity of solutions to (2) when  $\epsilon_h \gg \hbar$ . In particular, we will identify precisely the concentration phenomena that may occur and also show nonconcentration properties by propagation of second microlocal data. Note that, when written in nonsemiclassical terms, the regime we are interested in corresponds to the eigenvalue problem

$$-\Delta u_\lambda(x) + f(\lambda) V(x) u_\lambda(x) = \lambda^2 u_\lambda(x), \quad x \in \mathbb{T}^d, \quad \|u_\lambda\|_{L^2(\mathbb{T}^d)} = 1,$$

where  $1 \ll f(\lambda) \ll \lambda^2$ .

For the sake of simplicity, we will focus on the case of the rational torus  $\mathbb{T}^2$  and assume  $V \in \mathcal{C}^\infty(\mathbb{T}^2; \mathbb{R})$ . However, it is most likely that our analysis could be extended to more general completely integrable systems of dimension 2 following the approach of [Anantharaman et al. 2015]. As the small perturbation regime<sup>2</sup>  $0 \leq \epsilon_h \leq \hbar$  was studied in great detail in all the above references, here we will focus on the strong perturbation regime and we shall assume throughout the article that

$$\lim_{\hbar \rightarrow 0^+} \epsilon_h = 0 \quad \text{and} \quad \lim_{\hbar \rightarrow 0^+} \hbar \epsilon_h^{-1} = 0. \tag{4}$$

In order to state our results, we need some simple geometric preliminaries. Recall that the geodesics of  $\mathbb{T}^2$  are either closed or dense curves. For  $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2 - \{0\}$  and  $x \in \mathbb{T}^2$ , the geodesic  $s \mapsto x + s\xi$  is dense provided  $\xi_1$  and  $\xi_2$  are linearly independent over  $\mathbb{Q}$ ; otherwise it is periodic. We denote by  $\Omega_1 \subset \mathbb{R}^2 - \{0\}$  the set of  $\xi$  that generate a periodic geodesic and by  $\Omega_2$  its complement in  $\mathbb{R}^2 - \{0\}$ . Consider the average of  $V$  along geodesics:

$$\mathcal{I}(V)(x, \xi) := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T V(x + s\xi) ds.$$

---

<sup>2</sup>Note that, for the nonsemiclassical version, it means that  $f(\lambda) \leq 1$ .

Clearly,  $\mathcal{I}(V)$  is a zero-homogeneous function with respect to  $\xi$ . Moreover, a classical result by Kronecker implies

$$\mathcal{I}(V)(x, \xi) = \begin{cases} (1/L_\xi) \int_0^{L_\xi} V(x + s(\xi/\|\xi\|)) ds & \text{if } \xi \in \Omega_1, \\ \int_{\mathbb{T}^2} V(y) dy & \text{if } \xi \in \Omega_2, \end{cases}$$

where  $L_\xi$  denotes the length of any geodesic with velocity  $\xi$ . In particular, despite the fact that  $\mathcal{I}(V)$  is not continuous in general, one has  $\mathcal{I}(V)(\cdot, \xi) \in C^\infty(\mathbb{T}^2; \mathbb{R})$  for any  $\xi \in \mathbb{R}^2 - \{0\}$ , and  $\|\mathcal{I}(V)\|_{L^\infty(\mathbb{T}^2 \times \mathbb{R}^2)} \leq \|V\|_{L^\infty(\mathbb{T}^2)}$ .

Then, we define the set of critical geodesics:

$$\mathcal{C}(V) := \{x_0 \in \mathbb{T}^2 : \text{there exists } \xi \in \Omega_1 \text{ such that } \partial_x \mathcal{I}(V)(x_0, \xi) = 0\}. \tag{5}$$

Note that  $\mathcal{C}(V)$  is a union of closed geodesics of  $\mathbb{T}^2$ . For every closed geodesic  $\gamma$  of  $\mathbb{T}^2$ , we denote by  $\delta_\gamma$  the normalized Lebesgue measure along this closed geodesic. Then, we define  $\mathcal{N}(V)$  as the convex closure of the set of probability measures  $\delta_\gamma$ , where  $\gamma \subset \mathcal{C}(V)$ . With these conventions in mind, we can state our main result:

**Theorem 1.1.** *Suppose that  $d = 2$  and that (4) holds. Let  $(u_h)_{h \rightarrow 0^+}$  be a sequence satisfying (2). Then, for any accumulation point  $\nu$  of the sequence of probability measures*

$$\nu_h(dx) := |u_h(x)|^2 dx,$$

and for any closed geodesic  $\gamma$ , one has

$$\nu(\gamma) \neq 0 \implies \gamma \subset \mathcal{C}(V).$$

Moreover,  $\nu$  can be decomposed as

$$\nu = f dx + \nu_{\text{sing}},$$

where  $f \in L^1(\mathbb{T}^2)$  and where  $\nu_{\text{sing}} \in \mathcal{N}(V)$ .

Recall from the propagation properties of semiclassical measures [Gérard 1991; Zworski 2012] that any  $\nu$  as in Theorem 1.1 must a priori be a convex combination of the Lebesgue measure and of the measures  $\delta_\gamma$ , where  $\gamma$  runs over the set of all closed geodesics. This theorem shows that singular concentration along closed geodesics can only occur along certain closed orbits associated with critical points of the averages of  $V$  along closed geodesics. This result is sharp in the sense that Wunsch’s construction in [Anantharaman et al. 2015] shows that one can find quasimodes such that  $\nu(\gamma) = 1$  for a given closed geodesic. Despite these unavoidable concentration phenomena, Theorem 1.1 also shows that the accumulation points enjoy certain regularity properties. This extra regularity will come out from our analysis by making a second microlocalization of size  $\epsilon_h$  along rational directions, and it will be induced by certain Lagrangian tori associated to our problem. Note that these two aspects are close to the situation of Zoll manifolds treated in [Macià and Rivièrè 2016; 2017]. The main difference is that there exist infinitely many directions where the flow is periodic with periods tending to  $+\infty$ . We would like to treat these tori of periodic orbits as in these references, and this can be achieved via rescaling the variables along these rational directions; see Section 3D for more details. Finally, as we shall see in Sections 2 and 3, our analysis holds in the more general context of the time-dependent Schrödinger equation.

**Organization of the article.** Section 2 places our problem in the more general framework of the time-dependent Schrödinger equation associated with  $\widehat{P}_\epsilon(\hbar)$ : Theorem 1.1 becomes a direct consequence of the more general Theorem 2.1, which deals with the evolution problem. The proof of this result is obtained by characterizing time-dependent semiclassical measures for solutions to the Schrödinger equation. Following a strategy similar to that in [Anantharaman and Macià 2014; Macià 2010], such a characterization can be obtained by using two-microlocal techniques. In Section 3, we introduce the two-microlocal framework of our analysis that is needed to formulate our main results, Theorems 3.6 and 3.7. Section 4 presents several applications of these results. We first give the proof of Theorem 2.1; then we present a structure result for semiclassical measures of the evolution equation, Theorem 4.1, which we apply to compute the propagation of wave packet solutions (Proposition 4.3). This shows that Theorem 2.1 is sharp in some sense. The proofs of the two-microlocal statements of Section 3 are given in Section 5. Finally, the article contains two appendices. Appendix A contains the proof of a geometric result which already appeared in [Macià and Rivière 2016] and which we adapt to the context of  $\mathbb{T}^2$ . In Appendix B, we collect a few tools from semiclassical analysis.

In the following (except in Appendix B), we will always suppose that  $d = 2$  and that (4) holds even if part of the result holds in greater generality.

## 2. Semiclassical measures for the time-dependent Schrödinger equation

As was already mentioned, Theorem 1.1 is a consequence of our analysis of the time-dependent semiclassical Schrödinger equation:

$$i\hbar \partial_t v_\hbar = \widehat{P}_\epsilon(\hbar)v_\hbar, \quad v_\hbar|_{t=0} = u_\hbar \in L^2(\mathbb{T}^2), \quad \|u_\hbar\|_{L^2} = 1. \tag{6}$$

For the sake of simplicity, we shall focus on sequences of initial data oscillating at the frequency  $\hbar^{-1}$ . Thus, we will always assume the following properties hold:

$$\limsup_{\hbar \rightarrow 0} \|\mathbf{1}_{[R, \infty)}(-\hbar^2 \Delta)u_\hbar\|_{L^2(M)} \rightarrow 0 \quad \text{as } R \rightarrow \infty, \tag{7}$$

$$\limsup_{\hbar \rightarrow 0} \|\mathbf{1}_{[0, \delta]}(-\hbar^2 \Delta)u_\hbar\|_{L^2(M)} \rightarrow 0 \quad \text{as } \delta \rightarrow 0^+. \tag{8}$$

Fix now a sequence of time scales  $(\tau_\hbar)_{\hbar \rightarrow 0^+}$  such that

$$\lim_{\hbar \rightarrow 0^+} \tau_\hbar = +\infty.$$

We will deal with time-scaled solutions to the perturbed Schrödinger equation. More precisely, if  $v_\hbar$  is a solution to (6), then we shall study the behavior of

$$t \mapsto v_\hbar(\tau_\hbar t, \cdot).$$

As we will see below, the scale  $\tau_\hbar = \epsilon_\hbar^{-1}$  is critical for this problem, and Theorem 1.1 follows from the analysis of the time-dependent equation in the regime  $\tau_\hbar \gg \epsilon_\hbar^{-1}$ .

**2A. Time-dependent semiclassical measures.** For a given  $t$  in  $\mathbb{R}$ , we denote the Wigner distribution at time  $t$  by

$$\langle w_\hbar(t), a \rangle := \langle v_\hbar(t), \text{Op}_\hbar^w(a) v_\hbar(t) \rangle, \tag{9}$$

where  $\text{Op}_h^w(a)$  is an  $\hbar$ -pseudodifferential operator with principal symbol  $a \in C_c^\infty(T^*\mathbb{T}^2)$  — see [Appendix B](#). Above,  $v_h(t)$  denotes the solution at time  $t$  of (6) with initial conditions satisfying the oscillating assumptions (7) and (8). This quantity represents the distribution of the  $L^2$ -mass of the solution to (6) in the phase space  $T^*\mathbb{T}^2$ . According to [\[Macià 2009\]](#), we can extract a subsequence  $\hbar_n \rightarrow 0^+$  as  $n \rightarrow +\infty$  such that, for every  $a$  in  $C_c^\infty(T^*\mathbb{T}^2)$  and for every  $\theta$  in  $L^1(\mathbb{R})$ ,

$$\lim_{\hbar_n \rightarrow 0^+} \int_{\mathbb{R} \times T^*\mathbb{T}^2} \theta(t) \langle w_{\hbar_n}(t\tau_{\hbar_n}), a \rangle dt = \int_{\mathbb{R} \times T^*\mathbb{T}^2} \theta(t) a(x, \xi) \mu(t, dx, d\xi) dt,$$

where, for a.e.  $t$  in  $\mathbb{R}$ ,  $\mu(t)$  is a finite positive Radon measure on  $T^*\mathbb{T}^2$ . Recall also that, for a.e.  $t \in \mathbb{R}$ ,  $\mu(t)$  is in fact a *probability measure* which does not put any mass on the zero section, thanks to the frequency assumption (8). In other words,

$$\mu(t)(\mathring{T}^*\mathbb{T}^2) = 1 \quad \text{for a.e. } t \in \mathbb{R}, \tag{10}$$

where

$$\mathring{T}^*\mathbb{T}^2 := \{(x, \xi) \in T^*\mathbb{T}^2 : \xi \neq 0\}.$$

Moreover, for a.e.  $t$  in  $\mathbb{R}$ ,  $\mu(t)$  is *invariant by the geodesic flow*  $\varphi^s$  on  $T^*\mathbb{T}^2$ .

For instance,  $\mu(t)$  can be the normalized Lebesgue measure along a closed orbit of the geodesic flow. We will denote by  $\mathcal{M}(\tau, \epsilon)$  the set of accumulation points of the sequences  $(\mu_h)$ , where  $\mu_h(t, \cdot) := w_h(t\tau_h, \cdot)$ , as the sequence of initial data  $(u_h)$  varies among normalized sequences satisfying (7) and (8). Similarly, one can define  $\mathcal{N}(\tau, \epsilon)$  to be the set of accumulation points of the sequences  $(n_h)$  of time-dependent probability measures on  $\mathbb{T}^2$ ,  $n_h(t, dx) := |v_h(t\tau_h, x)|^2 dx$ , obtained by letting the initial data vary among sequences satisfying (7), (8). Using (7), one can verify that

$$\mathcal{N}(\tau, \epsilon) = \left\{ \int_{\mathbb{R}^2} \mu(t, x, d\xi) : \mu \in \mathcal{M}(\tau, \epsilon) \right\}. \tag{11}$$

**2B. Statement of the results.** In order to relate the time-dependent approach to the quasimode case, we can remark that, given a sequence of quasimodes  $(u_h)_{h \rightarrow 0^+}$  satisfying (2), we can always find a sequence of time scales  $(\tau_h)$  such that

$$\lim_{h \rightarrow 0} \tau_h \epsilon_h = +\infty,$$

and, for every  $t \in \mathbb{R}$ ,

$$\lim_{h \rightarrow 0} \|v_h(\tau_h t, \cdot) - e^{-i\tau_h t/(2\hbar)} u_h\|_{L^2(\mathbb{T}^2)} = 0,$$

where  $v_h$  denotes the solution to (6) with initial condition  $u_h$ . This choice of  $(\tau_h)$  ensures that any accumulation point  $\nu$  of the sequence of probability measures  $(|u_h|^2 dx)$  belongs to  $\mathcal{N}(\tau, \epsilon)$  (even though it is constant in  $t$ ), since it is also an accumulation point of  $(|v_h(\tau_h t, \cdot)|^2 dx)$ . In particular, [Theorem 1.1](#) follows from the more general statement:

**Theorem 2.1.** *Suppose that*

$$\lim_{h \rightarrow 0} \tau_h \epsilon_h = +\infty.$$

Let  $t \mapsto \nu(t)$  be an element of  $\mathcal{N}(\tau, \epsilon)$ . Then, for any closed geodesic  $\gamma$  not included inside  $\mathcal{C}(V)$  and for a.e.  $t$  in  $\mathbb{R}$ , one has

$$\nu(t)(\gamma) = 0.$$

Moreover,  $\nu(t)$  can be decomposed as

$$\nu(t) = f(t) dx + \nu_{\text{sing}}(t),$$

where, for a.e.  $t$  in  $\mathbb{R}$ ,  $f(t) \in L^1(\mathbb{T}^2)$  and  $\nu_{\text{sing}}(t) \in \mathcal{N}(V)$ .

The first step in the proof of this result is the partition of  $\mathbb{R}^2 - \{0\}$  into  $\varphi^s$ -invariant subsets that was used in [Macià 2010; Anantharaman and Macià 2014]. Recall that  $\Lambda \subset \mathbb{Z}^2$  is a primitive lattice of rank 1 provided that  $\dim \langle \Lambda \rangle = 1$  and that  $\langle \Lambda \rangle \cap \mathbb{Z}^2 = \Lambda$ , where  $\langle \Lambda \rangle$  is the linear subspace of  $\mathbb{R}^2$  spanned by  $\Lambda$ . We introduce the invariant set of rational covectors

$$\Omega_1 = \bigsqcup_{\Lambda \text{ rank-1 primitive}} \Lambda^\perp - \{0\},$$

and its complement  $\Omega_2$  inside  $\mathbb{R}^2 - \{0\}$ , which is still invariant. Observe that this is consistent with the conventions of the Introduction. Because of (10), we can decompose the measure as follows:

$$\mu(t) = \mu(t)|_{\mathbb{T}^2 \times \Omega_2} + \sum_{\Lambda \text{ rank-1 primitive}} \mu(t)|_{\mathbb{T}^2 \times \Lambda^\perp - \{0\}}. \tag{12}$$

As a consequence of the invariance by the geodesic flow, it can be verified that  $\mu(t)|_{\mathbb{T}^2 \times \Omega_2}$  is in fact independent of the  $x$ -variable. Hence, in order to prove Theorem 2.1, one only has to study the regularity of  $\mu(t)|_{\mathbb{T}^2 \times \Lambda^\perp - \{0\}}$  for every rank-1 primitive sublattice  $\Lambda$ . This will be achieved using two-microlocal tools adapted to this problem. The end of the proof of Theorem 2.1 is presented in Section 4A. For time scales  $\tau_h = \mathcal{O}(\epsilon_h^{-1})$ , we obtain a more precise result, in the sense that each component of the time-dependent semiclassical measure  $\mu(t)$  according to the partition (12) can be completely determined from the initial data that were used to generate it. Again, the relation with the sequence of initial data is elucidated using the class of two-microlocal semiclassical measures that will be introduced in the next section. A precise statement is given in Theorem 4.1, Section 4B.

Finally, in Section 4C, we provide explicit computations of semiclassical measures associated to wave-packets (Proposition 4.3) that yield:

(1) If  $\tau_h \epsilon_h \rightarrow 0$ , then

$$\{\delta_\gamma : \gamma \text{ periodic geodesic of } \mathbb{T}^2\} \subset \mathcal{N}(\tau, \epsilon).$$

(2) If  $\tau_h = \epsilon_h^{-1}$ , then

$$\{\delta_\gamma : \gamma \in \mathcal{C}(V)\} \subset \mathcal{N}(\tau, \epsilon).$$

### 3. Invariance and propagation of two-microlocal distributions

We now present our main result on the two-microlocal structure of solutions to the time-dependent Schrödinger equation along covectors in  $\Omega_1$ . In particular, we show how solutions of (6) can concentrate along rational covectors.

Before stating the result, we need some additional notation. For every primitive rank-1 lattice  $\Lambda$  of  $\mathbb{Z}^2$ , we set  $\mathbf{e}_\Lambda$  to be an element in  $\Lambda$  such that  $\mathbb{Z}\mathbf{e}_\Lambda = \Lambda$ , and  $\mathbf{e}_\Lambda^\perp$  to be the vector of same length which is directly orthogonal to  $\mathbf{e}_\Lambda$ . We define

$$L_\Lambda := \|\mathbf{e}_\Lambda\|.$$

We define two Hamiltonian maps associated to  $\Lambda$  as follows:

$$H_\Lambda(\xi) := \frac{1}{L_\Lambda} \langle \xi, \mathbf{e}_\Lambda \rangle \quad \text{and} \quad H_\Lambda^\perp(\xi) := \frac{1}{L_\Lambda} \langle \xi, \mathbf{e}_\Lambda^\perp \rangle.$$

Note that  $(H_\Lambda, H_\Lambda^\perp)$  defines a (nondegenerate) completely integrable system and that

$$\|\xi\|^2 = H_\Lambda(\xi)^2 + H_\Lambda^\perp(\xi)^2.$$

**3A. Two-microlocal distributions.** We aim at studying the concentration of solutions to (6) over  $\mathbb{T}^2 \times \Lambda^\perp$ , where  $\Lambda \subset \mathbb{Z}^2$  is a primitive rank-1 sublattice and where  $\Lambda^\perp$  denotes the set of covectors  $\xi$  such that  $H_\Lambda(\xi) = 0$ . For that purpose, we consider a two-microlocal scale  $\alpha_\hbar \rightarrow 0^+$  satisfying  $\hbar\alpha_\hbar^{-1} \rightarrow 0$  and we define the following two-microlocal Wigner distribution:

$$w_{\Lambda, \hbar}(t) : a \in C_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}) \mapsto \left\langle v_\hbar(t), \text{Op}_\hbar^w \left( a \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \right) v_\hbar(t) \right\rangle.$$

Above,  $\widehat{\mathbb{R}}$  is the compactified space  $\mathbb{R} \cup \{\pm\infty\}$ ,  $v_\hbar(t)$  is the solution of (6) at time  $t$ , and  $\text{Op}_\hbar^w(a)$  is a  $\hbar$ -pseudodifferential operator — see Appendix B.

**Remark 3.1.** Recall from (28) in Appendix B that the following useful relation holds:

$$\text{Op}_\hbar^w \left( a \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \right) = \text{Op}_{\hbar\alpha_\hbar^{-1}}^w (a(x, \alpha_\hbar \xi, H_\Lambda(\xi))),$$

and that we have made the assumption that  $\hbar\alpha_\hbar^{-1} \rightarrow 0$ . Therefore, the operators involved in the definition of  $w_{\Lambda, \hbar}$  are semiclassical pseudodifferential operators whose symbolic calculus enjoys a gain of  $\hbar\alpha_\hbar^{-1}$ .

**Remark 3.2.** The distributions  $w_{\Lambda, \hbar}$  were introduced in [Macià 2010; Anantharaman and Macià 2014] for the critical case  $\alpha_\hbar = \hbar$  under a slightly different form. There, the two microlocal variable  $\eta$  varies in the two-point compactification of  $\langle \Lambda \rangle$ . Of course, this is completely equivalent to our formulation for the two-dimensional torus, but turns out to be relevant when dealing with the higher-dimensional case. As we will see, the fact that the two-microlocal scale is asymptotically bigger than  $\hbar$  implies that the limiting objects are of a different nature than those obtained in [Macià 2010; Anantharaman and Macià 2014]. When  $\hbar\alpha_\hbar^{-1} \rightarrow 0$ , they are global variants on the torus of the two-scale semiclassical measures introduced in [Fermanian-Kammerer 2005] — see also [Anantharaman and Léautaud 2014] for a related construction on the torus, in a context related to that of [Anantharaman and Macià 2014].

Recall that we introduced a time scale  $\tau_\hbar \rightarrow \infty$ . From now on, we shall fix the two-microlocal scale as follows:

$$\alpha_\hbar := \begin{cases} 1/\tau_\hbar & \text{if } \tau_\hbar \epsilon_\hbar^{-1} \rightarrow 0, \\ \epsilon_\hbar & \text{otherwise.} \end{cases} \tag{13}$$

As we shall explain in [Section 5A](#), we can extract a subsequence  $\hbar_n \rightarrow 0^+$  such that, for any  $a \in \mathcal{C}_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$  and for any  $\theta \in L^1(\mathbb{R})$ ,

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}} \theta(t) \langle \omega_{\Lambda, \hbar_n}(t\tau_{\hbar_n}), a \rangle dt = \int_{\mathbb{R}} \theta(t) \left( \int_{T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}} a(x, \xi, \eta) \mu_\Lambda(t, dx, d\xi, d\eta) \right) dt,$$

where, for a.e.  $t$  in  $\mathbb{R}$ ,  $\mu_\Lambda(t)$  is an element of  $\mathcal{B}'$  for some Banach space  $\mathcal{B}$  that we will define in [Section 5A](#). We denote by  $\mathcal{M}_\Lambda(\tau, \epsilon)$  the set of accumulation points obtained in this manner for initial data varying among subsequences verifying (7) and (8). The main new result of this article describes some invariance and propagation properties of these quantities depending on the relative sizes of  $\tau_\hbar$  and  $\epsilon_\hbar$ .

For every primitive rank-1 sublattice, one has (see [Remark 5.3](#))

$$\mathcal{M}(\tau, \epsilon) = \left\{ \int_{\widehat{\mathbb{R}}} \mu_\Lambda(t, x, \xi, d\eta) : \mu_\Lambda \in \mathcal{M}_\Lambda(\tau, \epsilon) \right\}. \tag{14}$$

**3B. First properties.** Before proving our main results, we will verify a few preliminary results.

**Proposition 3.3.** *Let  $\mu_\Lambda(t)$  be an element of  $\mathcal{M}_\Lambda(\tau, \epsilon)$ . Then, for a.e.  $t$  in  $\mathbb{R}$ ,  $\mu_\Lambda(t)$  is a positive finite Radon measure concentrated on  $\mathring{T}^*\mathbb{T}^2 \times \widehat{\mathbb{R}}$ .*

In what follows, we write

$$\tilde{\mu}_\Lambda(t) := \mu_\Lambda(t) \llcorner_{\mathring{T}^*\mathbb{T}^2 \times \mathbb{R}}, \quad \tilde{\mu}^\Lambda(t) := \mu_\Lambda(t) \llcorner_{\mathring{T}^*\mathbb{T}^2 \times \{\pm\infty\}}.$$

Hence, we can split the two-microlocal measure as

$$\mu_\Lambda(t) = \tilde{\mu}_\Lambda(t) + \tilde{\mu}^\Lambda(t). \tag{15}$$

The measure  $\tilde{\mu}_\Lambda(t)$  describes in some sense the way the solutions of (6) concentrate in an  $\epsilon_\hbar$ -neighborhood of the rational direction  $\Lambda^\perp$ . We now give some other simple properties of these functionals which are analogous to the ones satisfied by time-dependent semiclassical measures [[Macià 2009](#)]. We shall also verify:

**Proposition 3.4.** *Let  $\mu_\Lambda(t) \in \mathcal{M}_\Lambda(\tau, \epsilon)$ . Then:*

- (1)  $\tilde{\mu}_\Lambda(t)$  is a (finite) positive measure on  $T^*\mathbb{T}^2 \times \mathbb{R}$  whose support is contained in  $\mathbb{T}^2 \times (\Lambda^\perp - \{0\}) \times \mathbb{R}$ .
- (2) For every  $a$  in  $\mathcal{C}_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ ,

$$\langle \tilde{\mu}_\Lambda(t), \xi \cdot \partial_x a \rangle = \langle \tilde{\mu}^\Lambda(t), \xi \cdot \partial_x a \rangle = 0.$$

Neither [Proposition 3.3](#), nor part (1) of [Proposition 3.4](#) uses that the functions used to generate  $\mu_\Lambda(t)$  are solutions to (6). This fact is only used in the second part of [Proposition 3.4](#). Note that all these properties follow from standard arguments which need to be slightly adapted in order to fit into the two-microlocal set-up — see [Section 5](#) for details.

**3C. Main results.** Consider the Hamiltonian flow  $\varphi_{H_\Lambda^\perp}$  associated with  $H_\Lambda^\perp$ . Note that, for a continuous function  $b$  on  $T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}$ , we can define the average along this  $L_\Lambda$ -periodic flow as

$$\mathcal{I}_\Lambda(b)(x, \xi, \eta) := \frac{1}{L_\Lambda} \int_0^{L_\Lambda} b(\varphi_{H_\Lambda^\perp}^s(x, \xi), \eta) ds.$$

A direct computation gives

$$\mathcal{I}_\Lambda(b)(x, \xi, \eta) = \frac{1}{L_\Lambda} \int_0^{L_\Lambda} b\left(x + s \frac{\mathbf{e}_\Lambda^\perp}{L_\Lambda}, \xi, \eta\right) ds = \sum_{k \in \Lambda} \hat{b}_k(\xi, \eta) e^{2i\pi k \cdot x},$$

provided  $b$  has the Fourier expansion

$$b(x, \xi, \eta) = \sum_{k \in \mathbb{Z}^2} \hat{b}_k(\xi, \eta) e^{2i\pi k \cdot x}.$$

Moreover, if  $\mathcal{I}(b)$  denotes the average of  $b$  along the geodesic flow

$$\varphi^s(x, \xi) = (x + s\xi, \xi)$$

on  $T^*\mathbb{T}^2$ , then the following holds:

$$\mathcal{I}(b)(x, \xi, \eta) = \mathcal{I}_\Lambda(b)(x, \xi, \eta), \quad \text{provided that } \xi \in \Lambda^\perp - \{0\}. \tag{16}$$

In the case where  $b$  only depends on  $x$ , as is the case with  $b = V$ , it is easy to check that  $\mathcal{I}_\Lambda(V)$  does not depend on  $\xi$  and therefore we can identify it with an element in  $C^\infty(\mathbb{T}^2; \mathbb{R})$ .

**Remark 3.5.** Part (2) of [Proposition 3.4](#) implies that  $\mu_\Lambda(t)$  is invariant under the geodesic flow  $\varphi^s$ . For  $b$  in  $C_c^\infty(T^*\mathbb{T}^2 \times \mathbb{R})$ , this observation combined with part (1) in [Proposition 3.4](#) and identity (16) implies that, for a.e.  $t$  in  $\mathbb{R}$ ,

$$\langle \mu_\Lambda(t), b \rangle = \langle \mu_\Lambda(t), \mathcal{I}_\Lambda(b) \rangle.$$

We shall use this property several times in our proof of [Theorem 3.6](#) below.

We need to define an auxiliary Hamiltonian function on  $\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}$

$$p_\Lambda^V\left(x, \sigma \frac{\mathbf{e}_\Lambda^\perp}{L_\Lambda}, \eta\right) := \frac{1}{2}\eta^2 + \mathcal{I}_\Lambda(V)(x). \tag{17}$$

Denote by  $\varphi_{p_\Lambda^V}^t$  the flow of the vector field on  $\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}$ :

$$\eta \frac{\mathbf{e}_\Lambda}{L_\Lambda} \cdot \partial_x - \frac{\mathbf{e}_\Lambda}{L_\Lambda} \cdot \partial_x \mathcal{I}_\Lambda(V) \partial_\eta.$$

This is the Hamiltonian vector field associated to  $p_\Lambda^V$  with respect to the symplectic form obtained by taking the push-forward of the canonical symplectic form on  $T^*\mathbb{T}^2$  via the diffeomorphism

$$T^*\mathbb{T}^2 \ni (x, \xi) \mapsto \left(x, H_\Lambda^\perp(x, \xi) \frac{\mathbf{e}_\Lambda^\perp}{L_\Lambda}, H_\Lambda(x, \xi)\right) \in \mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}. \tag{18}$$

The flow  $\varphi_{p_\Lambda^V}^t$  commutes with  $\varphi_{H_\Lambda^\perp}^s$  when acting on  $\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}$ .

We are now ready to state the main results of this article. The first one concerns the ‘‘compact’’ part of these two-microlocal distributions. Their possible behaviors are classified according to the limit of  $\tau_{\hbar \in \hbar}$ .

**Theorem 3.6** (invariance and propagation near  $\Lambda$ ). *Let  $\Lambda$  be a primitive rank-1 sublattice and let  $\mu_\Lambda$  be an element of  $\mathcal{M}_\Lambda(\tau, \epsilon)$  obtained as the limit of  $(w_{\Lambda, \hbar}(t\tau_\hbar))$ . Denote by  $\mu_\Lambda^0$  the limit of  $(w_{\Lambda, \hbar}(0))$ . The following results hold:*

(1) If  $\tau_h \epsilon_h \rightarrow 0$  as  $\hbar \rightarrow 0^+$ , then  $t \mapsto \tilde{\mu}_\Lambda(t)$  is continuous, and one has, for every  $a$  in  $C_c^0(\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R})$ ,

$$\tilde{\mu}_\Lambda(t)(a) = \tilde{\mu}_\Lambda^0(\mathcal{I}_\Lambda(a) \circ \varphi_{p_\Lambda^0}^t).$$

(2) If  $\tau_h \epsilon_h \rightarrow c > 0$  as  $\hbar \rightarrow 0^+$ , then  $t \mapsto \tilde{\mu}_\Lambda(t)$  is continuous, and one has, for every  $a$  in  $C_c^0(\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R})$ ,

$$\tilde{\mu}_\Lambda(t)(a) = \tilde{\mu}_\Lambda^0(\mathcal{I}_\Lambda(a) \circ \varphi_{p_\Lambda^c}^{ct}).$$

(3) If  $\tau_h \epsilon_h \rightarrow +\infty$  as  $\hbar \rightarrow 0^+$ , then one has, for a.e.  $t$  in  $\mathbb{R}$  and, for every  $a$  in  $C_c^0(\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R})$ ,

$$\text{for all } s \in \mathbb{R}, \quad \tilde{\mu}_\Lambda(t)(a) = \tilde{\mu}_\Lambda(t)(a \circ \varphi_{p_\Lambda^s}^s).$$

Equivalently, this theorem says that, besides invariance by the geodesic flow, the solutions of (6) satisfy some extra invariance properties in a shrinking neighborhood of the rational direction at least for times  $\tau_h \gg \epsilon_h^{-1}$ . For shorter times, the concentration in this shrinking neighborhood is completely determined by the initial data. The proof of this theorem is given in Section 5. Note that, when  $\tau_h \epsilon_h \rightarrow 0$ , the conclusion of part (1) holds even if  $\epsilon_h = \hbar$ ; this will be clear from the proof. Section 5.1 in [Anantharaman et al. 2015] provides explicit computations of two-microlocal semiclassical measures in that regime.

It is interesting to compare part (2) of Theorem 3.6 with its counterpart in [Anantharaman and Macià 2014], where the regime  $\epsilon_h = \hbar$  is studied in detail in any dimension (not only in the two-dimensional case analyzed here). First, the nature of the limiting object  $\tilde{\mu}_\Lambda$  is rather different in that setting. It is no longer a positive measure, but rather a measure taking values in the set of Wigner transforms of positive Hermitian trace-class operators on the space  $L^2(\mathbb{T}_\Lambda)$ .<sup>3</sup> As a result, time-dependent semiclassical measures are absolutely continuous with respect to the Lebesgue measures in the  $x$ -variable. In that setting, the role of the flow  $\varphi_{p_\Lambda^s}^s$  is played by the quantum flow  $e^{-is(D_\Lambda^2 + \mathcal{I}_\Lambda(V))}$  — see Corollary 25 in [Anantharaman and Macià 2014] for a precise statement.

The part at infinity satisfies an additional regularity property. Indeed, if we define

$$\mathcal{I}_0(a)(\xi, \eta) := \int_{\mathbb{T}^2} a(y, \xi, \eta) dy,$$

then the following holds:

**Theorem 3.7** (regularity at infinity). *Let  $\Lambda$  be a primitive rank-1 sublattice and let  $\mu_\Lambda(t)$  be an element of  $\mathcal{M}_\Lambda(\tau, \epsilon)$ . Then, one has, for every  $a$  in  $C_c^\infty(\mathbb{T}^2 \times \mathbb{R}^2 \times \widehat{\mathbb{R}})$  and for a.e.  $t$  in  $\mathbb{R}$ ,*

$$\langle \tilde{\mu}^\Lambda(t), \mathcal{I}_\Lambda(a) - \mathcal{I}_0(a) \rangle = 0.$$

*In particular, the measure  $\tilde{\mu}^\Lambda(t) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp \times \widehat{\mathbb{R}}}$  is constant in  $x$ .*

In other words, the part at infinity has no (nonzero) Fourier coefficients in the  $\Lambda$ -direction. As for Theorem 3.6, this result depends highly on the choice of two-microlocal scale we have fixed from the beginning, and other scalings would yield other properties. The first conclusion of this theorem is proved in Section 5. The last assertion follows from the invariance<sup>4</sup> of  $\tilde{\mu}^\Lambda(t)$  under the geodesic flow, which

<sup>3</sup>This space consists of those functions in  $L^2(\mathbb{T}^2)$  that are invariant by translations in the direction  $\Lambda^\perp$ .

<sup>4</sup>Recall also that  $\mu_\Lambda(t)$  is supported on  $\mathring{T}^* \mathbb{T}^2 \times \widehat{\mathbb{R}}$ .

implies that for every  $a \in C_c^0(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$

$$\langle \tilde{\mu}^\Lambda(t) \rfloor_{\mathbb{T}^2 \times \Lambda^\perp \times \widehat{\mathbb{R}}}, a \rangle = \langle \tilde{\mu}^\Lambda(t) \rfloor_{\mathbb{T}^2 \times \Lambda^\perp \times \widehat{\mathbb{R}}}, \mathcal{I}_\Lambda(a) \rangle = \langle \tilde{\mu}^\Lambda(t) \rfloor_{\mathbb{T}^2 \times \Lambda^\perp \times \widehat{\mathbb{R}}}, \mathcal{I}_0(a) \rangle.$$

Note also that the conclusion of [Theorem 3.7](#) holds in the regime  $\epsilon_\hbar = \hbar$  (in any dimension); see part (ii) of [Theorem 12](#) in [[Anantharaman and Macià 2014](#)].

**3D. Comparison with Zoll manifolds.** [Theorem 3.6](#) shares also a lot of similarities with our main result on semiclassical measures for perturbations of Zoll Laplacians in [[Macià and Rivière 2016](#), Section 2.2]. In that case, we were considering the semiclassical operator

$$-\frac{1}{2}\hbar^2 \Delta_g + \epsilon_\hbar^2 V,$$

where  $\Delta_g$  is the Laplace Beltrami operator associated to a certain Zoll metric (say the standard metric on the canonical sphere). In the present article, we are analyzing the semiclassical measures associated to the same Schrödinger operator  $\widehat{P}_\epsilon(\hbar)$ . Studying the “compact” part of elements inside  $\mathcal{M}_\Lambda(\tau, \epsilon)$  is equivalent to understanding the solutions of (6) near submanifolds

$$\mathbb{T}^2 \times \Lambda^\perp := \{(x, \xi) \in T^*\mathbb{T}^2 : H_\Lambda(\xi) = 0\},$$

where the geodesic flow is periodic as in the Zoll case. In order to make the comparison clearer and to justify the rescaling of order  $\epsilon_\hbar$ , we can rewrite our operator in a form which is very close to what we did in the Zoll framework; i.e.,

$$\widehat{P}_\epsilon(\hbar) = \frac{1}{2} \text{Op}_\hbar^w(H_\Lambda^\perp)^2 + \epsilon_\hbar^2 \text{Op}_\hbar^w\left(\frac{1}{2}\left(\frac{H_\Lambda}{\epsilon_\hbar}\right)^2 + V\right).$$

Thus, as in the Zoll case, we perturb in some sense a semiclassical operator  $\text{Op}_\hbar^w(H_\Lambda^\perp)^2$  associated to a “periodic” Hamiltonian flow and we obtain limit quantities which are invariant by the periodic flow and the Hamiltonian perturbation.

The main difference with the Zoll setting is that the perturbation depends on rescaled variables

$$\left(x, H_\Lambda^\perp(\xi), \frac{H_\Lambda(\xi)}{\epsilon_\hbar}\right) \in \mathbb{T}^2 \times \mathbb{R}^2 \simeq T^*\mathbb{T}^2.$$

For that reason, it is natural to test our Wigner distributions against symbols depending on these rescaled variables. Another notable difference with [[Macià and Rivière 2016](#)] is that, in the Zoll case, the critical time scale is of order  $\epsilon_\hbar^{-2}$ , while here, due to the use of rescaled variables, it is much shorter, i.e., of order  $\epsilon_\hbar^{-1}$ . Finally, in the Zoll case, a natural question was to discuss the case where the Radon transform of the perturbation identically vanishes [[Macià and Riviere 2017](#)]. Here, we emphasize that the  $H_\Lambda^\perp$ -average of the perturbation, namely  $\frac{1}{2}(H_\Lambda/\epsilon_\hbar)^2 + \mathcal{I}_\Lambda(V)$  cannot be equal to a constant for this choice of two-microlocal rescaling.

### 4. Applications of the two-microlocal results

We present some applications of the results of the preceding section.

**4A. Proof of Theorem 2.1.** Recall that only the structure of the terms  $\mu(t) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp - \{0\}}$  in the decomposition (12) needs to be clarified. Thanks to (14) and to Proposition 3.4, we deduce

$$\mu(t) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp - \{0\}} = \mu(t) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp} = \int_{\mathbb{R}} \tilde{\mu}_\Lambda(t, \cdot, d\eta) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp} + \int_{\{\pm\infty\}} \tilde{\mu}^\Lambda(t, \cdot, d\eta) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp}.$$

According to Theorem 3.7, the contribution from the part at infinity is independent of  $x$ . Hence, we are left with studying the regularity of the measures on  $\mathbb{T}^2$ :

$$\int_{\Lambda^\perp \times \mathbb{R}} \tilde{\mu}_\Lambda(t, \cdot, d\xi, d\eta).$$

The measure  $\tilde{\mu}_\Lambda$  is invariant under the Hamiltonian flow  $\varphi_{H_\Lambda^\perp}^t$  (see Remark 3.5) and, by part (3) of Theorem 3.6, it is also invariant under the Hamiltonian flow  $\varphi_{p_\Lambda^\vee}^t$ , which commutes with  $\varphi_{H_\Lambda^\perp}^t$ . Using Appendix A, which describes the regularity of bi-invariant measures, we can conclude the proof of Theorem 2.1. More specifically, part (1) follows from Proposition A.1 and part (2) from Corollary A.3.

**4B. Semiclassical measures up the critical time scale  $\tau_h = \epsilon_h^{-1}$ .** At the time scales up to the critical scale  $\epsilon_h^{-1}$ , we can completely determine  $\mu_t$  in terms of the initial data:

**Theorem 4.1.** *Let  $\mu \in \mathcal{M}(\tau, \epsilon)$ . Suppose that it is generated by some sequence of initial data  $(u_h)_{h \rightarrow 0^+}$ . For every rank-1 primitive lattice  $\Lambda$ , let  $\tilde{\mu}_\Lambda^0$  be the restriction to  $\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}$  of the two-microlocal measure associated with  $(u_h)_{h \rightarrow 0^+}$ , and denote by  $\mu^0$  the semiclassical measure of  $(u_h)_{h \rightarrow 0^+}$ :*

(1) *If  $\tau_h = \epsilon_h^{-1}$ , then, for every  $a \in \mathcal{C}_c^0(\mathbb{T}^2 \times \mathbb{R}^2)$ , the following holds:*

$$\begin{aligned} \int_{\mathbb{T}^2 \times \mathbb{R}^2} a(x, \xi) \mu(t, dx, d\xi) &= \int_{\mathbb{T}^2 \times \mathbb{R}^2} \mathcal{I}_0(a)(\xi) \mu^0(dx, d\xi) \\ &+ \sum_{\Lambda \text{ rank-1 primitive}} \int_{\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}} (\mathcal{I}_\Lambda(a) - \mathcal{I}_0(a))(\varphi_{p_\Lambda^\vee}^t(x, \xi, \eta)) \tilde{\mu}_\Lambda^0(dx, d\xi, d\eta). \end{aligned}$$

(2) *If  $\tau_h \epsilon_h \rightarrow 0$ , then the same result holds, provided we replace  $\varphi_{p_\Lambda^\vee}^t$  by  $\varphi_{p_\Lambda^\vee}^0$  in the formula above.*

The proof is as follows. Let  $\mu \in \mathcal{M}(\tau, \epsilon)$ , and decompose it as in (12). Using the lift property (14), we can further decompose  $\mu$  as follows:

$$\mu(t) = \mu(t) \llcorner_{\mathbb{T}^2 \times \Omega_2} + \sum_{\Lambda \text{ rank-1 primitive}} \int_{\{\pm\infty\}} \tilde{\mu}^\Lambda(t, d\eta) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp} + \sum_{\Lambda \text{ rank-1 primitive}} \int_{\mathbb{R}} \tilde{\mu}_\Lambda(t, \cdot, d\eta) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp}.$$

Thanks to the invariance by the geodesic flow and to Theorem 3.7, we can conclude one more time that the first two terms on the right-hand side of the equality are independent of  $x$ . Thanks to the second part of Theorem 3.6, we can also write

$$\tilde{\mu}_\Lambda(t) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}} = (\varphi_{p_\Lambda^\vee}^t)_* (\tilde{\mu}_\Lambda^0 \llcorner_{\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}}) \quad (\text{resp. } \tilde{\mu}_\Lambda(t) \llcorner_{\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}} = (\varphi_{p_\Lambda^\vee}^0)_* (\tilde{\mu}_\Lambda^0 \llcorner_{\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}})),$$

when  $\tau_h = \epsilon_h^{-1}$  (resp.  $\tau_h \epsilon_h \rightarrow 0$ ). The result follows from the fact that the zero Fourier coefficient of  $\mu(t)$  is itself equal to the zero Fourier coefficient of  $\mu^0$  thanks to the following adaptation of Proposition 29 from [Anantharaman and Macià 2014].

**Lemma 4.2.** *Suppose that*

$$\lim_{\hbar \rightarrow 0^+} \tau_{\hbar} \epsilon_{\hbar}^2 = 0.$$

*Let  $\mu$  be an element in  $\mathcal{M}(\tau, \epsilon)$  and let  $\mu^0$  be the semiclassical measure of the sequence of initial data used to generate  $\mu$ . Then, one has, for a.e.  $t$  in  $\mathbb{R}$ , and for every  $b \in C_c(\mathbb{R}^2)$ ,*

$$\int_{\mathbb{T}^2 \times \mathbb{R}^2} b(\xi) \mu(t, dx, d\xi) = \int_{\mathbb{T}^2 \times \mathbb{R}^2} b(\xi) \mu^0(dx, d\xi).$$

**4C. Propagation of wave packets.** An application of [Theorem 2.1](#) is the computation of semiclassical measures for wave-packet-type solutions to [\(6\)](#).

Let us first define wave-packet data on the torus. Take  $\rho \in C_c^\infty(\mathbb{R}^2)$  supported in a small neighborhood of the origin such that  $\|\rho\|_{L^2(\mathbb{R}^2)} = 1$ . Let  $(x_0, \xi_0) \in \hat{T}^*\mathbb{T}^2$  and set

$$U_{\hbar}^{x_0, \xi_0}(x) := \frac{1}{\sigma_{\hbar}} \rho\left(\frac{x - x_0}{\sigma_{\hbar}}\right) e^{i(\xi_0 \cdot x)/\hbar},$$

where  $\sigma_{\hbar} \rightarrow 0^+$  and  $\sigma_{\hbar} \gg \hbar$ . Finally, write

$$u_{\hbar}^{x_0, \xi_0}(x) = \sum_{k \in \mathbb{Z}^2} U_{\hbar}^{x_0, \xi_0}(x + k). \tag{19}$$

If the support of  $\rho$  is small enough, then

$$\|u_{\hbar}^{x_0, \xi_0}\|_{L^2(\mathbb{T}^2)} = 1.$$

These initial data concentrate around  $x_0$  and oscillate in the direction of  $\xi_0$ . Moreover, it is straightforward to check that  $(u_{\hbar}^{x_0, \xi_0})$  satisfies [\(7\)](#) and [\(8\)](#). We next compute the time-dependent semiclassical measure of the sequence  $(v_{\hbar}^{x_0, \xi_0})$  of solutions to [\(6\)](#) issued from the initial data  $(u_{\hbar}^{x_0, \xi_0})$ .

**Proposition 4.3.** *Suppose that the concentration scale  $(\sigma_{\hbar})$  satisfies  $\hbar(\epsilon_{\hbar} \sigma_{\hbar})^{-1} \rightarrow 0$  and that  $\xi_0 \in \Omega_1$ . Let  $\mu^{x_0, \xi_0} \in \mathcal{M}(\tau, \epsilon)$  be generated by the initial data  $(u_{\hbar}^{x_0, \xi_0})$ . Let  $\gamma(x, \xi_0)$  denote the geodesic in  $\mathbb{T}^2$  issued from  $(x, \xi_0)$  and  $\delta_{\gamma(x, \xi_0)}$  the uniform probability measure on that geodesic. The following hold:*

(1) *If  $\tau_{\hbar} \epsilon_{\hbar} \rightarrow 0$ , then*

$$\mu^{x_0, \xi_0}(t, dx, d\xi) = \delta_{\gamma(x_0, \xi_0)}(dx) \delta_{\xi_0}(d\xi).$$

(2) *If  $\tau_{\hbar} = \epsilon_{\hbar}^{-1}$ , then*

$$\mu^{x_0, \xi_0}(t, dx, d\xi) = \delta_{\gamma(x(t), \xi_0)}(dx) \delta_{\xi_0}(d\xi),$$

*where  $x(t)$  is the projection on  $\mathbb{T}^2$  of  $\varphi_{p_{\Lambda_{\xi_0}}^t}(x_0, \xi_0, 0)$  with  $\Lambda_{\xi_0} = \{\xi_0\}^\perp \cap \mathbb{Z}^2$ . If  $x_0$  is a critical point of  $\mathcal{I}_{\Lambda_{\xi_0}}(V)$  then  $x(t) = x_0$  for all  $t \in \mathbb{R}$ . In that case,  $\mu^{x_0, \xi_0}$  is also constant in time.*

*Proof.* [Lemma 4.2](#) ensures that  $\mu(t)$  is supported on  $\mathbb{T}^2 \times \langle \xi_0 \rangle$  for a.e.  $t \in \mathbb{R}$ . Therefore, by virtue of [\(14\)](#),

$$\mu(t) = \int_{\widehat{\mathbb{R}}} \mu_{\Lambda_{\xi_0}}(t, \cdot, d\eta) \llcorner_{\mathbb{T}^2 \times \langle \xi_0 \rangle},$$

where  $\mu_{\Lambda_{\xi_0}} \in \mathcal{M}_{\Lambda_{\xi_0}}(\tau, \epsilon)$  is generated by  $(u_{\hbar}^{x_0, \xi_0})$ . Let  $\mu_{\Lambda_{\xi_0}}^0$  be an accumulation point of  $(w_{\hbar, \Lambda_{\xi_0}}(0))$ . Since  $\hbar\sigma_{\hbar}^{-1} \ll \epsilon_{\hbar} \leq \tau_{\hbar}^{-1}$ , one can verify that, in every regime,

$$\mu_{\Lambda_{\xi_0}}^0(dx, d\xi, d\eta) = \delta_{x_0}(dx) \delta_{\xi_0}(d\xi) \delta_0(d\eta);$$

e.g., see the proof of Proposition 5.2 in [Anantharaman et al. 2015]. The result then follows from Theorem 2.1. □

### 5. Proof of the two-microlocal statements

From this point on, we fix a primitive sublattice  $\Lambda$  of  $\mathbb{Z}^2$  of rank 1 and we will proceed to the proofs of the results on two-microlocal distributions. Namely, we will first recall how to extract converging subsequences from the sequences  $(w_{\Lambda, \hbar}(t\tau_{\hbar}))_{\hbar \rightarrow 0^+}$ . Then, we will briefly recall how to adapt the proofs from [Anantharaman and Macià 2014] in order to prove Propositions 3.3 and 3.4. Finally, we will give the proofs of Theorems 3.6 and 3.7.

**5A. Extracting subsequences.** Recall that, following [Macià 2010; Anantharaman and Macià 2014; Anantharaman et al. 2015], we have introduced an auxiliary linear form whose invariance properties will be analyzed precisely. For every  $a \in C_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ , we have set

$$\langle w_{\Lambda, \hbar}(t\tau_{\hbar}), a \rangle := \left\langle v_{\hbar}(t\tau_{\hbar}), \text{Op}_{\hbar}^w \left( a \left( x, \xi, \frac{H_{\Lambda}(\xi)}{\alpha_{\hbar}} \right) \right) v_{\hbar}(t\tau_{\hbar}) \right\rangle,$$

where, recall,  $\alpha_{\hbar}$  is given by (13). It will be useful to keep in mind Remark 3.1 throughout this section.

**Remark 5.1.** We emphasize that, for  $a$  in  $C_c^\infty(T^*\mathbb{T}^2)$ , one has

$$\langle w_{\hbar}(t\tau_{\hbar}), a \rangle = \langle w_{\Lambda, \hbar}(t\tau_{\hbar}), a \rangle.$$

Our first step is to explain how to extract converging subsequences following more or less standard procedures [Gérard 1991; Macià 2009; Anantharaman and Macià 2014; Zworski 2012]. For the sake of completeness, we briefly recall it. For that purpose, we denote by

$$\mathcal{B} := C_0^D(\mathbb{T}^2 \times \mathbb{R}^2 \times \widehat{\mathbb{R}})$$

the space of  $C^D$  functions on  $\mathbb{T}^2 \times \mathbb{R}^2 \times \widehat{\mathbb{R}}$  all of whose derivatives tend to 0 at infinity. We choose  $D > 0$  large enough so that Theorem B.2 holds for functions in  $\mathcal{B}$ .

We endow this space with its natural topology of Banach spaces. According to Theorem B.2, one knows that, for every  $a$  in  $C_c^\infty(\mathbb{R} \times T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ , one has

$$|\langle w_{\Lambda, \hbar}(t\tau_{\hbar}), a(t) \rangle| \leq C \sum_{|\alpha| \leq D} (\hbar\alpha_{\hbar}^{-1})^{|\alpha|/2} \|\partial^\alpha a(t)\|_\infty. \tag{20}$$

Thus, the map  $t \mapsto w_{\Lambda, \hbar}(t\tau_{\hbar})$  defines a bounded sequence in  $L^1(\mathbb{R}, \mathcal{B})'$ , and, after extracting a subsequence, one finds that there exists  $\mu_{\Lambda}$  in  $L^1(\mathbb{R}, \mathcal{B})'$  such that, for every  $a$  in  $C_c^\infty(\mathbb{R} \times T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ , one has

$$\lim_{\hbar \rightarrow 0^+} \int_{\mathbb{R} \times T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}} a(t, x, \xi, \eta) w_{\Lambda, \hbar}(t\tau_{\hbar}), dx, d\xi, d\eta) dt = \int_{\mathbb{R} \times T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}} a(t, x, \xi, \eta) \mu_{\Lambda}(dt, dx, d\xi, d\eta).$$

Thanks to (20) and to the fact that  $\hbar\alpha_h^{-1} \rightarrow 0^+$ , recall that, for every  $\theta$  in  $C_c^\infty(\mathbb{R})$  and for every  $a$  in  $C_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ , one has

$$\left| \int_{\mathbb{R} \times T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}} \theta(t) a(x, \xi, \eta) \mu_\Lambda(dt, dx, d\xi, d\eta) \right| \leq C \|\theta\|_{L^1(\mathbb{R})} \|a\|_{C_0^0(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})}.$$

Hence,  $\mu_\Lambda$  is absolutely continuous with respect to the  $t$ -variable; i.e., for every  $\theta$  in  $L^1(\mathbb{R})$  and every  $a$  in  $C_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ , one has

$$\lim_{\hbar \rightarrow 0^+} \int_{\mathbb{R}} \theta(t) \langle w_{\Lambda, \hbar}(t\tau_\hbar), a \rangle dt = \int_{\mathbb{R}} \theta(t) \langle \mu_\Lambda(t), a \rangle dt.$$

Moreover, for a.e.  $t$  in  $\mathbb{R}$ ,  $\mu_\Lambda(t)$  is a finite Radon measure on  $T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}$ .

**5B. Proof of Proposition 3.3.** We already know that the linear functionals  $\mu_\Lambda$  are Radon measures. It remains to verify that they are positive. To see this, take  $a \in C_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$  such that  $a \geq 0$ . Using the Gårding inequality (Theorem 4.32 in [Zworski 2012]), we deduce that

$$\langle w_{\Lambda, \hbar}(t\tau_\hbar), a \rangle \geq \mathcal{O}(\hbar\alpha_h^{-1}) = o(1).$$

**Remark 5.2.** Note that the proof of the Gårding inequality in [Zworski 2012] is given in the case of  $\mathbb{R}^d$ . The extension to compact manifolds usually requires dealing with symbols that decay in  $\xi$  as we differentiate with respect to  $\xi$ . Yet, in the case of the torus, we can verify that this property remains true for an observable  $a$  all of whose derivatives are bounded (i.e., not necessarily decaying in  $\xi$ ) as in  $\mathbb{R}^d$ . For that purpose, one can start from the Gårding inequality on  $\mathbb{R}^d$  and apply the arguments of the proof of [Zworski 2012, Theorem 5.5], which shows  $L^2$ -boundedness of pseudodifferential of order 0 on  $\mathbb{T}^d$ .

After integrating against a test function  $\theta$  in  $L^1(\mathbb{R})$  and passing to the limit  $\hbar \rightarrow 0$ , one finds that, for a.e.  $t$  in  $\mathbb{R}$ ,

$$\langle \mu_\Lambda(t), a \rangle \geq 0.$$

This concludes the proof that  $\mu_\Lambda$  is a positive, finite Radon measure on  $T^*\mathbb{T}^2 \times \widehat{\mathbb{R}}$  and one sets  $\tilde{\mu}_\Lambda(t) = \mu_\Lambda(t)|_{T^*\mathbb{T}^2 \times \mathbb{R}}$  and  $\tilde{\mu}^\Lambda(t) = \mu_\Lambda(t)|_{T^*\mathbb{T}^2 \times \{\pm\infty\}}$ . Thanks to the frequency assumption (8), one has, for a.e.  $t$  in  $\mathbb{R}$ ,

$$\mu_\Lambda(t)(\{\xi=0\}) = 0. \tag{21}$$

**Remark 5.3.** Remark 5.1 implies that, for a.e.  $t$  in  $\mathbb{R}$ , the time-dependent semiclassical measure  $\mu(t)$  can be obtained by

$$\mu(t) = \int_{\widehat{\mathbb{R}}} \mu_\Lambda(t, \cdot, d\eta). \tag{22}$$

**5C. Proof of Proposition 3.4.** Concerning the support of  $\tilde{\mu}_\Lambda(t)$ , we let  $a$  be an element in  $C_c^\infty(T^*\mathbb{T}^2 \times \mathbb{R})$  whose support does not intersect  $\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}$ . Using Remark 3.1, one has

$$\text{Op}_\hbar^w \left( a \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \right) = \text{Op}_{\hbar\alpha_h^{-1}}^w (a(x, \alpha_\hbar \xi, H_\Lambda(\xi))).$$

Hence, this operator is equal to 0 when  $\hbar$  is small enough (thanks to our assumption on the support of  $a$ ). This concludes the proof of the first part of [Proposition 3.4](#).

Let us now discuss invariance by the geodesic flow, which is the only property that uses the particular form of  $v_\hbar(t\tau_\hbar)$  so far. Again, we start with the “compact” part and we fix  $a$  to be an element in  $\mathcal{C}_c^\infty(T^*\mathbb{T}^2 \times \mathbb{R})$ . Using composition rules for pseudodifferential operators, we write

$$\frac{d}{dt} \langle w_{\Lambda, \hbar}(t\tau_\hbar), a \rangle = \tau_\hbar \langle w_{\Lambda, \hbar}(t\tau_\hbar), \xi \cdot \partial_x a \rangle + \frac{i\tau_\hbar \epsilon_\hbar^2}{\hbar} \langle v_\hbar(t\tau_\hbar), [V, \text{Op}_{\hbar\alpha_\hbar^{-1}}^w(a(x, \alpha_\hbar \xi, H_\Lambda(\xi)))] v_\hbar(t\tau_\hbar) \rangle.$$

Using [Theorem B.3](#) (more specifically [Remark B.4](#)) one more time, we have

$$[V, \text{Op}_{\hbar\alpha_\hbar^{-1}}^w(a(x, \alpha_\hbar \xi, H_\Lambda(\xi)))] = -\frac{\hbar}{i\alpha_\hbar} \text{Op}_\hbar^w \left( \frac{\epsilon_\Lambda}{L_\Lambda} \cdot \partial_x V \partial_\eta a \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \right) + \mathcal{O}(\hbar^3(\alpha_\hbar)^{-3}).$$

Combining these two identities with the facts  $\hbar\alpha_\hbar^{-1} = o(1)$  and  $\epsilon_\hbar\alpha_\hbar^{-1} = \mathcal{O}(1)$ , we find that

$$\frac{d}{dt} \langle w_{\Lambda, \hbar}(t\tau_\hbar), a \rangle = \tau_\hbar \left( \left\langle w_{\Lambda, \hbar}(t\tau_\hbar), \xi \cdot \partial_x a - \frac{\epsilon_\hbar^2}{\alpha_\hbar} \frac{\epsilon_\Lambda}{L_\Lambda} \cdot \partial_x V \partial_\eta a \right\rangle + o(\hbar) \right).$$

Let now  $\theta$  be an element in  $\mathcal{C}_c^1(\mathbb{R})$ . Integrating the previous equality against  $\theta$  and integrating by parts, we find

$$\int_{\mathbb{R}} \theta(t) \left\langle w_{\Lambda, \hbar}(t\tau_\hbar), \xi \cdot \partial_x a - \frac{\epsilon_\hbar^2}{\alpha_\hbar} \frac{\epsilon_\Lambda}{L_\Lambda} \cdot \partial_x V \partial_\eta a \right\rangle dt = \mathcal{O}(\tau_\hbar^{-1}) + o(\hbar),$$

which implies the result for every  $a$  in  $\mathcal{C}_c^\infty(T^*\mathbb{T}^2 \times \mathbb{R})$  when we let  $\hbar$  go to 0. Note that we used the Calderón–Vaillancourt theorem ([Theorem B.2](#)) to bound the  $\epsilon_\hbar^2\alpha_\hbar^{-1}$  term on the left-hand side of this equality.

It now remains to treat the part at infinity. Let  $a$  be an element in  $\mathcal{C}_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$ . For every  $R \geq 1$  and for every smooth cutoff function near 0, we set

$$a^R(x, \xi, \eta) := a(x, \xi, \eta) \left( 1 - \chi \left( \frac{\eta}{R} \right) \right).$$

The same argument as before allows us to prove that, for every  $\theta$  in  $\mathcal{C}^1(\mathbb{R})$ , one has

$$\int_{\mathbb{R}} \theta(t) \left\langle w_{\Lambda, \hbar}(t\tau_\hbar), (\xi \cdot \partial_x a)^R - \frac{\epsilon_\hbar^2}{\alpha_\hbar} \frac{\epsilon_\Lambda}{L_\Lambda} \cdot \partial_x V \partial_\eta a^R \right\rangle dt = o(1).$$

Thus, we can take the limit  $\hbar \rightarrow 0$  and conclude the proof by letting  $R$  go to  $+\infty$ .

**5D. Invariance and propagation of two-microlocal distributions.** We now turn to the proofs of our main statements, namely [Theorems 3.6](#) and [3.7](#). Analogously to [[Anantharaman and Macià 2014](#)], we define the differential operators

$$D_\Lambda := \frac{1}{i} \frac{\epsilon_\Lambda}{L_\Lambda} \cdot \nabla \quad \text{and} \quad D_\Lambda^\perp := \frac{1}{i} \frac{\epsilon_\Lambda^\perp}{L_\Lambda} \cdot \nabla$$

associated with the Hamiltonians  $H_\Lambda$  and  $H_\Lambda^\perp$ . One has

$$-\Delta = (D_\Lambda^\perp)^2 + D_\Lambda^2. \tag{23}$$

Recall also that, for every smooth compactly supported function  $b$  on  $T^*\mathbb{T}^2$ , the Egorov theorem is exact for these operators and it tells us that

$$\text{Op}_\hbar^w(\mathcal{I}_\Lambda(b)) = \frac{1}{L_\Lambda} \int_0^{L_\Lambda} e^{isD_\Lambda^\perp} \text{Op}_\hbar^w(b) e^{-isD_\Lambda^\perp} ds. \tag{24}$$

and that

$$[D_\Lambda^\perp, \text{Op}_\hbar^w(\mathcal{I}_\Lambda(b))] = 0. \tag{25}$$

As mentioned before, this construction (which was originally presented in [Anantharaman and Macià 2014]) is reminiscent of the averaging argument of [Weinstein 1977] applied to certain one-dimensional tori that depend on  $\Lambda$ .

**5D1. Proof of Theorem 3.6.** Let  $a$  be an element in  $C_c^\infty(T^*\mathbb{T}^2 \times \mathbb{R})$ . We start our proof by computing the derivative of the two-microlocal Wigner distribution. One has

$$\frac{d}{dt} \langle w_{\Lambda, \hbar}(t\tau_\hbar), \mathcal{I}_\Lambda(a) \rangle = \frac{i\tau_\hbar}{\hbar} \langle v_\hbar(t\tau_\hbar), [\frac{1}{2}\hbar^2(D_\Lambda^\perp)^2 + \frac{1}{2}\hbar^2 D_\Lambda^2 + \epsilon_\hbar^2 V, \text{Op}_\hbar^w(a_{\Lambda, \hbar})] v_\hbar(t\tau_\hbar) \rangle,$$

where

$$a_{\Lambda, \hbar}(x, \xi) := \mathcal{I}_\Lambda(a) \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right).$$

Using (25), we deduce that

$$\frac{d}{dt} \langle w_{\Lambda, \hbar}(t\tau_\hbar), \mathcal{I}_\Lambda(a) \rangle = \frac{i\tau_\hbar}{\hbar} \langle v_\hbar(t\tau_\hbar), [\frac{1}{2}\hbar^2 D_\Lambda^2 + \epsilon_\hbar^2 V, \text{Op}_\hbar^w(a_{\Lambda, \hbar})] v_\hbar(t\tau_\hbar) \rangle.$$

Thanks to the commutation properties of the Weyl quantization from Remark B.4, one has

$$\begin{aligned} & \frac{d}{dt} \langle w_{\Lambda, \hbar}(t\tau_\hbar), \mathcal{I}_\Lambda(a) \rangle \\ &= \mathcal{O}(\tau_\hbar \epsilon_\hbar^2 \hbar^2 (\alpha_\hbar)^{-3}) \\ &+ \alpha_\hbar \tau_\hbar \left\langle v_\hbar(t\tau_\hbar), \text{Op}_\hbar^w \left( \frac{H_\Lambda(\xi)}{\alpha_\hbar} \frac{\epsilon_\Lambda \cdot \partial_x \mathcal{I}_\Lambda(a)(x, \xi, H_\Lambda(\xi)/\alpha_\hbar)}{L_\Lambda} - \frac{\epsilon_\hbar^2}{\alpha_\hbar^2} \partial_\eta \mathcal{I}_\Lambda(a) \frac{\epsilon_\Lambda \cdot \partial_x V}{L_\Lambda} \right) v_\hbar(t\tau_\hbar) \right\rangle. \end{aligned} \tag{26}$$

Our assumption  $\hbar \ll \epsilon_\hbar \ll \alpha_\hbar$  ensures that the remainder is in fact of order  $o(\hbar\tau_\hbar)$ .

We now distinguish three regimes.

First, we suppose that  $\epsilon_\hbar \tau_\hbar \rightarrow 0$  as  $\hbar \rightarrow 0^+$ . In particular,  $\alpha_\hbar = \tau_\hbar^{-1} \gg \epsilon_\hbar$ . Thanks to the Calderón–Vaillancourt theorem (Theorem B.2), we can verify that the last term in the right-hand side of equality (26) is in fact  $o(1)$  uniformly for  $t$  in  $\mathbb{R}$ . Letting  $\hbar \rightarrow 0$ , one finds that, for a.e.  $t$  in  $\mathbb{R}$ ,

$$\frac{d}{dt} \langle \mu_\Lambda(t), \mathcal{I}_\Lambda(a) \rangle = \left\langle \mu_\Lambda(t), \eta \frac{\epsilon_\Lambda}{L_\Lambda} \cdot \partial_x \mathcal{I}_\Lambda(a) \right\rangle.$$

Combining Proposition 3.4 with (21), one has then  $\langle \mu_\Lambda(t), a \rangle = \langle \mu_\Lambda^0, \mathcal{I}_\Lambda(a) \circ \varphi_{p_\Lambda^0}^t \rangle$  for a.e.  $t$  in  $\mathbb{R}$ , which proves point (1) of the theorem.

Suppose now that  $\tau_{\hbar\epsilon_{\hbar}} \rightarrow c > 0$ . Letting  $\hbar \rightarrow 0$ , the limit measure satisfies the following transport equation for all  $\theta \in C_c^1(\mathbb{R})$ :

$$-\int_{\mathbb{R}} \theta'(t) \langle \mu_{\Lambda}(t), \mathcal{I}_{\Lambda}(a) \rangle dt = c \int_{\mathbb{R}} \theta(t) \left\langle \mu_{\Lambda}(t), \eta \frac{\epsilon_{\Lambda} \cdot \partial_x \mathcal{I}_{\Lambda}(a)}{L_{\Lambda}} - \partial_{\eta} \mathcal{I}_{\Lambda}(a) \frac{\epsilon_{\Lambda} \cdot \partial_x V}{L_{\Lambda}} \right\rangle dt.$$

Using again Proposition 3.4 with (21), one deduces that

$$\partial_t \langle \mu_{\Lambda}(t), \mathcal{I}_{\Lambda}(a) \rangle = c \left\langle \mu_{\Lambda}(t), \eta \frac{\epsilon_{\Lambda} \cdot \partial_x \mathcal{I}_{\Lambda}(a)}{L_{\Lambda}} - \partial_{\eta} \mathcal{I}_{\Lambda}(a) \frac{\epsilon_{\Lambda} \cdot \partial_x \mathcal{I}_{\Lambda}(V)}{L_{\Lambda}} \right\rangle.$$

This proves point (2) of the theorem.

Finally, we suppose that  $\tau_{\hbar\epsilon_{\hbar}} \rightarrow +\infty$ . Let  $\theta$  be an element in  $C_c^1(\mathbb{R})$ . We integrate one more time equality (26) against  $\theta$ , and we make an integration by parts on the left-hand side of the equality. Then, we make use of the Calderón–Vaillancourt theorem (Theorem B.2) to bound the left-hand side. After letting  $\hbar$  go to 0, one finds that, for every  $\theta$  in  $C_c^1(\mathbb{R})$ ,

$$\int_{\mathbb{R}} \theta(t) \left\langle \mu_{\Lambda}(t), \eta \frac{\epsilon_{\Lambda} \cdot \partial_x \mathcal{I}_{\Lambda}(a)}{L_{\Lambda}} - \partial_{\eta} \mathcal{I}_{\Lambda}(a) \frac{\epsilon_{\Lambda} \cdot \partial_x \mathcal{I}_{\Lambda}(V)}{L_{\Lambda}} \right\rangle dt = 0,$$

where we used one more time Proposition 3.4 with (21) in order to replace  $V$  by its  $\Lambda$ -average  $\mathcal{I}_{\Lambda}(V)$ . This implies point (3) of the theorem.

**5D2. Proof of Theorem 3.7.** Let now  $a$  be an element in  $C_c^{\infty}(\mathbb{R}^2 \times \widehat{\mathbb{R}})$  and let  $k$  be an element in  $\Lambda - \{0\}$ . It suffices to show that

$$\langle \tilde{\mu}^{\Lambda}(t), e^{-2i\pi k \cdot x} a(\xi, \eta) \rangle = 0.$$

We fix  $\chi_1(\eta) \in C^{\infty}(\mathbb{R}, [0, 1])$  which is equal to 1 for  $\eta \geq 1$  and to 0 for  $\eta \leq \frac{1}{2}$ . For every  $R \geq 1$ , we set

$$a_{\pm}^{R,k}(x, \xi, \eta) := e^{-2i\pi k \cdot x} a(\xi, \eta) \chi_1\left(\pm \frac{\eta}{R}\right).$$

**Remark 5.4.** Let  $\theta$  be an element in  $C_c^1(\mathbb{R})$ . One has

$$\int_{\mathbb{R}} \theta(t) \frac{d}{dt} \left\langle w_{\Lambda, \hbar}(t\tau_{\hbar}), \frac{1}{\eta} a_{\pm}^{R,k} \right\rangle dt = - \int_{\mathbb{R}} \theta'(t) \left\langle w_{\Lambda, \hbar}(t\tau_{\hbar}), \frac{1}{\eta} a_{\pm}^{R,k} \right\rangle dt.$$

Thanks to the Calderón–Vaillancourt theorem (Theorem B.2), one knows that

$$\left\| \text{Op}_{\hbar}^w \left( \chi \left( \frac{H_{\Lambda}(\xi)}{R\alpha_{\hbar}} \right) a \left( \xi, \frac{H_{\Lambda}(\xi)}{\alpha_{\hbar}} \right) e^{-2i\pi k \cdot x} \frac{\alpha_{\hbar}}{H_{\Lambda}(\xi)} \right) \right\|_{L^2 \rightarrow L^2} = \mathcal{O}(R^{-1}).$$

Thus, one has

$$\int_{\mathbb{R}} \theta(t) \frac{d}{dt} \left\langle w_{\Lambda, \hbar}(t\tau_{\hbar}), \frac{1}{\eta} a_{\pm}^{R,k} \right\rangle dt = \mathcal{O}(R^{-1}).$$

In order to prove the proposition, we will now compute explicitly the derivative of  $\langle w_{\Lambda, \hbar}(t\tau_{\hbar}), \frac{1}{\eta} a_{\pm}^{R,k} \rangle$ . For that purpose, we need to compute the following bracket:

$$\left[ -\frac{\hbar^2 \Delta}{2} + \epsilon_{\hbar}^2 V, \text{Op}_{\hbar}^w \left( a_{\pm}^{R,k} \left( x, \xi, \frac{H_{\Lambda}(\xi)}{\alpha_{\hbar}} \right) \frac{\alpha_{\hbar}}{H_{\Lambda}(\xi)} \right) \right].$$

Using again (25), this commutator is in fact equal to

$$\left[ \frac{\hbar^2 D_\Lambda^2}{2} + \epsilon_\hbar^2 V, \text{Op}_\hbar^w \left( a_\pm^{R,k} \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \frac{\alpha_\hbar}{H_\Lambda(\xi)} \right) \right].$$

We split this commutator in two parts. Thanks to Remark B.4, one has

$$\left[ \frac{\hbar^2 D_\Lambda^2}{2}, \text{Op}_\hbar^w \left( a_\pm^{R,k} \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \frac{\alpha_\hbar}{H_\Lambda(\xi)} \right) \right] = -2\pi \hbar \alpha_\hbar \text{Op}_\hbar^w \left( \frac{\epsilon_\Lambda}{L_\Lambda} .k a_\pm^{R,k} \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \right).$$

For the other part of the commutator, we use one more time the commutation rule for pseudodifferential operators and the Calderón–Vaillancourt theorem (Theorem B.2). We find that

$$\left[ V, \text{Op}_\hbar^w \left( a_\pm^{R,k} \left( x, \xi, \frac{H_\Lambda(\xi)}{\alpha_\hbar} \right) \frac{\alpha_\hbar}{H_\Lambda(\xi)} \right) \right] = \mathcal{O}_{L^2 \rightarrow L^2}(\hbar \alpha_\hbar^{-1} R^{-1} + \hbar^3 \alpha_\hbar^{-3}).$$

As  $\hbar \epsilon_\hbar^{-1} \rightarrow 0$  and  $\epsilon_\hbar = \mathcal{O}(\alpha_\hbar)$ , we finally get that

$$\frac{d}{dt} \left\langle w_{\Lambda, \hbar}(t \tau_\hbar), \frac{1}{\eta} a_\pm^{R,k} \right\rangle = -\frac{2\pi \tau_\hbar \alpha_\hbar \epsilon_\Lambda .k}{L_\Lambda} \langle w_{\Lambda, \hbar}(t \tau_\hbar), a_\pm^{R,k} \rangle + \mathcal{O}(\tau_\hbar \epsilon_\hbar R^{-1}) + o(\tau_\hbar \hbar).$$

Let now  $\theta$  be an element in  $\mathcal{C}_c^1(\mathbb{R})$ . We integrate these expressions against  $\theta$ . Using Remark 5.4 and making the assumption that  $\limsup_{\hbar \rightarrow 0^+} \tau_\hbar \alpha_\hbar > 0$ , we obtain

$$\text{for all } k \in \Lambda - \{0\}, \quad \int_{\mathbb{R}} \theta(t) \langle w_{\Lambda, \hbar}(t \tau_\hbar), a_\pm^{R,k} \rangle dt = o(1) + \mathcal{O}(R^{-1}).$$

We now let  $\hbar$  go to 0, and we get that, for every  $R > 0$ ,

$$\text{for all } k \in \Lambda - \{0\}, \quad \int_{\mathbb{R}} \theta(t) \langle \mu_\Lambda(t), a_\pm^{R,k} \rangle dt = \mathcal{O}(R^{-1}).$$

To get the conclusion, we let  $R$  go to  $+\infty$ .

**Remark 5.5.** From this theorem, we deduce that, for every  $a(x, \xi, \eta)$  in  $\mathcal{C}_c^\infty(T^*\mathbb{T}^2 \times \widehat{\mathbb{R}})$  and for a.e.  $t$  in  $\mathbb{R}$ ,

$$\tilde{\mu}^\Lambda(t)(\mathcal{I}_\Lambda(a)) = \int_{T^*\mathbb{T}^2 \times \{\pm\infty\}} \hat{a}_0(\xi, \eta) \mu_\Lambda(t, d\xi, d\eta).$$

### Appendix A. Regularity of bi-invariant measures

In this appendix, we fix  $\Lambda$  a primitive sublattice of  $\mathbb{Z}^2$  of rank 1, and we aim at analyzing the regularity of the set of finite measures on  $T^*\mathbb{T}^2$  which are invariant by the Hamiltonian flows<sup>5</sup>  $\varphi_{H_\Lambda^\perp}^t$  and  $\varphi_{p_\Lambda^V}^t$ . We will now recall the results from Section 4 of [Macià and Rivière 2016] and explain how they can be adapted to the present framework. We refer the reader to this reference for the detailed proofs. We introduce the critical set in the direction of  $\Lambda$ ,

$$\text{Crit}_\Lambda(V) := \{(x, \xi) \in T^*\mathbb{T}^2 : H_\Lambda(\xi) = 0 \text{ and } \partial_x \mathcal{I}_\Lambda(V) = 0\}.$$

<sup>5</sup>By making a slight abuse of notation, we shall identify  $\varphi_{p_\Lambda^V}^t$ , a flow a priori defined on  $\mathbb{T}^2 \times \Lambda^\perp \times \mathbb{R}$ , to a flow on  $T^*\mathbb{T}^2$  via the diffeomorphism (18). Recall that  $\varphi_{H_\Lambda^\perp}^t$  and  $\varphi_{p_\Lambda^V}^t$  commute.

This is a closed subset of  $T^*\mathbb{T}^2$  which is invariant by the Hamiltonian flows  $\varphi_{H_\Lambda^\perp}^t$  and  $\varphi_{p_\Lambda^V}^t$ , and we introduce its complement

$$\mathcal{R}(\Lambda) := T^*\mathbb{T}^2 - \text{Crit}_\Lambda(V).$$

The map

$$\phi : \mathbb{R}^2 \times \mathcal{R}(\Lambda) \ni (s, t, x, \xi) \mapsto \varphi_{H_\Lambda^\perp}^s \circ \varphi_{p_\Lambda^V}^t(x, \xi) \in \mathcal{R}(\Lambda)$$

is a group action of  $\mathbb{R}^2$  on  $\mathcal{R}(\Lambda)$ . Moreover, for any  $(x_0, \xi_0) \in \mathcal{R}(\Lambda)$ , the map

$$\phi_{x_0, \xi_0} : \mathbb{R}^2 \ni (s, t) \mapsto \varphi_{H_\Lambda^\perp}^s \circ \varphi_{p_\Lambda^V}^t(x_0, \xi_0) \in \mathcal{R}(\Lambda)$$

is an immersion. Therefore, the stabilizer group  $G_{x_0, \xi_0}$  of  $(x_0, \xi_0)$  under  $\phi$  is discrete. This proves that the orbits of the action  $\phi$  are either diffeomorphic to the torus  $\mathbb{T}^2$ , to the cylinder  $\mathbb{T} \times \mathbb{R}$  or to  $\mathbb{R}^2$ . On the other hand, the moment map,

$$\Phi : \mathcal{R}(\Lambda) \ni (x, \xi) \mapsto (H_\Lambda^\perp(\xi), p_\Lambda^V(x, \xi)) \in \mathbb{R}^2,$$

is a submersion, and, for every  $(H, J) \in \Phi(\mathcal{R}(\Lambda))$ , the level set

$$\mathcal{L}_{(H, J)} := \Phi^{-1}(H, J)$$

is a smooth submanifold of  $\mathcal{R}(\Lambda)$  of dimension 2. To summarize, the pair  $(H_\Lambda^\perp, p_\Lambda^V)$  forms a completely integrable system on  $\mathcal{R}(\Lambda)$ , and the map  $\phi_{x_0, \xi_0}$  induces a diffeomorphism:

$$\text{for all } (x_0, \xi_0) \in \mathcal{R}(\Lambda), \quad \phi_{x_0, \xi_0} : \mathbb{R}^2 / G_{x_0, \xi_0} \rightarrow \mathcal{L}_{(H_0, J_0)}^{x_0, \xi_0} \quad \text{for } (H_0, J_0) := \Phi(x_0, \xi_0).$$

Here,  $\mathcal{L}_{(H_0, J_0)}^{x_0, \xi_0}$  denotes the connected component of  $\mathcal{L}_{(H_0, J_0)}$  that contains  $(x_0, \xi_0)$ . Therefore, if  $\mathcal{L}_{(H_0, J_0)}^{x_0, \xi_0}$  is compact then it is an embedded Lagrangian torus in  $T^*\mathbb{T}^2$ . In that case, we shall write

$$\mathbb{T}_{x_0, \xi_0}^2 := \mathbb{R}^2 / G_{x_0, \xi_0}.$$

In the following, we denote by  $\mathcal{R}_c(\Lambda)$  the set formed by those  $(x, \xi) \in \mathcal{R}(\Lambda)$  such that  $\mathcal{L}_{\Phi(x, \xi)}^{x, \xi}$  is compact. Mimicking the proof of Proposition 4.2 in [Macià and Rivière 2016], one can show that the following holds:

**Proposition A.1.** *Let  $\mu$  be a probability measure on  $\mathcal{R}(\Lambda)$  that is invariant by  $\varphi_{H_\Lambda^\perp}^t$  and  $\varphi_{p_\Lambda^V}^t$ . Set  $\bar{\mu} := \Phi_*\mu$ . Then, for every  $a \in C_c(\mathcal{R}(\Lambda))$ , one has*

$$\int_{\mathcal{R}(\Lambda)} a(x, \xi) \mu(dx, d\xi) = \int_{\Phi(\mathcal{R}(\Lambda))} \int_{\mathcal{L}_{(H, J)}} a(x, \xi) \lambda_{H, J}(dx, d\xi) \bar{\mu}(dH, dJ),$$

where, for  $(H, J) \in \Phi(\mathcal{R}(\Lambda))$ , the measure  $\lambda_{H, J}$  is a convex combination of the (normalized) Haar measures on the tori  $\mathcal{L}_{(H, J)}^{x_0, \xi_0}$  for  $(x_0, \xi_0) \in \mathcal{L}_{(H, J)} \cap \mathcal{R}_c(\Lambda)$ . In particular, for every  $(x, \xi)$  in  $\mathcal{R}(\Lambda)$ , one has

$$\mu(\{\varphi_{H_\Lambda^\perp}^s(x, \xi) : 0 \leq s \leq L_\Lambda\}) = 0.$$

An explicit formula for the restriction of the measure  $\lambda_{H,J}$  to a connected component  $\mathcal{L}_{(H,J)}^{x,\xi}$  with  $(x, \xi) \in \mathcal{R}_c(\Lambda) \cap \mathcal{L}_{(H,J)}$  is the following:

$$\int_{\mathcal{L}_{(H,J)}^{x_0,\xi_0}} a(x, \xi) \lambda_{H,J}(dx, d\xi) = c \int_{\mathbb{T}_{x_0,\xi_0}^2} a(\phi_{x_0,\xi_0}(s, t)) ds dt \tag{27}$$

for some constant  $c \in [0, 1]$ .

We will now discuss the regularity of the projections of bi-invariant measures following the proof from Section 4.2 in [Macià and Rivière 2016]. We denote by  $\Pi : T^*\mathbb{T}^2 \rightarrow \mathbb{T}^2$  the canonical projection. The main result from Section 4 in [Macià and Rivière 2016] is the following:

**Theorem A.2.** *Let  $\mu$  be a probability measure on  $\mathcal{R}(\Lambda)$  that is invariant by  $\varphi_{H_\Lambda}^t$  and  $\varphi_{p_\Lambda}^t$ . Then,  $\nu := \Pi_*\mu$  is a probability measure on  $\mathbb{T}^2$  that is absolutely continuous with respect to the Lebesgue measure.*

Denote by  $\mathcal{N}(\Lambda)$  the convex closure of the set of measures  $\delta_{\Pi \circ \Gamma}$ , where  $\Gamma \subset T^*\mathbb{T}^2$  ranges over the orbits of  $\varphi_{H_\Lambda}^t$  that are contained in  $\text{Crit}_\Lambda(V)$ . A direct consequence of the previous theorem is the following:

**Corollary A.3.** *The projection  $\nu := \Pi_*\mu$  of a probability measure  $\mu$  on  $T^*\mathbb{T}^2$  that is invariant by  $\varphi_{H_\Lambda}^t$  and  $\varphi_{p_\Lambda}^t$  can be decomposed as*

$$\nu = f \text{vol} + \alpha \nu_{\text{sing}},$$

where  $f \in L^1(\mathbb{T}^2)$ ,  $\alpha \in [0, 1]$  and  $\nu_{\text{sing}} \in \mathcal{N}(\Lambda)$ .

Note that, for a “generic” choice of  $V$ , the set of points  $x$  satisfying  $\partial_x \mathcal{I}_\Lambda(V) = 0$  consists of finitely many closed geodesics of  $\mathbb{T}^2$ . In particular,  $\nu_{\text{sing}}$  is a finite combination of measures carried by closed geodesics.

*Proof.* As it is simple to explain in the current framework, we briefly explain how the proof of Theorem 4.6 in [Macià and Rivière 2016] can be adapted to prove Theorem A.2— see also Lemma 2.1 in [Bialy and Polterovich 1989]. Recall that it is sufficient to fix some  $(x_0, \xi_0) \in \mathcal{R}_c(\Lambda)$  and to prove that the set of points where

$$\phi_{x_0,\xi_0} : (s, t) \in \mathbb{T}_{x_0,\xi}^2 \mapsto \Pi \circ \varphi_{H_\Lambda}^s \circ \varphi_{p_\Lambda}^t(x_0, \xi_0) \in \mathbb{T}^2$$

is not a local diffeomorphism is made of finitely many disjoint  $C^1$  closed curves. Such curves are called caustics. This can be proved as follows. One can verify that the points where we do not have a local diffeomorphism are defined by the points  $(s, t)$  satisfying

$$H_\Lambda(\phi_{x_0,\xi_0}(s, t)) = 0.$$

Note that, for every  $s$  in  $\mathbb{R}$ ,

$$H_\Lambda(\varphi_{p_\Lambda}^t(x_0, \xi_0)) = H_\Lambda(\phi_{x_0,\xi_0}(s, t)).$$

As  $(x_0, \xi_0)$  belongs to the  $\varphi_{p_\Lambda}^t$ -invariant set  $\mathcal{R}(\Lambda)$ , we know that

$$\partial_x \mathcal{I}_\Lambda(V)(\varphi_{p_\Lambda}^t(x_0, \xi_0)) \neq 0.$$

Thus, from the Hamilton–Jacobi equations, we deduce that there exists a small open neighborhood  $(t - \eta, t + \eta)$  of  $t$  such that, for every  $t' \in (t - \eta, t + \eta) - \{t\}$ ,

$$H_\Lambda \circ \varphi_{p_\Lambda}^{t'}(x_0, \xi_0) \neq 0.$$

In particular, there are only finitely many values of  $t$  such that  $H_\Lambda \circ \varphi_{p_\Lambda}^t(x_0, \xi_0) \neq 0$  and thus, there are only finitely many closed curves on  $\mathbb{T}_{x_0, \xi_0}^2$  where the map  $\phi_{x_0, \xi_0}$  is not a local diffeomorphism.  $\square$

### Appendix B. Background on semiclassical analysis

In this appendix, we give a brief reminder of semiclassical analysis and we refer to [Zworski 2012] (mainly Chapters 1 to 5) for a more detailed exposition. Given  $\hbar > 0$  and  $a$  in  $\mathcal{S}(\mathbb{R}^{2d})$  (the Schwartz class), one can define the Weyl quantization of  $a$  as follows:

$$\text{for all } u \in \mathcal{S}(\mathbb{R}^d), \quad \text{Op}_\hbar^w(a)u(x) := \frac{1}{(2\pi\hbar)^d} \iint_{\mathbb{R}^{2d}} e^{(i/\hbar)\langle x-y, \xi \rangle} a\left(\frac{1}{2}(x+y), \xi\right) u(y) dy d\xi.$$

This definition can be extended to any observable  $a$  with uniformly bounded derivatives, i.e., such that for every  $\alpha \in \mathbb{N}^{2d}$ , there exists  $C_\alpha > 0$  such that  $\sup_{x, \xi} |\partial^\alpha a(x, \xi)| \leq C_\alpha$ . More generally, we will use the convention, for every  $m \in \mathbb{R}$  and every  $k \in \mathbb{Z}$ ,

$$S^{m,k} := \left\{ (a_\hbar(x, \xi))_{0 < \hbar \leq 1} : \text{for all } (\alpha, \beta) \in \mathbb{N}^d \times \mathbb{N}^d, \sup_{(x, \xi) \in \mathbb{R}^{2d}; 0 < \hbar \leq 1} |\hbar^k \langle \xi \rangle^{-m} \partial_x^\alpha \partial_\xi^\beta a_\hbar(x, \xi)| < +\infty \right\},$$

where  $\langle \xi \rangle := (1 + \|\xi\|^2)^{1/2}$ . For such symbols,  $\text{Op}_\hbar^w(a)$  defines a continuous operator  $\mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}(\mathbb{R}^d)$  which acts by duality on  $\mathcal{S}'(\mathbb{R}^d)$ .

**Remark B.1.** We also have the following relation, which we use at different stages of our proof:

$$\text{for all } \delta > 0, \text{ for all } a \in S^{m,k}, \quad \text{Op}_\hbar^w(a(x, \xi)) = \text{Op}_{\hbar\delta}^w(a(x, \delta\xi)). \tag{28}$$

Among the above symbols, we distinguish the family of  $\mathbb{Z}^d$ -periodic symbols, which we denote by  $S_{\text{per}}^{m,k}$ . Note that any  $a$  in  $C^\infty(T^*\mathbb{T}^d)$  (with bounded derivatives) defines an element in  $S_{\text{per}}^{0,0}$ . Similarly to the proof of Theorem 4.19 in [Zworski 2012], one can verify that, for any  $a \in S_{\text{per}}^{m,k}$ ,

$$\text{Op}_\hbar^w(a)(e_k) = \sum_{q \in \mathbb{Z}^d} e_q \hat{a}_{q-k}(\pi\hbar(q+k)),$$

where  $e_k(x) := e^{2i\pi k \cdot x}$ , and  $\hat{a}_p(\xi) := \int_{\mathbb{T}^d} a(x, \xi) e^{-2i\pi p \cdot x} dx$ . In particular, for any  $a \in S_{\text{per}}^{m,k}$ , the operator  $\text{Op}_\hbar^w(a)$  maps trigonometric polynomials into a smooth  $\mathbb{Z}^d$ -periodic function, and more generally any smooth  $\mathbb{Z}^d$ -periodic function into a smooth  $\mathbb{Z}^d$ -periodic function. Thus, for every  $a$  in  $S_{\text{per}}^{m,k}$ , the operator  $\text{Op}_\hbar^w(a)$  acts by duality on the space of distributions  $\mathcal{D}'(\mathbb{T}^d)$ . An important feature of this quantization procedure is that it defines a bounded operator on  $L^2(\mathbb{T}^d)$  [Zworski 2012, Chapter 5]:

**Theorem B.2** (Calderón–Vaillancourt). *There exists a constant  $C_d > 0$  and an integer  $D > 0$  such that, for every  $a$  in  $S_{\text{per}}^{0,0}$ , one has, for every  $0 < \hbar \leq 1$ ,*

$$\| \text{Op}_\hbar^w(a) \|_{L^2(\mathbb{T}^d) \rightarrow L^2(\mathbb{T}^d)} \leq C_d \sum_{|\alpha| \leq D} \hbar^{|\alpha|/2} \| \partial^\alpha a \|_\infty.$$

Another important feature of the Weyl quantization procedure is the composition formula:

**Theorem B.3** (composition formula). *Let  $a \in S^{m_1, k_1}$  and  $b \in S^{m_2, k_2}$ . Then, one has, for any  $0 < \hbar \leq 1$ ,*

$$\text{Op}_\hbar^w(a) \circ \text{Op}_\hbar^w(b) = \text{Op}_\hbar^w(a \sharp_\hbar b)$$

*in the sense of operators from  $\mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}(\mathbb{R}^d)$ , where  $a \sharp_\hbar b$  has uniformly bounded derivatives, and, for every  $N \geq 0$ ,*

$$a \sharp_\hbar b \sim \sum_{k=0}^N \frac{1}{k!} \left(\frac{1}{2}i\hbar D\right)^k(a, b) + \mathcal{O}(\hbar^{N+1}),$$

*where  $D(a, b)(x, \xi) = (\partial_x \partial_v - \partial_y \partial_\xi)(a(x, \xi)b(y, v)) \big|_{y=x, v=\xi}$ .*

We refer to Chapter 4 of [Zworski 2012] for a detailed proof of this result. We observe that for  $N = 0$ , the coefficient is given by the symbol  $ab$ , and for  $N = 1$ , it is given by  $(\hbar/(2i))\{a, b\}$ , where  $\{\cdot, \cdot\}$  is the Poisson bracket. As before, we can restrict this result to the case of periodic symbols, and we can check that the composition formula remains valid for operators acting on  $C^\infty(\mathbb{T}^d)$ .

**Remark B.4.** We note that the formula for the composed symbols is quite symmetric, and we have in fact the following useful property; for every  $N \geq 0$ ,

$$a \sharp_\hbar b - b \sharp_\hbar a \sim \sum_{k=0}^N \frac{2}{(2k+1)!} \left(\frac{1}{2}i\hbar D\right)^{2k+1}(a, b) + \mathcal{O}(\hbar^{2N+3}).$$

Finally, note that, if  $b(\xi)$  is a polynomial in  $\xi$  of order  $\leq 2$ , one has the exact formula

$$a \sharp_\hbar b - b \sharp_\hbar a = \frac{\hbar}{2i}\{a, b\}.$$

### Acknowledgements

We warmly thank the referee for his careful reading and his useful suggestions regarding the results presented in this article.

### References

- [Anantharaman and Léautaud 2014] N. Anantharaman and M. Léautaud, “Sharp polynomial decay rates for the damped wave equation on the torus”, *Anal. PDE* **7**:1 (2014), 159–214. [MR](#) [Zbl](#)
- [Anantharaman and Macià 2014] N. Anantharaman and F. Macià, “Semiclassical measures for the Schrödinger equation on the torus”, *J. Eur. Math. Soc. (JEMS)* **16**:6 (2014), 1253–1288. [MR](#) [Zbl](#)
- [Anantharaman et al. 2015] N. Anantharaman, C. Fermanian-Kammerer, and F. Macià, “Semiclassical completely integrable systems: long-time dynamics and observability via two-microlocal Wigner measures”, *Amer. J. Math.* **137**:3 (2015), 577–638. [MR](#) [Zbl](#)
- [Anantharaman et al. 2016a] N. Anantharaman, M. Léautaud, and F. Macià, “Delocalization of quasimodes on the disk”, *C. R. Math. Acad. Sci. Paris* **354**:3 (2016), 257–263. [MR](#) [Zbl](#)
- [Anantharaman et al. 2016b] N. Anantharaman, M. Léautaud, and F. Macià, “Wigner measures and observability for the Schrödinger equation on the disk”, *Invent. Math.* **206**:2 (2016), 485–599. [MR](#) [Zbl](#)
- [Bialy and Polterovich 1989] M. L. Bialy and L. V. Polterovich, “Lagrangian singularities of invariant tori of Hamiltonian systems with two degrees of freedom”, *Invent. Math.* **97**:2 (1989), 291–303. [MR](#) [Zbl](#)

- [Bourgain 1993] J. Bourgain, “Eigenfunction bounds for the Laplacian on the  $n$ -torus”, *Internat. Math. Res. Notices* **1993**:3 (1993), 61–66. [MR](#) [Zbl](#)
- [Bourgain 2013] J. Bourgain, “Moment inequalities for trigonometric polynomials with spectrum in curved hypersurfaces”, *Israel J. Math.* **193**:1 (2013), 441–458. [MR](#) [Zbl](#)
- [Bourgain and Demeter 2015] J. Bourgain and C. Demeter, “The proof of the  $l^2$  decoupling conjecture”, *Ann. of Math. (2)* **182**:1 (2015), 351–389. [MR](#) [Zbl](#)
- [Bourgain et al. 2013] J. Bourgain, N. Burq, and M. Zworski, “Control for Schrödinger operators on 2-tori: rough potentials”, *J. Eur. Math. Soc. (JEMS)* **15**:5 (2013), 1597–1628. [MR](#) [Zbl](#)
- [Burq and Zworski 2004] N. Burq and M. Zworski, “Geometric control in the presence of a black box”, *J. Amer. Math. Soc.* **17**:2 (2004), 443–471. [MR](#) [Zbl](#)
- [Burq and Zworski 2012] N. Burq and M. Zworski, “Control for Schrödinger operators on tori”, *Math. Res. Lett.* **19**:2 (2012), 309–324. [MR](#) [Zbl](#)
- [Fermanian-Kammerer 2000] C. Fermanian-Kammerer, “Mesures semi-classiques 2-microlocales”, *C. R. Acad. Sci. Paris Sér. I Math.* **331**:7 (2000), 515–518. [MR](#) [Zbl](#)
- [Fermanian-Kammerer 2005] C. Fermanian-Kammerer, “Analyse à deux échelles d’une suite bornée de  $L^2$  sur une sous-variété du cotangent”, *C. R. Math. Acad. Sci. Paris* **340**:4 (2005), 269–274. [MR](#) [Zbl](#)
- [Fermanian-Kammerer and Gérard 2002] C. Fermanian-Kammerer and P. Gérard, “Mesures semi-classiques et croisement de modes”, *Bull. Soc. Math. France* **130**:1 (2002), 123–168. [MR](#) [Zbl](#)
- [Gérard 1991] P. Gérard, “Mesures semi-classiques et ondes de Bloch”, exposé 16 in *Séminaire sur les Équations aux Dérivées Partielles*, 1990–1991, École Polytech., Palaiseau, 1991. [MR](#) [Zbl](#)
- [Jakobson 1997] D. Jakobson, “Quantum limits on flat tori”, *Ann. of Math. (2)* **145**:2 (1997), 235–266. [MR](#) [Zbl](#)
- [Macià 2009] F. Macià, “Semiclassical measures and the Schrödinger flow on Riemannian manifolds”, *Nonlinearity* **22**:5 (2009), 1003–1020. [MR](#) [Zbl](#)
- [Macià 2010] F. Macià, “High-frequency propagation for the Schrödinger equation on the torus”, *J. Funct. Anal.* **258**:3 (2010), 933–955. [MR](#) [Zbl](#)
- [Macià 2011] F. Macià, “The Schrödinger flow in a compact manifold: high-frequency dynamics and dispersion”, pp. 275–289 in *Modern aspects of the theory of partial differential equations*, edited by M. Ruzhansky and J. Wirth, Oper. Theory Adv. Appl. **216**, Springer, 2011. [MR](#)
- [Macià and Rivière 2016] F. Macià and G. Rivière, “Concentration and non-concentration for the Schrödinger evolution on Zoll manifolds”, *Comm. Math. Phys.* **345**:3 (2016), 1019–1054. [MR](#) [Zbl](#)
- [Macià and Riviere 2017] F. Macià and G. Riviere, “Observability and quantum limits for the Schrödinger equation on the sphere”, preprint, 2017. [arXiv](#)
- [Miller 1996] L. Miller, *Propagation d’ondes semi-classiques à travers une interface et mesures 2-microlocales*, Ph.D. thesis, École Polytechnique, 1996.
- [Nier 1996] F. Nier, “A semi-classical picture of quantum scattering”, *Ann. Sci. École Norm. Sup. (4)* **29**:2 (1996), 149–183. [MR](#) [Zbl](#)
- [Vasy and Wunsch 2009] A. Vasy and J. Wunsch, “Semiclassical second microlocal propagation of regularity and integrable systems”, *J. Anal. Math.* **108** (2009), 119–157. [MR](#) [Zbl](#)
- [Weinstein 1977] A. Weinstein, “Asymptotics of eigenvalue clusters for the Laplacian plus a potential”, *Duke Math. J.* **44**:4 (1977), 883–892. [MR](#) [Zbl](#)
- [Wunsch 2008] J. Wunsch, “Spreading of Lagrangian regularity on rational invariant tori”, *Comm. Math. Phys.* **279**:2 (2008), 487–496. [MR](#) [Zbl](#)
- [Wunsch 2012] J. Wunsch, “Non-concentration of quasimodes for integrable systems”, *Comm. Partial Differential Equations* **37**:8 (2012), 1430–1444. [MR](#) [Zbl](#)
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics **138**, American Mathematical Society, Providence, RI, 2012. [MR](#) [Zbl](#)
- [Zygmund 1974] A. Zygmund, “On Fourier coefficients and transforms of functions of two variables”, *Studia Math.* **50** (1974), 189–201. [MR](#) [Zbl](#)

Received 29 Aug 2017. Revised 17 Jan 2018. Accepted 10 Apr 2018.

FABRICIO MACIÀ: [fabricio.macia@upm.es](mailto:fabricio.macia@upm.es)

*Universidad Politécnica de Madrid, ETSI Navales, Madrid, Spain*

GABRIEL RIVIÈRE: [gabriel.riviere@math.univ-lille1.fr](mailto:gabriel.riviere@math.univ-lille1.fr)

*Laboratoire Paul Painlevé (U.M.R. CNRS 8524), U.F.R. de Mathématiques, Université Lille 1, Villeneuve d'Ascq, France*

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

[patrick.gerard@math.u-psud.fr](mailto:patrick.gerard@math.u-psud.fr)

Université Paris Sud XI

Orsay, France

## BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy <a href="mailto:berti@sissa.it">berti@sissa.it</a>	Clément Mouhot	Cambridge University, UK <a href="mailto:c.mouhot@dpmms.cam.ac.uk">c.mouhot@dpmms.cam.ac.uk</a>
Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>	Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>
Alessio Figalli	ETH Zurich, Switzerland <a href="mailto:alessio.figalli@math.ethz.ch">alessio.figalli@math.ethz.ch</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>
Vaughan Jones	U.C. Berkeley & Vanderbilt University <a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Vadim Kaloshin	University of Maryland, USA <a href="mailto:vadim.kaloshin@gmail.com">vadim.kaloshin@gmail.com</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Richard B. Melrose	Massachusetts Inst. of Tech., USA <a href="mailto:rbb@math.mit.edu">rbb@math.mit.edu</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>
William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>	Maciej Zworski	University of California, Berkeley, USA <a href="mailto:zvorski@math.berkeley.edu">zvorski@math.berkeley.edu</a>

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 11 No. 8 2018

---

Invariant measure and long time behavior of regular solutions of the Benjamin–Ono equation	1841
MOUHAMADOU SY	
Rigidity of minimizers in nonlocal phase transitions	1881
OVIDIU SAVIN	
Propagation and recovery of singularities in the inverse conductivity problem	1901
ALLAN GREENLEAF, MATTI LASSAS, MATTEO SANTACESARIA, SAMULI SILTANEN and GUNTHER UHLMANN	
Quantitative stochastic homogenization and regularity theory of parabolic equations	1945
SCOTT ARMSTRONG, ALEXANDRE BORDAS and JEAN-CHRISTOPHE MOURRAT	
Hopf potentials for the Schrödinger operator	2015
LUIGI ORSINA and AUGUSTO C. PONCE	
Monotonicity of nonpluripolar products and complex Monge–Ampère equations with pre- scribed singularity	2049
TAMÁS DARVAS, ELEONORA DI NEZZA and CHINH H. LU	
On weak weighted estimates of the martingale transform and a dyadic shift	2089
FEDOR NAZAROV, ALEXANDER REZNIKOV, VASILY VASYUNIN and ALEXANDER VOLBERG	
Two-microlocal regularity of quasimodes on the torus	2111
FABRICIO MACIÀ and GABRIEL RIVIÈRE	
Spectral distribution of the free Jacobi process, revisited	2137
TAREK HAMDI	