

Permanence properties of amenable, transitive and faithful actions

Soyoung Moon*

Abstract

We study hereditary properties of the class of countable groups admitting an amenable, transitive and faithful action on a countable set. We consider mainly the case of amalgamated free products, and we show in particular that the double of amenable groups and the amalgamated free products of two amenable groups over a finite subgroup admit such actions.

1 Introduction

Let G be a countable group acting on a countable set X .

Definition 1.1. The action of G on X is *amenable* if there exists a G -invariant mean on X , i.e. a map $\mu : 2^X = \mathcal{P}(X) \rightarrow [0, 1]$ such that $\mu(X) = 1$, $\mu(A \cup B) = \mu(A) + \mu(B)$ for every pair of disjoint subsets A, B of X , and $\mu(gA) = \mu(A)$, $\forall g \in G, \forall A \subseteq X$.

A group is amenable if the action on itself by left multiplication is amenable.

The above definition is equivalent to the existence of a *Følner sequence*, i.e., a sequence $\{A_n\}_{n \geq 1}$ of finite non-empty subsets of X such that for every $g \in G$, one has

$$\lim_{n \rightarrow \infty} \frac{|A_n \Delta g \cdot A_n|}{|A_n|} = 0.$$

The class \mathcal{A} of countable groups admitting an amenable, transitive and faithful action on a countable set is introduced by Glasner and Monod in [2]. Whilst

*This work is supported by Swiss NSF grant 20 – 126689/1.

Received by the editors April 2010.

Communicated by A. Valette.

the class \mathcal{A} is closed under direct products, free products and extension of co-amenable subgroups [†] (Proposition 1.7. in [2]), it is not closed under amalgamated free products even if two factors are amenable groups. Indeed, viewing the group $SL_2(\mathbb{Z})$ as an amalgamated free product

$$SL_2(\mathbb{Z}) \simeq (\mathbb{Z}/6\mathbb{Z}) *_{(\mathbb{Z}/2\mathbb{Z})} (\mathbb{Z}/4\mathbb{Z}),$$

one can see the group $SL_2(\mathbb{Z}) \rtimes \mathbb{Z}^2$ as the amalgamated free product $G *_A H$ of $G = \mathbb{Z}/4\mathbb{Z} \rtimes \mathbb{Z}^2$ and $H = \mathbb{Z}/6\mathbb{Z} \rtimes \mathbb{Z}^2$ along $A = \mathbb{Z}/2\mathbb{Z} \rtimes \mathbb{Z}^2$. Notice that the three groups G, H and A are in \mathcal{A} since they are amenable, but the amalgamated free product is not in \mathcal{A} since the pair $(SL_2(\mathbb{Z}) \rtimes \mathbb{Z}^2, \mathbb{Z}^2)$ has relative Property (T) and \mathbb{Z}^2 does not have finite exponent (it follow from Lemma 4.3. in [2] that if $G \in \mathcal{A}$ and the pair (G, H) has relative Property (T), then H has finite exponent, i.e. there is an integer n such that for every $h \in H, h^n = 1$).

The purpose of this paper is to investigate some hereditary properties of the class \mathcal{A} in case of amalgamated free products. Our first result is (see Proposition 3):

Theorem 1. *Let G be a group such that G surjects onto an amenable group H . If A is a common subgroup of G and H such that $\pi|_A$ is injective and $[H : A] \geq 2$, then the amalgamated free product $G *_A H$ is in \mathcal{A} . In particular, if G is amenable, the double of G over A is in \mathcal{A} , for every subgroup A of G .*

In case of double of free groups on two generators over a cyclic subgroup $\mathbb{F}_2 *_Z \mathbb{F}_2$, it was proven in [4] that any finite index subgroup of such group is in \mathcal{A} . Let us mention that this result was generalized to amalgamated free products of any free groups over a cyclic subgroup (called cyclically pinched one-relator groups) $\mathbb{F}_n *_Z \mathbb{F}_m, \forall n, m \geq 2$ in [3].

Our second theorem concerns amalgamated free products over a finite subgroup (see Proposition 4). In particular we show:

Theorem 2. *Let G and H be countable groups and A be a common finite subgroup of G and H . If G is an infinite amenable group and there is a H -set Y such that the H -action is amenable and the action of A on Y is free, then $G *_A H$ is in \mathcal{A} .*

As corollaries, the amalgamated free product of an infinite amenable group and a residually finite group over a finite subgroup (for instance $SL_3(\mathbb{Z}) *_A G$ with any infinite amenable group G and a common finite subgroup A) is in \mathcal{A} (Corollary 8); and the amalgamated free product of two amenable groups over a finite subgroup is also contained in \mathcal{A} (Corollary 9).

Acknowledgement. I would like to thank Nicolas Monod for useful discussions and Alain Valette for his valuable comments on a previous version of this paper.

[†]A subgroup $H < G$ is co-amenable in G if the G -action on G/H is amenable.




Figure 1:

2 Hereditary properties of \mathcal{A}

2.1 Double of amenable groups

An amalgamated free product $G *_A H$ is called *double of G over A* if H is isomorphic to G and the amalgamated subgroup of H is given by the isomorphism. Such a group has a presentation $\langle G, \phi(G) \mid a = \phi(a), \forall a \in A \rangle$ where $\phi : G \rightarrow H$ is an isomorphism. As mentioned above, in general the amalgamated free product is not in \mathcal{A} even if the factors are amenable groups. But it is true if the amalgam is a double; or more generally, if one factor surjects onto the other factor which is amenable with some extra condition (Theorem 1):

Proposition 3. *Let H be an amenable group and let $\pi : G \twoheadrightarrow H$ be a group epimorphism and let $A < G$ be a subgroup such that $\pi|_A$ is injective and $[H : \pi(A)] \geq 2$. Then the amalgamated free product $G *_A H$ given by π , i.e.*

$$G *_A H = \langle G, H \mid \pi(a) = a, \forall a \in A \rangle$$

is in \mathcal{A} . In particular, if G is amenable, the double of G over A is in \mathcal{A} .

Proof. Let $\psi : G *_A H \rightarrow H$ be a homomorphism defined by $\psi(g) = \pi(g)$, and $\psi(h) = h$ for all $g \in G$ and $h \in H$. Let $\text{Ker}(\psi) = N$. Let X be the Bass-Serre tree of $G *_A H$ and $Y = N \backslash X$ be the quotient graph. Since N intersects A trivially, by the theory of Bass-Serre, N is isomorphic to the free product

$$N \simeq K * (*_i H_i),$$

where K is a free group isomorphic to the fundamental group $\pi_1(Y, T)$ of the graph Y relatively to the maximal tree T which is generated by g_y with $y \in O - T$ where O is an orientation of Y , and H_i is the intersection of N with a conjugate of some factor of $G *_A H$ (cf. Remarque in p. 61 in [7]). Thus in order to have $K \neq \{1\}$, it suffices that the quotient graph $Y = N \backslash X$ is not a tree. If there exist $x \in G$ and $h \in N$ such that $hH = xH$ and $hA \neq xA$ (see Figure 1), then the quotient graph Y will have a circuit of length 2.

Thus, it is sufficient to find $x \in G$ and $h \in N$ such that $x \in G \setminus A$ and $x^{-1}h \in H \setminus A$.

By assumption on the index of $\pi(A)$ in H , there exists $z \in H \setminus \pi(A)$ so that $z^{-1} \in H \setminus \pi(A)$. Let $x \in G \setminus A$ such that $\pi(x) = z$. Let $h = x\pi(x^{-1})$. Then $\psi(h) = \psi(x)\psi(\pi(x^{-1})) = \pi(x)\pi(x^{-1}) = 1$, so $h \in N$, and we have

$$x^{-1}h = x^{-1}x\pi(x^{-1}) = \pi(x^{-1}) = z^{-1} \in H \setminus A.$$

Therefore, the normal subgroup N is a free product $K * (*_i H_i)$ with $K \neq \{1\}$, and the quotient is the amenable group H . Recall that a non-trivial free product $G_1 * G_2$, where G_i is a free group for some $i \in \{1, 2\}$, is always in \mathcal{A} from Theorem 1.5 in [2]. Recall also that the class \mathcal{A} is closed under extension of co-amenable subgroups. It follows that N is co-amenable in $G *_A H$ and is in \mathcal{A} , thus $G *_A H$ is in \mathcal{A} . For the second statement, take $\pi = \text{Id}_G$. ■

2.2 Amalgamated free products over a finite subgroup

Definition 2.1. Let G, H be two countable groups and let A be a common finite subgroup of G and H . We say that the triple (G, H, A) is in the class \mathcal{A}' if there exist a G -action on X and a H -action on Y such that

- (i) the action $G \curvearrowright X$ is transitive;
- (ii) for every element g of $G \setminus A$, and h of $H \setminus A$, the sets

$$\text{supp}_A(g) = \{x \in X \mid Ax \cap gAx = \emptyset\}$$

$$\text{supp}_A(h) = \{x \in Y \mid Ax \cap hAx = \emptyset\}$$

are infinite;

- (iii) there exist Følner sequences $\{C_n\}_{n \geq 1}$ of $G \curvearrowright X$ and $\{D_n\}_{n \geq 1}$ of $H \curvearrowright Y$ such that

(iii)-1. $|C_n| = |D_n|, \forall n \geq 1$;

(iii)-2. the sets $\{A \cdot C_n\}_{n \geq 1}, \{A \cdot D_n\}_{n \geq 1}$ are pairwise disjoint;

- (iv) the action of A on X and Y are free.

Note that if $(G, H, A) \in \mathcal{A}'$ then $G \in \mathcal{A}$.

Proposition 4. *If $(G, H, A) \in \mathcal{A}'$ then $G *_A H \in \mathcal{A}$.*

Proof. Let X be a countable set carrying actions of G and H as in the definition. Since A acts freely in the two actions, conjugating we may assume that the G -action and the H -action coincide on A .

First of all, let us recall that the group $\text{Sym}(X)$ of permutations of X endowed with the topology of pointwise convergence [‡] is a Baire space. Recall that a subset $Y \subset \text{Sym}(X)$ is *meagre* if it is a union of countably many closed subsets with

[‡]A sequence $\alpha_n \in \text{Sym}(X)$ converges to $\alpha \in \text{Sym}(X)$ if for all finite subset F of X , there exists n_0 such that $\alpha_n|_F = \alpha|_F$, for all $n \geq n_0$.

empty interior; and *generic* or *dense* G_δ if its complement $Sym(X) \setminus Y$ is meagre. Baire's theorem states that in a complete metric space, the intersection of countably many dense open subsets is dense, in particular not empty. Thus in order to find a permutation α of X having the properties $\{P_i\}_{i \geq 1}$, it is enough to prove that the set

$$\mathcal{U}_i = \{\alpha \in Sym(X) \mid \alpha \text{ satisfies the property } P_i\}$$

is generic in $Sym(X)$ and take $\alpha \in \bigcap_{i \geq 1} \mathcal{U}_i$.

Now let $\{C_n\}_{n \geq 1}$ (respectively $\{D_n\}_{n \geq 1}$) be the Følner sequence for G (respectively for H) satisfying the condition (iii) as in Definition 2.1. Set $Z = \{\sigma \in Sym(X) \mid \sigma a = a\sigma, \forall a \in A\}$; this is a closed subset in $Sym(X)$ so it is a Baire space. For $\sigma \in Z$, denote by $H^\sigma = \sigma^{-1}H\sigma$. By universality, the amalgamated free product $G *_A H^\sigma$ acts on X by $g \cdot x = gx$ and $h \cdot x = \sigma^{-1}h\sigma x$ for all $g \in G$ and $h \in H$. We shall prove that the sets

$$\mathcal{O}_1 = \{\sigma \in Z \mid \text{the action } G *_A H^\sigma \text{ on } X \text{ is faithful}\},$$

and

$$\mathcal{O}_2 = \{\sigma \in Z \mid \text{there is a subsequence } \{n_k\} \text{ of } n \text{ such that } \sigma(C_{n_k}) = D_{n_k}\}$$

are generic in Z .

Indeed, for the genericity of \mathcal{O}_1 , we shall prove that for every non-trivial word $w \in G *_A H$, the set

$$\mathcal{V}_w = \{\sigma \in Z \mid w^\sigma = \text{Id}_X\}$$

is closed and of empty interior. It is clear that the set \mathcal{V}_w is closed. To prove that the set \mathcal{V}_w is of empty interior, let us consider the case where $w = ag_n h_n \cdots g_1 h_1$ with $a \in A$, $g_i \in G \setminus A$ and $h_i \in H \setminus A$ (the other three cases are similar). The corresponding element of $Sym(X)$ given by the action is $w^\sigma = ag_n \sigma^{-1} h_n \sigma \cdots g_1 \sigma^{-1} h_1 \sigma$. Let $\sigma \in \mathcal{V}_w$. Let $F \subset X$ be a finite subset. Choose $x_0 \notin F \cup \sigma(F)$ such that $Ax_0 \cap (F \cup \sigma(F)) = \emptyset$. Inductively on $1 \leq i \leq n$, we choose a new point $x_{4i-3} \in \text{supp}_A(h_i)$ such that Ax_{4i-3} and $h_i Ax_{4i-3}$ are outside of the finite set of all points defined before (this is possible by (ii) in Definition 2.1). Then we define

$$\sigma'(ax_{4i-4}) := ax_{4i-3} \text{ and } \sigma'(a\sigma^{-1}(x_{4i-3})) := a\sigma(x_{4i-4}),$$

for all $a \in A$. Then set $x_{4i-2} := h_i x_{4i-3}$. We choose again a new point $x_{4i-1} \in \text{supp}_A(g_i)$ such that Ax_{4i-1} and $g_i Ax_{4i-1}$ are outside of the finite set of all points considered so far. We then define

$$\sigma'(ax_{4i-1}) := ax_{4i-2} \text{ and } \sigma'(a\sigma^{-1}(x_{4i-2})) := a\sigma(x_{4i-1}),$$

for all $a \in A$. Then set $x_{4i} := g_i x_{4i-1}$.

Every point v on which σ' is defined verifies $\sigma' a(v) = a\sigma'(v)$, $\forall a \in A$. Indeed, let $a, a' \in A$. Then,

- $\sigma' a(a' x_{4i-4}) = \sigma'(aa' x_{4i-4}) = aa' x_{4i-3} = a(a' x_{4i-3}) = a\sigma'(a' x_{4i-4})$;
- $\sigma' a(a' \sigma^{-1}(x_{4i-3})) = \sigma'(aa' \sigma^{-1}(x_{4i-3})) = aa' \sigma(x_{4i-4}) = a(a' \sigma(x_{4i-4})) = a\sigma'(a' \sigma^{-1}(x_{4i-3}))$;

$$\begin{aligned} & \cdot \sigma' a(a' x_{4i-1}) = \sigma'(aa' x_{4i-1}) = aa' x_{4i-2} = a(a' x_{4i-2}) = a\sigma'(a' x_{4i-1}); \\ & \cdot \sigma' a(a' \sigma^{-1}(x_{4i-2})) = \sigma'(aa' \sigma^{-1}(x_{4i-2})) = aa' \sigma(x_{4i-1}) = a(a' \sigma(x_{4i-1})) = \\ & \quad a\sigma'(a' \sigma^{-1}(x_{4i-2})). \end{aligned}$$

By construction, the $4n + 1$ points obtained by the right subwords of $w^{\sigma'}$ are all distinct and in particular $w^{\sigma'} x_0 \neq x_0$. We then define σ' to be σ on every points except these finite points, so that $\sigma' \in Z \setminus \mathcal{V}_w$ satisfies $\sigma'|_F = \sigma|_F$. This concludes the genericity of \mathcal{O}_1 .

About the genericity of \mathcal{O}_2 , let us write $\mathcal{O}_2 = \bigcap_{N \in \mathbb{N}} \{\sigma \in \text{Sym}(X) \mid \text{there exists } m \geq N \text{ such that } \sigma(C_m) = D_m\}$. We shall show that for every $N \in \mathbb{N}$, the set $\mathcal{V}_N = \{\sigma \in \text{Sym}(X) \mid \forall m \geq N, \sigma(C_m) \neq D_m\}$ is of empty interior (the closedness is clear). Let $F \subset X$ be a finite subset and $\sigma \in \mathcal{V}_N$. Let $m \geq N$ large enough such that $A \cdot x_i \cap (F \cup \sigma(F)) = \emptyset$ and $A \cdot y_i \cap (F \cup \sigma(F)) = \emptyset$, for every $x_i \in C_m$ and $y_i \in D_m, \forall i$; this is possible by (iii)-2. of Definition 2.1. By (iv) of Definition 2.1, we have $|A \cdot x_i| = |A \cdot y_i|, \forall i$. We then define

$$\sigma'(ax_i) := ay_i \text{ and } \sigma'(a\sigma^{-1}(y_i)) := a\sigma(x_i),$$

for every $1 \leq i \leq |C_m| = |D_m|$ and $a \in A$. For all other points, we define σ' to be equal to σ so that $\sigma' \in Z \setminus \mathcal{V}_N$ and $\sigma'|_F = \sigma|_F$.

Let $\sigma \in \mathcal{O}_1 \cap \mathcal{O}_2$. Let $\{C_{n_k}\}_{k \geq 1}$ be the subsequence of $\{C_n\}_{n \geq 1}$ such that $\sigma(C_{n_k}) = D_{n_k}, \forall k \geq 1$. The sequence $\{C_{n_k}\}_{k \geq 1}$ is a Følner sequence for G , and for every $h \in H$, we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{|C_{n_k} \Delta h \cdot C_{n_k}|}{|C_{n_k}|} &= \lim_{k \rightarrow \infty} \frac{|C_{n_k} \Delta \sigma^{-1} h \sigma C_{n_k}|}{|C_{n_k}|} = \lim_{k \rightarrow \infty} \frac{|\sigma C_{n_k} \Delta h \sigma C_{n_k}|}{|C_{n_k}|} \\ &= \lim_{k \rightarrow \infty} \frac{|D_{n_k} \Delta h D_{n_k}|}{|D_{n_k}|} = 0. \end{aligned}$$

Thus the sequence $\{C_{n_k}\}_{k \geq 1}$ is a Følner sequence for $G *_A H^\sigma$, and therefore $G *_A H^\sigma$ is in \mathcal{A} . ■

Remark 2.1. 1. The condition (ii) in Definition 2.1 is trivially satisfied if the G -action on X is free (which implies that G is amenable).

2. About (iv) in Definition 2.1, this condition is used in the proof of the genericity of \mathcal{O}_1 and \mathcal{O}_2 where, given any two points x and y , we needed to have $|Ax| = |Ay|$ in order to define σ' such that $\sigma'(Ax) = Ay$.

The following lemma shows that given two infinite amenable groups, one can always find Følner sequences having the same cardinality. Precisely, we have:

Lemma 5. *Let G, H be infinite amenable groups. Then there exist Følner sequences $\{C_n\}_{n \geq 1}$ of G and $\{D_n\}_{n \geq 1}$ of H such that $|C_n| = |D_n|, \forall n \geq 1$.*

Proof. We shall show that for any $\varepsilon > 0$, any finite subset $F \subset G$ and any finite subset $E \subset H$, there exist a finite subset $C' \subset G$ and a finite subset $D' \subset H$ such that C' is (ε, F) -Følner and D' is (ε, E) -Følner verifying $|C'| = |D'|$.

Recall that a finite subset $A \subset G$ is (ε, F) -Følner if

$$|A \Delta gA| < \varepsilon|A|, \forall g \in F.$$

By amenability of G , there is a finite subset $C_0 \subset G$ which is (ε, F) -Følner. Let $\{D_n\}_{n \geq 1}$ be a Følner sequence of H . Let $n \gg 1$ large enough such that

- (1) D_n is $(\varepsilon/4, E)$ -Følner;
- (2) $|D_n| > \lambda|C_0|$, where $\lambda = \max \{8/\varepsilon, 2\}$.

By Euclidean division, there exist $d, r \in \mathbb{N}$ such that $|D_n| = d|C_0| + r$ with $r < |C_0|$. Let $g_1, \dots, g_d \in G$ such that $\{C_0g_i\}_i$ are pairwise disjoint. We put $C' := \bigsqcup_{i=1}^d C_0g_i$. Then C' is (ε, F) -Følner and $|C'| = d|C_0|$.

Now let $D' := D_n - \{x_1, \dots, x_r\}$ be a subset of D_n obtained from D_n by deleting any r elements of D_n . Then $|D'| = |D_n| - r = d|C_0| = |C'|$. We claim that D' is (ε, E) -Følner. Indeed, we have

$$\frac{|D_n|}{|D'|} = \frac{|D_n|}{d|C_0|} = \frac{d|C_0| + r}{d|C_0|} = 1 + \frac{r}{d|C_0|} < 1 + \frac{1}{d} \leq 2,$$

since $r < |C_0|$ and $|D_n| = d|C_0| + r > \lambda|C_0|$ so $d > \lambda - 1 \geq 1$ by definition of λ . In addition, we have

$$\frac{|D' \Delta D_n|}{|D_n|} = \frac{r}{|D_n|} < \frac{r}{\lambda|C_0|} < \frac{1}{\lambda} \leq \frac{\varepsilon}{8}.$$

Therefore,

$$\begin{aligned} \frac{|D' \Delta hD'|}{|D'|} &\leq \frac{|D_n|}{|D'|} \left(\frac{|D' \Delta D_n|}{|D_n|} + \frac{|D_n \Delta hD_n|}{|D_n|} + \frac{|hD_n \Delta hD'|}{|D_n|} \right) \\ &< 2 \left(\frac{\varepsilon}{8} + \frac{\varepsilon}{4} + \frac{\varepsilon}{8} \right) = \varepsilon, \end{aligned}$$

for every $h \in E$. ■

Lemma 6. *If there exist amenable actions of G and H , then there exist G -action and H -action such that the actions admit Følner sequences $\{C_n\}_{n \geq 1}$ for the G -action and $\{D_n\}_{n \geq 1}$ for the H -action with $|C_n| = |D_n|, \forall n \geq 1$.*

Proof. If $G \curvearrowright X$ amenably, we replace X by a disjoint union of infinitely many copies of X and use the same idea as in the proof of Lemma 5; if $C_0 \in X$ is (ε, F) -Følner, we put C' to be d copies of C_0 in d disjoint copies of X . ■

Now we are ready to prove Theorem 2:

Corollary 7. *Let G and H be countable groups and A be a common finite subgroup of G and H . If G is an infinite amenable group and there is a H -set Y such that the H -action is amenable and the action of A on Y is free, then $(G, H, A) \in \mathcal{A}'$.*

Proof. Let $\{C_n\}_{n \geq 1}$ be a Følner sequence of G . First of all, we can suppose that the sequence $\{A \cdot C_n\}_{n \geq 1}$ is pairwise disjoint. Indeed, if $\{C_n\}_{n \geq 1}$ is a Følner sequence of G , we define $\{C'_n\}_{n \geq 1}$ inductively on n ; let $C'_1 := C_1$ and for every $n \geq 1$, we choose $h_n \in G$ such that $AC_{n+1}h_n \cap (\cup_{i=1}^n AC'_i) = \emptyset$, and we set $C'_{n+1} := C_{n+1}h_n$. Moreover, if $\{D_n\}_{n \geq 1} \subset Y$ is a Følner sequence for H -action, we can also suppose that the sequence $\{A \cdot D_n\}_{n \geq 1}$ is pairwise disjoint. Indeed, let $Y_0 = \sqcup_{i \geq 1} Y_i$ where $Y_i = Y, \forall i \geq 1$ be a disjoint union of infinitely many copies of Y . Then the H -action on Y_0 is amenable with a Følner sequence $D_n \subset Y_n, \forall n \geq 1$ such that $\{A \cdot D_n\}_{n \geq 1}$ is pairwise disjoint. In addition, by Lemma 6 we can suppose that $|C_n| = |D_n|, \forall n \geq 1$, so that the condition (iii) in Definition 2.1 is verified.

Now we consider the H -action on $Y' := H \sqcup Y_0$. The action satisfies the condition (ii) of Definition 2.1 (for the H -action) since the H -action on itself is free, and the action of A on Y' is free (since the A -action on H is clearly free and the A -action on Y_0 is free by assumption). Finally let $G = X$. The G -action on itself is transitive and free. Thus the triple (G, H, A) is in \mathcal{A}' . ■

Corollary 8. *If G is infinite amenable group and H contains a finite index normal subgroup N with $N \cap A = \{1\}$, then $(G, H, A) \in \mathcal{A}'$.*

It follows from Corollary 7 by taking $Y = H/N$. So for example if H is residually finite, applying Proposition 4 we see that $G *_A H$ is in \mathcal{A} for every finite subgroup A of G and H . Besides, with $A = \{1\}$, we find a particular case of the result of Glasner-Monod; if G is amenable, then $G * H \in \mathcal{A}$ for every countable group H .

Corollary 9. *Let G, H be amenable groups and let A be a common finite subgroup of G and H . Then the amalgamated free product $G *_A H$ is in \mathcal{A} .*

Proof. The cases where G or H is infinite follow from Corollary 7 and Proposition 4. So let G and H be finite groups. From a result of Baumslag [1], if G and H are finite groups, then the amalgamated free product $G *_A H$ contains a free subgroup of finite index. So $G *_A H$ is in \mathcal{A} since \mathcal{A} is closed under the extension of co-amenable subgroup. ■

Remark 2.2. When G is a finitely generated group with polynomial growth, Lemma 5 can be strengthened. Indeed, in this case G admits Følner sequences of any prescribed size. More precisely, let $\{a_n\}_{n \geq 1}$ be a strictly ascending sequence of positive integers. Let G be an infinite finitely generated group with polynomial growth. Then G has a Følner sequence $\{F_n\}_{n \geq 1}$ such that $|F_n| = a_n, \forall n \geq 1$.

Proof. Let S be a finite symmetric generating set of G . Denote $B(k)$ the ball of radius k centered at 1 in the Cayley graph $\mathcal{G}(G, S)$. Let k_n such that $|B(k_n)| \leq a_n < |B(k_n + 1)|$. We choose a finite subset K_n such that $K_n \cap B(k_n) = \emptyset$ and $|K_n| = a_n - |B(k_n)|$, and set $F_n := B(k_n) \cup K_n, \forall n \geq 1$. Recall that the boundary

∂A of A is the set of edges (s, t) such that $s \in A$ and $t \notin A$. We have

$$\begin{aligned} \frac{|\partial F_n|}{|F_n|} &\leq \frac{|\partial B(k_n)|}{|F_n|} + \frac{|\partial K_n|}{|F_n|} \leq \frac{|\partial B(k_n)|}{|B(k_n)|} + \frac{|\partial K_n|}{|B(k_n)|} \\ &\leq \frac{|\partial B(k_n)|}{|B(k_n)|} + |S| \frac{|K_n|}{|B(k_n)|} \leq \frac{|\partial B(k_n)|}{|B(k_n)|} + |S| \frac{|B(k_n + 1)| - |B(k_n)|}{|B(k_n)|} \\ &\leq \frac{|\partial B(k_n)|}{|B(k_n)|} + |S| \frac{|\partial B(k_n)|}{|B(k_n)|} \leq (1 + |S|) \frac{|\partial B(k_n)|}{|B(k_n)|}. \end{aligned}$$

By a result of Pansu [6], a group with polynomial growth with degree d satisfies $\frac{|B(k)|}{k^d} \xrightarrow[k \rightarrow \infty]{} C$, for some $C > 0$. Thus in such a group, the sequence of all balls is a Følner sequence. This applies to our case and thus $|\partial F_n|/|F_n| \xrightarrow[k \rightarrow \infty]{} 0$. ■

2.3 Central extensions

From the idea of Lemma 7.3.1 in [5], we have:

Lemma 10. *Let $1 \rightarrow Z \rightarrow G \xrightarrow{p} Q \rightarrow 1$ be a central extension. Suppose that there is a co-amenable subgroup $H < Q$ such that $H \in \mathcal{A}$ (so Q is also in \mathcal{A}) and the central extension splits over H (i.e. the central extension $1 \rightarrow Z \rightarrow p^{-1}(H) \xrightarrow{p} H \rightarrow 1$ splits). Then $G \in \mathcal{A}$.*

Proof. Since the extension splits over H , the group $p^{-1}(H)$ is a semi-direct product of Z and H , but Z is central so $p^{-1}(H)$ is indeed isomorphic to the direct product $Z \times H$. Since $Z \in \mathcal{A}$ (Z is amenable) and $H \in \mathcal{A}$ by assumption, the group $p^{-1}(H)$ is in \mathcal{A} . Moreover, the map $G/p^{-1}(H) \rightarrow Q/H$ defined by $gp^{-1}(H) \mapsto p(g)H$ is G -equivariant and bijective, so the co-amenable of H in Q implies the co-amenable of $p^{-1}(H)$ in G , thus $G \in \mathcal{A}$. ■

Example 2.1. (Example 7.3.4. in [5]) For $p, q \geq 2$, a Torus knot group $\Gamma_{p,q}$ is the group with the presentation $\Gamma_{p,q} = \langle x, y | x^p = y^q \rangle$. There is a central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \Gamma_{p,q} \rightarrow \mathbb{Z}/p\mathbb{Z} * \mathbb{Z}/q\mathbb{Z} \rightarrow 1,$$

where $\mathbb{Z} \simeq \langle x^p = y^q \rangle$, $\mathbb{Z}/p\mathbb{Z} \simeq \langle x | x^p = 1 \rangle$ and $\mathbb{Z}/q\mathbb{Z} \simeq \langle y | y^q = 1 \rangle$.

The free product $\mathbb{Z}/p\mathbb{Z} * \mathbb{Z}/q\mathbb{Z}$ has a finite index free subgroup \mathbb{F} over which the central extension splits. Thus the group $\Gamma_{p,q}$ is in \mathcal{A} by Lemma 10.

Example 2.2. Let M be a 3-manifold which is constructed as a fiber bundle over a closed orientable surface with fiber a circle. Such a 3-manifold is an orientable Seifert fibred space. There is a central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \pi_1(M) \xrightarrow{p} \Gamma_g \rightarrow 1,$$

where Γ_g is the fundamental group of the closed orientable surface of genus $g \geq 2$. The derived subgroup Γ'_g is co-amenable in Γ_g and $\Gamma'_g \in \mathcal{A}$ since Γ'_g is free (since $[\Gamma_g : \Gamma'_g]$ is infinite). Thus we have a central extension that splits:

$$1 \rightarrow \mathbb{Z} \rightarrow p^{-1}(\Gamma'_g) \xrightarrow{p} \Gamma'_g \rightarrow 1.$$

Therefore $\pi_1(M)$ is in \mathcal{A} by Lemma 10.

References

- [1] G. Baumslag, *On the residual finiteness of generalised free products of nilpotent groups*, Bull. Amer. Math. Soc. **75** (1969), 305–316.
- [2] Y. Glasner and N. Monod, *Amenable actions, free products and a fixed point property*, Bull. Lond. Math. Soc. **39** (2007), no. 1, 138–150.
- [3] S. Moon, *Amenable actions of cyclically pinched one-relator group and generic property*, Preprint 2009, arXiv:0909.2824.
- [4] ———, *Amenable actions of amalgamated free products*, Groups, Geom. Dyn. **4** (2010), no. 2, 309–332.
- [5] P.-A. Cherix, M. Cowling, P. Jolissaint, P. Julg and A. Valette, *Groups with the Haagerup Property : Gromov's a -T-menability*, Progress in Mathematics, vol. 197, Birkhäuser Verlag, Basel, 2001.
- [6] P. Pansu, *Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un*, Ann. of Math. **129** (1989), 1–60.
- [7] J.-P. Serre, *Arbres, amalgames, SL_2* , vol. astésisque 46, Société Mathématique de France, 1977.

Institut de Mathématiques de Bourgogne
UMR 5584 du CNRS
Université de Bourgogne
9 avenue Alain Savary - BP 47870
21078 Dijon cedex
France
E-mail : soyoung.moon@u-bourgogne.fr