

Classification of simple quartics up to equisingular deformation

Çisem GÜNEŞ AKTAŞ

(Received September 4, 2015)

(Revised December 1, 2016)

ABSTRACT. We study complex spatial quartic surfaces with simple singularities up to equisingular deformations; as a first step, give a complete equisingular deformation classification of non-special simple quartic surfaces.

1. Introduction

1.1. Principal results. Throughout the paper, all algebraic varieties are over the field \mathbb{C} of complex numbers. A *quartic* is a surface in \mathbb{P}^3 of degree 4. We confine ourselves to *simple* quartics only, *i.e.*, those with **A–D–E** type singularities (see §3.1). Two such quartics are said to be *equisingular deformation equivalent* if they belong to the same deformation family in which the total Milnor number stays constant.

Four seems to be the highest degree where one can hope to obtain a complete equisingular deformation classification. Quartics with a non-simple singular point (which are typically rational or ruled) have been treated in A. Degtyarev [3, 4], where the complete classification is obtained and described in terms of lattice embeddings. An alternative description of some non-simple sets of singularities in terms of Dynkin diagrams can be found in T. Urabe [23, 24]. *Simple* quartics are *K3*-surfaces, and as such they can be studied by using the global Torelli theorem [21] and the surjectivity of the period map [13], combined with V. V. Nikulin’s theory of discriminant forms [19]. This approach was used by Urabe [25, 26], who showed that the total Milnor number μ of a simple quartic does not exceed 19 and listed (in terms of perturbations of Dynkin graphs) all realizable sets of singularities with the total Milnor number $\mu \leq 17$. From a slightly different perspective, also a worth mentioning is the study of simple *real* quartics up to *equivariant* equisingular deformation, see, e.g., the classification of nonsingular real quartics by V. Kharlamov [12] or the recent classification of the arrangements of the ten nodes of a real determinantal quartic by A. Degtyarev and I. Itenberg [7].

The author was supported by *TÜBİTAK-BİDEB 2211-Yurt İçi Yüksek Lisans Burs Programı* and the *TÜBİTAK* grant 114F325.

2010 *Mathematics Subject Classification*. Primary 14J28; Secondary 14J10, 14J17.

In the present paper, we start a systematic study of the equisingular stratification of the space of simple quartics. The principal difficulty here is the great number of strata (about 12 thousands); thus, as a first step, we confine ourselves to the so-called *non-special* quartics. The precise definition is rather technical, and we postpone it till §3.3. In a sense, the non-special quartics are an analogue of irreducible plane sextic curves admitting no dihedral coverings, cf. [1]; in a similar vein, we have the following geometric characterization, which is proved in §4.1.

THEOREM 1. *A simple quartic $X \subset \mathbb{P}^3$ is non-special if and only if*

$$H_1(X \setminus (\text{Sing } X \cup H)) = 0,$$

where $\text{Sing } X$ is the set of the singular points of X and H is a generic hyperplane section of X .

Still, the number of strata is too large to be listed explicitly, and; hence, in the existence part we adopt the approach of [1] and describe only those strata that are extremal with respect to degeneration. (Note that, unlike [23, 24, 25, 26], we do not introduce any artificial Dynkin graphs: all sets of singularities mentioned below and all perturbations thereof are indeed realized by quartics.) Recall that a set of simple singularities can be identified with a root system, i.e., a negative definite lattice generated by vectors of square -2 (see Dufree [8] and §3.1); the rank of this lattice is the total Milnor number of the quartic. By a *perturbation* of a set of simple singularities \mathbf{S} we mean any set of simple singularities \mathbf{S}' whose Dynkin graph is an induced subgraph of that of \mathbf{S} (see §4.2). Recall, further, that for a simple quartic $X \subset \mathbb{P}^3$, one has $\mu(X) \leq 19$ (see e.g., [25]); X is called *maximizing* if $\mu(X) = 19$.

Denote by $\mathcal{M}(\mathbf{S})$ the equisingular stratum of simple quartics with a given set of singularities \mathbf{S} . A connected component $\mathcal{D} \subset \mathcal{M}(\mathbf{S})$ is called *real* if it is preserved as a set under the complex conjugation map $\text{conj} : \mathbb{P}^3 \rightarrow \mathbb{P}^3$. Clearly, this property is independent of the choice of the coordinates in \mathbb{P}^3 , and all components of $\mathcal{M}(\mathbf{S})$ split into real and pairs of complex conjugate ones.

Our principal result is a complete description of the equisingular strata $\mathcal{M}_1(\mathbf{S})$ of non-special simple quartics.

THEOREM 2. *A set of singularities \mathbf{S} is realizable as the set of singularities of a non-special simple quartic if and only if \mathbf{S} can be obtained by a perturbation from one of the sets of singularities listed in Tables 1 and 2. The numbers (r, c) of, respectively, real and pairs of complex conjugate components of the strata $\mathcal{M}_1(\mathbf{S})$ with $\mu(\mathbf{S}) = 19$ are shown in Table 1. If \mathbf{S} is one of*

$$\mathbf{D}_6 \oplus 2\mathbf{A}_6, \quad \mathbf{D}_5 \oplus 2\mathbf{A}_6 \oplus \mathbf{A}_1, \quad 2\mathbf{A}_7 \oplus 2\mathbf{A}_2, \quad 3\mathbf{A}_6, \quad 2\mathbf{A}_6 \oplus 2\mathbf{A}_3$$

Table 1. The space $\mathcal{M}_1(\mathbf{S})$ with $\mu(\mathbf{S}) = 19$

Singularities	(r, c)	Singularities	(r, c)
$2\mathbf{E}_8 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$(1, 0)$	$\mathbf{A}_{18} \oplus \mathbf{A}_1$	$(1, 1)^2$
$\mathbf{E}_8 \oplus \mathbf{E}_7 \oplus \mathbf{A}_4$	$(1, 0)$	$\mathbf{A}_{17} \oplus \mathbf{A}_2$	$(1, 1)$
$\mathbf{E}_8 \oplus \mathbf{E}_6 \oplus \mathbf{D}_5$	$(1, 0)$	$\mathbf{A}_{16} \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$(1, 0)$
$\mathbf{E}_8 \oplus \mathbf{E}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_1$	$(1, 0)$	$\mathbf{A}_{15} \oplus 2\mathbf{A}_2$	$(0, 1)$
$\mathbf{E}_8 \oplus \mathbf{D}_7 \oplus 2\mathbf{A}_2$	$(1, 0)$	$\mathbf{A}_{14} \oplus \mathbf{A}_5$	$(0, 2)$
$\mathbf{E}_8 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_1$	$(1, 0)$	$\mathbf{A}_{14} \oplus \mathbf{A}_3 \oplus \mathbf{A}_2$	$(0, 2)$
$\mathbf{E}_8 \oplus \mathbf{A}_9 \oplus \mathbf{A}_2$	$(1, 0)$	$\mathbf{A}_{13} \oplus \mathbf{A}_6$	$(0, 2)$
$\mathbf{E}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_5$	$(1, 0)$	$\mathbf{A}_{13} \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$	$(1, 0)$
$\mathbf{E}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_1$	$(1, 0)$	$\mathbf{A}_{12} \oplus \mathbf{A}_6 \oplus \mathbf{A}_1$	$(1, 1)^2$
$\mathbf{E}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2$	$(0, 1)$	$\mathbf{A}_{12} \oplus \mathbf{A}_5 \oplus \mathbf{A}_2$	$(1, 1)^2$
$\mathbf{E}_8 \oplus 2\mathbf{A}_4 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$(1, 0)$	$\mathbf{A}_{12} \oplus \mathbf{A}_4 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$(0, 1)$
$\mathbf{E}_7 \oplus \mathbf{E}_6 \oplus \mathbf{A}_6$	$(1, 0)$	$\mathbf{A}_{12} \oplus \mathbf{A}_3 \oplus 2\mathbf{A}_2$	$(2, 0)$
$\mathbf{E}_7 \oplus \mathbf{A}_{12}$	$(1, 1)^2$	$\mathbf{A}_{11} \oplus \mathbf{A}_6 \oplus \mathbf{A}_2$	$(0, 2)$
$\mathbf{E}_7 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_2$	$(0, 1)$	$\mathbf{A}_{10} \oplus \mathbf{A}_9$	$(1, 1)^2$
$\mathbf{E}_7 \oplus \mathbf{A}_8 \oplus \mathbf{A}_4$	$(2, 0)^2$	$\mathbf{A}_{10} \oplus \mathbf{A}_8 \oplus \mathbf{A}_1$	$(0, 1)$
$\mathbf{E}_7 \oplus 2\mathbf{A}_6$	$(0, 1)$	$\mathbf{A}_{10} \oplus \mathbf{A}_7 \oplus \mathbf{A}_2$	$(0, 2)$
$\mathbf{E}_7 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$	$(1, 0)$	$\mathbf{A}_{10} \oplus \mathbf{A}_6 \oplus \mathbf{A}_3$	$(0, 2)$
$2\mathbf{E}_6 \oplus \mathbf{D}_7$	$(1, 0)$	$\mathbf{A}_{10} \oplus \mathbf{A}_6 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$(1, 0)$
$\mathbf{E}_6 \oplus \mathbf{D}_{13}$	$(1, 0)$	$\mathbf{A}_{10} \oplus \mathbf{A}_5 \oplus \mathbf{A}_4$	$(1, 0)$
$\mathbf{E}_6 \oplus \mathbf{D}_9 \oplus \mathbf{A}_4$	$(1, 0)$	$\mathbf{A}_{10} \oplus \mathbf{A}_4 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2$	$(0, 1)$
$\mathbf{E}_6 \oplus \mathbf{A}_{13}$	$(1, 0)$	$\mathbf{A}_9 \oplus \mathbf{A}_8 \oplus \mathbf{A}_2$	$(1, 1)$
$\mathbf{E}_6 \oplus \mathbf{A}_{12} \oplus \mathbf{A}_1$	$(1, 0)$	$\mathbf{A}_9 \oplus \mathbf{A}_6 \oplus 2\mathbf{A}_2$	$(1, 0)$
$\mathbf{D}_{15} \oplus 2\mathbf{A}_2$	$(1, 0)$	$\mathbf{A}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_5$	$(1, 1)$
$\mathbf{D}_{11} \oplus \mathbf{A}_6 \oplus \mathbf{A}_2$	$(0, 1)$	$\mathbf{A}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_1$	$(0, 1)$
$\mathbf{D}_9 \oplus \mathbf{A}_6 \oplus 2\mathbf{A}_2$	$(1, 0)$	$\mathbf{A}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2$	$(0, 3)$
$\mathbf{D}_7 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_2$	$(0, 1)$	$\mathbf{A}_7 \oplus 2\mathbf{A}_6$	$(0, 2)$
$\mathbf{D}_7 \oplus 2\mathbf{A}_6$	$(0, 1)$	$\mathbf{A}_7 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$	$(0, 1)$
$\mathbf{D}_7 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$	$(0, 1)$	$2\mathbf{A}_6 \oplus \mathbf{A}_5 \oplus \mathbf{A}_2$	$(2, 0)$
$\mathbf{D}_7 \oplus 2\mathbf{A}_4 \oplus 2\mathbf{A}_2$	$(1, 0)$	$2\mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$(0, 1)$
		$\mathbf{A}_6 \oplus 2\mathbf{A}_4 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2$	$(2, 0)$

Table 2. Extremal sets of singularities with $\mu(\mathbf{S}) = 18$

$\mathbf{E}_8 \oplus \mathbf{D}_{10}$	$\mathbf{D}_{14} \oplus \mathbf{A}_4$	$\mathbf{D}_6 \oplus 3\mathbf{A}_4$
$\mathbf{E}_8 \oplus \mathbf{D}_9 \oplus \mathbf{A}_1$	$\mathbf{D}_{10} \oplus \mathbf{A}_8$	$2\mathbf{D}_5 \oplus \mathbf{A}_8$
$2\mathbf{E}_7 \oplus 2\mathbf{A}_2$	$\mathbf{D}_{10} \oplus 2\mathbf{A}_4$	$2\mathbf{D}_5 \oplus 2\mathbf{A}_4$
$\mathbf{E}_7 \oplus \mathbf{D}_{11}$	$2\mathbf{D}_9$	$\mathbf{D}_5 \oplus \mathbf{A}_9 \oplus \mathbf{A}_4$
$\mathbf{E}_7 \oplus \mathbf{D}_9 \oplus \mathbf{A}_2$	$\mathbf{D}_9 \oplus \mathbf{A}_9$	$\mathbf{D}_5 \oplus \mathbf{A}_8 \oplus \mathbf{A}_5$
\mathbf{D}_{18}	$\mathbf{D}_9 \oplus \mathbf{A}_8 \oplus \mathbf{A}_1$	$\mathbf{D}_5 \oplus \mathbf{A}_5 \oplus 2\mathbf{A}_4$
$\mathbf{D}_{17} \oplus \mathbf{A}_1$		

then $\mathcal{M}_1(\mathbf{S})$ consists of two complex conjugate components; in all other cases, the stratum $\mathcal{M}_1(\mathbf{S})$ is connected.

Theorem 2 is proved in §4.2. In Table 1, we list the maximizing sets of singularities, which are all extremal. In most cases the transcendental lattice (the orthogonal complement of the Néron Severi lattice in $H_2(X)$) is unique in its genus; in six cases marked with ² in the table the genus consists of two lattices. Recall that each maximizing quartic surface is defined over an algebraic number field and this number of transcendental lattices is a lower bound for the degree of this field.

A complete list of all possible combinations of simple singularities realized by a complex quartic surface (not-necessarily non-special) was previously found by Yang [28]. His technique is also based on Nikulin's criterion of existence of lattices [19] and he also represents the result in terms of perturbations of certain extremal sets. We restrict our attention to non-special quartics and our extremal sets of singularities are extremal in this restricted class. For this reason, we can also assert that any perturbation of the set S of simple singularities is realized by a perturbation of simple quartics.

1.2. Contents of the paper. Our principal result, Theorem 2, is proved by a reduction to an arithmetical problem [7] (*cf.* also [5]), followed by Nikulin's theory of lattice extensions and Nikulin's existence theorem [19]. The principal novelty is a systematic usage of the Miranda–Morrison theory [15, 16, 17] computing the genus groups and a few other bits missing in [19] in the case of indefinite lattices.

In §2, based on Nikulin's work [19], we recall the basic notions and results about integral lattices, discriminant forms and lattice extensions; then, we outline the fundamentals of Miranda–Morrison's theory [17] which are used in §4.2. In §3, we discuss the relation between simple quartics and $K3$ -surfaces, explain the notion of abstract homological type, and recall the reduction of the classification problem to the arithmetical classification of abstract homological types. Finally, §4 is devoted to the proofs of our principal results: the proof of Theorem 1 is purely homotopy theoretical, whereas that of Theorem 2 depends essentially on the auxiliary material presented in §2 and §3.

1.3. Acknowledgements. I would like to express my gratitude to my advisor Alex Degtyarev for attracting my attention to the problem, motivating discussions, encouragement and infinite patience. I am also thankful to him for sharing his results (stated in Table 1) about the moduli space of maximizing non-special simple quartics.

I would like to thank the referee for his/her constructive comments and suggestions which significantly contributed to improving the quality of this paper.

2. Preliminaries

2.1. Finite quadratic forms. A *finite quadratic form* is a finite abelian group \mathcal{L} equipped with a map $q : \mathcal{L} \rightarrow \mathbb{Q}/2\mathbb{Z}$ satisfying $q(x+y) = q(x) + q(y) + 2b(x, y)$ and $q(nx) = n^2x$ for all $x, y \in \mathcal{L}$, where $b : \mathcal{L} \otimes \mathcal{L} \rightarrow \mathbb{Q}/\mathbb{Z}$ is a symmetric bilinear form (which is determined by q). To reduce the notation we write x^2 for $q(x)$ and $x \cdot y$ for $b(x, y)$. For a prime p , let $\mathcal{L}_p := \mathcal{L} \otimes \mathbb{Z}_p$, which is called the *p-primary part* of \mathcal{L} . Any finite quadratic form \mathcal{L} can be written as an orthogonal sum of its *p*-primary components \mathcal{L}_p , i.e., $\mathcal{L} = \bigoplus_p \mathcal{L}_p$ where the summation runs over all primes p . Denote by $\ell(\mathcal{L})$ the minimal number of generators of \mathcal{L} .

Consider a fraction $\frac{m}{n} \in \mathbb{Q}/2\mathbb{Z}$ with $\gcd(m, n) = 1$ and $mn \equiv 0 \pmod{2}$. By $\langle \frac{m}{n} \rangle$, we denote the finite non-degenerate (see §2.2) quadratic form on $\mathbb{Z}/n\mathbb{Z}$ generated by an element of square $\frac{m}{n}$ and of order n . For an integer $k \geq 1$, let $\mathcal{U}(2^k)$ and $\mathcal{V}(2^k)$ be the quadratic forms on $\mathbb{Z}/2^k\mathbb{Z} \oplus \mathbb{Z}/2^k\mathbb{Z}$ defined by the matrices

$$\mathcal{U}(2^k) = \begin{pmatrix} 0 & \frac{1}{2^k} \\ \frac{1}{2^k} & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{V}(2^k) = \begin{pmatrix} \frac{1}{2^{k-1}} & \frac{1}{2^k} \\ \frac{1}{2^k} & \frac{1}{2^{k-1}} \end{pmatrix}.$$

Nikulin [19] showed that any finite quadratic form can be written as an orthogonal sum of cyclic summands of the form $\langle \frac{m}{n} \rangle$ and copies of $\mathcal{U}(2^k)$ and $\mathcal{V}(2^k)$.

The *Brown invariant* of a finite quadratic form \mathcal{L} is the residue $\text{Br } \mathcal{L} \in \mathbb{Z}/8\mathbb{Z}$ defined by the Gauss sum

$$\exp\left(\frac{1}{4}i\pi \text{Br } \mathcal{L}\right) = |\mathcal{L}|^{-1/2} \sum_{x \in \mathcal{L}} \exp(i\pi x^2).$$

The Brown invariants of indecomposable *p*-primary blocks are as follows:

$$\text{Br} \left\langle \frac{2a}{p^{2s-1}} \right\rangle = 2 \left(\frac{a}{p} \right) - \left(\frac{-1}{p} \right) - 1, \quad \text{Br} \left\langle \frac{2a}{p^{2s}} \right\rangle = 0 \quad (\text{for } p \text{ odd, } s \geq 1 \text{ and } \gcd(a, p) = 1),$$

$$\text{Br} \left\langle \frac{a}{2^k} \right\rangle = a + \frac{1}{2}k(a^2 - 1) \pmod{8} \quad (\text{for } k \geq 1 \text{ and odd } a \in \mathbb{Z}),$$

$$\text{Br } \mathcal{U}(2^k) = 0,$$

$$\text{Br } \mathcal{V}(2^k) = 4k \pmod{8} \quad (\text{for all } k \geq 1).$$

A finite quadratic form is called *even* if $x^2 \equiv 0 \pmod{2}$ for all elements $x \in \mathcal{L}$ of order two; otherwise it is called *odd*. This definition implies that a

quadratic form is odd if and only if it contains $\langle \pm \frac{1}{2} \rangle$ as an orthogonal summand.

2.2. Integral lattices and discriminant forms. An (*integral*) *lattice* is a free abelian group L of finite rank with a symmetric bilinear form $b : L \otimes L \rightarrow \mathbb{Z}$. For short, we use the multiplicative notation $x \cdot y$ for $b(x, y)$ and x^2 for $b(x, x)$. A lattice L is called *even* if a^2 is an even integer for all $a \in L$. It is called *odd* otherwise. The determinant $\det(L)$ is defined to be the determinant of the Gram matrix of b in any basis of L . Since the transition matrix between any two integral bases has determinant ± 1 , the determinant of $\det(L) \in \mathbb{Z}$ of L is well defined. A lattice L is called *non-degenerate* if $\det(L) \neq 0$; it is called *unimodular* if $\det(L) = \pm 1$.

From now on all lattices considered are even and non-degenerate.

Given a lattice L , the form $b : L \otimes L \rightarrow \mathbb{Z}$ can be extended by linearity to a form $(L \otimes \mathbb{Q}) \otimes_{\mathbb{Q}} (L \otimes \mathbb{Q}) \rightarrow \mathbb{Q}$. If L is non-degenerate, the dual group $L^* := \text{Hom}(L, \mathbb{Z})$ can be identified with the subgroup

$$\{x \in L \otimes \mathbb{Q} \mid x \cdot y \in \mathbb{Z} \text{ for all } y \in L\}$$

Since the original bilinear form b on L is integer valued, L is a finite index subgroup of its dual. The quotient L^*/L is called the *discriminant group* of L and is denoted by \mathcal{L} or *disc* L . If $\{e_1, e_2, \dots, e_n\}$ is a basis set for L and $\{e_1^*, e_2^*, \dots, e_n^*\}$ is the dual basis for L^* , then the Gram matrix $[e_i \cdot e_j]$ is exactly the matrix of the homomorphism $\varphi : L \rightarrow L^*$, $x \mapsto [y \mapsto x \cdot y]$. Hence one has $|\mathcal{L}| = |\det(L)|$. Note that $x \cdot y \in \mathbb{Z}$ whenever $x \in L$ or $y \in L$. Thus, \mathcal{L} inherits from $L \otimes \mathbb{Q}$ a non-degenerate symmetric bilinear form $b_{\mathcal{L}} : \mathcal{L} \otimes \mathcal{L} \rightarrow \mathbb{Q}/\mathbb{Z}$; it is called the *discriminant form*. If L is even, this form $b_{\mathcal{L}}$ can be promoted to the quadratic extension $q_{\mathcal{L}} : \mathcal{L} \rightarrow \mathbb{Q}/2\mathbb{Z}$, $(x \bmod L) \mapsto (x^2 \bmod 2\mathbb{Z})$. Hence, the discriminant form of an even lattice is a finite quadratic form. Accordingly, given a prime p , we use the notation $\text{disc}_p L$ or \mathcal{L}_p for the p -primary part of \mathcal{L} , i.e., $\mathcal{L}_p = \mathcal{L} \otimes \mathbb{Z}_p$. Each discriminant group \mathcal{L} decomposes into orthogonal sum $\mathcal{L} = \bigoplus_p \mathcal{L}_p$ of its p -primary components. The *signature* of a non-degenerate lattice L is the pair (σ_+, σ_-) of its positive and negative inertia indices. Two non-degenerate integral lattices are said to have the same *genus* if their localizations over \mathbb{R} and over \mathbb{Z}_p are isomorphic. The following few statements give the relation between the genus of an even integral lattice and its discriminant form.

THEOREM 3 (Nikulin [19]). *The genus of an even integral lattice L is determined by its signature $(\sigma_+ L, \sigma_- L)$ and discriminant form $\text{disc } L$.*

The existence of an even integral lattice L with a given signature is given by Nikulin's existence theorem (see Theorem 5).

THEOREM 4 (van der Blij [27]). *For any non-degenerate even integral lattice L , one has $\text{Br } \mathcal{L} = \sigma_+ - \sigma_- \pmod{8}$.*

We denote by $g(L)$ the set of all isomorphism classes of all non-degenerate even integral lattices with the same genus as L . Each set $g(L)$ is known to contain finitely many isomorphism classes.

Given a prime p , we define the *determinant* $\det_p(\mathcal{L})$ as the determinant of the matrix of the quadratic form on \mathcal{L}_p in an appropriate basis (see [18] and [19] for details). Unless $p = 2$ one has $\det_p(\mathcal{L}) = u/|\mathcal{L}_p|$ where u is a well defined element of $u \in \mathbb{Z}_p^\times / (\mathbb{Z}_p^\times)^2$. If $p = 2$, the determinant $\det_2(\mathcal{L})$ is well defined only if \mathcal{L}_2 is even.

THEOREM 5 (Nikulin [19]). *Let \mathcal{L} be a finite quadratic form and let σ_\pm be a pair of integers. Then, the following four conditions are necessary and sufficient for the existence of an even integral lattice L whose signature is (σ_+, σ_-) and whose discriminant form is \mathcal{L} :*

- (1) $\sigma_\pm \geq 0$ and $\sigma_+ + \sigma_- \geq \ell(\mathcal{L})$.
- (2) $\sigma_+ - \sigma_- = \text{Br } \mathcal{L} \pmod{8}$.
- (3) For each $p \neq 2$, either $\sigma_+ + \sigma_- > \ell_p(\mathcal{L})$ or $\det_p(\mathcal{L}) \equiv (-1)^{\sigma_-} \cdot |\mathcal{L}| \pmod{(\mathbb{Z}_p^\times)^2}$.
- (4) Either $\sigma_+ + \sigma_- > \ell_2(\mathcal{L})$, or \mathcal{L}_2 is odd, or $\det_2(\mathcal{L}) \equiv \pm |\mathcal{L}| \pmod{(\mathbb{Z}_2^\times)^2}$.

2.3. Automorphisms of lattices. An *isometry* of integral lattices is a homomorphism of abelian groups preserving the forms. The group of auto-isometries of L is denoted by $O(L)$. There is a natural homomorphism $d : O(L) \rightarrow \text{Aut}(\mathcal{L})$, where $\text{Aut}(\mathcal{L})$ denotes the group of automorphisms of \mathcal{L} preserving the discriminant form q on \mathcal{L} . Obviously, one has $\text{Aut}(\mathcal{L}) = \prod_p \text{Aut}(\mathcal{L}_p)$, where the product runs over all primes. The restrictions of d to the p -primary components are denoted by $d_p : O(L) \rightarrow \text{Aut}(\mathcal{L}_p)$.

Given a vector u in L with $u \neq 0$, the *reflection* against its orthogonal hyperplane is the automorphism

$$r_u : L \rightarrow L$$

$$x \mapsto x - 2 \frac{(x \cdot u)}{u^2} u$$

The reflection r_u is well-defined whenever $u \in \left(\frac{u^2}{2}\right)L^*$. Note that $r_u^2 = \text{id}$, i.e., r_u is an involution. Each image $d_p(r_u) \in \text{Aut}(\mathcal{L}_p)$ is also a reflection (see §2.6). If $u^2 = \pm 1$ or $u^2 = \pm 2$, then the induced automorphism $d(r_u)$ is the identity.

2.4. Lattice extensions. For a non-degenerate even lattice S , an even integral lattice L containing S is called an *extension* of S . An *isomorphism* between

two extensions $L_1 \supset S$ and $L_2 \supset S$ is an isometry between L_1 and L_2 taking S to S . In particular, if the isomorphism $L_1 \rightarrow L_2$ restricts to id on S , the extensions L_1 and L_2 are called *strictly isomorphic*. For a given subgroup A of $O(S)$, we define *A-isomorphisms* of extensions of S as those which restrict to an element of A on S .

Recall that S is assumed to be non-degenerate, hence given a *finite index* extension $L \supset S$, one has $L \subset S^*$. Thus there are inclusions $S \subset L \subset L^* \subset S^*$ which imply $L/S \subset S^*/S = \mathcal{S}$. The subgroup $\mathcal{K} = L/S$ of \mathcal{S} is called the *kernel* of the finite index extension $L \supset S$. Since L is an even integral lattice, the discriminant quadratic form on \mathcal{S} restricts to zero on \mathcal{K} , i.e., \mathcal{K} is isotropic.

PROPOSITION 1 (Nikulin [19]). *Let S be a non-degenerate even lattice, and fix a subgroup $A \subset O(S)$. The map $L \mapsto \mathcal{K} = L/S \subset \mathcal{S}$ establishes a one-to-one correspondence between the set of A -isomorphism classes of finite index extensions $L \supset S$ and the set of A -orbits of isotropic subgroups $\mathcal{K} \subset \mathcal{S}$. Under this correspondence one has $L = \{x \in S^* \mid (x \bmod S) \in \mathcal{K}\}$ and $\mathcal{S} = \mathcal{K}^\perp / \mathcal{K}$.*

PROPOSITION 2 (Nikulin [19]). *Let $L \supset S$ be a finite index extension of a lattice S and let $\mathcal{K} \subset \mathcal{S}$ be its kernel. Then an auto-isometry $S \rightarrow S$ extends to L if and only if the induced automorphism of \mathcal{S} preserves \mathcal{K} .*

An extension $L \supset S$ is called *primitive* if L/S is torsion free. Following Nikulin [19], we confine ourselves to the special case where L is unimodular. If S is a primitive non-degenerate sublattice of a unimodular lattice L then S^\perp is also primitive in L and L is a finite index extension of $S \oplus S^\perp$. Furthermore, since $\text{disc } L = 0$, the kernel $\mathcal{K} \subset \mathcal{S} \oplus \mathcal{S}^\perp$ is the graph of an anti-isometry $\psi : \mathcal{S} \rightarrow \text{disc } S^\perp$. Hence the genus $g(S^\perp)$ is determined by the genera $g(S)$ and $g(L)$. Conversely, given a lattice $N \in g(S^\perp)$ and an anti-isometry $\psi : \mathcal{S} \rightarrow \mathcal{N}$ where \mathcal{N} is the discriminant group of N , the graph of ψ is an isotropic subgroup $\mathcal{K} \subset \mathcal{S} \oplus \mathcal{S}^\perp$ and the corresponding finite index extension $S \oplus N \hookrightarrow L$ is a unimodular primitive extension of S with $S^\perp \cong N$. Any anti-isometry $\psi : \mathcal{S} \rightarrow \text{disc } S^\perp$ induces a homomorphism $d^\psi : O(S) \rightarrow \text{Aut}(\mathcal{N})$. Recall that there is a natural homomorphism $d : O(N) \rightarrow \text{Aut}(\mathcal{N})$. Thus, since also an indefinite unimodular lattice is unique in its genus, we have the following theorem.

THEOREM 6 (Nikulin [19]). *Let L be an indefinite unimodular even lattice and $S \subset L$ a non-degenerate primitive sublattice. Fix a subgroup $A \subset O(S)$. Then the A -isomorphism class of a primitive extension $S \subset L$ is determined by*

- (1) *a choice of a lattice $N \in g(S^\perp)$ and*
- (2) *a choice of a double coset $c_N \in d^\psi(A) \backslash \text{Aut}(\mathcal{N}) / \text{Im } d$ (for a given N and some anti-isometry $\psi : \mathcal{S} \rightarrow \mathcal{N}$ inducing d^ψ).*

THEOREM 7 (Nikulin [19]). *Let L be an indefinite unimodular even lattice, $S \subset L$ a non-degenerate primitive sublattice and $\psi : \mathcal{S} \rightarrow \mathcal{N}$ the anti-isometry where, $N = S^\perp$. Then a pair of isometries $a_S \in O(S)$ and $a_N \in O(N)$ extends to L if and only if $d^\psi(a_S) = d(a_N)$.*

2.5. Miranda–Morrison’s theory. Let p be a prime. Define

$$\begin{aligned} \Gamma_p &:= \{\pm 1\} \times \mathbb{Q}_p^\times / (\mathbb{Q}_p^\times)^2, \\ \Gamma_0 &:= \{\pm 1\} \times \{\pm 1\} \subset \{\pm 1\} \times \mathbb{Q}^\times / (\mathbb{Q}^\times)^2. \end{aligned}$$

It is convenient to introduce the following subgroups related to Γ_p :

- $\Gamma_{p,0} := \{(1, 1), (1, u_p), (-1, 1), (-1, u_p)\} \subset \Gamma_p$; here, p is odd and u_p is the only nontrivial element of $\mathbb{Z}_p^\times / (\mathbb{Z}_p^\times)^2$,
- $\Gamma_{2,0} := \{(1, 1), (1, 3), (1, 5), (1, 7), (-1, 1), (-1, 3), (-1, 5), (-1, 7)\} \subset \Gamma_2$,
- $\Gamma_p^{++} := \{1\} \times \mathbb{Z}_p^\times / (\mathbb{Z}_p^\times)^2 \subset \Gamma_{p,0}$,
- $\Gamma_{2,2} := \{(1, 1), (1, 5)\} \subset \Gamma_2^{++}$,
- $\Gamma'_{2,0} := \Gamma_{2,0} / \Gamma_{2,2}$ (and $\Gamma'_{p,0} := \Gamma_{p,0}$ for $p \neq 2$),
- $\Gamma_0^{--} := \{(1, 1), (-1, -1)\} \subset \Gamma_0$.

Let, further,

$$\Gamma_{\mathbb{A},0} := \prod_p \Gamma_{p,0} \subset \Gamma_{\mathbb{A}} := \Gamma_{\mathbb{A},0} \cdot \sum_p \Gamma_p$$

where “ \cdot ” denotes the sum of the subgroups. Note that

$$\Gamma_{\mathbb{A}} = \left\{ (d_p, s_p) \in \prod_p \Gamma_p \mid (d_p, s_p) \in \Gamma_{p,0} \text{ for almost all } p \right\}$$

The natural map $\mathbb{Q}^\times / (\mathbb{Q}^\times)^2 \rightarrow \mathbb{Q}_p^\times / (\mathbb{Q}_p^\times)^2$ induces canonical maps

$$\varphi_p : \Gamma_0 \rightarrow \Gamma_{p,0}. \quad (2.1)$$

Let N be an indefinite lattice with $\text{rk}(N) \geq 3$. We will use certain subgroups $\Sigma_p^\sharp(N) \subset \Gamma_{p,0}$ and $\Sigma_p(N) \subset \Gamma_p$. In the notation of [17] (which slightly differs from the notation in [15, 16]), one has $\Sigma_p^\sharp(N) := \Sigma^\sharp(N \otimes \mathbb{Z}_p)$ and $\Sigma_p(N) := \Sigma(N \otimes \mathbb{Z}_p)$; we refer the reader to [17] (see chapter 7, section 4) for the precise definitions. The subgroups $\Sigma_p^\sharp(N)$ are computed explicitly in [17] (see Theorems 12.1, 12.2, 12.3 and 12.4 in chapter 7).

Another object defined in [17] (see chapter 8, sections 5, 6 and 7) is the \mathbb{F}_2 -module

$$E(N) := \Gamma_{\mathbb{A},0} / \prod_p \Sigma_p^\sharp(N) \cdot \Gamma_0. \quad (2.2)$$

This module is finite. Indeed, following [17] (see Definition 7.4 in chapter 8), we call a prime p *regular* with respect to N if $\Sigma_p^\sharp(N) = \Gamma_{p,0}$. An important point here is the fact that a prime p is regular unless $p \mid \det(N)$; thus, (2.2) reduces to finitely many primes p :

$$E(N) = \prod_{p \mid \det(N)} \Gamma_{p,0} \bigg/ \prod_{p \mid \det(N)} \Sigma_p^\sharp(N) \cdot \Gamma_0. \quad (2.3)$$

THEOREM 8 (Miranda–Morrison [17]). *Let N be a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$. Then there is an exact sequence*

$$O(N) \xrightarrow{d} \text{Aut}(\mathcal{N}) \xrightarrow{e} E(N) \rightarrow g(N) \rightarrow 1, \quad (2.4)$$

where $g(N)$ is the genus group of N .

A simplified version of (2.3) computing the numeric invariants

$$e_p(N) := [\Gamma_{p,0} : \Sigma_p^\sharp(N)] \quad \text{and} \quad \tilde{\Sigma}_p(N) := \varphi_p^{-1}(\Sigma_p^\sharp(N)) \subset \Gamma_0,$$

can be found in [15, 16]. This gives us the size of the group $E(N)$: one has

$$|E(N)| = \frac{e(N)}{[\Gamma_0 : \tilde{\Sigma}(N)]} \quad (2.5)$$

where

$$e(N) := \prod_p e_p(N), \quad \tilde{\Sigma}(N) := \bigcap_p \tilde{\Sigma}_p(N),$$

and the product and intersection run over all primes p or, equivalently, over all primes $p \mid \det(N)$.

The following theorem can be deduced from Theorems 6 and 8.

THEOREM 9 (Miranda–Morrison [15, 16]). *Let S be a primitive sublattice of an even unimodular lattice L such that $N := S^\perp$ is a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$. Then the strict isomorphism classes of primitive extensions $S \hookrightarrow L$ are in a canonical one-to-one correspondence with the group $E(N)$.*

As explained §2.4, given a unimodular lattice L and a primitive sublattice $S \subset L$, one has an anti-isometry $\psi : \mathcal{S} \rightarrow \mathcal{N}$ (where $N = S^\perp$), which induces a homomorphism $d^\psi : O(S) \rightarrow \text{Aut}(\mathcal{N})$. If N is indefinite and $\text{rk}(N) \geq 3$, then $d^\psi(O(S)) \subset \text{Aut}(\mathcal{N})$ is a normal subgroup with abelian quotient (see (2.4)) and we have a homomorphism $d^\perp : O(S) \rightarrow \text{Aut}(\mathcal{N}) \xrightarrow{e} E(N)$ independent of the choice of an anti-isometry ψ . The next statement follows from Theorems 8 and 6.

COROLLARY 1. *Let S be a primitive sublattice of an even unimodular lattice L such that $N := S^\perp$ is a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$ and let $A \subset O(S)$ be a subgroup. Then, the A -isomorphism classes of primitive extensions $S \hookrightarrow L$ are in a one-to-one correspondence with the \mathbb{F}_2 -module coker $d^\perp|_A$.*

2.6. Reflections. Recall that $\text{Aut}(\mathcal{N}) = \prod_p \text{Aut}(\mathcal{N}_p)$ where p runs over all primes. Let s be a prime and $\alpha \in \mathcal{N}_s$ such that

$$s^k \alpha = 0 \quad \text{and} \quad \alpha^2 = \frac{2u}{s^k} \bmod 2\mathbb{Z}, \quad \text{g.c.d}(u, s) = 1, \quad k \in \mathbb{N}. \quad (2.6)$$

We denote by \mathcal{N}_s^\dagger the set of all elements $\alpha \in \mathcal{N}_s$ satisfying (2.6) and let $\mathcal{N}^\dagger = \bigcup_s \mathcal{N}_s^\dagger$. Then one can define a map,

$$\mathcal{N}_s \rightarrow \mathbb{Z}/s^k\mathbb{Z}, \quad x \mapsto \frac{2(x \cdot \alpha)}{\alpha^2} \bmod s^k$$

where $\alpha \in \mathcal{N}_s^\dagger$. Thus, there is a reflection $r_\alpha \in \text{Aut } \mathcal{N}_s$ given by

$$r_\alpha : x \mapsto x - \frac{2(x \cdot \alpha)}{\alpha^2} \alpha.$$

If $\alpha^2 = \frac{1}{2} \bmod \mathbb{Z}$ and $2\alpha = 0$ then $r_\alpha = \text{id}$.

Let p be a prime and consider the homomorphism

$$\text{Aut}(\mathcal{N}) = \prod_p \text{Aut}(\mathcal{N}_p) \xrightarrow{\phi} \prod_p \Sigma_p(N) / \Sigma_p^\sharp(N)$$

which is the product of the epimorphisms

$$\phi_p : \text{Aut}(\mathcal{N}_p) \twoheadrightarrow \Sigma_p(N) / \Sigma_p^\sharp(N)$$

introduced in Miranda–Morrison [17] (see chapter 8, section 7). The images of the homomorphism ϕ_p can be computed on reflections as follows: For a prime s and an element $\alpha \in \mathcal{N}_s^\dagger$, the image of the reflection $r_\alpha \in \text{Aut}(\mathcal{N}_s)$ under ϕ_s is given by $\phi_s(r_\alpha) = (-1, us^k)$, see (2.6). If $s = 2$ and $\alpha^2 = 0 \bmod \mathbb{Z}$, then $\phi_s(r_\alpha)$ is only well-defined modulo Γ_2^{++} . If $s = 2$ and $\alpha^2 = \frac{1}{2} \bmod \mathbb{Z}$, then $\phi_s(r_\alpha)$ is well-defined modulo $\Gamma_{2,2}$. In these cases to determine the value of $\phi_s(r_\alpha)$, we need more information about α and N .

Given another prime p , we define the p -norm $|\alpha|_p \in \{\pm 1\}$ of $\alpha \in \mathcal{N}_s^\dagger$ by

$$|\alpha|_p := \begin{cases} \chi_p(s^k) & \text{if } s \neq p, \\ \chi_p(u) & \text{if } s = p, \end{cases}$$

where the homomorphism $\chi_p : \mathbb{Z}_p^\times / (\mathbb{Z}_p^\times)^2 \rightarrow \{\pm 1\}$ is defined as

$$\chi_p(u) := \begin{cases} \left(\frac{u}{p}\right) & \text{if } p \neq 2, \\ u \bmod 4 & \text{if } p = 2. \end{cases}$$

Note that $|\alpha|_2$ is undefined when $p = 2$ and $\alpha^2 = 0 \bmod \mathbb{Z}$. Following [1], given primes p, s and a vector $\alpha \in \mathcal{N}_s^\dagger$, we introduce the group

$$E_p(N) := \begin{cases} \{\pm 1\} & \text{if } p = 1 \bmod 4 \text{ and } e_p(N) \cdot |\tilde{\Sigma}_p(N)| = 8, \\ 1 & \text{otherwise,} \end{cases}$$

the map $\bar{\phi}_p : \mathcal{N}_s^\dagger \rightarrow E_p(N)$,

$$\bar{\phi}_p(\alpha) := \begin{cases} 1 & \text{if } E_p(N) = 1, \\ |\alpha|_p & \text{otherwise,} \end{cases}$$

and the map $\bar{\beta}_p : \mathcal{N}_s^\dagger \rightarrow \Gamma_0$,

$$\bar{\beta}_p(\alpha) := \begin{cases} (\delta_p(\alpha) \cdot |\alpha|_p, 1) & \text{if } p = 1 \bmod 4, \\ \delta_p(\alpha) \times |\alpha|_p & \text{otherwise,} \end{cases}$$

where the map $\delta_p : \mathcal{N}_s^\dagger \rightarrow \{\pm 1\}$ is defined by

$$\delta_p(\alpha) := (-1)^{\delta_{p,s}}$$

(here $\delta_{p,s}$ is the conventional Kronecker symbol). Note that we have the assignment

$$r_\alpha \mapsto (\delta_p(\alpha), |\alpha|_p) \in \Gamma'_{p,0}.$$

The following lemmas provide an explicit description for the group $E(N)$ and compute the image of the homomorphism e on the reflections r_α for the special case when N has one or two irregular primes.

LEMMA 1 (Akyol–Degtyarev [1]). *Let N be a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$, $\Sigma_2^\sharp(N) \supset \Gamma_{2,2}$, and assume that N has one irregular prime p . Then $E(N) = E_p(N)$ and $e(r_\alpha) = \bar{\phi}_p(\alpha)$ for any $\alpha \in \mathcal{N}^\dagger$.*

LEMMA 2 (Akyol–Degtyarev [1]). *Let N be a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$, $\Sigma_2^\sharp(N) \supset \Gamma_{2,2}$, and assume that N has two irregular primes p, q . Then*

$$E(N) = E_p(N) \times E_q(N) \times (\Gamma_0 / \tilde{\Sigma}_p(N) \cdot \tilde{\Sigma}_q(N)),$$

$$e(r_\alpha) = \bar{\phi}_p(\alpha) \times \bar{\phi}_q(\alpha) \times (\bar{\beta}_p(\alpha) \cdot \bar{\beta}_q(\alpha)),$$

for any $\alpha \in \mathcal{N}^\dagger$ such that $\alpha^2 \neq 0 \bmod \mathbb{Z}$ if $p = 2$ or $q = 2$.

COROLLARY 2 (Akyol–Degtyarev [1]). *Under the hypothesis of Lemma 2, assume, in addition, that $|E(N)| = |E_p(N)| = 2$. Then $E(N) = E_p(N)$ and $e(r_\alpha) = |\alpha|_p$ for any $\alpha \in \mathcal{N}^\dagger$.*

2.7. Positive sign structure. Let N be a non-degenerate lattice. The orthogonal projection of any maximal positive definite subspace in $N \otimes \mathbb{R}$ to any other such subspace is an isomorphism of vector spaces. Thus a choice of an orientation of one maximal positive definite subspace in $N \otimes \mathbb{R}$ defines a coherent orientation of any other. A choice of an orientation of a maximal positive definite subspace of $N \otimes \mathbb{R}$ is called a *positive sign structure*. We denote by $O^+(N)$ the subgroup of $O(N)$ consisting of the isometries preserving a positive sign structure. Obviously either $O^+(N) = O(N)$ or $O^+(N)$ is a subgroup of $O(N)$ of index 2. In the latter case, each element of $O(N) \setminus O^+(N)$ is called a *+disorienting* isometry of N . Following [17], we define the map $\det_+ : O(N) \rightarrow \{\pm 1\}$ as

$$\det_+(a) := \begin{cases} +1 & \text{if } a \text{ preserves the positive sign structure,} \\ -1 & \text{if } a \text{ reserves the positive sign structure.} \end{cases}$$

Note that $\text{Ker}(\det_+) = O^+(N)$.

PROPOSITION 3 (Miranda–Morrison [17]). *Let N be a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$. Then one has $\tilde{\Sigma}(N) \subset \Gamma_0^{--}$ if and only if $\det_+(a) = 1$ for all $a \in \text{Ker}[d : O(N) \rightarrow \text{Aut}(\mathcal{N})]$.*

Hence, if $\tilde{\Sigma} \subset \Gamma_0^{--}$, the map $\det_+ : \text{Im } d \rightarrow \{\pm 1\}$ is the descent of \det_+ which is well-defined by the proposition above. The following lemma computes the images of the function \det_+ on reflections.

LEMMA 3 (Akyol–Degtyarev [1]). *Let N be a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$, $\Sigma_2^\sharp(N) \supset \Gamma_{2,2}$, and assume that there exists a prime p such that $\tilde{\Sigma}_p(N) \subset \Gamma_0^{--}$. Then, for an element $\alpha \in \mathcal{N}^\dagger$ such that $r_\alpha \in \text{Im } d$ and $\alpha^2 \not\equiv 0 \pmod{\mathbb{Z}}$ if $p = 2$, one has $\det_+(r_\alpha) = \delta_p(\alpha) \cdot |\alpha|_p$.*

Defined in [16], we introduce the group

$$E^+(N) := \Gamma_{\mathbb{A},0} / \prod_p \Sigma_p^\sharp(N) \cdot \Gamma_0^{--}. \quad (2.7)$$

(Similar to (2.2) and (2.3) the actual computation reduces to finitely many primes $p \mid \det(N)$.) As in Theorem 8 there is an exact sequence

$$O^+(N) \xrightarrow{d} \text{Aut}(\mathcal{N}) \xrightarrow{e^+} E^+(N) \rightarrow g(N) \rightarrow 1.$$

The size of the group $E^+(N)$ is also computed in [16]: one replaces $[\Gamma_0 : \tilde{\Sigma}(N)]$ in (2.5) with $[\Gamma_0^{--} : \tilde{\Sigma}(N) \cap \Gamma_0^{--}]$. For an irregular prime p , we put $\tilde{\Sigma}_p^+(N) := \tilde{\Sigma}_p(N) \cap \Gamma_0^{--}$.

Given a unimodular even lattice L and a primitive sublattice $S \subset L$ such that $N := S^\perp$ is a non-degenerate indefinite lattice with $\text{rk}(N) \geq 3$, we have a well-defined homomorphism $d_+^\perp : O(S) \rightarrow E^+(N)$, cf. see §2.5 for the definition of d^\perp .

Let p and s be two irregular primes and choose an element $\alpha \in \mathcal{N}_s^\dagger$ as in (2.6), we introduce the group

$$E_p^+(N) := \begin{cases} E_p(N) & \text{if } p \equiv 1 \pmod{4}, \\ \Gamma_0 / \tilde{\Sigma}_p(N) \cdot \Gamma_0^{--} & \text{otherwise,} \end{cases}$$

the map $\bar{\phi}_p^+ : \mathcal{N}_s^\dagger \rightarrow E_p^+(N)$,

$$\bar{\phi}_p^+(\alpha) := \begin{cases} \bar{\phi}_p(\alpha) & \text{if } p \equiv 1 \pmod{4}, \\ \delta_p(\alpha) \cdot |\alpha|_p & \text{if } p \not\equiv 1 \pmod{4} \text{ and } E_p^+(N) \neq 1, \\ 1 & \text{if } p \not\equiv 1 \pmod{4} \text{ and } E_p^+(N) = 1, \end{cases}$$

and the map $\bar{\beta}_p^+ : \mathcal{N}_s^\dagger \rightarrow \Gamma_0^{--}$,

$$\bar{\beta}_p^+(\alpha) := \begin{cases} \delta_p(\alpha) \cdot |\alpha|_p & \text{if } p \equiv 1 \pmod{4}, \\ |\alpha|_p & \text{if } p \not\equiv 1 \pmod{4} \text{ and } E_p^+(N) \neq 1, \\ \text{proj}(\bar{\beta}_p(\alpha)) & \text{if } p \not\equiv 1 \pmod{4} \text{ and } E_p^+(N) = 1, \end{cases}$$

where $\text{proj} : \Gamma_0 \rightarrow \Gamma_0 / \tilde{\Sigma}_p(N) = \Gamma_0^{--}$ is the projection map. Next lemma computes the group $E^+(N)$ and the values of the homomorphism \mathbf{e}^+ on the reflections r_α

LEMMA 4 (Akyol–Degtyarev [1]). *Let N be a non-degenerate indefinite even lattice with $\text{rk}(N) \geq 3$, $\Sigma_2^\sharp(N) \supset \Gamma_{2,2}$ and assume that N has two irregular primes p, q . Then*

$$\begin{aligned} E^+(N) &= E_p^+(N) \times E_q^+(N) \times (\Gamma_0^{--} / \tilde{\Sigma}_p^+(N) \cdot \tilde{\Sigma}_q^+(N)) \\ \mathbf{e}^+(r_\alpha) &= \bar{\phi}_p^+(\alpha) \times \bar{\phi}_q^+(\alpha) \times (\bar{\beta}_p^+(\alpha) \cdot \bar{\beta}_q^+(\alpha)) \end{aligned}$$

for any $\alpha \in \mathcal{N}^\dagger$ such that $\alpha^2 \not\equiv 0 \pmod{\mathbb{Z}}$ if $p = 2$ or $q = 2$.

2.8. Root Systems. A root in a lattice L is an element $v \in L$ of square -2 . A root system is a negative definite lattice generated by its roots. Each root system splits uniquely into orthogonal sum of its irreducible components. As explained in [2], the irreducible root systems are \mathbf{A}_n , $n \geq 1$, \mathbf{D}_m , $m \geq 4$ and \mathbf{E}_6 , \mathbf{E}_7 , \mathbf{E}_8 . The corresponding discriminant forms are as follows:

$$\begin{aligned}
\text{disc } \mathbf{A}_n &= \left\langle -\frac{n}{n+1} \right\rangle, & \text{disc } \mathbf{D}_{2k+1} &= \left\langle -\frac{2k+1}{4} \right\rangle, \\
\text{disc } \mathbf{D}_{8k \pm 2} &= 2 \left\langle \mp \frac{1}{2} \right\rangle, & \text{disc } \mathbf{D}_{8k} &= \mathcal{U}(2), & \text{disc } \mathbf{D}_{8k+4} &= \mathcal{V}(2), \\
\text{disc } \mathbf{E}_6 &= \left\langle \frac{2}{3} \right\rangle, & \text{disc } \mathbf{E}_7 &= \left\langle \frac{1}{2} \right\rangle, & \text{disc } \mathbf{E}_8 &= 0.
\end{aligned}$$

Given a root system S , the group generated by reflections (defined by the roots of S) acts simply transitively on the set of Weyl chambers of S . The roots constituting a single Weyl chamber form a *standard basis* for S ; these roots are naturally identified with the vertices of the Dynkin graph $\Gamma := \Gamma_S$. Thus, one has an obvious homomorphism

$$\text{Sym}(\Gamma) \rightarrow O(S) \rightarrow \text{Aut}(\mathcal{S})$$

where $\text{Sym}(\Gamma)$ denotes the symmetries of Γ . Irreducible root systems correspond to connected Dynkin graphs. The following statement follows immediately from the classification of connected Dynkin graphs (see N. Bourbaki [2]).

LEMMA 5. *Let $\Gamma = \Gamma_S$ be the connected Dynkin graph of an irreducible root system S . Then one of the following holds:*

- (1) *If S is \mathbf{A}_1 , \mathbf{E}_7 or \mathbf{E}_8 , then $\text{Sym}(\Gamma) = 1$.*
- (2) *If S is \mathbf{D}_4 , then $\text{Sym}(\Gamma) = \mathbb{S}_3$.*
- (3) *For all other types, $\text{Sym}(\Gamma) = \mathbb{Z}/2\mathbb{Z}$.*

If S is \mathbf{A}_p , $p \geq 2$, \mathbf{D}_{2k+1} or \mathbf{E}_8 , then the only nontrivial symmetry of Γ induces $-\text{id}$ on \mathcal{S} . If S is \mathbf{E}_8 then $\mathcal{S} = 0$ and if S is \mathbf{A}_1 , \mathbf{A}_7 or \mathbf{D}_{2k} , the groups \mathcal{S} are \mathbb{F}_2 modules and $-\text{id} = \text{id}$ on $\text{Aut } \mathcal{S}$.

Further details on irreducible root systems can be found in N. Bourbaki [2].

3. Simple quartics

3.1. Quartics and K3-surfaces. A *quartic* is a surface $X \subset \mathbb{P}^3$ of degree four. A quartic is *simple* if all its singular points are simple, i.e., those of type **A**, **D**, **E**. Isomorphism classes of simple singularities are known to be in a one-to-one correspondence with those of irreducible root systems (see Dufree [8] for details). Hence, a set of simple singularities can be identified with a root system, the irreducible summands of the latter (see §2.8) correspond to the individual singularity points.

Let $X \subset \mathbb{P}^3$ be a simple quartic and consider its minimal resolution of singularities \tilde{X} . It is well known that \tilde{X} is a $K3$ -surface; hence, $H_2(\tilde{X}) \cong 2\mathbf{E}_8 \oplus 3\mathbf{U}$, where \mathbf{U} is the hyperbolic plane defined as $\mathbf{U} := \mathbb{Z}u_1 \oplus \mathbb{Z}u_2$, $u_1^2 = u_2^2 = 0$ and $u_1 \cdot u_2 = 1$. Note that $2\mathbf{E}_8 \oplus 3\mathbf{U}$ is the only even unimodular lattice of signature $(\sigma_+, \sigma_-) = (3, 19)$. We fix the notation $\mathbf{L}_X := H_2(\tilde{X})$ and $\mathbf{L} := 2\mathbf{E}_8 \oplus 3\mathbf{U}$.

For each simple singular point p of X the components of the exceptional divisor are smooth rational (-2) -curves spanning a root lattice in \mathbf{L}_X . These sublattices are obviously orthogonal and their orthogonal sum, which is identified with the set of singularities of X , is denoted by \mathbf{S}_X . The rank $\text{rk}(\mathbf{S}_X)$ equals the total Milnor number $\mu(X)$. Since $\sigma_-(\mathbf{L}) = 19$ and $\mathbf{S}_X \subset \mathbf{L}$ is negative definite, one has $\mu(X) \leq 19$ (see [25], cf., [20]). If $\mu(X) = 19$, the quartic is called *maximizing*. We introduce the following objects:

- $\mathbf{S}_X \subset \mathbf{L}_X$: the sublattice generated by the set of classes of exceptional divisors contracted by the blow-up map $\tilde{X} \rightarrow X$;
- $h_X \in \mathbf{L}_X$: the class of the pull-back of a generic plane section of X ;
- $\mathbf{S}_{X,h} = \mathbf{S}_X \oplus \mathbb{Z}h_X \subset \mathbf{L}_X$;
- $\tilde{\mathbf{S}}_X \subset \tilde{\mathbf{S}}_{X,h} \subset \mathbf{L}_X$: the primitive hulls of \mathbf{S}_X and $\mathbf{S}_{X,h}$, respectively, i.e., $\tilde{\mathbf{S}}_X := (\mathbf{S}_X \otimes \mathbb{Q}) \cap \mathbf{L}_X$ and $\tilde{\mathbf{S}}_{X,h} := (\mathbf{S}_{X,h} \otimes \mathbb{Q}) \cap \mathbf{L}_X$;
- $\omega_X \subset \mathbf{L}_X \otimes \mathbb{R}$: the oriented 2-subspace spanned by the real and imaginary parts of the class of a holomorphic 2-form on \tilde{X} (the *period* of \tilde{X}).

The triple $(\mathbf{S}_X, h_X, \mathbf{L}_X)$ is called the *homological type* of X .

3.2. Abstract homological types. As explained above the set of singularities of a quartic $X \subset \mathbb{P}^3$ can be viewed as a root lattice $\mathbf{S} \subset \mathbf{L}$.

DEFINITION 1. A *configuration* (extending a given set of singularities \mathbf{S}) is a finite index extension $\tilde{\mathbf{S}}_h \supset \mathbf{S}_h := \mathbf{S} \oplus \mathbb{Z}h$, $h^2 = 4$, satisfying the following conditions:

- (1) Each root $r \in (\mathbf{S} \otimes \mathbb{Q}) \cap \tilde{\mathbf{S}}_h$ with $r^2 = -2$ is in \mathbf{S} .
- (2) $\tilde{\mathbf{S}}_h$ does not have an element v with $v^2 = 0$ and $v \cdot h = 2$.

An *automorphism* of a configuration $\tilde{\mathbf{S}}_h$ is an auto-isometry of $\tilde{\mathbf{S}}_h$ preserving h . The group of automorphisms of $\tilde{\mathbf{S}}_h$ is denoted by $\text{Aut}_h(\tilde{\mathbf{S}}_h)$. One has the obvious inclusions $\text{Aut}_h(\tilde{\mathbf{S}}_h) \subset O(\tilde{\mathbf{S}}) \subset O(\mathbf{S})$, the latter is due to the condition (1) in Definition 1, since \mathbf{S} is recovered as the sublattice in $h^\perp \subset \tilde{\mathbf{S}}_h$ generated by roots.

DEFINITION 2. An *abstract homological type* extending a fixed set of singularities \mathbf{S} is an extension of $\mathbf{S}_h := \mathbf{S} \oplus \mathbb{Z}h$, $h^2 = 4$, to a lattice L isomorphic to $2\mathbf{E}_8 \oplus 3\mathbf{U}$, such that the primitive hull $\tilde{\mathbf{S}}_h$ of \mathbf{S}_h in L is a configuration.

An abstract homological type is uniquely determined by the triple $\mathcal{H} = (\mathbf{S}, h, \mathbf{L})$. An *isomorphism* between two abstract homological types $\mathcal{H}_i = (\mathbf{S}_i, h_i, \mathbf{L}_i)$, $i = 1, 2$, is an isometry $\mathbf{L}_1 \rightarrow \mathbf{L}_2$, taking h_1 and \mathbf{S}_1 to h_2 and \mathbf{S}_2 , respectively (as a set).

Given an abstract homological type $\mathcal{H} = (\mathbf{S}, h, \mathbf{L})$, we let $\tilde{\mathbf{S}} := (\mathbf{S} \otimes \mathbb{Q}) \cap \mathbf{L}$ and $\tilde{\mathbf{S}}_h := (\mathbf{S}_h \otimes \mathbb{Q}) \cap \mathbf{L}$ be the primitive hulls of \mathbf{S} and \mathbf{S}_h , respectively. Note that $\tilde{\mathbf{S}} = h_{\tilde{\mathbf{S}}_h}^\perp$, i.e., $\tilde{\mathbf{S}}$ is also the primitive hull of h^\perp . The orthogonal complement \mathbf{S}_h^\perp is a non-degenerate lattice with $\sigma_+ \mathbf{S}_h^\perp = 2$. It follows that all positive definite 2-subspaces in $\mathbf{S}_h^\perp \otimes \mathbb{R}$ can be oriented in a coherent way (see §2.7).

DEFINITION 3. An *orientation* of an abstract homological type $\mathcal{H} = (\mathbf{S}, h, \mathbf{L})$ is a choice θ of one of the coherent orientations of positive definite 2-subspaces of $\mathbf{S}_h^\perp \otimes \mathbb{R}$.

An *isomorphism* between two oriented abstract homological types $(\mathcal{H}_i, \theta_i)$, $i = 1, 2$, is an isomorphism $\mathcal{H}_1 \rightarrow \mathcal{H}_2$, taking θ_1 to θ_2 . An abstract homological type \mathcal{H} is called *symmetric* if (\mathcal{H}, θ) is isomorphic to $(\mathcal{H}, -\theta)$ for some orientation θ of \mathcal{H} , i.e., \mathcal{H} admits an automorphism reversing the orientation.

3.3. Classification of singular quartics. Due to Saint-Donat [22] and Urabe [25], a triple $\mathcal{H} = (\mathbf{S}, h, \mathbf{L})$ is isomorphic to the homological type $(\mathbf{S}_X, h_X, \mathbf{L}_X)$ of a simple quartic $X \subset \mathbb{P}^3$ if and only if \mathcal{H} is an abstract homological type in the sense of Definition 2. In this case, the oriented 2-subspace ω_X introduced in §3.1 defines an orientation of \mathcal{H} .

THEOREM 10 (see Theorem 2.3.1 in [7]). *The map sending a simple quartic surface $X \subset \mathbb{P}^3$ to its oriented homological type establishes a one to one correspondence between the set of equisingular deformation classes of quartics with a given set of simple singularities \mathbf{S} and the set of isomorphism classes of oriented abstract homological types extending \mathbf{S} . Complex conjugate quartics have isomorphic homological types that differ by the orientations.*

DEFINITION 4. A quartic X is called *non-special* if its homological type is primitive, i.e., $\mathbf{S}_h \subset \mathbf{L}$ is a primitive sublattice.

Note that the homological type $\mathcal{H} = (\mathbf{S}, h, \mathbf{L})$ is primitive if and only if $\tilde{\mathbf{S}}_h = \mathbf{S}_h$, in this case, one has $\text{disc } \tilde{\mathbf{S}}_h = \mathcal{S} \oplus \langle \frac{1}{4} \rangle$ and $\text{Aut}_h(\tilde{\mathbf{S}}_h) = O(\mathbf{S})$.

For a given set of simple singularities \mathbf{S} , the corresponding equisingular stratum of quartics is denoted by $\mathcal{M}(\mathbf{S})$. Our primary interest is the family $\mathcal{M}_1(\mathbf{S}) \subset \mathcal{M}(\mathbf{S})$ constituted by the non-special quartics with the set of singularities \mathbf{S} . More generally, since the kernel \mathcal{K} of the finite index extension $\mathbf{S}_h \subset \tilde{\mathbf{S}}_h$ is obviously invariant under equisingular deformations, one can

consider the strata $\mathcal{M}_*(\mathbf{S}) \subset \mathcal{M}(\mathbf{S})$ where the subscript $*$ is the sequence of invariant factors of the kernel \mathcal{K} .

4. Proofs

4.1. Proof of Theorem 1. Note that $X \setminus (\text{Sing } X \cup H) \cong \tilde{X} \setminus (E \cup H)$, where \tilde{X} is the minimal resolution of X and E is the exceptional divisor of the blow up $\tilde{X} \rightarrow X$. Recall that \mathbf{S}_X is the sublattice in $\mathbf{L}_X = H_2(\tilde{X})$ generated by the components of E (see §3.1). Thus, one has $H_2(E \cup H) = \mathbf{S}_X \oplus \mathbb{Z}h_X \subset \mathbf{L}_X = H_2(\tilde{X})$.

We have the following cohomology exact sequence of pair $(\tilde{X}, E \cup H)$:

$$\cdots \xrightarrow{j^*} H^2(\tilde{X}) \xrightarrow{i^*} H^2(E \cup H) \xrightarrow{\delta} H^3(\tilde{X}, E \cup H) \xrightarrow{j^*} \underbrace{H^3(\tilde{X})}_0 \rightarrow \cdots.$$

Hence, $H^3(\tilde{X}, E \cup H) = \text{coker } i^*$. By universal coefficients, since all groups involved are free, i^* is the adjoint of the map

$$i_* : H_2(E \cup H) \rightarrow H_2(\tilde{X}),$$

which is the inclusion $\mathbf{S}_{X,h} \hookrightarrow \mathbf{L}_X$. Thus, we have an exact sequence

$$0 \rightarrow H_2(E \cup H) \xrightarrow{i_*} H_2(\tilde{X}) \rightarrow \tilde{\mathbf{S}}_{X,h}/\mathbf{S}_{X,h} \oplus \mathbf{F} \rightarrow 0,$$

where \mathbf{F} is a finitely generated free abelian group. This sequence can be regarded as a free resolution of $\tilde{\mathbf{S}}_{X,h}/\mathbf{S}_{X,h} \oplus \mathbf{F}$ and, by the definition of derived functor, we have the following isomorphisms

$$\text{coker } i^* = \text{Ext}(\tilde{\mathbf{S}}_{X,h}/\mathbf{S}_{X,h} \oplus \mathbf{F}, \mathbb{Z}) = \text{Ext}(\tilde{\mathbf{S}}_{X,h}/\mathbf{S}_{X,h}, \mathbb{Z}).$$

Combining these observations with Poincaré–Lefschetz duality $H_1(\tilde{X} \setminus (E \cup H)) = H^3(\tilde{X}, E \cup H)$, we conclude that

$$H_1(\tilde{X} \setminus (E \cup H)) = \text{Ext}(\tilde{\mathbf{S}}_{X,h}/\mathbf{S}_{X,h}, \mathbb{Z}) \cong \tilde{\mathbf{S}}_{X,h}/\mathbf{S}_{X,h}$$

(the last isomorphism being not natural). In particular $H_1(\tilde{X} \setminus (E \cup H)) = 0$ if and only if $\mathbf{S}_{X,h} = \tilde{\mathbf{S}}_{X,h}$ i.e., if and only if X is non-special. \square

4.2. Proof of Theorem 2. For the reader's convenience, we divide the proof into three propositions; Theorem 2 is their immediate consequence.

PROPOSITION 4. *The realizable sets are all sets of singularities that can be obtained by a perturbation from either the 59 maximizing sets of singularities listed in Table 1 or 19 sets of singularities with the Milnor number 18 listed in Table 2.*

PROOF. According to Theorems 10 and Definition 4, a set of singularities \mathbf{S} is realized by a non-special quartic if and only if \mathbf{S} extends to a primitive homological type. Thus, we are interested in primitive extensions $\mathbf{S}_h \hookrightarrow \mathbf{L} = 3\mathbf{U} \oplus 3\mathbf{E}_8$. Since the homological type is primitive, one has $\text{disc } \tilde{\mathbf{S}}_h = \mathcal{S} \oplus \langle \frac{1}{4} \rangle$, and the realizable sets are easily found by using Nikulin's Existence Theorem (Theorem 5) applied to the genus of the transcendental lattice $\mathbf{T} := \mathbf{S}_h^\perp$, which is determined by \mathbf{S} , see §2.4. Implementing the algorithm in GAP [10], we found that 2872 sets of simple singularities are realized by non-maximal non-special quartics and 59 sets of simple singularities are realized by maximal non-special quartics. According to E. Looijenga [14], deformation classes of perturbations of an individual simple singularity of type \mathbf{S} are in a one-to-one correspondence with the isomorphism classes of primitive extensions $\mathbf{S}' \hookrightarrow \mathbf{S}$ of root lattices, see §2.8 and §2.4. As shown in [9], the latter is the case if and only if the Dynkin graph of \mathbf{S}' is an induced subgraph of that of \mathbf{S} . Hence, given a simple quartic X , any perturbation X to a simple quartic X' gives rise to a perturbation of the set of singularities \mathbf{S} of X to the set of singularities \mathbf{S}' of X' . Conversely, any induced subgraph of the Dynkin graph of a simple quartic X is that of an appropriate small perturbation X' of X . Proof of this statement repeats, almost literally, the proof of a similar theorem for plane sextic curves (see Proposition 5.1.1 in [6]). Accordingly, the list of 2872 sets of simple singularities realized by non-maximal non-special quartics is compared against the list of all perturbations of the 59 maximizing sets of singularities given in Table 1 and 19 sets of singularities with Milnor number 18 given in Table 2. The two lists coincide. \square

Let \mathbf{S} be one of the realizable sets of singularities and \mathbf{T} a representative of the genus $g(\mathbf{S}_h^\perp)$. By Theorem 10, the connected components of the space $\mathcal{M}_1(\mathbf{S})$ modulo the complex conjugation $\text{conj} : \mathbb{P}^3 \rightarrow \mathbb{P}^3$ are enumerated by the isomorphism classes of primitive homological types extending \mathbf{S} . We investigate these isomorphism classes separately for the maximizing case, *i.e.*, $\mu(\mathbf{S}) = 19$, and non-maximizing case, *i.e.*, $\mu(\mathbf{S}) \leq 18$.

If $\mu(\mathbf{S}) = 19$, the transcendental lattice \mathbf{T} is a positive definite sublattice of rank 2, and the numbers (r, c) of connected components of the space $\mathcal{M}_1(\mathbf{S})$ listed in Table 1 can easily be computed by Gauss theory of binary quadratic forms [11] (A. Degtyarev, private communication); details will appear elsewhere. Thus, throughout the rest of the proof we assume $\mu(\mathbf{S}) \leq 18$.

PROPOSITION 5. *For each realizable set of singularities \mathbf{S} with $\mu(\mathbf{S}) \leq 18$, the space $\mathcal{M}_1(\mathbf{S})/\text{conj}$ is connected.*

PROOF. If $\mu(\mathbf{S}) \leq 18$, then \mathbf{T} is an indefinite lattice with $\text{rk } \mathbf{T} \geq 3$ and we can apply Miranda–Morison's theory. We try to enumerate primitive homo-

logical types $\mathcal{H} = (\mathbf{S}, h, \mathbf{L})$ extending \mathbf{S} , *i.e.*, the primitive extensions $\mathbf{S}_h \hookrightarrow L$. Since the extension is primitive, $\tilde{\mathbf{S}}_h = \mathbf{S}_h$, one has $\text{disc } \tilde{\mathbf{S}}_h = \mathcal{S} \oplus \langle \frac{1}{4} \rangle$ and $\text{Aut}_h(\mathbf{S}_h) \cong O(\mathbf{S})$. Then we have a well-defined homomorphism $d^\perp : O(\mathbf{S}) \rightarrow E(\mathbf{T})$, and by Corollary 1,

$$\pi_0(\mathcal{M}_1(\mathbf{S})/\text{conj}) \cong \text{Coker}(d^\perp : O(\mathbf{S}) \rightarrow E(\mathbf{T})). \quad (4.1)$$

Thus, the space $\mathcal{M}_1(\mathbf{S})/\text{conj}$ is connected (equivalently, the primitive homological type extending \mathbf{S} is unique up to isomorphism) if and only if the map d^\perp is surjective, and we shall prove the latter statement is true.

Out of the 2872 sets of singularities realized by non-special non-maximizing quartics, for 2830 sets of singularities one gets $E(\mathbf{T}) = 1$ by using (2.5), and the assertion follows automatically.

For the remaining 42 cases, one has $|E(\mathbf{T})| \neq 1$. Among these, there are 18 set of singularities containing a point of type \mathbf{A}_4 and satisfying the hypothesis of Lemma 1 or Corollary 2 with $p = 5$. For these set of singularities one has $|E(\mathbf{T})| = 2$ and a nontrivial symmetry of any type \mathbf{A}_4 point maps to the generator $-1 \in E(\mathbf{T})$.

There are 8 sets of singularities containing a point of type \mathbf{A}_2 and satisfying the hypothesis of Lemma 2 with $p = 2$, $q = 3$. For these 8 cases, one has $|E(\mathbf{T})| = 2$ and a nontrivial symmetry of any type \mathbf{A}_2 point maps to the generator $-1 \in E(\mathbf{T})$. For the following 4 sets of singularities,

$$\mathbf{D}_9 \oplus \mathbf{A}_3 \oplus 3\mathbf{A}_2, \quad \mathbf{D}_7 \oplus \mathbf{D}_5 \oplus 3\mathbf{A}_2,$$

$$\mathbf{A}_{11} \oplus \mathbf{A}_3 \oplus 2\mathbf{A}_2, \quad \mathbf{A}_8 \oplus 2\mathbf{A}_3 \oplus 2\mathbf{A}_2,$$

one has $|E(\mathbf{T})| = 4$. Each of these 4 sets has two irregular primes $p = 2$, $q = 3$, and for all of them the homomorphism given by Lemma 2 is

$$\mathbf{e}(r_\alpha) = (\delta_2(\alpha) \cdot \delta_3(\alpha), |\alpha|_2 \cdot |\alpha|_3) \in \{\pm 1\} \times \{\pm 1\}.$$

A symmetry of any type \mathbf{A}_2 point and a transposition $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$ give rise to reflections $r_\alpha, r_\sigma \in \mathcal{T}$ with $\alpha^2 = \frac{2}{3}$ and $(\sigma)^2 = \frac{4}{3}$. The images $\mathbf{e}(r_\alpha) = (-1, -1)$ and $\mathbf{e}(r_\sigma) = (-1, 1)$ are linearly independent, thus generating the group $E(\mathbf{T})$.

The 9 sets of singularities listed in Table 3 still satisfy the assumptions of Lemma 2, which yields $|E(\mathbf{T})| = 2$. Also shown in the table are the irregular primes (p, q) , the homomorphism $\mathbf{e} : \text{Aut}(\mathcal{T}) \rightarrow E(\mathbf{T})$, and an automorphism of \mathbf{S} generating $E(\mathbf{T})$.

Finally, what remains are the three sets of singularities

$$\mathbf{D}_4 \oplus 2\mathbf{A}_4 \oplus 3\mathbf{A}_2, \quad 2\mathbf{A}_7 \oplus 2\mathbf{A}_2 \quad 2\mathbf{A}_4 \oplus 2\mathbf{A}_3 \oplus 2\mathbf{A}_2,$$

to which Lemmas 1, 2 or Corollary 2 can not be applied. For them, we compute the group $E(\mathbf{T})$ directly from the definition (2.2) which can be

Table 3. Extremal singularities

Singularities	(p, q)	$e : \text{Aut}(\mathcal{T}) \rightarrow E(\mathbf{T})$	generators of $E(\mathbf{T})$
$\mathbf{E}_8 \oplus 2\mathbf{A}_3 \oplus 2\mathbf{A}_2$	(2, 3)	$e(r_x) = \delta_2(\alpha) \cdot \delta_3(\alpha) \cdot \alpha _2 \cdot \alpha _3$	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$2\mathbf{E}_6 \oplus 2\mathbf{A}_3$	(2, 3)	$e(r_x) = \delta_2(\alpha) \cdot \delta_3(\alpha) \cdot \alpha _2 \cdot \alpha _3$	symmetry of \mathbf{A}_3
$\mathbf{D}_{11} \oplus \mathbf{A}_3 \oplus 2\mathbf{A}_2$	(2, 3)	$e(r_x) = \delta_2(\alpha) \cdot \delta_3(\alpha) \cdot \alpha _2 \cdot \alpha _3$	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$2\mathbf{D}_7 \oplus 2\mathbf{A}_2$	(2, 3)	$e(r_x) = \delta_2(\alpha) \cdot \delta_3(\alpha) \cdot \alpha _2 \cdot \alpha _3$	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$2\mathbf{D}_5 \oplus 2\mathbf{A}_4$	(2, 5)	$e(r_x) = \delta_2(\alpha) \cdot \delta_5(\alpha)$	$\mathbf{A}_4 \leftrightarrow \mathbf{A}_4$
$\mathbf{D}_5 \oplus \mathbf{A}_6 \oplus \mathbf{A}_3 \oplus 2\mathbf{A}_2$	(2, 3)	$e(r_x) = \delta_2(\alpha) \cdot \delta_3(\alpha) \cdot \alpha _2 \cdot \alpha _3$	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{A}_7 \oplus \mathbf{A}_4 \oplus \mathbf{A}_3 \oplus 2\mathbf{A}_2$	(2, 3)	$e(r_x) = \delta_2(\alpha) \cdot \delta_3(\alpha) \cdot \alpha _2 \cdot \alpha _3$	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$2\mathbf{A}_6 \oplus 2\mathbf{A}_3$	(2, 7)	$e(r_x) = \alpha _2 \cdot \alpha _7$	symmetry of \mathbf{A}_3
$2\mathbf{A}_6 \oplus 3\mathbf{A}_2$	(3, 7)	$e(r_x) = \delta_3(\alpha) \cdot \delta_7(\alpha) \cdot \alpha _3 \cdot \alpha _7$	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$

restated as

$$E(T) = \prod_{p \mid \det(T)} \Gamma_{p,0} \Big/ \prod_{p \mid \det(T)} \Sigma_p^\sharp(T) \cdot \varphi(\Gamma_0),$$

where we identify the inclusion $\Gamma_0 \hookrightarrow \Gamma_{\mathbb{A},0}$ with the product $\varphi := \prod_p \varphi_p$ (see 2.1). For example, for the case

$$2\mathbf{A}_4 \oplus 2\mathbf{A}_3 \oplus 2\mathbf{A}_2,$$

the computation can be summarized in the following table:

	$\Gamma_{5,0}$		$\Gamma_{3,0}$		$\Gamma'_{2,0}$	
generator of $\Sigma_5^\sharp(T)$	-1	-1	1	1	1	1
generator of $\Sigma_3^\sharp(T)$	1	1	-1	1	1	1
generator of $\Sigma_2^\sharp(T)$	1	1	1	1	1	1
$\varphi(-1, 1)$	-1	1	-1	1	-1	1
$\varphi(1, -1)$	1	1	1	-1	1	-1
	δ_5	$ \cdot _5$	δ_3	$ \cdot _3$	δ_2	$ \cdot _2$
a symmetry of \mathbf{A}_4	-1	-1	1	-1	1	1
a transposition $\mathbf{A}_4 \leftrightarrow \mathbf{A}_4$	-1	1	1	-1	1	1
a symmetry of \mathbf{A}_2	1	-1	-1	1	1	-1
a transposition $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$	1	-1	-1	-1	1	-1

The rank of the matrix composed by the 9 rows of the table (see, Remark 1) is $6 = \dim \Gamma'_{2,0} + \dim \Gamma_{3,0} + \dim \Gamma_{5,0}$, which implies that d^\perp is surjective.

For the remaining two cases

$$\mathbf{D}_4 \oplus 2\mathbf{A}_4 \oplus 3\mathbf{A}_2, \quad 2\mathbf{A}_7 \oplus 2\mathbf{A}_2,$$

the computation is almost literally the same. For $2\mathbf{A}_7 \oplus 2\mathbf{A}_2$, where $\Sigma_2^{\sharp}(\mathbf{T}) \not\subset \Gamma_{2,2}$, we have to modify $|\cdot|_2$ by replacing χ_2 with $\chi_2(u) = u \bmod 8 \in \{1, 3, 5, 7\} = \mathbb{Z}_2^{\times}/(\mathbb{Z}_2^{\times})^2$ and consider the full group $\Gamma_{2,0}$ instead of $\Gamma'_{2,0}$. \square

REMARK 1. *Here and below, when speaking about ranks and dimensions, we regard all groups Γ_* , E , E^+ , etc. as \mathbb{F}_2 -vector spaces. In particular, when computing the rank of a matrix, we need to switch from the multiplicative notation $\{1, -1\}$ to the additive $\{0, 1\}$.*

COROLLARY 3. *For all sets of singularities \mathbf{S} with $\mu(\mathbf{S}) \leq 18$, the corresponding transcendental lattice \mathbf{T} is unique in its genus, i.e., $g(\mathbf{T}) = 1$.*

PROPOSITION 6. *If \mathbf{S} is one of*

$$\mathbf{D}_6 \oplus 2\mathbf{A}_6, \quad \mathbf{D}_5 \oplus 2\mathbf{A}_6 \oplus \mathbf{A}_1, \quad 2\mathbf{A}_7 \oplus 2\mathbf{A}_2, \quad 3\mathbf{A}_6, \quad 2\mathbf{A}_6 \oplus 2\mathbf{A}_3$$

then $\mathcal{M}_1(\mathbf{S})$ consists of two complex conjugate components; in all other cases with $\mu(\mathbf{S}) \leq 18$, the stratum $\mathcal{M}_1(\mathbf{S})$ is connected.

PROOF. By Proposition 5, $\mathcal{M}_1(\mathbf{S})$ is connected if and only if the (unique) homological type extending \mathbf{S} is symmetric; otherwise $\mathcal{M}_1(\mathbf{S})$ consists of two complex conjugate components. By Theorem 7, the homological type is symmetric if and only if there is an isometry $a \in O(\mathbf{T})$ with $\det_+(a) = -1$ satisfying $d(a) \in d^{\psi}(O(\mathbf{S}))$, where d^{ψ} is the map induced by any anti-isometry $\psi: \mathcal{S} \oplus \langle \frac{1}{4} \rangle \rightarrow \mathcal{T}$. We consider separately the cases $|E(\mathbf{T})| = |E^+(\mathbf{T})|$ and $|E(\mathbf{T})| < |E^+(\mathbf{T})|$.

LEMMA 6. *If $|E(\mathbf{T})| = |E^+(\mathbf{T})|$, then $\mathcal{M}_1(\mathbf{S})$ is connected.*

PROOF. By definition, we have an exact sequence

$$0 \rightarrow \Gamma_0/\Gamma_0^{--} \cdot \tilde{\Sigma}(\mathbf{T}) \rightarrow E^+(\mathbf{T}) \rightarrow E(\mathbf{T}) \rightarrow 0. \quad (4.2)$$

Hence, $|E(\mathbf{T})| = |E^+(\mathbf{T})|$ if and only if $\tilde{\Sigma}(\mathbf{T}) \not\subset \Gamma_0^{--}$. Then, by Proposition 3, there exists a $+$ -disorienting isometry of \mathbf{T} inducing the identity on disc \mathbf{T} , and Theorem 7 applies. \square

LEMMA 7. *If $|E(\mathbf{T})| < |E^+(\mathbf{T})|$, then $\mathcal{M}_1(\mathbf{S})$ is connected if and only if $d_+^{\perp}: O(\mathbf{S}) \rightarrow E^+(\mathbf{T})$ is an epimorphism.*

PROOF. The non-trivial element of the kernel $K := \Gamma_0/\Gamma_0^{--} \cdot \tilde{\Sigma}(\mathbf{T}) \cong \{\pm 1\}$ in (4.2) is the image under the composed map $O(\mathbf{T}) \rightarrow \text{Aut}(\mathcal{T}) \rightarrow E^+(\mathbf{T})$ of any element $a \in O(\mathbf{T})$ with $\det_+(a) = -1$. Thus, $\mathcal{M}_1(\mathbf{S})$ is connected if and

only if $\text{Im}(d_+^\perp) \cap K \neq 0$, *i.e.*, $\text{rank } d_+^\perp > \text{rank } d^\perp$ (see Remark 1). On the other hand, Proposition 5 can be recast in the form $\text{rank } d^\perp = \dim E(\mathbf{T})$. Since $\dim E_+(\mathbf{T}) = \dim E(\mathbf{T}) + 1$, the statement follows. \square

Lemma 6 implies the connectedness of $\mathcal{M}_1(\mathbf{S})$ for 2721 sets of singularities. For the remaining 151 sets of singularities, one has $|E(\mathbf{T})| < |E^+(\mathbf{T})|$. For 118 of them, one has $|E(\mathbf{T})| = 1$ and $\tilde{\Sigma}_p(\mathbf{T}) \subset \Gamma_0^{--}$ for some prime p . Since $|E(\mathbf{T})| = 1$, the map $d : O(\mathbf{T}) \rightarrow \text{Aut}(\mathcal{T})$ is surjective and the isomorphism

$$\text{Aut}(\mathcal{T})/O^+(\mathbf{T}) = \Gamma_0/\Gamma_0^{--} \cdot \tilde{\Sigma}(\mathbf{T}) = E^+(\mathbf{T}) = \{\pm 1\}$$

(see (4.2)) is the descent of \det_+ , which is well-defined due Proposition 3. In most cases we can use Lemma 3 to show that there exists an element $a \in O(\mathbf{S})$ such that $\det_+(d^\psi(a)) = -1$. Namely,

- if $p = 2$, take for a a nontrivial symmetry of \mathbf{A}_2 , \mathbf{D}_5 , \mathbf{E}_6 or \mathbf{D}_9 ,
- if $p = 3$, take for a a nontrivial symmetry of \mathbf{A}_2 , \mathbf{D}_5 or \mathbf{A}_8 ,
- if $p = 7$, take for a a nontrivial symmetry of \mathbf{A}_2 .

The three sets of singularities

$$\mathbf{D}_6 \oplus 2\mathbf{A}_6, \quad \mathbf{D}_5 \oplus 2\mathbf{A}_6 \oplus \mathbf{A}_1, \quad 3\mathbf{A}_6$$

with $p = 7$ are exceptional (and are listed as such in the statement), as Lemma 3 implies $\det_+ \circ d^\psi \equiv 1$. Indeed, the image of d^ψ is generated by the reflections r_α with either

- $\alpha \in \mathcal{T}_7$, $\alpha^2 = \frac{6}{7}$ (a nontrivial symmetry of \mathbf{A}_6),
- $\alpha \in \mathcal{T}_7$, $\alpha^2 = \frac{12}{7}$ (a transposition $\mathbf{A}_6 \leftrightarrow \mathbf{A}_6$) or
- $\alpha \in \mathcal{T}_2$ (a nontrivial symmetry of \mathbf{D}_6 or \mathbf{D}_5).

One has $\delta_7(\alpha) = |\alpha|_7 = -1$ in the first two cases and $\delta_7(\alpha) = |\alpha|_7 = 1$ in the last one.

There are 30 other sets of singularities still satisfying the condition $\Gamma_{2,2} \subset \Sigma_2^\sharp(\mathbf{T})$ and having two irregular primes, so that we can apply Lemma 4. Among them, 13 sets of singularities contain two type \mathbf{A}_2 points and have $(p, q) = (2, 3)$ and $|E^+(\mathbf{T})| = 4$. A nontrivial symmetry of any type \mathbf{A}_2 point and a transposition $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$ map to two linearly independent elements generating $E^+(\mathbf{T})$. The remaining 17 sets of singularities are listed in Table 4, where we indicate the irregular primes (p, q) , order of $E^+ := E^+(\mathbf{T})$ and a collection of isometries of \mathbf{S} whose images generate $E^+(\mathbf{T})$. The set of singularities $\mathbf{S} = 2\mathbf{A}_6 \oplus 2\mathbf{A}_3$ marked as exceptional is one of the special cases listed in the statement. We have $|E^+(\mathbf{T})| = 4$ and the group $O(\mathbf{S})$ is generated by

- a nontrivial symmetry of \mathbf{A}_3 , mapped to $(1, 1) \in E^+(\mathbf{T})$,
- the transposition $\mathbf{A}_3 \leftrightarrow \mathbf{A}_3$, mapped to $(1, 1) \in E^+(\mathbf{T})$,

- a nontrivial symmetry of \mathbf{A}_6 , mapped to $(-1, 1) \in E^+(\mathbf{T})$ or
- the transposition $\mathbf{A}_6 \leftrightarrow \mathbf{A}_6$, mapped to $(-1, 1) \in E^+(\mathbf{T})$.

It follows that d_+^\perp is not surjective.

Finally, what remains are the 3 sets of singularities

$$\mathbf{D}_4 \oplus 2\mathbf{A}_4 \oplus 3\mathbf{A}_2, \quad 2\mathbf{A}_7 \oplus 2\mathbf{A}_2, \quad 2\mathbf{A}_4 \oplus 2\mathbf{A}_3 \oplus 2\mathbf{A}_2$$

to which Lemma 4 can not be applied and we need to compute the groups $E^+(\mathbf{T})$ directly from the definition (2.7). For

$$\mathbf{S} = 2\mathbf{A}_7 \oplus 2\mathbf{A}_2,$$

Table 4. Extremal singularities

Singularities	(p, q)	$ E^+ $	isometries of \mathbf{S} generating $E^+(\mathbf{T})$
$2\mathbf{E}_6 \oplus 2\mathbf{A}_3$	(2, 3)	4	symmetry of \mathbf{E}_6 ; $\mathbf{A}_3 \leftrightarrow \mathbf{A}_3$
$\mathbf{D}_9 \oplus \mathbf{A}_3 \oplus 3\mathbf{A}_2$	(2, 3)	8	symmetries of \mathbf{A}_2 , \mathbf{A}_3 ; $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{D}_8 \oplus \mathbf{A}_6 \oplus 2\mathbf{A}_2$	(2, 3)	2	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{D}_8 \oplus \mathbf{A}_3 \oplus 3\mathbf{A}_2$	(2, 3)	2	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{D}_7 \oplus \mathbf{D}_5 \oplus 3\mathbf{A}_2$	(2, 3)	8	symmetries of \mathbf{A}_2 , \mathbf{D}_5 ; $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{D}_7 \oplus \mathbf{D}_4 \oplus 3\mathbf{A}_2$	(2, 3)	2	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{D}_6 \oplus 2\mathbf{A}_4 \oplus 2\mathbf{A}_2$	(3, 5)	2	symmetry of \mathbf{A}_2
$2\mathbf{D}_5 \oplus 2\mathbf{A}_4$	(2, 5)	4	symmetry of \mathbf{D}_5 ; $\mathbf{A}_4 \leftrightarrow \mathbf{A}_4$
$\mathbf{D}_5 \oplus 2\mathbf{A}_4 \oplus 2\mathbf{A}_2 \oplus \mathbf{A}_1$	(3, 5)	4	symmetry of \mathbf{A}_2 ; $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{D}_4 \oplus 2\mathbf{A}_6 \oplus \mathbf{A}_2$	(2, 7)	2	symmetry of \mathbf{A}_6
$\mathbf{A}_{11} \oplus \mathbf{A}_3 \oplus 2\mathbf{A}_2$	(2, 3)	8	symmetries of \mathbf{A}_2 , \mathbf{A}_3 ; $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$\mathbf{A}_8 \oplus 2\mathbf{A}_3 \oplus 2\mathbf{A}_2$	(2, 3)	8	symmetries of \mathbf{A}_2 , \mathbf{A}_3 ; $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$2\mathbf{A}_6 \oplus 2\mathbf{A}_3$	(2, 7)	4	exceptional
$2\mathbf{A}_6 \oplus 3\mathbf{A}_2$	(3, 7)	4	symmetries of \mathbf{A}_2 , \mathbf{A}_6
$2\mathbf{A}_5 \oplus 2\mathbf{A}_4$	(3, 5)	2	symmetry of \mathbf{A}_4
$3\mathbf{A}_4 \oplus 2\mathbf{A}_2 \oplus 2\mathbf{A}_1$	(3, 5)	4	symmetry of \mathbf{A}_2 ; $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$
$2\mathbf{A}_4 \oplus 4\mathbf{A}_2$	(2, 3)	2	$\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$

which is the last exceptional case listed in statement, we have $\Sigma_2^\sharp(\mathbf{T}) \not\supset \Gamma_{2,2}$ and $|\cdot|_2$ needs to be modified by replacing χ_2 with $\chi_2(u) = u \bmod 8 \in \{1, 3, 5, 7\} = \mathbb{Z}_2^\times / (\mathbb{Z}_2^\times)^2$ and we have to consider the full group $\Gamma_{2,0}$ instead of $\Gamma'_{2,0}$. The computation can be summarized as follows:

	$\Gamma_{3,0}$		$\Gamma_{2,0}$		
generator of $\Sigma_3^\sharp(T)$	-1	-1	1	1	1
$\Sigma_2^\sharp(T) = \{1\}$	1	1	1	1	1
$\varphi(-1, -1)$	-1	-1	-1	-1	1

	δ_3	$ \cdot _3$	δ_2	$ \cdot _2$	
a symmetry of \mathbf{A}_2	-1	1	1	-1	-1
a transposition $\mathbf{A}_2 \leftrightarrow \mathbf{A}_2$	-1	-1	1	-1	-1
a symmetry of \mathbf{A}_7	1	1	-1	-1	1
a transposition $\mathbf{A}_7 \leftrightarrow \mathbf{A}_7$	1	-1	-1	-1	1

The rank of the matrix composed by the 7 rows of the table (see Remark 1) is $4 < \dim F_{3,0} + \dim F_{2,0}$, which implies that d_+^\perp is not surjective. For the other two cases, there are three irregular primes and the computation repeats literally that at the end of the proof of Proposition 5; in both cases, the map d_+^\perp turns out to be surjective. \square

References

- [1] Ayşegül Akyol and Alex Degtyarev, Geography of irreducible plane sextics, to appear in Proc. London Math. Soc, arXiv:1406.1491.
- [2] Nicolas Bourbaki, Lie groups and Lie algebras. Chapters 4–6, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley. MR **1890629** (**2003a**:17001)
- [3] Alex Degtyarev, Classification of quartic surfaces that have a nonsimple singular point, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1269–1290, 1337–1338. MR **1039964** (**91a**:14018)
- [4] Alex Degtyarev, Classification of quartics having a nonsimple singular point. II [MR1157141 (93e:14041)], Topology of manifolds and varieties, Adv. Soviet Math., vol. 18, Amer. Math. Soc., Providence, RI, 1994, pp. 23–54. MR 1296888
- [5] Alex Degtyarev, On deformations of singular plane sextics, J. Algebraic Geom. **17** (2008), no. 1, 101–135. MR **2357681** (**2008j**:14061)
- [6] Alex Degtyarev, Irreducible plane sextics with large fundamental groups, J. Math. Soc. Japan **61** (2009), no. 4, 1131–1169. MR **2588507** (**2011a**:14061)
- [7] Alex Degtyarev and Ilia Itenberg, On real determinantal quartics, Proceedings of the Gökova Geometry-Topology Conference 2010, Int. Press, Somerville, MA, 2011, pp. 110–128. MR 2931883
- [8] Alan H. Durfee, Fifteen characterizations of rational double points and simple critical points, Enseign. Math. (2) **25** (1979), no. 1–2, 131–163. MR **543555** (**80m**:14003)
- [9] E. B. Dynkin, Semisimple subalgebras of semisimple Lie algebras, Mat. Sbornik N.S. **30(72)** (1952), 349–462 (3 plates). MR 0047629 (13,904c)
- [10] The GAP Group, GAP—Groups, Algorithms, and Programming, Version 4.5.6, 2015.
- [11] Carl Friedrich Gauss, Disquisitiones arithmeticae, Springer-Verlag, New York, 1986, Translated and with a preface by Arthur A. Clarke, Revised by William C. Waterhouse, Cornelius Greither and A. W. Grootendorst and with a preface by Waterhouse. MR **837656** (**87f**:01105)

- [12] Viatcheslav Kharlamov, On the classification of nonsingular surfaces of degree 4 in \mathbf{RP}^3 with respect to rigid isotopies, *Funktsional. Anal. i Prilozhen.* **18** (1984), no. 1, 49–56. MR **739089** (85m:14034)
- [13] Vik. S. Kulikov, Surjectivity of the period mapping for $K3$ surfaces, *Uspehi Mat. Nauk* **32** (1977), no. 4(196), 257–258. MR 0480528 (58 #688)
- [14] Eduard Looijenga, The complement of the bifurcation variety of a simple singularity, *Invent. Math.* **23** (1974), 105–116. MR 0422675 (54 #10661)
- [15] Rick Miranda and David R. Morrison, The number of embeddings of integral quadratic forms. I, *Proc. Japan Acad. Ser. A Math. Sci.* **61** (1985), no. 10, 317–320. MR **834537** (87j:11031a)
- [16] Rick Miranda and David R. Morrison, The number of embeddings of integral quadratic forms. II, *Proc. Japan Acad. Ser. A Math. Sci.* **62** (1986), no. 1, 29–32. MR **839800** (87j:11031b)
- [17] Rick Miranda and David R. Morrison, Embeddings of integral quadratic forms, electronic, <http://www.math.ucsb.edu/~drm/manuscripts/eiqf.pdf>, 2009.
- [18] V. V. Nikulin, Finite groups of automorphisms of Kählerian $K3$ surfaces, *Trudy Moskov. Mat. Obshch.* **38** (1979), 75–137. MR **544937** (81e:32033)
- [19] V. V. Nikulin, Integer symmetric bilinear forms and some of their geometric applications, *Izv. Akad. Nauk SSSR Ser. Mat.* **43** (1979), no. 1, 111–177, 238. MR **525944** (80j:10031)
- [20] Ulf Persson, Double sextics and singular $K3$ surfaces, *Algebraic geometry, Sitges (Barcelona)*, 1983, *Lecture Notes in Math.*, vol. 1124, Springer, Berlin, 1985, pp. 262–328. MR **805337** (87i:14036)
- [21] I. I. Pjateckiĭ-Šapiro and I. R. Šafarevič, Torelli's theorem for algebraic surfaces of type $K3$, *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 530–572. MR 0284440
- [22] B. Saint-Donat, Projective models of $K - 3$ surfaces, *Amer. J. Math.* **96** (1974), 602–639. MR 0364263 (51 #518)
- [23] Tohsuke Urabe, Dynkin graphs and combinations of singularities on quartic surfaces, *Proc. Japan Acad. Ser. A Math. Sci.* **61** (1985), no. 8, 266–269. MR **816911** (87b:14017)
- [24] Tohsuke Urabe, Singularities in a certain class of quartic surfaces and sextic curves and Dynkin graphs, *Proceedings of the 1984 Vancouver conference in algebraic geometry, CMS Conf. Proc.*, vol. 6, Amer. Math. Soc., Providence, RI, 1986, pp. 477–497. MR **846037** (87h:14030)
- [25] Tohsuke Urabe, Elementary transformations of Dynkin graphs and singularities on quartic surfaces, *Invent. Math.* **87** (1987), no. 3, 549–572. MR **874036** (88b:14031)
- [26] Tohsuke Urabe, Tie transformations of Dynkin graphs and singularities on quartic surfaces, *Invent. Math.* **100** (1990), no. 1, 207–230. MR **1037147** (91b:14043)
- [27] F. van der Blij, An invariant of quadratic forms mod 8, *Nederl. Akad. Wetensch. Proc. Ser. A* **62** = *Indag. Math.* **21** (1959), 291–293. MR 0108467 (21 #7183)
- [28] Jin-Gen Yang, Enumeration of combinations of rational double points on quartic surfaces, *Singularities and complex geometry (Beijing, 1994)*, *AMS/IP Stud. Adv. Math.*, vol. 5, Amer. Math. Soc., Providence, RI, 1997, pp. 275–312. MR 1468283

Çisem Güneş Aktaş

Department of Mathematics

Bilkent University, 06800 Ankara, Turkey

E-mail: cisem@fen.bilkent.edu.tr