

# involve

a journal of mathematics

Bases for the global Weyl modules  
of  $\mathfrak{sl}_n$  of highest weight  $m\omega_1$

Samuel Chamberlin and Amanda Croan



# Bases for the global Weyl modules of $\mathfrak{sl}_n$ of highest weight $m\omega_1$

Samuel Chamberlin and Amanda Croan

(Communicated by Jim Haglund)

We utilize a theorem of B. Feigin and S. Loktev to give explicit bases for the global Weyl modules for the map algebras of the form  $\mathfrak{sl}_n \otimes A$  of highest weight  $m\omega_1$ . These bases are given in terms of specific elements of the universal enveloping algebra,  $U(\mathfrak{sl}_n \otimes A)$ , acting on the highest weight vector.

## 1. Introduction

Let  $\mathfrak{g}$  be a simple finite-dimensional complex Lie algebra. For the loop algebras  $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$ , the global Weyl modules were introduced by Chari and Pressley [2001]. Feigin and Loktev [2004] extended these global Weyl modules to the case where the Laurent polynomials above were replaced by the coordinate ring of a complex affine variety. Chari, Fourier and Khandaï [Chari et al. 2010] then generalized this definition to the map algebras,  $\mathfrak{g} \otimes A$ , where  $A$  is a commutative, associative, complex unital algebra. Feigin and Loktev [2004] also gave an isomorphism which explicitly determines the structure of the global Weyl modules for the map algebras of  $\mathfrak{sl}_n$  of highest weight  $m\omega_1$ .

The goal of this work is to use the structure isomorphism given by Feigin and Loktev to give nice bases for the global Weyl modules for the map algebras  $\mathfrak{sl}_n \otimes A$  of  $\mathfrak{sl}_n$  of highest weight  $m\omega_1$ . These bases will be given in terms of specific elements of  $U(\mathfrak{sl}_n \otimes A)$  acting on the highest weight vector. This was done in [Chamberlin 2011] in the case  $n = 2$ , but the case  $n > 2$  has not previously appeared in the literature.

## 2. Preliminaries

**2.1. The structure of  $\mathfrak{sl}_n$ .** Recall that  $\mathfrak{sl}_n$  is the Lie algebra of all complex traceless matrices. The Lie bracket is the commutator bracket given by  $[A, B] = AB - BA$ .

---

MSC2010: 17B10.

Keywords: Lie algebra, module, representation, Weyl.

Given any matrix  $[b_{i,j}]$ , define  $\varepsilon_k([b_{i,j}]) := b_{k,k}$ . For  $i \in \{1, \dots, n-1\}$ , define  $\alpha_i := \varepsilon_i - \varepsilon_{i+1}$ . Define

$$R^\pm := \{\pm(\alpha_i + \dots + \alpha_j) \mid 1 \leq i < j \leq n-1\}$$

to be the positive and negative roots respectively, and define  $R = R^+ \cup R^-$  to be the set of roots. Let  $e_{i,j}$  be the  $n \times n$  matrix with a one in the  $i$ -th row and  $j$ -th column and zeros in every other position. Define  $h_i := h_{\alpha_i} = e_{i,i} - e_{i+1,i+1}$ , for  $i \in \{1, \dots, n-1\}$ . Then  $\mathfrak{h} := \text{span}\{h_i \mid 1 \leq i \leq n\}$  is a Cartan subalgebra of  $\mathfrak{sl}_n$ . Given  $\alpha = \alpha_i + \dots + \alpha_j \in R^+$ , define  $x_\alpha := e_{i,j}$  and  $x_{-\alpha} := e_{j,i}$ . Then  $\{h_i, x_{\pm\alpha} \mid 1 \leq i \leq n-1, \alpha \in R\}$  is a Chevalley basis for  $\mathfrak{sl}_n$ . Given  $i \in \{1, \dots, n-1\}$ , define  $x_i := x_{\alpha_i} = e_{i,i+1}$  and  $x_{-i} := x_{-\alpha_i} = e_{i+1,i}$ . Note that, for all  $1 \leq i \leq n-1$ ,  $\text{span}\{x_{-i}, h_i, x_i\} \cong \mathfrak{sl}_2$ .

Define nilpotent subsuperalgebras  $\mathfrak{n}^\pm := \text{span}\{x_\alpha \mid \alpha \in R^\pm\}$  and note that  $\mathfrak{sl}_n = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ . Define the set of fundamental weights  $\{\omega_1, \dots, \omega_{n-1}\} \subset \mathfrak{h}^*$  by  $\omega_i(h_j) = \delta_{i,j}$  for all  $i, j \in \{1, \dots, n-1\}$ . Define  $P^+ := \text{span}_{\mathbb{Z}_{\geq 0}}\{\omega_1, \dots, \omega_{n-1}\}$  to be the set of dominant integral weights.

**2.2. Map algebras and Weyl modules.** For the remainder of this work fix a commutative, associative, complex unital algebra  $A$ . Define the map algebra of  $\mathfrak{sl}_n$  to be  $\mathfrak{sl}_n \otimes A$  with Lie bracket given by linearly extending the bracket

$$[z \otimes a, w \otimes b] = [z, w] \otimes ab$$

for all  $z, w \in \mathfrak{sl}_n$  and  $a, b \in A$ .

Define  $\mathbf{U}(\mathfrak{sl}_n \otimes A)$  to be the universal enveloping algebra of  $\mathfrak{sl}_n \otimes A$ .

As in [Chari et al. 2010] we define the global Weyl model for  $\mathfrak{sl}_n \otimes A$  of highest weight  $\lambda \in P^+$  to be the module generated by a vector  $w_\lambda$ , called the highest weight vector, with relations

$$(x \otimes a)w_\lambda = 0, \quad (h \otimes 1)w_\lambda = \lambda(h)w_\lambda, \quad (x_{-i} \otimes 1)^{\lambda(h_i)+1}w_\lambda = 0,$$

for all  $a \in A$ ,  $x \in \mathfrak{n}^+$ ,  $h \in \mathfrak{h}$ , and  $1 \leq i \leq n-1$ .

**2.3. Multisets.** Given any set  $S$ , define a multiset of elements of  $S$  to be a multiplicity function  $\chi : S \rightarrow \mathbb{Z}_{\geq 0}$ . Define  $\mathcal{F}(S) := \{\chi : S \rightarrow \mathbb{Z}_{\geq 0} : |\text{supp } \chi| < \infty\}$ . For  $\chi \in \mathcal{F}(S)$ , define  $|\chi| := \sum_{s \in S} \chi(s)$ . Notice that  $\mathcal{F}(S)$  is an abelian monoid under function addition. For  $\psi, \chi \in \mathcal{F}(S)$ , we write  $\psi \subseteq \chi$  if  $\psi(s) \leq \chi(s)$  for all  $s \in S$ . Define  $\mathcal{F}(\chi)(S) := \{\psi \in \mathcal{F}(S) \mid \psi \subseteq \chi\}$ . In the case  $S = A$ , the  $S$  will be omitted from the notation, so that  $\mathcal{F} := \mathcal{F}(A)$  and  $\mathcal{F}(\chi) = \mathcal{F}(\chi)(A)$ .

If  $\psi, \chi \in \mathcal{F}$  with  $\psi \in \mathcal{F}(\chi)$  we define  $\chi - \psi$  by standard function subtraction.

Also define  $\pi : \mathcal{F} - \{0\} \rightarrow A$  by

$$\pi(\psi) := \prod_{a \in A} a^{\psi(a)},$$

and extend  $\pi$  to  $\mathcal{F}$  by setting  $\pi(0) = 1$ . Define  $\mathcal{M} : \mathcal{F} \rightarrow \mathbb{Z}$  by

$$\mathcal{M}(\psi) := \frac{|\psi|!}{\prod_{a \in A} \psi(a)!}.$$

$\mathcal{M}(\psi) \in \mathbb{Z}$  for all  $\psi \in \mathcal{F}$  because if  $\text{supp } \psi = \{a_1, \dots, a_k\}$  then  $\mathcal{M}(\psi)$  is the multinomial coefficient

$$\binom{|\psi|}{\psi(a_1), \dots, \psi(a_k)}.$$

For  $s \in S$ , define  $\chi_s$  to be the characteristic function of the set  $\{s\}$ . Then, for all  $\chi \in \mathcal{F}(S)$ ,

$$\chi = \sum_{s \in S} \chi(s) \chi_s.$$

**2.4. The symmetric tensor space.** Given any vector space  $W$ , there is an action of the symmetric group  $S_k$  on  $W^{\otimes k} = W \otimes W \otimes \cdots \otimes W$  ( $k$  times) given by

$$\sigma(w_1 \otimes w_2 \otimes \cdots \otimes w_k) = w_{\sigma^{-1}(1)} \otimes w_{\sigma^{-1}(2)} \otimes \cdots \otimes w_{\sigma^{-1}(k)}, \quad \text{where } w_1, \dots, w_k \in W.$$

For any vector space  $W$ , define its  $k$ -th symmetric tensor space

$$S^k(W) = \text{span} \left\{ \sum_{\sigma \in S_k} \sigma(w_1 \otimes \cdots \otimes w_k) \mid w_1, \dots, w_k \in W \right\}.$$

Define  $V \cong \mathbb{C}^n$  to be an  $\mathfrak{sl}_n$ -module via left matrix multiplication and write the basis as  $v_1 := (1, 0, \dots, 0)$ , and for  $i \in \{1, \dots, n+m-1\}$ , set  $v_{i+1} := x_{-i} v_i$ . Then  $V \otimes A$  is an  $\mathfrak{sl}_n \otimes A$ -module under the action  $(z \otimes a)(w \otimes b) = zw \otimes ab$ .

Given  $\varphi_1, \dots, \varphi_n \in \mathcal{F}$  with  $k := \sum_{i=1}^n |\varphi_i|$ , define

$$w(\varphi_1, \dots, \varphi_n) := \bigotimes_{a_1 \in \text{supp } \varphi_1} (v_1 \otimes a_1)^{\otimes \varphi_1(a_1)} \otimes \cdots \otimes \bigotimes_{a_n \in \text{supp } \varphi_n} (v_n \otimes a_n)^{\otimes \varphi_n(a_n)} \in (V \otimes A)^{\otimes k}$$

and

$$v(\varphi_1, \dots, \varphi_n) := \sum_{\sigma \in S_k} \sigma(w(\varphi_1, \dots, \varphi_n)) \in S^k(V \otimes A).$$

We will need the following theorem:

**Theorem 2.4.1** [Feigin and Loktev 2004, Theorem 6]. *For all  $m \in \mathbb{N}$ ,  $W_A(m\omega_1) \cong S^m(V \otimes A)$  via the map given by*

$$w_{m\omega_1} \mapsto (v_1 \otimes 1)^{\otimes m}.$$

We will also need the following lemma:

**Lemma 2.4.2.** *Let  $\mathbf{B}$  be a basis for  $A$ . Then the set*

$$\mathfrak{B} := \left\{ v(\varphi_1, \dots, \varphi_n) \mid \varphi_1, \dots, \varphi_n \in \mathcal{F}(\mathbf{B}), \sum_{i=1}^n |\varphi_i| = m \right\}$$

is a basis for  $S^m(V \otimes A)$ .

*Proof.*  $\mathfrak{B}$  spans  $S^m(V \otimes A)$  because  $\mathbf{B}$  spans  $A$  and  $v_1, \dots, v_n$  span  $V$ .  $\mathfrak{B}$  is linearly independent because the set

$$\{(v_{j_1} \otimes b_1) \otimes \cdots \otimes (v_{j_m} \otimes b_m) \mid j_1, \dots, j_m \in \{1, \dots, n\}, b_1, \dots, b_m \in \mathbf{B}\}$$

is a basis for  $(V \otimes A)^{\otimes m}$  and hence is linearly independent.  $\square$

Given  $k \in \mathbb{N}$ , define  $\Delta^{k-1} : \mathbf{U}(\mathfrak{sl}_n \otimes A) \rightarrow \mathbf{U}(\mathfrak{sl}_n \otimes A)^{\otimes k}$  by extending the map  $\mathfrak{sl}_n \otimes A \rightarrow \mathbf{U}(\mathfrak{sl}_n \otimes A)^{\otimes k}$  given by

$$\Delta^{k-1}(z \otimes a) = \sum_{j=0}^{k-1} 1^{\otimes j} \otimes (z \otimes a) \otimes 1^{\otimes k-1-j}.$$

Note that  $\Delta^{k-1}(1) = 1^{\otimes k}$ , not  $k1^{\otimes k}$ .

Since  $V \otimes A$  is a  $\mathbf{U}(\mathfrak{sl}_n \otimes A)$ -module,  $(V \otimes A)^{\otimes m}$  is a left  $\mathbf{U}(\mathfrak{sl}_n \otimes A)$ -module with  $u$  acting as  $\Delta^{m-1}(u)$  followed by coordinatewise module actions. Moreover  $S^m(V \otimes A)$  is a submodule under this action. Thus  $S^m(V \otimes A)$  is a left  $\mathbf{U}(\mathfrak{sl}_n \otimes A)$ -module under this  $\Delta^{m-1}$  action.

**2.5.** For all  $i = 1, \dots, n-1$  and  $\chi, \varphi \in \mathcal{F}$ , recursively define  $q_i(\varphi, \chi) \in \mathbf{U}(\mathfrak{sl}_n \otimes A)$  as follows:

$$q_i(0, 0) := 1,$$

$$q_i(0, \chi) := -\frac{1}{|\chi|} \sum_{0 \neq \psi \in \mathcal{F}(\chi)} \mathcal{M}(\psi)(h_i \otimes \pi(\psi)) q_i(0, \chi - \psi),$$

$$q_i(\varphi, \chi) := -\frac{1}{|\varphi|} \sum_{\psi \in \mathcal{F}(\chi)} \sum_{d \in \text{supp } \varphi} \mathcal{M}(\psi)(x_{-i} \otimes d\pi(\psi)) q_i(\varphi - \chi_d, \chi - \psi).$$

Given  $\varphi_n, \dots, \varphi_1 \in \mathcal{F}$ , define

$$\begin{aligned} q(\varphi_1, \dots, \varphi_n) &:= q_{n-1}(\varphi_n, \varphi_{n-1}) q_{n-2}((|\varphi_n| + |\varphi_{n-1}|)\chi_1, \varphi_{n-2}) \\ &\quad \times \cdots \times q_2\left(\left(\sum_{j=3}^n |\varphi_j|\right)\chi_1, \varphi_2\right) q_1\left(\left(\sum_{k=2}^n |\varphi_j|\right)\chi_1, \varphi_1\right). \end{aligned}$$

**Remark.** The  $q_i(0, \chi)$  coincide with the  $p_i(\chi)$  defined in [Bagci and Chamberlin 2014].

### 3. Main theorem

The main result of this work is the theorem stated below.

**Theorem 3.0.1.** *Given a basis  $\mathbf{B}$  for  $A$  and  $m \in \mathbb{Z}_{>0}$ , the set*

$$\left\{ q(\varphi_1, \dots, \varphi_n) w_{m\omega_1} \mid \varphi_1, \dots, \varphi_n \in \mathcal{F}(\mathbf{B}), \sum_{i=1}^n |\varphi_i| = m \right\}$$

*is a basis for  $W_A(m\omega_1)$ .*

The proof of this theorem will be given after several lemmas and propositions.

#### 3.1. Necessary lemmas and propositions.

**Proposition 3.1.1.** *For all  $k \in \mathbb{N}$   $\Delta^k = (1^{\otimes k-1} \otimes \Delta^1) \circ \Delta^{k-1}$ .*

*Proof.* The case  $k = 1$  is trivial. For  $k \geq 2$  and  $u \in U(\mathfrak{sl}_n \otimes A)$  we have

$$\begin{aligned} (1^{\otimes k-1} \otimes \Delta^1)(\Delta^{k-1}(u)) &= (1^{\otimes k-1} \otimes \Delta^1)\left(\sum_{j=0}^{k-1} 1^{\otimes j} \otimes u \otimes 1^{\otimes k-1-j}\right) \\ &= (1^{\otimes k-1} \otimes \Delta^1)\left(\sum_{j=0}^{k-2} 1^{\otimes j} \otimes u \otimes 1^{\otimes k-1-j} + 1^{\otimes k-1} \otimes u\right) \\ &= \sum_{j=0}^{k-2} 1^{\otimes j} \otimes u \otimes 1^{\otimes k-2-j} \otimes \Delta^1(1) + 1^{\otimes k-1} \otimes \Delta^1(u) \\ &= \sum_{j=0}^{k-2} 1^{\otimes j} \otimes u \otimes 1^{\otimes k-2-j} \otimes 1 \otimes 1 + 1^{\otimes k-1} \otimes (u \otimes 1 + 1 \otimes u) \\ &= \sum_{j=0}^{k-2} 1^{\otimes j} \otimes u \otimes 1^{\otimes k-j} + 1^{\otimes k-1} \otimes u \otimes 1 + 1^{\otimes k-1} \otimes 1 \otimes u \\ &= \sum_{j=0}^k 1^{\otimes j} \otimes u \otimes 1^{\otimes k-j} = \Delta^k(u). \end{aligned}$$

□

Given  $\chi \in \mathcal{F}$  and  $k \in \mathbb{N}$ , define

$$\circ_k(\chi) = \left\{ \psi : \{1, \dots, k\} \rightarrow \mathcal{F}(\chi) \mid \sum_{j=1}^k \psi(j) = \chi \right\}.$$

**Lemma 3.1.2.** *For all  $i \in \{1, \dots, n-1\}$ ,*

$$\Delta^{k-1}(q_i(\varphi, \chi)) = \sum_{\substack{\psi \in \circ_k(\chi) \\ \phi \in \circ_k(\varphi)}} q_i(\phi(1), \psi(1)) \otimes \cdots \otimes q_i(\phi(k), \psi(k))$$

*Proof.* This can be proven by induction on  $k$ . The case  $k = 1$  is trivial. In the case  $k = 2$  the lemma becomes

$$\Delta^1(q_i(\varphi, \chi)) = \sum_{\substack{\psi \in \mathcal{F}(\chi) \\ \phi \in \mathcal{F}(\varphi)}} q_i(\phi, \psi) \otimes q_i(\varphi - \phi, \chi - \psi).$$

This can be proven by induction on  $|\varphi|$ . For  $k > 2$  use [Proposition 3.1.1](#). The details in the  $\mathfrak{sl}_2$  case can be found in [\[Chamberlin 2011\]](#). This can be extended to the  $\mathfrak{sl}_n$  case via the injection  $\Omega_i : \mathfrak{sl}_2 \otimes A \rightarrow \mathfrak{sl}_n \otimes A$  given by

$$\Omega_i(x^- \otimes a) = x_{-i} \otimes a, \quad \Omega_i(h \otimes a) = h_i \otimes a, \quad \Omega_i(x^+ \otimes a) = x_i \otimes a,$$

for all  $i \in \{1, \dots, n-1\}$  and  $a \in A$ .  $\square$

**Lemma 3.1.3.** *For all  $\varphi, \chi \in \mathcal{F}$  with  $|\varphi| + |\chi| > 1$  and all  $i \in \{1, \dots, n-1\}$ ,  $q_i(\varphi, \chi)(v_i \otimes 1) = 0$ .*

*Proof.* Assume that  $\varphi = 0$ . This case will proceed by induction on  $|\chi| > 1$ . If  $|\chi| = 2$  (so that  $\chi = \{a, b\}$  for some  $a, b \in A$ ) we have

$$\begin{aligned} q_i(0, \{a, b\})(v_i \otimes 1) &= [(h_i \otimes a) \otimes (h_i \otimes b) - (h_i \otimes ab)](v_i \otimes 1) \\ &= (h_i \otimes a) \otimes (v_i \otimes b) - (v_i \otimes ab) \\ &= (v_i \otimes ab) - (v_i \otimes ab) \\ &= 0. \end{aligned}$$

For the next case assume that  $|\chi| > 2$  then

$$q_i(0, \chi)(v_i \otimes 1) = -\frac{1}{|\chi|} \sum_{\emptyset \neq \psi \in \mathcal{F}(\chi)} \mathcal{M}(\psi)(h_i \otimes \pi(\psi)) q_i(\chi - \psi)(v_i \otimes 1) = 0$$

by induction. Now assume that  $|\varphi| = 1$  (or  $\varphi = \chi_b$  for some  $b \in A$ ). Then

$$\begin{aligned} q_i(\chi_b, \chi)(v_i \otimes 1) &= - \sum_{\psi \in \mathcal{F}(\chi)} \mathcal{M}(\psi)(x_{-i} \otimes b\pi(\psi)) q_i(0, \chi - \psi)(v_i \otimes 1) \\ &= -\mathcal{M}(\chi)(x_{-i} \otimes b\pi(\chi))(v_i \otimes 1) \\ &\quad - \sum_{a \in \text{supp } \chi} \mathcal{M}(\chi - \chi_a)(x_{-i} \otimes b\pi(\chi - \chi_a)) q_i(0, \chi_a)(v_i \otimes 1) \\ &= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) \\ &\quad - \sum_{a \in \text{supp } \chi} \mathcal{M}(\chi - \chi_a)(x_{-i} \otimes b\pi(\chi - \chi_a)) (-h_i \otimes a)(v_i \otimes 1) \\ &= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) + \sum_{a \in \text{supp } \chi} \mathcal{M}(\chi - \chi_a)(x_{-i} \otimes b\pi(\chi - \chi_a))(v_i \otimes a) \end{aligned}$$

$$\begin{aligned}
&= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) + \sum_{a \in \text{supp } \chi} \mathcal{M}(\chi - \chi_a)(v_{i+1} \otimes b\pi(\chi)) \\
&= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) + \sum_{a \in \text{supp } \chi} \frac{(|\chi| - 1)!}{\prod_{c \in \text{supp } (\chi - \chi_a)} (\chi - \chi_a)(c)!} (v_{i+1} \otimes b\pi(\chi)) \\
&= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) + \sum_{a \in \text{supp } \chi} \frac{(|\chi| - 1)!}{\prod_{\substack{c \in \text{supp } \chi \\ c \neq a}} \chi(c)! (\chi(a) - 1)!} (v_{i+1} \otimes b\pi(\chi)) \\
&= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) + \sum_{a \in \text{supp } \chi} \frac{\chi(a)}{|\chi|} \mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) \\
&= -\mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) + \mathcal{M}(\chi)(v_{i+1} \otimes b\pi(\chi)) \\
&= 0.
\end{aligned}$$

Finally assume that  $|\varphi| > 1$ . Then

$$\begin{aligned}
q_i(\varphi, \chi)(v_i \otimes 1) &= -\frac{1}{|\varphi|} \sum_{\psi \in \mathcal{F}(\chi)} \sum_{d \in \text{supp } \varphi} \mathcal{M}(\psi)(x_{-i} \otimes d\pi(\psi)) q_i(\varphi - \chi_d, \chi - \psi)(v_i \otimes 1) \\
&= -\frac{1}{|\varphi|} \sum_{\psi \in \mathcal{F}(\chi)} \sum_{d \in \text{supp } \varphi} \mathcal{M}(\psi) \\
&\quad \left( -\frac{1}{|\varphi| - 1} \sum_{\psi_1 \in \mathcal{F}(\chi - \psi)} \sum_{d_1 \in \text{supp } (\varphi - \chi_d)} \mathcal{M}(\psi_1) \right. \\
&\quad \left. (x_{-i} \otimes d\pi(\psi)) (x_{-i} \otimes d_1\pi(\psi_1)) q_i(\varphi - \chi_d - \chi_{d_1}, \chi - \psi - \psi_1) \right) (v_i \otimes 1) \\
&= 0,
\end{aligned}$$

because at least two  $x_{-i}$  terms act on a single  $v_i$  as 0.  $\square$

**Lemma 3.1.4.** *For all  $i \in \{1, \dots, n-1\}$  and  $\varphi, \chi \in \mathcal{F}$  with  $|\varphi| + |\chi| = k$  we have*

$$q_i(\varphi, \chi)(v_i \otimes 1)^{\otimes k} = (-1)^k v(0, \dots, 0, \chi, \varphi, 0, \dots, 0),$$

where  $\chi$  is in the  $i$ -th position and  $\varphi$  in the  $(i+1)$ -st.

*Proof.* We have  $q_i(\varphi, \chi)(v_i \otimes 1)^{\otimes k} i = \Delta^{k-1}(q_i(\varphi, \chi))(v_i \otimes 1)^{\otimes k}$ . By Lemma 3.1.2, this equals

$$\left( \sum_{\substack{\psi \in \circ_k(\chi) \\ \phi \in \circ_k(\varphi)}} q_i(\phi(1), \psi(1)) \otimes \cdots \otimes q_i(\phi(k), \psi(k)) \right) (v_i \otimes 1)^{\otimes k},$$

which can be rewritten as

$$\sum_{\substack{\psi \in \circ_k(\chi) \\ \phi \in \circ_k(\varphi)}} (q_i(\phi(1), \psi(1))(v_i \otimes 1)) \otimes \cdots \otimes (q_i(\phi(k), \psi(k))(v_i \otimes 1)).$$

By Lemma 3.1.3 we see that the only potentially nonzero terms in the sum are those for which  $|\phi(j)| + |\psi(j)| \leq 1$  for all  $j \in \{1, \dots, k\}$ . Since  $|\varphi| + |\chi| = k$  if we have  $|\psi(j)| + |\phi(j)| = 0$  for some  $j \in \{1, \dots, n-1\}$ , then there is a  $r \in \{1, \dots, n-1\}$  such that  $|\psi(r)| + |\phi(r)| > 1$ . So the only potentially nonzero terms in the sum are those for which  $|\phi(j)| + |\psi(j)| = 1$  for all  $j \in \{1, \dots, k\}$ . Suppose that  $\phi(j) = \chi_a$  and  $\psi(j) = 0$  for some  $j \in \{1, \dots, k\}$  and some  $a \in A$ . Then

$$q_i(\chi_a, 0)(v_i \otimes 1) = -(x_{-i} \otimes a)(v_i \otimes 1) = -(v_{i+1} \otimes a).$$

Suppose that  $\phi(j) = 0$  and  $\psi(j) = \chi_a$  for some  $j \in \{1, \dots, k\}$  and some  $a \in A$ . Then

$$q_i(0, \chi_a)(v_i \otimes 1) = -(h_i \otimes a)(v_i \otimes 1) = -(v_i \otimes a).$$

So  $-(v_{i+1} \otimes a)$  and  $-(v_i \otimes a)$  are the only possibilities for factors in the tensor product above. Since we are summing over all possible submultisets of  $\varphi$  and  $\chi$ , we have the result.  $\square$

**Lemma 3.1.5.** *For all  $m \in \mathbb{N}$  and all  $\varphi_1, \dots, \varphi_n \in \mathcal{F}$  with  $\sum_{i=1}^n |\varphi_i| = m$ ,*

$$q(\varphi_1, \dots, \varphi_n)(v_1 \otimes 1)^{\otimes m} = (-1)^{\sum_{j=1}^n j|\varphi_j|} v(\varphi_1, \dots, \varphi_n).$$

*Proof.* Since for all  $j \in \{1, \dots, n-1\}$  and  $k \in \{1, \dots, n\}$ ,

$$x_{-j} v_k = \delta_{j,k} v_{j+1}, \quad h_j v_k = \delta_{j,k} v_j - \delta_{j+1,k} v_{j+1},$$

so by Lemma 3.1.4 we have

$$\begin{aligned} q(\varphi_1, \dots, \varphi_n)(v_1 \otimes 1)^{\otimes m} &= q_{n-1}(\varphi_n, \varphi_{n-1}) q_{n-2}\left((|\varphi_n| + |\varphi_{n-1}|)\chi_1, \varphi_{n-2}\right) \\ &\quad \times \cdots \times q_1\left(\left(\sum_{j=2}^n |\varphi_j|\right)\chi_1, \varphi_1\right) (v_1 \otimes 1)^{\otimes m} \\ &= (-1)^m q_{n-1}(\varphi_n, \varphi_{n-1}) \cdots q_2\left(\left(\sum_{j=3}^n |\varphi_j|\right)\chi_1, \varphi_2\right) v\left(\varphi_1, \left(\sum_{j=2}^n |\varphi_j|\right)\chi_1, 0, \dots, 0\right) \\ &= (-1)^{|\varphi_1| + 2 \sum_{j=2}^n |\varphi_j|} q_{n-1}(\varphi_n, \varphi_{n-1}) \\ &\quad \times \cdots \times q_3\left(\left(\sum_{j=4}^n |\varphi_j|\right)\chi_1, \varphi_3\right) vi\left(\varphi_1, \varphi_2, \left(\sum_{j=3}^n |\varphi_j|\right)\chi_1, 0, \dots, 0\right) \end{aligned}$$

$$\begin{aligned}
 &= (-1)^{\sum_{j=1}^{n-2} j|\varphi_j|} q_{n-1}(\varphi_n, \varphi_{n-1}) v(\varphi_1, \dots, \varphi_{n-2}, (|\varphi_{n-1}| + |\varphi_n|)\chi_1, 0) \\
 &= (-1)^{\sum_{j=1}^n j|\varphi_j|} v(\varphi_1, \dots, \varphi_n).
 \end{aligned}$$

□

### 3.2. The proof of Theorem 3.0.1.

*Proof.* By Lemmas 3.1.5 and 2.4.2

$$\left\{ q(\varphi_1, \dots, \varphi_n)(v_1 \otimes 1)^{\otimes m} \mid \varphi_1, \dots, \varphi_n \in \mathcal{F}(\mathbf{B}), \sum_{i=1}^n |\varphi_i| = m \right\}$$

is a basis for  $S^m(V \otimes A)$ . Therefore by Theorem 2.4.1

$$\left\{ q(\varphi_1, \dots, \varphi_n)w_{m\omega_1} \mid \varphi_1, \dots, \varphi_n \in \mathcal{F}(\mathbf{B}), \sum_{i=1}^n |\varphi_i| = m \right\}$$

is a basis for  $W_A(m\omega_1)$ .

□

## References

- [Bagci and Chamberlin 2014] I. Bagci and S. Chamberlin, “Integral bases for the universal enveloping algebras of map superalgebras”, *J. Pure Appl. Algebra* **218**:8 (2014), 1563–1576. [MR](#) [Zbl](#)
- [Chamberlin 2011] S. H. Chamberlin, *Integral bases for the universal enveloping algebras of map algebras*, Ph.D. thesis, University of California, Riverside, 2011, <http://search.proquest.com/docview/883977727>. [MR](#)
- [Chari and Pressley 2001] V. Chari and A. Pressley, “Weyl modules for classical and quantum affine algebras”, *Represent. Theory* **5** (2001), 191–223. [MR](#) [Zbl](#)
- [Chari et al. 2010] V. Chari, G. Fourier, and T. Khandai, “A categorical approach to Weyl modules”, *Transform. Groups* **15**:3 (2010), 517–549. [MR](#) [Zbl](#)
- [Feigin and Loktev 2004] B. Feigin and S. Loktev, “Multi-dimensional Weyl modules and symmetric functions”, *Comm. Math. Phys.* **251**:3 (2004), 427–445. [MR](#) [Zbl](#)

Received: 2015-06-29 Accepted: 2015-08-25

|  |  |
|--|--|
| <a href="mailto:samuel.chamberlin@park.edu">samuel.chamberlin@park.edu</a> | <i>Department of Information Systems, Computer Science and Mathematics, Park University, 8700 NW River Park Drive #30, Parkville, 64152, United States</i> |
| <a href="mailto:amanda.croan@park.edu">amanda.croan@park.edu</a>           | <i>Department of Information Systems, Computer Science and Mathematics, Park University, 8700 NW River Park Drive #30, Parkville, 64152, United States</i> |

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

|                      |   |                        |   |
|----------------------|---|------------------------|---|
| Colin Adams          | Williams College, USA                     | Suzanne Lenhart        | University of Tennessee, USA              |
| John V. Baxley       | Wake Forest University, NC, USA           | Chi-Kwong Li           | College of William and Mary, USA          |
| Arthur T. Benjamin   | Harvey Mudd College, USA                  | Robert B. Lund         | Clemson University, USA                   |
| Martin Bohner        | Missouri U of Science and Technology, USA | Gaven J. Martin        | Massey University, New Zealand            |
| Nigel Boston         | University of Wisconsin, USA              | Mary Meyer             | Colorado State University, USA            |
| Amarjit S. Budhiraja | U of North Carolina, Chapel Hill, USA     | Emil Minchev           | Ruse, Bulgaria                            |
| Pietro Cerone        | La Trobe University, Australia            | Frank Morgan           | Williams College, USA                     |
| Scott Chapman        | Sam Houston State University, USA         | Mohammad Sal Moslehian | Ferdowsi University of Mashhad, Iran      |
| Joshua N. Cooper     | University of South Carolina, USA         | Zuhair Nashed          | University of Central Florida, USA        |
| Jem N. Corcoran      | University of Colorado, USA               | Ken Ono                | Emory University, USA                     |
| Toka Diagana         | Howard University, USA                    | Timothy E. O'Brien     | Loyola University Chicago, USA            |
| Michael Dorff        | Brigham Young University, USA             | Joseph O'Rourke        | Smith College, USA                        |
| Sever S. Dragomir    | Victoria University, Australia            | Yuval Peres            | Microsoft Research, USA                   |
| Behrouz Emamizadeh   | The Petroleum Institute, UAE              | Y.-F. S. Pétermann     | Université de Genève, Switzerland         |
| Joel Foisy           | SUNY Potsdam, USA                         | Robert J. Plemmons     | Wake Forest University, USA               |
| Errin W. Fulp        | Wake Forest University, USA               | Carl B. Pomerance      | Dartmouth College, USA                    |
| Joseph Gallian       | University of Minnesota Duluth, USA       | Vadim Ponomarenko      | San Diego State University, USA           |
| Stephan R. Garcia    | Pomona College, USA                       | Bjorn Poonen           | UC Berkeley, USA                          |
| Anant Godbole        | East Tennessee State University, USA      | James Propp            | U Mass Lowell, USA                        |
| Ron Gould            | Emory University, USA                     | Józeph H. Przytycki    | George Washington University, USA         |
| Andrew Granville     | Université Montréal, Canada               | Richard Rebarber       | University of Nebraska, USA               |
| Jerrold Griggs       | University of South Carolina, USA         | Robert W. Robinson     | University of Georgia, USA                |
| Sat Gupta            | U of North Carolina, Greensboro, USA      | Filip Saidak           | U of North Carolina, Greensboro, USA      |
| Jim Haglund          | University of Pennsylvania, USA           | James A. Sellers       | Penn State University, USA                |
| Johnny Henderson     | Baylor University, USA                    | Andrew J. Sterge       | Honorary Editor                           |
| Jim Hoste            | Pitzer College, USA                       | Ann Trenk              | Wellesley College, USA                    |
| Natalia Hritonenko   | Prairie View A&M University, USA          | Ravi Vakil             | Stanford University, USA                  |
| Glenn H. Hurlbert    | Arizona State University, USA             | Antonia Vecchio        | Consiglio Nazionale delle Ricerche, Italy |
| Charles R. Johnson   | College of William and Mary, USA          | Ram U. Verma           | University of Toledo, USA                 |
| K. B. Kulasekera     | Clemson University, USA                   | John C. Wierman        | Johns Hopkins University, USA             |
| Gerry Ladas          | University of Rhode Island, USA           | Michael E. Zieve       | University of Michigan, USA               |

### PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# involve

2017

vol. 10

no. 4

|   |     |
|---|-----|
| New algorithms for modular inversion and representation by the form<br>$x^2 + 3xy + y^2$  | 541 |
| CHRISTINA DORAN, SHEN LU AND BARRY R. SMITH   |     |
| New approximations for the area of the Mandelbrot set   | 555 |
| DANIEL BITTNER, LONG CHEONG, DANTE GATES AND HIEU D.<br>NGUYEN  |     |
| Bases for the global Weyl modules of $\mathfrak{sl}_n$ of highest weight $m\omega_1$  | 573 |
| SAMUEL CHAMBERLIN AND AMANDA CROAN  |     |
| Leverage centrality of knight's graphs and Cartesian products of regular<br>graphs and path powers  | 583 |
| ROGER VARGAS, JR., ABIGAIL WALDRON, ANIKA SHARMA,<br>RIGOBERTO FLÓREZ AND DARREN A. NARAYAN   |     |
| Equivalence classes of $\mathrm{GL}(p, \mathbb{C}) \times \mathrm{GL}(q, \mathbb{C})$ orbits in the flag variety of<br>$\mathfrak{gl}(p+q, \mathbb{C})$ | 593 |
| LETICIA BARCHINI AND NINA WILLIAMS  |     |
| Global sensitivity analysis in a mathematical model of the renal interstitium   | 625 |
| MARIEL BEDELL, CLAIRE YILIN LIN, EMMIE ROMÁN-MELÉNDEZ<br>AND IOANNIS SGOURALIS  |     |
| Sums of squares in quaternion rings   | 651 |
| ANNA COOKE, SPENCER HAMBLEN AND SAM WHITFIELD   |     |
| On the structure of symmetric spaces of semidihedral groups   | 665 |
| JENNIFER SCHAEFER AND KATHRYN SCHLECHTWEG   |     |
| Spectrum of the Laplacian on graphs of radial functions   | 677 |
| RODRIGO MATOS AND FABIO MONTENEGRO  |     |
| A generalization of Eulerian numbers via rook placements  | 691 |
| ESTHER BANAIAN, STEVE BUTLER, CHRISTOPHER COX, JEFFREY<br>DAVIS, JACOB LANDGRAF AND SCARLITTE PONCE   |     |
| The $H$ -linked degree-sum parameter for special graph families   | 707 |
| LYDIA EAST KENNEY AND JEFFREY SCOTT POWELL  |     |