

involve

a journal of mathematics

Interpolation on Gauss hypergeometric functions with an application

Hina Manoj Arora and Swadesh Kumar Sahoo



Interpolation on Gauss hypergeometric functions with an application

Hina Manoj Arora and Swadesh Kumar Sahoo

(Communicated by Kenneth S. Berenhaut)

We use some standard numerical techniques to approximate the hypergeometric function

$${}_2F_1[a, b; c; x] = 1 + \frac{ab}{c}x + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{x^2}{2!} + \dots$$

for a range of parameter triples (a, b, c) on the interval $0 < x < 1$. Some of the familiar hypergeometric functional identities and asymptotic behavior of the hypergeometric function at $x = 1$ play crucial roles in deriving the formula for such approximations. We also focus on error analysis of the numerical approximations leading to monotone properties of quotients of gamma functions in parameter triples (a, b, c) . Finally, an application to continued fractions of Gauss is discussed followed by concluding remarks consisting of recent works on related problems.

1. Introduction and preliminaries

For a complex number z and $c \neq 0, -1, -2, -3, \dots$, the *hypergeometric series* is defined by

$$1 + \sum_{n=1}^{\infty} \frac{(a)_n (b)_n}{(c)_n (1)_n} z^n.$$

Here $(a)_n$ denotes the shifted factorial notation defined, in terms of the gamma function, by

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)} = \begin{cases} a(a+1) \cdots (a+n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0, a \neq 0. \end{cases}$$

Note that the hypergeometric series defines an analytic function, denoted by the symbol ${}_2F_1[a, b; c; z]$, in $|z| < 1$. As quoted in the historical remarks in [Anderson et al. 1997, 1.55, p. 24], the concept of hypergeometric series was first introduced

MSC2010: primary 65D05; secondary 33B15, 33B20, 33C05, 33F05.

Keywords: interpolation, hypergeometric function, gamma function, error estimate.

by J. Wallis in 1656 to refer to a generalization of the geometric series. Less than a century later, Euler extensively studied the analytic properties of the hypergeometric function and found, for instance, its integral representation; see [Anderson et al. 1997, Theorem 1.19(2)]. Gauss made his first contribution to the subject in 1812. Due to the outstanding contribution made by Gauss to the field, the hypergeometric function is also sometimes known as the *Gauss hypergeometric function*. Most elementary functions which are solutions to certain differential equations can be written in terms of the Gauss hypergeometric functions. One can easily verify by using the Frobenius technique that the function ${}_2F_1[a, b; c; z]$ is one of the solutions of the *hypergeometric differential equation* [Andrews et al. 1999; Beals and Wong 2010; Rainville 1960]

$$z(1-z)w'' + (c - (a+b+1)z)w' - abw = 0.$$

We refer to [Rainville 1943; 1960] for Kummer's 24 solutions to the hypergeometric differential equation, and to [Beals and Wong 2010] for related applications. The asymptotic behavior of ${}_2F_1[a, b; c; z]$ near $z = 1$ reveals that

$${}_2F_1[a, b; c; 1] = \frac{\Gamma(c-a-b)\Gamma(c)}{\Gamma(c-a)\Gamma(c-b)} < \infty, \quad \text{valid for } \operatorname{Re}(c-a-b) > 0. \quad (1-1)$$

Interpolating polynomials for elementary real functions such as trigonometric functions, logarithmic functions, exponential functions, etc. have already been derived in undergraduate texts in numerical analysis; see for instance [Atkinson 1978]. These elementary functions are in fact hypergeometric functions with specific parameters a, b, c ; see for instance [Andrews et al. 1999; Rainville 1960]. Most of such polynomial approximations are computed when the functional values at the given boundary points are possible. Hence the asymptotic behavior (1-1) of the hypergeometric function near $z = 1$ motivates us to construct interpolating polynomials for real hypergeometric functions ${}_2F_1[a, b; c; x]$, $a, b, c \in \mathbb{R}$, $c \notin \{0, -1, -2, -3, \dots\}$, of a real variable x using several numerical techniques in the interval $[0, 1]$; however, the interval may be extended to $[-1, 1]$ as the hypergeometric series in x is convergent for $|x| < 1$ and it has a certain asymptotic behavior near -1 as well, with suitable choices of the parameters a, b, c ; see for instance [Rainville 1960, Theorem 26]. More precisely, when we compute an interpolating polynomial $p_n(x)$ of a hypergeometric function ${}_2F_1[a, b; c; x]$ on $[0, 1]$ we take the value ${}_2F_1[a, b; c; 1]$ in the sense that the hypergeometric function defined at $x = 1$ by means of its asymptotic behavior at $x = 1$; see (1-1). Several hypergeometric functional identities also play a crucial role in determining functional values at the interpolating points.

The following lemmas are useful in describing the error analysis for the interpolating polynomials that we obtained in this paper. Our subsequent paper(s) in this series will cover the study of interpolating polynomials using other techniques.

Lemma 1.1 [Anderson et al. 1997, Lemma 1.33(1), p. 13; see also Lemma 1.35(2)]. *If $a, b, c \in (0, \infty)$, then ${}_2F_1[a, b; c; x]$ is strictly increasing on $[0, 1)$. In particular, if $c > a + b$ then for $x \in [0, 1]$ we have*

$${}_2F_1[a, b; c; x] \leq \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}.$$

Lemma 1.2 [Anderson et al. 1997, Lemma 2.16(2), p. 36]. *The gamma function $\Gamma(x)$ is a log-convex function on $(0, \infty)$. In other words, the logarithmic derivative, $\Gamma'(x)/\Gamma(x)$, of the gamma function is increasing on $(0, \infty)$.*

Note that in all the plots in this paper, graphs drawn in blue represent the original functions and graphs drawn in red represent interpolating polynomials.

2. Linear interpolation on ${}_2F_1[a, b; c; x]$

For performing linear interpolation of the function ${}_2F_1[a, b; c; x] = f(x)$, we consider the end points $x_0 = 0$ and $x_1 = 1$ of the interval $[0, 1]$. The functional values at these points are respectively $f(0) = 1$ and $f(1)$, described in (1-1). Hence, the equation of the segment of the straight line joining 0 and 1 is

$$P_l(x) = f(x_0) + \frac{x - x_0}{x_1 - x_0} (f(x_1) - f(x_0)) = \frac{\Gamma(c)\Gamma(c - a - b) - \Gamma(c - a)\Gamma(c - b)}{\Gamma(c - a)\Gamma(c - b)} x + 1,$$

when $c - a - b > 0$ and $c \neq 0, -1, -2, -3, \dots$. The polynomial $P_l(x)$ represents the linear interpolation of ${}_2F_1[a, b; c; x]$ interpolating at 0 and 1.

Using Lemma 1.1, we obtain the following error estimate:

Lemma 2.1. *Let $a, b, c \in (-2, \infty)$ with $c - a - b > 2$. The deviation of the given function $f(x) = {}_2F_1[a, b; c; x]$ from the approximating function $P_l(x)$ for all values of $x \in [0, 1]$ is estimated by*

$$|E_l(f, x)| = |f(x) - P_l(x)| \leq \frac{|a(a + 1)b(b + 1)|}{8} \frac{\Gamma(c)\Gamma(c - a - b - 2)}{\Gamma(c - a)\Gamma(c - b)}.$$

Proof. Maximizing

$$|E_l(f, x)| = \frac{1}{2} x(1 - x) |f''(x)|$$

in $[0, 1]$ yields

$$\frac{1}{8} (1 - 0)^2 \max_{0 \leq x \leq 1} |f''(x)|,$$

where $f(x) = {}_2F_1[a, b; c; x]$. The following well-known derivative formula is useful:

$$\frac{d}{dx} {}_2F_1[a, b; c; x] = \frac{ab}{c} {}_2F_1[a + 1, b + 1; c + 1; x]. \tag{2-1}$$

The proof follows from (1-1), Lemma 1.1, (2-1), and the fact that

$$\Gamma(x + 1) = x\Gamma(x). \quad \square$$




Figure 1. Linear interpolation of ${}_2F_1[1, 2; 6; x]$ at 0 and 1.

nodes x_i	0	0.25	0.5	0.75	1
actual values ${}_2F_1[1, 2; 6; x_i]$	1	1.0936	1.2149	1.3843	1.6667
polynomial approx. by $P_l(x_i)$	1	1.1667	1.3333	1.5000	1.6667
validity of error bounds by $E_l(f, x_i)$	0	$0.0731 < 1.25$	$0.1184 < 1.25$	$0.1157 < 1.25$	0

Table 1. Comparison of the functional and linear polynomial values.

Remark 2.2. It follows from Lemma 2.1 that there is no error for the choices $a = 0$, $a = -1$, $b = 0$, $b = -1$. In other words, for these choices $E_l(f, x)$ vanishes.

Figure 1 shows linear interpolation of the hypergeometric function at 0 and 1, whereas Table 1 compares the values of the hypergeometric function up to four decimal places with its interpolating polynomial values in the interval $[0, 1]$ for the choice of parameters $a = 1$, $b = 2$ and $c = 6$. Figure 1 and Table 1 also indicate errors at various points within the unit interval except at the end points.

3. Quadratic interpolation on ${}_2F_1[a, b; c; x]$

Let the three points in consideration for quadratic interpolation be $x_0 = 0$, $x_1 = 0.5$ and $x_2 = 1$. The functional values at $x_0 = 0$ and $x_2 = 1$ can be found easily in terms of the parameters but the functional value at $x_1 = 0.5$ can be obtained through different identities involving hypergeometric functions ${}_2F_1[a, b; c; x]$ dealing with various constraints on the parameters a, b, c . This section consists of two subsections and in each subsection the method to obtain the functional value of ${}_2F_1[a, b; c; x]$ at $x = 0.5$ uses three different identities. Finally, we compare the resultant interpolations. In fact we observe that the interpolating polynomial remains unchanged in two cases, although the approaches are different (see the subsection on page 630 for more details).

Quadratic interpolation on ${}_2F_1[a, 1 - a; c; x]$. This section deals with the value ${}_2F_1[a, b; c; \frac{1}{2}]$, where $a + b = 1$ due to the following identity of Bailey [1935,

p. 11] (see also [Rainville 1960, p. 69]):

$$\begin{aligned}
 & {}_2F_1\left[a, 1-a; c; \frac{1}{2}\right] \\
 &= \frac{2^{1-c} \Gamma(c) \Gamma\left(\frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}(c+a)\right) \Gamma\left(\frac{1}{2}(1+c-a)\right)} = \frac{\Gamma\left(\frac{1}{2}c\right) \Gamma\left(\frac{1}{2}(1+c)\right)}{\Gamma\left(\frac{1}{2}(c+a)\right) \Gamma\left(\frac{1}{2}(1+c-a)\right)}, \quad (3-1)
 \end{aligned}$$

where c is a positive integer. It follows from (3-1) that

$$\Gamma\left(\frac{1}{2}c\right)\Gamma\left(\frac{1}{2}(1+c)\right) = 2^{1-c}\sqrt{\pi}\Gamma(c), \quad (3-2)$$

since $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$. In this case, we obtain

$$\begin{aligned}
 f(x_0) &= f(0) = {}_2F_1[a, 1-a; c; 0] = 1, \\
 f(x_1) &= f(0.5) = {}_2F_1\left[a, 1-a; c; \frac{1}{2}\right] = \frac{\Gamma\left(\frac{1}{2}c\right) \Gamma\left(\frac{1}{2}(1+c)\right)}{\Gamma\left(\frac{1}{2}(c+a)\right) \Gamma\left(\frac{1}{2}(1+c-a)\right)}, \\
 f(x_2) &= f(1) = {}_2F_1[a, 1-a; c; 1] = \frac{\Gamma(c)\Gamma(c-1)}{\Gamma(c-a)\Gamma(c+a-1)} \quad (c > 1).
 \end{aligned}$$

Consider the well-known Lagrange fundamental polynomials

$$\begin{aligned}
 L_0(x) &= \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)}, \\
 L_1(x) &= \frac{(x-x_0)(x-x_2)}{(x_1-x_0)(x_1-x_2)}, \\
 L_2(x) &= \frac{(x-x_0)(x-x_1)}{(x_2-x_0)(x_2-x_1)}.
 \end{aligned}$$

Then the quadratic interpolation of $f(x) = {}_2F_1[a, 1-a; c; x]$ becomes

$$\begin{aligned}
 P_{q_3}(x) &= f(x_0)L_0(x) + f(x_1)L_1(x) + f(x_2)L_2(x) \\
 &= (2x^2 - 3x + 1) + (-4x^2 + 4x) \frac{\Gamma\left(\frac{1}{2}c\right) \Gamma\left(\frac{1}{2}(1+c)\right)}{\Gamma\left(\frac{1}{2}(c+a)\right) \Gamma\left(\frac{1}{2}(1+c-a)\right)} \\
 &\quad + (2x^2 - x) \frac{\Gamma(c)\Gamma(c-1)}{\Gamma(c-a)\Gamma(c+a-1)}.
 \end{aligned}$$

This leads to the following result.

Theorem 3.1. *Let $a, b, c \in \mathbb{R}$ be such that $c > 1$. Then*

$$\begin{aligned}
 P_{q_1}(x) &= \left(2 - \frac{4\Gamma\left(\frac{1}{2}c\right)\Gamma\left(\frac{1}{2}(1+c)\right)}{\Gamma\left(\frac{1}{2}(c+a)\right)\Gamma\left(\frac{1}{2}(1+c-a)\right)} + \frac{2\Gamma(c)\Gamma(c-1)}{\Gamma(c-a)\Gamma(c+a-1)}\right)x^2 \\
 &\quad + \left(\frac{4\Gamma\left(\frac{1}{2}c\right)\Gamma\left(\frac{1}{2}(1+c)\right)}{\Gamma\left(\frac{1}{2}(c+a)\right)\Gamma\left(\frac{1}{2}(1+c-a)\right)} - \frac{\Gamma(c)\Gamma(c-1)}{\Gamma(c-a)\Gamma(c+a-1)} - 3\right)x + 1.
 \end{aligned}$$

is a quadratic interpolation of ${}_2F_1[a, 1-a; c; x]$ in $[0, 1]$.




Figure 2. The quadratic interpolation of ${}_2F_1[0.9, 0.1; 1.5; x]$ at 0, 0.5, and 1.

Remark 3.2. It is evident that when $a = 0, 1$, then $P_{q_1}(x) = {}_2F_1[a, 1 - a; c; x] = 1$ for all $x \in [0, 1]$ and for all $c > 1$. Moreover, for all $c > 1$, we have the following three natural observations:

- (i) If $-1 < a < 0$, then $P_{q_1}(x)$ and ${}_2F_1[a, 1 - a; c; x]$ decrease together in $[0, 1]$.
- (ii) If $0 < a < 1$, then $P_{q_1}(x)$ and ${}_2F_1[a, 1 - a; c; x]$ increase together in $[0, 1]$.
- (iii) If $1 < a < 2$, then $P_{q_1}(x)$ and ${}_2F_1[a, 1 - a; c; x]$ decrease together in $[0, 1]$.

Indeed, these follow from derivative test. More observations are stated later while estimating the error (see Remark 3.10).

An interpolating polynomial $P_{q_1}(x)$ of ${}_2F_1[a, 1 - a; c; x]$ for certain choices of parameters a and c is as shown in Figure 2.

Remark 3.3. Note that in Theorem 3.1, the parameter c cannot be chosen such that $c \leq \frac{1}{2}(a + b + 1)$ since the choice $b = 1 - a$ results in $c \leq 1$, which is a contradiction to the assumption that $c > 1$. In particular, $c \neq \frac{1}{2}(a + b + 1)$ in Theorem 3.1, which is the negation of a constraint that will be considered in the next subsection.

Quadratic interpolation on ${}_2F_1[a, b; \frac{1}{2}(a + b + 1); x]$. In this section, $f(x) = {}_2F_1[a, b; c; x]$, $c = \frac{1}{2}(a + b + 1)$, is first interpolated using the following quadratic transformation obtained from [Andrews et al. 1999, (3.1.3)]; see also [Rainville 1960, Theorem 2.5].

Lemma 3.4. *If $\frac{1}{2}(a + b + 1)$ is a positive integer, and if $|x| < 1$ and $|4x(1 - x)| < 1$, then*

$${}_2F_1[a, b; \frac{1}{2}(a + b + 1); x] = {}_2F_1[\frac{1}{2}a, \frac{1}{2}b; \frac{1}{2}(a + b + 1); 4x(1 - x)]. \tag{3-3}$$

If we choose $x = 0.5$ then the right-hand side of (3-3) computes the asymptotic behavior of the hypergeometric function at 1. Hence the functional value at $x = 0.5$ of the function $f(x) = {}_2F_1[a, b; \frac{1}{2}(a + b + 1); x]$ can be obtained with the help of (1-1). Due to Lemma 3.4 and (1-1), in this case, the constraints on the parameters are computed as

- $a + b < 1$;
- $a + b \neq -(2n + 1)$ for $n \in \mathbb{N} \cup \{0\}$.

One can easily obtain that

$$f(x_0) = {}_2F_1\left[a, b; \frac{1}{2}(a+b+1); 0\right] = 1;$$

$$f(x_1) = {}_2F_1\left[a, b; \frac{1}{2}(a+b+1); \frac{1}{2}\right] = \frac{\sqrt{\pi} \Gamma\left(\frac{1}{2}(a+b+1)\right)}{\Gamma\left(\frac{1}{2}(a+1)\right)\Gamma\left(\frac{1}{2}(b+1)\right)},$$

$$f(x_2) = {}_2F_1\left[a, b; \frac{1}{2}(a+b+1); 1\right] = \frac{\Gamma\left(\frac{1}{2}(1-a-b)\right)\Gamma\left(\frac{1}{2}(a+b+1)\right)}{\Gamma\left(\frac{1}{2}(a+1-b)\right)\Gamma\left(\frac{1}{2}(b+1-a)\right)} = \frac{\cos\left(\frac{\pi}{2}(b-a)\right)}{\cos\left(\frac{\pi}{2}(b+a)\right)},$$

where $f(x_2)$ is obtained by the well-known Euler’s reflection formula (in nonintegral variable x)

$$\Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin(\pi x)}.$$

This leads to the additional constraints on the parameters

$$\begin{aligned} a + b \neq 1 \pm 2n \quad \text{and} \quad a - b \neq -1 \pm 2n, \quad n \in \mathbb{Z} \quad \text{or} \\ a + b \neq -1 \pm 2n \quad \text{and} \quad a - b \neq 1 \pm 2n, \quad n \in \mathbb{Z}. \end{aligned} \tag{3-4}$$

(These constraints may be relaxed when one does not use Euler’s reflection formula!)

Thus, the first quadratic interpolation of $f(x) = {}_2F_1\left[a, b; \frac{1}{2}(a+b+1); x\right]$ becomes

$$\begin{aligned} P_{q_2}(x) &= f(x_0)L_0(x) + f(x_1)L_1(x) + f(x_2)L_2(x) \\ &= (2x^2 - 3x + 1) + (-4x^2 + 4x) \frac{\sqrt{\pi} \Gamma\left(\frac{1}{2}(a+b+1)\right)}{\Gamma\left(\frac{1}{2}(a+1)\right)\Gamma\left(\frac{1}{2}(b+1)\right)} \\ &\quad + (2x^2 - x) \frac{\cos\left(\frac{\pi}{2}(b-a)\right)}{\cos\left(\frac{\pi}{2}(b+a)\right)}. \end{aligned}$$

This leads to the following result.

Theorem 3.5. *Let $a, b \in \mathbb{R}$ and $n \in \mathbb{N} \cup \{0\}$ be such that $a + b \neq -(2n + 1)$ and $a + b < 1$. If either $a + b \neq 1 \pm 2n$ and $a - b \neq -1 \pm 2n$, or $a + b \neq -1 \pm 2n$ and $a - b \neq 1 \pm 2n$ hold, then*

$$\begin{aligned} P_{q_2}(x) &= \left(2 - \frac{4\sqrt{\pi} \Gamma\left(\frac{1}{2}(a+b+1)\right)}{\Gamma\left(\frac{1}{2}(a+1)\right)\Gamma\left(\frac{1}{2}(b+1)\right)} + \frac{2 \cos\left(\frac{\pi}{2}(b-a)\right)}{\cos\left(\frac{\pi}{2}(b+a)\right)} \right) x^2 \\ &\quad + \left(\frac{4\sqrt{\pi} \Gamma\left(\frac{1}{2}(a+b+1)\right)}{\Gamma\left(\frac{1}{2}(a+1)\right)\Gamma\left(\frac{1}{2}(b+1)\right)} - \frac{\cos\left(\frac{\pi}{2}(b-a)\right)}{\cos\left(\frac{\pi}{2}(b+a)\right)} - 3 \right) x + 1 \end{aligned}$$

is a quadratic interpolation of ${}_2F_1\left[a, b; \frac{1}{2}(a+b+1); x\right]$ in $[0, 1]$.

Secondly, we discuss quadratic interpolation of the same function ${}_2F_1[a, b; c; x]$, $c = \frac{1}{2}(a + b + 1)$, in $[0, 1]$, but using a different hypergeometric identity. Finally, we observe that both the interpolations are same except at a minor difference in one of the constraints.

Recall the transformation formula [Rainville 1960, Theorem 20, p. 60]:

Lemma 3.6. *If $|x| < 1$ and $|x/(1 - x)| < 1$, then we have*

$${}_2F_1[a, b; c; x] = (1 - x)^{-a} {}_2F_1\left[a, c - b; c; \frac{-x}{1 - x}\right].$$

Note that $-x/(1 - x) = -1$ for $x = 0.5$. This suggests that to find the value $f(0.5) = 2^a {}_2F_1[a, c - b; c; -1]$ we can use the following identity [Rainville 1960, Theorem 26, p. 68]; see also [Beals and Wong 2010].

Lemma 3.7. *Let $a', b' \in \mathbb{R}$. If $1 + a' - b' \neq \{0, -1, -2, -3, \dots\}$ and $b' < 1$, then we have*

$${}_2F_1[a', b'; a' - b' + 1; -1] = \frac{\Gamma(a' - b' + 1)\Gamma(\frac{1}{2}a' + 1)}{\Gamma(a' + 1)\Gamma(\frac{1}{2}a' - b' + 1)}.$$

Comparison of the parameters $a' = a$, $b' = c - b$ and $a' - b' + 1 = c$ leads to

$${}_2F_1[a, c - b; c; -1] = \frac{\Gamma(a - c + b + 1)\Gamma(\frac{1}{2}a + 1)}{\Gamma(a + 1)\Gamma(\frac{1}{2}a - c + b + 1)} \tag{3-5}$$

with the constraints

- $2c = a + b + 1$;
- $c \neq \{0, -1, -2, -3, \dots\} \iff a + b \neq -(2n + 1), n \in \mathbb{N} \cup \{0\}$;
- $c - b < 1 \iff a - b < 1$.

Under these conditions, (3-5) leads to

$$\begin{aligned} f(x_1) = f(0.5) &= {}_2F_1\left[a, b; \frac{1}{2}(a + b + 1); \frac{1}{2}\right] = 2^a \frac{\Gamma(\frac{1}{2}(a + b + 1))\Gamma(\frac{1}{2}a + 1)}{\Gamma(a + 1)\Gamma(\frac{1}{2}(b + 1))} \\ &= \frac{2^{a-1} \Gamma(\frac{1}{2}(a + b + 1)) \Gamma(\frac{1}{2}a)}{\Gamma(a) \Gamma(\frac{1}{2}(b + 1))} = \frac{\sqrt{\pi} \Gamma(\frac{1}{2}(a + b + 1))}{\Gamma(\frac{1}{2}(a + 1))\Gamma(\frac{1}{2}(b + 1))}, \end{aligned}$$

where the last equality holds by (3-2). Also as discussed at the beginning of this subsection, we have

$$\begin{aligned} f(x_0) = f(0) &= {}_2F_1\left[a, b; \frac{1}{2}(a + b + 1); 0\right] = 1, \\ f(x_2) = f(1) &= {}_2F_1\left[a, b; \frac{1}{2}(a + b + 1); 1\right] = \frac{\cos(\frac{\pi}{2}(b - a))}{\cos(\frac{\pi}{2}(b + a))}, \quad a + b < 1, \end{aligned}$$

with additional constraints obtained in (3-4) (here also (3-4) may be relaxed!).




Figure 3. The quadratic interpolation of ${}_2F_1[0.1, 0.3; 0.7; x]$ at 0, 0.5, and 1.

Thus, the second quadratic interpolation of $f(x) = {}_2F_1[a, b; \frac{1}{2}(a + b + 1); x]$ remains same as the first quadratic interpolation obtained in [Theorem 3.5](#) but with an additional constraint $a - b < 1$. This shows that the quadratic interpolation obtained by [Theorem 3.5](#) is stronger than what was discussed so far using [Lemmas 3.6](#) and [3.7](#). A quadratic interpolation of ${}_2F_1[a, b; \frac{1}{2}(a + b + 1); x]$ is shown in [Figure 3](#).

Error estimates. The error estimate in quadratic interpolation of ${}_2F_1[a, b; c; x]$ interpolating at 0, 0.5, 1 in $[0, 1]$ is formulated as below:

Lemma 3.8. Let $P_q(x)$ be a quadratic interpolation of $f(x) = {}_2F_1[a, b; c; x]$ interpolating at 0, 0.5, 1 in $[0, 1]$. If $a, b, c \in (-3, \infty)$ with $c - a - b > 3$, then the deviation of $f(x)$ from $P_q(x)$ is estimated by

$$|E_q(f, x)| = |f(x) - P_q(x)| \leq \frac{1}{6}M |a(a + 1)(a + 2)b(b + 1)(b + 2)| \frac{\Gamma(c)\Gamma(c - a - b - 3)}{\Gamma(c - a)\Gamma(c - b)}$$

for all values of $x \in [0, 1]$, where M is defined by

$$M := \begin{cases} \frac{1}{12}(3 - \sqrt{3})(-1 + \frac{1}{6}(3 - \sqrt{3}))(-1 + \frac{1}{3}(3 - \sqrt{3})), & x < \frac{1}{2}, \\ -\frac{1}{12}(3 + \sqrt{3})(-1 + \frac{1}{6}(3 + \sqrt{3}))(-1 + \frac{1}{3}(3 + \sqrt{3})), & x > \frac{1}{2}. \end{cases} \tag{3-6}$$

Proof. We need to estimate

$$\max_{0 \leq x \leq 1} \frac{1}{6} |x(x - 0.5)(x - 1)| \max_{0 \leq x \leq 1} |f'''(x)|,$$

where $f(x) = {}_2F_1[a, b; c; x]$. Note that

$$\max_{0 \leq x \leq 1} |x(x - 0.5)(x - 1)| = M (\approx 0.0481125 \dots)$$

by (3-6). We apply the well known derivative formula (2-1) to maximize $|f'''(x)|$, $0 \leq x \leq 1$. The proof follows from (1-1), [Lemma 1.1](#), (2-1), and the fact that

$$\Gamma(x + 1) = x\Gamma(x). \quad \square$$

The following result is an immediate consequence of [Lemma 3.8](#) which estimates the difference $E_{q_1}(f, x) = {}_2F_1[a, 1 - a; c; x] - P_{q_1}(x)$ in $[0, 1]$.

Corollary 3.9. *Let $a, c \in \mathbb{R}$ be such that $-3 < a < 4$ and $c > 4$. Then the deviation of ${}_2F_1[a, 1 - a; c; x]$ from $P_{q_1}(x)$ is estimated by*

$$|E_{q_1}(f, x)| = |f(x) - P_{q_1}(x)| \leq \frac{1}{6}M |a(a+1)(a+2)(1-a)(2-a)(3-a)| \frac{\Gamma(c)\Gamma(c-4)}{\Gamma(c-a)\Gamma(c+a-1)}$$

for all values of $x \in [0, 1]$, where M is obtained by [\(3-6\)](#).

Remark 3.10. It follows from [Corollary 3.9](#) that there is no error for any of the choices $a = -2, -1, 0, 1, 2, 3$. In other words, for any of these choices, $E_{q_1}(f, x)$ vanishes.

Similarly, as a consequence of [Lemma 3.8](#), we obtain:

Corollary 3.11. *Let $a, b \in \mathbb{R}$ be such that $-7 < a + b < -5$. Then the deviation of ${}_2F_1[a, b; \frac{1}{2}(a+b+1); x]$ from $P_{q_2}(x)$ is estimated by*

$$|E_{q_2}(f, x)| = |f(x) - P_{q_2}(x)| \leq \frac{1}{6}M |a(a+1)(a+2)b(b+1)(b+2)| \frac{\Gamma(\frac{1}{2}(a+b+1))\Gamma(\frac{1}{2}(-a-b-5))}{\Gamma(\frac{1}{2}(b-a+1))\Gamma(\frac{1}{2}(a-b+1))}$$

for all values of $x \in [0, 1]$, where M is obtained by [\(3-6\)](#).

Remark 3.12. It follows from [Corollary 3.11](#) that since $E_{q_2}(f, x)$ vanishes for the choices $a = -2, -1, 0$ and $b = -2, -1, 0$, there is no error for these choices of the parameters a and b .

Now we describe a slightly deeper analysis on the error obtained in [Corollary 3.9](#) through the following lemma, which is a consequence of [Lemma 1.2](#). A similar analysis can be described for [Corollary 3.11](#).

Lemma 3.13. *Let $a, c \in \mathbb{R}$ be such that $c > 4$. If either $1 < a < 4$ or $-3 < a < 0$ holds, then the quotient*

$$\frac{\Gamma(c)\Gamma(c-4)}{\Gamma(c-a)\Gamma(c+a-1)}$$

decreases when c increases.

Proof. We use [Lemma 1.2](#). Since $c - a > c - 4 > 0$, on one hand we have

$$\frac{\Gamma'(c-4)}{\Gamma(c-4)} - \frac{\Gamma'(c-a)}{\Gamma(c-a)} < 0.$$

On the other hand, since $c < c + a - 1$, we have

$$\frac{\Gamma'(c)}{\Gamma(c)} - \frac{\Gamma'(c+a-1)}{\Gamma(c+a-1)} < 0.$$

Thus, if

$$g(c) = \frac{\Gamma(c)\Gamma(c-4)}{\Gamma(c-a)\Gamma(c+a-1)},$$

it follows that

$$\begin{aligned} \frac{g'(c)}{g(c)} &= \frac{\Gamma'(c)}{\Gamma(c)} + \frac{\Gamma'(c-4)}{\Gamma(c-4)} - \frac{\Gamma'(c-a)}{\Gamma(c-a)} - \frac{\Gamma'(c+a-1)}{\Gamma(c+a-1)} \\ &= \left(\frac{\Gamma'(c-4)}{\Gamma(c-4)} - \frac{\Gamma'(c-a)}{\Gamma(c-a)} \right) + \left(\frac{\Gamma'(c)}{\Gamma(c)} - \frac{\Gamma'(c+a-1)}{\Gamma(c+a-1)} \right) < 0. \end{aligned}$$

By the definition of the gamma function, obviously, one can see that $\Gamma(x) > 0$ for $x > 0$. This shows that $g(c) > 0$ and hence $g'(c) < 0$. Thus, $g(c)$ decreases for $1 < a < 4 < c$.

For $c > 4$, if $-3 < a < 0$ holds then we consider the rearrangement

$$\frac{g'(c)}{g(c)} = \left(\frac{\Gamma'(c)}{\Gamma(c)} - \frac{\Gamma'(c-a)}{\Gamma(c-a)} \right) + \left(\frac{\Gamma'(c-4)}{\Gamma(c-4)} - \frac{\Gamma'(c+a-1)}{\Gamma(c+a-1)} \right)$$

and show that $g'(c)/g(c) < 0$. □

Using Mathematica or other similar tools, one can see that [Lemma 3.13](#) even holds true for the remaining range $0 \leq a \leq 1$. This suggests the following conjecture.

Conjecture 3.14. *Let $a, c \in \mathbb{R}$ be such that $0 \leq a \leq 1$ and $c > 4$. Then the quotient*

$$\frac{\Gamma(c)\Gamma(c-4)}{\Gamma(c-a)\Gamma(c+a-1)}$$

decreases when c increases.

Thus, we observe that when $c > 4$ increases then the error $E_{q_1}(f, x)$ estimated in [Corollary 3.9](#) decreases (see also [Figures 4](#) and [5](#)).

[Figures 4](#) and [5](#) describe the quadratic interpolation of the hypergeometric functions ${}_2F_1[a, 1-a, c, x]$ at 0, 0.5 and 1, whereas [Tables 2](#) and [3](#) compare the values of the hypergeometric function up to four decimal places with its interpolating




Figure 4. The error estimate $E_{q_1}(f, x)$ when $a = 3.9$ and c increases from 4.5 to 6.5.




Figure 5. The error estimate $E_{q_1}(f, x)$ when $a = 0.9$ and c increases from 4.1 to 6.1.

nodes x_i	0	0.25	0.5	0.75	1
actual values ${}_2F_1[3.9, -2.9; 5; x_i]$	1	0.5372	0.2516	0.0998	0.0367
polynomial approx. by $P_{q_1}(x_i)$	1	0.5591	0.2516	0.0775	0.0367
validity of error bounds by $E_{q_1}(f, x_i)$	0	$0.0219 < 0.0274$	0	$0.0223 < 0.0274$	0
actual values ${}_2F_1[3.9, -2.9; 6; x_i]$	1	0.6027	0.3358	0.1724	0.0845
polynomial approx. by $P_{q_1}(x_i)$	1	0.6163	0.3358	0.1585	0.0845
validity of error bounds by $E_{q_1}(f, x_i)$	0	$0.0136 < 0.0158$	0	$0.0139 < 0.0158$	0

Table 2. Comparison of the functional and quadratic polynomial values.

nodes x_i	0	0.25	0.5	0.75	1
actual values ${}_2F_1[0.9, 0.1; 5; x_i]$	1	1.0047	1.0099	1.0158	1.0227
polynomial approx. by $P_{q_1}(x_i)$	1	1.0046	1.0099	1.0160	1.0227
validity of error bounds by $E_{q_1}(f, x_i)$	0	$0.0001 < 0.0016$	0	$0.0002 < 0.0016$	0
actual values ${}_2F_1[0.9, 0.1; 6; x_i]$	1	1.0039	1.0082	1.0128	1.0182
polynomial approx. by $P_{q_1}(x_i)$	1	1.0038	1.0082	1.0129	1.0182
validity of error bounds by $E_{q_1}(f, x_i)$	0	$0.0001 < 0.0004$	0	$0.0001 < 0.0004$	0

Table 3. Comparison of the functional and quadratic polynomial values.

polynomial values in the interval $[0, 1]$ for the choice of parameters $a = 3.9, c = 5$ and $a = 0.9, c = 6$ respectively. Figures 4 and 5 and Tables 2 and 3 also indicate errors at various points within the unit interval except at the interpolating points at $x = 0, 0.5, 1$.

The error estimate $|E_{q_2}(f, x)|$ for the function ${}_2F_1[a, b; \frac{1}{2}(a + b + 1); x]$ can be analyzed in a similar way, and hence we omit the proof.

4. An application

In this section, we briefly consider interpolation of a continued fraction that converges to a quotient of two hypergeometric functions. Gauss used the contiguous relations to give several ways to write a quotient of two hypergeometric functions as a continued fraction. For instance, it is well known that

$$\frac{{}_2F_1[a + 1, b; c + 1; x]}{{}_2F_1[a, b; c; x]} = \frac{1}{1 + \frac{\frac{(a-c)b}{c(c+1)}x}{1 + \frac{\frac{(b-c-1)(a+1)}{(c+1)(c+2)}x}{1 + \frac{\frac{(a-c-1)(b+1)}{(c+2)(c+3)}x}{1 + \frac{\frac{(b-c-2)(a+2)}{(c+3)(c+4)}x}{1 + \dots}}}}, \quad |x| < 1. \quad (4-1)$$

On one hand, if we adopt the basic linear interpolation method that we discussed in Section 2 (that is, linear interpolation directly) to the function

$$g(x) = \frac{{}_2F_1[a + 1, b; c + 1; x]}{{}_2F_1[a, b; c; x]}$$

at $x_0 = 0$ and $x_1 = 1$, we obtain the linear interpolation of the above continued fraction in the form

$$R_l(x) = g(x_0) + \frac{x - x_0}{x_1 - x_0}(g(x_1) - g(x_0)) = 1 + \left(\frac{b}{c - b}\right)x, \quad c - b > a,$$

since $g(x_0) = 1$ and $g(x_1) = c/(c - b)$. For the choice $a = 1, b = 2, c = 6$, this approximation is also shown in Figure 6.

On the other hand, an application of linear interpolation of ${}_2F_1[a, b; c; x]$ obtained in Section 2 leads to the following approximation of the above continued




Figure 6. Approximation of ${}_2F_1[a+1, b; c+1; x]/{}_2F_1[a, b; c; x]$ through $R_l(x)$.




Figure 7. Approximation of ${}_2F_1[a+1, b; c+1; x]/{}_2F_1[a, b; c; x]$ through $R_r(x)$.

fraction in terms of ratio of polynomial approximation (we call this *rational interpolation*):

$$\begin{aligned}
 R_r(x) &= \frac{1}{P_l(x)} \left(\frac{\Gamma(c+1)\Gamma(c-a-b) - \Gamma(c-a)\Gamma(c-b+1)}{\Gamma(c-a)\Gamma(c-b+1)} x + 1 \right) \\
 &= \frac{[c\Gamma(c)\Gamma(c-a-b)/(c-b) - \Gamma(c-a)\Gamma(c-b)]x + \Gamma(c-a)\Gamma(c-b)}{[\Gamma(c)\Gamma(c-a-b) - \Gamma(c-a)\Gamma(c-b)]x + \Gamma(c-a)\Gamma(c-b)} \\
 &= 1 + \frac{b}{c-b} \left[\frac{\Gamma(c-a-b)\Gamma(c)x}{[\Gamma(c)\Gamma(c-a-b) - \Gamma(c-a)\Gamma(c-b)]x + \Gamma(c-a)\Gamma(c-b)} \right],
 \end{aligned}$$

where $c - a - b > 0$. For the choice $a = 1, b = 2, c = 6$, this approximation is also shown in [Figure 7](#).

Observe that

$$R_r(x_0) = 1 = R_l(x_0) \quad \text{and} \quad R_r(x_1) = \frac{c}{c-b} = R_l(x_1)$$

and hence R_r also interpolates the continued fraction under consideration at 0 and 1. Further we observe that both the approximations $R_l(x)$ and $R_r(x)$ of the continued

fraction are easy to obtain and the first approximation, i.e., $R_l(x)$, is in a simpler form than $R_r(x)$, as expected. Now, it would be interesting to know which one would give the best approximation to the continued fraction under consideration. With the special choice $a = 1, b = 2, c = 6$, we see from Figures 6 and 7 that $R_l(x)$ is a better approximation than $R_r(x)$. One may ask: does it happen for arbitrary parameters a, b, c ? Since $R_l(x) = R_r(x)$ if and only if $\Gamma(c)\Gamma(c - a - b) = \Gamma(c - a)\Gamma(c - b)$, the answer to this question is yes except when $\Gamma(c)\Gamma(c - a - b) = \Gamma(c - a)\Gamma(c - b)$.

This leads to the following result:

Theorem 4.1. *Let $R_l(x)$ and $R_r(x)$ be respectively the linear interpolation and the rational interpolation of the quotient ${}_2F_1[a + 1, b; c + 1; x]/{}_2F_1[a, b; c; x]$ (equivalently, of the continued fraction (4-1)). Then $R_l(x)$ and $R_r(x)$ coincide with each other if and only if $\Gamma(c)\Gamma(c - a - b) = \Gamma(c - a)\Gamma(c - b)$ holds for $c - a - b > 0$.*

5. Concluding remarks and future scope

Recall that, in this paper, we use some standard interpolation techniques to approximate the hypergeometric function

$${}_2F_1[a, b; c; x] = 1 + \frac{ab}{c}x + \frac{a(a + 1)b(b + 1)}{c(c + 1)} \frac{x^2}{2!} + \dots$$

for a range of parameter triples (a, b, c) on the interval $0 < x < 1$. Some of the familiar hypergeometric functional identities and asymptotic behavior of the hypergeometric function at $x = 1$ played crucial roles in deriving the formula for such approximations. One can expect similar formulae using other well-known interpolations and obtain better approximations for the hypergeometric function; however, we discuss such results in an upcoming manuscript(s). Different numerical methods for the computation of the confluent and Gauss hypergeometric functions were studied recently in [Pearson et al. 2017]. Such investigation may be extended to the q -analog of the hypergeometric functions, namely, Heine’s basic hypergeometric functions; for instance refer to [Chen and Fu 2011] for similar discussions.

We also focus on error analysis of the numerical approximations leading to monotone properties of quotients of gamma functions in parameter triples (a, b, c) . Monotone properties of the gamma function and its quotients in different forms are of recent interest to many researchers; see for instance [Alzer 1993; Anderson and Qiu 1997; Bustoz and Ismail 1986; Chen and Zhou 2014; Giordano and Laforgia 2001; Gautschi 1959; Luo et al. 2017; Mortici and Dumitrescu 2017]. In this paper, we also studied and stated a conjecture (see Conjecture 3.14) related to monotone properties of quotients of gamma functions to analyze the error estimate of the numerical approximations under consideration.

Finally, an application to continued fractions of Gauss is also discussed. Approximations of continued fractions in different forms are also attractive to many researchers; see [Lu et al. 2017; 2016].

Acknowledgements

This work was carried out when Arora was in internship at IIT Indore during the summer of 2014. The authors would like to thank the referee and the editor for their valuable remarks on this paper.

References

- [Alzer 1993] H. Alzer, “Some gamma function inequalities”, *Math. Comp.* **60**:201 (1993), 337–346. [MR](#) [Zbl](#)
- [Anderson and Qiu 1997] G. D. Anderson and S.-L. Qiu, “A monotoneity property of the gamma function”, *Proc. Amer. Math. Soc.* **125**:11 (1997), 3355–3362. [MR](#) [Zbl](#)
- [Anderson et al. 1997] G. D. Anderson, M. K. Vamanamurthy, and M. K. Vuorinen, *Conformal invariants, inequalities, and quasiconformal maps*, Wiley, New York, 1997. [MR](#) [Zbl](#)
- [Andrews et al. 1999] G. E. Andrews, R. Askey, and R. Roy, *Special functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge Univ. Press, 1999. [MR](#) [Zbl](#)
- [Atkinson 1978] K. E. Atkinson, *An introduction to numerical analysis*, Wiley, New York, 1978. [MR](#) [Zbl](#)
- [Bailey 1935] W. N. Bailey, *Generalized hypergeometric series*, Cambridge Tracts in Math. and Math. Phys. **32**, Cambridge Univ. Press, 1935. [Zbl](#)
- [Beals and Wong 2010] R. Beals and R. Wong, *Special functions*, Cambridge Studies in Advanced Mathematics **126**, Cambridge Univ. Press, 2010. [MR](#) [Zbl](#)
- [Bustoz and Ismail 1986] J. Bustoz and M. E. H. Ismail, “On gamma function inequalities”, *Math. Comp.* **47**:176 (1986), 659–667. [MR](#) [Zbl](#)
- [Chen and Fu 2011] S. H. L. Chen and A. M. Fu, “A $2n$ -point interpolation formula with its applications to q -identities”, *Discrete Math.* **311**:16 (2011), 1793–1802. [MR](#) [Zbl](#)
- [Chen and Zhou 2014] B. Chen and H. Zhou, “On completely monotone of an arbitrary real parameter function involving the gamma function”, *Appl. Math. Comput.* **242** (2014), 658–663. [MR](#) [Zbl](#)
- [Gautschi 1959] W. Gautschi, “Some elementary inequalities relating to the gamma and incomplete gamma function”, *J. Math. and Phys.* **38**:1-4 (1959), 77–81. [MR](#) [Zbl](#)
- [Giordano and Laforgia 2001] C. Giordano and A. Laforgia, “Inequalities and monotonicity properties for the gamma function”, *J. Comput. Appl. Math.* **133**:1-2 (2001), 387–396. [MR](#) [Zbl](#)
- [Lu et al. 2016] D. Lu, L. Song, and C. Ma, “A quicker continued fraction approximation of the gamma function related to the Windschitl’s formula”, *Numer. Algorithms* **72**:4 (2016), 865–874. [MR](#) [Zbl](#)
- [Lu et al. 2017] D. Lu, X. Liu, and T. Qu, “Continued fraction approximations and inequalities for the gamma function by Burnside”, *Ramanujan J.* **42**:2 (2017), 491–500. [MR](#) [Zbl](#)
- [Luo et al. 2017] S. Luo, J. Wei, and W. Zou, “On a transcendental equation involving quotients of gamma functions”, *Proc. Amer. Math. Soc.* **145**:6 (2017), 2623–2637. [MR](#) [Zbl](#)
- [Mortici and Dumitrescu 2017] C. Mortici and S. Dumitrescu, “Efficient approximations of the gamma function and further properties”, *Comput. Appl. Math.* **36**:1 (2017), 677–691. [MR](#) [Zbl](#)

- [Pearson et al. 2017] J. W. Pearson, S. Olver, and M. A. Porter, “Numerical methods for the computation of the confluent and Gauss hypergeometric functions”, *Numer. Algorithms* **74**:3 (2017), 821–866. [MR](#) [Zbl](#)
- [Rainville 1943] E. D. Rainville, *Intermediate course in differential equations*, Wiley, New York, 1943.
- [Rainville 1960] E. D. Rainville, *Special functions*, Macmillan, New York, 1960. [MR](#) [Zbl](#)

Received: 2016-11-21 Revised: 2017-07-07 Accepted: 2017-07-21

hina.arora.256@gmail.com

*Discipline of Electrical Engineering,
Indian Institute of Technology, Indore, India*

hina.arora@stonybrook.edu

*Department of Applied Mathematics & Statistics,
Stony Brook University, Stony Brook, NY, United States*

swadesh@iiti.ac.in

*Discipline of Mathematics, Indian Institute of Technology,
Indore, India*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2018 is US \$190/year for the electronic version, and \$250/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

involve

2018

vol. 11

no. 4

Modeling of breast cancer through evolutionary game theory	541
KE'YONA BARTON, CORBIN SMITH, JAN RYCHTÁŘ AND TSVETANKA SENDOVA	
The isoperimetric problem in the plane with the sum of two Gaussian densities	549
JOHN BERRY, MATTHEW DANNENBERG, JASON LIANG AND YINGYI ZENG	
Finiteness of homological filling functions	569
JOSHUA W. FLEMING AND EDUARDO MARTÍNEZ-PEDROZA	
Explicit representations of 3-dimensional Sklyanin algebras associated to a point of order 2	585
DANIEL J. REICH AND CHELSEA WALTON	
A classification of Klein links as torus links	609
STEVEN BERES, VESTA COUFAL, KAIA HLAVACEK, M. KATE KEARNEY, RYAN LATTANZI, HAYLEY OLSON, JOEL PEREIRA AND BRYAN STRUB	
Interpolation on Gauss hypergeometric functions with an application	625
HINA MANOJ ARORA AND SWADESH KUMAR SAHOO	
Properties of sets of nontransitive dice with few sides	643
LEVI ANGEL AND MATT DAVIS	
Numerical studies of serendipity and tensor product elements for eigenvalue problems	661
ANDREW GILLETTE, CRAIG GROSS AND KEN PLACKOWSKI	
Connectedness of two-sided group digraphs and graphs	679
PATRECK CHIKWANDA, CATHY KRILOFF, YUN TECK LEE, TAYLOR SANDOW, GARRETT SMITH AND DMYTRO YEROSHKIN	
Nonunique factorization over quotients of PIDs	701
NICHOLAS R. BAETH, BRANDON J. BURNS, JOSHUA M. COVEY AND JAMES R. MIXCO	
Locating trinomial zeros	711
RUSSELL HOWELL AND DAVID KYLE	