

involve

a journal of mathematics

Darboux calculus

Marco Aldi and Alexander McCleary



Darboux calculus

Marco Aldi and Alexander McCleary

(Communicated by Kenneth S. Berenhaut)

We introduce a formalism to analyze partially defined functions between ordered sets. We show that our construction provides a uniform and conceptual approach to all the main definitions encountered in elementary real analysis including Dedekind cuts, limits and continuity.

1. Introduction

Following the pioneering work of Bolzano and Weierstrass, “ (ε, δ) -definitions” are at the heart of textbook presentations of elementary analysis; see, e.g., [Rudin 1953]. While with practice the motivated student quickly becomes proficient in this language, it is natural to ask if fundamental notions such as limits, continuity and integrals could perhaps be defined more conceptually.

In the present paper we develop a rather general framework, which we refer to as *Darboux calculus*, whose specialization to the context of real analysis provides a unified and conceptual approach to all the main definitions encountered in, say, single variable calculus. Our starting point is the observation that the completeness of the ordered set of extended real numbers $\widehat{\mathbb{R}} = \{\pm\infty\} \cup \mathbb{R}$ is equivalent to the validity of the following.

Lemma 1.1. *Let \mathcal{O} be a (partially) ordered set, let $S \subseteq \mathcal{O}$ be any subset and let $\psi : S \rightarrow \widehat{\mathbb{R}}$ be an order-preserving function. Then the set of order-preserving functions $f : \mathcal{O} \rightarrow \widehat{\mathbb{R}}$ whose restriction to S coincides with ψ has a maximum and a minimum.*

In particular, such an order-preserving function ψ singles out a distinguished subset $\text{Dar}(\psi) \subseteq \mathcal{O}$, the *Darboux set of ψ* , of elements on which the maximum and minimum extensions of ψ coincide. Equivalently, $\text{Dar}(\psi)$ can be thought of as the subset to which ψ extends canonically. We denote this canonical extension by ex_ψ .

The prototypical example of this construction is provided by the Darboux integral. Let \mathcal{O} denote the set of all bounded functions on an interval $[a, b] \subseteq \mathbb{R}$, let S be the

MSC2010: 06A06, 06A11, 18B35, 26A06, 97I10.

Keywords: partially ordered sets, Kan extensions, foundations of real analysis.

subset of step functions and let ψ be the function that to each step function assigns its integral defined naively in terms of signed areas of rectangles. In this case, as shown in [Example 7.9](#) below, $\text{Dar}(\psi)$ coincides with the set of Darboux integrable functions on $[a, b]$ and ex_ψ is the Darboux integral.

This approach to the Darboux integral exemplifies the philosophy of this paper: naturally occurring pairs (\mathcal{X}, φ) consisting of a class \mathcal{X} of $\widehat{\mathbb{R}}$ -valued functions and an order-preserving function $\varphi : \mathcal{X} \rightarrow \widehat{\mathbb{R}}$ are of the form $(\text{Dar}(\psi), \text{ex}_\psi)$ for a suitable order-preserving function ψ defined on a subset $\mathcal{S} \subseteq \mathcal{X}$ of functions that “obviously belong to \mathcal{X} ”.

For instance, let \mathcal{O} be the set of all sequences of real numbers, let \mathcal{S} be the subset of sequences that are eventually constant and let ψ be the function that to each sequence $\eta \in \mathcal{S}$ assigns the only value that η attains infinitely many times. Then, as shown in [Example 7.5](#) below, $\text{Dar}(\psi)$ coincides with the set of convergent (possibly to $\pm\infty$) sequences and $\text{ex}_\psi(f) = \lim_n f(n)$ for every $f \in \text{Dar}(\psi)$. The advantage here is that instead of having to come up with a clever (ε, δ) -definition of limit of a sequence we only need to prescribe the obvious limit of an eventually constant sequence and the formalism of Darboux calculus automatically takes care of the general case.

Similarly, let \mathcal{O} be the set of all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ and fix $x_0 \in \mathbb{R}$. It is shown in [Example 7.6](#) that if \mathcal{S} denotes the set of all functions that are constant on some open neighborhood of x_0 and ψ is the function that to each $\eta \in \mathcal{S}$ assigns $\psi(\eta) = \eta(x_0)$, then $\text{Dar}(\psi)$ is the set of functions that are continuous at x_0 and $\text{ex}_\psi(f) = f(x_0)$ for all $f \in \text{Dar}(\psi)$. Once again, given as only input the set of functions that are obviously continuous at x_0 , our machinery returns the set of functions that are continuous at x_0 as output. We view this as an intuitive alternative to the standard (ε, δ) -definition of continuity.

The statement of [Lemma 1.1](#) holds more generally if $\widehat{\mathbb{R}}$ is replaced with any ordered set that is complete in the sense that every subset has a least upper bound and a greatest lower bound. Furthermore, the inclusion $\iota : \mathcal{S} \hookrightarrow \mathcal{O}$ can be replaced with an arbitrary embedding of ordered sets. In fact, the reader familiar with category theory will easily recognize the maximum and minimum extensions of ψ in [Lemma 1.1](#) as, respectively, the right and left Kan extensions [[Mac Lane 1971](#)] (assuming they exist) of ψ along ι . Similarly, the Darboux set of ψ can be thought of as the equalizer of the left and right Kan extensions. Here we are implicitly using the standard interpretation of an ordered set \mathcal{O} as a category whose objects are the elements $x \in \mathcal{O}$ and such that $\text{Hom}(x, y)$ consists of a single element if $x \leq y$ and is empty otherwise. From the vantage point of category theory, the present paper can be summarized as the observation that equalizers of left and right Kan extensions arise naturally in elementary analysis. While some of our propositions and theorems are particular instances of much more general results about left and

right Kan extensions, we choose to give self-contained proofs in the case of ordered sets. In this way, we hope to provide evidence of the effectiveness of Darboux calculus as a stand-alone approach to the foundations of analysis that might be one day used to teach the subject at the undergraduate level.

An example of the flexibility of categorical thinking in this context comes from looking at the Yoneda embedding of an ordered set \mathcal{O} into the set of order-preserving functions from \mathcal{O} to the unique (up to a unique isomorphism) nontrivial ordered set with two elements. As it turns out, the Darboux set of the identity function of the image of the Yoneda embedding essentially coincides with the Dedekind–MacNeille completion of \mathcal{O} . While the idea of understanding Dedekind cuts in terms of presheaves is not new, see, e.g., [Taylor 1999], our emphasis is on the fact that Darboux sets are not only effective in isolating interesting classes of \mathbb{R} -valued functions but can be used to construct \mathbb{R} itself! In fact we show that with a little more effort, the field structure of \mathbb{R} can also be recovered from that of \mathbb{Q} in terms of Kan extensions. Our exposition appears to be somewhat more succinct, direct and self-contained than previous treatments of elementary analysis based on category theory; see, e.g., [Univalent Foundations 2013; Taylor 2010; Edalat and Lieutier 2004]. It would be interesting to carry out a detailed comparison between these approaches and the one presented here.

The paper is organized as follows. Section 2 contains basic material on ordered sets and order-preserving functions. In Section 3 we introduce the main concepts used in this paper, including Darboux sets and Darboux extensions. Section 4 is devoted to the notion of completeness defined here in terms of extensions of partially defined order-preserving functions. As we show, our definition, which we refer to as Darboux completeness, is in fact equivalent to the more familiar notion of Dedekind completeness. In Sections 5–6 we discuss the Yoneda embedding and the Darboux completion of an arbitrary ordered set. In particular in Section 5 we use Darboux extensions to prove that completely integrally closed subgroups of automorphisms of a complete ordered set lift to automorphisms of the completion, a result that we use to construct the field operations on \mathbb{R} . Our strategy here can be thought of as a Darboux-theoretic version of the approach used in [Fuchs 1963] to establish similar results directly at the level of Dedekind cuts. Once the real numbers are constructed, in Section 7 we shift our attention to ordered sets of \mathbb{R} -valued functions. We prove that an \mathbb{R} -valued function f has limit with respect to some filter basis F (in the sense that each ε -neighborhood of the limit contains the image $f(\mathcal{S})$ of some $\mathcal{S} \in F$) if and only if f is in the Darboux set of the partial function defined by assigning to each function constant on some $\mathcal{S} \in F$ the only value that it attains on \mathcal{S} . This characterization of convergence with respect to a filter basis yields at once Darboux-theoretic formulations of several (ε, δ) -definitions such as limits of sequences, limits of functions of one real variable and continuity. After discussing

Darboux integrability (after which the general notion of Darboux set is modeled), we use Darboux calculus to prove a theorem which simultaneously generalizes the usual linearity theorems for limits, continuous functions and integrals. In fact, all the major theorems of elementary real analysis (e.g., the intermediate value theorem, the extreme value theorem and the fundamental theorem of calculus) can be proved conceptually using the language of Darboux calculus. We hope to come back to this point elsewhere and ultimately provide an exhaustive and fully self-contained treatment of elementary real analysis in the language of this paper.

2. Preliminaries on ordered sets

Definition 2.1. A *(partially) ordered set* is a set \mathcal{O} together with a reflexive, anti-symmetric, and transitive relation, which we denote by \leq .

Example 2.2. If \mathcal{O} is an ordered set, every subset $S \subseteq \mathcal{O}$ inherits an induced order. For every $x, y \in \mathcal{O}$ such that $x \leq y$, the *interval with endpoints x and y* is the (ordered) subset $[x, y]$ of all $z \in \mathcal{O}$ such that $x \leq z \leq y$.

Example 2.3. A *discrete* set is an ordered set with the trivial order with respect to which $x \leq y$ if and only if $x = y$. If \mathcal{O} is an ordered set, we denote by $|\mathcal{O}|$ its underlying discrete set.

Remark 2.4. If \mathcal{O} is an ordered set, we denote by \mathcal{O}^{op} the *opposite* ordered set such that $|\mathcal{O}^{\text{op}}| = |\mathcal{O}|$ and $x \leq y$ in \mathcal{O}^{op} if and only if $y \leq x$ in \mathcal{O} .

Example 2.5. Given two ordered sets $\mathcal{O}_1, \mathcal{O}_2$, we denote by $\mathcal{O}_1 \times \mathcal{O}_2$ the ordered set such that $|\mathcal{O}_1 \times \mathcal{O}_2| = |\mathcal{O}_1| \times |\mathcal{O}_2|$ with order such that $(x_1, x_2) \leq (y_1, y_2)$ if and only if $x_1 \leq y_1$ and $x_2 \leq y_2$.

Definition 2.6. Let \mathcal{O} and \mathcal{P} be ordered sets. The set of *order-preserving functions from \mathcal{O} to \mathcal{P}* is

$$\text{OP}(\mathcal{O}, \mathcal{P}) = \{f : |\mathcal{O}| \rightarrow |\mathcal{P}| \mid f(x) \leq f(y) \text{ if } x \leq y\}.$$

We view $\text{OP}(\mathcal{O}, \mathcal{P})$ as an ordered set such that $f \leq g$ if and only if $f(x) \leq g(x)$ for all $x \in \mathcal{O}$. We use the shorthand notation $\text{OP}(\mathcal{O}) = \text{OP}(\mathcal{O}, \mathcal{O})$. We also say that $f \in \text{OP}(\mathcal{O}, \mathcal{P})$ is an *embedding* if for any $x, y \in \mathcal{O}$, $f(x) \leq f(y)$ implies $x \leq y$. An *isomorphism* is a surjective embedding. Given an ordered set \mathcal{O} , we denote by $\text{Aut}(\mathcal{O})$ the group of all isomorphisms in $\text{OP}(\mathcal{O})$.

Definition 2.7. If \mathcal{O} is an ordered set, we define its *augmentation* to be the ordered set $\widehat{\mathcal{O}}$ such that

- (1) $|\widehat{\mathcal{O}}| = |\mathcal{O}| \cup \{-\infty, +\infty\}$;
- (2) the canonical inclusion of $|\mathcal{O}|$ into $|\widehat{\mathcal{O}}|$ defines an embedding of \mathcal{O} into $\widehat{\mathcal{O}}$;
- (3) $\widehat{\mathcal{O}} = [-\infty, +\infty]$.

Definition 2.8. Let \mathcal{O} and \mathcal{P} be ordered sets. A *partial function* $\psi : \mathcal{O} \rightarrow \mathcal{P}$ from \mathcal{O} to \mathcal{P} is an order-preserving function $\psi : \text{dom}(\psi) \rightarrow \mathcal{P}$ defined on an ordered subset $\text{dom}(\psi) \subseteq \mathcal{O}$ called the *domain* of ψ . The ordered set $\text{im}(\psi) = \psi(\text{dom}(\psi))$ is called the *image* of ψ . An *extension* of ψ to \mathcal{O} is an order-preserving function $f : \mathcal{O} \rightarrow \mathcal{P}$ whose restriction $f|_{\text{dom}(\psi)}$ to $\text{dom}(\psi)$ coincides with ψ .

Example 2.9. Let \mathcal{O} be an ordered set and let $\mathbf{1}$ be the unique (up to a unique isomorphism) ordered set with one element. Then \mathcal{O} is canonically identified with $\text{OP}(\mathbf{1}, \mathcal{O})$.

Definition 2.10. Let \mathcal{O} and \mathcal{P} be ordered sets. A set Ψ of partial functions from \mathcal{O} to \mathcal{P} is *compatible* if for any $\psi', \psi'' \in \Psi$, the restrictions of ψ' and ψ'' to $\text{dom}(\psi') \cap \text{dom}(\psi'')$ coincide. If Ψ is compatible, we define its *common extension* to be the partial function $\psi : \mathcal{O} \rightarrow \mathcal{P}$ such that

$$\text{dom}(\psi) = \bigcup_{\psi' \in \Psi} \text{dom}(\psi')$$

and $\psi(x) = \psi'(x)$ for every $x \in \text{dom}(\psi')$ and for every $\psi' \in \Psi$.

Remark 2.11. Let \mathcal{O} and \mathcal{P} be ordered sets. If ψ is the common extension of a compatible set Ψ of partial functions from \mathcal{O} to \mathcal{P} , then $f : \mathcal{O} \rightarrow \mathcal{P}$ is an extension of ψ to \mathcal{O} if and only if it is an extension of ψ' to \mathcal{O} for each $\psi' \in \Psi$.

3. Darboux sets and Darboux extensions

Definition 3.1. Let \mathcal{O} and \mathcal{P} be ordered sets. A partial function $\psi : \mathcal{O} \rightarrow \mathcal{P}$ is *extremizable* if there exist extensions $\text{lex}_\psi, \text{uex}_\psi : \mathcal{O} \rightarrow \mathcal{P}$ of ψ to \mathcal{O} such that $\text{lex}_\psi \leq f \leq \text{uex}_\psi$ for all extensions $f : \mathcal{O} \rightarrow \mathcal{P}$ of ψ to \mathcal{O} . If this is the case, we call lex_ψ and uex_ψ the *lower and upper extensions* of ψ , respectively.

Remark 3.2. Let $\psi : \mathcal{O} \rightarrow \mathcal{P}$ be an extremizable partial function. If $x \in \text{dom}(\psi)$, then $\text{lex}_\psi(x) = \text{uex}_\psi(x)$. Therefore, if $f : \mathcal{O} \rightarrow \mathcal{P}$ is an order-preserving function such that $\text{lex}_\psi \leq f \leq \text{uex}_\psi$, then f is automatically an extension of ψ .

Example 3.3. Let \mathcal{O} and \mathcal{P} be ordered sets and let $\psi : \mathcal{O} \rightarrow \widehat{\mathcal{P}}$ be a partial function such that $\text{dom}(\psi) = \{x\}$. Then ψ is extremizable. Moreover, $\text{lex}_\psi(y) = \psi(x)$ if $x \leq y$ and $\text{lex}_\psi(y) = -\infty$ otherwise. Similarly, $\text{uex}_\psi(y) = \psi(x)$ if $y \leq x$ and $\text{uex}_\psi(y) = +\infty$ otherwise.

Definition 3.4. Let \mathcal{O} and \mathcal{P} be ordered sets. For each extremizable partial function $\psi : \mathcal{O} \rightarrow \mathcal{P}$, we define the *Darboux set* of ψ to be

$$\text{Dar}(\psi) = \{x \in \mathcal{O} \mid \text{lex}_\psi(x) = \text{uex}_\psi(x)\}.$$

Moreover, we denote by $\text{ex}_\psi : \mathcal{O} \rightarrow \mathcal{P}$ the *Darboux extension* of ψ , i.e., the restriction of uex_ψ (or equivalently of lex_ψ) to $\text{Dar}(\psi)$.

Definition 3.5. Let \mathcal{O}, \mathcal{P} be ordered sets and let ψ be a partial function from \mathcal{O} to \mathcal{P} . We say that $x \in \mathcal{O}$ is ψ -bounded if $y \leq x \leq z$ for some $y, z \in \text{dom}(\psi)$. We denote the set of ψ -bounded elements of \mathcal{O} by $\mathbf{B}(\psi)$. We say that ψ is *encompassing* if every element of \mathcal{O} is ψ -bounded. Moreover, for each extremizable $\psi : \mathcal{O} \rightarrow \mathcal{P}$ we define the *bounded Darboux set* of ψ to be the subset $\mathbf{BDar}(\psi)$ of all ψ -bounded elements of $\text{Dar}(\psi)$.

Remark 3.6. Let \mathcal{O}, \mathcal{P} be ordered sets and let ψ be the common extension of a compatible set Ψ of partial functions from \mathcal{O} to \mathcal{P} . If any $\psi' \in \Psi$ is encompassing, then $\text{dom}(\psi') \subseteq \text{dom}(\psi)$ implies that ψ is also encompassing.

Remark 3.7. Let \mathcal{O} and \mathcal{P} be ordered sets and let Ψ be a compatible set of extremizable partial functions from \mathcal{O} to \mathcal{P} . If the common extension ψ of Ψ is also extremizable, then [Remark 2.11](#) implies that $f \in [\text{lex}_\psi, \text{uex}_\psi]$ if and only if $f \in [\text{lex}_{\psi'}, \text{uex}_{\psi'}]$ for each $\psi' \in \Psi$. In particular,

$$\{\text{uex}_\psi\} = \bigcap_{\psi' \in \Psi} [\text{uex}_\psi, \text{uex}_{\psi'}] \quad \text{and} \quad \{\text{lex}_\psi\} = \bigcap_{\psi' \in \Psi} [\text{lex}_{\psi'}, \text{lex}_\psi].$$

Remark 3.8. Let \mathcal{O} and \mathcal{P} be ordered sets and let ψ be an extremizable partial function from \mathcal{O} to \mathcal{P} . If $f : \mathcal{O} \rightarrow \mathcal{P}$ is an extension of ex_ψ to \mathcal{O} , then its restriction to $\text{dom}(\psi)$ coincides with ψ and thus $\text{lex}_\psi \leq f \leq \text{uex}_\psi$. Since by construction lex_ψ and uex_ψ restrict to ex_ψ on $\text{Dar}(\psi)$, it follows that the set of extensions of ex_ψ to \mathcal{O} coincides with the set of extensions of ψ to \mathcal{O} . In particular, $\text{Dar}(\text{ex}_\psi) = \text{Dar}(\psi)$ and $\text{ex}_{\text{ex}_\psi} = \text{ex}_\psi$.

Definition 3.9. Let $\mathcal{O}_1, \mathcal{O}_2$ and \mathcal{O}_3 be ordered sets. The partial functions $\psi_1 : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ and $\psi_2 : \mathcal{O}_2 \rightarrow \mathcal{O}_3$ are *composable* if $\text{dom}(\psi_2) \cap \text{im}(\psi_1)$ is nonempty. If this is the case, their *composition* is the partial function $\psi_2 \circ \psi_1 : \mathcal{O}_1 \rightarrow \mathcal{O}_3$ such that $(\psi_2 \circ \psi_1)(x) = \psi_2(\psi_1(x))$ for each x in

$$\text{dom}(\psi_2 \circ \psi_1) = \{x \in \text{dom}(\psi_1) \mid \psi_1(x) \in \text{dom}(\psi_2)\}.$$

Proposition 3.10. Let $\mathcal{O}_1, \mathcal{O}_2$ and \mathcal{O}_3 be ordered sets. Let $\psi_1 : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ and $\psi_2 : \mathcal{O}_2 \rightarrow \mathcal{O}_3$ be partial functions such that

- (i) $\text{dom}(\psi_2) \subseteq \text{im}(\psi_1)$;
- (ii) ψ_1, ψ_2 and $\psi_2 \circ \psi_1$ are extremizable.

Then

- (1) $\text{lex}_{\psi_2 \circ \psi_1} \leq \text{lex}_{\psi_2} \circ \text{lex}_{\psi_1} \leq \text{uex}_{\psi_2} \circ \text{uex}_{\psi_1} \leq \text{uex}_{\psi_2 \circ \psi_1}$;
- (2) $\text{ex}_{\psi_1}(\text{Dar}(\psi_2 \circ \psi_1) \cap \text{Dar}(\psi_1)) \subseteq \text{Dar}(\psi_2)$;
- (3) $(\text{ex}_{\psi_2} \circ \text{ex}_{\psi_1})(x) = \text{ex}_{\psi_2 \circ \psi_1}(x)$ for all $x \in \text{Dar}(\psi_1) \cap \text{Dar}(\psi_2 \circ \psi_1)$.

Proof. Item (1) is a consequence of the fact that $\text{uex}_{\psi_2} \circ \text{uex}_{\psi_1}$ and $\text{lex}_{\psi_2} \circ \text{lex}_{\psi_1}$ are extensions of $\psi_2 \circ \psi_1$ to \mathcal{O}_1 . If $x \in \text{Dar}(\psi_2 \circ \psi_1) \cap \text{Dar}(\psi_1)$, then (1) implies

$$\text{ex}_{\psi_2 \circ \psi_1}(x) = \text{lex}_{\psi_2}(\text{ex}_{\psi_1}(x)) = \text{uex}_{\psi_2}(\text{ex}_{\psi_1}(x)),$$

which proves (2) and (3). \square

Remark 3.11. Since $\text{dom}(\psi_2) = \mathcal{O}_2$ implies $\text{lex}_{\psi_2} = \psi_2 = \text{uex}_{\psi_2}$, we know that $\text{Dar}(\psi_2 \circ \psi_1) \subseteq \text{Dar}(\psi_1)$ whenever the partial function ψ_2 in the statement of [Proposition 3.10](#) is an embedding and thus $\text{ex}_{\psi_1}(\text{Dar}(\psi_2 \circ \psi_1)) \subseteq \text{Dar}(\psi_2)$.

Lemma 3.12. *Let $\mathcal{O}, \mathcal{P}, \mathcal{P}'$ be ordered sets and let $\psi : \mathcal{O} \rightarrow \text{OP}(\mathcal{P}, \mathcal{P}')$ be an extremizable partial function. For each $p \in \mathcal{P}$, let $\text{ev}_p : \text{OP}(\mathcal{P}, \mathcal{P}') \rightarrow \mathcal{P}'$ be the order-preserving function that to each $f : \mathcal{P} \rightarrow \mathcal{P}'$ assigns its evaluation $\text{ev}_p(f) = f(p)$ at p . If $\text{ev}_p \circ \psi : \mathcal{O} \rightarrow \mathcal{P}'$ is extremizable for every $p \in \mathcal{P}$, then*

$$\text{ev}_p \circ \text{uex}_{\psi} = \text{uex}_{\text{ev}_p \circ \psi} \quad \text{and} \quad \text{ev}_p \circ \text{lex}_{\psi} = \text{lex}_{\text{ev}_p \circ \psi}.$$

Proof. Using [Proposition 3.10](#), $\text{ev}_p \circ \text{uex}_{\psi} = \text{uex}_{\text{ev}_p} \circ \text{uex}_{\psi} \leq \text{uex}_{\text{ev}_p \circ \psi}$. Consider the order-preserving function $g : \mathcal{O} \rightarrow \text{OP}(\mathcal{P}, \mathcal{P}')$ such that $(g(x))(p) = \text{uex}_{\text{ev}_p \circ \psi}$ for every $x \in \mathcal{O}$ and for every $p \in \mathcal{P}$. Then $(g(\eta))(p) = \text{uex}_{\text{ev}_p \circ \psi}(\eta) = (\psi(\eta))(p)$ for every $\eta \in \text{dom}(\psi)$. Therefore, $g \leq \text{uex}_{\psi}$ and thus $\text{uex}_{\text{ev}_p \circ \psi} = \text{ev}_p \circ g \leq \text{ev}_p \circ \text{uex}_{\psi}$. Hence, $\text{ev}_p \circ \text{uex}_{\psi} = \text{uex}_{\text{ev}_p \circ \psi}$. The second equality is proved in a similar way. \square

Lemma 3.13. *Let $\mathcal{O}, \mathcal{P}_1, \mathcal{P}_2$ be ordered sets, let $\psi : \mathcal{O} \rightarrow \mathcal{P}_1 \times \mathcal{P}_2$ be a partial function and for $i = 1, 2$ let $\pi_i : \mathcal{P}_1 \times \mathcal{P}_2 \rightarrow \mathcal{P}_i$ be the (order-preserving) projection onto the respective factor. Then ψ is extremizable if and only if $\pi_i \circ \psi : \mathcal{O} \rightarrow \mathcal{P}_i$ is extremizable for each $i = 1, 2$. If this is the case, then $\pi_i \circ \text{uex}_{\psi} = \text{uex}_{\pi_i \circ \psi}$ and $\pi_i \circ \text{lex}_{\psi} = \text{lex}_{\pi_i \circ \psi}$ for each $i = 1, 2$.*

Proof. Assume that ψ is extremizable. Then $\pi_i \circ \text{lex}_{\psi}$ and $\pi_i \circ \text{uex}_{\psi}$ are extensions of the partial function $\pi_i \circ \psi : \mathcal{O} \rightarrow \mathcal{P}_i$ (with domain $\text{dom}(\psi)$) for each $i = 1, 2$. Furthermore, if $f_1 : \mathcal{O} \rightarrow \mathcal{P}_1$ and $f_2 : \mathcal{O} \rightarrow \mathcal{P}_2$ are, respectively, extensions of $\pi_1 \circ \psi$ and $\pi_2 \circ \psi$, then $(f_1, f_2) : \mathcal{O} \rightarrow \mathcal{P}_1 \times \mathcal{P}_2$ is an extension of ψ . By assumption, this implies $\text{lex}_{\psi} \leq (f_1, f_2) \leq \text{uex}_{\psi}$ and thus

$$\pi_i \circ \text{lex}_{\psi} \leq f_i \leq \pi_i \circ \text{uex}_{\psi}$$

for each $i = 1, 2$. Hence $\pi_i \circ \psi$ is extremizable, $\pi_i \circ \text{uex}_{\psi} = \text{uex}_{\pi_i \circ \psi}$ and $\pi_i \circ \text{lex}_{\psi} = \text{lex}_{\pi_i \circ \psi}$ for each $i = 1, 2$. Conversely, assume that $\pi_1 \circ \psi$ and $\pi_2 \circ \psi$ are extremizable. Then $(\text{lex}_{\pi_1 \circ \psi}, \text{lex}_{\pi_2 \circ \psi})$ and $(\text{uex}_{\pi_1 \circ \psi}, \text{uex}_{\pi_2 \circ \psi})$ are both extensions of $\psi = (\pi_1 \circ \psi, \pi_2 \circ \psi)$. Moreover, if $f : \mathcal{O} \rightarrow \mathcal{P}_1 \times \mathcal{P}_2$ is any extension of ψ , then $\pi_i \circ f$ is an extension of $\pi_i \circ \psi$ for each $i = 1, 2$. Since $f = (\pi_1 \circ f, \pi_2 \circ f)$, this implies

$$(\text{lex}_{\pi_1 \circ \psi}, \text{lex}_{\pi_2 \circ \psi}) \leq f \leq (\text{uex}_{\pi_1 \circ \psi}, \text{uex}_{\pi_2 \circ \psi})$$

and thus ψ is extremizable with lower extension equal to $(\text{lex}_{\pi_1 \circ \psi}, \text{lex}_{\pi_2 \circ \psi})$ and upper extension equal to $(\text{uex}_{\pi_1 \circ \psi}, \text{uex}_{\pi_2 \circ \psi})$. \square

Remark 3.14. Let \mathcal{O} be a nonempty ordered set and let $\emptyset : \mathcal{O} \rightarrow \mathcal{O}$ be the empty partial function of \mathcal{O} , i.e., the unique partial function from \mathcal{O} to itself whose domain is the empty set. Since the set of extensions of \emptyset to \mathcal{O} coincides with $\text{OP}(\mathcal{O})$, if \emptyset is extremizable, then in particular $\text{lex}_{\emptyset} \leq x \leq \text{uex}_{\emptyset}$ for every constant function $x : \mathcal{O} \rightarrow \mathcal{O}$. In other words, the lower and upper Darboux extensions of the empty partial function are constant and (with a slight abuse of notation) $\mathcal{O} = [\text{lex}_{\emptyset}(\mathcal{O}), \text{uex}_{\emptyset}(\mathcal{O})]$.

4. Darboux-complete ordered sets

Definition 4.1. An ordered set \mathcal{P} is *Darboux complete* if every partial function from $\widehat{\mathcal{P}}$ to itself is extremizable.

Example 4.2. Since by [Example 3.3](#) each partial function $f : \widehat{\mathcal{O}} \rightarrow \widehat{\mathcal{O}}$ is extremizable, the empty ordered set \emptyset is Darboux complete.

Lemma 4.3. Let S be a nonempty subset of a Darboux-complete ordered set \mathcal{P} . If $\text{id}_S : \widehat{\mathcal{P}} \rightarrow \widehat{\mathcal{P}}$ is the identity function on S and $J = [\text{uex}_{\text{id}_S}(-\infty), \text{lex}_{\text{id}_S}(+\infty)]$, then

- (1) $S \subseteq J$;
- (2) J is the intersection of all intervals of $\widehat{\mathcal{P}}$ that contain S .

Proof. Since \mathcal{P} is Darboux complete, lex_{id_S} and uex_{id_S} exist. For every $s \in S$

$$\text{uex}_{\text{id}_S}(-\infty) \leq \text{uex}_{\text{id}_S}(s) = \text{lex}_{\text{id}_S}(s) \leq \text{lex}_{\text{id}_S}(+\infty),$$

which implies (1). If $x, y \in \widehat{\mathcal{P}}$ are such that $S \subseteq [x, y]$, let $\psi : \widehat{\mathcal{P}} \rightarrow \widehat{\mathcal{P}}$ be the partial function with domain \widehat{S} whose restriction to S is the identity and such that $\psi(-\infty) = x$ and $\psi(+\infty) = y$. Then

$$x = \text{uex}_{\psi}(-\infty) \leq \text{uex}_{\text{id}_S}(-\infty) \leq \text{lex}_{\text{id}_S}(+\infty) \leq \text{lex}_{\psi}(+\infty) = y. \quad \square$$

Proposition 4.4. Let \mathcal{P} be an ordered set. The following are equivalent:

- (1) \mathcal{P} is Darboux complete.
- (2) Every partial function with codomain $\widehat{\mathcal{P}}$ is extremizable.
- (3) For every ordered set \mathcal{O} , every partial function with codomain $\text{OP}(\mathcal{O}, \widehat{\mathcal{P}})$ is extremizable.

Proof. Assume that \mathcal{P} is Darboux complete. Let ψ be a partial function from an ordered set \mathcal{O} to $\widehat{\mathcal{P}}$ and let $x \in \mathcal{O}$. Consider the subsets

$$S_x = \{\psi(y) \mid y \leq x \text{ and } y \in \text{dom}(\psi)\} \subseteq \widehat{\mathcal{P}}, \quad (1)$$

$$S^x = \{\psi(y) \mid x \leq y \text{ and } y \in \text{dom}(\psi)\} \subseteq \widehat{\mathcal{P}} \quad (2)$$

together with their identity functions $\text{id}_{\mathcal{S}_x}, \text{id}_{\mathcal{S}^x} : \widehat{\mathcal{P}} \rightarrow \widehat{\mathcal{P}}$. Define $l, u : \mathcal{O} \rightarrow \widehat{\mathcal{P}}$ such that

$$l(x) = \text{lex}_{\text{id}_{\mathcal{S}_x}}(+\infty) \quad \text{and} \quad u(x) = \text{uex}_{\text{id}_{\mathcal{S}^x}}(-\infty)$$

for all $x \in \mathcal{O}$. To see that l and u are indeed order-preserving, assume that $x, y \in \mathcal{O}$ are such that $x \leq y$. Since $\mathcal{S}_x \subseteq \mathcal{S}_y$, $\text{lex}_{\text{id}_{\mathcal{S}_y}}$ is an extension of $\text{id}_{\mathcal{S}_x}$ and thus l is order-preserving. Similarly, u is order-preserving because $\mathcal{S}^y \subseteq \mathcal{S}^x$ implies that the restriction of $\text{lex}_{\text{id}_{\mathcal{S}_x}}$ to \mathcal{S}^y coincides with $\text{id}_{\mathcal{S}^y}$. Moreover l is an extension of ψ to \mathcal{O} since for every $x \in \text{dom}(\psi)$, $\mathcal{S}_x \subseteq [-\infty, \psi(x)]$ and [Lemma 4.3](#) implies

$$\psi(x) = \text{lex}_{\text{id}_{\mathcal{S}_x}}(\psi(x)) \leq l(x) \leq \psi(x).$$

On the other hand, $\psi(y) = f(y) \leq f(x)$ for any extension f of ψ to \mathcal{O} and for any $\psi(y) \in \mathcal{S}_x$. Therefore, $\mathcal{S}_x \subseteq [-\infty, f(x)]$ and thus (using again [Lemma 4.3](#)), $l(x) \leq f(x)$. Together with a similar argument involving u , this proves (2). Assume that (2) holds and let $\mathcal{O}, \mathcal{O}'$ be arbitrary ordered sets. Consider the canonical embedding α that to each partial function $\psi : \mathcal{O}' \rightarrow \text{OP}(\mathcal{O}, \widehat{\mathcal{P}})$ assigns the partial function $\alpha(\psi) : \mathcal{O}' \times \mathcal{O} \rightarrow \widehat{\mathcal{P}}$ such that $(\alpha(\psi))(x', x) = (\psi(x'))(x)$ for all $(x', x) \in \text{dom}(\alpha(\psi)) = \text{dom}(\psi) \times \mathcal{O}$. The Darboux completeness of \mathcal{P} ensures that $\alpha(\psi)$ is extremizable and thus $\text{lex}_{\alpha(\psi)} \leq \alpha(f) \leq \text{uex}_{\alpha(\psi)}$ for each extension f of ψ to \mathcal{O}' . Since the restriction of α to the subset of order-preserving functions $\mathcal{O}' \rightarrow \text{OP}(\mathcal{O}, \widehat{\mathcal{P}})$ is an isomorphism, $\text{lex}_{\psi} = \alpha^{-1}(\text{lex}_{\alpha(\psi)})$ and $\text{uex}_{\psi} = \alpha^{-1}(\text{uex}_{\alpha(\psi)})$, which proves (3). [Example 2.9](#) shows that (1) is a particular case of (3), which concludes the proof. \square

Remark 4.5. Let \mathcal{P} be an ordered set. Assume \mathcal{P} is a Darboux-complete ordered set, and $\mathcal{S} \subseteq \mathcal{P}$ is nonempty and bounded, i.e., $\mathcal{S} \subseteq [x, y]$ for some $x, y \in \mathcal{P}$. Then [Lemma 4.3](#) implies that $\text{lex}_{\text{id}_{\mathcal{S}}}(+\infty)$ and $\text{uex}_{\text{id}_{\mathcal{S}}}(-\infty)$ are respectively the least upper bound $\sup(\mathcal{S})$ and the greatest lower bound $\inf(\mathcal{S})$ of \mathcal{S} . Therefore, \mathcal{P} is Dedekind complete. Conversely, suppose that the least upper bound and the greatest lower bound of every nonempty bounded subset of \mathcal{P} exist. Given any partial function $\psi : \widehat{\mathcal{P}} \rightarrow \widehat{\mathcal{P}}$, let \mathcal{S}_x and \mathcal{S}^x be defined as in (1) and (2) respectively. Then the same argument as in the proof of [Proposition 4.4](#) shows that ψ is extremizable with $\text{lex}_{\psi}(x) = \sup(\mathcal{S}_x)$ and $\text{uex}_{\psi}(x) = \inf(\mathcal{S}^x)$ for all $x \in \mathcal{O}$. Hence, \mathcal{P} is Darboux complete if and only if \mathcal{P} is Dedekind complete. While these two notions of completeness are equivalent, the point of view of this paper is that Darboux completeness allows for a more direct and conceptual route to the foundations of elementary analysis.

Corollary 4.6. *Let \mathcal{O} be an ordered set, let \mathcal{P} be a Darboux-complete ordered set and let N be a positive integer. Every encompassing partial function from \mathcal{O} to \mathcal{P}^N is extremizable.*

Proof. By [Lemma 3.13](#), it suffices to prove the $N = 1$ case. Let $\varphi : \mathcal{O} \rightarrow \mathcal{P}$ be encompassing and let $\iota : \mathcal{P} \rightarrow \widehat{\mathcal{P}}$. By assumption, for each $z \in \mathcal{O}$ there exist

$x, y \in \text{dom}(\varphi)$ such that $x \leq z \leq y$ and thus

$$\varphi(x) = \iota(\varphi(x)) \leq \text{lex}_{\iota \circ \varphi}(z) \leq \text{uex}_{\iota \circ \varphi}(z) \leq \iota(\psi(y)) = \varphi(y).$$

Therefore, $\text{lex}_{\iota \circ \varphi}$ and $\text{uex}_{\iota \circ \varphi}$ have their image contained in \mathcal{P} and thus are extensions of φ to \mathcal{O} . Moreover, $\text{lex}_{\iota \circ \varphi}(x) \leq f(x) \leq \text{uex}_{\iota \circ \varphi}(x)$ for every extension f of φ to \mathcal{O} and for every $x \in \mathcal{P}$. Hence φ is extremizable and $\text{lex}_{\varphi}(x) = \text{lex}_{\iota \circ \varphi}(x)$, $\text{uex}_{\varphi}(x) = \text{uex}_{\iota \circ \varphi}(x)$ for all $x \in \mathcal{O}$. \square

Example 4.7. We define the *free cocompletion* of an ordered set \mathcal{O} to be the ordered set $\mathcal{O}^\vee = \text{OP}(\mathcal{O}^{\text{op}}, \widehat{\mathcal{O}})$. Let $\mathcal{P} = \mathcal{O}^\vee \setminus \{\pm\infty\}$, where $\pm\infty$ denotes the constant function such that $\text{im}(\pm\infty) = \pm\infty$. Combining [Example 4.2](#) with [Proposition 4.4](#) shows that every partial function with codomain $\mathcal{O}^\vee = \widehat{\mathcal{P}}$ is extremizable. Using [Proposition 4.4](#) again, we conclude that \mathcal{P} is Darboux complete.

Example 4.8. Let \mathcal{O} be an ordered set, let \mathcal{P} be a Darboux-complete ordered set and let \mathcal{S} be a nonempty subset of \mathcal{O} . Furthermore, let $\psi_{\mathcal{S}} : \text{OP}(\mathcal{O}, \mathcal{P}) \rightarrow \widehat{\mathcal{P}}$ be the partial function with domain the subset of functions that are constant on \mathcal{S} and such that $\psi_{\mathcal{S}}(f) = f(x)$ for every $f \in \text{dom}(\psi_{\mathcal{S}})$ and every $x \in \mathcal{S}$. For each $x \in \mathcal{S}$, ev_x coincides with $\psi_{\mathcal{S}}$ on $\text{dom}(\psi_{\mathcal{S}})$ and thus $\text{ev}_x \in [\text{lex}_{\psi_{\mathcal{S}}}, \text{uex}_{\psi_{\mathcal{S}}}]$. In particular, if $f : \mathcal{O} \rightarrow \mathcal{P}$ is in the Darboux set of $\psi_{\mathcal{S}}$, then $\text{ev}_x \circ f = \text{ev}_y \circ f$ for every $x, y \in \mathcal{S}$, i.e., f is constant on \mathcal{S} . Hence, $\text{dom}(\psi_{\mathcal{S}}) = \text{Dar}(\psi_{\mathcal{S}})$.

Remark 4.9. Using the notation of [Example 4.8](#), assume furthermore that \mathcal{O} is discrete. For every order-preserving function $f : \mathcal{O} \rightarrow \mathcal{P}$ and for every $y \in \mathcal{P}$, let $f_y \in \text{dom}(\psi_{\mathcal{S}})$ be the function whose restriction to $\mathcal{O} \setminus \mathcal{S}$ coincides with f and such that $f_y(x) = y$ for all $x \in \mathcal{S}$. In particular, if there exists $y, z \in \mathcal{P}$ such that $f(x) \in [y, z]$ for all $x \in \mathcal{S}$, then $f \in [f_y, f_z]$ and thus $[\text{lex}_{\psi_{\mathcal{S}}}, \text{uex}_{\psi_{\mathcal{S}}}] \subseteq [y, z]$. Moreover, [Corollary 4.6](#) implies that the restriction $\varphi_{\mathcal{S}} : \text{B}(\psi_{\mathcal{S}}) \rightarrow \mathcal{P}$ of $\psi_{\mathcal{S}}$ to $\text{B}(\psi_{\mathcal{S}})$ is extremizable.

5. Completely integrally closed subgroups

Proposition 5.1. Let $\mathcal{O}, \mathcal{O}'$ be ordered sets, let \mathcal{P} be a Darboux-complete ordered set and consider the composition of ordered functions $\mu : \text{OP}(\widehat{\mathcal{P}}) \times \text{OP}(\widehat{\mathcal{P}}) \rightarrow \text{OP}(\widehat{\mathcal{P}})$ defined by setting $\mu(\varphi, \varphi') = \varphi \circ \varphi'$ for all $\varphi, \varphi' \in \text{OP}(\widehat{\mathcal{P}})$. If $\psi : \mathcal{O} \rightarrow \text{OP}(\widehat{\mathcal{P}})$ and $\psi' : \mathcal{O}' \rightarrow \text{OP}(\widehat{\mathcal{P}})$ are partial functions with images in $\text{Aut}(\widehat{\mathcal{P}})$, then

$$\mu \circ (\text{uex}_{\psi} \times \text{uex}_{\psi'}) = \text{uex}_{\mu \circ (\psi \times \psi')} \quad \text{and} \quad \mu \circ (\text{lex}_{\psi} \times \text{lex}_{\psi'}) = \text{lex}_{\mu \circ (\psi \times \psi')}.$$

Proof. Since $\mu \circ (\text{uex}_{\psi} \times \text{uex}_{\psi'})$ is an extension of $\mu \circ (\psi \times \psi')$ to $\mathcal{O} \times \mathcal{O}'$, we have $\mu \circ (\text{uex}_{\psi} \times \text{uex}_{\psi'}) \leq \text{uex}_{\mu \circ (\psi \times \psi')}$. On the other hand, if $\eta \in \text{dom}(\psi)$ is fixed, then

$$(\psi(\eta))^{-1} \circ \text{uex}_{\mu \circ (\psi \times \psi')}(\eta, \eta') = \psi'(\eta')$$

for every $\eta' \in \text{dom}(\psi')$. Using the assumption that ψ' is extremizable, it follows that

$$(\psi(\eta))^{-1} \circ \text{uex}_{\mu \circ (\psi \times \psi')}(\eta, x') \leq \text{uex}_{\psi'}(x')$$

and thus

$$\begin{aligned} \text{uex}_{\mu \circ (\psi \times \psi')}(\eta, x') &\leq \psi(\eta) \circ \text{uex}_{\psi'}(x') \\ &= (\mu \circ (\text{uex}_{\psi} \times \text{uex}_{\psi'}))(\eta, x') \leq \text{uex}_{\mu \circ (\psi \times \psi')}(\eta, x') \end{aligned}$$

for all $(\eta, x') \in \text{dom}(\psi) \times \mathcal{O}'$. Setting $q = (\text{uex}_{\psi'}(x'))(p)$ yields

$$\text{ev}_p \circ \text{uex}_{\mu \circ (\psi \times \psi')}(\eta, x) = (\psi(\eta))(q) = \text{uex}_{\text{ev}_q \circ \psi}(\eta)$$

for every $p \in \widehat{\mathcal{P}}$ and for every $\eta \in \text{dom}(\psi)$. [Lemma 3.12](#) then implies

$$\begin{aligned} \text{ev}_p \circ \text{uex}_{\mu \circ (\psi \times \psi')}(x, x') &\leq \text{uex}_{\text{ev}_q \circ \psi}(x) = \text{ev}_q \circ \text{uex}_{\psi}(x) \\ &= \text{ev}_p \circ \mu \circ (\text{uex}_{\psi} \times \text{uex}_{\psi'})(x, x') \end{aligned}$$

for all $p \in \widehat{\mathcal{P}}$ and for all $(x, x') \in \mathcal{O} \times \mathcal{O}'$. This proves the first half of the proposition, the second equality is proved in a similar way. \square

Remark 5.2. Given any ordered set \mathcal{O} , the ordered set $\text{OP}(\mathcal{O})$ of order-preserving functions $f : \mathcal{O} \rightarrow \mathcal{O}$ is a monoid with respect to composition.

Definition 5.3. Let \mathcal{O} be an ordered set. A subgroup (that is, a submonoid closed under inverses) \mathcal{A} of $\text{OP}(\mathcal{O})$ is *completely integrally closed* if for every $a, a' \in \mathcal{A}$, $a^n \leq a'$ for all $n \in \mathbb{N}$ implies $a \leq \text{id}_{\mathcal{O}}$.

Remark 5.4. Completely integrally closed subgroups are a particular instance of the more general notion of (abstract) completely integrally closed ordered groups, which plays a key role in the classical study [[Fuchs 1963](#)] of embeddings in Dedekind-complete ordered groups. The remainder of this section can be thought of as an alternate construction of these embeddings formulated in the equivalent language of Darboux-complete ordered sets. Our main application is the self-contained construction of the field structure on the ordered set of real numbers described in [Section 6](#).

Proposition 5.5. *Let \mathcal{P} be a Darboux-complete ordered set. If \mathcal{A} is a completely integrally closed subgroup of $\text{OP}(\widehat{\mathcal{P}})$, then $\text{BDar}(\text{id}_{\mathcal{A}})$ is a subgroup of $\text{OP}(\widehat{\mathcal{P}})$.*

Proof. Since $\text{lex}_{\text{id}_{\mathcal{A}}} \circ \mu$ and $\text{uex}_{\text{id}_{\mathcal{A}}} \circ \mu$ are extensions of $\mu \circ (\text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{A}})$ to $(\text{OP}(\widehat{\mathcal{P}}))^2$, we obtain

$$\text{lex}_{\mu \circ (\text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{A}})} \leq \text{lex}_{\text{id}_{\mathcal{A}}} \circ \mu \leq \text{uex}_{\text{id}_{\mathcal{A}}} \circ \mu \leq \text{uex}_{\mu \circ (\text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{A}})}. \quad (3)$$

By [Proposition 5.1](#), we conclude that these inequalities restrict to equalities on $(\text{Dar}(\text{id}_{\mathcal{A}}))^2$. Hence $\text{Dar}(\text{id}_{\mathcal{A}})$ is closed under composition. We conclude that $\text{Dar}(\text{id}_{\mathcal{A}})$, which contains the submonoid \mathcal{A} of $\text{OP}(\widehat{\mathcal{P}})$, is itself a submonoid of

$\text{OP}(\widehat{\mathcal{P}})$. Given $\varphi_1, \varphi_2 \in \text{BDar}(\text{id}_{\mathcal{A}})$, by definition there exist $a_i, a'_i \in \mathcal{A}$ such that $a_i \leq \varphi_i \leq a'_i$ for $i = 1, 2$. Therefore $a_1 \circ a_2 \leq \varphi_1 \circ \varphi_2 \leq a'_1 \circ a'_2$ and thus $\text{BDar}(\psi)$ is also a submonoid. In order to construct inverses, consider the partial function $\psi : (\text{OP}(\widehat{\mathcal{P}}))^{\text{op}} \rightarrow \text{OP}(\widehat{\mathcal{P}})$ with domain \mathcal{A} and such that $\psi(a) = a^{-1}$ for every $a \in \mathcal{A}$. By [Proposition 5.1](#),

$$\text{lex}_{\psi}(\varphi) \circ \varphi = \text{lex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi) \leq \text{uex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)$$

for all $\varphi \in \text{Dar}(\text{id}_{\mathcal{A}})$. Since $\text{im}(\psi) = \mathcal{A}$,

$$\text{id}_{\mathcal{A}} \circ \mu(\psi \times \text{id}_{\mathcal{A}}) = \mu(\psi \times \text{id}_{\mathcal{A}})$$

and thus, using [Proposition 3.10](#),

$$\text{lex}_{\psi}(\varphi) \circ \varphi \leq \text{lex}_{\text{id}_{\mathcal{A}}}(\text{lex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)) \leq \text{lex}_{\text{id}_{\mathcal{A}}}(\text{uex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)). \quad (4)$$

Let $\varphi \in \text{BDar}(\text{id}_{\mathcal{A}})$, $a \in \mathcal{A}$ such that $a \leq \varphi$, and $a' \in \mathcal{A}$ such that $a' \leq \text{uex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)$. Then we have

$$a \circ a' \leq a \circ \text{uex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi) \leq a \circ \text{uex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(a, \varphi) = \varphi.$$

Iterating the same argument with a replaced by $a \circ (a')^{n-1}$ yields $a \circ (a')^n \leq \varphi$ for all $n \in \mathbb{N}$. Since \mathcal{A} is completely integrally closed, this implies $a' \leq \text{id}_{\widehat{\mathcal{P}}}$. Together with a similar argument involving $\text{lex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}$, we conclude that

$$\text{lex}_{\text{id}_{\mathcal{A}}}(\text{uex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)) \leq \text{id}_{\mathcal{P}} \leq \text{uex}_{\text{id}_{\mathcal{A}}}(\text{lex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)).$$

Therefore, applying $\text{uex}_{\text{id}_{\mathcal{A}}}$ to both sides of (4) and using [Proposition 3.10](#) yields

$$\text{id}_{\widehat{\mathcal{P}}} \leq \text{uex}_{\text{id}_{\mathcal{A}}}(\text{lex}_{\mu \circ (\psi \times \text{id}_{\mathcal{A}})}(\varphi, \varphi)) \leq \text{uex}_{\text{id}_{\mathcal{A}}}(\text{lex}_{\psi}(\varphi)) \circ \varphi \leq \text{uex}_{\text{id}_{\mathcal{A}}}(\text{id}_{\widehat{\mathcal{P}}}) = \text{id}_{\widehat{\mathcal{P}}},$$

where the last equality follows from the fact that $\text{id}_{\widehat{\mathcal{P}}}$ is an element of \mathcal{A} . Hence φ has a left inverse. A similar argument shows that it has right inverse and concludes the proof. \square

Corollary 5.6. *Let \mathcal{P} be a Darboux-complete ordered set and let $\mathcal{A} \subseteq \text{OP}(\widehat{\mathcal{P}})$ be a commutative completely integrally closed subgroup. Then $\text{BDar}(\text{id}_{\mathcal{A}})$ is a commutative group.*

Proof. Let $\mu' : \text{OP}(\widehat{\mathcal{P}}) \times \text{OP}(\widehat{\mathcal{P}}) \rightarrow \text{OP}(\widehat{\mathcal{P}})$ denote composition in reverse order; i.e., $\mu'(\varphi, \varphi') = \varphi' \circ \varphi$ for all $\varphi, \varphi' \in \text{OP}(\widehat{\mathcal{P}})$. Since the restrictions of $\text{lex}_{\text{id}_{\mathcal{A}}} \circ \mu'$ and $\text{uex}_{\text{id}_{\mathcal{A}}} \circ \mu'$ to $\mathcal{A} \times \mathcal{A}$ coincide with $\mu \circ (\text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{A}})$, we obtain

$$\text{lex}_{\mu \circ (\text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{A}})} \leq \text{lex}_{\text{id}_{\mathcal{A}}} \circ \mu' \leq \text{uex}_{\text{id}_{\mathcal{A}}} \circ \mu' \leq \text{uex}_{\mu \circ (\text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{A}})}.$$

Together with (3), this implies the commutativity of the monoid $\text{Dar}(\text{id}_{\mathcal{A}})$, which contains $\text{BDar}(\text{id}_{\mathcal{A}})$. \square

6. The Darboux completion

Remark 6.1. Let \mathcal{O} be an ordered set. For each $x \in \mathcal{O}$, let $\delta_x : \mathcal{O}^{\text{op}} \rightarrow \widehat{\mathcal{O}}$ be the partial function such that $\text{dom}(\delta_x) = \{x\}$ and $\delta_x(x) = +\infty$. Then $\text{lex}_{\delta_x}(y) = +\infty$ if and only if $y \leq x$. Let us define $Y(x) = \text{lex}_{\delta_x}$ for every $x \in \mathcal{O}$. If $f \in \mathcal{O}^\vee$ then $f(x) = +\infty$ if and only if $Y(x) \leq f$. Moreover, $f \leq g$ in \mathcal{O}^\vee if and only if $Y(x) \leq f$ implies $Y(x) \leq g$. In particular, $Y(x) \leq Y(y)$ if and only if $x \leq y$. Hence the assignment $x \mapsto Y(x)$ defines an order-preserving embedding $Y : \mathcal{O} \rightarrow \mathcal{O}^\vee$ called the *Yoneda embedding of \mathcal{O}* .

Proposition 6.2. *Let \mathcal{O} be an ordered set, let $\varphi : \mathcal{O}^\vee \rightarrow \mathcal{O}^\vee$ be the identity function of the image of the Yoneda embedding of \mathcal{O} and let $\text{Dar}(\mathcal{O})$ denote the Darboux set of φ . Then*

- (1) $\text{lex}_\varphi = \text{id}_{\mathcal{O}^\vee}$;
- (2) *if $g \in \text{OP}(\text{Dar}(\mathcal{O}))$ restricts to the identity on $Y(\mathcal{O})$, then $g = \text{id}_{\text{Dar}(\mathcal{O})}$;*
- (3) $\text{uex}_\varphi(\mathcal{O}^\vee) \subseteq \text{Dar}(\mathcal{O})$;
- (4) *the empty partial function of $\text{Dar}(\mathcal{O})$ is extremizable.*

Proof. Since $\text{id}_{\mathcal{O}^\vee}$ restricts to φ on $Y(\mathcal{O})$, we know $\text{lex}_\varphi(f) \leq f$ for every $f \in \mathcal{O}^\vee$. On the other hand, $Y(x) \leq f$ implies $Y(x) = \text{lex}_\varphi(Y(x)) \leq \text{lex}_\varphi(f)$. Using [Remark 6.1](#), this proves (1). Item (2) follows immediately from (1) and the definition of $\text{Dar}(\mathcal{O})$. [Proposition 3.10](#) and (1) yield

$$\text{uex}_\varphi = \text{uex}_\varphi \circ \text{lex}_\varphi \leq \text{uex}_\varphi \circ \text{uex}_\varphi \leq \text{uex}_{\varphi \circ \varphi} = \text{uex}_\varphi,$$

which readily implies (3). Since $+\infty \leq \text{uex}_\varphi(+\infty) \leq +\infty$, we have $+\infty \in \text{Dar}(\mathcal{O})$. If $-\infty \notin \text{Dar}(\mathcal{O})$, then by [Remark 6.1](#) there exists $x \in \mathcal{O}$ such that $Y(x) \leq \text{uex}_\varphi(-\infty)$. By [Lemma 4.3](#) this implies $x \leq y$ for all $y \in \mathcal{O}$. Therefore, the empty partial function of $\text{Dar}(\mathcal{O})$ is extremizable, $\text{uex}_{\mathcal{O}}(\mathcal{O}) = +\infty$ and $\text{lex}_{\mathcal{O}}(\mathcal{O})$ is the function that takes the value $-\infty$ on the complement of a set of cardinality at most 1. \square

Definition 6.3. Using the notation of [Proposition 6.2](#) and [Remark 3.14](#), we define the *Darboux completion of an ordered set \mathcal{O}* to be the ordered set

$$\text{Dar}'(\mathcal{O}) = \text{Dar}(\mathcal{O}) \setminus \{\text{lex}_{\mathcal{O}}(\text{Dar}(\mathcal{O})), \text{uex}_{\mathcal{O}}(\text{Dar}(\mathcal{O}))\}.$$

Corollary 6.4. *The Darboux completion of an ordered set is Darboux complete.*

Proof. Let \mathcal{O} be an ordered set and let $\iota : \text{Dar}(\mathcal{O}) \rightarrow \mathcal{O}^\vee$ be the inclusion. For any partial function $\psi : \text{Dar}(\mathcal{O}) \rightarrow \text{Dar}(\mathcal{O})$, [Example 4.7](#) ensures that $\iota \circ \psi$ is extremizable. By [Proposition 6.2](#), $\text{uex}_\varphi \circ \text{uex}_{\iota \circ \psi}$ and $\text{uex}_\varphi \circ \text{lex}_{\iota \circ \psi}$ are order-preserving functions in $\text{OP}(\text{Dar}(\mathcal{O}))$ that restrict to ψ on $\text{dom}(\psi)$. On the other hand, $\text{lex}_{\iota \circ \psi} \leq \iota \circ g \leq \text{uex}_{\iota \circ \psi}$ for any extension g of ψ to $\text{Dar}(\mathcal{O})$. Since $\text{uex}_\varphi \circ \iota \circ g = g$, this implies

$\text{uex}_\varphi \circ \text{lex}_{\psi'} \leq g \leq \text{uex}_\varphi \circ \text{uex}_{\psi'}$ and thus ψ is extremizable. This concludes the proof, since by construction $\text{Dar}(\mathcal{O})$ is canonically isomorphic to $\widehat{\text{Dar}'(\mathcal{O})}$. \square

Remark 6.5. From now on we use the Yoneda embedding to canonically identify \mathcal{O} with a subset of $\text{Dar}(\mathcal{O}) \subseteq \mathcal{O}^\vee$. In particular, this provides a canonical embedding of $\text{OP}(\mathcal{O})$ into the set of partial functions $\text{Dar}(\mathcal{O}) \rightharpoonup \text{Dar}(\mathcal{O})$.

Example 6.6. We define the set of *real numbers* to be the Darboux completion \mathbb{R} of the ordered set \mathbb{Q} of rational numbers. Moreover, $\text{Dar}(\mathbb{Q})$ is canonically identified with the set of *extended real numbers* $\widehat{\mathbb{R}} = \mathbb{R} \cup \{\pm\infty\}$.

Example 6.7. Let $\mathbb{Q}_{>0} \subseteq \mathbb{Q}$ be the ordered set of positive rational numbers and let $\mathbb{R}_{>0} = \text{Dar}'(\mathbb{Q}_{>0})$. Extending each function in $\text{Dar}'(\mathbb{Q}_{>0})$ by $+\infty$ to $\mathbb{Q} \setminus \mathbb{Q}_{>0}$ yields a canonical embedding of $\mathbb{R}_{>0}$ into \mathbb{R} whose image consists of real numbers that are greater than $Y(0)$. Moreover, the composition of this embedding with the canonical embedding of $\mathbb{Q}_{>0}$ into $\text{Dar}'(\mathbb{Q}_{>0})$ coincides with the restriction of the canonical embedding of \mathbb{Q} into \mathbb{R} . Keeping in mind the above canonical identifications, it makes sense to write equalities such as $\mathbb{Q}_{>0} = \mathbb{Q} \cap \mathbb{R}_{>0}$.

Remark 6.8. Let \mathcal{O} be an ordered set, let \mathcal{P} be a complete ordered set and let $\psi : \text{Dar}(\mathcal{O}) \rightarrow \widehat{\mathcal{P}}$ be an embedding with domain $Y(\mathcal{O})$ and inverse $\psi' : \widehat{\mathcal{P}} \rightarrow \text{Dar}(\mathcal{O})$. Since $\text{uex}_{\psi'} \circ \text{uex}_\psi$ restricts to the identity on $Y(\mathcal{O})$, it is equal to $\text{id}_{\text{Dar}(\mathcal{O})}$ by [Proposition 6.2](#). Therefore, $\text{uex}_\psi : \text{Dar}(\mathcal{O}) \rightarrow \widehat{\mathcal{P}}$ is an embedding. In particular, it can attain the values $\pm\infty$ at most once, which implies that uex_ψ restricts to an embedding $f : \text{Dar}'(\mathcal{O}) \rightarrow \mathcal{P}$. By [Remark 4.5](#) this implies that $\text{Dar}'(\mathcal{O})$ also satisfies the universal property of the Dedekind–MacNeille completion of \mathcal{O} and is therefore canonically isomorphic to it. In particular, this shows that our definition of \mathbb{R} is canonically isomorphic to the ordered set \mathbb{R}' of Dedekind cuts of \mathbb{Q} . In fact, in this case it is easy to see directly that $\text{uex}_{\psi'} : \widehat{\mathbb{R}'} \rightarrow \widehat{\mathbb{R}}$ is injective since it maps the cut associated to a rational number x to $Y(x)$ and $(\text{uex}_{\psi'}(C))^{-1}(+\infty) = C$ for any irrational cut C .

Proposition 6.9. *There exists a canonical embedding $\alpha : \text{Aut}(\mathcal{O}) \rightarrow \text{Aut}(\text{Dar}(\mathcal{O}))$. Moreover, α is a group homomorphism.*

Proof. Let $\varphi \in \text{Aut}(\mathcal{O})$. Using the convention of [Remark 6.5](#), we may think of φ as a partial function $\text{Dar}(\mathcal{O}) \rightharpoonup \text{Dar}(\mathcal{O})$. Then by [Remark 6.8](#)

$$\text{lex}_{\varphi^{-1}} \circ \text{uex}_\varphi = \text{id}_{\text{Dar}(\mathcal{O})} = \text{uex}_\varphi \circ \text{lex}_{\varphi^{-1}}.$$

This implies that uex_φ is invertible and $\text{uex}_\varphi \leq \text{lex}_\varphi \circ \text{lex}_{\varphi^{-1}} \circ \text{uex}_\varphi \leq \text{lex}_\varphi$. Therefore, $\text{ex}_\varphi \in \text{Aut}(\text{Dar}(\mathcal{O}))$. Let $\alpha(\varphi) = \text{ex}_\varphi$ for all $\varphi \in \text{Aut}(\mathcal{O})$. Combining [Remark 6.8](#) and [Proposition 3.10](#), we conclude that α is an injective group homomorphism and the proposition is proved. \square

Example 6.10. Addition in \mathbb{Q} defines an embedding $\lambda : \mathbb{Q} \rightarrow \text{Aut}(\mathbb{Q})$ such that $(\lambda(r))(s) = r + s$ for all $r, s \in \mathbb{Q}$. Composing with α we obtain an embedding $\beta : \mathbb{Q} \rightarrow \text{Aut}(\widehat{\mathbb{R}})$. Since every (order-preserving) automorphism of $\widehat{\mathbb{R}}$ necessarily fixes $\pm\infty$, we have a canonical identification of $\text{Aut}(\widehat{\mathbb{R}})$ with $\text{Aut}(\mathbb{R})$. In particular, $(\beta(x))(\pm\infty) = \pm\infty$ for all $x \in \mathbb{Q}$.

Proposition 6.11. \mathbb{R} is canonically isomorphic to $\text{BDar}(\text{id}_{\beta(\mathbb{Q})})$.

Proof. Considering the embedding β constructed in [Example 6.10](#) as a partial function $\mathbb{R} \rightarrow \text{OP}(\mathbb{R})$ (which is extremizable by [Proposition 4.4](#)), we obtain order-preserving functions $\text{lex}_\beta, \text{uex}_\beta : \mathbb{R} \rightarrow \text{OP}(\widehat{\mathbb{R}})$. The order-preserving function $\text{ev}_0 : \text{BDar}(\text{id}_{\beta(\mathbb{Q})}) \rightarrow \mathbb{R}$ is surjective by [Remark 6.8](#) since $\text{ev}_0 \circ \text{lex}_\beta$ and $\text{ev}_0 \circ \text{uex}_\beta$ both restrict to the identity on \mathbb{Q} . Since $\text{lex}_\beta \circ \text{ev}_0$ and $\text{uex}_\beta \circ \text{ev}_0$ both restrict to the identity on $\beta(\mathbb{Q})$, they both equal the identity on $\text{BDar}(\text{id}_{\beta(\mathbb{Q})})$. Therefore ev_0 is invertible with inverse ex_β . \square

Remark 6.12. Combining [Proposition 6.11](#) with [Corollary 5.6](#), we conclude that \mathbb{R} has a canonical structure of commutative group. Alternatively, this structure can be understood as follows. Let $+$ be the addition operation on \mathbb{Q} , thought of as a partial function $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$. Since $(\text{ex}_\beta(r))(s) = r + s$ for all $r, s \in \mathbb{Q}$, we obtain

$$\text{lex}_+(x, y) \leq (\text{ex}_\beta(x))(y) \leq \text{uex}_+(x, y) \quad (5)$$

for all $x, y \in \mathbb{R}$. On the other hand, for every $r \in \mathbb{Q}$ both $\text{ex}_\beta(r)^{-1} \circ \text{uex}_+(r, -)$ and $\text{ex}_\beta(r)^{-1} \circ \text{lex}_+(r, -)$ restrict to the identity of \mathbb{Q} . By [Remark 6.8](#), this implies $\text{lex}_+(r, -) = \text{ex}_\beta(r) = \text{uex}_+(r, -)$ for all $r \in \mathbb{Q}$ and thus $\text{lex}_\beta(x) \leq \text{lex}_+(x, -) \leq \text{uex}_+(x, -) \leq \text{uex}_\beta(x)$ for all $x \in \mathbb{R}$. Hence the inequalities of (5) are actually equalities for all $x, y \in \mathbb{R}$.

Remark 6.13. A similar argument shows that the multiplication on $\mathbb{Q}_{>0}$ thought of as a partial function $\bullet : \mathbb{R}_{>0} \times \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ defines a partial function $\gamma : \mathbb{R}_{>0} \rightarrow \text{Aut}(\mathbb{R}_{>0})$ such that $\text{dom}(\gamma) = \mathbb{Q}_{>0}$ and $(\text{ex}_\gamma(x))(y) = \text{ex}_\bullet(x, y)$ for all $x, y \in \mathbb{R}_{>0}$.

Theorem 6.14. $(\mathbb{R}_{>0}, \text{ex}_+, \text{ex}_\bullet)$ is a semifield.

Proof. Let $\psi : (\mathbb{R}_{>0})^3 \rightarrow \mathbb{R}_{>0}$ be the partial function with domain $(\mathbb{Q}_{>0})^3$ and such that $\psi(r, s, t) = r(s + t)$ for all $r, s, t \in \mathbb{Q}_{>0}$. Since $\text{ex}_\bullet(r, \text{ex}_+(s, t)) = \psi(r, s, t)$ for all $r, s, t \in \mathbb{Q}_{>0}$,

$$\text{lex}_\psi(x, y, z) \leq \text{ex}_\bullet(x, \text{ex}_+(y, z)) \leq \text{uex}_\psi(x, y, z) \quad (6)$$

for all $x, y, z \in \mathbb{R}_{>0}$. On the other hand, since $\gamma(s+t)$ agrees with both $\text{lex}_\psi(-, s, t)$ and $\text{uex}_\psi(-, s, t)$ on $\mathbb{Q}_{>0}$ for all $s, t \in \mathbb{Q}_{>0}$, they also agree on $\mathbb{R}_{>0}$. Using that $(\gamma(s+t))(x) = \text{ex}_\bullet(x, s+t) = (\text{ex}_\gamma(x))(s+t)$, we obtain

$$\text{lex}_+(y, z) \leq (\text{ex}_\gamma(x))^{-1} \text{lex}_\psi(x, y, z) \leq (\text{ex}_\gamma(x))^{-1} \text{uex}_\psi(x, y, z) \leq \text{uex}_+(y, z).$$

Since $\text{ex}_\bullet(x, \text{ex}_+(y, z)) = (\text{ex}_\gamma(x))(\text{ex}_+(y, z))$ for all $x, y, z \in \mathbb{R}_{>0}$, we conclude that the inequalities of (6) are in fact equalities. It follows from the distributivity of \bullet over $+$ on $\mathbb{Q}_{>0}$ that $\text{ex}_+(\text{ex}_\bullet(r, t), \text{ex}_\bullet(s, t)) = \psi(r, s, t)$ for all $r, s, t \in \mathbb{Q}_{>0}$. Hence, ex_\bullet distributes over ex_+ on $\mathbb{R}_{>0}$ and the theorem is proved. \square

Remark 6.15. A standard argument shows that ex_\bullet can be canonically extended to an operation \cdot on \mathbb{R} (which is not order-preserving) in such a way that $(\mathbb{R}, \text{ex}_+, \cdot)$ is a field. With a slight abuse of notation, from now on we write $+$ for ex_+ . Since $(\beta(x))(\pm\infty) = \pm\infty$ for all $x \in \mathbb{Q}$, we set $x + (\pm\infty) = \pm\infty$ for all $x \in \mathbb{R}$.

Remark 6.16. For each ordered set \mathcal{O} , the set $\text{OP}(\mathcal{O}, \mathbb{R})$ inherits a canonical structure of \mathbb{R} -algebra with operations defined pointwise on \mathcal{O} . In particular, $f_1 \leq f_2$ implies $f_1 + f_3 \leq f_2 + f_3$ for any $f_1, f_2, f_3 \in \text{OP}(\mathcal{O}, \mathbb{R})$ and $f_1 f_3 \leq f_2 f_3$ whenever $0 \geq f_3$.

7. Limits and integrals

Definition 7.1. A *filter basis* on an ordered set \mathcal{O} is a collection F of nonempty subsets of \mathcal{O} that is closed under finite intersections. To each filter basis F of \mathcal{O} we associate the partial function $\psi_F : \text{OP}(\mathcal{O}, \mathbb{R}) \rightarrow \widehat{\mathbb{R}}$ such that

$$\text{dom}(\psi_F) = \bigcup_{S \in F} \text{dom}(\psi_S),$$

where ψ_S is defined as in [Example 4.8](#) and $\psi_F(f) = \psi_S(f)$ for each $f \in \text{dom}(\psi_S)$ and for each $S \in F$.

Definition 7.2. Let \mathcal{O} be a discrete set and let F be a filter basis on \mathcal{O} . An order-preserving function $f : \mathcal{O} \rightarrow \mathbb{R}$ is *F-convergent* if there exists $\lim_F(f) \in \widehat{\mathbb{R}}$ such that for every $\varepsilon > 0$ there exists $S \in F$ such that $f(x) \in [\lim_F(f) - \varepsilon, \lim_F(f) + \varepsilon]$ for all $x \in S$.

Theorem 7.3. Let \mathcal{O} be a discrete set and let F be a filter basis on \mathcal{O} . An order-preserving function $f : \mathcal{O} \rightarrow \mathbb{R}$ is *F-convergent* if and only if $f \in \text{Dar}(\psi_F)$. Moreover, $\text{ex}_{\psi_F}(f) = \lim_F(f)$ for all $f \in \text{Dar}(\psi_F)$.

Proof. Assume that $f \in \text{Dar}(\psi_F)$. Then by [Remark 2.11](#) for every $\varepsilon > 0$ there exist $S', S'' \in F$ such that $[\text{ex}_{\psi_F}(f), \text{ex}_{\psi_{S'}}(f)] \subseteq [\text{ex}_{\psi_F}(f), \text{ex}_{\psi_F}(f) + \varepsilon]$ and $[\text{ex}_{\psi_{S''}}(f), \text{ex}_{\psi_F}(f)] \subseteq [\text{ex}_{\psi_F}(f) - \varepsilon, \text{ex}_{\psi_F}(f)]$. Therefore, setting $S = S' \cap S''$ we obtain

$$\text{ex}_{\psi_F}(f) - \varepsilon \leq \text{lex}_{\psi_S}(f) \leq f(x) \leq \text{uex}_{\psi_S}(f) \leq \text{ex}_{\psi_F}(f) + \varepsilon$$

for every $x \in S$. Hence f is *F-convergent* and $\lim_F(f) = \text{ex}_{\psi_F}(f)$. Conversely, if f is *F-convergent*, for every $\varepsilon > 0$ there exists $S \in F$ such that

$$\lim_F(f) - \varepsilon \leq f(x) \leq \lim_F(f) + \varepsilon$$

for all $x \in \mathcal{S}$. Using [Remark 4.9](#), this implies

$$[\text{lex}_{\psi_F}(f), \text{uex}_{\psi_F}(f)] \subseteq [\text{lex}_{\psi_{\mathcal{S}}}(f), \text{uex}_{\psi_{\mathcal{S}}}(f)] \subseteq [\lim_F(f) - \varepsilon, \lim_F(f) + \varepsilon]$$

for every $\varepsilon > 0$. Hence, $f \in \text{Dar}(\psi_F)$ and $\lim_F(f) = \text{ex}_{\psi_F}(f)$. \square

Remark 7.4. In terms of the philosophy outlined in [Section 1](#), the functions in $\text{dom}(\psi_F)$, i.e., the functions that are constant on some element of F , are “obviously F -convergent” and ψ_F is their “obvious limit”. Feeding the machinery of Darboux calculus with this information results in a construction of general F -convergent functions that is alternative to the (ε, δ) -definition given in [Definition 7.2](#).

Example 7.5. Let \mathbb{N} be the set of natural numbers with its usual order. Let $\mathcal{O} = |\mathbb{N}|$ and let $F = \{\mathbb{N} \setminus [1, n]\}_{n \in \mathbb{N}}$. Then $\text{OP}(\mathcal{O}, \mathbb{R})$ is the set of all sequences, $\text{dom}(\psi_F)$ is the set of sequences that are eventually constant and $\psi_F(f)$ is the function that to such a sequence assigns its obvious limit, i.e., the only value that f attains infinitely many times. Moreover, $\text{Dar}(\psi_F)$ is precisely the set of all convergent sequences (including those converging to $\pm\infty$) and ex_{ψ_F} is their limit.

Example 7.6. Let $\mathcal{O} = |\mathbb{R}|$, let $x_0 \in \mathbb{R}$ and let F be the collection of all subsets of the form $[x_0 - \delta, x_0 + \delta]$ for some $\delta > 0$. Then $\text{OP}(\mathcal{O}, \mathbb{R})$ is the set of all real-valued functions of one real variable, $\text{dom}(\psi_F)$ is the subset of functions that are constant in a neighborhood of x_0 and $\psi_F(f) = f(x_0)$ for all $f \in \text{dom}(\psi_F)$. Moreover, $\text{Dar}(\psi_F)$ is precisely the set of all functions that are continuous at x_0 and $\text{ex}_{\psi_F}(f) = f(x_0)$ for all $f \in \text{dom}(\psi_F)$.

Example 7.7. In the notation of [Example 7.6](#), we could also consider F to be the collection of all subsets of the form $[x_0 - \delta, x_0 + \delta] \setminus \{x_0\}$ for some $\delta > 0$. Then $f \in \text{Dar}(\psi_F)$ if and only if f has a limit at x_0 , in which case $\text{ex}_{\psi_F}(f)$ equals the limit. We leave the obvious variations leading to left and right limits to the reader.

Definition 7.8. We denote by $\text{Int}(\mathcal{O})$ the ordered set of all intervals with endpoints in \mathcal{O} with order given by $[a, b] \leq [c, d]$ if and only if $c \leq a \leq b \leq d$. We write $\text{int}(\mathcal{O})$ for the collection of nonempty subsets of \mathcal{O} of the form $[x, z] \setminus \{x, z\}$, for some $[x, z] \in \text{Int}(\mathcal{O})$ (also ordered by inclusion).

Example 7.9. For any $J \in \text{Int}(\mathbb{R})$ let $m : \text{int}(J) \rightarrow \mathbb{R}_{>0}$ be the order-preserving function defined by $m([x, z] \setminus \{x, z\}) = z - x$ whenever $x \leq z$ and 0 otherwise. Given $J \in \text{Int}(\mathbb{R})$, let $\text{Par}(J)$ be the collection of *partitions* of J , i.e., finite collections $P \subseteq \text{int}(J)$ of mutually disjoint subsets such that $J \setminus \bigcup_{I \in P} I$ is finite. For each $I \in \text{int}(J)$, let $\psi_I : \text{OP}(|J|, \mathbb{R}) \rightarrow \mathbb{R}$ be the partial function associated to the nonempty subset I as in [Example 4.8](#). In particular, the set of all bounded \mathbb{R} -valued functions on J coincides with

$$\mathcal{O} = B(\psi_J) = \bigcap_{I \in P} B(\psi_I)$$

for any $P \in \text{Par}(J)$. Let $\varphi_I : B(\psi_I) \rightarrow \mathbb{R}$ be the extremizable partial function defined as in [Remark 4.9](#) for each $I \in \text{int}(J)$ and let $U_P, L_P : \mathcal{O} \rightarrow \mathbb{R}$ be the order-preserving functions defined by

$$U_P = \sum_{I \in P} m(I) \text{uex}_{\varphi_I} \quad \text{and} \quad L_P = \sum_{I \in P} m(I) \text{lex}_{\varphi_I}.$$

By [Remark 4.9](#) it is clear that for each $f \in \mathcal{O}$, $U_P(f)$ coincides with the usual *upper Darboux sums* of f and $L_P(f)$ coincides with the usual *lower Darboux sums* of f ; see, e.g., [\[Rudin 1953\]](#). On the other hand, for each $P \in \text{Par}(J)$ consider the partial function $\varphi_P : \mathcal{O} \rightarrow \mathbb{R}$ defined by

$$\varphi_P(f) = \sum_{I \in P} m(I) \varphi_I(f) \quad (7)$$

for each element f of

$$\text{dom}(\varphi_P) = \bigcap_{I \in P} \text{dom}(\varphi_I).$$

Since φ_P is clearly encompassing, it is also extremizable by [Corollary 4.6](#). Moreover,

$$\text{lex}_{\varphi_P} \leq L_P \leq U_P \leq \text{uex}_{\varphi_P}, \quad (8)$$

as each term in the above chain of inequalities restricts to φ_P on $\text{dom}(\varphi_P)$. Since each function in \mathcal{O} attains only finitely many values, $\text{dom}(\varphi_P) \cong \mathbb{R}^N$ for some integer N . In particular, [Corollary 4.6](#) ensures that if $\rho_P : \mathcal{O} \rightarrow \text{dom}(\varphi_P)$ denotes the identity function on $\text{dom}(\varphi_P)$ then ρ_P is extremizable. Therefore

$$\text{uex}_{\varphi_P} \leq \varphi_P \circ \text{uex}_{\rho_P} = \sum_{I \in P} m(I) (\varphi_I \circ \text{uex}_{\rho_P}) \leq \sum_{I \in P} m(I) \text{uex}_{\varphi_I \circ \rho_P} = U_P. \quad (9)$$

Combined with an analogous estimate for lex_{φ_P} and [\(8\)](#), [\(9\)](#) shows that $\text{uex}_{\varphi_P} = U_P$ and $\text{lex}_{\varphi_P} = L_P$. Since $m(I) = m(I_1) + m(I_2)$ whenever $I \setminus (I_1 \cup I_2)$ is finite and $I \subseteq I'$ implies $\varphi_I(f) = \varphi_{I'}(f)$ for each $f \in \text{dom}(\varphi'_I)$, we have for each $f \in \text{dom}(\varphi_P) \cap \text{dom}(\varphi_{P'})$

$$\varphi_P(f) = \sum_{I \in P} m(I) \varphi_I(f) = \sum_{I \in P} \sum_{I' \in P'} m(I \cap I') \varphi_{I \cap I'}(f) = \sum_{I' \in P'} m(I') \varphi_{I'}(f) = \varphi_{P'}(f).$$

Let φ be the common extension of the compatible set $\{\varphi_P\}_{P \in \text{Par}(J)}$. In particular, $\text{dom}(\varphi)$ is the set of *step functions on J* , i.e., the set of all functions on $|J|$ that are constant on each interval of some partition of J . Combining [Remark 3.6](#) with [Corollary 4.6](#) we conclude that φ is encompassing and thus extremizable. By [Remark 3.7](#), an order-preserving function $g : \mathcal{O} \rightarrow \mathbb{R}$ restricts to φ on $\text{dom}(\varphi)$ if and only if

$$g \in \bigcap_{P \in \text{Par}(J)} [\text{lex}_{\varphi_P}, \text{uex}_{\varphi_P}] = \bigcap_{P \in \text{Par}(J)} [L_P, U_P].$$

Hence, lex_φ and uex_φ coincide with the lower and upper integrals of f on J , respectively. In particular, $\text{Dar}(\varphi)$ coincides with the set of functions on J that are integrable in the sense of Riemann and ex_φ is the Riemann integral. This is in fact the motivating example for the philosophy of [Section 1](#): to define the Riemann integral it is sufficient to feed the “obvious” definition for step functions, given by (7), into the machinery of Darboux calculus to automatically obtain the correct general definition.

Theorem 7.10 (linearity). *Let $\psi : \text{OP}(\mathcal{O}, \mathbb{R}) \rightarrow \mathbb{R}$ be a partial function such that*

- (1) *ψ is encompassing;*
- (2) *$\text{dom}(\psi)$ is an \mathbb{R} -linear subspace of $\text{OP}(\mathcal{O}, \mathbb{R})$;*
- (3) *ψ is an \mathbb{R} -linear transformation.*

Then for every $f_1, f_2 \in \text{OP}(\mathcal{O}, \mathbb{R})$ and for every nonnegative $a_1, a_2 \in \mathbb{R}$

$$\text{uex}_\psi(a_1 f_1 + a_2 f_2) \leq a_1 \text{uex}_\psi(f_1) + a_2 \text{uex}_\psi(f_2) \quad (10)$$

and similarly

$$a_1 \text{lex}_\psi(f_1) + a_2 \text{lex}_\psi(f_2) \leq \text{lex}_\psi(a_1 f_1 + a_2 f_2). \quad (11)$$

Moreover

$$-\text{lex}_\psi(f) = \text{uex}_\psi(-f) \quad (12)$$

for every $f \in \text{OP}(\mathcal{O}, \mathbb{R})$. In particular, $\text{Dar}(\psi)$ is an \mathbb{R} -linear subspace of $\text{OP}(\mathcal{O}, \mathbb{R})$ and ex_ψ is \mathbb{R} -linear.

Proof. Since ψ is encompassing, it is extremizable. By the additivity of ψ , the assignment $f_1 \mapsto \text{uex}_\psi(f_1 + \eta_2) - \psi(\eta_2)$ coincides with ψ on $\text{dom}(\psi)$ for each fixed $\eta_2 \in \text{dom}(\psi)$. Therefore

$$\text{uex}_\psi(f_1 + \eta_2) \leq \text{uex}_\psi(f_1) + \psi(\eta_2) \quad (13)$$

for every $f_1 \in \text{OP}(\mathcal{O}, \mathbb{R})$ and for every $\eta_2 \in \text{dom}(\psi)$. In particular

$$\text{uex}_\psi(f_1) + \psi(\eta_2) = \text{uex}_\psi((f_1 + \eta_2) + (-\eta_2)) + \psi(\eta_2) \leq \text{uex}_\psi(f_1 + \eta_2)$$

and thus the inequality in (13) is actually an equality. Therefore, for each $f_1 \in \text{OP}(\mathcal{O}, \mathbb{R})$ the assignment $f_2 \mapsto \text{uex}_\psi(f_1 + f_2) - \text{uex}_\psi(f_1)$ coincides with ψ on $\text{dom}(\psi)$. This proves (10) when $a_1 = a_2 = 1$. Since ψ is compatible with scalar multiplication, a is a positive real number and the assignment $f \mapsto a^{-1} \text{uex}_\psi(af)$ coincides with ψ on $\text{dom}(\psi)$. Therefore, $\text{uex}_\psi(af) \leq a \text{uex}_\psi(f)$, which in turn implies

$$a \text{uex}_\psi(f) = a \text{uex}_\psi(a^{-1}(af)) \leq \text{uex}_\psi(af).$$

As a result, $a \text{uex}_\psi(f) = \text{uex}_\psi(af)$ for every positive real number a and for every $f \in \text{OP}(\mathcal{O}, \mathbb{R})$. This proves (10) and (11) is proved similarly. Since ψ is odd,

the assignments $f \mapsto -\text{lex}_\psi(-f)$ and $f \mapsto -\text{uex}_\psi(-f)$ both restrict to ψ on $\text{dom}(\psi)$ and thus

$$\text{lex}_\psi(f) \leq -\text{uex}_\psi(-f) \leq -\text{lex}_\psi(-f) \leq \text{uex}_\psi(f) \quad (14)$$

for every $f \in \text{OP}(\mathcal{O}, \mathbb{R})$. The first inequality in (14) implies $\text{uex}_\psi(-f) \leq -\text{lex}_\psi(f)$, while the last inequality of (14) implies $-\text{lex}_\psi(f) \leq \text{uex}_\psi(-f)$ and thus (12). The last statement is a straightforward consequence of (10)–(12). \square

Example 7.11. Specializing Theorem 7.10 to Examples 7.5–7.9 we obtain the well-known linearity theorems for limits of sequences, continuous functions, limits of functions of real variable and integrals.

References

- [Edalat and Lieutier 2004] A. Edalat and A. Lieutier, “Domain theory and differential calculus (functions of one variable)”, *Math. Structures Comput. Sci.* **14**:6 (2004), 771–802. [MR](#) [Zbl](#)
- [Fuchs 1963] L. Fuchs, *Partially ordered algebraic systems*, Pergamon, Oxford, 1963. [MR](#) [Zbl](#)
- [Mac Lane 1971] S. Mac Lane, *Categories for the working mathematician*, Graduate Texts in Math. **5**, Springer, 1971. [MR](#) [Zbl](#)
- [Rudin 1953] W. Rudin, *Principles of mathematical analysis*, McGraw-Hill, New York, 1953. [MR](#) [Zbl](#)
- [Taylor 1999] P. Taylor, *Practical foundations of mathematics*, Cambridge Studies in Advanced Math. **59**, Cambridge Univ. Press, 1999. [MR](#) [Zbl](#)
- [Taylor 2010] P. Taylor, “A lambda calculus for real analysis”, *J. Log. Anal.* **2** (2010), art. id. 5. [MR](#) [Zbl](#)
- [Univalent Foundations 2013] The Univalent Foundations Program, *Homotopy type theory: univalent foundations of mathematics*, Inst. Advanced Study, Princeton, NJ, 2013. [MR](#) [Zbl](#)

Received: 2016-05-06

Revised: 2017-12-20

Accepted: 2018-05-22

maldi2@vcu.edu

*Department of Mathematics and Applied Mathematics,
Virginia Commonwealth University, Richmond, VA,
United States*

mcclearyaj@vcu.edu

*Department of Mathematics, Colorado State University,
Fort Collins, CO, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

| | | | |
|----------------------|---|------------------------|---|
| Colin Adams | Williams College, USA | Chi-Kwong Li | College of William and Mary, USA |
| Arthur T. Benjamin | Harvey Mudd College, USA | Robert B. Lund | Clemson University, USA |
| Martin Bohner | Missouri U of Science and Technology, USA | Gaven J. Martin | Massey University, New Zealand |
| Nigel Boston | University of Wisconsin, USA | Mary Meyer | Colorado State University, USA |
| Amarjit S. Budhiraja | U of N Carolina, Chapel Hill, USA | Frank Morgan | Williams College, USA |
| Pietro Cerone | La Trobe University, Australia | Mohammad Sal Moslehian | Ferdowsi University of Mashhad, Iran |
| Scott Chapman | Sam Houston State University, USA | Zuhair Nashed | University of Central Florida, USA |
| Joshua N. Cooper | University of South Carolina, USA | Ken Ono | Emory University, USA |
| Jem N. Corcoran | University of Colorado, USA | Yuval Peres | Microsoft Research, USA |
| Toka Diagana | Howard University, USA | Y.-F. S. Pétermann | Université de Genève, Switzerland |
| Michael Dorff | Brigham Young University, USA | Jonathon Peterson | Purdue University, USA |
| Sever S. Dragomir | Victoria University, Australia | Robert J. Plemmons | Wake Forest University, USA |
| Joel Foisy | SUNY Potsdam, USA | Carl B. Pomerance | Dartmouth College, USA |
| Errin W. Fulp | Wake Forest University, USA | Vadim Ponomarenko | San Diego State University, USA |
| Joseph Gallian | University of Minnesota Duluth, USA | Bjorn Poonen | UC Berkeley, USA |
| Stephan R. Garcia | Pomona College, USA | József H. Przytycki | George Washington University, USA |
| Anant Godbole | East Tennessee State University, USA | Richard Rebarber | University of Nebraska, USA |
| Ron Gould | Emory University, USA | Robert W. Robinson | University of Georgia, USA |
| Sat Gupta | U of North Carolina, Greensboro, USA | Javier Rojo | Oregon State University, USA |
| Jim Haglund | University of Pennsylvania, USA | Filip Saidak | U of North Carolina, Greensboro, USA |
| Johnny Henderson | Baylor University, USA | Hari Mohan Srivastava | University of Victoria, Canada |
| Glenn H. Hurlbert | Arizona State University, USA | Andrew J. Sterge | Honorary Editor |
| Charles R. Johnson | College of William and Mary, USA | Ann Trenk | Wellesley College, USA |
| K. B. Kulasekera | Clemson University, USA | Ravi Vakil | Stanford University, USA |
| Gerry Ladas | University of Rhode Island, USA | Antonia Vecchio | Consiglio Nazionale delle Ricerche, Italy |
| David Larson | Texas A&M University, USA | John C. Wierman | Johns Hopkins University, USA |
| Suzanne Lenhart | University of Tennessee, USA | Michael E. Zieve | University of Michigan, USA |

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019

vol. 12

no. 3

| | |
|---|-----|
| Darboux calculus | 361 |
| MARCO ALDI AND ALEXANDER MCCLEARY | |
| A countable space with an uncountable fundamental group | 381 |
| JEREMY BRAZAS AND LUIS MATOS | |
| Toeplitz subshifts with trivial centralizers and positive entropy | 395 |
| KOSTYA MEDYNETS AND JAMES P. TALISSE | |
| Associated primes of h -wheels | 411 |
| COREY BROOKE, MOLLY HOCH, SABRINA LATO, JANET STRIULI AND BRYAN WANG | |
| An elliptic curve analogue to the Fermat numbers | 427 |
| SKYE BINEGAR, RANDY DOMINICK, MEAGAN KENNEY, JEREMY ROUSE AND ALEX WALSH | |
| Nilpotent orbits for Borel subgroups of $SO_5(k)$ | 451 |
| MADELEINE BURKHART AND DAVID VELLA | |
| Homophonic quotients of linguistic free groups: German, Korean, and Turkish | 463 |
| HERBERT GANGL, GIZEM KARAALI AND WOOHYUNG LEE | |
| Effective moments of Dirichlet L -functions in Galois orbits | 475 |
| RIZWANUR KHAN, RUOYUN LEI AND DJORDJE MILIĆEVIĆ | |
| On the preservation of properties by piecewise affine maps of locally compact groups | 491 |
| SERINA CAMUNGOL, MATTHEW MORISON, SKYLAR NICOL AND ROSS STOKKE | |
| Bin decompositions | 503 |
| DANIEL GOTSHALL, PAMELA E. HARRIS, DAWN NELSON, MARIA D. VEGA AND CAMERON VOIGT | |
| Rigidity of Ulam sets and sequences | 521 |
| JOSHUA HINMAN, BORYS KUCA, ALEXANDER SCHLESINGER AND ARSENIY SHEYDVASSER | |