

involve

a journal of mathematics

Orbigraphs: a graph-theoretic analog to Riemannian orbifolds

Kathleen Daly, Colin Gavin, Gabriel Montes de Oca,
Diana Ochoa, Elizabeth Stanhope and Sam Stewart



Orbigraphs: a graph-theoretic analog to Riemannian orbifolds

Kathleen Daly, Colin Gavin, Gabriel Montes de Oca,
Diana Ochoa, Elizabeth Stanhope and Sam Stewart

(Communicated by Kenneth S. Berenhaut)

A Riemannian orbifold is a mildly singular generalization of a Riemannian manifold that is locally modeled on \mathbb{R}^n modulo the action of a finite group. Orbifolds have proven interesting in a variety of settings. Spectral geometers have examined the link between the Laplace spectrum of an orbifold and the singularities of the orbifold. One open question in this field is whether or not a singular orbifold and a manifold can be Laplace isospectral. Motivated by the connection between spectral geometry and spectral graph theory, we define a graph-theoretic analog of an orbifold called an orbigraph. We obtain results about the relationship between an orbigraph and the spectrum of its adjacency matrix. We prove that the number of singular vertices present in an orbigraph is bounded above and below by spectrally determined quantities, and show that an orbigraph with a singular point and a regular graph cannot be cospectral. We also provide a lower bound on the Cheeger constant of an orbigraph.

1. Introduction

A Riemannian orbifold is a mildly singular generalization of a Riemannian manifold. A point in an n -dimensional manifold is contained in a neighborhood that is homeomorphic to \mathbb{R}^n . A point in an n -dimensional orbifold is contained in a neighborhood that is homeomorphic to a quotient of \mathbb{R}^n under the action of a finite group. Two useful examples of orbifolds to consider are the \mathbb{Z}_n -football (Figure 1, left) and the \mathbb{Z}_n -teardrop (Figure 1, right):

Example 1. Let \mathbb{Z}_n act on a 2-dimensional sphere by rotations generated by a $2\pi/n$ -radian rotation about an axis passing through the center of the sphere. The quotient of the sphere under this action is the \mathbb{Z}_n -football. Points lying on the intersection of the sphere with the axis of rotation are fixed by all rotations. The images in the \mathbb{Z}_n -football of these points are the conical points at the north and

MSC2010: primary 05C50, 05C20; secondary 60J10.

Keywords: graph spectrum, regular graph, directed graph, orbifold.




Figure 1. Left: football obtained by 180-degree rotation of sphere.
Right: teardrop orbifold.

south poles of the football. If the local lift of a point in an orbifold has nontrivial isotropy, the point is called a *singular point* in the orbifold. The singular set of the \mathbb{Z}_n -football consists of the cone points at its north and south poles.

Example 2. The \mathbb{Z}_n -teardrop is topologically a 2-sphere except for a single point whose neighborhood is locally modeled on the cone $\mathbb{R}^2/\mathbb{Z}_n$, where \mathbb{Z}_n acts by rotations around a fixed point. Thus the \mathbb{Z}_n -teardrop's singular set consists of the isolated cone point. Thurston [1979] showed that unlike the \mathbb{Z}_n -football, the \mathbb{Z}_n -teardrop cannot be obtained as the quotient of a manifold under a smooth, discrete group action.

Introduced by Satake [1956] under the name *V-manifold*, and later renamed and studied as orbifolds by Thurston [1979], orbifolds have proven interesting in a variety of settings; see [Adem et al. 2007; Gordon 2012; Hodgson and Tysk 1993], for example. Of particular interest are results relating the eigenvalue spectrum of the Laplace operator on a Riemannian orbifold (an orbifold endowed with a suitably invariant Riemannian metric) to the singular set of the orbifold. For example, in the presence of a curvature hypothesis, one of us [Stanhope 2005] showed that the Laplace spectrum constrains the structure of the singular set. One fundamental orbifold spectral geometry question that remains open is whether or not the Laplace spectrum actually detects the presence of singular points.

Brooks [1991; 1999] proposes viewing k -regular graphs as combinatorial analogs of smooth manifolds. The infinite k -regular tree T_k is viewed as the graph-theoretic version of the universal cover of a finite k -regular graph. A finite k -regular graph Γ is studied as the quotient of T_k by the fundamental group of Γ in analogy to the study of quotients of the universal cover of a manifold under the action of a discrete cocompact group of isometries acting freely. In this setting Brooks obtains several results including a characterization of Ramanujan graphs, a partial converse to Sunada's theorem, and links between the spectrum of a k -regular graph and the graph's diameter and girth.

Following Brooks' analogy, observe that the action of a discrete, cocompact group of isometries which is not free yields a quotient space that is an orbifold rather than a manifold. Given the successful examination of orbifolds from the




Figure 2. Left: a small 3-orbigraph. Right: a 3-orbigraph with 7 vertices.

perspective of spectral geometry, we seek to extend Brooks’ analogy one step further by first proposing a graph-theoretic analog of an orbifold and, second, applying the lens of spectral graph theory to orbifold graphs. References in the literature to an orbifold-like class of graphs are limited. Brooks [1999] himself describes an “orbifold graph” as a quotient of a k -regular graph under a nonfree group action. He offers orbifold graphs as a motivating idea, but chooses to “avoid entering into the technicalities of ‘orbifold graphs’.” Juan-Pineda, Lafont, Millan-Vossler and Pallekonda [Juan-Pineda et al. 2011] describe an analogy between orbifolds and objects from Bass–Serre theory [Bass 1993] called *graphs-of-groups*. Although the present work has its roots in the ideas of Brooks, the graphs that we examine here can be viewed as a generalization of the edge-index graph of a graph-of-groups.

We define an *orbigraph* to be a member of the following class of weighted, directed graphs.

Definition 3. An *orbigraph of degree k (k -orbigraph)* is a finite, weighted, directed graph Ω where the adjacency matrix A of Ω satisfies the following:

- (i) $A_{ij} \in \mathbb{Z}_{\geq 0}$.
- (ii) $\sum_j A_{ij} = k$.
- (iii) $A_{ij} > 0$ if and only if $A_{ji} > 0$.

Figure 2 shows two examples of orbigraphs.

Remark 4. All orbigraphs discussed below will be assumed to be connected unless noted otherwise. Condition (iii) in Definition 3 implies that a connected orbigraph must be strongly connected. Nonzero diagonal entries in the adjacency matrix of an orbigraph correspond to weighted loops in the orbigraph.

In Section 2 below we demonstrate the analogy between orbigraphs and orbifolds through the following three points:

- (a) The local structure of a vertex in a k -orbigraph is that of the quotient of a k -regular graph, just as the local structure of a k -dimensional orbifold is the quotient of a k -dimensional manifold.

(b) Some vertices in an orbigraph have the same local structure as a vertex in a regular graph and some do not. This leads us to the definition of regular and singular vertices in an orbigraph—an essential piece of the analogy between orbifolds and orbigraphs.

(c) We show that some orbigraphs can be obtained as the quotient of a finite regular graph under an equitable partition and some cannot. This mirrors the fundamental fact from the geometric setting that orbifolds are divided into two classes: those that are covered by a manifold (like the football) and those that are not (like the teardrop). Indeed, the presence of singular objects that are not merely quotients of regular objects saves the study of orbifolds and orbigraphs from being simply a reduced version of a known field of study.

[Section 3](#) connects orbigraphs to the theory of Markov chains. In [Section 4](#) Markov chain methods are used to obtain a graph-theoretic characterization of when an orbigraph can be obtained as the quotient of a finite regular graph, and when it cannot. This characterization makes it easy to generate examples of orbigraphs with these properties, facilitating our later examination of how spectral results for orbifolds carry over to the orbigraph setting. Also using Markov chain methods we provide a lower bound on the Cheeger constant of a k -orbigraph in terms of k and the size of its vertex set. This adds a third family to the list in [\[Chung 2005\]](#) of families of directed graphs that satisfy similar bounds. It would be interesting to know if the bound presented here is sharp, or if an improved bound could be used to obtain a strong upper bound on the convergence of random walks on orbigraphs. Our examination of the Cheeger constant on orbigraphs is the topic of [Section 5](#).

In [Section 6](#) we follow the philosophy of Brooks and ask questions from the spectral geometry of orbifolds in the orbigraph setting. The orbigraph spectrum discussed here is the list of eigenvalues of the adjacency matrix of an orbigraph. Because the analogy between orbifolds and orbigraphs established in [Section 2](#) is strong, the questions carry over naturally and we obtain several interesting results:

(a) We show that the spectrum does not detect whether or not an orbigraph can be obtained as the quotient of a finite k -regular graph. The analogous question for orbifolds is still an open problem in spectral geometry.

(b) The number of singular points in an orbigraph can be bounded both above and below by spectrally determined quantities. In the geometric setting one can seek spectral bounds on the number of components of the singular set. In dimension 2, the fifth author and Proctor [\[Proctor and Stanhope 2010\]](#) obtained a result of this type under a curvature hypothesis.

(c) The spectrum of an orbigraph detects the presence of singular points. As mentioned above, this question is still open in the orbifold setting.

2. Orbigraphs as discrete orbifolds

2.1. Local structure of a k -orbigraph. The local structure of an orbigraph is that of a quotient of a k -regular graph. There are multiple ways to define the quotienting process for graphs. Here quotient graphs will be formed with respect to an equitable partition. The definition given below uses the approach of Barrett, Francis and Webb [Barrett et al. 2017] to extend the definition of an equitable partition from the familiar setting of simple graphs to the more general setting of weighted directed graphs. We also follow the thorough treatment of the simple graph case in Chapter 5 of [Godsil 1993].

In what follows let $w(u, v)$ denote the weight of directed edge (u, v) .

Definition 5. Let Γ be a graph (possibly directed, weighted, or both) and

$$\mathcal{P} = \{V_1, V_2, \dots, V_m\}$$

be a partition of its vertices:

- (a) We say \mathcal{P} is an *equitable partition* if for all pairs i, j the number $\sum_{v \in V_j} w(u, v)$ is the same for each element u in V_i .
- (b) Given an equitable partition \mathcal{P} on Γ , the weighted directed graph with adjacency matrix $A_{ij} = \sum_{v \in V_j} w(u, v)$, u in V_i , is called the *quotient graph* of Γ with respect to \mathcal{P} and will be denoted by Γ/\mathcal{P} .

Remark 6. If a group G acts on a simple graph Γ by automorphisms, the vertex orbits of the action form an equitable partition of the vertex set of Γ . This type of equitable partition is called an *orbit partition*. In this case the quotient graph will be written Γ/G .

To discuss the local structure of an orbigraph we introduce further terms from graph theory. Note that an undirected edge $\{v, w\}$ of weight n in a graph will be viewed as being equivalent to a pair of weight- n directed edges (v, w) and (w, v) , and vice versa.

- Definition 7.**
- (a) The *k -star graph* is the complete bipartite graph $K_{1,k}$ and will be denoted by S_k . The vertex with degree k in S_k is the *central vertex* of S_k .
 - (b) The *neighborhood* of a vertex v in an undirected graph Γ is the subgraph of Γ including the vertex v , all vertices w adjacent to v , and all edges $\{v, w\}$.
 - (c) The *out-neighborhood* of a vertex v in a directed graph Δ is the directed subgraph of Δ including vertex v , all vertices w at which edges initiating at v terminate, and all directed edges (v, w) with initial vertex v .

Because the neighborhood of each vertex in a simple k -regular graph is S_k , we view a simple k -regular graph as the graph-theoretic analog of a k -dimensional manifold.




Figure 3. Out-neighborhoods of the central vertex in quotients of S_3 .

Let G be a group of graph automorphisms of S_k and form the quotient graph S_k/G . The central vertex c of S_k/G is the vertex in S_k/G associated to the element of the orbit partition on S_k containing the central vertex of S_k . The out-neighborhood of c in S_k/G is a weighted star graph with between 1 and k edges. The sum of the weights over all edges in the out-neighborhood of c is k .

Example 8. There are only three different weighted, directed graphs that arise as quotients of S_3 by a group of graph automorphisms. Figure 3 illustrates the out-neighborhoods of the central vertex in each of these three quotients.

Because all row sums in the adjacency matrix of a k -orbigraph Ω are k , the out-neighborhood of a vertex v in Ω is identical to the outgoing neighborhood of the central vertex in some quotient of a k -star. In this way, a k -star quotient provides the local model of the neighborhood of a point in an orbigraph. Our interest in the local structure of an orbigraph at a vertex is in the number of outgoing edges and the weights of those edges. The terminal point of an outgoing edge is not important. Because of this the out-neighborhood of a vertex with a loop is taken with the loop “undone”. For example, vertex v_1 in Figure 2, left, is locally modeled on the middle graph in Figure 3.

To complete our analogy between the local structure of orbifolds and the local structure of orbigraphs we observe that requirement (iii) in Definition 3 corresponds to the fact that if local neighborhoods U, V in an orbifold satisfy $U \cap V \neq \emptyset$ then we also have $V \cap U \neq \emptyset$.

2.2. Singular points in an orbigraph. The key feature of the study of orbifolds that distinguishes it from manifold theory is the presence of orbifold singular points. We define a singular vertex in an orbigraph in the following way.

Definition 9. A vertex v of an orbigraph is *singular* if any outgoing edge from v has weight greater than 1. A vertex that is not singular is called *regular*.

We see that regular graphs contain no singular vertices, as required by our analogy between regular graphs and manifolds.

Example 10. Both vertices in the orbigraph in Figure 2, left, are singular. Vertices v_1, v_4 and v_6 in the orbigraph in Figure 2, right, are singular, and the rest are regular.




Figure 4. Graph diagram of K_4 .

In contrast to the orbifold setting, singular points in an orbigraph are not marked with an isotropy group. However we can quantify the extent to which a vertex v is singular by noting the number of outgoing edges from v that have weight greater than 1. We can also consider the list of weights of outgoing edges from v . As mentioned in the [Introduction](#), graphs-of-groups offer an alternative graph-theoretic interpretation of orbifolds. A graph-of-groups, in contrast to an orbigraph, has vertices that are marked with a group in a way that is analogous to an orbifold isotropy group.

2.3. Good and bad orbigraphs. In [Example 1](#) we saw that the football orbifold is the quotient of a sphere under the smooth action of a finite group. In [Example 2](#) it was asserted that the teardrop orbifold cannot be obtained as a quotient in this manner. Orbifolds that can be written as the quotient of a manifold under a smooth, discrete group action are called *good*. Otherwise they are called *bad*. Following these ideas we define *good* and *bad* orbigraphs as follows.

Definition 11. A k -orbigraph Ω is said to be *good* if it can be obtained as the quotient of a finite k -regular graph Γ via an equitable partition on Γ . If an orbigraph is not good it is called *bad*.

Example 12. The orbigraph in [Figure 2](#), left, is good because it is the quotient of the complete graph K_4 , as presented in [Figure 4](#), by the group \mathbb{Z}_3 generated by a $2\pi/3$ -radian rotation about the center vertex. The orbigraph in [Figure 2](#), right, is bad. This follows from [Theorem 20](#) below and the observation that the product of edge weights along cycle $(v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_1)$ is 2, while the product of edge weights along the reverse cycle $(v_1, v_7, v_6, v_5, v_4, v_3, v_2, v_1)$ is 4.

The analogy with the covering theory of topological spaces is further strengthened by the following two lemmas.

Lemma 13. *If Ω is a k -orbigraph and \mathcal{P} is an equitable partition on the vertices of Ω , then Ω/\mathcal{P} is a k -orbigraph.*

Proof. Let A denote the adjacency matrix of Ω/\mathcal{P} , where $\mathcal{P} = \{V_1, V_2, \dots, V_m\}$, and let $w_\Omega(\cdot, \cdot)$ denote the weight function on directed edges in Ω . Because Ω is an orbigraph, we know $w_\Omega(u, v)$ is a nonnegative integer for all vertices u, v

in Ω . Hence $A_{ij} = \sum_{v \in V_j} w_\Omega(u, v)$, for any $u \in V_i$, is a nonnegative integer. Fixing $i \in \{1, 2, \dots, m\}$, and taking u some element of V_i , consider the i -th row sum of A :

$$\sum_j A_{ij} = \sum_j \sum_{v \in V_j} w_\Omega(u, v) = \sum_{v \in \Omega} w_\Omega(u, v) = k.$$

Finally suppose $A_{ij} > 0$. Then there must a $j \in \{1, 2, \dots, m\}$ for which any $u \in V_i$ has $w_\Omega(u, v) > 0$ for some $v \in V_j$. Because Ω is an orbigraph, we must also have $w_\Omega(v, u) > 0$. Thus $A_{ji} > 0$. \square

Definition 14. We say that an orbigraph Ω_1 covers an orbigraph Ω_2 if there is an equitable partition \mathcal{P} of the vertices of Ω_1 such that $\Omega_1/\mathcal{P} = \Omega_2$.

Lemma 15. *The covering relation is transitive.*

Proof. Suppose Ω_1 is an orbigraph with equitable partition \mathcal{P}_1 such that $\Omega_1/\mathcal{P}_1 = \Omega_2$, and Ω_2 has an equitable partition \mathcal{P}_2 such that $\Omega_2/\mathcal{P}_2 = \Omega_3$. We need to show there is an equitable partition \mathcal{P}_3 of Ω_1 such that $\Omega_1/\mathcal{P}_3 = \Omega_3$. For $i = 1, 2$ let A_i denote the adjacency matrix of orbigraph Ω_i , and P_i denote the characteristic matrix corresponding to partition \mathcal{P}_i . By a straightforward modification of [Godsil 1993, Lemma 2.1, p. 77] to the setting of weighted, directed graphs we have that $A_1 P_1 = P_1 A_2$ and $A_2 P_2 = P_2 A_3$. Thus $A_1 P_1 P_2 = P_1 A_2 P_2 = P_1 P_2 A_3$. We conclude $P_1 P_2$ defines an equitable partition on Ω_1 with quotient orbigraph Ω_3 . \square

As a consequence of the previous two lemmas we obtain the following.

Corollary 16. *The quotient of any good orbigraph must also be good.*

3. Orbigraphs and Markov chains

The fact that the row sum of the adjacency matrix of an orbigraph is constant provides an immediate connection between orbigraphs and Markov chains. Following [Kelly 1979], we review ideas from the theory of Markov chains and introduce notation that will be used hereafter. Matrix A will denote the adjacency matrix of a k -orbigraph Ω with n vertices. Define $P = (1/k)A$. Matrix P is the transition matrix of a stationary Markov chain, as all entries of P lie in the interval $[0, 1]$ and all rows of P sum to 1. Because the adjacency matrix of a k -orbigraph has right eigenvalue k (to see this consider the eigenvector with all entries equal to 1), P has right eigenvalue 1 and stationary distribution vector $\pi = (\pi_1, \pi_2, \dots, \pi_n)$, with $\sum_{k=1}^n \pi_k = 1$, for which $\pi P = \pi$. By Remark 4 we know Ω is strongly connected so π is the unique stationary distribution of P .

Our first result connecting orbigraphs to Markov chains is a bound on the minimal entry of π in terms of the degree and number of vertices of an orbigraph.

Lemma 17. *Let π_m be a minimal entry in stationary distribution π . Then*

$$\pi_m \geq \frac{1}{nk^{n-1}}.$$

Proof. Let π_M denote a maximal entry in π and let c be the minimal nonzero value that appears as an entry in matrix P . Because Ω is strongly connected, there is a path of length $\ell < n$ from the M -th vertex to the m -th vertex of Ω . This implies that $(P^\ell)_{Mm}$ is nonzero. Using this and the fact that $\pi P = \pi$, we have

$$\pi_m = \sum_{k=1}^n (P^\ell)_{km} \pi_k \geq (P^\ell)_{Mm} \pi_M \geq c^\ell \pi_M \geq c^{n-1} \pi_M.$$

Because P is the transition matrix associated to an orbigraph, we have $c \geq 1/k$. Also, we know that $\pi_M \geq 1/n$ because the sum of the entries of π is 1. Thus $\pi_m \geq c^{n-1} \pi_M \geq 1/(nk^{n-1})$ as required. \square

Here we relate the stationary distribution of a good orbigraph to that of its finite regular cover.

Lemma 18. *Let Γ be a k -regular graph with N vertices, $\mathcal{P} = \{V_1, V_2, \dots, V_n\}$ be an equitable partition of the vertices of Γ , and P be the transition matrix of the orbigraph Γ/\mathcal{P} . Let $|V_i|$ denote the number of vertices in partition element V_i . The stationary distribution of P is the n -tuple π , where $\pi_i = (1/N)|V_i|$.*

Proof. Let Q denote the transition matrix obtained by scaling the adjacency matrix of Γ by $1/k$. The result follows from the observation that the stationary distribution of Q is the N -tuple $(1/N, 1/N, \dots, 1/N)$ and [Godsil 1993, Lemma 2.2, p. 78]. \square

4. Characterizing good and bad orbigraphs

We use the Markov chain methods and notation from Section 3 to provide a quick way to distinguish good orbigraphs from bad orbigraphs.

Definition 19. An orbigraph Ω satisfies the *balanced cycle condition* if the product of the edge weights along each directed cycle $v_1, v_2, \dots, v_l, v_1$ in Ω equals the product of the edge weights along the reverse directed cycle $v_1, v_l, v_{l-1}, \dots, v_1$.

Theorem 20. *An orbigraph is good if and only if it satisfies the balanced cycle condition.*

A stationary Markov chain is said to satisfy the *detailed balance equations* if

$$\pi_i P_{ij} = \pi_j P_{ji} \quad \text{for all } i, j = 1, 2, \dots, n.$$

The Markov chain analog of the balanced cycle condition from Definition 19 is called the *Kolmogorov criterion*. In particular, an orbigraph satisfies the balanced cycle condition if and only if the corresponding Markov chain satisfies the Kolmogorov criterion. We can now state a needed lemma.

Lemma 21. *A stationary Markov chain satisfies the detailed balance equations if and only if it satisfies the Kolmogorov criterion.*

Proof. This follows from combining Theorems 1.2 and 1.7 in [Kelly 1979]. \square

Proof of Theorem 20. Suppose Ω is a good orbigraph. This implies $\Omega = \Gamma/\mathcal{P}$, where Γ is a k -regular graph and $\mathcal{P} = \{V_1, V_2, \dots, V_n\}$ is an equitable partition on Γ . Scaling the adjacency matrix of Γ by $1/k$ yields the symmetric transition matrix Q of a Markov chain. We relate the stationary distribution of Q to the stationary distribution of P , the transition matrix of Ω , by Lemma 18. In particular $\pi_i = (1/N)|V_i|$, where π denotes the stationary distribution of P and N is the number of vertices in Γ .

The following computation confirms that P satisfies the detailed balance equations:

$$\begin{aligned} \pi_j P_{ji} &= \frac{1}{N} |V_j| P_{ji} = \frac{1}{N} |V_j| \sum_{k \in V_i} Q_{jk} = \frac{1}{N} \sum_{l \in V_j} \sum_{k \in V_i} Q_{lk} \\ &= \frac{1}{N} \sum_{k \in V_i} \sum_{l \in V_j} Q_{kl} = \frac{1}{N} |V_i| \sum_{l \in V_j} Q_{il} = \pi_i P_{ij}. \end{aligned}$$

(The argument closely follows that of [Tian and Kannan 2006, Theorem 2.16], which is given in the setting of lumpable Markov chains. It makes essential use of the fact that \mathcal{P} is an equitable partition and that Q is a symmetric matrix.) The fact that Ω satisfies the balanced cycle condition now follows from Lemma 21.

Now suppose Ω is an orbigraph that satisfies the balanced cycle condition. By Lemma 21, P and π satisfy the detailed balance equations $\pi_i P_{ij} = \pi_j P_{ji}$. Multiplying by k on both sides gives $\pi_i A_{ij} = \pi_j A_{ji}$. Because A has all nonnegative integer entries, π will have all nonnegative rational entries. Thus there is an integer m for which $m\pi = (d_1, d_2, \dots, d_n)$ is a vector of nonnegative integers. This allows us to write

$$d_i A_{ij} = d_j A_{ji}, \tag{1}$$

an equality of products of nonnegative integers.

We now build a finite k -regular cover Γ of Ω . Let X be the set of nonzero, nondiagonal entries of A . Let $Y = \{A_{11} + 1, A_{22} + 1, \dots, A_{nn} + 1\}$. Let c be the least common multiple of the integers in $X \cup Y$. For each $i = 1, 2, \dots, n$ we take V_i to be a set of cd_i vertices. The disjoint union $V_1 \sqcup V_2 \sqcup \dots \sqcup V_n$ forms the vertex set of Γ and gives the needed vertex partition \mathcal{P} of Γ .

It remains to specify adjacency in Γ in such a way that $\Gamma/\mathcal{P} = \Omega$. Suppose $i \neq j$. For the quotient $\Gamma/\mathcal{P} = \Omega$ to be valid, each vertex in V_i must be adjacent to A_{ij} vertices in V_j , and each vertex in V_j must be adjacent to A_{ji} vertices in V_i . Thus the number of edges with one vertex in V_i and one vertex in V_j , which we will denote by $e_{\{i,j\}}$, is simultaneously $A_{ij}|V_i|$ and $A_{ji}|V_j|$. The adapted detailed

balance equations from (1) show that this requirement follows from our choice for the sizes of V_i and V_j as

$$A_{ij}|V_i| = A_{ij}cd_i = A_{ji}cd_j = A_{ji}|V_j|.$$

Because A_{ij} divides $|V_j|$ and A_{ji} divides $|V_i|$, we can distribute the $e_{\{i,j\}}$ edges connecting V_i and V_j with exactly A_{ij} edges adjacent to each vertex in V_i and exactly A_{ji} edges adjacent to each vertex in V_j . Because $A_{ii} + 1$ divides $|V_i|$, we can require that all elements of V_i are adjacent to exactly A_{ii} other elements of V_i . This completes the adjacency relations for Γ .

By construction we observe $\Gamma/\mathcal{P} = \Omega$. The degree of a vertex v in Γ is $\sum_{j=1} A_{ij} = k$; thus Γ is k -regular. Should Γ fail to be connected, any connected component Γ' of Γ will satisfy $\Gamma'/\mathcal{P} = \Omega$. \square

Remark 22. Corollary 16 and Theorem 20 imply that if an orbigraph Ω satisfies the balanced cycle condition then so does any orbigraph quotient of Ω . This stands in contrast to [Tian and Kannan 2006, Example 2.17].

5. Bounding the Cheeger constant of an orbigraph

Chung [2005] defined a Cheeger constant for directed graphs and obtained lower bounds on the Cheeger constant for both regular and Eulerian directed graphs. Using R to denote a k -regular directed graph on n vertices and E an Eulerian directed graph with m edges, Chung showed

$$h(R) \geq \frac{2}{kn} \quad \text{and} \quad h(E) \geq \frac{2}{m}. \quad (2)$$

Here we apply Chung's methods to obtain a lower bound on the Cheeger constant of an orbigraph. We use notation from Section 3.

Define a function F from Ω to the nonnegative real numbers by

$$F(i, j) = \pi_i P_{ij},$$

where i and j are vertices in Ω . This function is an example of a *circulation* on Ω ; see [Chung 2005, Lemma 3.1]. Letting S range over all nonempty proper subsets of the vertex set of Ω , the Cheeger constant $h(\Omega)$ of Ω is defined as

$$h(\Omega) = \inf_S \frac{\sum_{i \in S, j \notin S} F(i, j)}{\min\{\sum_{j \in S} F(j), \sum_{j \in \bar{S}} F(j)\}},$$

where $F(j) = \sum_{i, i \rightarrow j} F(i, j)$ and \bar{S} is the set of vertices of Ω that are not in S .

We have the following lower bound on the Cheeger constant of Ω .

Proposition 23. *Let Ω be a k -orbigraph with n vertices. Then*

$$h(\Omega) \geq \frac{2}{n^2 k n}.$$

Proof. We begin by bounding the numerator in the expression defining the Cheeger constant (let π_m denote a minimal entry in π):

$$\sum_{i \in S, j \notin S} F(i, j) = \sum_{i \in S, j \notin S} \pi_i P_{ij} \geq \sum_{i \in S, j \notin S} \pi_m P_{ij} \geq \frac{1}{nk^n}.$$

The last inequality follows from [Lemma 17](#) and the observation that the smallest possible nonzero value for an entry in P is $1/k$.

To bound the denominator first observe that $\sum_{j \in S} F(j)$ is no greater than the sum of the columns in P associated to the vertices in S . It is similar for $\sum_{j \in \bar{S}} F(j)$. Since the total sum of the entries in P is n , we have

$$\sum_{j \in S} F(j) + \sum_{j \in \bar{S}} F(j) \leq n.$$

Thus $\min \left\{ \sum_{v \in S} F(v), \sum_{v \in \bar{S}} F(v) \right\} \leq n/2$.

We see that for any choice of S the quotient in the definition of the Cheeger constant must be greater than or equal to $2/(n^2 k^n)$, completing the proof. \square

Remark 24. Chung uses the inequalities in [\(2\)](#) to obtain convergence bounds for a type of random walk on regular and Eulerian directed graphs. The presence of n in the exponent in the denominator of the orbigraph bound makes it too weak to obtain a similar orbigraph result. It would be interesting to see if a better bound on the Cheeger constant of an orbigraph, should one exist, would allow a convergence result similar to the regular and Eulerian cases.

6. Spectral results for orbigraphs

Because different matrices can be associated to a given graph, a variety of graph spectra are examined in spectral graph theory. Here the *spectrum* of an orbigraph Ω is defined to be the list of eigenvalues of the adjacency matrix of Ω with each eigenvalue repeated according to its multiplicity. We will write the spectrum of an orbigraph with n vertices as a multiset $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$. The study of the spectral properties of directed graphs is relatively new and has yielded interesting applications, as well as directed graph analogs of familiar graph-theoretical results, including Cheeger's inequality; see [\[Chung 2005; Langville and Meyer 2006\]](#), for example. We focus on developing results that relate the spectrum of an orbigraph to its orbigraph structure.

Remark 25. Just as with k -regular graphs, the spectral radius of a k -orbigraph is k . In addition, the number of eigenvalues in the spectrum of an orbigraph (counting multiplicity) is equal to the number of vertices in the orbigraph.

Lemma 26. *Suppose orbigraph Ω_1 covers orbigraph Ω_2 . Then the spectrum of Ω_2 is contained in the spectrum of Ω_1 as multisets.*




Figure 5. The left and center orbigraphs are cospectral. The left orbigraph is bad. The center orbigraph is good as it is covered by the right-most graph using the indicated partition.

Proof. This follows from the argument in Lemma 2.2 of Chapter 5 in [Godsil 1993], adjusted to allow the graph carrying the equitable partition to be a weighted, directed graph. \square

Corollary 27. Any orbigraph with complex eigenvalues must be bad.

Proof. This follows from Lemma 26 and the fact that regular graphs have real eigenvalues. \square

Theorem 28. The spectrum of an orbigraph does not distinguish good orbigraphs from bad orbigraphs.

Proof. The orbigraph on the left in Figure 5 and the orbigraph in the center of the figure both have spectrum $\{-2, 0, 1, 3\}$. However the orbigraph on the left is bad and the orbigraph in the center is good. To see that the left orbigraph is bad, apply Theorem 20 and the fact that the product of the edge weights along cycle (v_1, v_2, v_3, v_4) is not equal to the product of the edge weights of this cycle reversed. The center orbigraph is good because it is covered by the 3-regular graph on the right side of Figure 5 using the indicated equitable partition. \square

In the following lemma a directed edge from vertex v_1 to vertex v_2 of weight w is considered to contribute w -many different ways to move from v_1 to v_2 . The length spectrum of a graph is the finite list of nonnegative integers where the m -th number in the list counts the number of closed walks of length m present in the graph.

Lemma 29. The eigenvalue spectrum of an orbigraph determines and is determined by the length spectrum of the orbigraph.

Proof. Let Ω be a k -orbigraph, A its adjacency matrix, and w_m the number of closed walks in Ω of length m . We know that

$$w_m = \text{tr}(A^m) \quad (3)$$

because the diagonal of A^m counts the number of closed walks of length m . However

$$\text{tr}(A^m) = \sum_{i=1}^n \lambda_i^m.$$

Thus the eigenvalue spectrum of Ω uniquely determines the length spectrum of Ω , and conversely by Newton's identities [Mead 1992] the length spectrum of Ω uniquely determines the eigenvalue spectrum of Ω . \square

We now prove that the number of singular points in an orbigraph is bounded above and below by spectrally determined quantities.

Theorem 30. *Let Ω be a k -orbigraph with n vertices. If s is the number of singular points in Ω , then we have*

$$\frac{\sum_{i=1}^n \lambda_i^2 - nk}{k^2 - k} \leq s \leq \sum_{i=1}^n \lambda_i^2 - nk,$$

where λ_i are the eigenvalues of the adjacency matrix A of Ω .

Proof. First note that $\sum_{i=1}^n \lambda_i^2 = \text{tr}(A^2)$ and by Lemma 29 this quantity counts the number of closed walks of length 2 in Ω . A given vertex v in Ω has outgoing edges with weights summing to k , each of which is matched by at least one incoming edge. This implies the number of closed walks of length 2 starting at v is at least k . Observing that there are n vertices in Ω , we obtain $\text{tr}(A^2) \geq nk$. Now suppose v_1 is a singular vertex in Ω . This vertex has at least one outgoing edge (v_1, v_2) of weight greater than 1. Edge (v_1, v_2) contributes at least one closed walk of length 2, beginning and ending at v_2 , that has not yet been counted. We conclude that $\text{tr}(A^2) \geq nk + s$; thus $s \leq \sum_{i=1}^n \lambda_i^2 - nk$.

For the lower bound, note that each singular vertex v_i contributes $A_{ji}(A_{ij} - 1)$ extra (i.e., beyond the initial k length-2 paths) length-2 paths based at v_j . Thus the total number of extra paths contributed by vertex v_i is $\sum_{v_i \sim v_j} A_{ji}(A_{ij} - 1)$. We bound this quantity in terms of k :

$$\sum_{v_i \sim v_j} A_{ji}(A_{ij} - 1) \leq \sum_{v_i \sim v_j} k(A_{ij} - 1) = k \sum_{v_i \sim v_j} A_{ij} - \sum_{v_i \sim v_j} k \leq k^2 - k.$$

Hence each singular vertex contributes at most $k^2 - k$ extra walks of length 2, so $s(k^2 - k) \geq \sum_{i=1}^n \lambda_i^2 - nk$. Isolating s in this inequality completes the proof. \square

Remark 31. The orbigraph with adjacency matrix kI_n , where I_n denotes the $n \times n$ identity matrix, achieves the lower bound in Theorem 30 for all choices of k and n . Thus this lower bound is sharp in k and n .

Corollary 32. *Suppose Ω is a k -orbigraph with n vertices. Then Ω is isomorphic to a k -regular graph if and only if*

$$\sum_i \lambda_i^2 - nk = 0 \quad \text{and} \quad \sum_i \lambda_i = 0.$$

Proof. A simple k -regular graph Ω has no self loops; thus [Lemma 29](#) implies $\sum_i \lambda_i = 0$. Viewing each edge $\{v_i, v_j\}$ in Ω as two directed edges, (v_i, v_j) and (v_j, v_i) , we see each vertex in Ω has exactly k closed walks of length 2. Therefore $\sum_i \lambda_i^2 = nk$.

Conversely, assume that Ω is an orbigraph such that $\sum_i \lambda_i^2 = nk$ and $\sum_i \lambda_i = 0$. Then by [Theorem 30](#), we have $s \leq 0$. As $s \geq 0$ we see $s = 0$. Thus the outgoing edges of each vertex in Ω all have weight 1. The second condition implies Ω has no loops. By combining pairs of directed edges (v_i, v_j) and (v_j, v_i) into a single undirected edge $\{v_i, v_j\}$, we obtain a simple k -regular graph. \square

In the smooth setting it is not known if a manifold can have the same Laplace spectrum as a nonmanifold orbifold. We can resolve this question in the setting of orbigraphs.

Corollary 33. *A regular graph and an orbigraph with one or more singular points cannot be cospectral.*

Proof. Suppose regular graph Γ and orbigraph Ω are cospectral and that Ω contains $s \geq 1$ singular points. By [Remark 25](#) the largest eigenvalue in the shared spectrum of Γ and Ω is the degree of regularity of each graph. Denote this largest eigenvalue by k . In addition the shared spectrum implies that each graph has the same number of vertices n . By the forward direction of [Corollary 32](#), the fact that Γ is k -regular implies $\sum_i \lambda_i^2 - nk = 0$ and $\sum_i \lambda_i = 0$. However the backwards direction of [Corollary 32](#) implies $s = 0$, a contradiction. \square

Acknowledgements

This work was supported in part by the John S. Rogers Science Research Program and the Early Research Program at Lewis & Clark College. The authors would like to thank Omar Lopez and Yung-Pin Chen for foundational work with orbigraphs and discussions of Markov processes, respectively. We also thank the reviewer for helpful suggestions.

References

- [Adem et al. 2007] A. Adem, J. Leida, and Y. Ruan, *Orbifolds and stringy topology*, Cambridge Tracts in Math. **171**, Cambridge Univ. Press, 2007. [MR](#) [Zbl](#)
- [Barrett et al. 2017] W. Barrett, A. Francis, and B. Webb, “Equitable decompositions of graphs with symmetries”, *Linear Algebra Appl.* **513** (2017), 409–434. [MR](#) [Zbl](#)

- [Bass 1993] H. Bass, “Covering theory for graphs of groups”, *J. Pure Appl. Algebra* **89**:1-2 (1993), 3–47. [MR](#) [Zbl](#)
- [Brooks 1991] R. Brooks, “The spectral geometry of k -regular graphs”, *J. Anal. Math.* **57** (1991), 120–151. [MR](#) [Zbl](#)
- [Brooks 1999] R. Brooks, “Non-Sunada graphs”, *Ann. Inst. Fourier (Grenoble)* **49**:2 (1999), 707–725. [MR](#) [Zbl](#)
- [Chung 2005] F. Chung, “Laplacians and the Cheeger inequality for directed graphs”, *Ann. Comb.* **9**:1 (2005), 1–19. [MR](#) [Zbl](#)
- [Godsil 1993] C. D. Godsil, *Algebraic combinatorics*, Chapman & Hall, New York, 1993. [MR](#) [Zbl](#)
- [Gordon 2012] C. Gordon, “Orbifolds and their spectra”, pp. 49–71 in *Spectral geometry* (Hanover, NH, 2010), edited by A. H. Barnett et al., Proc. Sympos. Pure Math. **84**, Amer. Math. Soc., Providence, RI, 2012. [MR](#) [Zbl](#)
- [Hodgson and Tysk 1993] C. Hodgson and J. Tysk, “Eigenvalue estimates and isoperimetric inequalities for cone-manifolds”, *Bull. Austral. Math. Soc.* **47**:1 (1993), 127–143. [MR](#) [Zbl](#)
- [Juan-Pineda et al. 2011] D. Juan-Pineda, J.-F. Lafont, S. Millan-Vossler, and S. Pallekonda, “Algebraic K -theory of virtually free groups”, *Proc. Roy. Soc. Edinburgh Sect. A* **141**:6 (2011), 1295–1316. [MR](#) [Zbl](#)
- [Kelly 1979] F. P. Kelly, *Reversibility and stochastic networks*, Wiley, Chichester, UK, 1979. [MR](#) [Zbl](#)
- [Langville and Meyer 2006] A. N. Langville and C. D. Meyer, *Google’s PageRank and beyond: the science of search engine rankings*, Princeton Univ. Press, 2006. [MR](#) [Zbl](#)
- [Mead 1992] D. G. Mead, “Newton’s identities”, *Amer. Math. Monthly* **99**:8 (1992), 749–751. [MR](#) [Zbl](#)
- [Proctor and Stanhope 2010] E. Proctor and E. Stanhope, “Spectral and geometric bounds on 2-orbifold diffeomorphism type”, *Differential Geom. Appl.* **28**:1 (2010), 12–18. [MR](#) [Zbl](#)
- [Satake 1956] I. Satake, “On a generalization of the notion of manifold”, *Proc. Nat. Acad. Sci. USA* **42** (1956), 359–363. [MR](#) [Zbl](#)
- [Stanhope 2005] E. Stanhope, “Spectral bounds on orbifold isotropy”, *Ann. Global Anal. Geom.* **27**:4 (2005), 355–375. [MR](#) [Zbl](#)
- [Thurston 1979] W. P. Thurston, “The geometry and topology of three-manifolds”, lecture notes, Princeton Univ., 1979, available at <http://www.msri.org/publications/books/gt3m>.
- [Tian and Kannan 2006] J. P. Tian and D. Kannan, “Lumpability and commutativity of Markov processes”, *Stoch. Anal. Appl.* **24**:3 (2006), 685–702. [MR](#) [Zbl](#)

Received: 2017-09-02

Revised: 2018-11-10

Accepted: 2019-01-01

daly_kathleen@bah.com

Booz Allen Hamilton, Beaver Creek, OH, United States

cgvavin@908devices.com

908 Devices, Campbell, CA, United States

gabem@uoregon.eduDepartment of Mathematics, University of Oregon,
Eugene, OR, United Statesdochoa@lclark.eduDepartment of Mathematical Sciences, Lewis & Clark College,
Portland, OR, United Statesstanhope@lclark.eduDepartment of Mathematical Sciences, Lewis & Clark College,
Portland, OR, United Statessams@umn.eduSchool of Mathematics, University of Minnesota,
Minneapolis, MN, United States

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

| | | | |
|----------------------|---|------------------------|---|
| Colin Adams | Williams College, USA | Chi-Kwong Li | College of William and Mary, USA |
| Arthur T. Benjamin | Harvey Mudd College, USA | Robert B. Lund | Clemson University, USA |
| Martin Bohner | Missouri U of Science and Technology, USA | Gaven J. Martin | Massey University, New Zealand |
| Nigel Boston | University of Wisconsin, USA | Mary Meyer | Colorado State University, USA |
| Amarjit S. Budhiraja | U of N Carolina, Chapel Hill, USA | Frank Morgan | Williams College, USA |
| Pietro Cerone | La Trobe University, Australia | Mohammad Sal Moslehian | Ferdowsi University of Mashhad, Iran |
| Scott Chapman | Sam Houston State University, USA | Zuhair Nashed | University of Central Florida, USA |
| Joshua N. Cooper | University of South Carolina, USA | Ken Ono | Emory University, USA |
| Jem N. Corcoran | University of Colorado, USA | Yuval Peres | Microsoft Research, USA |
| Toka Diagana | Howard University, USA | Y.-F. S. Pétermann | Université de Genève, Switzerland |
| Michael Dorff | Brigham Young University, USA | Jonathon Peterson | Purdue University, USA |
| Sever S. Dragomir | Victoria University, Australia | Robert J. Plemmons | Wake Forest University, USA |
| Joel Foisy | SUNY Potsdam, USA | Carl B. Pomerance | Dartmouth College, USA |
| Errin W. Fulp | Wake Forest University, USA | Vadim Ponomarenko | San Diego State University, USA |
| Joseph Gallian | University of Minnesota Duluth, USA | Bjorn Poonen | UC Berkeley, USA |
| Stephan R. Garcia | Pomona College, USA | József H. Przytycki | George Washington University, USA |
| Anant Godbole | East Tennessee State University, USA | Richard Rebarber | University of Nebraska, USA |
| Ron Gould | Emory University, USA | Robert W. Robinson | University of Georgia, USA |
| Sat Gupta | U of North Carolina, Greensboro, USA | Javier Rojo | Oregon State University, USA |
| Jim Haglund | University of Pennsylvania, USA | Filip Saidak | U of North Carolina, Greensboro, USA |
| Johnny Henderson | Baylor University, USA | Hari Mohan Srivastava | University of Victoria, Canada |
| Glenn H. Hurlbert | Arizona State University, USA | Andrew J. Sterge | Honorary Editor |
| Charles R. Johnson | College of William and Mary, USA | Ann Trenk | Wellesley College, USA |
| K. B. Kulasekera | Clemson University, USA | Ravi Vakil | Stanford University, USA |
| Gerry Ladas | University of Rhode Island, USA | Antonia Vecchio | Consiglio Nazionale delle Ricerche, Italy |
| David Larson | Texas A&M University, USA | John C. Wierman | Johns Hopkins University, USA |
| Suzanne Lenhart | University of Tennessee, USA | Michael E. Zieve | University of Michigan, USA |

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019

vol. 12

no. 5

| | |
|---|-----|
| Orbigraphs: a graph-theoretic analog to Riemannian orbifolds | 721 |
| KATHLEEN DALY, COLIN GAVIN, GABRIEL MONTES DE OCA, DIANA OCHOA, ELIZABETH STANHOPE AND SAM STEWART | |
| Sparse neural codes and convexity | 737 |
| R. AMZI JEFFS, MOHAMED OMAR, NATCHANON SUAYSOM, ALEINA WACHTEL AND NORA YOUNGS | |
| The number of rational points of hyperelliptic curves over subsets of finite fields | 755 |
| KRISTINA NELSON, JÓZSEF SOLYMOSSI, FOSTER TOM AND CHING WONG | |
| Space-efficient knot mosaics for prime knots with mosaic number 6 | 767 |
| AARON HEAP AND DOUGLAS KNOWLES | |
| Shabat polynomials and monodromy groups of trees uniquely determined by ramification type | 791 |
| NAIOMI CAMERON, MARY KEMP, SUSAN MASLAK, GABRIELLE MELAMED, RICHARD A. MOY, JONATHAN PHAM AND AUSTIN WEI | |
| On some edge Folkman numbers, small and large | 813 |
| JENNY M. KAUFMANN, HENRY J. WICKUS AND STANISŁAW P. RADZISZOWSKI | |
| Weighted persistent homology | 823 |
| GREGORY BELL, AUSTIN LAWSON, JOSHUA MARTIN, JAMES RUDZINSKI AND CLIFFORD SMYTH | |
| Leibniz algebras with low-dimensional maximal Lie quotients | 839 |
| WILLIAM J. COOK, JOHN HALL, VICKY W. KLIMA AND CARTER MURRAY | |
| Spectra of Kohn Laplacians on spheres | 855 |
| JOHN AHN, MOHIT BANSIL, GARRETT BROWN, EMILEE CARDIN AND YUNUS E. ZEYTUNCU | |
| Pairwise compatibility graphs: complete characterization for wheels | 871 |
| MATTHEW BEAUDOUIN-LAFON, SERENA CHEN, NATHANIEL KARST, DENISE SAKAI TROXELL AND XUDONG ZHENG | |
| The financial value of knowing the distribution of stock prices in discrete market models | 883 |
| AYELET AMIRAN, FABRICE BAUDOIN, SKYLYN BROCK, BEREND COSTER, RYAN CRAVER, UGONNA EZEAKA, PHANUEL MARIANO AND MARY WISHART | |