

ON THE GROUP OF CONFORMAL TRANSFORMATIONS OF A COMPACT RIEMANNIAN MANIFOLD. III

CHUAN-CHIH HSIUNG

1. Introduction

Let g_{ij} , R_{hijk} , $R_{ij} = R^k{}_{ijk}$ be respectively the metric, Riemann and Ricci tensors of a Riemannian manifold M^n of dimension n , and denote

$$(1.1) \quad P = R^{hijk}R_{hijk}, \quad Q = R^{ij}R_{ij}.$$

Throughout this paper all Latin indices take the values $1, \dots, n$ unless stated otherwise, and repeated indices imply summation. In a recent paper [2] the author proved

Theorem 1. *Suppose that a compact Riemannian manifold $M^n (n > 2)$ with constant scalar curvature $R = g^{ij}R_{ij}$ admits an infinitesimal nonhomothetic conformal transformation v , and let L_v be the operator of the infinitesimal transformation v . If*

$$(1.2) \quad a^2L_vP + b(2a + nb)L_vQ = \text{const.},$$

where a and b are constants such that

$$(1.3) \quad c \equiv 4a^2 + 2(n-2)ab + n(n-2)b^2 > 0,$$

then M^n is isometric to a sphere.

In particular, when $a = 0$ or $b = 0$, Theorem 1 is reduced to a result of Yano [4], which is a generalization of some results of Lichnerowicz [3] and the author [1]. Yano pointed out that condition (1.3) is equivalent to that a and b are not both zero.

Very recently, Yano and Sawaki [5] obtained the following theorem similar to Theorem 1:

Theorem 2. *Suppose that a compact Riemannian manifold $M^n (n > 2)$ with constant R admits an infinitesimal nonhomothetic conformal transformation v . If*

$$(1.4) \quad L_vL_v[(n-2)a^2P + 4b(2a+b)Q] \leq 0,$$

where a and b are constants such that $a + b \neq 0$, then M^n is isometric to a sphere.

The purpose of this paper is to generalize both Theorems 1 and 2 to

Theorem 3. Suppose that a compact Riemannian manifold M^n ($n > 2$) with constant R admits an infinitesimal nonhomothetic conformal transformation v . If

$$(1.5) \quad L_v L_v \left(a^2 P + \frac{c - 4a^2}{n - 2} Q \right) \leq 0,$$

where

$$(1.6) \quad c \equiv 4a^2 + (n - 2) \left[2a \sum_{i=1}^4 b_i + (b_1 - b_2 + b_3 - b_4 + b_5 - b_6)^2 - 2b_1 b_3 - 2b_2 b_4 + 2b_5 b_6 + (n - 1) \sum_{i=1}^6 b_i^2 \right] > 0,$$

a and b 's being constants, then M^n is isometric to a sphere.

It is obvious that Theorem 3 is reduced to a generalization of Theorem 1 when $b_2 = \dots = b_6 = 0$, and to Theorem 2 when

$$b_1 = \dots = b_4 = b/(n - 2), \quad b_5 = b_6 = 0.$$

We need the following theorem of Yano [5] to prove Theorem 3:

Theorem 4. Suppose that a compact orientable Riemannian manifold M^n ($n > 2$) with constant R admits an infinitesimal nonhomothetic conformal transformation v so that

$$(1.7) \quad L_v g_{ij} = 2\phi g_{ij}, \quad \phi = \text{const.},$$

and let ∇ denote the operator of covariant derivation of M^n with respect to g_{ij} . If

$$(1.8) \quad \int_{M^n} T_{ij} \phi^i \phi^j dA \geq 0,$$

where

$$(1.9) \quad T_{ij} = R_{ij} - \frac{R}{n} g_{ij},$$

$\phi^i = \nabla^i \phi = g^{ij} \nabla_j \phi$, and dA is the element of area of M^n at a point, then M^n is isometric to a sphere.

2. Lemmas

Throughout this section M^n will always denote a compact orientable Riemannian manifold of dimension $n > 2$. Let Δ be the Laplace-Beltrami operator on M^n . Then, for any scalar field f on M^n ,

$$(2.1) \quad \Delta f = -\nabla^i \nabla_j f.$$

Thus we have

$$(2.2) \quad \int_{M^n} \Delta f \, dA = 0$$

from the well-known Green's formula :

$$(2.3) \quad \int_{M^n} \nabla^i \xi_i \, dA = 0,$$

where ξ_i is any vector field on M^n .

Lemma 1. *If a nonconstant scalar field ϕ on a manifold M^n satisfies $\Delta\phi = k\phi$, where k is constant, then k is positive.*

Proof. From equations (2.1), (2.2) we obtain

$$(2.4) \quad \begin{aligned} 0 &= \int_{M^n} \Delta(\phi^2) \, dA = 2 \int_{M^n} (\phi \Delta\phi - \phi^i \phi_i) \, dA \\ &= 2 \int_{M^n} (k\phi^2 - \phi^i \phi_i) \, dA, \end{aligned}$$

which gives Lemma 1 immediately.

Lemma 2. *Let v be an infinitesimal conformal transformation on M^n so that*

$$(2.5) \quad L_v g_{ij} = 2\phi g_{ij}.$$

Then

$$(2.6) \quad \begin{aligned} L_v R_{hijk} &= 2\phi R_{hijk} - g_{hk} \nabla_j \phi_i + g_{hj} \nabla_k \phi_i \\ &\quad - g_{ij} \nabla_k \phi_h + g_{ik} \nabla_j \phi_h, \end{aligned}$$

$$(2.7) \quad L_v R_{ij} = g_{ij} \Delta\phi - (n-2) \nabla_j \phi_i,$$

$$(2.8) \quad L_v R = 2(n-1) \Delta\phi - 2R\phi.$$

Lemma 2 can be proved by a straightforward computation.

Lemma 3. *If M^n has constant R and admits an infinitesimal nonhomothetic conformal transformation v so that (1.7) holds, then*

$$(2.9) \quad \Delta\phi = R\phi/(n-1),$$

$$(2.10) \quad R > 0.$$

Equation (2.9) follows from equation (2.8) due to the constancy of R , and equation (2.10) from Lemma 1.

Lemma 4 (Yano and Sawaki [5]). *If M^n admits an infinitesimal conformal transformation v so that equation (2.5) holds, then, for any scalar field f on M^n ,*

$$(2.11) \quad \int_{M^n} \phi f \, dA = -\frac{1}{n} \int_{M^n} L_v f \, dA.$$

Proof. Substituting $f v_i$ for ξ_i in the Green's formula (2.3) we obtain

$$(2.12) \quad \int_{M^n} f \nabla^i v_i \, dA = - \int_{M^n} v_i \nabla^i f \, dA = - \int_{M^n} L_v f \, dA.$$

On the other hand, since

$$L_v g_{ij} = \nabla_i v_j + \nabla_j v_i,$$

from equation (2.5) we have $\nabla^i v_i = n\phi$, which and equation (2.12) yield the required equation (2.11) immediately.

3. Proof of Theorem 3

On the manifold M^n consider the covariant tensor field of order 4:

$$(3.1) \quad \begin{aligned} W_{hijk} = & aT_{hijk} + b_1 g_{hk} T_{ij} - b_2 g_{hj} T_{ik} + b_3 g_{ij} T_{hk} \\ & - b_4 g_{ik} T_{hj} + b_5 g_{hi} T_{jk} - b_6 g_{jk} T_{hi}, \end{aligned}$$

where

$$(3.2) \quad T_{hijk} = R_{hijk} - \frac{R}{n(n-1)} (g_{ij} g_{hk} - g_{ik} g_{hj}),$$

and a and b 's are constants satisfying (1.6). Then

$$(3.3) \quad W^{hijk} W_{hijk} = a^2 P + \frac{c - 4a^2}{n-2} Q - \frac{1}{n} \left(\frac{2a^2}{n-1} + \frac{c - 4a^2}{n-2} \right) R^2,$$

where c is defined by (1.6). From equation (3.3) it follows that

$$(3.4) \quad L_v (W^{hijk} W_{hijk}) = L_v \left((a^2 P + \frac{c - 4a^2}{n-2} Q) \right).$$

By assuming the infinitesimal nonhomothetic conformal transformation v to

be defined by (1.7), from equations (3.1), (3.2), (1.9), (2.5), (2.6), (2.7), (2.9) we can easily obtain

$$\begin{aligned}
 L_v W_{hijk} &= 2a\phi R_{hijk} - [a + (n - 2)b_1]g_{hk}\nabla_j\phi_i \\
 &+ [a + (n - 2)b_2]g_{hj}\nabla_k\phi_i - [a + (n - 2)b_3]g_{ij}\nabla_k\phi_h \\
 &+ [a + (n - 2)b_4]g_{ik}\nabla_j\phi_h - (n - 2)b_5g_{hi}\nabla_k\phi_j + (n - 2)b_6g_{jk}\nabla_i\phi_h \\
 &- \frac{\phi R}{n(n - 1)}g_{ij}g_{hk}[4a + (3n - 4)(b_1 + b_3)] \\
 (3.5) \quad &+ \frac{\phi R}{n(n - 1)}g_{ik}g_{hj}[4a + (3n - 4)(b_2 + b_4)] \\
 &- \frac{3n - 4}{n(n - 1)}b_5\phi Rg_{hi}g_{jk} + \frac{3n - 4}{n(n - 1)}b_6\phi Rg_{jk}g_{hi} + 2b_1\phi g_{nk}R_{ij} \\
 &- 2b_2\phi g_{hj}R_{ik} + 2b_3\phi g_{ij}R_{hk} - 2b_4\phi g_{ik}R_{hj} + 2b_5\phi g_{hi}R_{jk} \\
 &- 2b_6\phi g_{jk}R_{ki} .
 \end{aligned}$$

Multiplying both sides of equation (3.5) by W^{hijk} and making use of equations (3.1), (3.2), (1.9), (3.3), (1.6) and $R^i_{ijk} = 0$, an elementary but lengthy calculation yields

$$(3.6) \quad W^{hijk}L_v W_{hijk} = 2\phi W^{hijk}W_{hijk} - cT^{ij}\nabla_j\phi_i .$$

By substituting equation (3.6) in the well-known formula

$$(3.7) \quad L_v(W^{hijk}W_{hijk}) + 2W^{hijk}L_v W_{hijk} - 8\phi W^{hijk}W_{hijk},$$

we thus have

$$(3.8) \quad \phi L_v(W^{hijk}W_{hijk}) = -4\phi^2 W^{hijk}W_{hijk} - 2c\phi T^{ij}\nabla_j\phi_i .$$

Since the manifold M^n is of constant R , it is known that for $n < 2$

$$(3.9) \quad \nabla^j R_{ij} = 0 ,$$

and therefore

$$(3.10) \quad \nabla^j T_{ij} = 0 .$$

Thus

$$(3.11) \quad \nabla^j(T_{ij}\phi^i) = T_{ij}\phi^i\phi^j + \phi T_{ij}\nabla^j\phi^i .$$

Without loss of generality we may assume our manifold M^n to be orientable, since otherwise we need only to take an orientable twofold covering space of

M^n . Substituting equation (3.8) in equation (3.11), integrating the resulting equation over the manifold M^n and using equation (2.3) we obtain

$$(3.12) \quad 2c \int_{M^n} T_{ij} \phi^i \phi^j dA = \int_{M^n} \phi L_v(W^{hijk} W_{hijk}) dA \\ + 4 \int_{M^n} \phi^2 W^{hijk} W_{hijk} dA .$$

On the right side of equation (3.12), the second integral is nonnegative since its integrand is so, and the first integral is equal to, by Lemma 4 and equations (3.3), (1.5),

$$-\frac{1}{n} \int_{M^n} L_v L_v(W^{hijk} W_{hijk}) dA = -\frac{1}{n} \int_{M^n} L_v L_v \left(a^2 P + \frac{c - 4a^2}{n - 2} Q \right) dA \geq 0 .$$

Hence the integral on the left side of equation (3.12) is nonnegative, and Theorem 3 follows from Theorem 4 immediately.

References

- [1] C. C. Hsiung, *On the group of conformal transformations of a compact Riemannian manifold*, Proc. Nat. Acad. Sci. U.S.A. **54** (1965) 1509-1513.
- [2] —, *On the group of conformal transformations of a compact Riemannian manifold. II*, Duke Math. J. **34** (1967) 337-341.
- [3] A. Lichnerowicz, *Sur les transformations conformes d'une variété riemannienne compacte*, C. R. Acad. Sci. Paris **259** (1964) 697-700.
- [4] K. Yano, *On Riemannian manifolds with constant scalar curvature admitting a conformal transformation group*, Proc. Nat. Acad. Sci. U.S.A. **55** (1966) 472-476.
- [5] K. Yano & S. Sawaki, *Riemannian manifolds admitting a conformal transformation group*, J. Differential Geometry **2** (1968) 185-190.

LEHIGH UNIVERSITY