

## AN INTEGRAL FORMULA FOR IMMERSIONS IN EUCLIDEAN SPACE

ROBERT B. GARDNER

### 1. Introduction

This paper derives a general rigidity theorem and an integral formula for immersions of a compact oriented riemannian manifold without boundary in a euclidean space. The formula is applied to a volume-preserving immersion to establish a simple geometric criterion that the immersion be isometric. As the integral formula has a formal resemblance to one derived by Chern and Hsiung in [1], we conclude the paper with some remarks about that work.

### 2. Notations and conventions

Let  $M$  be a compact oriented  $m$ -dimensional riemannian manifold without boundary with metric  $ds^{2*}$ , and let

$$X: M \rightarrow R^{m+n}$$

be an immersion in an  $(m + n)$ -dimensional euclidean space  $R^{m+n}$ . As such  $M$  admits a second riemannian metric,

$$ds^2 = dX \cdot dX .$$

We fix the range of indices so that the capital Latin indices run from 1 to  $m + n$ , the small Greek indices from 1 to  $m$ , and the small Latin indices from  $m + 1$  to  $m + n$ .

Matters being so, we choose orthonormal coframes  $\{\tau^{a*}\}$  for  $ds^{2*}$  on  $M$  which diagonalize  $ds^2$  with respect to  $ds^{2*}$ . Thus

$$ds^{2*} = \Sigma(\tau^{a*})^2 , \quad ds^2 = \Sigma g_a(\tau^{a*})^2 ,$$

and the first invariants of the pair of metrics are the elementary symmetric functions in the functions  $g_a$ .

Next we choose a family of orthonormal frames  $\{e_a\}$  on  $X(M)$  in  $R^{m+n}$  in such a way that  $\{e_a\}$  are unit tangent vectors of  $X(M)$  and the pull back of the dual coframe  $\{\tau^A\}$  satisfies

$$\tau^a = h_a \tau^{a*} ,$$

where  $h_\alpha = (g_\alpha)^{1/2}$ . As such the volume elements of  $ds^2$  and  $ds^{2\sharp}$  are respectively

$$dV = \tau^1 \wedge \dots \wedge \tau^m, \quad dV^\sharp = \tau^{1\sharp} \wedge \dots \wedge \tau^{m\sharp}.$$

The pull back of the structure equations

$$\begin{aligned} de_A &= \Sigma \varphi_A^\beta e_B, \\ d\tau^B &= \Sigma \tau^A \wedge \varphi_A^B, \\ d\varphi_A^B &= \Sigma \varphi_A^C \wedge \varphi_C^B \end{aligned}$$

of  $R^{m+n}$  give rise to a skew-symmetric matrix of linear differential forms

$$\varphi_\alpha^\beta = \Sigma \Gamma_{\alpha\gamma}^\beta \tau^\gamma,$$

called the Levi-Civita connection for  $ds^2$ , and a vector of quadratic differential forms

$$\Sigma \tau^\alpha \odot \varphi_\alpha^a = \Sigma A_{\alpha\beta}^a \tau^\alpha \odot \tau^\beta,$$

called the vector-valued second fundamental form.

The exterior differential equations

$$\begin{aligned} d\tau^{\alpha\sharp} &= \Sigma \tau^{\alpha\sharp} \wedge \varphi_\tau^{\alpha\sharp}, \\ \varphi_\tau^{\alpha\sharp} &= -\varphi_\alpha^{\tau\sharp} \end{aligned}$$

define a unique skew-symmetric matrix of linear differential forms

$$\varphi_\alpha^{\beta\sharp} = \Sigma \Gamma_{\alpha\gamma}^{\beta\sharp} \tau^\gamma,$$

called the Levi-Civita connection for  $ds^{2\sharp}$ . This matrix allows us to introduce a covariant differentiation with respect to  $ds^{2\sharp}$ . Thus, if  $f$  is a function we introduce  $f_{;\alpha}$  by

$$df = \Sigma f_{;\alpha} \tau^{\alpha\sharp};$$

if  $w = \Sigma a_\alpha \tau^{\alpha\sharp}$  is a linear differential form then we introduce  $a_{\alpha;\beta}$  by

$$da_\alpha - \Sigma a_\tau \varphi_\alpha^{\tau\sharp} = \Sigma a_{\alpha;\beta} \tau^{\beta\sharp};$$

if  $Q = \Sigma b_{\alpha\beta} \tau_\alpha^{\alpha\sharp} \odot \tau^{\beta\sharp}$  is a quadratic differential form then we introduce  $b_{\alpha\beta;\gamma}$  by

$$\begin{aligned} db_{\alpha\beta} - \Sigma \varphi_\alpha^{\gamma\sharp} b_{\gamma\beta} - \Sigma b_{\alpha\tau} \varphi_\tau^{\beta\sharp} \\ = \Sigma b_{\alpha\beta;\gamma} \tau^{\gamma\sharp}. \end{aligned}$$

Finally we introduce the Hodge mapping defined with respect to  $ds^{2\sharp}$ , which is the linear mapping  $*_\sharp$  characterized by

$$*_\sharp(\tau^{\alpha\sharp}) = (-1)^{\alpha-1} \tau^{1\sharp} \wedge \dots \wedge \tau^{\alpha-1\sharp} \wedge \tau^{\alpha+1\sharp} \wedge \dots \wedge \tau^{m\sharp}.$$

As such if  $w = \Sigma a_\alpha \tau^{\alpha\sharp}$  is a linear differential form then  $d *_\sharp w$  is an exact  $m$ -form, and a short calculation proves that

$$d *_\sharp w = \Sigma a_{\alpha;\alpha} \tau^{1\sharp} \wedge \dots \wedge \tau^{m\sharp} = \Sigma a_{\alpha;\alpha} dV^\sharp.$$

We recall that if  $w = df$ , where  $f$  is a real-valued function, then

$$d *_{\sharp} df = \Delta_{\sharp}(f)dV ,$$

where  $\Delta_{\sharp}(f)$  is the Laplacian of  $f$  taken with respect to the metric  $ds^{2\sharp}$ .

These operations make sense in the case that  $ds^{2\sharp} = ds^2$ , and we will denote the Laplacian with respect to  $ds^2$  by  $\Delta$ .

### 3. The integral formula

Let 0 denote a choice of origin in  $R^{m+n}$ ; then the linear differential form

$$\Omega = \Sigma(X \cdot e_{\alpha})\tau^{\alpha} = \frac{1}{2}X \cdot dX$$

is defined independent of the particular family of the orthonormal frames  $\{e_{\alpha}\}$  and orthonormal coframes  $\{\tau^{\alpha}\}$ , and hence induces a globally defined differential form on  $M$ . As such Stokes' theorem applies to yield the integral formula

$$(3.1) \quad 0 = \int_M d *_{\sharp} \Omega = \int_M \Delta_{\sharp}(\frac{1}{2}X \cdot X)dv .$$

The explicit expression of the resulting integral formula is simplified by the introduction of the vector

$$(3.2) \quad \begin{aligned} h^* &= \Sigma A_{\alpha\alpha}^{\alpha} h_{\alpha}^2 e_{\alpha} + \Sigma(\Gamma_{\alpha\alpha}^{\beta} - \Gamma_{\alpha\alpha}^{\beta\sharp}) h_{\alpha}^2 e_{\beta} \\ &+ \Sigma(h_{\alpha} \delta_{\alpha}^{\beta})_{;\beta} e_{\beta} . \end{aligned}$$

The naturality of this vector is apparent from the following proposition.

**Proposition 3.3.** *Let  $a$  be any fixed vector in  $R^{m+n}$ ; then*

$$(3.3) \quad \Delta_{\sharp}(a \cdot X) = a \cdot h^* .$$

*Proof.* Utilizing the structure equations, we have

$$\begin{aligned} d(a \cdot X) &= \Sigma(a \cdot e_{\alpha}) h_{\alpha} \tau^{\alpha\sharp} , \\ d(a \cdot e_{\alpha}) h_{\alpha} - \Sigma \varphi_{\alpha}^{\beta\sharp}(a \cdot e_{\beta}) h_{\beta} & \\ &= \Sigma(a \cdot e_i) A_{\alpha\gamma}^i h_{\alpha} h_{\gamma} \tau^{\gamma\sharp} + \Sigma(a \cdot e_{\beta})(\Gamma_{\alpha\gamma}^{\beta} - \Gamma_{\alpha\gamma}^{\beta\sharp}) h_{\alpha} h_{\gamma} \tau^{\gamma\sharp} \\ &+ \Sigma(a \cdot e_{\beta}) h_{\gamma} h_{\alpha} \Gamma_{\alpha\gamma}^{\beta\sharp} \tau^{\gamma\sharp} , \end{aligned}$$

and hence contracting the coefficients on  $\alpha$  and  $\gamma$  gives (3.3) as claimed.

In particular this last Proposition is true if  $ds^{2\sharp} = ds^2$ . In this case the vector characterized by the last proposition will be denoted by  $h$ . We note that

$$(3.4) \quad h = \Sigma A_{\alpha\alpha}^i e_i ,$$

which is the mean curvature vector of the immersion.

With this preparation the integral formula obtained from (3.1) may be stated as follows.

**Theorem 3.4.** *Let  $M$  be a compact oriented manifold without boundary endowed with the riemannian metric  $ds^{2\sharp} = \Sigma(\tau^{\alpha\sharp})^2$ , and let*

$$X: M \rightarrow R^{m+n}$$

*be an immersion with induced metric  $ds^2 = \Sigma g_\alpha(\tau^{\alpha\sharp})^2$ , then*

$$(3.5) \quad 0 = \int_M (\Sigma g_\alpha + X \cdot h^*) dV^\sharp .$$

*Proof.* Since

$$\begin{aligned} d(X \cdot e_\alpha)h_\alpha - (X \cdot e_r)h_r\varphi_\alpha^{\sharp*} &= \tau^\alpha h_\alpha + \Sigma(X \cdot e_r)\varphi_r^\alpha h_\alpha + \Sigma(X \cdot e_i)\varphi_i^\alpha h_\alpha \\ &\quad + (X \cdot e_\alpha)dh_\alpha - \Sigma(X \cdot e_r)h_r\varphi_\alpha^{\sharp*} \\ &= g_\alpha \tau^{\alpha\sharp} + \Sigma(X \cdot e_r)(\varphi_r^\alpha - \varphi_\alpha^{\sharp*})h_\alpha \\ &\quad + \Sigma(X \cdot e_r)(dh_\alpha \delta_r^\alpha - h_r\varphi_\alpha^{\sharp*})h_\alpha \\ &\quad + \Sigma(X \cdot e_i)\varphi_i^\alpha h_\alpha , \end{aligned}$$

we have

$$\begin{aligned} (\Sigma(X \cdot e_\alpha)h_\alpha)_{,\alpha} &= \Sigma g_\alpha + \Sigma(X \cdot e_\alpha)(\Gamma_{rr}^\alpha - \Gamma_{rr}^{\alpha\sharp})g_r \\ &\quad + \Sigma(X \cdot e_r)(h_\alpha \delta_r^\alpha)_{,\alpha} + \Sigma(X \cdot e_i)A_{\alpha\alpha}^i g_\alpha \\ &= \Sigma g_\alpha + X \cdot h^* , \end{aligned}$$

which gives (3.5) by integration.

We note that applying the formula to the special case, where  $ds^{2\sharp} = ds^2$ , gives

$$(3.6) \quad 0 = \int_M (m + X \cdot h) dV ,$$

which is a classical formula of Minkowski.

#### 4. Applications to volume-preserving immersions

**Theorem 4.1.** *Let  $X: M \rightarrow R^{m+n}$  be an immersion of a compact oriented riemannian manifold without boundary. Then among all volume-preserving diffeomorphisms, the isometries are characterized as those for which the integral*

$$- \int_M X \cdot h^* dV$$

attains the minimal value of  $m$  times the value of  $\text{vol. } M$ .

*Proof.* By Newton's inequality, the hypothesis of volume-preserving implies

$$\frac{1}{m} \Sigma g_\alpha \geq (\Pi g_\alpha)^{1/m} = 1 ,$$

or

$$(4.2) \quad \Sigma g_\alpha - m \geq 0$$

with equality if and only if

$$(4.3) \quad g_\alpha = 1 \quad (1 \leq \alpha \leq m) .$$

As such subtraction of (3.5) from (3.6), together with the hypothesis that  $dV^* = dV$ , gives

$$0 = \int_M [(\Sigma g_\alpha - m) + X \cdot (h^* - h)] dV ,$$

but then (4.2) implies

$$\int_M X \cdot (h^* - h) dV \leq 0 ,$$

or

$$\int_M X \cdot h^* dV^* \leq \int_M X \cdot h dV = -m \text{ vol } M .$$

If this maximum is achieved, then the integral formula becomes

$$0 = \int_M (\Sigma g_\alpha - m) dV ,$$

and hence (4.2) forces

$$\Sigma g_\alpha - m = 0 ,$$

and the equality statement (4.3) implies that the immersion is an isometry.

**Corollary 4.4.** *Let  $X: M \rightarrow R^{m+n}$  be a volume-preserving immersion of a compact oriented riemannian manifold without boundary. Then*

$$h^* = h$$

*if and only if the immersion is isometric.*

**5. A general rigidity theorem**

Now consider the situation that the metric  $ds^{2\#}$  comes from a second immersion. Thus we have the picture

$$\begin{array}{ccc}
 M & \xrightarrow{X} & R^{m+n} \\
 & \searrow X^\# & \\
 & & R^{m+n}
 \end{array}$$

with  $ds^2 = dX \cdot dX$  and  $ds^{2\#} = dx^\# \cdot dx^\#$ .

**Theorem 5.** *A necessary and sufficient condition that two immersions of a compact oriented manifold without boundary differ by a translation is that*

$$h^* = h_\# ,$$

where  $h^*$  is defined by (3.2), and  $h_\#$  is the mean curvature vector of the  $X^\#$  immersion.

*Proof.* By Proposition 3.3 we have

$$\Delta_\#(X - X^\#) \cdot a = (h^* - h_\#) \cdot a .$$

Therefore  $X - X^\# = \text{constant}$  if and only if  $h^* = h_\#$ .

As a corollary we obtain the rigidity theorem that two isometric immersions of a compact oriented riemannian manifold without boundary differ by a translation if and only if they have the same mean curvature vectors. In the case of hypersurfaces this was a problem proposed by Minkowski.

**6. Remarks on the paper of Chern and Hsiung**

The integral formula in [1] was derived for volume-preserving diffeomorphisms between compact submanifolds of euclidean space without boundaries. One of the basic tools in [1] was the observation that Gårding's inequality applies to a classical mixed invariant of two positive definite quadratic forms. We will now show that a direct calculation of the mixed invariant allows us to deduce their inequality from Newton's inequality. C. C. Hsiung has pointed out that this is done by a different method in [2].

Let  $V$  be an  $n$ -dimensional real vector space, and  $\text{Hom}(V, V)$  the real vector space of all  $n \times n$  matrices with real coefficients. Then for  $X, Y \in \text{Hom}(V, V)$  we introduce functions  $P^i(X, Y)$  for  $1 \leq i \leq n - 1$  by

$$\det(X + tY) = \det X + tP^1(X, Y) + \dots + t^{n-1}P^{n-1}(X, Y) + t^n \det Y .$$

In particular

$$P^1(X, Y) = \frac{d}{dt} \det(X + tY)|_{t=0} = \langle [X + tY], d(\det) \rangle(X),$$

where  $[X + tY]$  is the tangent vector to the curve  $X + tY$  in  $\text{Hom}(V, V)$ , and  $\langle \cdot, \cdot \rangle$  is the canonical bilinear pairing between the tangent and cotangent spaces of  $\text{Hom}(V, V)$  at  $X$ .

If we introduce the natural coordinates

$$\pi_{ij} : \text{Hom}(V, V) \rightarrow R$$

defined for  $X = (X_{lm})$  by  $\pi_{ij}(X) = X_{ij}$ , then

$$\begin{aligned} d(\det)|_X &= \sum \frac{\partial \det X}{\partial \pi_{ij}} d\pi_{ij}|_X \\ &= \text{trace}(\text{cofactor } X \cdot dX), \end{aligned}$$

and

$$\begin{aligned} \langle [X + tY], dX \rangle &= \frac{d}{dt} \pi_{ij}(X + tY)|_{t=0} \\ &= (\pi_{ij}(Y)) = Y. \end{aligned}$$

Therefore by linearity

$$P^1(X, Y) = \text{trace}(\text{cofactor } X \cdot Y).$$

If  $X$  is non-singular, then

$$\text{cofactor } X = (\det X)X^{-1},$$

and hence the classical mixed invariant of the pair  $X, Y$  utilized by Chern and Hsiung in [1] is

$$(6.1) \quad Y_X = \frac{P^1(X, Y)}{n \det X} = \frac{1}{n} \text{trace}(X^{-1} \cdot Y).$$

The basic inequality used in [1] is thus equivalent to the fact that positive definite symmetric matrices  $X, Y$  satisfy

$$\frac{1}{n} \text{trace}(X^{-1} \cdot Y) \geq \left( \frac{\det Y}{\det X} \right)^{1/n}$$

with equality if and only if  $Y$  is congruent by an orthogonal matrix to a multiple of  $X$ . By diagonalizing  $Y$  with respect to  $X$  this is an immediate consequence of Newton's inequality.

Utilizing the explicit expression (6.1) of the mixed invariant, Donald Singley has proved that the integral formula in [1] may be generalized to immersions of compact riemannian manifolds without boundary by the integral formula

$$0 = \int_M d * *_{\sharp}^{-1} * \Omega .$$

### References

- [ 1 ] S. S. Chern & C. C. Hsiung, *On the isometry of compact submanifolds in Euclidean space*, Math. Ann. **149** (1963) 278–285.
- [ 2 ] B. H. Rhodes, *On some inequalities of Gårding*, Acad. Roy. Belg. Bull. Cl. Sci. (5) **52** (1966) 594–599.

COLUMBIA UNIVERSITY