

## The automorphism group of Leech lattice and elliptic modular functions

Dedicated to Professor Hiroshi Nagao on his 60th birthday

By Takeshi KONDO

(Received May 28, 1984)

### Introduction.

As usual, we denote by  $\cdot 0$  the automorphism group of Leech lattice which is an even unimodular lattice in 24-dimensional Euclidean space [1]. So  $\cdot 0$  has a natural 24-dimensional representation  $\rho_0$  over the rational number field. In this paper, Frame shapes of conjugacy classes of  $\cdot 0$  with respect to  $\rho_0$ , the list of which is given in Table I of Appendix, will play a central role. For the definition of Frame shape, see § 1.2.

Let  $\mathcal{F}$  be the set of all elliptic modular functions  $f(z)$  satisfying the following conditions:

- (1)  $f(z)$  is a modular function with respect to a discrete subgroup  $\Gamma$  of  $SL(2, \mathbf{R})$  containing  $\Gamma_0(N)$  for some integer  $N$  (i. e.  $f\left(\frac{az+b}{cz+d}\right) = f(z)$  for any  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$  and meromorphic on the upper half plane and at all cusps of  $\Gamma$ ),
- (2) the genus of  $\Gamma$  is zero and  $f(z)$  is a generator of a function field for  $\Gamma$ ,
- (3)  $f(z)$  has a Fourier expansion of the form  $f(z) = 1/q + \sum_{n=0}^{\infty} a_n q^n$  ( $q = e^{2\pi iz}$ ).

Now the main result of this paper is to show that various "transformations" (cf. § 1.1) of Frame shapes of  $\cdot 0$  yield functions of  $\mathcal{F}$  (Th. 3.2, 3.4, 3.5 and Table II~IV in Appendix). Furthermore, an application of this result is as follows: Let  $G$  be a finite group which has a  $d$ -dimensional representation  $\rho$  over the rational number field where  $d$  is a divisor of 24. For each of such many (not all) pairs  $(G, \rho)$ , we can construct a mapping from  $G$  to  $\mathcal{F}$

$$G \ni \sigma \longmapsto j_\sigma(z) \in \mathcal{F}$$

such that all coefficients  $a_k(\sigma)$  ( $k \geq 1$ ) of a Fourier expansion  $j_\sigma(z) = 1/q + \sum_{k=0}^{\infty} a_k(\sigma) q^k$  are generalized characters of  $G$  (Th. 4.6, 4.8 and 4.10). Such a mapping is called a *moonshine* of  $G$ . A moonshine of Fischer-Griess's Monster is constructed in a remarkable paper of Conway-Norton [2] and other examples of moonshines can be found in Queen [10] and Koike [4]. Constructions of moon-

shines in this paper are rather elementary compared with those of Conway-Norton-Queen. For examples of pairs  $(G, \rho)$  which does not yield a moonshine, we refer readers to Remark 4.4.

The author is very grateful to Prof. Masao Koike for many valuable discussions and suggestions.

#### NOTATIONS.

$\mathbf{Z}$  the ring of rational integers

$\mathbf{Q}$  the field of rational numbers

$\mathbf{R}$  the field of real numbers

$$SL(2, \mathbf{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbf{Z} \text{ and } ad - bc = 1 \right\}$$

$$SL(2, \mathbf{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbf{R} \text{ and } ad - bc = 1 \right\}$$

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbf{Z}) \mid c \equiv 0 \pmod{N} \right\}$$

$\langle \dots \rangle$  = a group generated by  $\dots$ .

For the notations of conjugacy classes of  $\cdot 0$ , see the first paragraph of Appendix.

### §1. Generalized permutations and Frame shapes.

#### 1.1. A symbol

$$\prod_t t^{r_t} = 1^{r_1} 2^{r_2} \dots \quad (r_t \in \mathbf{Z})$$

is called a *generalized permutation* if  $r_t \neq 0$  for only a finite number of positive integers  $t$ . For a generalized permutation  $\pi = \prod_t t^{r_t}$ , set

$$\deg \pi = \sum_t t r_t,$$

$$\operatorname{sgn} \pi = \prod_t (-1)^{(t-1)r_t}.$$

Obviously  $\deg \pi$  and  $\operatorname{sgn} \pi$  are generalizations of degree and sign of a permutation on a finite set.

Now we will define some transformations of a generalized permutation. Let  $r$  be a positive integer and  $\pi = \prod_t t^{r_t}$  be a generalized permutation. Then define

$$\pi/r = \prod_t (rt)^{r_t/r}, \quad \text{where } r|r_t \text{ for any } t,$$

$$\pi \circ r = \prod_t t^{r_t/(r+1)} (rt)^{r_t/(r+1)}, \quad \text{where } r+1|r_t \text{ for any } t,$$

$$\pi \circ (r/1) = \prod_t (rt)^{r_t/(r-1)} t^{-r_t/(r-1)}, \quad \text{where } r-1|r_t \text{ for any } t.$$

These are called the *r-th harmonic*, the *r-transformation* and the *(r/1)-transformation* of  $\pi$  respectively. All of these transformations have the same degree as  $\pi$ .

We note that (2/1)-transformation can be defined for all generalized permutations.

Let  $\eta(z)$  be Dedekind eta-function :

$$\eta(z) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n), \quad q = e^{2\pi iz}.$$

For a generalized permutation  $\pi = \prod_t t^{r_t}$ , we put

$$(1.1) \quad \eta_{\pi}(z) = \prod_t \eta(tz)^{r_t}.$$

The meaning of the transformations defined above consists in considering the transformation of functions  $\eta_{\pi}(z)$  :

$$\eta_{\pi}(z) \longmapsto \eta_{\pi^*}(z) \quad \pi^* = \pi/r, \quad \pi \circ (r/1) \quad \text{or} \quad \pi \circ r.$$

1.2. Let  $G$  be a finite group and

$$G \ni \sigma \longmapsto \rho(\sigma) \in GL(d, \mathbf{Q})$$

be a  $d$ -dimensional representation of  $G$  over the rational number field  $\mathbf{Q}$ . Then we will assign to every element (or every conjugacy class) of  $G$  a generalized permutation of degree  $d$  as follows. The characteristic polynomial  $\det(xI_d - \rho(\sigma))$  ( $I_d$  = the identity matrix of degree  $d$ ) of  $\rho(\sigma)$  ( $\sigma \in G$ ) can be written in the form

$$\prod_t (x^t - 1)^{r_t} \quad (r_t \in \mathbf{Z})$$

where  $t$  ranges over all positive integers dividing the order of  $G$ . Then a generalized permutation  $\prod_t t^{r_t}$  of degree  $d$  is called *Frame shape* of an element  $\sigma$  with respect to the representation  $\rho$ . We also refer to Frame shape of a conjugacy class of  $G$  (w.r.t.  $\rho$ ), as two conjugate elements of  $G$  yield the same Frame shape.

REMARK 1.1. If a representation  $\rho$  is a permutation representation of  $G$  (i.e. every  $\rho(\sigma)$  is a permutation matrix), the Frame shape of  $\sigma$  w.r.t.  $\rho$  is just a cycle decomposition of a permutation corresponding to  $\rho(\sigma)$ . Thus Frame shape can be regarded as a generalization of a cycle decomposition of an element in a permutation group.

REMARK 1.2. If  $G$  has no subgroup of index 2 and so  $\det \rho(\sigma) = 1$  for all  $\sigma \in G$ , we have  $\text{sgn } \pi = 1$  for all Frame shapes  $\pi$  of conjugacy classes of  $G$ , because  $\det \rho(\sigma) = \text{sgn } \pi$ .

REMARK 1.3. A generalized permutation is not always a Frame shape. For example, a generalized permutation  $1.2^{-2}4$  is not a Frame shape, as  $(x-1)(x^4-1)/(x^2-1)^2$  is not a polynomial.

## §2. A class of elliptic modular functions.

2.1. As in the introduction, let  $\mathcal{F}$  be the set of all elliptic modular functions  $f(z)$  having the following properties:

(1)  $f(z)$  is a modular function with respect to a discrete subgroup  $\Gamma$  of  $SL(2, \mathbf{R})$  containing some  $\Gamma_0(N)$ ,

(2) the genus of  $\Gamma$  is zero and  $f(z)$  is a generator of a function field for  $\Gamma$  and

(3)  $f(z)$  has a Fourier expansion of the form  $f(z) = 1/q + \sum_{n=0}^{\infty} a_n q^n$  ( $q = e^{2\pi iz}$ ). For simplicity, we call  $\Gamma$  in (1) and (2) a group for  $f(z)$  and also  $f(z)$  a Hauptmodule for  $\Gamma$ . Clearly the well known modular invariant  $j(z)$  belongs to  $\mathcal{F}$  and  $\Gamma_0(1) = SL(2, \mathbf{Z})$  is a group for  $j(z)$ . Other examples of  $f(z) \in \mathcal{F}$  and a group for  $f(z)$  can be found in Table 3 of [2] which is very useful in this paper. In these examples, a group for  $f(z) \in \mathcal{F}$  is the one obtained by adjoining to  $\Gamma_0(N)$  some of its Atkin-Lehner's involutions  $W_Q$

$$W_{Q,N} = W_Q = \begin{pmatrix} aQ & b \\ cN & dQ \end{pmatrix} \quad a, b, c, d \in \mathbf{Z}$$

where  $Q \parallel N$ , i.e.  $Q$  is a divisor of  $N$  with  $(Q, N/Q) = 1$  and  $\det W_Q = Q$ . As in [2] and [10], we use the notations

$$N+Q_1, Q_2, \dots, \quad N-, \quad N+$$

which denote

$$\langle \Gamma_0(N), W_{Q_1}, W_{Q_2}, \dots \rangle, \quad \Gamma_0(N), \quad \langle \Gamma_0(N), W_Q \mid Q \parallel N \rangle$$

respectively.

LEMMA 2.1. Let  $\eta_\pi(z)$  be a function defined by (1.1) for a generalized permutation  $\pi$ . Assume that

(1)  $\deg \pi = -2A$ ,

(2)  $\eta_\pi(z)$  is a modular function w.r.t. a discrete subgroup  $\Gamma$  of  $SL(2, \mathbf{R})$  containing some  $\Gamma_0(N)$ ,

(3)  $\Gamma_{i\infty} = \{M(i\infty) = i\infty \mid M \in \Gamma\}$  is generated by  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ ,

(4)  $z = i\infty$  is the unique pole of  $\eta_\pi(z)$  among all inequivalent cusps of  $\Gamma$ . Then  $\eta_\pi(z) \in \mathcal{F}$  and  $\Gamma$  is a group for  $\eta_\pi(z)$ .

PROOF. The condition (1) means that  $\eta_\pi(z)$  has a Fourier expansion of the form  $1/q + \sum_{n=0}^{\infty} a_n q^n$  and the condition (3) shows that  $q$  can be taken as a local parameter of  $\eta_\pi(z)$  at  $z = i\infty$  and so  $z = i\infty$  is a pole of  $\eta_\pi(z)$  of order 1. Let  $\mathfrak{R}$  be a Riemann surface corresponding to  $\Gamma$ , i.e.  $\mathfrak{R} = \Gamma \backslash \mathfrak{H}^*$  where  $\mathfrak{H}^*$  is a union of the upper half plane and the set of all cusps of  $\Gamma$ . Since  $\eta_\pi(z)$  has no

poles on the upper half plane, the condition (4) means that  $z=i\infty$  is the unique pole of  $\eta_\pi(z)$  on  $\mathfrak{H}$  and so  $\eta_\pi(z)$  gives an isomorphism from  $\mathfrak{H}$  onto the Riemann sphere. Thus the genus of  $\mathfrak{H}$  is zero and  $\eta_\pi(z)$  is a generator of a function field of  $\mathfrak{H}$ . This completes the proof of Lemma 2.1.

2.2. Here we mention the well known transformation formula of Dedekind eta-function :

$$(2.1) \quad \eta\left(\frac{az+b}{cz+d}\right) = v(M)(cz+d)^{1/2}\eta(z) \quad \text{for } M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbf{Z})$$

where  $v(M)^{24}=1$ . An explicit formula of  $v(M)$  was given by Petersson [6; Th. 2 of Chap. 4] :

$$(2.2) \quad v(M) = \begin{cases} \left(\frac{d}{c}\right)^* \exp\left\{\frac{\pi i}{12}[(a+d)c - bd(c^2-1) - 3c]\right\} & \text{if } c \text{ is odd} \\ \left(\frac{c}{d}\right)_* \exp\left\{\frac{\pi i}{12}[(a+d)c - bd(c^2-1) + 3d - 3 - 3cd]\right\} & \text{if } c \text{ is even} \end{cases}$$

where, by using Jacobi symbol  $\left(\frac{n}{m}\right)$ , we put

$$\left(\frac{c}{d}\right)^* = \left(\frac{c}{|d|}\right) \quad \text{and} \quad \left(\frac{c}{d}\right)_* = \left(\frac{c}{|d|}\right)(-1)^e, \quad e = \frac{\text{sgn } c - 1}{2} \cdot \frac{\text{sgn } d - 1}{2},$$

$$\left(\frac{0}{\pm 1}\right)^* = 1, \quad \left(\frac{0}{1}\right)_* = 1 \quad \text{and} \quad \left(\frac{0}{-1}\right)_* = -1.$$

Now we give some formulas which are useful for our calculations of  $\eta_\pi(z)$  :

$$(2.3) \quad \eta\left(z + \frac{1}{2}\right) = e^{\pi i/24} \eta(2z)^3 / \eta(z)\eta(4z).$$

(2.4) If  $2|N_0|N$  and  $(2N_0, N) = N_0$ ,

$$\eta(2N_0z/(Nz+1)) = v(M)(Nz+1)^{1/2}\eta((N_0z+1)/2),$$

$$M = \begin{pmatrix} 2 & -1 \\ N/N_0 & (N_0-N)/2N_0 \end{pmatrix} \in SL(2, \mathbf{Z}).$$

$$(2.5) \quad \eta\left(z + \frac{1}{3}\right)\eta\left(z + \frac{2}{3}\right) = e^{\pi i/12} \eta(3z)^4 / \eta(z)\eta(9z).$$

(2.6) If  $3|N_0|N$ ,  $(3N_0, N) = N_0$  and  $N/N_0 \equiv \varepsilon \pmod{3}$  ( $\varepsilon = \pm 1$ ),

$$\eta(3N_0z/(Nz+1)) = v(M)(Nz+1)^{1/2}\eta((N_0z+\varepsilon)/3),$$

$$M = \begin{pmatrix} 3 & -\varepsilon \\ N/N_0 & (N_0-N\varepsilon)/3 \end{pmatrix} \in SL(2, \mathbf{Z}).$$

(2.7) If  $W_Q = \begin{pmatrix} aQ & b \\ cN & dQ \end{pmatrix}$  is an Atkin-Lehner's involution of  $\Gamma_0(N)$ ,

$$\eta(tW_Q(z)) = v(M)(Q, t)^{-1/2}(cNz + dQ)^{1/2}\eta((Qt/(Q, t)^2)z),$$

$$\text{where } M = \begin{pmatrix} a(Q, t) & bt/(Q, t) \\ cN(Q, t)/Qt & dQ/(Q, t) \end{pmatrix} \in SL(2, \mathbf{Z}).$$

(2.3) and (2.5) are obtained by direct computations. (2.4), (2.6) and (2.7) follow from (2.1).

LEMMA 2.2 (M. Newmann [9; Th. 1]). *Let  $\pi = \prod_{t|N} t^{r_t}$  be a generalized permutation and  $\eta_\pi(z) = \prod_{t|N} \eta(tz)^{r_t}$ , where  $t$  ranges over all positive divisors of some integer  $N$ . Assume that*

- (0)  $\sum_t r_t = 0,$
- (1)  $\sum_t r_t t \equiv 0 \pmod{24},$
- (2)  $\sum_t r_t N/t \equiv 0 \pmod{24}$
- (3) *the number  $\prod_{t|N} t^{r_t}$  is a rational square.*

Then  $\eta_\pi(z)$  is a modular function w.r.t.  $\Gamma_0(N)$ .

A proof of Lemma 2.2 can be done by using (2.1) and (2.2).

LEMMA 2.3. *Let  $\pi = \prod_t t^{r_t}$  be a generalized permutation of degree 24 and  $r > 1$  be an integer with  $r | r_t$  for any  $t$ . Assume that*

- (1)  $\eta_\pi(z)^{-1} \in \mathfrak{F},$
- (2)  $\prod_t t^{r_t/r}$  is a rational square.

Then  $\eta_{\pi/r}(z)^{-1} \in \mathfrak{F}$ , where  $\pi/r$  is the  $r$ -th harmonic of  $\pi$ .

PROOF. Let  $f(z) = \eta_\pi(z)^{-1}$  and  $g(z) = \eta_{\pi/r}(z)^{-1}$ . Then we have

$$(*) \quad g(z) = f(rz)^{1/r}.$$

If  $\Gamma$  is a group for  $f(z) \in \mathfrak{F}$ ,  $f(rz)$  is a modular function w.r.t.  $\Gamma_1 = \begin{pmatrix} r & 0 \\ 0 & 1 \end{pmatrix}^{-1} \Gamma \begin{pmatrix} r & 0 \\ 0 & 1 \end{pmatrix}$  and  $[\mathbf{C}(g(z)) : \mathbf{C}(f(rz))] = r$ . Also we have, by (\*),

$$(**) \quad g(Mz) = \delta g(z) \quad (\delta^r = 1) \quad \text{for any } M \in \Gamma_1$$

and, in particular,

$$(***) \quad g\left(z + \frac{1}{r}\right) = e^{-2\pi i/r} g(z) \quad \text{for } \begin{pmatrix} 1 & 1/r \\ 0 & 1 \end{pmatrix} \in \Gamma_1.$$

Now let  $\Gamma_2 = \{M \in \Gamma_1 | g(Mz) = g(z)\}$ . Then, by (\*\*) and (\*\*\*), we must have  $[\Gamma_1 : \Gamma_2] = r$  and so  $\mathbf{C}(g(z))$  is a function field for  $\Gamma_2$ . By the assumption (2) and

Lemma 2.2,  $\Gamma_2$  contains some  $\Gamma_0(N)$ . Thus  $g(z) \in \mathcal{F}$ . This completes the proof of Lemma 2.3.

LEMMA 2.4. For a generalized permutation  $\pi = \prod_t t^{r_t}$  and  $T = \begin{pmatrix} 1 & 1/2 \\ 0 & 1 \end{pmatrix} \in SL(2, \mathbf{R})$ , define

$$\pi \circ T = \prod_{t: \text{even}} t^{r_t} \prod_{t: \text{odd}} (2t)^{3r_t} t^{-r_t} (4t)^{-r_t}.$$

If  $\eta_\pi(z) \in \mathcal{F}$ , we have  $\mathcal{F} \ni \eta_{\pi \circ T}(z) = -\eta_\pi(Tz)$ .

PROOF. If  $\eta_\pi(z)$  is a modular function w.r.t.  $\Gamma$ ,  $\eta_{\pi \circ T}(z)$  is a modular function w.r.t.  $T^{-1}\Gamma T$ . From this we get  $\mathcal{F} \ni \eta_{\pi \circ T}(z)$  if  $\eta_\pi(z) \in \mathcal{F}$ . The equality  $\eta_{\pi \circ T}(z) = -\eta_\pi(Tz)$  follows from (2.3), q.e.d.

### § 3. Frame shapes of conjugacy classes of $\cdot 0$ .

3.1. The automorphism group  $\cdot 0$  of Leech lattice has a natural 24-dimensional representation  $\rho_0$  over  $\mathbf{Q}$ . It is not difficult to compute Frame shape of every conjugacy class of  $\cdot 0$  by using the character values of the representation  $\rho_0$  and the power mapping of conjugacy classes of  $\cdot 0$  [11; Table 1]. The list of Frame shapes of conjugacy classes of  $\cdot 0$  is given in Table I of Appendix. The following observation of the list may be useful (cf. [2; p. 315]):

THEOREM 3.1. Let  $\pi = \prod_t t^{r_t}$  be a Frame shape of  $\cdot 0$ . Then  $\sum_t r_t$  is even and if  $\sum_t r_t = 0$ , we have  $\eta_\pi(z)^{-1} \in \mathcal{F}$  (= a class of elliptic modular functions defined in § 2).

PROOF. By inspection of Table I of Appendix,  $\sum_t r_t$  is even. It can be seen from Table 3 of [2] that, if  $\sum_t r_t = 0$ ,  $\eta_\pi(z)^{-1} \in \mathcal{F}$ .

REMARK 3.1. Let  $\sum_t r_t = 2k \neq 0$  and  $N$  be a product of L.C.M. and G.C.D. of  $\{t | r_t \neq 0\}$ . Then  $\eta_\pi(z)$  is a cusp form or an Eisenstein series of level  $N$  and weight  $k$  with some character, according as  $r_t \geq 0$  for any  $t$  or not (cf. [3], [5] and [8]).

The following theorem is one of the main results of this paper:

THEOREM 3.2. Let  $\pi = \prod_t t^{r_t}$  be a Frame shape of  $\cdot 0$  and  $r > 1$  be an integer with  $r-1 | 24$  and  $r-1 | r_t$  for any  $t$ . Then we have

$$(3.1) \quad \eta_{\pi \circ (r/1)}(z)^{-1} \in \mathcal{F} \quad (\pi \circ (r/1) = (r/1)\text{-transformation of } \pi \text{ (§ 1.1)}),$$

except for the following cases:

| $r$ | <i>classes</i>        |
|-----|-----------------------|
| 3   | $\pm 4C, 4F, 8D, 12M$ |
| 5   | $2C, 4B, 6I, 8B$      |
| 7   | $4F$                  |
| 13  | $2C$                  |

In these exceptional cases, we have  $\text{sgn}(\pi^{1/(r-1)}) = -1$ , where  $\pi^{1/(r-1)} = \prod_t t^{r_t/(r-1)}$ .

REMARK 3.2. The case  $r=2$  of Th. 3.2 is a part of a theorem of Conway-Norton-Queen [2], [10] which says that a mapping

$$\cdot 0 \ni \sigma \longmapsto \eta_{\sigma \circ (2/1)}(z)^{-1} = q^{-1} \prod_{n=0}^{\infty} \prod_{i=1}^{24} (1 - \varepsilon_i(\sigma) q^{2n-1})$$

is a moonshine of  $\cdot 0$ , where  $\varepsilon_i(\sigma)$  ( $1 \leq i \leq 24$ ) are eigenvalues of  $\rho_0(\sigma)$ . In Table II of Appendix, we will give  $\sigma \circ (2/1)$  and a group for  $\eta_{\sigma \circ (2/1)}(z)^{-1}$  for each  $\sigma$ . This table can be also found in Queen [10; Table 1], but, in Queen's table, some conjugacy classes of  $\cdot 0$  are missing and, in our table, a group for  $\eta_{\sigma \circ (2/1)}(z)^{-1}$  ( $\sigma \in \cdot 0$ ) is described more explicitly than in Queen's table.

REMARK 3.3. Notations being as in Th. 3.2, let  $G$  be a finite group with no subgroup of index 2 and  $\rho$  be a  $24/(r-1)$ -dimensional representation over  $\mathbf{Q}$ . If  $\text{sgn}(\pi^{1/(r-1)}) = -1$ ,  $\pi^{1/(r-1)}$  is not a Frame shape of  $G$  w.r.t.  $\rho$ , because  $\text{sgn}(\pi^{1/(r-1)}) = -1$  means that the determinant of a linear transformation with Frame shape  $\pi^{1/(r-1)}$  is  $-1$ .

PROOF OF THEOREM 3.2. This is done by using Table I of Appendix and examining (3.1) in case by case for each conjugacy class of  $\cdot 0$ . Here we will give a proof of the case  $r=3$ . (Also the case  $r=2$  (cf. Remark 3.2) can be dealt with quite similarly, and other cases  $r>3$  are rather easy to be examined.)

First of all, we see from Table I of Appendix that, if  $\pi$  is a Frame shape of  $\cdot 0$ ,  $\pi \circ (3/1)$  is

- (1) a Frame shape of  $\cdot 0$ ,
- (2) for some  $r$ , the  $r$ -th harmonic of a Frame shape  $\prod_t t^{r_t}$  of  $\cdot 0$  such that  $\sum_t r_t = 0$  and  $\prod_t t^{r_t/r}$  is a rational square, or
- (3) one of the following generalized permutations:

| classes of $\pi$ | $\pi \circ (3/1)$                         |  |
|------------------|---|--|
| $\pm 3A$         | $9^6 1^6 / 3^{12}$ ,                      | $2^6 3^{12} 18^6 / 1^6 6^{12} 9^6$   |
| $\pm 4C$         | $3^6 \cdot 12^2 / 1^2 \cdot 4^2$ ,        | $1^2 6^3 12^2 / 2^3 3^2 4^2$   |
| $4F$             | $12^3 / 4^3$                              |  |
| $\pm 6A$         | $1^2 2^2 9^2 18^2 / 3^4 6^4$ ,            | $2^4 3^4 18^4 / 1^2 6^8 9^2$   |
| $6B$             | $4^3 6^6 36^3 / 2^3 12^6 18^3$            |  |
| $8D$             | $4 \cdot 24^2 / 8^2 12$                   |  |
| $\pm 12A$        | $3^4 2^3 36^2 / 1^2 9^2 12^4$ ,           | $1^2 4^2 6^4 9^2 36^2 / 2^2 3^4 12^4 18^2$                                     |
| $12B$            | $4^2 6^2 36^2 / 2 \cdot 12^4 18$          |  |
| $12C$            | $2 \cdot 4 \cdot 18 \cdot 36 / 6^2 12^2$  |  |
| $12M$            | $36 / 12$                                 |  |
| $\pm 15B$        | $1 \cdot 5 \cdot 9 \cdot 45 / 3^2 15^2$ , | $2 \cdot 3^2 10^2 15 \cdot 18 \cdot 90 / 1 \cdot 5 \cdot 6^2 9 \cdot 30^2 45$  |
| $\pm 21A$        | $3^2 7 \cdot 63 / 1 \cdot 3 \cdot 21^2$ , | $1 \cdot 6^2 9 \cdot 14 \cdot 21^2 126 / 2 \cdot 3^2 7 \cdot 18 \cdot 42^2 63$ |
| $24A$            | $6^2 8 \cdot 72 / 2 \cdot 18 \cdot 24^2$  |  |



If we have the case (1) or (2), we can conclude from Th. 3.1 and Lemma 2.3 that  $\eta_{\pi \circ (3/1)}(z)^{-1} \in \mathcal{F}$ . So suppose we have the case (3).

Classes  $\pm 4C, 4F, 8D, 12M$ ; These classes are exceptional ones in Th. 3.2 and then we have  $\text{sgn}(\pi^{1/(r-1)}) = -1$ .

Classes  $\pm 6A$ ; We see from Table 3 of [2] that  $\pi \circ (3/1)$  is a Hauptmodule for  $18+$  or  $18+9$ .

Classes  $12B$  or  $12C$ ; By Table 3 of [2],  $\pi \circ (3/1)$  is the 2nd-harmonic of a Hauptmodule for  $18+9$  or  $18+$ . Then (3.1) follows from Lemma 2.3.

Classes  $+3A, -12A$  or  $+15B$ ;  $\pi \circ (3/1)$  is a Hauptmodule for  $9+, 36+$  or  $45+$  respectively by Table 3 of [2].

Now conjugacy classes  $-3A, 6B, +12A, -15B, \pm 21A$  and  $24A$  remain to be examined. Since  $\pi \circ (3/1)$  for  $6B$  or  $24A$  is the 2nd-harmonic of  $-3A$  or  $+12A$  respectively, it is sufficient to see (Lemma 2.3) that, for five classes  $-3A, +12A, -5A$  and  $\pm 21A, \eta_{\pi \circ (3/1)}(z)^{-1} \in \mathcal{F}$ . Among these classes, we will prove (3.1) for the class  $-3A$ , as other classes can be also dealt with quite similarly.

Let  $\pi$  be the Frame shape of the class  $-3A$  and let

$$f(z) = \eta_{\pi \circ (3/1)}(z)^{-1},$$

$$M = W_2 \begin{pmatrix} 1 & 0 \\ 6 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1/3 \\ 0 & 1 \end{pmatrix} \quad \text{and}$$

$$\Gamma = \langle \Gamma_0(18), M \rangle,$$

where  $W_2 = \begin{pmatrix} 18 & 1 \\ 18 & 2 \end{pmatrix}$  is an Atkin-Lehner's involution of  $\Gamma_0(18)$  (cf. § 2.2). Firstly it follows from Lemma 2.2 that  $f(z)$  is a modular function w. r. t.  $\Gamma_0(18)$  and then we see that, by using (2.2), (2.4), (2.6) and (2.7),  $f(Mz) = f(z)$ . Thus  $f(z)$  is a modular function w. r. t.  $\Gamma$ . Now we will apply Lemma 2.1 to show that  $f(z) \in \mathcal{F}$  and  $\Gamma$  is a group for  $f(z)$ . A representative of inequivalent cusps of  $\Gamma_0(18)$  is

$$0, 1/2, \pm 1/3, \pm 1/6, 1/9, 1/18$$

and these cusps are divided into two classes under  $\Gamma$ :

$$0 \sim -1/6 \sim 1/6 \sim 1/9 \quad \text{and} \quad 1/3 \sim -1/3 \sim 1/2 \sim 1/18 \sim i\infty.$$

Since, by (2.7),  $f(W_{18}(z)) = c/f(z)$  ( $c = \text{constant}$  and  $W_{18} = \begin{pmatrix} 0 & -1 \\ 18 & 0 \end{pmatrix}$ , an Atkin-Lehner's involution of  $\Gamma_0(18)$ ), we have  $f(1/18) = f(i\infty) = \infty$  and  $f(0) = 0$ . Thus  $z = i\infty$  is the unique pole of  $f(z)$  among inequivalent cusps of  $\Gamma$ . Then Lemma 2.1 yields that  $f(z) \in \mathcal{F}$  and  $\Gamma$  is a group for  $f(z)$ . This completes the proof of Th. 3.2.

In Table III of Appendix, we will give the list of  $\pi \circ (3/1)$  and groups for

$\eta_{\pi \circ (3/1)}(z)^{-1} \in \mathcal{F}$ . Also we will give the list of  $\pi \circ (r/1)$  ( $r > 3$ ) in Table IV, together with the list of  $\pi \circ s$ ,  $s$ -transformations of  $\pi$ .

**3.2.** In this paragraph, we give theorems analogous to Th. 3.2 for  $r$ -th harmonics and  $r$ -transformations. For that purpose, it is convenient to introduce the “ghost classes” of  $\cdot 0$ . We call the following generalized permutations “ghost classes” of  $\cdot 0$ :

| Name of classes | Frame shapes                        |
|-----------------|-------------------------------------|
| $\pm 9Z$        | $3^2 9^2, \quad 6^2 18^2 / 3^2 9^2$ |
| $16Z$           | $8.16$                              |
| $18Z$           | $6.18$                              |
| $\pm 25Z$       | $25/1, \quad 1.50/2.25$             |

We refer to [2],[3] and [8] for these classes. In [2], the class  $-25Z$  is denoted by  $50Z$ .

It is easy to see the following

**THEOREM 3.3.** *For all Frame shapes  $\pi$  of the ghost classes of  $\cdot 0$ , we have  $\eta_{\pi \circ (2/1)}(z)^{-1} \in \mathcal{F}$ . Also we have  $\eta_{\pi \circ (3/1)}(z)^{-1} \in \mathcal{F}$  for Frame shapes of the classes  $\pm 9Z$ .*

**THEOREM 3.4.** *Let  $s > 1$  be an integer with  $s+1 | 24$ . For each Frame shape  $\pi = \prod_t t^{r_t}$  of  $\cdot 0$  with  $s+1 | r_t$  for any  $t$ , the  $s$ -transformation  $\pi \circ s$  of  $\pi$  is a Frame shape of  $\cdot 0$  or that of a ghost class of  $\cdot 0$  except for the following cases:*

| $s$ | classes         |
|-----|-----------------|
| 2   | $4E, \quad 12F$ |
| 3   | $2B, \quad 6H$  |
| 5   | $4E$            |
| 11  | $2B$            |

*For these exceptional classes,  $\pi \circ s$  is an  $r$ -th harmonic of a Frame shape of  $\cdot 0$  for some  $r | 24$  and also we have  $\eta_{\pi \circ s}(z)^{-1} \in \mathcal{F}$  and  $\eta_{\pi \circ s \circ (2/1)}(z)^{-1} \in \mathcal{F}$ .*

The proof of this theorem is done just by inspection of Table I of Appendix and we don't need any facts from the theory of elliptic modular functions other than Lemma 2.3 (cf. Table IV of Appendix). Finally we have the following theorem for  $r$ -th harmonics of Frame shapes of  $\cdot 0$ :

**THEOREM 3.5.** *Let  $r > 1$  be an integer and  $\pi = \prod_t t^{r_t}$  be a Frame shape of  $\cdot 0$  with  $r | r_t$  for any  $t$ .*

- (1) *Let  $\sum_t r_t = 0$ . If  $\prod_t t^{r_t/r}$  is a rational square, we have  $\eta_{\pi/r}(z)^{-1} \in \mathcal{F}$ .*
- (2) *Let  $\sum_t r_t \neq 0$ . Then one of the following holds:*
  - (i)  *$\pi/r$  is a Frame shape of a conjugacy class or ghost class of  $\cdot 0$ ,*
  - (ii)  *$(\sum_t r_t)/r$  is odd, or*
  - (iii)  *$r=4$  and  $\pi=1^8 4^8 / 2^8$ , a Frame shape of the class  $-4A$ .*

Also the proof of this theorem is obtained by inspection of Table I of Ap-

pendix and Lemma 2.3.

**§ 4. Some examples of moonshines for finite groups.**

**4.1.** In this paragraph, we collect some lemmas on group characters.

LEMMA 4.1. Let  $G$  be a finite group and  $\sigma \mapsto \rho(\sigma)$  ( $\sigma \in G$ ) be an  $n$ -dimensional representation of  $G$  over the complex number field. Let  $\varepsilon_1(\sigma), \varepsilon_2(\sigma), \dots, \varepsilon_n(\sigma)$  be eigenvalues of  $\rho(\sigma)$ . Define functions  $a_k(\sigma)$  ( $1 \leq k < \infty$ ) and  $b_k(\sigma)$  ( $1 \leq k \leq n$ ) as follows:

$$\{(1 - \varepsilon_1(\sigma)q) \cdots (1 - \varepsilon_n(\sigma)q)\}^{-1} = \sum_{k=0}^{\infty} a_k(\sigma)q^k,$$

$$(1 + \varepsilon_1(\sigma)q) \cdots (1 + \varepsilon_n(\sigma)q) = \sum_{k=0}^n b_k(\sigma)q^k$$

where  $q$  is a variable. Then  $a_k(\sigma)$  and  $b_k(\sigma)$  are characters of  $G$ .

PROOF. It can be easily seen that, if  $V$  is a representation space of  $G$ ,  $a_k(\sigma)$  (resp.  $b_k(\sigma)$ ) is a character of the representation which  $\rho$  induces on the space of symmetric (resp. anti-symmetric) tensors of  $V$  of degree  $k$ , q. e. d.

LEMMA 4.2. Let  $G$  be a finite group and  $\sigma \mapsto \rho(\sigma)$  ( $\sigma \in G$ ) be a representation of  $G$  over  $\mathbf{Q}$ . For an element  $\sigma$  of  $G$  with Frame shape  $\prod_t t^{r_t}$ , define a function  $\chi_k(\sigma)$  as follows:

$$\eta_\sigma(z)^{-1} = q^{-m/24} \left( \sum_{k=0}^{\infty} \chi_k(\sigma)q^k \right),$$

where  $m = \sum_t r_t t$  is the degree of  $\rho$  and  $\eta_\sigma(z)$  is a function defined by (1.1). Then  $\chi_k(\sigma)$  is an ordinary character of  $G$ .

PROOF. Let

$$\xi(q) = \left( \prod_t (1 - q^t)^{r_t} \right)^{-1} = \sum_k a_k(\sigma)q^k.$$

Then, by Lemma 4.1,  $a_k(\sigma)$  is an ordinary character of  $G$ , as we have clearly

$$\prod_t (1 - q^t)^{r_t} = (1 - \varepsilon_1(\sigma)q) \cdots (1 - \varepsilon_m(\sigma)q)$$

where  $\varepsilon_1(\sigma), \dots, \varepsilon_m(\sigma)$  are eigenvalues of  $\rho(\sigma)$ . Then it follows from  $\eta_\sigma(z)^{-1} = \prod_{n=1}^{\infty} \xi(q^n)$  that the coefficients  $\chi_k(\sigma)$  of  $\eta_\sigma(z)^{-1}$  are ordinary characters, q. e. d.

LEMMA 4.3. Let  $\mathfrak{S}_n$  be the symmetric group of degree  $n$  and  $\xi(q) = \sum_{k=0}^{\infty} c_k q^k$  ( $c_0 = 1$ ) be a formal power series of  $q$  with non-negative integral coefficients, i. e.  $0 \leq c_k \in \mathbf{Z}$  ( $k = 1, 2, 3, \dots$ ). For an element  $\sigma$  of  $\mathfrak{S}_n$  with a cycle decomposition  $\prod_t t^{r_t}$ , define functions  $d_k(\sigma)$  as follows:

$$\prod_t \xi(q^t)^{r_t} = \sum_{k=0}^{\infty} d_k(\sigma)q^k.$$

Then  $d_k(\sigma)$  are characters of  $\mathfrak{S}_n$ .

PROOF. We may assume that  $\xi(q)$  is a polynomial, as  $d_k(\sigma)$  ( $1 \leq k \leq n$ ) are clearly the same as the ones which are obtained from a polynomial  $\sum_{k=0}^n c_k q^k$ . Thus we may assume  $\xi(q) = 1 + c_1 q + \dots + c_n q^n$ . Let

$$m = c_1 + c_2 + \dots + c_n$$

and  $x_0, x_1, \dots, x_m$  be  $m+1$  independent variables. For  $\sigma = \prod_t t^{r_t} \in \mathfrak{S}_n$ , define a function  $\chi_{k_0 k_1 \dots k_m}(\sigma)$  on  $\mathfrak{S}_n$  as follows:

$$\prod_t (x_0^t + \dots + x_m^t)^{r_t} = \sum_{k_0 + \dots + k_m = n} \chi_{k_0 k_1 \dots k_m}(\sigma) x_0^{k_0} x_1^{k_1} \dots x_m^{k_m}.$$

It is well known [5; §5.2] that  $\chi_{k_0 \dots k_m}(\sigma)$  is a character of  $\mathfrak{S}_n$  which is induced from the principal character of the subgroup  $\mathfrak{S}_{k_0} \times \mathfrak{S}_{k_1} \times \dots \times \mathfrak{S}_{k_m}$  of  $\mathfrak{S}_n$ . Now, by putting,

$$\begin{aligned} x_0 &= 1, & x_1 &= \dots = x_{c_1} = q, \\ x_{c_1+1} &= \dots = x_{c_1+c_2} = q^2, \\ &\dots \\ x_{c_1+\dots+c_{n-1}} &= \dots = x_m = q^n, \end{aligned}$$

we see that  $d_k(\sigma)$  is a linear combination of several induced characters  $\chi_{k_0 \dots k_m}(\sigma)$  with non-negative coefficients. Thus  $d_k(\sigma)$  is a character of  $\mathfrak{S}_n$ , q.e.d.

LEMMA 4.4. Let  $d > 1$  be an integer and let

$$\eta(2z)\eta(dz)/\eta(z)\eta(2dz) = q^{(1-d)/24} (1 + \sum_k c_k q^k).$$

Then we have  $c_k \geq 0$  ( $1 \leq k < \infty$ ).

PROOF. If  $k \leq 2dn$ , the  $c_k$  are the same as the coefficients of  $q$ -expansion of

$$\begin{aligned} & \prod_{h=1}^{dn} (1 - q^{2h}) \prod_{h=1}^{2n} (1 - q^{dh}) / \prod_{h=1}^{2dn} (1 - q^h) \prod_{h=1}^n (1 - q^{2dh}) \\ &= \left( \prod_{h=1}^n (1 - q^{d(2h-1)}) / (1 - q^{2h-1}) \right) \prod_{h=n}^{dn-1} (1 - q^{2h+1})^{-1}. \end{aligned}$$

Clearly all coefficients of  $q$ -expansion of the right hand side are non-negative.

4.2. Let  $G$  be a finite group and  $\mathfrak{F}$  be a class of elliptic modular functions defined in the introduction. A mapping from  $G$  to  $\mathfrak{F}$

$$G \ni \sigma \longmapsto j_\sigma(z) \in \mathfrak{F}$$

is said to satisfy *Moonshine condition* or simply to be a *moonshine* if every coefficient  $a_k(\sigma)$  ( $k \geq 1$ ) of a Fourier expansion of  $j_\sigma(z) = 1/q + \sum_{k=0}^\infty a_k(\sigma)q^k$  is a generalized character of  $G$ . Furthermore, if every coefficient  $a_k(\sigma)$  ( $k \geq 1$ ) of  $j_\sigma(z)$  is an ordinary character of  $G$ , this moonshine is called *proper*.

REMARK 4.1. In the above moonshine condition, the constant term  $a_0(\sigma)$  of a Fourier expansion of  $j_\sigma(z)$  need not necessarily to be any generalized character. But, in all moonshines which appear in this paper,  $a_0(\sigma)$  will be also a generalized character.

Let  $d > 1$  be a divisor of 24 and  $\sigma \mapsto \rho(\sigma)$  ( $\sigma \in G$ ) be a  $d$ -dimensional representation of  $G$  over  $\mathbb{Q}$ .

LEMMA 4.5. Let  $G$ ,  $\rho$  and  $d$  be as above and  $\pi = \prod_h h^{d_h}$  be a generalized permutation of degree  $24/d$ . For every element  $\sigma$  of  $G$  with Frame shape  $\sigma = \prod_t t^{r_t}$ , we put

$$j_\sigma^\pi(z) = \prod_t \left( \prod_h \eta(htz)^{d_h} \right)^{-r_t}.$$

Then  $j_\sigma^\pi(z)$  has a Fourier expansion of the form

$$(*) \quad j_\sigma^\pi(z) = \frac{1}{q} + \sum_{k=0}^{\infty} a_k(\sigma) q^k$$

and  $a_k(\sigma)$  ( $k=0, 1, 2, \dots$ ) are generalized characters of  $G$ .

PROOF. It is clear that  $j_\sigma^\pi(z)$  has a Fourier expansion of the form (\*). For each  $h$ , the coefficients of  $\prod_t \eta(htz)^{r_t}$  are generalized characters of  $G$  by Lemma 4.2. From this, the second statement follows, q.e.d.

Now we ask when we have  $j_\sigma^\pi(z) \in \mathcal{F}$ , i.e. a mapping  $\sigma \mapsto j_\sigma^\pi(z)$  is a moonshine.

THEOREM 4.6. Let  $G$ ,  $\rho$  and  $d$  be as above. Assume that,

(#) for any element  $\sigma$  of  $G$  with a Frame shape  $\prod_t t^{r_t}$ , a generalized permutation  $\prod_t t^{24r_t/d}$  of degree 24 is a Frame shape of  $\cdot 0$ .

(1) Let  $d^+ = 1 + (24/d)$ . Then a mapping

$$G \ni \sigma \longmapsto j_{d^+, \sigma}^+(z) = \left( \prod_t (\eta(d^+tz) / \eta(tz))^{r_t} \right)^{-1}$$

is a moonshine of  $G$ , where  $\prod_t t^{r_t}$  is a Frame shape of  $\sigma$ .

(2) Let  $d^- = (24/d) - 1$ . Then a mapping

$$G \ni \sigma \longmapsto j_{d^-, \sigma}^-(z) = \left( \prod_t (\eta(2tz)\eta(2d^-tz) / \eta(tz)\eta(d^-tz))^{r_t} \right)^{-1}$$

is also a moonshine of  $G$ , where  $\prod_t t^{r_t}$  is a Frame shape of  $\sigma$ .

PROOF. (1) Let  $\pi^+$  be a generalized permutation  $d^+/1$ . Then we have  $j_{d^+, \sigma}^+(z) = j_\sigma^{\pi^+}(z)$  in the notation of Lemma 4.5. Thus the coefficients of a Fourier expansion of  $j_{d^+, \sigma}^+(z)$  are generalized characters by Lemma 4.5. On the other hand, we have

$$j_{d^+, \sigma}^+(z) = \eta_{\sigma' \circ (d^+/1)}(z)^{-1}$$

where  $\sigma' = \prod_t t^{24r_t/d}$  and  $\sigma' \circ (d^+/1)$  is a  $(d^+/1)$ -transformation of  $\sigma'$ . Then it follows from (#) and Th. 3.2 that  $j_{d^+, \sigma}^+(z) \in \mathcal{F}$ .

(2) Let  $\pi^-$  be a generalized permutation  $(2 \cdot 2d^-)/(1 \cdot d^-)$ . Then we have, in

the notation of Lemma 4.5,

$$j_{\bar{a}, \sigma}^-(z) = j_{\bar{\sigma}}^-(z),$$

and also

$$j_{\bar{a}, \sigma}^-(z) = \eta_{\sigma' \circ a^{-1} \circ (2/1)}(z)^{-1}.$$

Then (2) follows from Lemma 4.5, (#) and Th. 3.3, q.e.d.

REMARK 4.2. Moonshines in Th. 4.6 are not proper, i.e. Fourier coefficients of  $j_{\bar{a}, \sigma}^+(z)$  and  $j_{\bar{a}, \sigma}^-(z)$  are not necessarily ordinary characters. Some examples of proper moonshines will be given in the next paragraph §4.3.

Let  $p$  be a prime with  $p+1|24$  and  $d>1$  be an integer with  $d|24$ . Many  $d$ -dimensional rational representations of  $SL(2, p)$  satisfy the condition (#) of Th. 4.6. In the following, we will give examples of such representations and Frame shapes w.r.t. them for  $p=5, 7$ .

$SL(2, 5)$

|   |           |                   |          |                              |
|---|-----------|-------------------|----------|------------------------------|
| $\pm 1A$  | $2A$      | $\pm 3A$          | $\pm 5A$ |                              |
| $1^4$   | $2^2$     | 1.3               | 5/1      | absolutely irreducible       |
| $1^6$   | $1^2 2^2$ | $3^2$             | 1.5      | a permutation representation |
| $1^6$   | $2^4/1^2$ | $3^2$             | 1.5      | absolutely irreducible       |
| $\left\{ \begin{array}{l} 1^4 \\ 2^4/1^4 \end{array} \right.$ | $4^2/2^2$ | 1.3               | 5/1      | absolutely irreducible       |
|   |           | 2.6/1.3           | 1.10/2.5 |                              |
| $\left\{ \begin{array}{l} 1^4 \\ 2^4/1^4 \end{array} \right.$ | $4^2/2^2$ | $3^2/1^2$         | 5/1      | not absolutely irreducible   |
|   |           | $1^2 6^2/2^2 3^2$ | 1.10/2.5 |                              |
| $\left\{ \begin{array}{l} 1^6 \\ 2^6/1^6 \end{array} \right.$ | $4^3/2^3$ | $3^2$             | 1.5      | not absolutely irreducible   |
|   |           | $6^2/3^2$         | 2.10/1.5 |                              |

$SL(2, 7)$

|  |           |                   |                    |           |                            |
|--|-----------|-------------------|--------------------|-----------|----------------------------|
| $\pm 1A$   | $2A$      | $\pm 3AB$         | $4AB$              | $\pm 7AB$ |                            |
| $1^6$  | $1^2 2^2$ | $3^2$             | 2.4                | 7/1       | absolutely irreducible     |
| $1^6$  | $2^4/1^2$ | $3^2$             | $1^2 4^2/2^2$      | 7/1       | not absolutely irreducible |
| $1^8$  | $2^4$     | $1^2 3^2$         | $4^2$              | 1.7       | permutation representation |
| $1^8$  | $2^4$     | $3^3/1$           | $4^2$              | 1.7       | absolutely irreducible     |
| $\left\{ \begin{array}{l} 1^8 \\ 2^8/1^8 \end{array} \right.$          | $4^4/2^4$ | $3^3/1$           | $8^2/4^2$          | 1.7       | absolutely irreducible     |
|  |           | $1.6^3/2.3^3$     | 2.14/1.7           |           |                            |
| $\left\{ \begin{array}{l} 1^8 \\ 2^8/1^8 \end{array} \right.$          | $4^4/2^4$ | $1^2 3^2$         | $8^2/4^2$          | 1.7       | not absolutely irreducible |
|  |           | $2^2 6^2/1^2/3^2$ | 2.14/1.7           |           |                            |
| $\left\{ \begin{array}{l} 1^{12} \\ 2^{12}/1^{12} \end{array} \right.$ | $8^3/4^3$ | $3^4$             | $4^6/2^6$          | $7^2/1^2$ | not absolutely irreducible |
|  |           | $6^4/3^4$         | $1^2 14^2/2^2 7^2$ |           |                            |

Notations: For  $SL(2, p) \ni \sigma$ , if homomorphic image of  $\sigma$  in  $PSL(2, p)$  is of order  $n$ , conjugate class of  $\sigma$  is denoted by  $nA$  or  $\pm nA$ , according as  $\sigma$  and  $-\sigma$  are conjugate in  $SL(2, p)$  or not. And  $nAB$  (resp.  $\pm nAB$ ) expresses that there exist two conjugate classes  $nA$  and  $nB$  (resp.  $\pm nA$  and  $\pm nB$ ) of  $PSL(2, p)$  of order  $n$  with the same Frame shapes.

REMARK 4.3. Let  $\rho$  be one of representations of  $SL(2, 5)$  of degree 6 in the above table. Then if  $\sigma$  is a class of order 5 and so its Frame shape is 1.5, we have  $j_{6, \sigma}^+(z) = \eta(z)/\eta(25z)$  which is a ghost element of Monster's moonshine. Similarly another ghost element  $\eta(2z)\eta(25z)/\eta(z)\eta(50z)$  of Monster's moonshine [2] also appears in the moonshines  $\sigma \mapsto j_{4, \sigma}^-(z)$  which are obtained from any one of 4-dimensional representations of  $SL(2, 5)$ .

REMARK 4.4.  $SL(2, 9)$  has representations of degree 4 and 6 with the following Frame shapes :

|   |           |          |                   |      |          |
|---|-----------|----------|-------------------|------|----------|
| $\pm 1A$  | $2A$      | $\pm 3A$ | $\pm 3B$          | $4A$ | $\pm 5A$ |
| $\left\{ \begin{array}{l} 1^4 \\ 2^4/1^4 \end{array} \right.$ | $4^2/2^2$ | 1.3      | $3^2/1^2$         | 8/4  | 5/1      |
| $1^6$   | $1^{2^2}$ | 2.6/1.3  | $1^{2^6}/2^2 3^2$ | 2.4  | 1.10/2.5 |
|   |           | $1^3$    | $3^2$             |      | 1.5      |

The one of degree 6 is a natural permutation representation of  $PSL(2, 9) \simeq \mathfrak{A}_6$  (=the Alternating group of degree 6). The representation of degree 4 satisfies the condition (#) of Th. 4.6, while the one of degree 6 does not, as  $(1^3 3)^4 = 1^{12} 3^4$  is not a Frame shape of  $\cdot 0$ . It is easy to see that, if  $\sigma = (1^3 3)^4$ ,  $\eta_{\sigma \circ (5/1)}(z)^{-1}$  does not satisfy the second condition in the definition of  $\mathfrak{F}$ . Thus the representation of degree 6 does not yield a moonshine. Similarly the permutation representation of Mathieu group  $M_{12}$  of degree 12 also does not yield a moonshine. In fact, there are elements of  $M_{12}$  with cycle decompositions  $1^4 2^2$  and  $1^2 2.8$ . These permutations do not satisfy (#) of Th. 4.6 and it can be shown that, if  $\sigma$  is one of these permutations,  $\eta_{\sigma \circ (3/1)}(z)^{-1} \notin F$ , where  $\sigma^2 = 1^8 4^4$  or  $1^4 2^2 8^2$ .

4.3. In this paragraph, we give some examples of proper moonshines for Mathieu group  $M_{24}$  and  $PSL(2, p)$  ( $p+1|24$ ).

LEMMA 4.7. Let  $\sigma = \prod_i t^{r_i}$  be a cycle decomposition (=Frame shape) of an element  $\sigma$  of  $M_{24}$  w.r.t. the natural permutation representation of  $M_{24}$ . Then  $(2/1)$ -transformation of  $\sigma$  is a Frame shape of  $\cdot 0$ . More explicitly, we have the following table:

|                      |               |            |            |            |        |               |            |        |           |                   |       |           |
|----------------------|---------------|------------|------------|------------|--------|---------------|------------|--------|-----------|-------------------|-------|-----------|
| $\sigma$             | $1^8$         | $1^8 2^8$  | $2^{12}$   | $1^6 3^6$  | $3^8$  | $1^4 2^2 4^4$ | $2^4 4^4$  | $4^6$  | $1^4 5^4$ | $1^2 2^2 3^2 6^2$ | $6^4$ | $1^3 7^3$ |
| $\sigma \circ (2/1)$ | $-2A$         | $4A$       | $2B$       | $-3A$      | $-3D$  | $8C$          | $8A$       | $4E$   | $-5B$     | $12E$             | $6H$  | $-7B$     |
| $\sigma$             | $1^2 2.4.8^2$ | $2^2 10^2$ | $1^2 11^2$ | $2.4.6.12$ | $12^2$ | $1.2.7.14$    | $1.3.5.15$ | $3.21$ | $1.23$    |                   |       |           |
| $\sigma \circ (2/1)$ | $16B$         | $10C$      | $-11B$     | $24C$      | $12L$  | $28B$         | $-30B$     | $-21C$ | $-23AB$   |                   |       |           |

where the second line denotes conjugacy classes of  $\cdot 0$  with Frame shapes  $\sigma \circ (2/1)$ .

PROOF. This can be seen immediately from Table I of Appendix.

THEOREM 4.8. For an element  $\sigma$  of  $M_{24}$  with a cycle decomposition  $\prod_i t^{r_i}$ , we put

$$j_\sigma(z) = \prod_t (\eta(2tz)^2 / \eta(tz)\eta(4tz))^{rt}.$$

Then a mapping

$$M_{24} \ni \sigma \longmapsto j_\sigma(z)$$

is a proper moonshine of  $M_{24}$ .

PROOF. Let  $\sigma' = \sigma \circ (2/1) \circ (2/1)$ . Then we have

$$\sigma' = \prod_t t^{rt}(4t)^{rt}(2t)^{-2rt}$$

and

$$j_\sigma(z) = \eta_{\sigma'}(z)^{-1}.$$

Then it follows from Lemma 4.7 and Th. 3.2 that  $j_\sigma(z) \in \mathcal{F}$ . Furthermore, we see from Lemma 4.3 and 4.4 that this moonshine is proper, q.e.d.

LEMMA 4.9. Let  $p$  be a prime with  $p+1 \mid 24$  and  $\rho_p$  be a permutation representation of  $PSL(2, p)$  of degree  $p+1$  on a projective line over  $F_p$ , a finite field of  $p$  elements. For an element  $\sigma$  of  $PSL(2, p)$ , let  $\sigma = \prod_t t^{rt}$  be a Frame shape of  $\sigma$  w.r.t.  $\rho_p$ . Then a generalized permutation  $\prod_t t^{24rt/(p+1)}$  of degree 24 is a Frame shape of  $M_{24}$ .

PROOF. It is easy to check this for each  $p=2, 3, 5, 7, 11$  and  $23$ . See the table in §4.2 for  $p=5, 7$ .

THEOREM 4.10. Notations being as in Lemma 4.9, we put, for  $PSL(2, p) \ni \sigma$ ,

$$j_{p,\sigma}(z) = \prod_t (\eta(2tz)\eta(dtz) / \eta(2dtz)\eta(tz))^{rt}$$

where  $d=24/(p+1)+1$  and  $\prod_t t^{rt}$  is a Frame shape of  $\sigma$ . Then a mapping

$$\sigma \longmapsto j_{p,\sigma}(z)$$

is a proper moonshine of  $PSL(2, p)$ .

PROOF. Let  $\sigma' = \prod_t t^{24rt/(p+1)}$ . Then we have

$$\sigma' \circ (d/1) \circ (2/1) = \prod_t (2dt)^{rt} t^{rt} (2t)^{-rt} (dt)^{-rt}$$

and so

$$j_{p,\sigma}(z) = \eta_{\sigma' \circ (d/1) \circ (2/1)}(z)^{-1}.$$

By Lemma 4.9,  $\sigma'$  is a Frame shape of  $M_{24}$  and then, by Lemma 4.7 and Th. 3.2,  $j_{p,\sigma}(z) \in \mathcal{F}$ . By Lemma 4.3 and 4.4, this moonshine is proper, q.e.d.

REMARK 4.5.  $j_\sigma(z)$  and  $j_{23,\sigma}(z)$  being in Th. 4.8 and 4.10 respectively, we have  $j_\sigma(z) = j_{23,\sigma}(z)$ . Since  $PSL(2, 23)$  is a subgroup of  $M_{24}$  and the embedding is unique up to conjugation, a moonshine  $\sigma \mapsto j_{23,\sigma}(z)$  of  $PSL(2, 23)$  is a restriction of a moonshine  $\sigma \mapsto j_\sigma(z)$  of  $M_{24}$ .



4.4. Here we will make two remarks on Th. 4.6, 4.8 and 4.10. In these theorems, we constructed moonshine of a finite group  $G$  by using a representation of  $G$  of degree  $d$  satisfying the condition (#) of Th. 4.6 and one of transformations  $d^+/1$ ,  $d^-\circ(2/1)$  and  $(2/1)\circ(2/1)$  (in this case,  $d=24$ ) of “degree”  $24/d$ . These are, however, not all transformations we can use to construct moonshines. In fact, for  $d=4$  or  $6$ , we can also use the following transformations of “degree”  $24/d$ :

$$\begin{aligned} d=4 & \quad 5\circ(2/1)\circ(2/1), & (3/1)\circ(4/1) \\ d=6 & \quad 3\circ(2/1)\circ(2/1), & (3/1)\circ(3/1). \end{aligned}$$

But these transformations do not always yield a moonshine. For example, for a representation of  $SL(2, 5)$  or  $SL(2, 7)$  in which a Frame shape  $2^4/1^2$  appears (§ 4.2), a transformation  $3\circ(2/1)\circ(2/1)$  does not yield a moonshine, because we can easily see that  $\pi=(2^{16}/1^8)\circ 3\circ(2/1)\circ(2/1)=2^8 6^8 8^4 24^4/1^2 3^2 4^{10} 12^{10}$  but  $\eta_\pi(z)^{-1} \notin \mathfrak{F}$ .

The second remark is that  $j_\sigma(z)$  in Th. 4.8 can be related to some even lattice.

Let  $V$  be a 24-dimensional vector space over  $\mathbf{Q}$  and  $e_i$  ( $1 \leq i \leq 24$ ) be a basis of  $V$ . Furthermore let  $(u, v)$  ( $u, v \in V$ ) be an inner product of  $V$  with  $(e_i, e_j) = 2\delta_{ij}$ . Set  $L = \sum_{i=1}^{24} \mathbf{Z}e_i \in V$ . Then  $L$  is an even lattice of  $V$  on which the Mathieu group  $M_{24}$  acts in such a way that  $e_i^\sigma = e_{\sigma(i)}$  ( $\sigma \in M_{24}$ ). For each  $\sigma \in M_{24}$ , we put

$$L_\sigma = \{v \in L \mid v^\sigma = v\}$$

and

$$\Theta_\sigma(z) = \sum_{v \in L_\sigma} e^{\pi i(v, v)z} \quad (\Theta\text{-series of } L_\sigma).$$

THEOREM 4.11.  $j_\sigma(z)$  being as in Th. 4.8, we have

$$(*) \quad \Theta_\sigma(z) = j_\sigma(z)^2 \eta_\sigma(2z)$$

where  $\prod_t t^{r_t}$  is a cycle decomposition of  $\sigma$  and  $\eta_\sigma(z) = \prod_t \eta(tz)^{r_t}$ .

PROOF. Let  $\theta(z) = \sum_{x \in \mathbf{Z}} e^{2\pi i x^2 z}$ . It is easy to see that  $\Theta_\sigma(z) = \prod_t \theta(tz)^{r_t}$ . Then (\*) follows from the identity  $\theta(z) = \eta(2z)^5 / \eta(z)^2 \eta(4z)^2$ , q. e. d.

**Appendix. Table I~IV.**

For conjugacy classes of  $\cdot 0$ , we use the following notations in § 3~§ 4 and Table I~IV. The heading column  $nA, nB, \dots$  of Table I are the Atlas names of conjugacy classes of the Conway’s simple group  $\cdot 1$  (=the factor group of  $\cdot 0$  by its center  $\langle \pm 1 \rangle$ ) [11; Table 1], i.e. conjugacy classes of  $\cdot 1$  of order  $n$  are named  $nA, nB, \dots$  in descending order of their centralizer sizes.

Case (1). If the inverse image in  $\cdot 0$  of a class  $nX$  ( $X=A, B, \dots$ ) of  $\cdot 1$  is a conjugacy class of  $\cdot 0$ , this class is also denoted by  $nX$ .

Case (2). If the inverse image in  $\cdot 0$  of a class  $nX$  of  $\cdot 1$  consists of two conjugacy classes of  $\cdot 0$ , these classes are denoted by  $+nX$  and  $-nX$ .

Table I; In case (1), Frame shape of a class  $nX$  is written after the heading column and in case (2), firstly Frame shape of  $+nX$  and then that of  $-nX$  are written. For a Frame shape  $\pi = \prod_t t^{r_t}$  with  $\sum_t r_t = 0$ , a group for  $\eta_\pi(z)^{-1}$  is given in parenthesis after the Frame shape by using notations of Table 2 and 3 of [2] (cf. also §2.1 of this paper).

Table II~IV; Let  $\pi$  be a Frame shape of  $\cdot 0$ . If  $\pi \circ (r/1)$  (resp.  $\pi \circ s$ ) is a Frame shape of a conjugacy class  $nX$  ( $+nX$  or  $-nX$ ) of  $\cdot 0$ ,  $\pi \circ (r/1)$  (resp.  $\pi \circ s$ ) is also denoted by  $nX$  ( $nX$  or  $-nX$ ). Note that  $+$  of  $+nX$  is omitted. And if  $\pi \circ (r/1)$  (resp.  $\pi \circ s$ ) is the  $m$ -th harmonic of a Frame shape of a class  $nX$  ( $+nX$  or  $-nX$ ),  $\pi \circ (r/1)$  (resp.  $\pi \circ s$ ) is denoted by  $nX/m$  ( $nX/m$  or  $-nX/m$ ).

Some of  $(r/1)$ -transformations are expressed by generalized permutations with symbols (?). These are exceptional classes in Th. 3.2.

In Table II and III, groups for  $\eta_{\pi \circ (2/1)}(z)^{-1}$  or  $\eta_{\pi \circ (3/1)}(z)^{-1}$  are given in parenthesis after  $\pi \circ (2/1)$  or  $\pi \circ (3/1)$ . Then the following notations are used:

$$T = \begin{pmatrix} 1 & 1/2 \\ 0 & 1 \end{pmatrix}, \quad S = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}, \quad T_n = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix},$$

$$\left(\frac{1}{h}n\right) = \begin{pmatrix} 1 & 1/h \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}, \quad \left(n\frac{1}{h}\right) = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix} \begin{pmatrix} 1 & 1/h \\ 0 & 1 \end{pmatrix},$$

$W_Q$  = an Atkin-Lehner's involution of  $\Gamma_0(N)$  for some  $N$ .

A notation like " $N+Q, \left(h\frac{1}{n}\right), \dots$ " denotes  $\langle \Gamma_0(N), W_Q, \begin{pmatrix} 1 & 1/h \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}, \dots \rangle$ .

## References

- [1] J. H. Conway, Three Lectures on Exceptional Groups (Chap. VII of Finite Simple Groups, edited by M. B. Powell and G. Higman), Academic Press, London and New York, 1971.
- [2] J. H. Conway and S. P. Norton, Monstrous moonshine, Bull. London Math. Soc., 11 (1979), 308-339.
- [3] M. Koike, On Mackay's conjecture, Nagoya Math. J., 95 (1984), 85-90.
- [4] M. Koike, Moonshine for  $PSL_2(p)$ , preprint.
- [5] M. Koike, a personal communication (a table of explicit expressions of  $\eta_\pi(z)$  as Eisenstein series).
- [6] M. I. Knopp, Modular Functions in Analytic Number Theory, Markham Mathematics Series, 1970.
- [7] D. E. Littlewood, The Theory of Group Characters and Matrix Representations of Groups, 2nd Edition, Oxford Univ. Press, 1950.
- [8] G. Mason,  $M_{24}$  and certain automorphic forms, preprint.
- [9] M. Newman, Construction and application of a class of modular functions II, Proc. London Math. Soc. (3), 9 (1959), 373-387.
- [10] L. Queen, Modular functions arising from some finite groups, Math. Comp., 37 (1981), 547-580.
- [11] R. A. Wilson, The maximal subgroups of Conway's group  $Co_1$ , J. Algebra, 85 (1983), 144-165.

Table I. Frame shapes of conjugacy classes of  $\cdot 0$ .

|    |                             |                             |                                     |
|----|-----------------------------|-----------------------------|-------------------------------------|
| 1A | $1^{24}$ ,                  |                             | $2^{24}/1^{24}$ (2-)                |
| 2A | $1^8 2^8$ ,                 |                             | $2^{16}/1^8$                        |
| 2B |                             | $4^{12}/2^{12}$ (4/2-)      |                                     |
| 2C |                             | $2^{12}$                    |                                     |
| 3A | $3^{12}/1^{12}$ (3-),       |                             | $1^{12} 6^{12}/2^{12} 3^{12}$ (6+6) |
| 3B | $1^6 3^6$ ,                 |                             | $2^6 6^6/1^6 3^6$ (6+3)             |
| 3C | $3^9/1^3$ ,                 |                             | $1^3 6^9/2^3 3^9$ (6-)              |
| 3D | $3^8$ ,                     |                             | $6^8/3^8$ (6/3-)                    |
| 4A | $4^8/1^8$ (4-),             |                             | $1^8 4^8/2^8$                       |
| 4B |                             | $4^8/2^4$                   |                                     |
| 4C | $1^4 2^2 4^4$ ,             |                             | $2^6 4^4/1^4$                       |
| 4D |                             | $2^4 4^4$                   |                                     |
| 4E |                             | $8^6/4^6$ (8/4-)            |                                     |
| 4F |                             | $4^6$                       |                                     |
| 5A | $5^6/1^6$ (5-),             |                             | $1^6 10^6/2^6 5^6$ (10+10)          |
| 5B | $1^4 5^4$ ,                 |                             | $2^4 10^4/1^4 5^4$ (10+5)           |
| 5C | $5^5/1$ ,                   |                             | $1 \cdot 10^5/2 \cdot 5^5$ (10-)    |
| 6A | $3^4 6^4/1^4 2^4$ (6+2),    |                             | $1^4 6^8/2^8 3^4$ (6-)              |
| 6B |                             | $2^6 12^6/4^6 6^6$ (12/2+6) |                                     |
| 6C | $1^4 2 \cdot 6^5/3^4$ ,     |                             | $2^5 3^4 6/1^4$                     |
| 6D | $2 \cdot 6^5/1^5 3$ (6-),   |                             | $1^5 3 \cdot 6^4/2^4$               |
| 6E | $1^2 2^2 3^2 6^2$ ,         |                             | $2^4 6^4/1^2 3^2$                   |
| 6F | $3^3 6^3/1 \cdot 2$ ,       |                             | $1 \cdot 6^6/2^2 3^3$               |
| 6G |                             | $2^3 6^3$                   |                                     |
| 6H |                             | $12^4/6^4$ (12/6-)          |                                     |
| 6I |                             | $6^4$                       |                                     |
| 7A | $7^4/1^4$ (7-),             |                             | $1^4 14^4/2^4 7^4$ (14+14)          |
| 7B | $1^3 7^3$ ,                 |                             | $2^3 14^3/1^3 7^3$ (14+7)           |
| 8A |                             | $8^4/2^4$ (8/2-)            |                                     |
| 8B |                             | $2^4 8^4/4^4$               |                                     |
| 8C | $2^2 8^4/1^4 4^2$ (8-),     |                             | $1^4 8^4/2^2 4^2$                   |
| 8D |                             | $8^4/4^2$                   |                                     |
| 8E | $1^2 2 \cdot 4 \cdot 8^2$ , |                             | $2^3 4 \cdot 8^2/1^2$               |
| 8F |                             | $4^2 8^2$                   |                                     |

Table I (continued)

|     |                               |  |
|-----|-------------------------------|--|
| 9A  | $9^3/1^3$ (9-),               | $1^3 18^3/2^3 9^3$ (18+18)                     |
| 9B  | $9^3/3$ ,                     | $3.18^3/6.9^3$ (18-)                           |
| 9C  | $1^3 9^3/3^2$ ,               | $2^3 3^2 18^3/1^3 6^2 9^3$ (18+9)              |
| 10A | $5^2 10^2/1^2 2^2$ (10+2),    | $1^2 10^4/2^4 5^2$ (10-)                       |
| 10B |                               | $2^3 20^3/4^3 10^3$ (20/2+10)                  |
| 10C |                               | $4^2 20^2/2^2 10^2$ (20/2+5)                   |
| 10D | $1^2 2.10^3/5^2$ ,            | $2^3 5^2 10/1^2$                               |
| 10E | $2.10^3/1^3 5$ (10-),         | $1^3 5.10^2/2^2$                               |
| 10F |                               | $2^2 10^2$                                     |
| 11A | $1^2 11^2$ ,                  | $2^2 22^2/1^2 11^2$ (22+11)                    |
| 12A | $1^4 12^4/3^4 4^4$ (12+12),   | $2^4 3^4 12^4/1^4 4^4 6^4$ (12+4)              |
| 12B |                               | $2^2 12^4/4^4 6^2$ (12-)                       |
| 12C |                               | $6^2 12^2/2^2 4^2$ (12/2+2)                    |
| 12D | $1.12^3/3^3 4$ (12-),         | $2.3^3 12^3/1.4.6^3$                           |
| 12E | $4^2 12^2/1^2 3^2$ (12+3),    | $1^2 3^2 4^2 12^2/2^2 6^2$                     |
| 12F |                               | $4^3 24^3/8^3 12^3$ (24/4+6)                   |
| 12G |                               | $4^2 12^2/2.6$                                 |
| 12H | $2^3 6.12^2/1.3.4^2$ ,        | $1.2^2 3.12^2/4^2$                             |
| 12I | $1^2 4.6^2 12/3^2$ ,          | $2^2 3^2 4.12/1^2$                             |
| 12J |                               | 2.4.6.12                                       |
| 12K | $2^2 3.12^3/1^3 4.6^2$ (12-), | $1^3 12^3/2.3.4.6$                             |
| 12L |                               | $24^2 12^2$ (24/12-)                           |
| 12M |                               | $12^2$   |
| 13A | $13^2/1^2$ (13-),             | $1^2 26^2/2^2 13^2$ (26+26)                    |
| 14A |                               | $2^2 28^2/4^2 14^2$ (28/2+14)                  |
| 14B | 1.2.7.14,                     | $2^2 14^2/1.7$                                 |
| 15A | $1^3 15^3/3^3 5^3$ (15+15),   | $2^3 3^3 5^3 30^3/1^3 6^3 10^3 15^3$ (30+6,10) |
| 15B | $3^2 15^2/1^2 5^2$ (15+5),    | $1^2 5^2 6^2 30^2/2^2 3^2 10^2 15^2$ (30+5,6)  |
| 15C | $15^2 3^2$ (15/3-),           | $3^2 30^2/6^2 15^2$ (30/3+10)                  |
| 15D | 1.3.5.15,                     | $2.6.10.30/1.3.5.15$ (30+3,5)                  |
| 15E | $1^2 15^2/3.5$ ,              | $2^2 3.5.30^2/1^2 6.10.15^2$ (30+15)           |
| 16A |                               | $2^2 16^2/4.8$                                 |
| 16B | $2.16^2/1^2 8$ (16-),         | $1^2 16^2/2.8$                                 |
| 18A | $9.18/1.2$ (18+2),            | $1.18^2/2^2 9$ (18-)                           |
| 18B | $2.3.18^2/1^2 6.9$ (18-),     | $1^2 9.18/2.3$                                 |
| 18C | $1.2.18^2/6.9$ ,              | $2^2 9.18/1.6$                                 |

Table I (continued)

|     |                                     |  |
|-----|-------------------------------------|--|
| 20A | $1^2 20^2 / 4^2 5^2$ (20+20),       | $2^2 5^2 20^2 / 1^2 4^2 10^2$ (20+4)             |
| 20B | 4.20                                |  |
| 20C | 1.2.10.20/4.5,                      | $2^2 5.20/1.4$                                   |
| 21A | $1^2 21^2 / 3^2 7^2$ (21+21),       | $2^2 3^2 7^2 42^2 / 1^2 6^2 14^2 21^2$ (42+6,14) |
| 21B | 7.21/1.3 (21+3),                    | 1.3.14.42/2.6.7.21 (42+3,14)                     |
| 21C | 3.21                                | 6.42/3.21 (42/3+7)                               |
| 22A | 2.22                                |  |
| 23A | 1.23,                               | 2.46/1.23 (46+23)                                |
| 23B | 1.23,                               | 2.46/1.23 (46+23)                                |
| 24A | $2^2 24^2 / 6^2 8^2$ (24/2+12)      |  |
| 24B | $1^2 4.6.24^2 / 2.3^2 8^2$ (24+24), | $2.3^2 4.24^2 / 1^2 6.8^2$ (24+8)                |
| 24C | 8.24/2.6 (24/2+3)                   |  |
| 24D | 12.24/4.8 (24/4+2)                  |  |
| 24E | 2.6.8.24/4.12                       |  |
| 24F | 1.4.6.24/3.8,                       | 2.3.4.24/1.8                                     |
| 26A | 2.52/4.26 (52/2+26)                 |  |
| 28A | 4.28/1.7 (28+7),                    | 1.4.7.28/2.14                                    |
| 28B | 4.56/8.28 (56/4+14)                 |  |
| 30A | 1.2.15.30/3.5.6.10 (30+2,15),       | $2^2 3.5.30^2 / 1.6^2 10^2$ (30+15)              |
| 30B | 2.10.12.60/4.6.20.30 (60/2+5,6)     |  |
| 30C | 6.60/12.30 (60/6+10)                |  |
| 30D | 1.6.10.15/3.5,                      | 2.3.5.30/1.15                                    |
| 30E | 2.30/3.5 (30+15),                   | 2.3.5.30/6.10                                    |
| 33A | 3.33/1.11 (33+11),                  | 1.6.11.66/2.3.22.33 (66+6,11)                    |
| 35A | 1.35/5.7 (35+35),                   | 2.5.7.70/1.10.14.35 (70+10,14)                   |
| 36A | 1.36/4.9 (36+36),                   | 2.9.36/1.4.18 (36+4)                             |
| 39A | 1.39/3.13 (39+39),                  | 2.3.13.78/1.6.26.39 (78+6,26)                    |
| 39B | 1.39/3.13 (39+39),                  | 2.3.13.78/1.6.26.39 (78+6,26)                    |
| 40A | 2.40/8.10 (40/2+20)                 |  |
| 42A | 4.6.14.84/2.12.28.42 (84/2+6,14)    |  |
| 60A | 3.4.5.60/1.12.15.20 (60+12,15),     | 1.4.6.10.15.60/2.3.5.12.20.30 (60+4,15)          |

Table II. (2/1)-transformations of Frame shapes of  $\cdot 0$ .

|    |  |  |
|----|--|--|
| 1A | -1A (2-),  | $1^{24}4^{24}/2^{48}$ (4+)                                     |
| 2A | 4A (4-),   | $1^84^{16}/2^{24}$ (4-)  |
| 2B | $-1A^\circ(2/1)/2$ (16+16, $(\frac{1}{2}8)$ )                        |  |
| 2C | 2B (8+ $(\frac{1}{2}4)$ )  |  |
| 3A | -3A (6+6),   | $2^{24}3^{12}12^{12}/1^{12}4^{12}6^{24}$ ((6+6) <sup>T</sup> ) |
| 3B | -3B (6+3),   | $1^63^64^612^6/2^{12}6^{12}$ (12+)                             |
| 3C | -3C (6-),  | $2^63^912^9/1^34^36^{18}$ (12+4)                               |
| 3D | -3D (18+9, $(-\frac{1}{3}6)$ ),                                      | $-1A^\circ(2/1)/3$ (36+4, 9, $(\frac{1}{3}12)$ )               |
| 4A | $1^88^8/2^84^8$ (8+),  | $2^{16}8^8/1^84^{16}$ (8+8 <sup>T</sup> )                      |
| 4B | $-2A^\circ(2/1)/2$ (8-)  |  |
| 4C | 8C (8-),   | $1^44^28^4/2^{10}$ (8-)  |
| 4D | 8A (16+ $(\frac{1}{2}8)$ )   |  |
| 4E | $-1A^\circ(2/1)/4$ (64+64, $(\frac{1}{4}16)$ )                       |  |
| 4F | 4E (32+ $(-\frac{1}{4}8)$ , $(\frac{1}{2}16)$ )                      |  |
| 5A | -5A (10+10),   | $2^{12}5^620^6/1^64^610^{12}$ ((10+10) <sup>T</sup> )          |
| 5B | -5B (10+5),  | $1^44^45^420^4/2^810^8$ (20+)                                  |
| 5C | -5C (10-),   | $2^25^520^5/1.4.10^{10}$ (20+4)                                |
| 6A | 12A (12+12),   | $2^43^412^4/1^44^46^4$ (12+12 <sup>T</sup> )                   |
| 6B | $-3A^\circ(2/1)/2$ (24+24, $(\frac{1}{2}12)$ ) <sup>ST12</sup>       |  |
| 6C | $2^33^4.4.12^5/1^46^9$ (12+12 <sup>TW4</sup> ),                      | $1^44^56^312/2^93^4$ (12+12 <sup>TW4</sup> )                   |
| 6D | $1^53.4.12^5/2^66^6$ (12+12),  | $2^912^4/1^53.4^46^3$ (12+12 <sup>T</sup> )                    |
| 6E | 12E (12+3),  | $1^23^24^412^4/2^66^6$ (12+3)                                  |
| 6F | 12D (12-),   | $2^33^312^6/1.4^26^9$ (12-)                                    |
| 6G | $-3B/2$ (24+3, $(\frac{1}{2}12)$ )                                   |  |
| 6H | $-1A^\circ(2/1)/6$ (144+144, $(\frac{1}{6}24)$ , $(\frac{1}{3}48)$ ) |  |
| 6I | 6H (72+9, $(-\frac{1}{6}12)$ , $(-\frac{1}{3}24)$ )                  |  |
| 7A | -7A (14+14),   | $2^87^428^4/1^44^414^8$ ((14+14) <sup>T</sup> )                |
| 7B | -7B (14+7),  | $1^34^37^328^3/2^614^6$ (28+)                                  |
| 8A | $4A^\circ(2/1)/2$ (32+32, $(\frac{1}{2}16)$ )                        |  |
| 8B | $-4A^\circ(2/1)/2$ (32+32 <sup>T8</sup> , $(\frac{1}{2}16)$ )        |  |
| 8C | $1^44^416^4/2^68^6$ (16+),   | $2^616^4/1^48^6$ (16+16 <sup>T</sup> )                         |
| 8D | $-2A^\circ(2/1)/2$ (16-)   |  |
| 8E | 16B (16-),   | $1^24^216^2/2^58$ (16-)  |
| 8F | 4A/4 (64+ $(-\frac{1}{4}16)$ )                                       |  |
| 9A | -9A (18+18),   | $2^69^336^3/1^34^318^6$ ((18+18) <sup>T</sup> )                |
| 9B | -9B (18-),   | $6^39^336^3/3.12.18^6$ (36+4)                                  |
| 9C | -9C (18+9),  | $1^34^36^49^336^3/2^63^212^218^6$ (36+)                        |

Table II (continued)

|     |   |   |
|-----|---|---|
| 10A | 20A (20+20),  | $2^6 5^2 20^4 / 1^2 4^4 10^6$ (20+20 <sup>T</sup> )   |
| 10B | $-5A^\circ (2/1)/2$ ((40+40, $(\frac{1}{2}20)$ ) <sup>ST</sup> 20)  |   |
| 10C | $-5B^\circ (2/1)/2$ (80+5, 16, $(\frac{1}{2}40)$ )  |   |
| 10D | $2 \cdot 4 \cdot 5^2 20^3 / 1^2 10^5$ (20+20 <sup>TW</sup> 4),  | $1^2 4^3 10 \cdot 20 / 2^5 5^2$ (20+20 <sup>TW</sup> 4)   |
| 10E | $1^3 4 \cdot 5 \cdot 20^3 / 2^4 10^4$ (20+20),  | $2^5 20^2 / 1^3 4^2 5 \cdot 10$ (20+20 <sup>T</sup> )   |
| 10F | 10C (40+5, $(\frac{1}{2}20)$ )  |   |
| 11A | -11A (22+11),   | $1^2 4^2 11^2 44^2 / 2^4 22^4$ (44+)  |
| 12A | $2^4 3^4 4^4 24^4 / 1^4 6^4 8^4 12^4$ (24+8, $W_3(\frac{1}{2}12)$ ),  | $1^4 4^8 6^8 24^4 / 2^8 3^4 8^4 12^8$ (24+24, $W_3(\frac{1}{2}12)$ )                              |
| 12B | $-6A^\circ (2/1)/2$ (24+W <sub>3</sub> <sup>T</sup> 12)   |   |
| 12C | 24A (48+48, $(\frac{1}{2}24)$ )   |   |
| 12D | $2 \cdot 3^3 4 \cdot 24^3 / 1 \cdot 6^3 8 \cdot 12^3$ (24+8),   | $1 \cdot 4^2 6^6 24^3 / 2^2 3^3 8 \cdot 12^6$ (24+8 <sup>T</sup> )                                |
| 12E | $1^2 3^2 8^2 24^2 / 2^2 4^2 6^2 12^2$ (24+),  | $2^4 6^4 8^2 24^2 / 1^2 3^2 4^4 12^4$ (24+3 <sup>T</sup> , 8 <sup>T</sup> )                       |
| 12F | $-3A^\circ (2/1)/4$ (96+96, $(\frac{1}{4}24)$ , $(\frac{1}{2}48)$ )   |   |
| 12G | $-6E^\circ (2/1)/2$ (24+W <sub>3</sub> <sup>T</sup> )   |   |
| 12H | $1 \cdot 3 \cdot 4^5 24^2 / 2^4 6^2 8^2 12$ (24+W <sub>3</sub> <sup>T</sup> 12),  | $4^4 6 \cdot 24^2 / 1 \cdot 2 \cdot 3 \cdot 8^2 12^2$ (24+W <sub>3</sub> <sup>T</sup> 12)         |
| 12I | $2^2 8 \cdot 3^2 12 \cdot 24 / 1^2 4 \cdot 6^4$ (24+W <sub>3</sub> <sup>T</sup> ),  | $1^2 4 \cdot 8 \cdot 6^2 24 / 2^4 3^2 12$ (24+W <sub>3</sub> <sup>T</sup> )                       |
| 12J | 24C (48+3, $(\frac{1}{2}24)$ )  |   |
| 12K | $1^3 4^3 6^3 24^3 / 2^5 3 \cdot 8 \cdot 12^5$ (24+24),  | $2^4 3 \cdot 24^3 / 1^3 8 \cdot 12^4$ (24+24 <sup>T</sup> )                                       |
| 12L | $-1A^\circ (2/1)/12$ (576+ $(\frac{1}{6}96)$ , $(\frac{1}{3}192)$ , $(\frac{1}{4}144)$ , 576; $Z_2 \times Z_2 \times D_8$ ) |   |
| 12M | 12L (288+ $(-\frac{1}{12}24)$ , $(-\frac{1}{6}48)$ , $(-\frac{1}{4}72)$ , $(-\frac{1}{2}144)$ )                             |   |
| 13A | -13A (26+26),   | $2^4 13^2 52^2 / 1^2 4^2 26^4$ ((26+26) <sup>T</sup> )  |
| 14A | $-7A^\circ (2/1)/2$ ((56+56, $(\frac{1}{2}2)$ ) <sup>ST</sup> 28)   |   |
| 14B | 28A (28+7),   | $1 \cdot 4^2 7 \cdot 28^2 / 2^3 14^3$ (28+7)  |
| 15A | -15A (30+6, 10),  | $1^3 4^3 6^6 10^6 15^3 60^3 / 2^6 3^3 5^3 12^3 20^3 30^6$ ((30+6, 10) <sup>T</sup> )              |
| 15B | -15B (30+5, 6),   | $2^4 3^2 10^4 12^2 15^2 60^2 / 1^2 4^2 5^2 6^4 20^2 30^4$ ((30+5, 6) <sup>T</sup> )               |
| 15C | -15C (90+9, 10, $(-\frac{1}{3}30)$ ),   | $-5A^\circ (2/1)/3$ ((90+9, 10, $(-\frac{1}{3}30)$ ) <sup>T</sup> )                               |
| 15D | -15D (30+3, 5),   | $1 \cdot 3 \cdot 4 \cdot 5 \cdot 12 \cdot 15 \cdot 20 \cdot 60 / 2^2 6^2 10^2 30^2$ (60+)         |
| 15E | -15E (30+15),   | $1^2 4^2 6^2 10^2 15^2 60^2 / 2^4 3 \cdot 5 \cdot 12 \cdot 20 \cdot 30^4$ ((30+15) <sup>T</sup> ) |
| 16A | $-8C^\circ (2/1)/2$ (64+ $(\frac{1}{2}32)$ , $64^T 16$ )  |   |
| 16B | $1^2 4 \cdot 8 \cdot 32^2 / 2^3 16^3$ (32+),  | $2^3 8 \cdot 32^2 / 1^2 4 \cdot 16^3$ (32+32 <sup>T</sup> )                                       |
| 18A | 36A (36+36),  | $2^3 9 \cdot 36^2 / 1 \cdot 4^2 18^3$ (36+36 <sup>T</sup> )                                       |
| 18B | $1^2 4 \cdot 6^2 9 \cdot 36^2 / 2^3 3 \cdot 12 \cdot 18^3$ (36+36),   | $2^3 3 \cdot 36 / 1^2 4 \cdot 6 \cdot 9$ (36+36 <sup>T</sup> )                                    |
| 18C | $4 \cdot 6 \cdot 9 \cdot 36^2 / 1 \cdot 12 \cdot 18^3$ (36+36 <sup>W</sup> 4 <sup>T</sup> ),                                | $1 \cdot 4^2 6 \cdot 36 / 2^3 9 \cdot 12$ (36+36 <sup>W</sup> 4 <sup>T</sup> )                    |
| 20A | $2^2 4^2 5^2 40^2 / 1^2 8^2 10^2 20^2$ (40+40, $W_5(\frac{1}{2}20)$ ),  | $1^2 4^4 10^4 40^2 / 2^4 5^2 8^2 20^4$ (40+8, $W_5(\frac{1}{2}20)$ )                              |
| 20B | $-5B/4$ (160+5, $(-\frac{1}{4}40)$ , $(\frac{1}{2}80)$ )  |   |
| 20C | $4^2 5 \cdot 40 / 1 \cdot 8 \cdot 10^2$ (40+W <sub>5</sub> $(\frac{1}{2}20)$ ),   | $1 \cdot 4^3 10 \cdot 40 / 2^3 5 \cdot 8 \cdot 20$ (40+W <sub>5</sub> $(\frac{1}{2}20)$ )         |

Table II (continued)

|      |   |  |
|------|---|--|
| 21A  | -21A (42+6,14),   | $1^2 4^2 6^4 14^4 21^2 84^2 / 2^4 3^2 7^2 12^2 28^2 42^4$ ((42+6,14) <sup>T</sup> )  |
| 21B  | -21B (42+3,14),   | $2^2 6^2 7 \cdot 21 \cdot 28 \cdot 84 / 1 \cdot 3 \cdot 4 \cdot 12 \cdot 14^2 42^2$ ((42+3,14) <sup>T</sup> )  |
| 21C  | -21C (126+7,9,(- $\frac{1}{3}$ 42)),  | $3 \cdot 12 \cdot 21 \cdot 84 / 6^2 42^2$ (252+4,9,7,( $\frac{1}{3}$ 84))  |
| 22A  | -11A/2 (88+11,( $\frac{1}{2}$ 44))  |  |
| 23AB | -23A (46+23),   | $1 \cdot 4 \cdot 23 \cdot 92 / 2^2 46^2$ (92+)   |
| 24A  |   | $12A^\circ (2/1)/2$ (96+32, $96^T 24$ , ( $\frac{1}{3}$ 48))   |
| 24B  | $2^3 3^2 8^3 12^2 48^2 / 1^2 4^2 6^3 16^2 24^3$ (48+48,16 <sup>T</sup> ),   | $1^2 6^3 8^3 48^2 / 2^3 3^2 16^2 24^3$ (48+16,48 <sup>T</sup> )  |
| 24C  |   | $12E^\circ (2/1)/2$ (96+3,32, ( $\frac{1}{2}$ 48))   |
| 24D  |   | $12A/4$ (192+192, ( $\frac{1}{4}$ 48); D <sub>8</sub> )  |
| 24E  |   | $-12E^\circ (2/1)/2$ (96+3, $32^T 24$ , ( $\frac{1}{2}$ 48))   |
| 24F  | $2 \cdot 3 \cdot 8^2 12 \cdot 48 / 1 \cdot 4 \cdot 6^2 16 \cdot 24$ (48+W <sub>3</sub> ( $\frac{1}{2}$ 24)),  | $1 \cdot 6 \cdot 8^2 48 / 2^2 3 \cdot 16 \cdot 24$ (48+W <sub>3</sub> ( $\frac{1}{2}$ 24))   |
| 26A  |   | $-13A^\circ (2/1)/2$ ((104+104, ( $\frac{1}{2}$ 52)) <sup>ST</sup> 52)   |
| 28A  | $1 \cdot 7 \cdot 8 \cdot 56 / 2 \cdot 4 \cdot 14 \cdot 28$ (56+),   | $2^2 8 \cdot 14^2 56 / 1 \cdot 4^2 7 \cdot 28^2$ (56+7 <sup>T</sup> ,8 <sup>T</sup> )  |
| 28B  |   | $-7A^\circ (2/1)/4$ ((224+224, ( $\frac{1}{4}$ 56), ( $\frac{1}{2}$ 112)) <sup>ST</sup> 112)   |
| 30A  | 60A (60+12,15),   | $1 \cdot 4^4 6^3 10^3 15 \cdot 60^2 / 2^3 3 \cdot 5 \cdot 12^2 20^2 30^3$ (60+12 <sup>T</sup> ,15 <sup>T</sup> )   |
| 30B  |   | $-15B^\circ (2/1)/2$ ((120+5,24, ( $\frac{1}{2}$ 60)) <sup>ST</sup> 60)  |
| 30C  |   | $-5A^\circ (2/1)/6$ ((360+360, ( $\frac{1}{6}$ 60), ( $\frac{1}{3}$ 120), ( $\frac{1}{2}$ 180)) <sup>ST</sup> 60)  |
| 30D  | $3 \cdot 4 \cdot 5 \cdot 60 / 2 \cdot 6 \cdot 10 \cdot 30$ (60+12,15),  | $4 \cdot 6^2 10^2 60 / 2 \cdot 3 \cdot 5 \cdot 12 \cdot 20 \cdot 30$ (60+12 <sup>T</sup> ,15 <sup>T</sup> )  |
| 30E  | $2 \cdot 3 \cdot 5 \cdot 12 \cdot 20 \cdot 30 / 1 \cdot 6^2 10^2 15$ (60+15, TW <sub>5</sub> ),   | $1 \cdot 4 \cdot 6 \cdot 10 \cdot 15 \cdot 60 / 2^2 3 \cdot 5 \cdot 30^2$ (60+15, TW <sub>5</sub> )  |
| 33A  | -33A (66+6,11),   | $2^2 3 \cdot 12 \cdot 22^2 33 \cdot 136 / 1 \cdot 4 \cdot 6^2 11 \cdot 44 \cdot 66^2$ ((66+6,11) <sup>T</sup> )  |
| 35A  | -35A (70+10,14),  | $1 \cdot 4 \cdot 10^2 14^2 35 \cdot 140 / 2^2 5 \cdot 7 \cdot 20 \cdot 28 \cdot 70^2$ ((70+10,14) <sup>T</sup> )   |
| 36A  | $2 \cdot 4 \cdot 9 \cdot 72 / 1 \cdot 8 \cdot 18 \cdot 36$ (72+72, W <sub>9</sub> ( $\frac{1}{2}$ 36)),   | $1 \cdot 4^2 18^2 72 / 2^2 8 \cdot 9 \cdot 36^2$ (72+8, W <sub>9</sub> ( $\frac{1}{2}$ 36))  |
| 39AB | -39A (78+6,26),   | $1 \cdot 4 \cdot 6^2 26^2 39 \cdot 156 / 2^2 3 \cdot 12 \cdot 13 \cdot 52 \cdot 78^2$ ((78+6,26) <sup>T</sup> )  |
| 40A  |   | $20A^\circ (2/1)/2$ (160+32, $160^T 40$ , ( $\frac{1}{2}$ 40))   |
| 42A  |   | $-21A^\circ (2/1)/2$ ((168+21,56, ( $\frac{1}{2}$ 84)) <sup>ST</sup> 84)   |
| 60A  | $1 \cdot 6 \cdot 8 \cdot 10 \cdot 15 \cdot 20 \cdot 48 \cdot 120 / 2 \cdot 3 \cdot 4 \cdot 5 \cdot 24 \cdot 30 \cdot 40 \cdot 60$ (120+15,120, W <sub>3</sub> ( $\frac{1}{2}$ 60)), | $2^2 3 \cdot 5 \cdot 8 \cdot 12 \cdot 20^2 30^2 48 \cdot 120 / 1 \cdot 4^2 6^2 10^2 15 \cdot 24 \cdot 40 \cdot 60^2$ (120+15,24, W <sub>3</sub> ( $\frac{1}{2}$ 60)) |

Table III. (3/1)-transformations of Frame shapes of ·0.

|    |                          |  |
|----|--------------------------|--|
| 1A | 3A (3-),                 | -3A (6+6)  |
| 2A | 6A (6+2),                | -6A (6-)   |
| 2B | 6B (12/2+6)              |  |
| 2C | 3A/2 (6/2-)              |  |
| 3A | $1^6 9^6 / 3^{12}$ (9+), | $2^6 3^{12} 18^6 / 1^6 6^{12} 9^6$ (18+W <sub>2</sub> (6(- $\frac{1}{3}$ ))) |
| 3B | 9A (9-),                 | -9A (18+18)  |
| 3D | 3A/3 (9/3-),             | -3A/3 (18/3+6)   |



Table III (continued)

|     |  |  |
|-----|--|--|
| 4A  | 12A (12+12),   | -12A (12+4)  |
| 4B  | 12B (12-)  |  |
| 4C  | $3^2 6 \cdot 12^2 / 1^2 2 \cdot 4^2$ (?),                                | $1^2 6^3 12^2 / 2^3 3^2 4^2$ (?)   |
| 4D  | 12C (12/2+2)   |  |
| 4E  | 12F (24/4+6)   |  |
| 4F  | $12^3 / 4^3$ (?)   |  |
| 5A  | 15A (15+15),   | -15A (30+6,10)   |
| 5B  | 15B (15+5),  | -15B (30+5,6)  |
| 6A  | $1^2 2^2 9^2 18^2 / 3^4 6^4$ (18+),                                      | $2^4 3^4 18^4 / 1^2 6^8 9^2$ (18+9)  |
| 6B  | $-3A \circ (3/1) / 2$  | $(72 + (\frac{1}{2}36), W_{72}(-\frac{1}{3}24); D_8)$  |
| 6E  | 18A (18+2),  | -18A (18-)   |
| 6H  | $-3A/6$  | (36/6+6)   |
| 6I  | $3A/6$   | (18/6-)  |
| 7A  | 21A (21+21),   | -21A (42+6,14)   |
| 8A  | 24A (24/2+12)  |  |
| 8B  | $-12A/2$   | (24/2+4)   |
| 8C  | 24B (24+24),   | -24B (24+8)  |
| 8D  | $4 \cdot 24^2 / 8^2 12$ (?)  |  |
| 8F  | 24D (24/4+2)   |  |
| 10A | 30A (30+2,15),   | -30A (30+15)   |
| 10C | 30B (60/2+5,6)   |  |
| 10F | 15B (30/2+5)   |  |
| 11A | 33A (33+11),   | -33A (66+6,11)   |
| 12A | $3^4 4^2 36^2 / 1^2 9^2 12^4$ (36+9, $W_{36}(12(\frac{1}{3}))$ ),        | $1^2 4^2 6^4 9^2 36^2 / 2^2 3^4 12^4 18^2$ (36+)   |
| 12B | $(-6A) \circ (2/1) / 2$  | (36/2+9)   |
| 12C | $6A \circ (2/1) / 2$   | (36/2+)  |
| 12E | 36A (36+36),   | -36A (36+4)  |
| 12L | $-3A/12$   | (72/12+6)  |
| 12M | $36/12$  | (?)  |
| 13A | 39AB (39+39),  | -39AB (78+6,26)  |
| 14A | 42A (84/2+6,14)  |  |
| 15B | $1 \cdot 5 \cdot 9 \cdot 45 / 3^2 15^2$ (45+),                           | $2 \cdot 3^2 10 \cdot 15^2 18 \cdot 90 / 1 \cdot 5 \cdot 6^2 9 \cdot 30^2 45$ (90+5, $W_2(30(\frac{1}{3})); D_8$ )   |
| 15C | $15A/3$ (45/3+15),   | -15A/3 (90/3+6,10)   |
| 20A | 60A (60+12,15),  | -60A (60+4,15)   |
| 21A | $3^2 7 \cdot 63 / 1 \cdot 9 \cdot 21^2$ (63+9, $W_7(21(\frac{1}{3}))$ ), | $1 \cdot 6^2 9 \cdot 14 \cdot 21^2 126 / 2 \cdot 3^2 7 \cdot 18 \cdot 42^2 63$ (126+9, 126, $W_7(42(\frac{1}{3}))$ ) |
| 24A | $12A \circ (3/1) / 2$  | $(144+9, W_{16}(48(\frac{1}{3})), (\frac{1}{2}36); Z_2 \times Z_2 \times Z_2)$                                       |

Table IV. Other transformations of Frame shapes of  $\cdot 0$ .

| 2-transf. |           | (4/1)-transf.                                | 5-transf.    | (7/1)-transf.  |
|-----------|-----------|--|--------------|----------------|
| 1A        | 2A, 4A    | 4A, $4A^\circ(2/1)$                          | 1A 5B, -5B   | 7A, -7A        |
| 2B        | 8A        | $4A^\circ(2/1)/2$                            | 2B 10C       | 14A            |
| 2C        | 4D        | 8A   | 2C 10F       | 7A/2           |
| 3A        | 6A, 12A   | 12A, $12A^\circ(2/1)$                        | 3A 15B, -15B | 21A, -21A      |
| 3B        | 6E, 12E   | 12E, $12E^\circ(2/1)$                        | 3B 15D, -15D | 21B, -21B      |
| 3C        | 6F, 12D   | 12D, $12D^\circ(2/1)$                        | 4E -5B/4     | 28B            |
| 4E        | 4A/4      | $4A^\circ(2/1)$                              | 4F 20B       | 28/4 (?)       |
| 4F        | 8F        | 4A/4   | 5A 25Z, -25Z | 35A, -35A      |
| 5A        | 10A, 20A  | 20A, $20A^\circ(2/1)$                        | 6B 30B       | 42A            |
| 6B        | 24A       | $12A^\circ(2/1)/2$                           |              |                |
| 6G        | 12J       | 24C  | 7-transf.    |                |
| 7B        | 14B, 28A  | 28A, $28A^\circ(2/1)$                        | 1A 7B, -7B   | (9/1)-transf.  |
| 9A        | 18A, 36A  | 36A, $36A^\circ(2/1)$                        | 2A 14B, -14B | 9A, -9A        |
| 10B       | 40A       | $20A^\circ(2/1)/2$                           | 3D 21C, -21C | 18A, -18A      |
| 12F       | 12A/4     | $12A^\circ(2/1)/4$                           | 4A 28A, -28A | 9A/3, -9A/3    |
| 15A       | 30A, 60A  | 60A, $60A^\circ(2/1)$                        |              | 36A, -36A      |
|           | 3-transf. |  | 11-transf.   |                |
| 1A        | 3B, -3B   | (5/1)-transf.                                |              | (13/1)-transf. |
| 2A        | 6E, -6E   | 5A, -5A                                      | 1A 11A, -11A | 13A, -13A      |
| 2B        | -3B/2     | 10A, -10A                                    | 2B -11A/2    | 26A            |
| 2C        | 6G        | 10B  | 2C 22A       | 26/2 (?)       |
| 3A        | 9A, -9A   | $10^3/2^3$ (?)                               | 3A 33A, -33A | 39A, -39A      |
| 3D        | 9Z, -9Z   | 15A, -15A                                    | 23-transf.   |                |
| 4A        | 12E, -12E | 15C, -15C                                    | 1A 23A, -23A | (25/1)-transf. |
| 4B        | 12G       | 20A, -20A                                    |              | 25Z, -25Z      |
| 4D        | 12J       | $2 \cdot 20^2/4^2 10$ (?)                    |              |                |
| 5B        | 15D, -15D | 10A/2  |              |                |
| 6A        | 18A, -18A | 25Z, -25Z                                    |              |                |
| 6H        | -3B/6     | 30A, -30A                                    |              |                |
| 6I        | 18Z       | -5A/6  |              |                |
| 7A        | 21B, -21B | 30/6 (?)                                     |              |                |
| 8A        | 24C       | 35A, -35A                                    |              |                |
| 8B        | 24E       | 20A/2  |              |                |
| 12A       | 36A, -36A | $4 \cdot 10 \cdot 40/2 \cdot 8 \cdot 20$ (?) |              |                |
|           |           | 60A, -60A                                    |              |                |

Takeshi KONDO

Department of Mathematics  
 College of Arts and Sciences  
 University of Tokyo  
 Komaba, Meguro-ku  
 Tokyo 153, Japan