

**On the differentiability and the representation of one-parameter semi-group of linear operators.**

By Kôzaku YOSIDA.

(Received Oct. 25, 1947.)

1. *The theorems.* Let  $\{U_t\}$ ,  $0 \leqq t < \infty$ , be a one-parameter semi-group of linear (=additive, continuous) operators from a complex Banach space  $E$  to  $E$ :

$$U_t U_s = U_{t+s}, U_0 = I \quad (= \text{the identity operator}), \tag{1.1}$$

such that

$$\sup_t \|U_t\| \leqq 1, \tag{1.2}$$

$$\lim_{t \rightarrow t_0} U_t x = U_{t_0} x \quad (\text{lim} = \text{strong limit}), 0 \leqq t_0 < \infty, x \in E. \tag{1.3}$$

The purpose of the present note is to prove the following two theorems<sup>1)</sup>.

*Theorem 1.* If we denote by  $D$  the totality of  $x$  for which

$$\text{weak } \lim_{h \downarrow 0} h^{-1}(U_h - I)x = Ax \tag{1.4}$$

exists, then  $D$  is dense in  $E$ . Moreover  $A$  is a closed additive operator from  $D$  to  $E$  with the properties:

$$\text{for any } x \in D, \lim_{h \rightarrow 0} h^{-1}(U_{t+h} - U_t)x = AU_t x = U_t Ax, \tag{1.5}$$

there exists a sequence  $\{I_n\}$  of linear operators each commutative with every  $U_t$  and  $A$  such that i) the range  $R(I_n) = \{I_n x; x \in E\} \subseteq D$ ,  $AI_n = n(I_n - I)$ , ii)  $\|I_n\| \leqq 1$ ,  $\lim_{n \rightarrow \infty} I_n x = x$ , iii)  $U_t x = \lim_{n \rightarrow \infty} \exp(tAI_n)x = \lim_{n \rightarrow \infty} \sum_{m=0}^{\infty} (m!)^{-1} (tAI_n)^m x$  uniformly for  $t$  in any finite interval<sup>2)</sup>, (1.6)

$$\|(A - nI)x\| \geqq n \|x\| \quad (n=1, 2, \dots) \text{ for } x \in D \text{ and the range } R(A - nI) \text{ coincides with } E \quad (n=1, 2, \dots), \tag{1.7}$$

let, by (1.7),  $y_n$  be the unique solution of  $(A - nI)y_n = y$  ( $n=1, 2, \dots$ ), then  $\lim_{n \rightarrow \infty} A(-ny_n) = \lim_{n \rightarrow \infty} (-n(y + ny_n)) = Ay$  for  $y \in D$ . (1.8)

*Theorem 2.* Let, conversely,  $A$  be an additive operator from a dense

linear subset  $D$  of  $E$  to  $E$  such that (1.7) and (1.8) are satisfied. Then there exists a one-parameter semi-group  $U_t$  which satisfies (1.1)—(1.3) and (1.5).

We have, thus, a characterisation of the differential quotient  $A$  of the one-parameter semi-group of linear operators. This may be applied to an operator-theoretical treatment of temporally homogeneous stochastic process<sup>3)</sup>. As an application of the theorem 1, we give a new proof of Stone's theorem<sup>4)</sup> (see 4 below).

2. *Proof of the theorem 1. The differentiability.* We may define the integral

$$C_\varphi \cdot x = \int_0^\infty \varphi(s) U_s x ds \quad (2.1)$$

for complex-valued continuous function  $\varphi(s)$  following after S. Bochner, G. Birkhoff, I. Gelfand, B. J. Pettis and other authors<sup>5)</sup>. Let  $\varphi(s)$  satisfy

$$\lim_{h \rightarrow 0} \int_h^\infty \left| \frac{\varphi(s-h) - \varphi(s)}{h} + \varphi'(s) \right| ds = 0. \quad (2.2)$$

We have, by (1.1)

$$\begin{aligned} \frac{1}{h} (U_h - I) C_\varphi x &= \frac{1}{h} \int_0^\infty \varphi(s) U_h U_s x ds - \frac{1}{h} \int_0^\infty \varphi(s) U_s x ds \\ &= \int_h^\infty \frac{\varphi(s-h) - \varphi(s)}{h} U_s x ds - \frac{1}{h} \int_0^h \varphi(s) U_s x ds. \end{aligned}$$

Thus, by (2.2) and  $U_0 = I$ ,

$$\lim_{h \rightarrow 0} \frac{1}{h} (U_h - I) C_\varphi x \text{ exists and } = C_{-\varphi'} \cdot x - \varphi(0)x. \quad (2.3)$$

Such  $C_\varphi x$  is dense in  $E$ , since for any  $\delta > 0$

$$\varphi_\delta(s) = \delta \exp(-\delta s) \quad (2.4)$$

satisfies (2.2) and moreover, for any  $x \in E$ ,  $\lim_{\delta \rightarrow \infty} C_{\varphi_\delta} \cdot x = x$ .

We have, by (1.4), for  $x \in D$ ,

$$\begin{aligned} \text{weak } \lim_{h \rightarrow 0} \frac{1}{h} (U_h - I) U_t x &= \text{weak } \lim_{h \rightarrow 0} \frac{1}{h} (U_{t+h} - U_t) x \\ &= U_t \cdot \text{weak } \lim_{h \rightarrow 0} \frac{1}{h} (U_h - I) x. \end{aligned}$$

Hence  $U_t \cdot D \subseteq D$  and  $AU_t x = U_t Ax$  for any  $x \in D$ , viz.  $A$  is commutative with every  $U_t$ , and

$$\begin{cases} \text{the right weak derivative } D^+ U_t x \text{ exists and} \\ = AU_t x = U_t Ax \text{ for any } x \in D. \end{cases} \quad (2.5)$$

Hence, by the continuity (1.3) we have, for any  $f \in E^*$  (=the conjugate space of  $E$ )

$$f(U_t x) - f(x) = \int_0^t D^+ f(U_s x) ds = \int_0^t f(U_s Ax) ds = f\left(\int_0^t U_s Ax ds\right)$$

and therefore

$$U_t x - x = \int_0^t U_s Ax ds \quad \text{for } x \in D. \quad (2.6)$$

Thus we have the strong differentiability (1.5).

*The closedness of  $A$ .* Let  $x_n \in D$  ( $n=1, 2, \dots$ ) and let  $\lim_{n \rightarrow \infty} x_n = x$ ,  $\lim_{n \rightarrow \infty} Ax_n = z$ . Then, by (2.6),

$$U_t x - x = \int_0^t U_s z ds.$$

*The representation (1.6).* Put

$$I_n = C\varphi_n \quad (n=1, 2, \dots), \quad (2.7)$$

then, by (2.3) and (2.4), i) and ii) are surely satisfied. In fact, we have by (1.2) and (2.4)

$$\|I_n\| \leq \int_0^\infty n \exp(-ns) ds = 1. \quad (2.8)$$

Thus, since

$$AI_n = n(I_n - I) \quad (2.9)$$

by (2.3), we have

$$\exp(tAI_n) = \sum_{m=0}^{\infty} \frac{(tAI_n)^m}{m!} = \exp(tn(I_n - I)), \quad 0 \leq t < \infty. \quad (2.10)$$

Thus

$$\|\exp(tAI_n)\| = \|\exp(tnI_n) \exp(-tnI)\| \leq \exp(tn) \exp(-tn) = 1. \quad (2.11)$$

Since  $AI_n$  is commutative with each  $U_t$ , we have, for  $x \in D$ ,

$$\begin{aligned}
\|U_t x - \exp(tAI_n)x\| &= \left\| \int_0^t \frac{d}{ds} \left( (\exp(t-s)AI_n)U_s x \right) ds \right\| \\
&= \left\| \int_0^t (\exp(t-s)AI_n) U_s (A-AI_n)x ds \right\| \\
&\leq \int_0^t \| (A-AI_n)x \| ds \quad \text{by (1.2) and (2.11)} \\
&= t \| (A-AI_n)x \|. \quad (2.12)
\end{aligned}$$

We have

$$AI_n = I_n Ax \quad \text{for } x \in D, \quad (2.13)$$

for  $A$  is a closed operator commutative with each  $U_t$ . Thus, by (2.12),

$$\|U_t x - \exp(tAI_n)x\| \leq t \|(I - I_n)Ax\| \quad \text{for } x \in D. \quad (2.14)$$

Since  $U_t$  and  $\exp(tAI_n)$  are both of norm  $\leq 1$ , we have iii) by the fact that  $D$  is dense in  $E$ .

*The proof of (1.7).* We first show that  $R(A-nI)$  is dense in  $E$ . If otherwise, there exists  $f \in E^*$ ,  $f \neq 0$ , such that  $f(Ax-nx) = 0$  on  $D$ . Thus, by  $U_t D \subseteq D$  we have  $f(AU_t x) = nf(U_t x)$  and hence

$$\frac{d}{dt} f(U_t x) = nf(U_t x).$$

Therefore, by  $f(U_0 x) = f(x)$ , we obtain  $f(U_t x) = f(x) \exp(nt)$ . This is a contradiction. Proof. By  $f \neq 0$  and by the fact  $D$  is dense in  $E$  there exists  $x \in D$  such that  $f(x) \neq 0$ . Then  $f(x) \exp(nt)$  is unbounded in  $t$  when  $t \rightarrow \infty$ , contrary to  $|f(U_t x)| \leq \|f\| \cdot \|U_t\| \cdot \|x\| \leq \|f\| \cdot \|x\|$ . Next we show that

$$\|(A-nI)x\| \geq n\|x\| \quad \text{for } x \in D.$$

Assume the contrary and let  $\|(A-nI)x\| = a < n$  for a certain  $x \in D$  with  $\|x\| = 1$ . Let  $f \in E^*$  be such that  $f(x) = 1$ ,  $\|f\| = 1$ . Then, by

$$\frac{d}{dt} U_t x = U_t Ax = nU_t x + U_t(A-nI)x,$$

we obtain

$$\begin{cases} \frac{d}{dt} \varphi(t) = n\varphi(t) + \psi(t), & \text{where} \\ \varphi(t) = f(U_t x), & \psi(t) = f(U_t(A-nI)x). \end{cases}$$

Since  $\varphi(0)=1$  we have

$$\varphi(t) = \exp(nt) \left( \int_0^t \exp(-nt) \phi(t) dt + 1 \right)$$

and hence, by  $|\phi(t)| \leq \|f\| \cdot \|U_t\| \cdot \|(A-nI)x\| \leq a$ ,

$$|\varphi(t)| \geq \exp(nt) (1 - an^{-1}(1 - \exp(-nt))).$$

Thus  $\varphi(t)$  is unbounded in  $t$  when  $t \rightarrow \infty$ , contrary to  $|\varphi(t)| \leq \|f\| \cdot \|U_t\| \cdot \|x\| \leq 1$ . Therefore, for any  $y \in E$ , there exists a sequence  $\{x_h\} \subseteq D$  such that  $\lim_{h \rightarrow \infty} (A-nI)x_h = y$ . Because of  $\|(A-nI)(x_h - x_k)\| \geq n \|x_h - x_k\|$ ,  $\{x_h\}$  is a Cauchy sequence. Let  $\lim_{h \rightarrow \infty} x_h = x$ , then  $\lim_{h \rightarrow \infty} (A-nI)x_h = y$  and by the closedness of  $A$  we have  $y = (A-nI)x$ .

The proof of (1.8). We have, by (2.9),  $AI_n y - nI_n y = -ny$  and hence

$$-ny_n = I_n y. \quad (2.15)$$

Thus (1.6) and (2.13) imply (1.8).

3. Proof of the theorem 2. By (1.7), the operator  $J_n$  defined by

$$J_n y = -ny_n \quad (n=1, 2, \dots) \quad (3.1)$$

satisfies

$$\|J_n\| \leq 1. \quad (3.2)$$

Since

$$AJ_n y = A(-ny_n) = -nA y_n = -n(y + ny_n) = n(J_n - I)y, \quad (3.3)$$

we have, by (3.2),

$$\|\text{ext}(tAJ_n)y\| = \|\exp(ntJ_n) \text{ext}(-ntI)y\| \leq \exp(nt) \exp(-nt) \cdot \|y\|,$$

hence the linear operator defined by

$$U_t^{(n)} = \exp(tAJ_n) \quad (3.4)$$

satisfies

$$\|U_t^{(n)}\| \leq 1, \quad (3.5)$$

$$U_t^{(n)} x - x = \int_0^t U_s^{(n)} AJ_n x ds, \quad (3.6)$$

$$\lim_{h \rightarrow 0} h^{-1} (U_{t+h}^{(n)} - U_t^{(n)}) x = U_t^{(n)} AJ_n x. \quad (3.7)$$

We have

$$J_n J_m = J_m J_n, \quad (3.8)$$

for, by (3.1),

$$J_n = -n^{-1}(A - nI)^{-1}. \quad (3.9)$$

Thus  $AJ_n$  is commutative with  $U_t^{(m)}$  and hence

$$\begin{aligned} \|(U_t^{(m)} - U_t^{(n)})x\| &= \left\| \int_0^t \frac{d}{ds} \left( (\exp(t-s)AJ_n)U_s^{(m)}x \right) ds \right\| \\ &= \left\| \int_0^t (\exp(t-s)AJ_n)U_s^{(m)}(AJ_m - AJ_n)x ds \right\| \\ &= \int_0^t \|(AJ_m - AJ_n)x\| ds \quad \text{by (3.5) and (3.7)} \\ &= t \|(AJ_m - AJ_n)x\|. \end{aligned}$$

Therefore, by (1.8),

$$U_t y = \lim_{n \rightarrow \infty} U_t^{(n)} y \quad (y \in D) \quad (3.10)$$

exists uniformly for  $t$  in any finite interval. Since  $D$  is dense in  $E$  and since we have (3.5), we see that the limit  $U_t^{(n)}$  exists for all  $y \in E$  and that  $U_t$  satisfies (1.1)—(1.3). Hence, by letting  $n \rightarrow \infty$  in (3.6), we obtain

$$U_t y - y = \int_0^t U_s A y ds, \quad y \in D. \quad (3.11)$$

4. *Stone's theorem.* If  $E$  is a Hilbert space and if  $U_t$  is, for any  $t \geq 0$ , a unitary operator, then  $H = -iA$  ( $i = \sqrt{-1}$ ) is self-adjoint and

$$U_t = \int_{-\infty}^{\infty} \exp(i\lambda t) dE(\lambda), \quad \text{where } H = \int_{-\infty}^{\infty} \lambda dE(\lambda). \quad (4.1)$$

This theorem due to M. H. Stone may be obtained as follows<sup>6)</sup>. Put  $U_{-t} = U_t^{-1} = U_t^*$  for  $t \geq 0$ , then  $\{U_t\}$ ,  $-\infty < t < \infty$ , is a one-parameter group of unitary operators strongly continuous in  $t$ . Thus it is easy to see, by (1.5),

$$\frac{dU_t x}{dt} = iHU_t = iU_t Hx \quad \text{for } x \in D, \quad -\infty < t < \infty. \quad (1.5)'$$

Hence, if  $x, y \in D$ ,

$$(Hx, y) = \frac{1}{i} \left( \frac{d}{dt} (U_t x, y) \right)_{t=0} = \frac{1}{i} \left( \frac{d}{dt} (x, U_{-t} y) \right)_{t=0} = (x, Hy), \quad (4.2)$$

which shows that  $H$  is a symmetric operator. We proved in 2 that  $R(H+iI) = E$  ((1.7)) by letting  $t \rightarrow \infty$ . Letting  $t \rightarrow -\infty$ , similar argument shows that  $R(H-iI) = E$  also. Therefore the Cayley transform of

$H$  is unitary and hence  $H$  is self-adjoint. Let  $H = \int_{-\infty}^{\infty} \lambda dE(\lambda)$  be its spectral resolution, then, as in (3.9),  $I_n = (I - n^{-1}A)^{-1}$  and hence

$$\begin{aligned} AI_n &= n(I_n - I) = n\left((I - in^{-1}H)^{-1} - I\right) = \int_{-\infty}^{\infty} n\left((1 - i\lambda n^{-1})^{-1} - 1\right) dE(\lambda) \\ &= \int_{-\infty}^{\infty} i\lambda(1 - i\lambda n^{-1})^{-1} dE(\lambda). \end{aligned}$$

Hence

$$\begin{aligned} U_t x &= \lim_{n \rightarrow \infty} \exp(tAI_n)x = \lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} \exp\left(it\lambda(1 - i\lambda n^{-1})^{-1}\right) dE(\lambda)x \\ &= \int_{-\infty}^{\infty} \exp(i\lambda t) dE(\lambda)x. \end{aligned}$$

Mathematical Institute,  
Nagoya University.

#### References.

- 1) Cf. I. Gelfand: C. R. URSS, **25** (1939), 713-718 and N. Dunford—I. E. Segal: *Bullet. Amer. Math. Soc.*, **52** (1946), 911-914.
- 2) According to N. Dunford and I. E. Segal's paper referred to in 1), E. Hille (*Proc. Nat. Acad. Sci.*, **28** (1942), 175-178, 421-424) obtained the representation  $U_t x = \lim_{n \rightarrow \infty} \exp(tn(U_n - I))x$ . Hille's paper is not available to the author.
- 3) The question of the characterization of  $A$  together with differentiability of  $U_t$  is proposed to the author by Dr. K. Itô in connection with his theory of stochastic differential equations. See his forthcoming paper in *Jap. J. of Math.*
- 4) *Proc. Nat. Acad. Sci.*, **16** (1930), 172-175. See also J. von Neumann: *Ann. of Math.*, **33** (1932), 567-573. Proofs of Stone's theorem are given by many authors: F. Riesz, B. von Sz. Nagy and H. Nakano.
- 5) S. Bochner: *Fund. Math.*, **20** (1933), 262-276. G. Birkhoff: *Trans. Amer. Math. Soc.* **38** (1935), 357-378. I. Gelfand: *Commun. Inst. Sci. Math. et Mech. Univ. Kharkoff*, **13** (1936), 35-40. B. J. Pettis: *Trans. Amer. Math. Soc.*, **44** (1938), 277-304.
- 6) In the case of Stone's theorem, we may replace (1.3) by the separability of  $\{U_t x; -\infty < t < \infty\}$  and the weak measurability of  $U_t$ . This may be carried out by virtue of N. Dunford's theorem in *Ann. of Math.*, **39** (1938), 567-573. This fact is, however, already proved by J. von Neumann in another way. See his paper referred to in 4).

*Added during the proof.* Meanwhile, E. Hille kindly communicated me that he also obtained essentially the same results as above by a different method. See his paper in *C. R.*, 8 September (1947) and *PNAS notes* referred to in 2).