# A NOTE ON THE PRODUCT OF MEROMORPHIC FUNCTIONS AND ITS DERIVATIVES

# KIT-WING YU

#### Abstract

It is shown that if f is an even or odd transcendental meromorphic function and if c is any even meromorphic function which does not vanish identically and satisfies T(r,c) = o(T(r,f)) as  $r \to +\infty$ , then ff' - c has infinitely many zeros.

#### 1. Introduction and our main results

In 1959, W. K. Hayman [5] proved that

**THEOREM A.** If *n* is an integer greater than or equal to 3 and *f* is a transcendental meromorphic function, then  $f^n f'$  takes every non-zero complex number infinitely many times.

Later, he conjectured [6] that this remains valid for the cases n = 1, 2. In 1979, E. Mues [7] proved the case n = 2 and the conjecture was proven by A. Eremenko and W. Bergweiler [2] in 1995 and independently by H. H. Chen and M. L. Fang [3].

In 1994, Yik-Man Chiang asked W. Bergweiler whether ff' - c has infinitely many zeros if f is a transcendental meromorphic function and if c is a meromorphic function which does not vanish identically and satisfies T(r, c) = o(T(r, f))as  $r \to +\infty$ . In [8], Q. D. Zhang studied the value distribution of  $\varphi(z)f(z)f'(z)$ and obtained the following theorem.

THEOREM B. If f is a transcendental meromorphic function and  $\varphi$  is a nonzero meromorphic function such that  $T(r, \varphi) = S(r, f)$  as  $r \to +\infty$ , then

$$T(r,f) < \frac{9}{2}\overline{N}(r,f) + \frac{9}{2}\overline{N}\left(r,\frac{1}{\varphi f f' - 1}\right) + S(r,f).$$

By this, we have

2000 Mathematics Subject Classfication: Prelimary 30D35. Key words: derivatives, meromorphic functions, zeros. Received July 21, 2000; revised January 5, 2001. COROLLARY A. If  $\delta(\infty, f) > 7/9$ , then ff' - c has infinitely many zeros.

In 1995, W. Bergweiler and A. Eremenko [2] used some results from iteration theory to show that ff' - c has infinitely many zeros if c is a non-zero constant. Later, W. Bergweiler [1] answered this question affirmatively in the case that f is of finite order and c is a polynomial. Actually, he showed that

THEOREM C. If f is a transcendental meromorphic function of finite order and c is a polynomial, then ff' - c has infinitely many zeros.

In the following discussion, we assume that f is a transcendental meromorphic function. From Corollary A, we can further assume that  $\delta(\infty, f) \le 7/9$  and we shall show the following result.

**THEOREM.** Let f be a transcendental meromorphic function and c be a meromorphic function which does not vanish identically and satisfies T(r,c) = o(T(r,f)) as  $r \to +\infty$ . Then ff' - c or ff' + c has infinitely many zeros.

COROLLARY. Suppose that f is an even or odd transcendental meromorphic function and c is an even meromorphic function. Suppose further that there exists a sequence  $\{z_1, -z_1, z_2, -z_2, \ldots,\}$  which are zeros of  $\varphi f f' + 1$  but not zeros or poles of f. Then ff' - c has infinitely many zeros.

Here, we assume that the readers are familiar with the basic concepts of the Nevanlinna value distribution theory and the notations m(r, f), N(r, f),  $\overline{N}(r, f)$ , T(r, f) and etc., see e.g., [4].

## 2. A lemma

LEMMA. Suppose that f is a non-constant meromorphic function and that  $\varphi$  is a non-vanishing meromorphic function such that  $T(r, \varphi) = o(T(r, f))$  as  $r \to +\infty$ . Then for any finite non-zero distinct complex numbers a and b and any positive integer k such that  $\varphi f^{(k)} \neq \text{constant}$ , we have

$$T(r,f) < N\left(r,\frac{1}{f}\right) + N\left(r,\frac{1}{\varphi f^{(k)} - a}\right) + N\left(r,\frac{1}{\varphi f^{(k)} - b}\right)$$
$$- N(r,f) - N\left(r,\frac{1}{(\varphi f^{(k)})'}\right) + S(r,f)$$

as  $r \to +\infty$ .

*Proof.* First of all, we have

(1) 
$$m\left(r,\frac{1}{\varphi f}\right) \le m\left(r,\frac{1}{\varphi f^{(k)}}\right) + m\left(r,\frac{f^{(k)}}{f}\right) + O(1).$$

From

$$\begin{split} m\!\left(r,\frac{1}{\varphi f}\right) &= T(r,\varphi f) - N\!\left(r,\frac{1}{\varphi f}\right) + O(1),\\ m\!\left(r,\frac{1}{\varphi f^{(k)}}\right) &= T(r,\varphi f^{(k)}) - N\!\left(r,\frac{1}{\varphi f^{(k)}}\right) + O(1), \end{split}$$

and (1), we have

(2) 
$$T(r,\varphi f) \le N\left(r,\frac{1}{\varphi f}\right) + T(r,\varphi f^{(k)}) - N\left(r,\frac{1}{\varphi f^{(k)}}\right) + m\left(r,\frac{f^{(k)}}{f}\right) + O(1).$$

By the second fundamental theorem,

(3) 
$$T(r,\varphi f^{(k)}) < N\left(r,\frac{1}{\varphi f^{(k)}}\right) + N\left(r,\frac{1}{\varphi f^{(k)}-a}\right) + N\left(r,\frac{1}{\varphi f^{(k)}-b}\right) - N_1(r) + S(r,\varphi f^{(k)})$$

as  $r \to +\infty$ , where, as usual,  $N_1(r)$  is defined as

$$N_1(r) = 2N(r, \varphi f^{(k)}) - N(r, (\varphi f^{(k)})') + N\left(r, \frac{1}{(\varphi f^{(k)})'}\right).$$

Let  $z_0$  be a pole of order  $p \ge 1$  of f. Then  $f^{(k)}$  and  $f^{(k+1)}$  have a pole of order k+p and k+p+1 at  $z_0$  respectively. Thus  $2(k+p) - (k+p+1) = k+p-1 \ge p$  and

(4) 
$$N_1(r) \ge N(r, f) + N\left(r, \frac{1}{(\varphi f^{(k)})'}\right) + S(r, f).$$

It is clear that  $S(r, f^{(k)}) = S(r, f)$  and  $m\left(r, \frac{f^{(k)}}{f}\right) = S(r, f)$ . Thus by (2), (3) and (4),

$$\begin{split} T(r,\varphi f) < N\bigg(r,\frac{1}{\varphi f}\bigg) + N\bigg(r,\frac{1}{\varphi f^{(k)}-a}\bigg) + N\bigg(r,\frac{1}{\varphi f^{(k)}-b}\bigg) \\ - N(r,f) - N\bigg(r,\frac{1}{(\varphi f^{(k)})'}\bigg) + S(r,f) \end{split}$$

as  $r \to +\infty$ . Since  $T(r, \varphi) = o(T(r, f))$  as  $r \to +\infty$ , we have the desired result.

# 3. Proofs of the theorem and the corollary

*Proof of the theorem.* Let  $\varphi = 1/c$ ,  $F = (1/2)f^2$ , k = 1, a = 1 and b = -1. Then by the above lemma, we have

$$\begin{split} 2T(r,f) &< 2N\left(r,\frac{1}{f}\right) + N\left(r,\frac{1}{\varphi f f' - 1}\right) + N\left(r,\frac{1}{\varphi f f' + 1}\right) \\ &- 2N(r,f) - N\left(r,\frac{1}{(\varphi f f')'}\right) + S(r,f) \end{split}$$

as  $r \to +\infty$ .

By the assumption that  $\delta(\infty, f) \leq 7/9$ , we have

$$\frac{N\!\left(r,\!\frac{1}{\varphi f f'-1}\right)+N\!\left(r,\!\frac{1}{\varphi f f'+1}\right)}{2T(r,f)}>0$$

as  $r \to +\infty$ . Hence, the result of the theorem follows.

*Proof of the corollary.* If

$$\frac{N\left(r,\frac{1}{\varphi f f'-1}\right)}{2T(r,f)} > 0$$

as  $r \to +\infty$  outside a set of finite linear measure, then  $\varphi f f' - 1$  has infinitely many zeros and thus f f' - c has infinitely many zeros.

Let  $z_0 \neq 0$  be a zero of  $\varphi ff' + 1$ . Since f is even or odd, f' is odd or even. Therefore, ff' is an odd function. Now we have  $\varphi(-z_0)f(-z_0)f'(-z_0) - 1 = -\varphi(z_0)f(z_0)f'(z_0) - 1 = 0$  and thus  $-z_0$  is a zero of  $\varphi ff' - 1$ . Hence, the desired result follows.

### 4. Remarks

1. In [8], Q. D. Zhang showed that

$$2T(r,f) < \overline{N}(r,f) + 2\overline{N}\left(r,\frac{1}{f}\right) + \overline{N}\left(r,\frac{1}{\varphi f f' - 1}\right) + S(r,f).$$

From this, it is immedicate that if  $\delta(0, f) + (1/2)\delta(\infty, f) > 1/2$ , then ff' - c has infinitely many zeros, where  $c = 1/\varphi$ .

2. In [9], Z. F. Zhang and G. D. Song showed that if  $a(z) \neq 0$  and T(r,a) = S(r, f) as  $r \to +\infty$ , n, k are positive integers, where  $n \ge 2$ , then  $f(f^{(k)})^n - a(z)$  has infinitely many zeros.

Acknowledgement. The author wishes to thank the referee for many valuable comments.

#### REFERENCES

- W. BERGWEILER, On the product of a meromorphic function and its derivative, Bull. Hong Kong Math. Soc., 1 (1996), 97–101.
- [2] W. BERGWEILER AND A. EREMENKO, On the singularities of the inverse of a meromorphic function of finite order, Rev. Mat. Iberoamericana, 11 (1995), 355–373.
- [3] H. H. CHEN AND M. L. FANG, On the value distribution of  $f^n f'$ , Science in China Ser. A, 25 (1995), 121–127.
- [4] W. K. HAYMAN, Meromorphic Functions, Oxford Math. Monogr., Clarendon Press, Oxford, 1964.
- [5] W. K. HAYMAN, Picard values of meromorphic functions and their derivatives, Ann. of Math., 70 (1959), 9-42.
- [6] W. K. HAYMAN, Research Problems in Function Theory, The Athlone Press, London, 1967.
- [7] E. MUES, Über ein Problem von Hayman, Math. Z., 164 (1979), 239-259.
- [8] Q. D. ZHANG, On the value distribution of  $\varphi(z) f'(z)$ , Acta Math. Sinica, **37** (1994), 91–97 (in Chinese).
- [9] Z. F. ZHANG AND G. D. SONG, On the zeros of  $f(f^{(k)})^n a(z)$ , Chinese Ann. Math. Ser. A, **19** (1998), 275–282.

DEPARTMENT OF MATHEMATICS THE HONG KONG UNIVERSITY OF SCIENCE & TECHNOLOGY CLEAR WATER BAY, KOWLOON, HONG KONG e-mail: makwing@ust.hk

(CURRENT ADDRESS) RM. 205, KWAI SHUN HSE., KWAI FONG EST., K.C., HONG KONG e-mail: kitwing@hotmail.com