

## A “DOUBLE” CAUCHY-SCHWARZ TYPE INEQUALITY

Manolis Magiropoulos and Dimitri Karayannakis

**Abstract.** A “double” version of the C-S inequality in any complex pre-Hilbert space is given, along with some numerical applications.

A substantial part of the mathematical folklore in the frame of inner-product spaces involves applications and/or extensions of the Cauchy-Schwarz (C-S) inequality

$$|\langle e, f \rangle| \leq \|e\| \|f\|,$$

where by  $\|\cdot\|$  we indicate the norm induced by the inner-product  $\langle \cdot, \cdot \rangle$ .

For the case of a (real) Hilbert space M. Lambrou (Univ. of Crete) indicated to the second author, by personal communication, the following “double” version of the C-S inequality.

$$|\langle e, f \rangle| |\langle e, g \rangle| \leq \frac{1}{2} \{ \|f\| \|g\| + |\langle f, g \rangle| \} \|e\|^2. \quad (*)$$

(Note that if  $f, g$  are considered to be linearly dependent we simply obtain the C-S inequality.)

The fact that the R.H.S. of (\*) provides a better bound than the “natural”  $\|e\|^2 \|f\| \|g\|$  is evident by use of the C-S inequality itself. That (\*) gives, in certain cases, a much better bound for  $|\langle e, f \rangle| |\langle e, g \rangle|$  becomes clear from the following example.

Let  $f = \sin x$ ,  $g = \cos x$ ,  $e = 1/x$  be considered as members of the classical Hilbert space  $L^2[\alpha, \beta]$ ,  $\alpha > 0$ . Then

$$|\langle e, f \rangle| |\langle e, g \rangle| \leq \left( \int_{\alpha}^{\beta} \frac{dx}{x^2} \right) \left( \int_{\alpha}^{\beta} \sin^2 x dx \right)^{1/2} \left( \int_{\alpha}^{\beta} \cos^2 x dx \right)^{1/2}.$$

On the other hand (\*) provides the bound

$$\frac{1}{2} \left( \int_{\alpha}^{\beta} \frac{dx}{x^2} \right) \left[ \left( \int_{\alpha}^{\beta} \sin^2 x dx \right)^{1/2} \left( \int_{\alpha}^{\beta} \cos^2 x dx \right)^{1/2} + \left| \int_{\alpha}^{\beta} \sin x \cos x dx \right| \right].$$

Subtracting the (\*)-bound from the C-S-bound one has

$$\frac{1}{8} \left( \frac{1}{\alpha} - \frac{1}{\beta} \right) [(\beta - \alpha + \sin 2\alpha - \sin 2\beta)^{1/2} (\beta - \alpha + \sin 2\beta - \sin 2\alpha)^{1/2} - |\cos 2\beta - \cos 2\alpha|],$$

which increases to  $+\infty$  for a (number of) suitable limit behavior of  $\alpha$  or  $\beta$ .

In the present work we present an elementary proof of (\*) for any pre-Hilbert space over the complex field (which naturally also covers the real case), along with a few applications of (\*).

Theorem 1. Let  $e, f, g$  be elements of a complex pre-Hilbert space  $(H, \langle, \rangle)$ ; then

$$2|\langle e, f \rangle| |\langle e, g \rangle| \leq \{\|f\| \|g\| + |\langle f, g \rangle|\} \|e\|^2.$$

Proof. Based on a previous remark, let  $f, g$  be linearly independent, and let  $e = \lambda f + \mu g$ . W.L.O.G. we may also assume that  $\|e\| = \|f\| = \|g\| = 1$ . If  $f \perp g$  the L.H.S. of (\*) becomes  $2|\lambda| |\mu|$  whereas the R.H.S. becomes  $|\lambda|^2 + |\mu|^2$  and we are done. If  $f$  and  $g$  are not orthogonal to each other, by the Gram-Schmidt construction we obtain  $e = kf + sh$  with  $h = (g - cf)(1 - |c|^2)^{-1/2}$  with  $c = \langle g, f \rangle \neq 0, 1$ . Then  $h \perp f$  and  $\|h\| = 1$ . It is easily seen that we may consider  $c > 0$  since, otherwise, by switching from  $f$  to  $(cf)/|c|$  we find ourselves in an equivalent position. Then  $|\kappa|^2 + |s|^2 = 1$  with  $\kappa = \langle e, f \rangle$  and  $s = \langle e, h \rangle$ . In a similar manner we may assume that  $\kappa \geq 0$ . Then

$$|\langle e, f \rangle| |\langle e, g \rangle| = \kappa |\kappa c + s(1 - c^2)^{1/2}|.$$

But

$$\begin{aligned} |\kappa c + s(1 - c^2)^{1/2}|^2 &= \kappa^2 c^2 + 2\text{Re}(\kappa c(1 - c^2)^{1/2} s) + |s|^2(1 - c^2) \\ &\leq (\kappa c + |s|(1 - c^2)^{1/2})^2. \end{aligned}$$

Thus,

$$|\langle e, f \rangle| |\langle e, g \rangle| \leq \kappa^2 c + \kappa(1 - \kappa^2)^{1/2}(1 - c^2)^{1/2}. \tag{**}$$

Using the first derivative criterion, etc. for local extremes, it can be easily seen that the R.H.S. of (\*\*) is bounded by  $(1 + c)/2$  and we are done.

It remains now to prove (\*) for the case  $e \notin \text{sp}\{f, g\}$ . Then,  $e = e_1 + e_2$  with  $e_1 \in \text{sp}\{f, g\}$  and  $e_2 \perp \text{sp}\{f, g\}$ . The L.H.S. of (\*) becomes

$$2|\langle e, f \rangle| |\langle e, g \rangle| \leq \{\|f\| \|g\| + |\langle f, g \rangle|\} \|e_1\|^2,$$

because of the first part. Since  $\|e_1\| \leq \|e\|$  we obtain the announced result. Q.E.D.

In case  $(H, \langle \cdot, \cdot \rangle)$  is Hilbert, we can generalize (\*) as follows.

**Theorem 2.** For any projection  $P$  and any vectors  $f, g$

$$2|\langle Pf, g \rangle| \equiv 2|\langle Pf, Pg \rangle| \leq \|f\| \|g\| + |\langle f, g \rangle|.$$

**Proof.** Let  $Q = I - P$ . Then

$$\|f\|^2 = \|Pf\|^2 + \|Qf\|^2, \quad \|g\|^2 = \|Pg\|^2 + \|Qg\|^2.$$

Since

$$\langle f, g \rangle = \langle Pf, Pg \rangle + \langle Qf, Qg \rangle,$$

by the classical Schwarz inequality we have

$$|\langle f, g \rangle| \geq |\langle Pf, Pg \rangle| - \|Qf\| \cdot \|Qg\|.$$

Therefore, for the assertion, it suffices to prove that

$$|\langle Pf, Pg \rangle| + \|Qf\| \cdot \|Qg\| \leq \sqrt{(\|Pf\|^2 + \|Qf\|^2)(\|Pg\|^2 + \|Qg\|^2)}.$$

Then, using the classical Schwarz inequality once more, it suffices to prove

$$\|Pf\| \cdot \|Pg\| + \|Qf\| \cdot \|Qg\| \leq \sqrt{(\|Pf\|^2 + \|Qf\|^2)(\|Pg\|^2 + \|Qg\|^2)}$$

which is nothing but the classical Cauchy inequality.

**Remark.** The result of (\*) in the case where  $(H, \langle \cdot, \cdot \rangle)$  of Theorem 1 is considered complete corresponds to the case where  $\text{rank}(P) = 1$ .

We turn now to a couple of applications of Theorem 1 starting with a complex Hilbert space  $(H, \langle \cdot, \cdot \rangle)$ .

(i) Notice first that if  $f$  and  $g$  in  $H$  are mutually orthogonal then (\*) reduces to

$$2|\langle e, f \rangle| |\langle e, g \rangle| \leq \|e\|^2 \|f\| \|g\| \quad (***)$$

If now  $e \in H$  and  $\{f_i\}$ ,  $i \in I$ , an orthogonal family in  $H$ , we can consider finite products of the Fourier coefficients of  $e$  w.r.t.  $\{f_i\}$  namely

$$\prod_{j \in J} |\langle e, f_j \rangle|,$$

where  $J \subset I$  a finite set with at least two elements (in which case, naturally, we also impose  $\dim H \geq 2$ ). It is evident that the C-S inequality would have provided the crude (upper) bound  $\|e\|^n$ , where  $n$  is the cardinality of  $J$ .

In view of (\*\*\*) though, we obtain the following far better bound, namely:

$$\prod_{j \in J} |\langle e, f_j \rangle| \leq \begin{cases} 2^{-n/2} \|e\|^n, & \text{if } n \text{ is even;} \\ 2^{-(n-1)/2} \|e\|^n, & \text{if } n \text{ is odd.} \end{cases}$$

(ii) For another application of (\*) let us employ the pre-Hilbert space  $L^1(0, +\infty)$  and consider its (linearly independent) elements

$$\frac{\sin x}{x}, \quad \frac{\cos 2x}{1+x^2} \quad \text{and} \quad \frac{1}{1+x^2}$$

in the roles of  $e$ ,  $f$  and  $g$ , respectively. The direct calculation of  $I = \langle e, f \rangle$  is a rather painful experience within the techniques of contour integration, or even by tracing it in suitable tables.

On the other hand by elementary contour integration and/or by reference to [1], we have

$$\langle e, g \rangle = \frac{\pi}{2}(1 - e^{-1}), \quad \|f\| = \left[ \frac{\pi}{8}(1 + 5e^{-4}) \right]^{1/2}, \quad \|g\| = \frac{\pi^{1/2}}{2},$$

$$\langle f, g \rangle = \frac{3}{4}\pi e^{-2}, \quad \text{and} \quad \|e\|^2 = \frac{\pi}{2}.$$

Thanks to (\*) we obtain the following (strict) upper bound

$$I < \frac{\pi}{8(1 - e^{-1})} \left[ \left( \frac{1}{2} + \frac{5}{2}e^{-4} \right)^{1/2} + 3e^{-2} \right] < 0.7112.$$

Reference

1. I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products*, ed. A. Jeffrey, Academic Press, 1980.

Manolis Magiropoulos  
Technological & Educational Institute of Crete  
Heraklion, Greece  
email: mageir@stef.teiher.gr

Dimitri Karayannakis  
Technological & Educational Institute of Crete  
Heraklion, Greece  
email: dkar@stef.teiher.gr