

Bibliography

- [1] Y. Akizuki and S. Nakano, Note on Kodaira–Spencer’s proof of Lefschetz theorems, Proc. Japan Acad. **30** (1954), 266–272.
- [2] M. Artin, Algebraization of formal moduli II: Existence of the formal modifications, Ann. of Math., **91** (1970), 88–135.
- [3] S. Bando and Y.-T. Siu, Stable sheaves and Einstein-Hermitian metrics, in *Geometry and Analysis on Complex Manifold, Festschrift for Prof. S. Kobayashi’s 60th Birthday* (T. Mabuchi, J. Noguchi and T. Ochiai, eds.), 1994, World Scientific, pp. 39–50.
- [4] C. Bănică and O. Stănişilă, *Algebraic methods in the global theory of complex spaces* Translated from the Romanian version (Editura Academiei, 1974); John Wiley & Sons, 1976.
- [5] A. Borel and J.-P. Serre, Le théorème de Riemann–Roch (d’après Grothendieck), Bull. Soc. Math. France **86** (1958), 97–136.
- [6] *Algebraic Geometry, Bowdoin, 1985* (S. Bloch ed.), Proc. of Symp. in Pure Math., **46** (1987) Amer. Math. Soc.
- [7] E. Cattani, A. Kaplan, W. Schmid, Degeneration of Hodge structures, Ann. of Math., **123** (1986), 457–535.
- [8] S. D. Cutkosky, Zariski decomposition of divisors on algebraic varieties, Duke Math. J., **53** (1986), 149–156.
- [9] V. I. Danilov, The geometry of toric varieties, Russian Math. Surveys, **33** (1978), 97–154.
- [10] P. Deligne, Théorie de Hodge, II, Publ. Math. IHES., **40** (1972), 5–57.
- [11] P. Deligne and L. Illusie, Relèvements modulo p^2 et décomposition du complexe de de Rham, Invent. Math., **89** (1987), 247–270.
- [12] S. Donaldson, Anti-self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles, Proc. London Math. Soc., **50** (1985), 1–26.
- [13] H. Esnault and E. Viehweg, Logarithmic De Rham complexes and vanishing theorems, Invent. Math., **86** (1986), 161–194.
- [14] O. Forster and K. Knorr, Ein Beweis des Grauertschen Bildgarbensatzes nach Ideen von B. Malgrange, Manuscripta Math., **5** (1971), 19–44.
- [15] *The Birational Geometry of Degenerations* (R. Friedman and D. Morrison eds.), Prog. in Math., **29** (1983) Birkhäuser.
- [16] J. Frisch, Points de platitude d’un morphisme d’espaces analytiques complexes, Invent. Math., **4** (1967), 118–138.
- [17] A. Fujiki, On the blowing down of analytic spaces, Publ. RIMS. Kyoto Univ., **10** (1975), 473–507.
- [18] ———, Closedness of the Douady spaces of compact Kähler spaces, Publ. RIMS. Kyoto Univ., **14** (1978), 1–52.
- [19] Y. Fujimoto, Logarithmic transformations of elliptic fiber spaces, J. Math. Kyoto Univ., **28** (1988), 91–110.
- [20] T. Fujita, On Zariski problem, Proc. Japan Acad. Ser. A, **55** (1979), 106–110.
- [21] ———, The sheaf of relative canonical forms of a Kähler fibre space over a curve, Proc. Japan Acad., **54** (1978), 183–184.
- [22] ———, On Kähler fiber spaces over a curve, J. Math. Soc. Japan, **30** (1978), 779–794.

- [23] ———, On L -dimension of coherent sheaves, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, **28** (1981), 215–236; Correction **29**, 719–720.
- [24] ———, Semipositive line bundles, *J. Fac. Sci. Univ. Tokyo, Sec. IA*, **30** (1983), 353–378.
- [25] ———, Zariski decomposition and canonical rings of elliptic threefolds, *J. Math. Soc. Japan*, **38** (1986), 19–37.
- [26] ———, Approximating Zariski decomposition of big line bundles, *Kodai Math. J.*, **17** (1994), 1–3.
- [27] ———, On Kodaira energy of polarized log varieties, *J. Math. Soc. Japan*, **48** (1996), 1–12.
- [28] H. Grauert, *Ein Theorem der analytischen Garbentheorie und die Modulräume komplexer Strukturen*, *Publ. Math. IHES.*, **5** (1960).
- [29] H. Grauert and R. Remmert, Bilder und Urbilder analytischer Garben, *Ann. of Math.*, **68** (1958), 393–443.
- [30] H. Grauert and O. Riemenschneider, Verschwindungssätze für analytische Kohomologiegruppen auf komplexen Räumen, *Invent. Math.*, **11** (1970), 263–292.
- [31] ———, Kählersche Mannigfaltigkeiten mit hyper- q -konvexem Rand, in *Problems in analysis* (Lectures Sympos. in honor of S. Bochner, Princeton Univ., 1969, R. Gunning ed.), Princeton Univ. Press, 1970, pp. 61–79.
- [32] P. Griffiths, Periods of integrals on algebraic manifolds I, II and III, *Amer. J. Math.*, **90** (1968), 568–626; ibid. **90** (1968), 805–865; *Publ. Math. IHES.*, **38** (1970), 125–180.
- [33] A. Grothendieck and J. Dieudonné *Éléments de géométrie algébrique II, Étude globale élémentaire de quelques classes de morphismes*, *Publ. Math. IHES.*, **8**, (1961).
- [34] ———, *Éléments de géométrie algébrique III, Étude cohomologique des faisceaux cohérents I*, *Publ. Math. IHES.*, **11**, (1961).
- [35] G. Harder and M. Narasimhan, On the cohomology groups of moduli space of vector bundles on curves, *Math. Ann.*, **212** (1975), 215–248.
- [36] R. Hartshorne, Ample vector bundles, *Publ. Math. IHES.*, **29** (1966), 63–94.
- [37] ———, *Residues and Duality*, Lecture Notes in Math., **20**, Springer-Verlag, 1966.
- [38] ———, *Algebraic geometry*, Graduate Texts in Mathematics, No. **52**, Springer-Verlag 1977.
- [39] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero, I, II, *Ann. of Math.*, **79** (1964), 109–326.
- [40] ———, Desingularization of complex-analytic varieties, in *Actes du Congrès International des Mathématiciens* (Nice, 1970), Tome 2, Gauthier-Villars, 1971, pp. 627–631.
- [41] ———, Flattening theorem in complex-analytic geometry, *Amer. J. Math.*, **97** (1975), 503–547.
- [42] S. Iitaka, Deformation of compact complex surfaces I, II, and III, in *Global Analysis, papers in honor of K. Kodaira*, Princeton Univ. Press 1969, pp. 267–272; *J. Math. Soc. Japan*, **22** (1970), 247–261; ibid. **23** (1971), 692–705.
- [43] ———, On D -dimensions of algebraic varieties, *J. Math. Soc. Japan*, **23** (1971), 356–373.
- [44] ———, *Algebraic Geometry – an Introduction to Birational Geometry of Algebraic Varieties*, Graduate Texts in Mathematics, No. **76**, Springer 1981.
- [45] ———, Basic structure of algebraic varieties, in [46], pp. 303–316.
- [46] *Algebraic Varieties and Analytic Varieties* (S. Iitaka ed.), Adv. Stud. in Pure Math., **1** (1983) Kinokuniya and North-Holland.
- [47] S. Ishii, Simultaneous canonical models of deformations of isolated singularities, in *Algebraic Geometry and Analytic Geometry* ICM-90 Sato. Conf. Proc. (1991) Springer, pp. 81–100.
- [48] M. Kashiwara, The asymptotic behavior of a variation of polarized Hodge structure, *Publ. RIMS. Kyoto Univ.*, **21** (1985), 853–875.
- [49] M. Kashiwara and T. Kawai, The Poincaré lemma for variations of polarized Hodge structure, *Publ. RIMS. Kyoto Univ.*, **23** (1987), 345–407.
- [50] Y. Kawamata, Characterization of abelian varieties, *Compo. Math.*, **43** (1981), 253–276.
- [51] ———, A generalization of Kodaira–Ramanujam’s vanishing theorem, *Math. Ann.*, **261** (1982), 43–46.

- [52] ———, Kodaira dimension of algebraic fiber spaces over curves, *Invent. Math.*, **66** (1982), 57–71.
- [53] ———, Kodaira dimension of certain algebraic fiber spaces, *J. Fac. Sci. Univ. Tokyo Sect. IA*, **30** (1983), 1–24.
- [54] ———, The cone of curves of algebraic varieties, *Ann. Math.*, **119** (1984), 603–633.
- [55] ———, Pluricanonical systems on minimal algebraic varieties, *Invent. Math.*, **79** (1985), 567–588.
- [56] ———, Minimal models and the Kodaira dimension of algebraic fiber spaces, *J. Reine Angew. Math.*, **363** (1985), 1–46.
- [57] ———, The Zariski decomposition of log-canonical divisors, in [6], pp. 425–433.
- [58] ———, Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces, *Ann. of Math.*, **127** (1988), 93–163.
- [59] ———, Abundance theorem for minimal threefolds, *Invent. Math.*, **108** (1992), 229–246.
- [60] ———, Deformation of canonical singularities, *J. Amer. Math. Soc.*, **12** (1999), 85–92.
- [61] Y. Kawamata, K. Matsuda and K. Matsuki, Introduction to the minimal model problem, in [111], pp. 283–360.
- [62] G. Kempf, F. Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal Embeddings*, Lecture Notes in Math., **339**, Springer, 1973.
- [63] R. Kiehl and J. Verdier, Ein einfacher Beweis des Kohärenzsatzes von Grauert, *Math. Ann.*, **195** (1971), 24–50.
- [64] S. Kleiman, Toward a numerical theory of ampleness, *Ann. of Math.*, **84** (1966), 293–344.
- [65] F. Knöller, 2-dimensionale singularitäten und differentialformen, *Math. Ann.*, **206** (1973), 205–213.
- [66] S. Kobayashi, First Chern class and holomorphic tensor fields, *Nagoya Math. J.*, **77** (1980), 5–11.
- [67] K. Kodaira, On a differential-geometric method in the theory of analytic stacks, *Proc. Nat. Acad. Sci. USA* **39** (1953), 1268–1273.
- [68] ———, On the Kähler varieties of restricted type (An intrinsic characterization of algebraic varieties), *Ann. of Math.*, **60** (1954), 28–48.
- [69] ———, On complex analytic surfaces II, III, *Ann. of Math.*, **77** (1963), 563–626; ibid. **78** (1963), 1–40.
- [70] ———, On the structure of compact complex analytic surfaces, I, *Amer. J. Math.*, **86** (1964), 751–798.
- [71] J. Kollar, Higher direct images of dualizing sheaves I, II, *Ann. of Math.*, **123** (1986), 11–42; ibid. **124** (1986), 171–202.
- [72] ———, Subadditivity of the Kodaira dimension: Fibers of general type, in [111], pp. 361–398.
- [73] J. Kollar and S. Mori, Classification of three dimensional flips, *J. Amer. Math. Soc.*, **5** (1992), 533–703.
- [74] J. Kollar et al, *Flips and Abundance for Algebraic Threefolds*, Astérisque **211**, Soc. Math. de France, 1992.
- [75] M. Levine, Pluri-canonical divisors on Kähler manifolds, and II, *Invent. Math.*, **74** (1983), 293–903; *Duke Math. J.*, **52** (1985), 61–65.
- [76] T. Luo, A note on the Hodge index theorem, *Manusc. Math.*, **67** (1990), 17–20.
- [77] K. Matsuda, On the numerically fixed parts of line bundles, *Proc. Japan Acad. Ser. A*, **61** (1985), 219–221.
- [78] V. B. Mehta and A. Ramanathan, Semi-stable sheaves on projective varieties and their restriction to curves, *Math. Ann.*, **258** (1982), 213–224.
- [79] ———, Restriction of stable sheaves and representations of the fundamental group, *Invent. Math.*, **77** (1984), 163–172.
- [80] M. Miyanishi, Projective degeneration of surfaces according to S. Tsunoda, in [111], pp. 415–447.

- [81] Y. Miyaoka, Deformation of a morphism along a foliation, in [6], pp. 245–268.
- [82] ———, The Chern classes and the Kodaira dimension of a minimal variety, in [111], pp. 449–476.
- [83] ———, On the Kodaira dimension of minimal threefolds, Math. Ann., **281** (1988), 325–332.
- [84] ———, Abundance conjecture for 3-folds: case $\nu = 1$, Compo. Math., **68** (1988), 203–220.
- [85] S. Mori, Projective manifolds with ample tangent bundles, Ann. of Math., **110** (1979), 593–606.
- [86] ———, Threefolds whose canonical bundles are not numerically effective, Ann. of Math., **116** (1982), 133–176.
- [87] ———, On 3-dimensional terminal singularities, Nagoya Math. J., **98** (1985), 43–66.
- [88] ———, Classification of higher dimensional varieties, in [6], pp. 269–331.
- [89] ———, Flip theorem and the existence of minimal models for 3-folds, J. Amer. Math. Soc., **1** (1988), 117–253.
- [90] S. Mori et al, *Birational Geometry of Algebraic Varieties, Open Problems*, Proc. of the XXIIrd international symposium, Division of Math., Aug. 22–Aug. 27, 1988 Katata, The Taniguchi Foundation.
- [91] A. Moriwaki, Semi-ampleness of the numerically effective part of Zariski-decomposition, J. Math. Kyoto Univ., **26** (1986), 461–481.
- [92] ———, Torsion freeness of higher direct images of canonical bundle, Math. Ann., **276** (1987), 385–398.
- [93] ———, Problem of Zariski decomposition, in [90], pp. 52–56.
- [94] I. Nakamura, Complex parallelisable manifolds and their small deformations, J. Diff. Geom., **10** (1975), 85–112.
- [95] S. Nakano, Vanishing theorems for weakly 1-complete manifolds, and II, in *Number theory, Algebraic Geometry and Commutative Algebra, in honor of Y. Akizuki* (Y. Kusunoki et al, eds), Kinokuniya (1973), pp. 169–179; Publ. RIMS. Kyoto Univ., **10** (1974), 101–110.
- [96] N. Nakayama, Invariance of the plurigenera of algebraic varieties under minimal model conjectures, Topology, **25** (1986), 237–251.
- [97] ———, Hodge filtrations and the higher direct images of canonical sheaves, Invent. Math., **85** (1986), 217–221.
- [98] ———, The lower semi-continuity of the plurigenera of complex varieties, in [111], pp. 551–590.
- [99] ———, Remarks on \mathbb{Q} -factorial singularities, preprint Univ. Tokyo 1986.
- [100] ———, Open problems, in [90], pp. 61–62.
- [101] ———, Zariski decomposition problem for pseudo-effective divisors, in Proc. of the Meeting and Workshop *Algebraic Geometry and Hodge Theory* vol. 1 (K. Konno, M.-H. Saito, S. Usui eds.), Hokkaido Univ. Tech. Rep. ser. in Math. #16 (1990), pp. 189–217.
- [102] ———, On smooth exceptional curves in threefolds, J. Fac. of Sci. Univ. Tokyo Sec. IA, **37** (1990), 511–525.
- [103] ———, A counterexample to the Zariski-decomposition conjecture (Japanese), in Proc. of the Workshop *Hodge Theory and Algebraic Geometry* at Hokkaido Univ. June 1994, Infinite Analysis Lecture Note No. **10** Kyoto Univ. (1994), pp. 96–101.
- [104] ———, Zariski-decomposition and abundance, preprint RIMS-1142, Kyoto Univ. 1997.
- [105] ———, Invariance of the plurigenera of algebraic varieties, preprint RIMS-1191, Kyoto Univ. 1998.
- [106] ———, Normalized tautological divisors of semi-stable vector bundles, preprint RIMS-1214, Kyoto Univ. 1998.
- [107] M. S. Narasimhan and C. S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface, Ann. of Math., **82** (1965), 540–567.
- [108] K. Nishiguchi, Kodaira dimension is not necessarily lower semicontinuous under degenerations of surfaces, Math. Ann., **263** (1983), 377–383.
- [109] ———, Degeneration of K3 surfaces, J. Math. Kyoto Univ., **28** (1988), 267–300.

- [110] T. Oda, *Convex Bodies and Algebraic Geometry* (Japanese), Kinokuniya 1985; English version: *Convex Bodies and Algebraic Geometry – An Introduction to the Theory of Toric Varieties*, *Ergebnisse der Math.*, (3) **15**, Springer-Verlag, 1988.
- [111] *Algebraic Geometry, Sendai, 1985* (T. Oda ed.), *Adv. Stud. in Pure Math.*, **10** (1987) Kinokuniya and North-Holland.
- [112] T. Oomae, Master's thesis, Dept. of Math. Kyoto Univ. 1998.
- [113] M. Reid, Canonical 3-folds, in *Géométrie Algébrique Angers 1979* (A. Beauville ed.), (1980) Alphen aan den Rijn, pp. 273–310.
- [114] ———, Minimal model of canonical 3-folds, in [46], pp. 131–180.
- [115] ———, Projective morphisms according to Kawamata, preprint, Univ. Warwick 1983.
- [116] J.-P. Ramis and G. Ruget, Complexe dualisant et théorèmes de dualité en géométrie analytique complexe, *Publ. Math. I.H.E.S.*, **38** (1970), 77–91.
- [117] J.-P. Ramis, G. Ruget, and J.-L. Verdier, Dualité relative en géométrie analytique complexe, *Invent. Math.*, **13** (1971), 261–283.
- [118] R. Remmert and K. Stein, Über die wesentlichen Singularitäten analytischer Mengen, *Math. Ann.*, **126**, (1953) 263–306.
- [119] M. Saito, Modules de Hodge polarisables, *Publ. RIMS. Kyoto Univ.*, **24** (1988), 849–995.
- [120] ———, Mixed Hodge modules, *Publ. RIMS. Kyoto Univ.*, **26** (1990), 221–333.
- [121] ———, Decomposition theorem for proper Kähler morphisms, *Tohoku Math. J.*, **42** (1990), 127–148.
- [122] ———, On Kollar's conjecture, in *Several Complex Variables and Complex Geometry*, Proceedings of Symp. in Pure Math. **52**, Part **2** (1991), Amer. Math. Soc., pp. 509–517.
- [123] F. Sakai, Degeneracy of holomorphic maps with ramification, *Invent. Math.*, **26** (1974), 213–229.
- [124] ———, Kodaira dimensions of complements of divisors, in *Complex Analysis and Algebraic Geometry* (W. L. Baily, Jr. and T. Shioda eds.), Iwanami Shoten and Cambridge Univ. Press, 1977, pp. 239–257.
- [125] I. Satake, Toric bundles over abelian schemes, in *Algebraic cycles and related topics* (Kitasakado, 1994), World Sci. Publ., 1995, pp. 43–49.
- [126] W. Schmid, Variation of Hodge structures: The singularities of period mapping, *Invent. Math.*, **22** (1973), 211–319.
- [127] J. P. Serre, Faisceaux algébriques cohérents, *Ann. Math.*, **61** (1955), 197–278.
- [128] ———, Géométrie algébrique et géométrie analytique, *Ann. Inst. Fourier*, **6** (1956), 1–42.
- [129] Y.-T. Siu, Noetherianness of rings of holomorphic functions on Stein compact series, *Proc. Amer. Math. Soc.*, **21** (1969), 483–489.
- [130] ———, Invariance of plurigenera, *Invent. Math.*, **134** (1998), 661–673.
- [131] V. Shokurov, The non-vanishing theorem, *Izv. Akad. Nauk SSSR Ser. Mat.*, **49** (1985), 635–651; *Math. USSR. Izv.*, **26** (1985), 591–604.
- [132] ———, 3-fold log flips, *Izv. Ross. Akad. Nauk Ser. Mat.*, **56** (1992), 105–203; English transl. in *Russian Acad. Sci. Izv. Math.*, **40** (1993), 95–202.
- [133] S. Suzuki, Birational geometry of birational pairs, *Comment. Math. Univ. St. Pauli*, **32** (1983), 85–106.
- [134] E. Szabó, Divisorial log terminal singularities, *J. Math. Sci. Univ. Tokyo*, **1** (1994), 631–639.
- [135] K. Takegoshi, Higher direct images of canonical sheaves tensorized with semi-positive vector bundles by proper Kähler morphisms, *Math. Ann.*, **303** (1995), 389–416.
- [136] P. Thullen, Über die wesentlichen Singularitäten analytischer Funktionen und Flächen in Räumen von n komplexen Veränderlichen, *Math. Ann.*, **111**, (1935) 137–157.
- [137] S. Tsunoda, Degeneration of surfaces, in [111], pp. 755–764.
- [138] S. Tsunoda and M. Miyanishi, The structure of open algebraic surfaces II, in [141], pp. 499–544.
- [139] K. Ueno, *Classification Theory of Algebraic Varieties and Compact Complex Spaces*, Lecture Notes in Math., **439** Springer, 1975.

- [140] ———, Degeneration of elliptic surfaces and certain non-Kähler manifolds, in [141], pp. 545–566.
- [141] *Classification of Algebraic and Analytic Manifolds* Proc. Katata Sympo. 1982 (K. Ueno ed.), Prog. in Math., **39** (1983) Birkhäuser.
- [142] K. Uhlenbeck and S. T. Yau, On the existence of Hermitian-Yang-Mills connections in stable vector bundles; and its note, Comm. Pure and Appl. Math., **39** (1986), S257–S293; ibid. **42** (1989), S703–S707.
- [143] J. Varouchas, Stabilité de la classe des variétés kähleriennes pour les certains morphismes propres, Invent. Math., **77** (1984), 117–128.
- [144] J.-L. Verdier, Dualité dans la cohomologie des espaces localement compacts, Séminaire Bourbaki, 18e année 65/66, no. **300**.
- [145] ———, Base change for twisted inverse image of coherent sheaves, in *Algebraic Geometry* (Internat. Colloq., Tata Inst. Fund. Res., Bombay, 1968), Oxford Univ. Press, 1969 pp. 393–408.
- [146] E. Viehweg, Vanishing theorems, J. Reine. Angew. Math., **335** (1982), 1–8.
- [147] ———, Weak positivity and the additivity of the Kodaira dimension for certain fiber spaces, in [46], pp. 329–353.
- [148] ———, Weak positivity and the additivity of the Kodaira dimension II – the local Torelli map, in [141], pp. 567–589.
- [149] ———, Weak positivity and the stability of certain Hilbert points, Invent. Math., **96** (1989), 636–667.
- [150] S. T. Yau, On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equations, I, Comm. Pure and Appl. Math., **31** (1978), 339–411.
- [151] O. Zariski, The theorem of Riemann-Roch for high multiples of an effective divisor on an algebraic surface, Ann. of Math., **76** (1962), 560–615.