

On the Insufficiency of Linear Diagrams for Syllogisms

OLIVER LEMON and IAN PRATT

Abstract In Volume 33:1 of the *Notre Dame Journal of Formal Logic*, a system for diagramming syllogistic inferences using straight line segments is presented by Englebretsen. In light of recent research on the representational power of diagrammatic representation systems by the authors, we point out some problems with the proposal, and indeed, with any proposal for representing logically possible situations diagrammatically. We shall first outline the proposed linear diagrammatic system of Englebretsen, and then show by means of counterexamples that it is inadequate as a representation scheme for general logical inferences (the task for which the system is intended). We also show that modifications to the system fail to remedy the problems. The considerations we present are not limited to the particular proposal of Englebretsen; we thus draw a more general moral about the use of spatial relations in representation systems.

1 Diagrammatic representation systems Diagrammatic representation systems are of increasing interest for at least two reasons. Philosophically, diagram systems interest those concerned with the nature of representation itself—in particular, those who argue that too much attention has been given to sequential symbol systems. These writers claim that diagrams represent by *analogy* or *surrogacy*—in virtue of sharing structure with the domains that they represent (see Barwise and Shimojima [2], Cummins [4], and Swoyer [21]). Practically, diagrammatic representations are frequently used in visual interfaces to databases, programming languages, and in logic teaching. Each of these domains demands careful consideration of the formal properties of the diagrammatic systems in question. For both of these reasons we propose to investigate the expressive power of one proposed diagram system and to determine its utility in reasoning tasks.

2 The diagrammatic system LD In [5], Englebretsen presents us with a system for diagramming syllogistic inferences. In this system, individuals are represented

Received September 24, 1997; revised October 31, 1999




Figure 1: Some philosophers are vegetarians; Descartes is a philosopher; no dogs are philosophers; all dalmatians are dogs; all dogs are carnivores.

by labeled points and sets are represented by labeled straight line segments. Relationships between sets and individuals or between sets and sets are then represented by incidence relations between the corresponding points and line segments. The author explains how such a representation system can be used to carry out syllogistic reasoning tasks and illustrates his explanation with many examples. We shall call the basic system of [5] (that is, without representation of relations or pronouns) “LD” for “linear diagrams”. Figure 1 shows an example of a diagram of LD together with its intended interpretation.

More formally, we may reconstruct LD as follows. A linear diagram is a finite set of *dots* and *dashes*. A dot is simply a labeled point in the plane and a dash is a finite line segment with a dot attached to its right terminus. The restriction to right termini is Englebretsen’s, not ours: Englebretsen does not consider vertical dashes, but we may assume that these are labeled by dots at their upper terminus. The terminology of *dots* and *dashes* is ours, not Englebretsen’s: it is introduced simply to avoid confusion with the infinitely many unmarked points and lines in the space occupied by an LD diagram and changes nothing of substance.

LD is to be interpreted as follows.

Definition 2.1 (Linear diagrams, [5])

1. Dashes represent sets.
2. Any dot which is not the right terminus of some dash represents an individual.
3. The presence of a dot on a dash (other than its right terminus) indicates that the individual represented by the dot is a member of the set represented by the dash.
4. That two dashes intersect represents the fact that the sets they represent have a nonempty intersection.
5. That two dashes do not intersect represents the fact that the sets they represent have an empty intersection.
6. That a dash l lies within a dash l' represents the fact that the set represented by l is a subset of the set represented by l' .

Actually, Englebretsen further insists that dashes representing a set L and not- L (the complement of L) must be parallel. However, given that line segments rather than lines are employed in the system, it is unclear precisely what is gained by insisting that line segments representing complementary sets be parallel (as opposed to merely nonintersecting). On the other hand, it is not clear that any harm is done either.

Note that there is some ambiguity regarding the representation of individuals as dots. It is not clear from [5] whether two coincident dots must represent the same individual; nor is it clear whether one individual can be represented by multiple (in particular, noncoincident) dots. Englebretsen claims that

“ . . . identity statements are . . . easily diagrammed by our method. A proposition of the form ‘a is (identical to) b’ . . . means that ‘a’ and ‘b’ label the same point.” ([5], p. 47)

It is unclear how to interpret this pronouncement. Certainly, Englebretsen cannot wish to claim that every *point* in the plane can represent no more than one individual. For “some As are Bs” is to be represented by line intersection and that intersection, though a single point, may contain more than one individual. However, in the counterexamples which we present below, we take no particular stance as regards these issues.

Inference in LD is to be carried out, as is usual with diagrammatic representations, by enumeration of cases. That is, a conclusion follows from a set of premises if all ways of diagramming the premises result in a diagram depicting the conclusion. Clearly, it follows from the premises of Figure 1 (some philosophers are vegetarians, Descartes is a philosopher, no dogs are philosophers, all dalmatians are dogs, all dogs are carnivores) that all dalmatians are carnivores, since this fact must hold (by part 6 of Definition 2.1) however, exactly, the diagram of the premises is drawn. Equally clearly, it does not follow that Descartes is a carnivore, because the diagram shows us a way for the premises to be true and the conclusion false.

While there has been much recent research in the logical analysis of diagrammatic reasoning (Allwein and Barwise [1], [2], Glasgow et al. [8], [16], Shin [17], Stenning and Oberlander [20]) we believe that some basic properties of diagrammatic representation systems deserve more attention, particularly when their suitability for performing logical inferences is in question. In particular, the following two properties of representation systems are of central importance.

1. For every representation of the system there is some possible situation of which it is true (*self-consistency*).
2. Every possible situation has some representation true of it.

The inference system of Euler’s Circles (see, e.g., Hammer [10]) has been shown [15] to exhibit the first, but not the second, of these properties, thus making it unsuitable for syllogistic inference in general. In particular, since not all logical possibilities can be represented in the system, attempts to diagram some situations will lead to incorrect inferences. In the present paper we perform a similar analysis for the proposal to use linear diagrams in syllogistic inference.

Englebretsen’s stated aim is to design a diagram system which is simple but which also avoids the expressive limitations imposed by the geometry of closed plane

figures (e.g., Euler’s Circles)—limitations which, according to Englebretsen, are often overlooked by those in the *diagrammatic reasoning* community. Thus,

“The geometric restrictions on closed plane figures which prevent *perspicuous* representations involving more than four terms using simple continuous figures do not apply to the still simpler linear figures.” ([5], p. 47, our italics)

“. . . the major advantage of line diagrams is their ability to represent inferences involving relatively large numbers of terms . . .” ([5], p. 46)

We are not told which geometric restrictions Englebretsen has in mind (the above claims are not proven formally); he merely mentions a four-term limit on the use of closed plane figures, due to geometric restrictions from the use of the plane as a representational medium. The reference given, (Gardner [7]), mentions only a *virtual four term limit* based on psychological rather than geometrical restrictions, which suggests that the restrictions Englebretsen seeks to avoid are practical in nature. Roughly, the problem seems to be that, when diagrammatic systems are used to represent large numbers of premises, the result tends to look like a plate of spaghetti.

We shall show in Sections 3 and 4 that, contrary to Englebretsen’s assertions, the system LD is subject to geometrical constraints which compromise its utility for logical inference, regardless of considerations of perspicuity and readability. In earlier work ([14, 15]), it has been demonstrated that other diagrammatic representation systems, based on the representation of sets by areas, fall victim to similar problems. Thus, Englebretsen’s claim to have overcome the expressive limitations imposed by plane geometry through the use of line segments rather than areas, cannot be maintained.

3 First counterexample to the correctness of LD Let $P_i (1 \leq i \leq 3)$ be individuals and $L_j (1 \leq j \leq 3)$ sets. Consider the following situation.

Individual P_i is a member of the set L_j if and only if $i \neq j (1 \leq i \leq 3, 1 \leq j \leq 3)$.

It is clear that this situation corresponds to a finite set of statements of the forms “ P is an L ” and “ P is not an L ”, and thus falls within the purview of LD. Indeed, it can be diagrammed as shown in Figure 3. Here, the individuals P_i are represented by dots $p_i (1 \leq i \leq 3)$ and the sets L_i are represented by the dashes $l_i (1 \leq i \leq 3)$.




Figure 2: A linear diagram.

But now let P_4 be a fourth individual and consider above the situation but augmented by

Individual P_4 is a member of the all the sets $L_j (1 \leq j \leq 3)$.

To see why this augmented situation cannot be diagrammed in LD, suppose that the new individual is represented by a dot p_4 . Then the four dots p_i must all be distinct (since no two may lie on exactly the same dashes l_j). It follows that the dashes l_1 and l_2 are collinear, since they both contain the distinct dots p_3 and p_4 ; by similar reasoning, l_1 and l_3 must be collinear, so all the dashes lie on some common line λ (say).

Orient the diagram so that λ is horizontal. Let us write $p < p'$ to indicate that dot p is to the left of dot p' (with the obvious interpretation for $p \leq p'$). Finally, let \dot{l}_j stand for the right terminus of the dash l_j ($1 \leq j \leq 3$). It follows from what we are told about L_1 that either $p_2, p_3, p_4 \leq \dot{l}_1 < p_1$ or $p_1 < p_2, p_3, p_4 \leq \dot{l}_1$. Whence, from what we are told about L_2 , either $p_2 < p_3, p_4 \leq \dot{l}_1 < p_1$ or $p_1 < p_3, p_4 < p_2 \leq \dot{l}_1$. Either way, p_3 lies between p_1 and p_2 , which contradicts what we are told about L_3 .

Note that this type of behavior means that LD could be used to make invalid inferences. For instance, suppose we omit from the above (augmented) situation the fact that individual P_1 does not belong to set L_1 . Then the above geometrical argument shows that all ways of diagramming the remaining facts will force the dot p_1 to lie on the dash l_1 , thus inviting the inference that P_1 belongs to L_1 . Of course, this inference would be invalid.

There is a way in which Englebretsen might save his system from the problem just raised. He could point out that the restriction to *straight* line segments is inessential. To be sure, the original motivation for using lines to represent sets was to avoid the unreadable diagrams resulting from region-based representation systems; but perhaps there is a middle way. For example, one might represent sets as connected chains of straight line segments, or even as arbitrary semi-algebraic curves. Since we can only guess at the possibilities here, we shall assume only that the plane figures used to represent sets are semi-algebraic curves. This generalization of Englebretsen's system, which we call *curved LD*, will now be investigated.

4 Second counterexample We now investigate *curved LD*—the generalization of LD employing semi-algebraic curves instead of straight line segments. Henceforth, we use the term *dash* to refer to semi-algebraic curves in a diagram, with dots at one of their endpoints, where again, we interpret dashes as representing sets. Where two sets have a nonempty intersection, we consider two cases for the curves which represent them; where the curves are only allowed to cross at one point (the *single-crossings* case) and where multiple crossings are permitted.

Let P_i ($1 \leq i \leq 5$) be individuals and L_{jk} ($1 \leq j < k \leq 5$) sets. Consider the following situation (we label it S).

Individual P_i is a member of the set L_{jk} if and only if $i = j$ or $i = k$.

No L_{jk} s are $L_{j'k}$ s if $\{j, k\} \cap \{j', k'\} = \emptyset$.

It is clear that this situation corresponds to a finite set of statements of the forms “ P is an L ”, “ P is not an L ”, and “No L s are L 's”, and thus falls within the purview of LD; let us see how we might represent it.

Let each individual P_i be represented by a dot p_i ($1 \leq i \leq 5$) and each set L_{ij} by a dash l_{ij} ($1 \leq i < j \leq 5$). Then the five dots p_i must all be distinct (since no two may lie on exactly the same dashes l_{kj}). Now, each dash l_{ij} ($1 \leq i < j \leq 5$) contains the dots p_i and p_j , so that every pair of the five points is to be joined by some dash (a semi-algebraic curve). In addition, the semi-algebraic curves l_{jk} and $l_{j'k'}$ may not intersect if they do not share one of the p_i .

In our first case (where curves are only allowed to cross each other *once* if their corresponding sets intersect), we have the result that the above situation S cannot be represented in curved LD, for in any such representation the dots p_i ($1 \leq i \leq 5$) and the sections of the dashes l_{ij} lying between p_i and p_j ($1 \leq i < j \leq 5$) constitute a drawing (that is, a plane embedding) of the $\alpha_{i,j}$ ($1 \leq i < j \leq 5$), which is the well-known nonplanar graph K_5 (see Bollobás [3]). Thus the situation S cannot be diagrammed in curved LD. It follows that, for any of the statements in the above situation, its negation will, according to the LD inference procedure, be implied by the others. Of course, such an inference would be invalid. Figure 3 shows one attempt to realize these premises in this version of curved LD. Any other attempt, in the single-crossings case, would fail similarly. Again, we have shown that there are consistent sets of statements falling within the purview of the system of Linear Diagrams, which cannot be represented by any Linear Diagram.




Figure 3: An unsuccessful attempt to diagram the second counterexample using curved LD. According to the premises, lines l_{35} and l_{24} should not cross. The nonplanarity result shows that any attempt to diagram the premises must result in a some such disallowed crossing.

This result holds for the case where curves are only allowed to cross once, if at all. The more general case is where curves may cross multiple times, if they cross at all. The result appears to be true for this case too, although it has proven difficult to demonstrate, so we leave the case open with the following conjecture.

Conjecture 4.1 *Let v_1, \dots, v_5 be distinct points in the plane and let $\alpha_{i,j}$ ($1 \leq i < j \leq 5$) be semi-algebraic curves, each with endpoints v_i and v_j . Then, for some arcs $\alpha_{i,j}, \alpha_{i',j'}$ we have $|\alpha_{i,j} \cap \alpha_{i',j'}| \neq \emptyset$ and $\{i, j\} \cap \{i', j'\} = \emptyset$.*

5 Conclusion We have shown that the proposed system of [5] for diagramming syllogistic inferences does not manage, by employing lines rather than regions, to avoid important geometric limitations on plane figures. The above examples show

that the diagram system cannot perform the representational task set for it. Thus, using the proposed representation system, or indeed slight generalizations of the proposal (employing semi-algebraic curves in place of straight line segments), would lead to mistakes in logical inferences.

The study of the system LD and its variants illustrates a general point about the representational use of spatial relations; that use of such relations is only appropriate in the representation of similarly constrained structures (for example, trivially, spatial objects and relations). The use of space in representations of more abstract structures, such as sets or models (for example, [9, 10]), is thus to be approached with some caution.

The second counterexample relies, in the single-crossings case, on the assumption that curves representing properties cross only once, if at all. We conjecture that the result extends to the multiple-crossings case. However, any system of diagrammatic reasoning which relied for its logical soundness on the drawing of multiply-crossing arcs could hardly be considered an intuitive or practical one.

Acknowledgments The support of the Leverhulme Trust (grant number FAQ/120/AQ) and of the British Council's British-German Academic Research Collaboration Programme (Project 720) is gratefully acknowledged.

REFERENCES

- [1] Allwein, G., and J. Barwise, eds., *Logical Reasoning with Diagrams*, Oxford University Press, Oxford, 1996. [Zbl 0870.03002](#) [MR 97c:03012](#) 2
- [2] Barwise, J., and A. Shimojima, "Surrogate reasoning," *Cognitive Studies: Bulletin of Japanese Cognitive Science Society*, vol. 4 (1995), pp. 7–27. 1, 2
- [3] Bollobás, B., *Graph Theory: An Introductory Course*, Springer, New York, 1979. [Zbl 0411.05032](#) [MR 80j:05053](#) 4
- [4] Cummins, R., *Representations, Targets, and Attitudes*, The MIT Press, Cambridge, 1996. 1
- [5] Englebretsen, G., "Linear diagrams for syllogisms (with relationals)," *Notre Dame Journal of Formal Logic*, vol. 33 (1992), pp. 37–69. [Zbl 0775.03001](#) [MR 94c:03014](#) 2, 2, 2.1, 2, 2, 2, 5
- [6] Englebretsen, G., *Something to Reckon With*, University of Ottawa Press, Ottawa, 1996.
- [7] Gardner, M., *Logic Machines and Diagrams*, Harvester Press, Sussex, 1958. [Zbl 0082.33701](#) [MR 87b:03024](#) 2
- [8] Glasgow, J., N. H. Narayanan, and B. Chandrasekaran, eds., *Diagrammatic Reasoning: Cognitive and Computational Perspectives*, AAAI Press/The MIT Press, Cambridge, 1995. 2
- [9] Hammer, E. M., "Reasoning with sentences and diagrams," *Notre Dame Journal of Formal Logic*, vol. 35 (1994) pp. 73–87. [Zbl 0801.03004](#) [MR 95d:03049](#) 5
- [10] Hammer, E. M., *Logic and Visual Information*, Studies in Logic, Language, and Computation, CSLI Publications and FoLLI, Stanford, 1995. [Zbl 0930.03002](#) [MR 97a:03001](#) 2, 5

- [11] Lemon, O., review of *Logic and Visual Information* by E. M. Hammer (CSLI Publications), *Journal of Logic, Language, and Information*, vol. 6 (1997), pp. 213–16.
- [12] Lemon, O., “Comparing the efficacy of lenses, lines, and logics,” *Logic, Language, and Diagrams*, edited by Barker-Plummer, Beaver, di Luzio, and van Benthem, CSLI Publications, Stanford, 2000 (in press).
- [13] Lemon, O., M. de Rijke, and A. Shimojima, “Efficacy of diagrammatic reasoning,” *Journal of Logic, Language, and Information*, vol. 8 (1999), pp. 265–71.
[Zbl 0941.03538](#)
- [14] Lemon, O., and I. Pratt, “Logical and diagrammatic reasoning: the complexity of conceptual space,” pp. 430–35 in *19th Annual Conference of the Cognitive Science Society*, edited by M. Shafto and P. Langley, Lawrence Erlbaum Associates, New Jersey, 1997.
[2](#)
- [15] Lemon, O., and I. Pratt, “Spatial logic and the complexity of diagrammatic reasoning,” *Machine Graphics and Vision*, Special Issue on Diagrammatic Representation and Reasoning, vol. 6 (1997), pp. 89–108. [2](#), [2](#)
- [16] Shimojima, A., “Reasoning with diagrams and geometrical constraints,” *Logic, Language and Computation*, Lecture Notes, CSLI, vol. 1 (1996), pp. 527–40.
[Zbl 0875.68786](#) [MR 1 396 556](#) [2](#)
- [17] Shin, S., *The Logical Status of Diagrams*, Cambridge University Press, Cambridge, 1995. [Zbl 0829.03002](#) [MR 95j:03014](#) [2](#)
- [18] Shin, S., and O. Lemon, “Diagrams,” *Stanford Encyclopedia of Philosophy*, forthcoming.
- [19] Stenning, K., and O. Lemon, “Aligning logical and psychological perspectives on diagrammatic reasoning,” *Artificial Intelligence Review*, forthcoming. [Zbl 0978.68112](#)
- [20] Stenning, K., and J. Oberlander, “A cognitive theory of graphical and linguistic reasoning: logic and implementation,” *Cognitive Science*, vol. 19 (1995), pp. 97–140. [2](#)
- [21] Swoyer, C., “Structural representation and surrogative reasoning,” *Synthese*, vol. 87 (1991), pp. 449–508. [MR 1 119 284](#) [1](#)

*CSLI**Stanford University**Room 102 Cordura Hall**210 Panama Street**Stanford CA 94305-4115**email: lemon@csli.stanford.edu**Department of Computer Science**University of Manchester**UNITED KINGDOM**email: ipratt@cs.man.ac.uk*