

GENERICALLY RATIONAL POLYNOMIALS

Dedicated to Professor Y. Nakai on his sixtieth birthday

MASAYOSHI MIYANISHI*) AND TOHRU SUGIE

(Received February 13, 1979)

Introduction. Let k be an algebraically closed field of characteristic zero. Let $k[x, y]$ be a polynomial ring of two variables and let $A_k^2 = \text{Spec}(k[x, y])$. Embed A_k^2 into the projective plane P_k^2 as the complement of a line l_∞ . Let $f \in k[x, y]$ be an irreducible polynomial, let F_α be the curve on A_k^2 defined by $f = \alpha$ for every $\alpha \in k$ and let C_α be the closure of F_α in P_k^2 . Then the set $\Lambda(f) := \{C_\alpha; \alpha \in k \cup (\infty)\}$ is a linear pencil on P_k^2 defined by f , where $C_\infty = dl_\infty$, d being the degree of f . The set $\Lambda_0(f) := \{F_\alpha; \alpha \in k\}$ is called *the linear pencil on A_k^2 defined by f* . The polynomial f is called *generically rational* when the general members of $\Lambda(f)$ (or $\Lambda_0(f)$) are irreducible rational curves. Since the algebraic function field $k(x, y)$ of one variable over the subfield $k(f)$ then has genus 0, Tsen's theorem says that f is generically rational if and only if f is a field generator in the sense of Russell [9, 10], i.e., there is an element $g \in k(x, y)$ such that $k(x, y) = k(f, g)$. If f is a generically rational polynomial, we can associate with f a non-negative integer n , where $n+1$ is the number of places at infinity of a general member F_α of $\Lambda_0(f)$, i.e., the number of places of F_α whose centers lie outside A_k^2 .

If $d \geq 1$, the pencil $\Lambda(f)$ has base points situated outside A_k^2 . Let $\varphi: W \rightarrow P_k^2$ be the shortest succession of quadratic transformations with centers at the base points (including infinitely near base points) of $\Lambda(f)$ such that the proper transform Λ' of $\Lambda(f)$ by φ has no base points. Then the linear pencil Λ' defines a morphism $\rho: W \rightarrow P_k^1$, whose general fibers are the proper transforms of general members of $\Lambda(f)$; thence they are nonsingular rational curves by virtue of Bertini's theorem. Moreover, W contains in a canonical way an open subset isomorphic to A_k^2 . A generically rational polynomial f is said to be of *simple type* if the morphism ρ has $n+1$ cross-sections contained in the boundary set $W - A_k^2$ (cf. Definition 1.8, below).

If $n=0$, a generically rational polynomial f is sent to one of the coordinates x, y of A_k^2 by a biregular automorphism of A_k^2 (cf. Abhyankar-Moh's theorem [1, 4]); hence f is of simple type. If $n=1$, a generically rational polynomial is always of simple type (cf. Theorem 2.3, below). However, if $n > 1$, a generi-

*) Supported by Grant-in-Aid for Scientific Research

cally rational polynomial is not necessarily of simple type, as clarified by Saito [13]. In effect, Saito determined in [12, 13] the standard forms of generically rational polynomials for $n=1, 2$ by analytic methods.

One of the purposes of the present article is to determine the standard forms of generically rational polynomials of simple type for arbitrary $n \geq 0$ by means of purely algebraic methods (cf. the results in §§2, 3).

The notations which we use frequently in this article are the following:

Let R be a k -algebra. Then R^* is the multiplicative group consisting of invertible elements of R ; R^* is called the unit group of R ; if R is a finitely generated, normal k -algebra then R^*/k^* is a finitely generated abelian group (cf. [3]). For elements x, y of R^* , we denote $x \sim y$ if $xy^{-1} \in k^*$.

Let V be a nonsingular projective surface. Let D_1, D_2 be divisors on V . Then $(D_1 \cdot D_2)$ is the intersection number (or multiplicity) of D_1, D_2 on V . Let C be a curve on V and let P be a point on C . Then $\text{mult}_P C$ is the multiplicity of C at P . If D_1 and D_2 are divisors, locally effective at P , then $i(D_1, D_2; P)$ is the local intersection multiplicity of D_1, D_2 at P . For an effective divisor D , we denote by $|D|$ the underlying reduced curve. (*)

Let $\varphi: W \rightarrow V$ be a birational morphism of nonsingular rational surfaces. If D is a divisor on V then $\varphi^*(D)$ (or $\varphi'(D)$, resp.) denotes the total transform (or the proper transform, resp.) of D by φ . Similarly, if Λ is a linear pencil on V then $\varphi'(\Lambda)$ denotes the proper transform of Λ by φ .

Let $\rho: V \rightarrow B$ be a surjective morphism from a nonsingular projective surface onto a nonsingular curve, whose general fibers are nonsingular rational curves; we call ρ a P^1 -fibration. An irreducible curve Γ on V is called a *cross-section* (or *quasi-section*, resp.) if $(\Gamma \cdot C) = 1$ (or $(\Gamma \cdot C) \geq 1$, resp.) for a general fiber C of ρ .

The ground field k is always assumed to be an algebraically closed field of characteristic zero. The affine space (or the projective space, resp.) of dimension n defined over k is denoted by A_k^n (or P_k^n , resp.).

1. Standard compactifications of A_k^2

1.1. Lemma (cf. Gizatullin [2]). *Let $\varphi: V \rightarrow B$ be a surjective morphism from a nonsingular projective surface V onto a nonsingular complete curve B such that almost all fibers are isomorphic to P_k^1 . Let $F = n_1 C_1 + \cdots + n_r C_r$ be a singular fiber of φ , where C_i is an irreducible curve, $C_i \neq C_j$ if $i \neq j$, and $n_i > 0$. Then we have:*

- (1) *The greatest common divisor (n_1, \dots, n_r) of n_1, \dots, n_r is 1; $\text{Supp}(F) = \bigcup_{i=1}^r C_i$ is connected.*

(*) The readers are warned of not confusing $|D|$ with the complete linear system determined by D . In the present article, we do not use the symbol $|D|$ to signify the latter meaning.

- (2) For $1 \leq i \leq r$, C_i is isomorphic to \mathbf{P}_k^1 and $(C_i^2) < 0$.
- (3) For $i \neq j$, $(C_i \cdot C_j) = 0$ or 1.
- (4) For three distinct indices i, j and l , $C_i \cap C_j \cap C_l = \phi$.
- (5) One of C_i 's, say C_1 , is an exceptional component, i.e., an exceptional curve of the first kind. If $\tau: V \rightarrow V_1$ is the contraction of C_1 , then φ factors as $\varphi: V \xrightarrow{\tau} V_1 \xrightarrow{\varphi_1} B$, where $\varphi_1: V_1 \rightarrow B$ is a fibration by \mathbf{P}^1 .
- (6) If one of n_i 's, say n_1 , equals 1 then there is an exceptional component among C_i 's with $2 \leq i \leq r$.

1.2. A generalization of Lemma 1.1 is the following

Lemma. Let V be a nonsingular projective surface and let Λ be an irreducible linear pencil on V such that general members of Λ are rational curves. Let \mathfrak{B} be the set of points of V which are base points of Λ . Let $F = n_1 C_1 + \dots + n_r C_r$, be a reducible member of Λ such that $r \geq 2$, where C_i is an irreducible component, $C_i \neq C_j$ if $i \neq j$, and $n_i > 0$. Then the following assertions hold true:

- (1) If $C_i \cap \mathfrak{B} = \phi$ then C_i is isomorphic to \mathbf{P}_k^1 and $(C_i^2) < 0$.
- (2) If $C_i \cap C_j \neq \phi$ for $i \neq j$ and $C_i \cap C_j \cap \mathfrak{B} = \phi$ then $C_i \cap C_j$ consists of a single point where C_i and C_j intersect each other transversally.
- (3) For three distinct indices i, j, l , if $C_i \cap C_j \cap C_l \cap \mathfrak{B} = \phi$ then $C_i \cap C_j \cap C_l = \phi$.
- (4) Assume that $(C_i^2) < 0$ whenever $C_i \cap \mathfrak{B} \neq \phi$. Then the set $S := \{C_i; C_i \text{ is an irreducible component of } F \text{ such that } C_i \cap \mathfrak{B} = \phi\}$ is nonempty, and there is an exceptional component in the set S .
- (5) With the same assumption as in (4) above, if a component of S , say C_1 , has multiplicity $n_1 = 1$ then there exists an exceptional component in S other than C_1 .

Proof. Let $\rho: \tilde{V} \rightarrow V$ be the shortest succession of quadratic transformations with centers at base points (including infinitely near base points) of Λ such that the proper transform $\tilde{\Lambda}$ of Λ by ρ has no base points. Then, by Bertini's theorem, general members of $\tilde{\Lambda}$ are isomorphic to \mathbf{P}_k^1 . The assertions (1), (2) and (3) are then apparently true by Lemma 1.1. We shall prove the assertions (4) and (5), assuming that $\mathfrak{B} \neq \phi$. Let $P \in \mathfrak{B}$. Set $P_0 := P$, and let P_1, \dots, P_{s-1} exhaust infinitely near base points of Λ such that P_i is an infinitely near point of P_{i-1} of order one for $1 \leq i \leq s-1$. For $1 \leq i \leq s$, let $\sigma_i: V_i \rightarrow V_{i-1}$ be the quadratic transformation of V_{i-1} with center at P_{i-1} , where $V_0 := V$, and let $\sigma = \sigma_1 \cdots \sigma_s$. Then σ factors ρ , i.e., $\rho = \sigma \cdot \bar{\rho}$. Let $E_i := (\sigma_{i+1} \cdots \sigma_s)'(\sigma_i^{-1}(P_{i-1}))$ for $1 \leq i < s$ and let $E_s := \sigma_s^{-1}(P_{s-1})$. Let $E_i := \bar{\rho}'(E_i')$ for $1 \leq i \leq s$. It is clear that $E_i' \cong E_i$ and $(E_i'^2) \geq (E_i^2)$ for $1 \leq i \leq s$, and that $(E_i^2) < -1$ for $1 \leq i < s$ and $(E_s^2) = -1$. Moreover, E_s is not contained in any member of $\tilde{\Lambda}$; indeed, if otherwise, $\tilde{\Lambda}$ would have yet a base point on E_s , which contradicts the choice of points P_1, \dots, P_{s-1} . The member \tilde{F} of $\tilde{\Lambda}$ corresponding to F of Λ

may contain some (not necessarily all) of E_1, \dots, E_{s-1} . After the above process made for every point of \mathfrak{B} and every sequence of infinitely near base points $\{P_0, P_1, \dots, P_{s-1}\}$ as above, we know that if we write $\tilde{F}=(n_1\tilde{C}_1+\dots+n_r\tilde{C}_r)+(m_1D_1+\dots+m_tD_t)$ with $\tilde{C}_i=\rho'(C_i)$ for $1\leq i\leq r$ then we have;

1° if $C_i\in S$ then $\tilde{C}_i\cong C_i$ and $(\tilde{C}_i^2)=(C_i^2)$,

2° if $C_i\notin S$ then $(\tilde{C}_i^2)\leq -2$,

3° $(D_i^2)\leq -2$ for $1\leq i\leq t$.

Then the assertions (4) and (5) follow from the assertions (5) and (6) of Lemma 1.1. Q.E.D.

1.3. Let V be a nonsingular projective surface containing an open subset U , which is isomorphic to A_k^2 . Since U is affine, the boundary set $V-U$ is connected and purely of codimension 1. Write $V-U=\bigcup_{i=1}^r C_i$, where C_i is an irreducible component. We assume that the boundary curve $V-U$ has only normal crossings as singularities. Then we have the following

Lemma (cf. Ramanujam [8]). *Let V and C_i 's be as above. Then the following assertions hold true:*

(1) *For every i , C_i is isomorphic to P_k^1 .*

(2) *For $i\neq j$, $(C_i\cdot C_j)=0$ or 1.*

(3) *For three distinct indices i, j and l , $C_i\cap C_j\cap C_l=\phi$.*

(4) *There is no circular chain (or loop) $\{C_{i_1}, \dots, C_{i_s}\}$ such that $(C_{i_j}\cdot C_{i_{j+1}})=1$ for $1\leq j<s$ and $(C_{i_s}\cdot C_{i_1})=1$.*

V is then called a *normal compactification* of A_k^2 . V is called a *minimal normal compactification* if the following additional condition is satisfied:

(5) *If an irreducible component, say C_1 , of $V-U$ is an exceptional curve of the first kind, then at least three other components of $V-U$ meet C_1 .*

1.4. Let V be a normal compactification of A_k^2 . Then the dual graph of the boundary curve $V-U$ is defined by assigning a vertex \circ to each irreducible component and by connecting two vertices by an edge (like $\circ-\circ$) if two corresponding components meet each other. The condition (4) of Lemma 1.3 says that the dual graph of $V-U$ is a *tree*. The dual graph is said to be *linear* if it is a tree and each vertex has at most two branches. Then we have the following.

Lemma (cf. Ramanujam [8]). *Let V be a minimal normal compactification of A_k^2 . Then the dual graph of the boundary curve is linear.*

The dual graphs of all possible minimal normal compactifications of A_k^2 were classified by Morrow [7]. An algebraic proof of the above lemma of Ramanujam and also of Morrow's result (even over the ground field of positive characteristic) was given by S. Mori [6]. However, we use this result only in

one place (cf. 3.11) in this article.

1.5. Let $k[x, y]$ be a polynomial ring in two variables x, y and let f be an irreducible polynomial of degree $d > 0$ in $k[x, y]$. Let $\Lambda(f)$ be the linear pencil on \mathbf{P}_k^2 defined by f . Then $\Lambda(f)$ has base points on the line l_∞ at infinity. Set $V_0 := \mathbf{P}_k^2$. Let $\varphi: \tilde{V} \rightarrow V_0$ be the shortest succession of quadratic transformations with centers at base points (including infinitely near base points) of $\Lambda(f)$ such that the proper transform $\tilde{\Lambda}$ of $\Lambda(f)$ by φ has no base points. Let $\tilde{\rho}: \tilde{V} \rightarrow \mathbf{P}_k^1$ be the surjective morphism defined by $\tilde{\Lambda}$. Then each irreducible (exceptional) curve arising in the process φ of quadratic transformations is either a quasi-section of $\tilde{\rho}$ or contained in a fiber of $\tilde{\rho}$. Let $\tilde{\Gamma}_1, \dots, \tilde{\Gamma}_p$ exhaust all irreducible (exceptional) curves arising in the process φ , which are quasi-sections of $\tilde{\rho}$. Then it is clear that every $\tilde{\Gamma}_i$ is isomorphic to \mathbf{P}_k^1 .

Note that \tilde{V} contains in a canonical way an open subset \tilde{U} which is isomorphic to \mathbf{A}_k^2 and that \tilde{V} is a normal compactification of $\tilde{U} \cong \mathbf{A}_k^2$. Let $\psi: \tilde{V} \rightarrow V$ be a contraction of all possible exceptional (contractible) components of $\tilde{V} - \tilde{U}$, which are contained in the fibers of $\tilde{\rho}$. Then V is a nonsingular projective surface containing an open subset U , which is isomorphic to \mathbf{A}_k^2 . However, V is not necessarily a normal compactification of \mathbf{A}_k^2 (cf. [5; Example 2.4.4, p. 122]). Moreover, there exists a surjective morphism $\rho: V \rightarrow \mathbf{P}_k^1$ such that $\tilde{\rho} = \rho \cdot \psi$. We set $\Gamma_i = \psi(\tilde{\Gamma}_i)$ for $1 \leq i \leq p$ and denote by S_∞ the fiber of ρ which corresponds to the member dl_∞ of $\Lambda(f)$. If we identify U with $\text{Spec}(k[x, y])$, there exists an inhomogeneous coordinate u of \mathbf{P}_k^1 such that the point $\rho(S_\infty)$ is given by $u = \infty$ and $S_\alpha \cap U = F_\alpha$ for each $\alpha \in k$, where S_α is the fiber of ρ lying over the point $u = \alpha$.

1.6. Assume that f is generically rational, i.e., general members of $\Lambda(f)$ are rational curves. Then, with the above notations, S_∞ is isomorphic to \mathbf{P}_k^1 . Let S_1, \dots, S_r be all fibers of ρ such that $F_i := S_i \cap U$ is a reducible curve for $1 \leq i \leq r$. Let m_i be the number of irreducible components of F_i for $1 \leq i \leq r$. On the other hand, $\Gamma_1, \dots, \Gamma_p$ exhaust all quasi-sections of ρ which are contained in $V - U$, while they are not necessarily nonsingular. Let δ_j be the degree of the morphism $\rho|_{\Gamma_j}: \Gamma_j \rightarrow \mathbf{P}_k^1$. Note that $\psi|_{\tilde{\Gamma}_j}: \tilde{\Gamma}_j \rightarrow \Gamma_j$ is the desingularization of Γ_j . For $1 \leq j \leq p$, let $\nu_j = \sum_Q (e_Q - 1)$, where e_Q is the ramification index of $\tilde{\rho}|_{\tilde{\Gamma}_j}$ at a point $Q \in \tilde{\Gamma}_j$ and the summation ranges over all points Q of $\tilde{\Gamma}_j$ such that $\tilde{\rho}(Q) \neq \rho(S_\infty)$. Then we have the following.

Lemma (cf. Saito [13], Suzuki [15]). *With the assumptions and the notations as above, we have:*

$$(1) \quad p+r-1 = \sum_{i=1}^r m_i; \quad (2) \quad n+1 = p + \sum_{j=1}^p \nu_j, \text{ where } n+1 \text{ is the number of}$$

places at infinity of a general member F_∞ of $\Lambda_0(f)$ (cf. Introduction.)

Proof. (1) We note that S_1, \dots, S_r are all singular fibers of ρ . Indeed, suppose that S is a singular fiber of ρ other than S_1, \dots, S_r . Then S has only one irreducible component C which meets U . Let μ be the multiplicity of C in S . Since $\mu C \cap U = S \cap U$ is defined by an equation $f = \alpha$ with some $\alpha \in k$, we have $f - \alpha = \lambda g^\mu$ for some irreducible polynomial $g \in k[x, y]$ and $\lambda \in k^*$. The assertion (6) of Lemma 1.1 implies that $\mu > 1$. Then f is reducible, which contradicts the choice of f . Let S be an irreducible fiber of ρ . Then we have

$$\chi(V) = \chi(S) \cdot \chi(\mathbf{P}_k^1) + \sum_{i=1}^r (\chi(S_i) - \chi(S)),$$

where $\chi(\)$ denotes the Euler number (cf. Šafarevič [11; p. 58]). Let m'_i be the number of irreducible components of S_i . Since the dual graph of S_i is a tree, it is easy to show that $\chi(S_i) = 2m'_i - (m'_i - 1) = m'_i + 1$. On the other hand, it is easy to see from construction of V that the Picard group $\text{Pic}(V)$ is a free abelian group of rank $1 + p + \sum_{i=1}^r (m'_i - m_i)$. Since V is rational, $\chi(V)$ is then equal to $2 + 1 + p + \sum_{i=1}^r (m'_i - m_i)$. Thus, from the above equality, we obtain

$$3 + p + \sum_{i=1}^r (m'_i - m_i) = 4 + \sum_{i=1}^r (m'_i - 1),$$

whence follows the first equality.

(2) For each j ($1 \leq j \leq p$), $S_\infty \cap \Gamma_j$ consists of a single point, which is also a one-place point. For, if otherwise, $\tilde{V} - \tilde{U}$ would contain a circular chain of irreducible components, which is a contradiction by the assertion of Lemma 1.3. Then there is only one point on $\tilde{\Gamma}_j$ lying over the point $\rho(S_\infty)$, where the ramification index of $\tilde{\rho}|_{\tilde{\Gamma}_j}$ equals δ_j . By Hurwitz's formula, we obtain $\delta_j - 1 = \nu_j$ for every j ($1 \leq j \leq p$). On the other hand, note that if S is a general fiber of ρ , S has $n + 1$ distinct points outside of $S \cap U$, among which δ_j points lie on Γ_j for each j ($1 \leq j \leq p$). Therefore, we have the second equality. Q.E.D.

1.7. Lemma. *Let f be a generically rational polynomial in $k[x, y]$. Then, with the notations as above, the following conditions are equivalent:*

- (1) $n = 0$;
- (2) $p = 1$;
- (3) $r = 0$, i.e., the curve F_∞ on A_k^2 defined by $f = \alpha$ is irreducible for every $\alpha \in k$;
- (4) f is sent to one of coordinates x, y by a biregular automorphism of $A_k^2 := \text{Spec}(k[x, y])$.

Proof. Note that $p > 0$. Then the implication (1) \Rightarrow (2) is clear. Suppose

$p=1$. Then $r=0$ by the first equality of Lemma 1.6, because $m_i \geq 2$ for every i . This implies the assertion (3) (cf. the proof of the first equality of Lemma 1.6). Suppose the condition (3) is satisfied. Then V is a relatively minimal rational ruled surface, whence V is a Hirzebruch surface $F_a = \text{Proj}(\mathcal{O}_{P^1} \otimes \mathcal{O}_{P^1}(a))$ with $a \geq 0$. Let M be the minimal section of F_a , i.e., M is the cross-section of ρ with $(M^2) = -a$; when $a=0$ we take a cross-section of ρ as M . Since $\text{Pic}(V)$ is a free abelian group generated by M and the fiber S_∞ of ρ , the unique quasi-section Γ of ρ contained in $V-U$ is linearly equivalent to a divisor of the form $\alpha M + \beta S_\infty$, where α and β are non-negative integers such that $\beta \geq \alpha a$. Since $U \cong \mathbb{A}_k^1$ and $V-U = \Gamma \cup S_\infty$, we know that Γ and S_∞ also generate $\text{Pic}(V)$. Then $\alpha=1$, and Γ is a cross-section. This implies that $S \cap U$ is isomorphic to \mathbb{A}_k^1 for every fiber S of ρ other than S_∞ . Hence the curve F_0 defined by $f=0$ is isomorphic to \mathbb{A}_k^1 . Now, the assertion (4) follows from Abhyankar-Moh's theorem (cf. Abhyankar-Moh [1], Miyanishi [4]). The implication (4) \Rightarrow (1) is clear. Q.E.D.

1.8. DEFINITION. *Let f be a generically rational polynomial in $k[x, y]$. f is said to be of simple type if the equality $p=n+1$ holds.*

The second equality of Lemma 1.6 shows that f is of simple type if and only if $\Gamma_1, \dots, \Gamma_p$ are all cross-sections of ρ .

Lemma. *Let f be a generically rational polynomial in $k[x, y]$ such that a general member of $\Lambda_0(f)$ has two places at infinity, i.e., $n=1$. Then f is of simple type. Moreover, $\Lambda_n(f)$ has only one reducible fiber, which has two irreducible components.*

Proof. By Lemma 1.7, we have $p \geq 2$. Then, by the second equality of Lemma 1.6, we must have $p=2$, whence f is of simple type. Then the first equality of Lemma 1.6 implies that $r=1$ and $m_1=2$. Q.E.D.

1.9. Let f be a generically rational polynomial in $k[x, y]$. We assume in this paragraph that f is of simple type, i.e., $p=n+1$. Then every $\Gamma_j (1 \leq j \leq p)$ is a nonsingular rational curve. We assume, furthermore, that $p \geq 2$, i.e., $n \geq 1$. Since U is isomorphic to \mathbb{A}_k^2 , two distinct Γ_j and Γ_l do not meet each other on any fiber S of ρ other than S_∞ ; indeed, if otherwise, $\tilde{V}-\tilde{U}$ would contain a circular chain of irreducible components. Note also that S_1, \dots, S_r exhaust all singular fibers of ρ .

Let Δ be an irreducible component of $S_i-S_i \cap U$ (if it exists at all) for some $i (1 \leq i \leq r)$. If there is a sequence of irreducible components $\{\Delta_1, \dots, \Delta_t\}$ of $S_i-S_i \cap U$ such that $\Delta_l = \Delta, \Delta_l \cap \Delta_{l+1} \neq \emptyset$ for $1 \leq l < t$ and $\Delta_l \cap \Gamma_j \neq \emptyset$ for some $j (1 \leq j \leq n+1)$, we say that Δ is connected to Γ_j . Since $V-U$ is a connected curve, every irreducible component Δ of $S_i-S_i \cap U$ is connected to some $\Gamma_j (1 \leq j \leq n+1)$, while Δ is not connected to two distinct Γ_j and Γ_l ; indeed, if

otherwise, $\tilde{V}-\tilde{U}$ would contain a circular chain of irreducible components. For each pair (i, j) with $1 \leq i \leq r$ and $1 \leq j \leq n+1$, let E_{ij} be the union of all irreducible components Δ of $S_i - S_i \cap U$, which are connected to Γ_j . Then E_{ij} is a connected curve if it is not empty.

Set $P_\infty := \rho(S_\infty)$ and $P_i := \rho(S_i)$ for $1 \leq i \leq r$. Then the points P_∞ and P_i 's are determined by $u = \infty$ and $u = c_i (1 \leq i \leq r)$, respectively, with respect to the inhomogeneous coordinate u of \mathbf{P}_k^1 (cf. the paragraph 1.5). Set $Z := \mathbf{P}_k^1 - \{P_\infty, P_1, \dots, P_r\}$. Then $\rho^{-1}(Z)$ with the morphism $\rho: \rho^{-1}(Z) \rightarrow Z$ is a trivial \mathbf{P}^1 -bundle over Z , i.e., there exists a Z -isomorphism $\eta: \rho^{-1}(Z) \xrightarrow{\sim} Z \times \mathbf{P}_k^1$. We may assume that there exist points Q_1, \dots, Q_{n+1} on \mathbf{P}_k^1 such that $\eta^{-1}(Z \times \{Q_j\}) = \Gamma_j \cap \rho^{-1}(Z)$ for $1 \leq j \leq n+1$. Now, choose an inhomogeneous coordinate v on \mathbf{P}_k^1 (=the fiber of ρ) such that Q_{n+1} is defined by $v = \infty$ and Q_j is defined by $v = d_j$ for $1 \leq j \leq n$; in the sequel, we set $Q_\infty := Q_{n+1}$ and $\Gamma_\infty := \Gamma_{n+1}$. Let $p_2: Z \times \mathbf{P}_k^1 \rightarrow \mathbf{P}_k^1$ be the second projection. Then the morphism $p_2 \circ \eta: \rho^{-1}(Z) \rightarrow \mathbf{P}_k^1$ (or equivalently saying, the inclusion of the subfield $k(v)$ into $k(x, y)$) defines a linear pencil L on V without fixed components such that $\Gamma_1, \dots, \Gamma_{n+1}$ are contained in distinct members $\Xi_1, \dots, \Xi_n, \Xi_\infty := \Xi_{n+1}$ of L , respectively.

For each $i (1 \leq i \leq r)$, let S_{i1}, \dots, S_{im_i} exhaust all irreducible components of S_i such that $S_{ia} \cap U \neq \emptyset$ for $1 \leq a \leq m_i$. Set $F_{ia} := S_{ia} \cap U$ and let f_{ia} be an irreducible polynomial in $k[x, y]$ such that F_{ia} is defined by $f_{ia} = 0$, where $1 \leq a \leq m_i$. Then, for each $i (1 \leq i \leq r)$, we have

$$u - c_i = f - c_i = \lambda_i \cdot \prod_{a=1}^{m_i} (f_{ia})^{\alpha_{ia}},$$

where $\lambda_i \in k^*$ and α_{ia} is a positive integer. Then we have:

1.9.1. $k[x, y, (\prod_{i=1}^r \prod_{a=1}^{m_i} f_{ia})^{-1}] = k[u, v, \prod_{i=1}^r (u - c_i)^{-1}, \prod_{j=1}^n (v - d_j)^{-1}].$

1.9.2. *If $n \geq 2$, the linear pencil L has no base points which lie outside S_∞ .*

Proof. Suppose that Q is a base point of L with $Q \notin S_\infty$. Then it is clear that $Q \in S_i$ for some $i (1 \leq i \leq r)$. Note that, for every $j (1 \leq j \leq n+1)$, the underlying curve of Ξ_j is contained in the union of Γ_j and the underlying curves of S_i 's $(1 \leq i \leq r)$. Since $n+1 \geq 3$, this implies that either there are at least two components of S_i intersecting one of the cross-sections Γ_j 's, or there are at least three components of S_i passing through the point Q , which contradicts the assertion (4) of Lemma 1.1.

1.9.3. *After a suitable modification^(*) of V with centers at points of S_∞ , we may*

(*) a succession of quadratic transformations with centers at points and contractions of exceptional curves of the first kind.

assume that L has no base points on S_∞ .

Proof. If S_∞ is not contained in any member of L then L has no base points on S_∞ . Suppose that L has a base point Q on S_∞ . Let $s:=i(\Xi, \Xi'; Q)$ for distinct general members Ξ and Ξ' of L . Set $Q_0:=Q$, and let Q_1, \dots, Q_{s-1} be infinitely near points of Q_0 , which lie on (the proper transforms of) Ξ , such that Q_i is an infinitely near point of Q_{i-1} of order one for $1 \leq i < s$. Let $\sigma_i: V_i \rightarrow V_{i-1}$ be the quadratic transformation of V_{i-1} with center at Q_{i-1} , where $V_0:=V$, and let $\sigma = \sigma_1 \cdots \sigma_s$. Let $E_i := (\sigma_{i+1} \cdots \sigma_s)'(\sigma_i^{-1}(Q_{i-1}))$ for $1 \leq i < s$, and let $E_s := \sigma_s^{-1}(Q_{s-1})$. Let $S'_\infty := \sigma'(S_\infty)$, let $\Gamma'_j := \sigma'(\Gamma_j)$ for $1 \leq j \leq n+1$, and let $L' := \sigma'(L)$. Then we have the following dual graph of $\sigma^{-1}(S_\infty)$:



Since $n+1 \geq 2$, some member $\Xi_j (1 \leq j \leq n+1)$ does not contain S_∞ . Moreover, since $(\Gamma_j \cdot S_\infty) = 1$, we know that Q_0, Q_1, \dots, Q_{s-1} exhaust all base points (including infinitely near base points) of L centered on S_∞ . Therefore, L' has no base points centered on S_∞ . Thus E_s is a cross-section of L' , and $S'_\infty \cup E_1 \cup \dots \cup E_{s-1}$ is contained in some member of L' . Let $\bar{\sigma}: V_s \rightarrow \bar{V}$ be the contraction of $S'_\infty, E_1, \dots, E_{s-1}$. Now, replace V by \bar{V} . Then the above modification $\bar{\sigma} \cdot \sigma^{-1}: V \rightarrow \bar{V}$ changes only S_∞ . Q.E.D.

1.9.4. Assume that $r \geq 2$. Then, for every $j (1 \leq j \leq n+1)$, at most one of E_{1j}, \dots, E_{rj} is a nonempty set.

Proof. Suppose that (at least) two of E_{1j}, \dots, E_{rj} , say E_{1j} and E_{2j} , are nonempty sets. By 1.9.3, we may assume that L has no base points on S_∞ . Then V is a normal compactification of U . Note that $(S_\infty^2) = 0$ and no exceptional components are contained in $\bigcup_{i=1}^r E_{ij}$. By contracting (possible) exceptional components in $V-U$, we would obtain a minimal normal compactification of \mathbf{A}_k^2 , for which the dual graph of the boundary curve is not linear. This contradicts Lemma 1.4. Q.E.D.

1.9.5. For distinct pairs (i, j) and (i', j') , we have $E_{ij} \cap E_{i'j'} = \emptyset$. Furthermore, we have $S_i - S_i \cap U = \bigcup_{j=1}^{n+1} E_{ij}$ for every $i (1 \leq i \leq r)$.

Proof. Clear from the construction and the above arguments.

1.9.6. Assume that $n \geq 2$. With V modified so that L has no base points (cf. 1.9.3), L defines a \mathbf{P}^1 -fibration $\tau: V \rightarrow \mathbf{P}_k^1$ such that S_∞ is a cross-section of τ .

1.9.7. Assume that $n \geq 1$. After the modification of V so that L has no base points on S_∞ , we call V a *standard compactification of A_k^2 with respect to* a generically rational polynomial f of simple type. Then we have the following

Lemma. *Let f be a generically rational polynomial of simple type in $k[x, y]$ with $n \geq 1$. Let V and \bar{V} be standard compactifications of $A_k^2 := \text{Spec}(k[x, y])$ with respect to f . Let $\rho: V \rightarrow P_k^1$ (or $\bar{\rho}: \bar{V} \rightarrow P_k^1$, resp.) be the P^1 -fibration of V (or \bar{V} , resp.) defined by f . Then there exists an isomorphism $\theta: V \rightarrow \bar{V}$ such that $\rho = \bar{\rho} \cdot \theta$.*

Proof. V (or \bar{V} , resp.) contains an open subset U (or \bar{U} , resp.) isomorphic to A_k^2 . The identity morphism $id.: U \xrightarrow{\sim} \bar{U}$ extends to a birational mapping $\theta: V \rightarrow \bar{V}$ such that $\rho = \bar{\rho} \cdot \theta$. We shall show that θ is an isomorphism. Let \mathfrak{B} and $\bar{\mathfrak{B}}$ be the generic fibers of ρ and $\bar{\rho}$, respectively. Then both \mathfrak{B} and $\bar{\mathfrak{B}}$ are complete normal models of the algebraic function field $k(x, y)$ of one variable over $\mathfrak{k} = k(f)$. Hence \mathfrak{B} and $\bar{\mathfrak{B}}$ are isomorphic to P_k^1 . Let $\Gamma_1, \dots, \Gamma_{n+1}$ (or $\bar{\Gamma}_1, \dots, \bar{\Gamma}_{n+1}$, resp.) be the cross-sections of ρ (or $\bar{\rho}$, resp.) contained in $V - U$ (or $\bar{V} - \bar{U}$, resp.). Since $\mathfrak{B} - \{\Gamma_1 \cap \mathfrak{B}, \dots, \Gamma_{n+1} \cap \mathfrak{B}\}$ is identified to $\bar{\mathfrak{B}} - \{\bar{\Gamma}_1 \cap \bar{\mathfrak{B}}, \dots, \bar{\Gamma}_{n+1} \cap \bar{\mathfrak{B}}\}$ under θ , \mathfrak{B} is isomorphic to $\bar{\mathfrak{B}}$ under θ , and we may assume that $\theta(\Gamma_j \cap \mathfrak{B}) = \bar{\Gamma}_j \cap \bar{\mathfrak{B}}$ for $1 \leq j \leq n+1$. Then $\theta: \Gamma_j \rightarrow \bar{\Gamma}_j$ is an isomorphism for $1 \leq j \leq n+1$. Let S_1, \dots, S_r (or $\bar{S}_1, \dots, \bar{S}_r$, resp.) be reducible fibers of ρ (or $\bar{\rho}$, resp.). Define \bar{E}_{ij} ($1 \leq i \leq r$ and $1 \leq j \leq n+1$) for \bar{V} in the same fashion as for V ; note that $\bigcup_{i=1}^r \bigcup_{j=1}^{n+1} \bar{E}_{ij}$ contains no exceptional curves of the first kind. Let S_∞ (or \bar{S}_∞ , resp.) be the fiber of ρ (or $\bar{\rho}$, resp.) contained in $V - U$ (or $\bar{V} - \bar{U}$, resp.). If θ is not biregular, $V - U$ contains an irreducible curve T which becomes an exceptional curve of the first kind after a succession of quadratic transformations with centers at points in $V - U$. Since every component in $\bigcup_{i=1}^r \bigcup_{j=1}^{n+1} E_{ij}$ has self-intersection multiplicity ≤ -2 , T must be S_∞ . However, if S_∞ is contracted after a sequence of quadratic transformations, $\bar{\Gamma}_1, \dots, \bar{\Gamma}_{n+1}$ meet each other at one point on \bar{S}_∞ , which is a contradiction. Hence θ is biregular. Similarly, θ^{-1} is biregular. Thus θ is an isomorphism such that $\rho = \bar{\rho} \cdot \theta$.

Q.E.D.

1.10. In the second section we use the following

Lemma (cf. Miyanishi [3,5]). *Let X be a nonsingular affine surface defined by an affine k -domain A . Assume that the following conditions are satisfied:*

- (1) *A is a unique factorization domain and $A^* = k^*$.*
- (2) *There exist nonsingular irreducible curves C_1 and C_2 on X such that $C_1 \cap C_2 = \{P\}$, and C_1 and C_2 intersect each other transversally at P .*
- (3) *C_1 (resp. C_2) has only one place at infinity.*

(4) Let a_2 be a prime element of A defining the curve C_2 . Then $a_2 - \alpha$ is a prime element of A for all $\alpha \in k$.

(5) There is a nonsingular projective surface V containing X as an open subset such that the closure \bar{C}_2 of C_2 in V is nonsingular and $(a_2)_0$ (=the zero part of the divisor of a_2) = \bar{C}_2 . Then X is isomorphic to A_k^2 , and the curves C_1 and C_2 are sent to the axes of a suitable coordinate system of A_k^2 .

1.11. Here, we recall a general result due to Russell [9; Cor. 3.7] on a generically rational polynomial.

Proposition. Let f be a generically rational polynomial in $k[x, y]$. Then there are at most two points (including infinitely near points) of the curve $f=0$ on the line at infinity l_∞ . In particular, the degree form of f has at most two distinct irreducible factors.

2. Generically rational polynomials with $n=1$

In this section, f is a generically rational, irreducible polynomial in $k[x, y]$ with $n=1$. Then f is of simple type and $\Lambda_0(f)$ has only one reducible member consisting of two irreducible components (cf. Lemma 1.8). Let $f=c_1$ be the reducible member of $\Lambda_0(f)$. We use the notations of the previous section with due modifications.

2.1. Lemma. With the notations of 1.9, we have one of the following cases:

(1) Case $F_{11} \cap F_{12} \neq \phi$. Then $(S_{11} \cdot S_{12})=1$ and $F_{1a} \cong A_k^1$ for $a=1, 2$.

(2) Case $F_{11} \cap F_{12} = \phi$. Then $(S_{11} \cdot S_{12})=0$, and one of F_{11} and F_{12} , say F_{11} , is isomorphic to A_k^1 and the other, say F_{12} , is isomorphic to $A_k^1 - (\text{one point})$.

Proof. (1) Suppose $F_{11} \cap F_{12} \neq \phi$. Then $(S_{11} \cdot S_{12})=1$ (cf. Lemma 1.1, (3)). Since $E_{11} \cap E_{12} = \phi$, we may assume that $(\Gamma_a \cup E_{1a}) \cap S_{1a} \neq \phi$ for $a=1, 2$. Since the dual graph of S_1 is a tree, it is easy to see that $F_{1a} \cong A_k^1$ for $a=1, 2$.

(2) Suppose $F_{11} \cap F_{12} = \phi$. Then one of S_{11} and S_{12} , say S_{12} , intersects both $\Gamma_1 \cup E_{11}$ and $\Gamma_2 \cup E_{12}$, and the other one, say S_{11} , intersects only one of $\Gamma_1 \cup E_{11}$ and $\Gamma_2 \cup E_{12}$. Then $(S_{11} \cdot S_{12})=0$, $F_{11} \cong A_k^1$ and $F_{12} \cong A_k^1 - (\text{one point})$. Q.E.D.

2.2. Lemma. With the notations of 1.9 and after a suitable change of coordinates of $k[x, y]$, we have one of the following cases:

(1) $f_{11}=x$ and $f_{12}=y$.

(2) $f_{11}=x$ and $f_{12}=x^l y + P(x)$, where $l > 0$ and $P(x) \in k[x]$ with $\deg P(x) < l$ and $P(0) \neq 0$.

Proof. (1) Suppose $F_{11} \cap F_{12} \neq \phi$. Then, by virtue of Lemmas 1.7 and 1.10, we may change coordinates of $k[x, y]$ so that $f_{11}=x$ and $f_{12}=y$.

(2) Suppose $F_{11} \cap F_{12} = \phi$. Then, by virtue of Abhyankar-Moh's Theorem

(cf. Lemma 1.7), we may assume that $f_{11}=x$. Taking account of 1.9.1, we have only to prove the following assertion:

Let $g \in k[x, y]$ be an irreducible polynomial such that $g \notin k[x]$ and $g(0, y) \neq 0$. Assume that we have the following identification of rings,

$$k[x, y, x^{-1}, g^{-1}] = k[u, v, u^{-1}, v^{-1}].$$

Then g is of the form:

$$g = \lambda(x^l y + P(x)), \text{ where } \lambda \in k^*, l > 0 \text{ and } P(x) \in k[x] \\ \text{with } \deg P(x) < l \text{ and } P(0) \neq 0.$$

Indeed, by comparison of the unit groups of both rings, we have: $u \sim x^\alpha g^\beta$ and $v \sim x^\gamma g^\delta$, where $\alpha, \beta, \gamma, \delta \in \mathbf{Z}$ such that $\alpha\delta - \beta\gamma = \pm 1$. This implies that g satisfies

$$x^a g^b y = G(x, g), \text{ where } G(x, g) \in k[x, g].$$

If $b > 0$ then we have a relation: $gh(x, y) \in k[x]$ with $h(x, y) \in k[x, y]$, which implies $g \in k[x]$, a contradiction. Hence $b = 0$. Write

$$g = A_0(x)y^M + \dots + A_M(x) \text{ and } G(x, g) = B_0(x)g^N + \dots + B_N(x),$$

where $A_i(x), B_j(x) \in k[x]$ ($0 \leq i \leq M$ and $0 \leq j \leq N$) and $A_0(x)B_0(x) \neq 0$. Then we have

$$x^a y = B_0(x)A_0(x)^N y^{NM} + \dots,$$

whence $M = 1$ and $A_0(x) \sim x^l$. Therefore, g is written in the stated form after replacing y by $y + C(x)$. Q.E.D.

2.3. Theorem (cf. Saito [12; p. 332], Sugie [14]). *Let f be a generically rational, irreducible polynomial in $k[x, y]$ with $n = 1$. Then, after a suitable change of coordinates, f is reduced to either one of the following two forms:*

- (1) $f \sim x^\alpha y^\beta + 1$, where $\alpha > 0, \beta > 0$ and $(\alpha, \beta) = 1$.
- (2) $f \sim x^\alpha (x^l y + P(x))^\beta + 1$, where $\alpha, \beta, l > 0, (\alpha, \beta) = 1$ and $P(x) \in k[x]$ with $\deg P(x) < l$ and $P(0) \neq 0$.

Proof. Clear by Lemma 2.2.

2.4. Proposition. *Assume that the curve C_0 on \mathbf{P}_k^2 defined by $f = 0$ intersects l_∞ in only one point P . Let $d_1 := \text{mult}_P C_0$. Then $d_1 < d = \text{the degree of } f$. Moreover, there exists a birational automorphism ρ of \mathbf{P}_k^2 such that ρ induces a biregular automorphism on $A_k^2 := \mathbf{P}_k^2 - l_\infty$ and that the proper transform C'_0 of C_0 by ρ intersects l_∞ in two points with $(C'_0 \cdot l_\infty) \leq d_1$.*

The existence of an automorphism ρ such that $\rho|_{A_k^2}$ is a biregular automorphism and that the proper transform $\rho'(C_0)$ intersects l_∞ in two points

follows from the above theorem. However, the above proposition is proved by a constructive method depending on Lemma 1.2, which allows us to determine more explicitly an automorphism ρ with the stated properties. For more details, the readers will be referred to Miyanishi [5; Chap. II, §6].

2.5. It is clear that the integers α and β in Theorem 2.3 are the multiplicities of components S_{11} and S_{12} of the fiber S_1 , respectively, in the standard compactification of A_k^2 with respect to f . On the other hand, Lemma 1.9.7 implies that the standard compactification of A_k^2 with respect to f is obtained by assuming that f is written in the form stated in Theorem 2.3. By a straightforward computation, we know that the dual graphs of E_{11} and E_{12} are linear. For details, see [5; *ibid.*].

3. Generically rational polynomials of simple type with $n > 1$

In this section, f is a generically rational, irreducible polynomial of simple type with $n > 1$. In the paragraphs 3.1~3.4, we consider the case $r=1$; in the paragraphs 3.5~3.8, we consider the case where $r \geq 2$ and $m_i < n$ for every i ($1 \leq i \leq r$); in the paragraphs 3.9~3.12, we consider the case where $r=2$, $m_1 = n$ and $m_2=2$.

3.1. We shall consider the case $r=1$. For convenience's sake, we simplify the notations as follows: $\Sigma := S_1$; $\Sigma_a := S_{1a}$, $F_a := F_{1a}$, $f_a := f_{1a}$, $\alpha_a := \alpha_{1a}$ for $1 \leq a \leq m_1$, where $m_1 = p = n+1$; $E_j := E_{1j}$ for $1 \leq j \leq n+1$; we may assume that $c_1=1$ (cf. 1.9.) Note that the pencil L on the standard compactification V of A_k^2 with respect to f has no base points; hence L defines a P^1 -fibration $\tau: V \rightarrow P_k^1$. Let Ξ be a general fiber of τ . Since $(S_\infty \cdot \Xi) = 1$, we have $(\Sigma \cdot \Xi) = 1$. This implies that there exists an irreducible component Δ of Σ such that $(\Delta \cdot \Xi) = 1$, i.e., Δ is a cross-section of τ , and the other components of Σ are contained in the fibers Ξ_1, \dots, Ξ_{n+1} and possibly one other fiber of L , as will be seen below. First, we shall prove the following:

Lemma. *With the assumptions and the notations as above, we have one of the following cases:*

(1) $\Delta \cap U \neq \emptyset$. We may assume that $\Delta = \Sigma_{n+1}$. Then $\Xi \cap U \cong A_k^1$ and $F_a \cong A_k^1$ for $1 \leq a \leq n$; Σ_a ($1 \leq a \leq n$) belongs to one and only one of Ξ_1, \dots, Ξ_{n+1} , while none of Ξ_1, \dots, Ξ_{n+1} contains two of Σ_a 's; thus, after a suitable change of inhomogeneous coordinate v of P_k^1 (=the fiber of ρ) and a suitable change of indices, we may assume that Σ_a belongs to Ξ_a for $1 \leq a \leq n$; $F_{n+1} = \Delta - (\Delta \cap \Xi_{n+1}) \cup \bigcup_{j=1}^n ((\Gamma_j \cup E_j) \cap \Delta)$, where $\Delta \cong P_k^1$; $\alpha_{n+1} = 1$.

(2) $\Delta \cap U = \emptyset$. We may assume that Δ is a component of E_{n+1} . Then $\Xi \cap U \cong A_k^1$; Σ_a ($1 \leq a \leq n+1$) belongs to one and only one of Ξ_1, \dots, Ξ_n ; none of

Σ_a 's belongs to Ξ_{n+1} ; thus, after a suitable change of indices, we may assume that Σ_a belongs to Ξ_a for $1 \leq a \leq n$ and Σ_{n+1} belongs to either one of Ξ_j 's ($1 \leq j \leq n$), say Ξ_1 , or none of them; $F_1 \cong A_k^1$ for $2 \leq a \leq n$ and $F_{n+1} \cong A_k^1$. Moreover, (i) $F_1 \cong A_k^1$ either if Σ_{n+1} is a component of Ξ_1 and $\Sigma_1 \cap \Sigma_{n+1} = \emptyset$ or if Σ_{n+1} belongs to none of Ξ_j 's ($1 \leq j \leq n$), and (ii) $F_1 \cong A_k^1$ if Σ_{n+1} is a component of Ξ_1 and $\Sigma_1 \cap \Sigma_{n+1} \neq \emptyset$.

Proof. (1) Suppose $\Delta \cap U \neq \emptyset$. As in the statement, we may assume that $\Delta = \Sigma_{n+1}$. Then $\Xi \cap U = \Xi - \Xi \cap S_\infty$, whence $\Sigma \cap U \cong A_k^1$. Since $(\Xi \cdot \Sigma_a) = 0$ for $1 \leq a \leq n$, Σ_a belongs to the fibers of L . Hence we may assume that one of Ξ_1, \dots, Ξ_{n+1} , say Ξ_{n+1} , contains none of Σ_a 's ($1 \leq a \leq n$). Then the inhomogeneous coordinate v of P_k^1 (=the fiber of ρ) is an element of $k[x, y]$; indeed, the polar divisor of v on V has no irreducible components meeting the open set $U = \text{Spec}(k[x, y])$. Since $\Xi \cap U \cong A_k^1$, we may assume, after a suitable change of coordinates in $k[x, y]$, that $v = x$ (cf. Abhyankar-Moh's Theorem). Then the curve $v = \alpha$ on U is isomorphic to A_k^1 for every $\alpha \in k$. This implies that Σ_a ($1 \leq a \leq n$) belongs to one and only one of Ξ_1, \dots, Ξ_n , and that none of Ξ_j 's ($1 \leq j \leq n$) contains two of Σ_a 's. Note that $\Gamma_j \cup E_j$ is contained in Ξ_j for $1 \leq j \leq n+1$. It is now clear that $F_{n+1} = \Delta - (\Delta \cap \Xi_{n+1}) \cup \bigcup_{j=1}^n ((\Gamma_j \cup E_j) \cap \Delta)$, where $(\Gamma_j \cup E_j) \cap \Delta$ is a single point if it is nonempty and $\Delta \cong P_k^1$. Since $(\Xi \cdot \Sigma) = 1$, the multiplicity α_{n+1} of Δ in Σ equals 1.

(2) Suppose $\Delta \cap U = \emptyset$. After a suitable change of the inhomogeneous coordinate v , we may assume that Δ is a component of E_{n+1} . We have $(\Xi \cdot \Sigma_a) = 0$ for $1 \leq a \leq n+1$ and $(\Xi_j \cdot \Delta) = 1$ for $1 \leq j \leq n$. Hence $\Gamma_j \cup E_j$, which is contained in Ξ_j , is connected to Δ by means of one or more of Σ_a 's for $1 \leq j \leq n$. This implies that Ξ_{n+1} contains at most one of Σ_a 's and Ξ_j ($1 \leq j \leq n$) contains at most two of Σ_a 's. Suppose that Ξ_{n+1} contains one of Σ_a 's, say Σ_{n+1} . Then Ξ_j ($1 \leq j \leq n$) contains one and only one of Σ_a 's ($1 \leq a \leq n$). After a suitable change of indices, we may assume that Ξ_j contains Σ_j for $1 \leq j \leq n$. Then it is easy to see that $v - d_j \sim f_j^{\beta_j} f_{n+1}^{-\gamma}$ for $1 \leq j \leq n$, where $\beta_j > 0$ ($1 \leq j \leq n$) and $\gamma > 0$; γ is independent of j . Since $u - 1 \sim f_1^{\alpha_1} \cdots f_{n+1}^{\alpha_{n+1}}$, the identification of rings in 1.9.1 implies (by computing the unit groups of the rings on both hand sides) that the following matrix A is unimodular,

$$A = \begin{pmatrix} \alpha_1 & \cdots & \alpha_n & \alpha_{n+1} \\ \beta_1 & & & -\gamma \\ & \ddots & & \vdots \\ 0 & & \beta_n & -\gamma \end{pmatrix},$$

However, $\det A = (-1)^n \{ \beta_1 \cdots \beta_n \alpha_{n+1} + \gamma \sum_{i=1}^n \beta_1 \cdots \beta_{i-1} \alpha_i \beta_{i+1} \cdots \beta_n \}$, whence A is not unimodular because $n \geq 2$. Therefore, Ξ_{n+1} contains none of Σ_a 's ($1 \leq a \leq n+1$). Then, after a suitable change of indices, we may assume that Σ_j is used

to connect $\Gamma_j \cup E_j$ to Δ for $1 \leq j \leq n$. The remaining component Σ_{n+1} belongs to one of Ξ_j 's ($1 \leq j \leq n$), say Ξ_1 , or none of them. It is then clear that $F_a \cong \mathbf{A}_k^1$ for $2 \leq a \leq n$. If either Σ_{n+1} belongs to none of Ξ_j 's ($1 \leq j \leq n$) or Σ_{n+1} is a component of Ξ_1 and $\Sigma_1 \cap \Sigma_{n+1} = \phi$, then $F_1 \cong \mathbf{A}_k^1$. If Σ_{n+1} is a component of Ξ_1 and $\Sigma_1 \cap \Sigma_{n+1} \neq \phi$, then $F_1 \cong \mathbf{A}_k^1$. In any case, $F_{n+1} \cong \mathbf{A}_k^1$. Since $\Xi \cap U = \Xi - (\Xi \cap S_\infty) \cup (\Xi \cap \Delta)$, it is isomorphic to \mathbf{A}_k^1 . Q.E.D.

3.2. Lemma. *With the assumptions and the notations of Lemma 3.1, after a suitable change of coordinates of $k[x, y]$, we have one of the following four cases:*

(1) Case $\Delta \cap U \neq \phi$:

$$f_j \sim x - d_j \quad (1 \leq j \leq n) \quad \text{and}$$

$$f_{n+1} \sim y \cdot \prod_{j=1}^n (x - d_j)^{\varepsilon_j} + P(x), \quad \varepsilon_j \geq 0, \quad P(x) \in k[x],$$

where $\varepsilon_j = 0$ whenever $(\Gamma_j \cup E_j) \cap \Delta = \phi$, and where $\varepsilon_j > 0$ and $P(d_j) \neq 0$ whenever $(\Gamma_j \cup E_j) \cap \Delta \neq \phi$. Moreover, $\alpha_{n+1} = 1$.

(2) Case $\Delta \cap U = \phi$ and $\Sigma_{n+1} \not\subset |\Xi_j|^{(*)}$ for $1 \leq j \leq n$:

$$f_{n+1} \sim x \quad \text{and} \quad f_j \sim x^l(x^t y + P(x)) - d_j \quad \text{for} \quad 1 \leq j \leq n,$$

where $l > 0, t \geq 0$ and $P(x) \in k[x]; \deg P(x) < t$ and $P(0) \neq 0$ if $t > 0$, and $P(x) = 0$ if $t = 0$. Moreover, $\alpha_{n+1} = 1$.

(3) Case $\Delta \cap U = \phi, \Sigma_{n+1} \subset |\Xi_1|$ and $\Sigma_1 \cap \Sigma_{n+1} \neq \phi$:

$$f_{n+1} \sim x, \quad f_1 \sim y \quad \text{and} \quad f_j \sim x^l y + d_1 - d_j \quad \text{for} \quad 2 \leq j \leq n,$$

where $l > 0$. Moreover, $\alpha_{n+1} - \alpha_1 l = \pm 1$.

(4) Case $\Delta \cap U = \phi, \Sigma_{n+1} \subset |\Xi_1|$ and $\Sigma_1 \cap \Sigma_{n+1} = \phi$:

$$f_{n+1} \sim x, \quad f_1 \sim x^t y + P(x) \quad \text{and} \quad f_j \sim x^l(x^t y + P(x)) + d_1 - d_j \quad \text{for} \quad 2 \leq j \leq n,$$

where $l, t > 0$ and $P(x) \in k[x]$ with $\deg P(x) < t$ and $P(0) \neq 0$. Moreover, $\alpha_{n+1} - l\alpha_1 = \pm 1$.

Proof. (1) Since none of Σ_a 's ($1 \leq a \leq n+1$) is contained in Ξ_{n+1} we have $v \in k[x, y]$, and since $\Xi \cap U \cong \mathbf{A}_k^1$, we may assume that $v = x$ (cf. Abhyankar-Moh's Theorem). Then, each fiber Ξ_j ($1 \leq j \leq n$) of L (or τ) has only one component Σ_j which intersects U , and $\Sigma_j \cap U = F_j$. Since Ξ_j corresponds to the value $v = d_j$, we have $f_j \sim x - d_j$. Now, the curve F_{n+1} , which is defined by $f_{n+1} = 0$, is written in the form:

(*) For an effective divisor D on V , we denote by $|D|$ the underlying reduced curve of D .

$$f_{n+1} = P_0(x)y^N + \cdots + P_N(x),$$

where $P_0(x), \dots, P_N(x) \in k[x]$ and $P_0(x) \neq 0$. Since F_{n+1} meets the curve $x = \alpha$ transversally in a single point for $\alpha \in k$ such that $\alpha \neq d_j$ ($1 \leq j \leq n$) or $\alpha = d_j$ with $F_j \cap F_{n+1} \neq \phi$. Hence we know that $N=1$ and $P_0(x) \sim \prod_{j=1}^n (x-d_j)^{\varepsilon_j}$, where $\varepsilon_j=0$ if $F_j \cap F_{n+1} \neq \phi$, i.e., $(\Gamma_j \cup E_j) \cap \Delta = \phi$. If $F_j \cap F_{n+1} = \phi$, i.e., $(\Gamma_j \cup E_j) \cap \Delta \neq \phi$, we have $\varepsilon_j > 0$ and $P_1(d_j) \neq 0$. Thus, we obtain the stated form for f_{n+1} . On the other hand, we have:

$$u-1 \sim f_1^{\alpha_1} \cdots f_{n+1}^{\alpha_{n+1}} \quad \text{and} \quad v-d_j \sim f_j^{\beta_j} \quad \text{with} \quad \beta_j > 0 \quad \text{for} \quad 1 \leq j \leq n.$$

The identification of rings in 1.9.1 implies that the following matrix A is unimodular,

$$A = \begin{pmatrix} \alpha_1 & \cdots & \alpha_n & \alpha_{n+1} \\ \beta_1 & & & 0 \\ & \ddots & & \vdots \\ 0 & & \ddots & \beta_n \\ & & & 0 \end{pmatrix}.$$

Since $\det A = (-1)^n \alpha_{n+1} \beta_1 \cdots \beta_n$, we have $\beta_1 = \cdots = \beta_n = \alpha_{n+1} = 1$.

(2) Since $\Sigma_{n+1} \not\subset |\Xi_j|$ for $1 \leq j \leq n$, there is one more reducible fiber Ξ_0 , other than Ξ_j 's ($1 \leq j \leq n+1$), such that

$$\Xi_0 = T + l\Sigma_{n+1} + Z,$$

where $(T \cdot S_\infty) = 1$, $(T \cdot \Sigma_{n+1}) \leq 1$, $|Z| \subset E_{n+1}$ if $Z > 0$, and $l > 0$. Then T_0 ($:= T \cap U \cong \mathbf{A}_k^1$ if $(T \cdot \Sigma_{n+1}) = 1$ and $T_0 \cong \mathbf{A}_k^1$ if $(T \cdot \Sigma_{n+1}) = 0$). Since $\Xi \cap U \cong \mathbf{A}_k^1$, we may assume, by virtue of Theorem 2.3 that $f_{n+1} = x$ and T_0 is defined by $y = 0$ (if $T \cong \mathbf{A}_k^1$) or $x^t y + P(x) = 0$ (if $T \cong \mathbf{A}_k^1$), where $t > 0$ and $P(x) \in k[x]$ with $\deg P(x) < t$ and $P(0) \neq 0$. Moreover, we may assume that the fiber Ξ_0 corresponds to the value $v = 0$, where $v \in k[x, y]$ because $|\Xi_{n+1}| \cap U = \phi$. Since Ξ_j corresponds to the value $v = d_j$ and $F_j = |\Xi_j| \cap U$ for $1 \leq j \leq n$, we have $f_j \sim x^t(x^t y + P(x)) - d_j$. We obtain $\alpha_{n+1} = 1$ by the same argument as in the case (1).

(3) The fiber Ξ_1 is, then, written in the form:

$$\Xi_1 = \Sigma_1 + l\Sigma_{n+1} + Z,$$

where $(\Sigma_1 \cdot \Delta) = 1$, $(\Sigma_1 \cdot \Sigma_{n+1}) = 1$, $|Z| = E_1 \cup \Gamma_1$, and $l > 0$. Since $F_1 \cong F_{n+1} \cong \mathbf{A}_k^1$, we may assume that $f_{n+1} = x$ and $f_1 = y$ (cf. Lemma 1.10). Since $v \in k[x, y]$ and the fiber Ξ_1 corresponds to the value $v = d_1$, we may assume that $v = x^t y + d_1$. Then we have $f_j \sim v - d_j = x^t y + d_1 - d_j$ for $2 \leq j \leq n$, because $\Sigma_j \subset |\Xi_j|$ and $(\Sigma_j \cdot \Delta) = 1$. On the other hand, we have:

$$u-1 \sim f_1^{\alpha_1} \cdots f_{n+1}^{\alpha_{n+1}}, \quad v-d_1 \sim f_1 f_{n+1}^l$$

and $v-d_j \sim f_j$ for $2 \leq j \leq n$.

By the same reasoning as in the case (1), we have the following unimodular matrix

$$A = \begin{pmatrix} \alpha_1 \cdots \alpha_n & \alpha_{n+1} \\ 1 & l \\ & 1 & 0 \\ & & \ddots & \\ 0 & & & \ddots & 0 \\ & & & & 1 \end{pmatrix}.$$

Since $\det A = (-1)^n(\alpha_{n+1} - \alpha_1 l)$, we have $\alpha_{n+1} - \alpha_1 l = \pm 1$.

(4) The fiber Ξ_1 is, then, written in the form:

$$\Xi_1 = \Sigma_1 + l\Sigma_{n+1} + Z,$$

where $(\Xi_1 \cdot \Delta) = 1$, $(\Sigma_1 \cdot \Sigma_{n+1}) = 0$, $\Sigma_{n+1} \cap |Z| \neq \emptyset$, $|Z| = E_1 \cup \Gamma_1$ and $l > 0$. Since $F_{n+1} \cong A_k^1$, we may assume that $f_{n+1} = x$. Since $v \in k[x, y]$ and $\Xi \cap U \cong A_k^1$, v is a generically rational polynomial with $n = 1$, and the curve $v = d_1$ is a reducible member of $\Lambda_0(v)$ with two mutually non-intersecting components. By virtue of Lemma 2.2, we may set

$$f_1 = x^t y + P(x), \quad t > 0, \quad P(x) \in k[x]$$

with $\deg P(x) < t$ and $P(0) \neq 0$.

Then we may assume that $v - d_1 = f_1 f_{n+1}^l = x^l(x^t y + P(x))$. Hence $f_j \sim v - d_j = x^l(x^t y + P(x)) + d_1 - d_j$, for $2 \leq j \leq n$. We obtain $\alpha_{n+1} - l\alpha_1 = \pm 1$ by the same argument as in the case (3). Q.E.D.

3.3. Theorem. *Let f be a generically rational, irreducible polynomial of simple type in $k[x, y]$ with $n > 1$ and only one reducible member in $\Lambda_0(f)$. Then, after a suitable change of coordinates in $k[x, y]$, f is reduced to one of the following four polynomials:*

(1) $f \sim \left(\prod_{j=1}^n (x - d_j)^{\alpha_j} \right) \cdot (y \cdot \prod_{j=1}^n (x - d_j)^{\varepsilon_j} + P(x)) + 1,$

where d_1, \dots, d_n are mutually distinct elements in k and $P(x) \in k[x]$; $\alpha_j > 0$ and $\varepsilon_j \geq 0$ for $1 \leq j \leq n$; $P(d_j) \neq 0$ if $\varepsilon_j > 0$.

(2) $f \sim x \cdot \prod_{j=1}^n (x^l(x^t y + P(x)) - d_j)^{\alpha_j} + 1,$

where $l > 0$, $t \geq 0$ and $P(x) \in k[x]$; $\deg P(x) < t$ and $P(0) \neq 0$ if $t > 0$ and $P(x) = 0$ if $t = 0$; α_j 's and d_j 's are as in the case (1).

(3) $f \sim x^\beta y^{\alpha_1} \cdot \prod_{j=2}^n (x^l y - d_j)^{\alpha_j} + 1,$

where d_2, \dots, d_n are mutually distinct elements in k^* ; $\beta > 0$, $l > 0$ and $\alpha_j > 0$ for

$$1 \leq j \leq n; \beta - \alpha_1 l = \pm 1.$$

$$(4) \quad f \sim x^\beta \cdot (x^t y + P(x))^{\alpha_1} \cdot \prod_{j=2}^n (x^l (x^t y + P(x)) - d_j)^{\alpha_j + 1},$$

where $t > 0$ and $P(x) \in k[x]$ with $\deg P(x) < t$ and $P(0) \neq 0$; β, l, α_j 's and d_j 's are as in the case (3).

Proof. Follows easily from Lemma 3.2.

3.4. By comparison of the unit groups of two rings, which are connected to each other by the identification of rings as in 1.9.1, we can prove the following:

Proposition. Let f_1, \dots, f_{n+1} ($n \geq 2$) be mutually distinct irreducible polynomials in $k[x, y]$. Assume that we have the identification of rings,

$$k[x, y, (f_1 \cdots f_{n+1})^{-1}] = k[u, v, u^{-1}, (\prod_{j=1}^n (v - d_j))^{-1}],$$

where $u = f_1^{\alpha_1} \cdots f_{n+1}^{\alpha_{n+1}}$ with $\alpha_i > 0$ ($1 \leq i \leq n+1$), and d_1, \dots, d_n are mutually distinct elements of k . Then, after a suitable change of indices and a suitable change of variables u and v , we are reduced to the following case:

$$v - d_1 \sim f_1^{\beta_1} f_2^{\beta_2} \quad \text{and} \quad v - d_j \sim f_{j+1} \quad \text{for} \quad 2 \leq j \leq n,$$

where $\beta_1, \beta_2 \geq 0$ and $\alpha_1 \beta_2 - \alpha_2 \beta_1 = \pm 1$.

3.5. Next, we shall consider the case where $r \geq 2$ and $m_i < n$ for every i ($1 \leq i \leq r$). Here, we retain the notations in 1.9. Since $m_i \geq 2$, we have $n \geq 3$.

Lemma. With the assumptions and the notations as above, we have:

(1) For every i ($1 \leq i \leq r$), $E_{ij} \cap \Xi = \emptyset$ for $1 \leq j \leq n+1$, where Ξ is a general fiber of the \mathbf{P}^1 -fibration $\tau: V \rightarrow \mathbf{P}_k^1$ (cf. 1.9.6), and there exists an irreducible component Δ_i among S_{ia} 's ($1 \leq a \leq m_i$) such that $(\Delta_i \cdot \Xi) = 1$, i.e., Δ_i is a cross-section of τ ; moreover, $E_{ij} \cup \Gamma_j \subset |\Xi_j|$ for $1 \leq i \leq r$ and $1 \leq j \leq n+1$.

(2) $\Xi \cap U \cong \mathbf{A}_k^1$.

(3) For every i ($1 \leq i \leq r$), $F_{ia} \cong \mathbf{A}_k^1$ if $S_{ia} \neq \Delta_i$ ($1 \leq a \leq m_i$) and $\Delta_i \cap U = \Delta_i - \bigcup_{j=1}^{n+1} ((\Gamma_j \cup E_{ij}) \cap \Delta_i)$.

(4) One of Ξ_j 's ($1 \leq j \leq n+1$), say Ξ_{n+1} , contains none of S_{ia} 's ($1 \leq i \leq r$ and $1 \leq a \leq m_i$). Then, every Ξ_j ($1 \leq j \leq n$) contains one and only one of S_{ia} 's ($1 \leq i \leq r$ and $1 \leq a \leq m_i$) with multiplicity 1. After a suitable change of indices, we may assume that, for every i ($1 \leq i \leq r$), we have:

$$\Delta_i = S_{i1} \quad \text{and} \quad S_{ia} \subset |\Xi_b|,$$

where $b := \gamma(i, a) = (m_1 - 1) + \cdots + (m_{i-1} - 1) + (a - 1)$ for $2 \leq a \leq m_i$.

Proof. (1) Suppose that, for some i ($1 \leq i \leq r$), S_i contains an irreducible component Δ such that $\Delta \cap U = \phi$ and $(\Delta \cdot \Xi) = 1$, i.e., Δ is a cross-section of τ . Suppose, for convenience's sake, that $\Delta \subset E_{i,n+1}$. Then $(\Gamma_j \cup E_{ij}) \cap \Xi = \phi$ for $1 \leq j \leq n$, whence $\Gamma_j \cup E_{ij} \subset |\Xi_j|$. Since Δ is a cross-section of τ , every $\Gamma_j \cup E_{ij}$ ($1 \leq j \leq n$) is connected to Δ by one or more components of S_{ia} 's ($1 \leq a \leq m_i$). However, since $m_i < n$, this is impossible. This implies that there exists a component Δ_i among S_{ia} 's ($1 \leq a \leq m_i$) such that $(\Delta_i \cdot \Xi) = 1$, i.e., Δ_i is a cross-section. Then $(\Gamma_j \cup E_{ij}) \cap \Xi = \phi$ for $1 \leq j \leq n+1$, whence $(\Gamma_j \cup E_{ij}) \subset |\Xi_j|$.

(2) It is now clear that $\Xi \cap U = \Xi - (\Xi \cap S_\infty) \cong A^k$.

(3) Suppose that $S_{ia} \neq \Delta_i$ ($1 \leq a \leq m_i$). Since $S_{ia} \cap \Xi = \phi$, S_{ia} is a component of some Ξ_j ($1 \leq j \leq n+1$). If $S_{ia} \cap \Delta_i \neq \phi$, then $S_{ia} \cap (\Gamma_j \cup E_{ij}) \neq \phi$; indeed, if otherwise, $S_{ia} \subset U$, which is a contradiction. Hence $F_{ia} \cong A^k$. If $S_{ia} \cap \Delta_i = \phi$ then $E_{ij} \neq \phi$ and $E_{ij} \cap S_{ia} \neq \phi$, whence $F_{ia} \cong A^k$. It is now clear that $\Delta_i \cap U = \Delta_i - \bigcup_{j=1}^{n+1} ((\Gamma_j \cup E_{ij}) \cap \Delta_i)$.

(4) Since Δ_i ($1 \leq i \leq r$) is a cross-section of τ , Δ_i is not contained in any one of Ξ_j 's ($1 \leq j \leq n$). The remaining components of S_{ia} 's ($1 \leq i \leq r$ and $1 \leq a \leq m_i$) are contained in the union of Ξ_1, \dots, Ξ_{n+1} . Since $\sum_{i=1}^r (m_i - 1) = n$ (cf. Lemma 1.6), one of Ξ_j 's, say Ξ_{n+1} , contains none of S_{ia} 's. Then the inhomogeneous coordinate v is an element of $k[x, y]$. Since $\Xi \cap U \cong A^k$, we may assume that $v = x$ (cf. Abhyankar-Moh's Theorem). Then $|\Xi_j| \cap U$, which corresponds to the value $v = d_j$, is irreducible for $1 \leq j \leq n$. Hence, every Ξ_j ($1 \leq j \leq n$) contains one and only one of S_{ia} 's. The remaining assertion is easy to prove.

Q.E.D

3.6. Lemma. *With the assumptions and the notations as in Lemma 3.5, we have:*

- (1) $f_{ia} \sim x - d_b$ with $b = \gamma(i, a)$, for $1 \leq i \leq r$ and $2 \leq a \leq m_i$.
- (2) $f_{i1} \sim y \cdot \prod_{j=1}^n (x - d_j)^{\varepsilon_j} + P_i(x)$ with $\varepsilon_j \geq 0$ and $P_i(x) \in k[x]$, where $\varepsilon_j = 0$ if $(\Gamma_j \cup E_{ij}) \cap \Delta_i = \phi$, and $\varepsilon_j > 0$ and $P_i(d_j) \neq 0$ if $(\Gamma_j \cup E_{ij}) \cap \Delta_i \neq \phi$.
- (3) $\alpha_{i1} = 1$ for every i ($1 \leq i \leq r$).

Proof. (1) As in the proof of Lemma 3.5, we may assume that $v = x$. Then, since $S_{ia} \subset |\Xi_b|$ with $b = \gamma(i, a)$ and $S_{ia} = |\Xi_b| \cap U$, we have $f_{ia} \sim x - d_b$.

(2) f_{i1} can be determined by the same argument as used in the proof of Lemma 3.2, the assertion (1). We only note that Δ_i meets (or does not meet, resp.) the unique component among S_{ia} 's contained in Ξ_j if and only if $(\Gamma_j \cup E_{ij}) \cap \Delta_i = \phi$ (or $(\Gamma_j \cup E_{ij}) \cap \Delta_i \neq \phi$).

(3) We have:

$$u - c_i \sim \prod_{a=1}^{m_i} f_{ia}^{\alpha_{ia}} \quad \text{for } 1 \leq i \leq r$$

and $v - d_j \sim f_{ia}$ with $j = \gamma(i, a)$, for $1 \leq j \leq n$.

Let $A_i (1 \leq i \leq r)$ and A be the following square matrices of size m_i and $n+r$, respectively;

$$A_i = \begin{pmatrix} \alpha_{i1} & \cdots & \alpha_{im_i} \\ 1 & & \\ & \ddots & 0 \\ 0 & & \ddots & \\ & & & 1 \end{pmatrix} \quad \text{and} \quad A = \begin{pmatrix} A_1 & & & 0 \\ & \ddots & & \\ & & \ddots & \\ 0 & & & A_r \end{pmatrix}.$$

The identification of rings in 1.9.1 implies that A is a unimodular matrix. Hence $\alpha_{i1}=1$ for $1 \leq i \leq r$. Q.E.D.

3.7. Theorem. *Let f be a generically rational, irreducible polynomial of simple type in $k[x, y]$ with $n > 1$, $r \geq 2$ and $n > m_i$ for every $i (1 \leq i \leq r)$; indeed, $n \geq 3$. Then, after a suitable change of coordinates in $k[x, y]$, f has one of the following presentations:*

$$f \sim \left(\prod_{a=2}^{m_i} (x - d_{e(a)})^{\alpha_{ia}} \right) \cdot (y \cdot \prod_{j=1}^n (x - d_j)^{\varepsilon_j} + P_i(x)) + c_i \quad \text{for } 1 \leq i \leq r,$$

where d_1, \dots, d_n are mutually distinct elements of k , $c_i \in k^*$; $\alpha_{ia} > 0$, $\varepsilon_j \geq 0$ and $e(a) := \gamma(i, a)$ (cf. Lemma 3.5); $P_i(d_j) \neq 0$ if $\varepsilon_j > 0$; $\varepsilon_j > 0$ if $j \neq e(a)$ for all $a (1 \leq a \leq m_i)$.

Proof. Follows easily from Lemmas 3.5 and 3.6.

3.8. Finally, we shall consider the case where $r \geq 2$ and $m_i = n$ for some $i (1 \leq i \leq r)$. We may assume that $m_1 = n$. Lemma 1.6 then implies $r = 2$, $m_1 = n$ and $m_2 = 2$. We consider first the case $n \geq 3$; the case $n = 2$ will be treated in the paragraphs 3.11 and 3.12. The \mathbf{P}^1 -fibration $\rho: V \rightarrow \mathbf{P}_k^1$ has two singular fibers S_1 and S_2 , where $S_1 \cap U$ has n irreducible components and $S_2 \cap U$ has 2 irreducible components. Here we retain the notations in 1.9. Let $\Delta_i (i = 1, 2)$ be the irreducible component in S_i such that $(\Delta_i \cdot \Xi) = 1$, where Δ_2 must be one of S_{21} and S_{22} , say S_{21} , (cf. the proof of the assertion (1) in Lemma 3.5).

Lemma. *Assume that $n \geq 3$. With the assumptions and the notations as above, we have one of the following two cases:*

(I) *Case $\Delta_1 \cap U \neq \emptyset$. Then we have the same situation as stated in Lemma 3.5 with $r = 2$.*

(II) *Case $\Delta_1 \cap U = \emptyset$. We may assume that $\Delta_1 \subset E_{1, n+1}$. Then the following assertions hold true:*

(1) *For every $j (1 \leq j \leq n)$, $\Gamma_j \cup E_{1j} \subset |\Xi_j|$; $S_{1a} (1 \leq a \leq n)$ belongs to one and only one of Ξ_j 's ($1 \leq j \leq n$); after a change of indices, we may assume that S_{1a} is contained in Ξ_a with multiplicity 1 for $1 \leq a \leq n$; $F_{1a} \cong \mathbf{A}_*^1$ for $1 \leq a \leq n$.*

(2) *$\Xi \cap U \cong \mathbf{A}_*^1$, where Ξ is a general fiber of the \mathbf{P}^1 -fibration $\tau: V \rightarrow \mathbf{P}_k^1$.*

(3) *S_{22} belongs to one of Ξ_j 's ($1 \leq j \leq n+1$), and $S_{22} \not\subset |\Xi_{n+1}|$; we may*

assume that $S_{22} \subset |\Xi_1|$; $F_{22} \cong A_k^1$ and $F_{21} = S_{21} - \bigcup_{j=1}^{n+1} (\Gamma_j \cup E_{2j}) \cap S_{21}$.

Proof. (I) If $\Delta_1 \cap U \neq \phi$, we have only to follow the arguments in Lemma 3.5.

(II) Assume that $\Delta_1 \cap U = \phi$. We may assume that $\Delta_1 \subset E_{1,n+1}$. Since $E_{1j} \cap \Xi = \phi$, we have $\Gamma_j \cup E_{1j} \subset |\Xi_j|$ for $1 \leq j \leq n$. Since $(\Gamma_j \cup E_{1j}) \cap \Delta_1 = \phi$, $\Gamma_j \cup E_{1j}$ is connected to Δ_1 by one and only one of S_{1a} 's for $1 \leq j \leq n$. After a suitable change of indices a , we may assume that $S_{1a} \subset |\Xi_a|$. Then it is clear that S_{1a} is a component of Ξ_a with multiplicity 1 and that $F_{1a} \cong A_k^1$ for $1 \leq a \leq n$. This proves the assertion (1).

(2) Since $\Xi \cap U = \Xi - (\Xi \cap S_\infty) \cup (\Xi \cap \Delta_1)$, $\Xi \cap U \cong A_k^1$.

(3) Suppose that S_{22} belongs to none of Ξ_j 's ($1 \leq j \leq n+1$). Then $S_{22} \subset U$, which is a contradiction. Hence S_{22} belongs to one of Ξ_j 's ($1 \leq j \leq n+1$). Suppose that $S_{22} \subset |\Xi_{n+1}|$. Then, in the identification of rings in 1.9.1, we have:

$$u - c_1 \sim \prod_{a=1}^n f_{1a}^{\alpha_{1a}}, \quad u - c_1 \sim f_{21}^{\alpha_{21}} f_{22}^{\alpha_{22}}$$

$$v - d_j \sim f_{1j} f_{22}^\beta \quad \text{for } 1 \leq j \leq n,$$

where $\beta > 0$, β is independent of j . By comparison of the unit groups of the rings of both hand sides in 1.9.1, we know that the following matrix A is unimodular,

$$A = \begin{pmatrix} \alpha_{11} \cdots \alpha_{1n} & 0 & 0 \\ 0 & \cdots & 0 & \alpha_{21} & \alpha_{22} \\ 1 & & & 0 & -\beta \\ & \ddots & & \vdots & \vdots \\ 0 & & \ddots & \vdots & \vdots \\ & & & 1 & 0 & -\beta \end{pmatrix}.$$

Since $\det A = -\beta \alpha_{21} \cdot (\sum_{i=1}^n \alpha_{1i})$ and $n \geq 3$, A is not a unimodular matrix, which is a contradiction. Hence $S_{22} \not\subset |\Xi_{n+1}|$. Then we may assume that $S_{22} \subset |\Xi_1|$. Then $F_{22} \cong A_k^1$ (cf. the proof of the assertion (3) of Lemma 3.5), and $F_{21} = S_{21} - \bigcup_{j=1}^{n+1} (\Gamma_j \cup E_{2j}) \cap S_{21}$. Q.E.D.

3.9. Lemma. *With the assumptions and the notations as in Lemma 3.8, assume that $\Delta_1 \cap U = \phi$. Then, after a suitable change of coordinates in $k[x, y]$, we have:*

- (1) $f_{22} \sim x$, $f_{11} \sim x^l y + P(x)$, and $f_{1j} \sim x(x^l y + P(x)) + d_1 - d_j$ for $2 \leq j \leq n$, where $l > 0$, $P(x) \in k[x]$, $\deg P(x) < l$ and $P(0) \neq 0$.
- (2) $\alpha_{11} = \alpha_{21} = 1$.

Proof. Since $F_{22} \cong A_k^1$, we may assume that $f_{22} = x$ (cf. Abhyankar-Moh's Theorem). Since $|\Xi_{n+1}| \cap U = \phi$, the inhomogeneous coordinate v is an element of $k[x, y]$. Since $\Xi \cap U \cong A_k^1$, v is a generically rational polynomial with $n=1$,

and the curve $v = d_1$ is the unique reducible member of $\Lambda_0(v)$ with two disjoint components F_{22} and F_{11} . Since $(S_{11} \cdot \Delta_1) = 1$, the fiber Ξ_1 is of the form:

$$\Xi_1 = S_{11} + tS_{22} + Z \quad \text{with} \quad |Z| = \Gamma_1 \cup E_{11}.$$

On the other hand, since $F_{11} \cap F_{22} = \phi$, f_{11} is written in the form:

$$f_{11} = x'y + P(x),$$

where $l > 0$ and $P(x) \in k[x]$ with $\deg P(x) < l$ and $P(0) \neq 0$. Then we have:

$$\begin{aligned} v - d_j &\sim x^t(x'y + P(x)) + d_1 - d_j \quad \text{for} \quad 1 \leq j \leq n, \\ v - d_j &\sim f_{1j} \quad \text{for} \quad 2 \leq j \leq n. \end{aligned}$$

Since $u - c_1 \sim \prod_{a=1}^n f_{1a}^{\alpha_a}$ and $u - c_2 \sim f_{21}^{\alpha_{21}} f_{22}^{\alpha_{22}}$, we have the following unimodular matrix,

$$A = \begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} & 0 & 0 \\ 0 & \cdots & 0 & \alpha_{21} & \alpha_{22} \\ 1 & & 0 & 0 & t \\ & \ddots & & & \\ 0 & & & & 1 \end{pmatrix}.$$

Since $\det A = \alpha_{11} \alpha_{21} t = 1$, we have $\alpha_{11} = \alpha_{21} = t = 1$. Q.E.D.

3.10. Theorem. *Let f be a generically rational, irreducible polynomial of simple type in $k[x, y]$ with $n \geq 3$, $r = 2$, $m_1 = n$ and $m_2 = 2$. Then, after a suitable change of coordinates in $k[x, y]$, f is reduced to a polynomial of one of the following types:*

(I) *f is a polynomial of the type given in Theorem 3.7, where $r = 2$, $m_1 = n$ and $m_2 = 2$;*

(II) $f \sim (x'y + P(x)) \cdot \prod_{j=2}^n (x(x'y + P(x)) - d_j)^{\alpha_j} + c,$

where $l > 0$, $\alpha_j > 0$ ($2 \leq j \leq n$), $P(x) \in k[x]$ with $\deg P(x) < l$ and $P(0) \neq 0$, $c \in k^*$, and d_j 's ($2 \leq j \leq n$) are mutually distinct elements in k^* .

Proof. Follows from Lemmas 3.8 and 3.9.

3.11. We shall consider the remaining case: $n = 2$ and $m_1 = m_2 = 2$. As in 3.8, let Δ_i be an irreducible component of S_i such that $(\Delta_i \cdot \Xi) = 1$, where $i = 1, 2$. We shall prove the following:

Lemma. *With the assumptions and the notations as above, we have either $\Delta_1 \cup U \neq \phi$ or $\Delta_2 \cap U \neq \phi$.*

Proof. Suppose that $\Delta_i \cap U = \phi$ for $i = 1, 2$. We may assume that $\Delta_1 \subset E_{13}$ and $\Gamma_i \cup E_{1i} \cup S_{1i} \subset |\Xi_i|$ for $i = 1, 2$, (cf. the proof of Lemma 3.8); the com-

ponent S_{1_i} has multiplicity 1 in Ξ_i for $i=1, 2$. Suppose $\Delta_2 \subset E_{23}$. Then $E_{13} \neq \phi$ and $E_{23} \neq \phi$, which contradicts the property 1.9.4; this is the only one place where we have to depend on Lemma 1.4. Therefore, $\Delta_2 \not\subset E_{23}$. We may assume that $\Delta_2 \subset E_{22}$, $\Gamma_1 \cup E_{21} \cup S_{21} \subset |\Xi_1|$ and $\Gamma_3 \cup E_{23} \cup S_{22} \subset |\Xi_3|$. Then we have: $u - c_i \sim f_{i1}^{\alpha_{i1}} f_{i2}^{\alpha_{i2}}$ for $i=1, 2$, $v - d_1 \sim f_{11} f_{21} f_{22}^{-\gamma}$ and $v - d_2 \sim f_{12} f_{22}^{-\gamma}$, where $\gamma > 0$. Hence we have the following unimodular matrix,

$$A = \begin{pmatrix} \alpha_{11} & \alpha_{12} & 0 & 0 \\ 0 & 0 & \alpha_{21} & \alpha_{22} \\ 1 & 0 & 1 & -\gamma \\ 0 & 1 & 0 & -\gamma \end{pmatrix}.$$

Since $\det A = -\alpha_{11}\alpha_{22} - \gamma(\alpha_{11}\alpha_{21} + \alpha_{12}\alpha_{21}) < -1$, this is a contradiction. Q.E.D.

Therefore, we have two cases: $\Delta_i \cap U \neq \phi$ for $i=1, 2$; $\Delta_1 \cap U = \phi$ and $\Delta_2 \cap U \neq \phi$. The case $\Delta_1 \cap U \neq \phi$ and $\Delta_2 \cap U = \phi$ is reduced to the second case. In both cases, we are reduced to the same situation as in Lemma 3.8 with $r=2$ and $m_i=2$ ($i=1, 2$). Therefore, we have:

3.12. Theorem. *Theorem 3.10 is valid in the case $n=2$.*

3.13. According to Russell [10], a polynomial f in $k[x, y]$ is said to be a *good* field generator if there exists a polynomial g in $k[x, y]$ such that $k(x, y) = k(f, g)$; otherwise, f is said to be a *bad* field generator. The observations in §§2, 3 imply the following:

Theorem. *Let f be a generically rational polynomial of simple type in $k[x, y]$. Then there exists a generically rational polynomial g with at most two places at infinity such that $k(x, y) = k(f, g)$. In particular, f is a good field generator.*

We note that a field generator is not necessarily good. An example of a bad field generator was given in [10].

References

[1] S.S. Abhyankar and T.T. Moh: *Embeddings of the line in the plane*, J. Reine Angew. Math. **276** (1975), 148-166.
 [2] M.H. Gizatullin: *On affine surfaces that can be completed by a nonsingular rational curve*, Izv. Akad. Nauk SSSR, Ser. Mat. **34** (1970), 778-802; Math. USSR-Izvestija **4** (1970), 787-810.
 [3] M. Miyanishi: *An algebraic characterization of the affine plane*, J. Math. Kyoto Univ. **15** (1975), 169-184.
 [4] M. Miyanishi: *Analytic irreducibility of certain curves on a nonsingular affine rational surface*, Proc. Internat. Sympos. on Algebraic Geometry (Kyoto), Kinokuniya, Tokyo, 1977, 575-587.
 [5] M. Miyanishi: *Lectures on curves on rational and unirational surfaces*, Tata

- Institute of Fundamental Research, Bombay, 1978.
- [6] S. Mori: *Embeddings of A^2 into rational ruled surfaces*, Kōkyuroku **183** (1973), Res. Inst. Math. Analysis, Kyoto Univ., (in Japanese).
 - [7] J.A. Morrow: *Minimal normal compactifications of C^2* , Rice Univ. Studies, 1973, 97–112.
 - [8] C.P. Ramanujam: *A topological characterization of the affine plane as an algebraic variety*, Ann. of Math. **94** (1971), 69–88.
 - [9] P. Russell: *Field generators in two variables*, J. Math. Kyoto Univ. **15** (1975), 555–571.
 - [10] P. Russell: *Good and bad field generators*, J. Math. Kyoto Univ. **17** (1977), 319–331.
 - [11] I.R. Šafarevič et al.: *Algebraic surfaces*, Proc. Steklov Inst. Math. **75** (1965).
 - [12] H. Saito: *Fonctions entières qui se réduisent à certains polynômes*. I, Osaka J. Math. **9** (1972), 293–332.
 - [13] H. Saito: *Fonctions entières qui se réduisent à certains polynômes*. II, Osaka J. Math. **14** (1977), 649–674.
 - [14] T. Sugie: *Certain affine plane curves with two places at infinity*, Thesis for Master's degree, Osaka Univ., 1978, (in Japanese).
 - [15] M. Suzuki: *Propriétés topologiques des polynômes de deux variables complexes et automorphismes algébriques de l'espace C^2* , J. Math. Soc. Japan **26** (1974), 241–257.

Masayoshi Miyanishi
Department of Mathematics
Osaka University
Toyonaka, Osaka 560
Japan

Tohru Sugie
Department of Mathematics
Kyoto University
Kyoto, 606
Japan