

INJECTIVE PAIRS IN PERFECT RINGS

MITSUO HOSHINO and TAKESHI SUMIOKA

(Received April 21, 1997)

Throughout this note, rings are associative rings with identity and modules are unitary modules. Sometimes, we use the notation ${}_A X$ (resp. X_A) to signify that the module X considered is a left (resp. right) A -module. For each pair of subsets X and M of a ring A , we set $\ell_X(M) = \{a \in X \mid aM = 0\}$ and $r_M(X) = \{a \in M \mid Xa = 0\}$.

Following Baba and Oshiro [1], we call a pair (eA, Af) of a right ideal eA and a left ideal Af in a ring A an i -pair if (a) e and f are local idempotents; (b) eA_A and ${}_A Af$ have essential socles; and (c) $\text{soc}(eA_A) \cong fA/fJ$ and $\text{soc}({}_A Af) \cong Ae/Je$, where J is the Jacobson radical of A .

Generalizing a result of Fuller [3], Baba and Oshiro [1] showed that for a local idempotent e in a semiprimary ring A , eA_A is injective if and only if there exists a local idempotent f in A such that (eA, Af) is an i -pair in A and $r_{Af}(\ell_{eA}(M)) = M$ for every submodule M of Af_{fAf} , and that for an i -pair (eA, Af) in a semiprimary ring A the following are equivalent: (1) eAe is artinian; (2) Af_{fAf} is artinian; and (3) both eA_A and ${}_A Af$ are injective.

Our aim is to extend the results mentioned above to perfect rings. Following Harada [4], we call a module L_A M -simple-injective if for any submodule N of M_A every $\theta : N_A \rightarrow L_A$ with $\text{Im } \theta$ simple can be extended to some $\phi : M_A \rightarrow L_A$. For a local idempotent e in a left perfect ring A , we will show that eA_A is A -simple-injective if and only if there exists a local idempotent f in A such that (eA, Af) is an i -pair in A and $r_{Af}(\ell_{eA}(M)) = M$ for every submodule M of Af_{fAf} , and that eA_A is injective if it is A -simple-injective and has finite Loewy length. We will show also that for an i -pair (eA, Af) in a left perfect ring A the following are equivalent: (1) eAe is artinian; (2) Af_{fAf} is artinian; and (3) both eA_A and ${}_A Af$ are injective.

1. Localization and injective objects

Let \mathcal{A} and \mathcal{B} be abelian categories, $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{A}$ covariant functors, and $\varepsilon : \mathbf{1}_{\mathcal{A}} \rightarrow GF$ and $\delta : FG \rightarrow \mathbf{1}_{\mathcal{B}}$ homomorphisms of functors, where $\mathbf{1}_{\mathcal{A}} : \mathcal{A} \rightarrow \mathcal{A}$ and $\mathbf{1}_{\mathcal{B}} : \mathcal{B} \rightarrow \mathcal{B}$ are identity functors. We assume the conditions: (a) $\delta_F \circ F\varepsilon = \text{id}_F$; (b) $G\delta \circ \varepsilon_G = \text{id}_G$; (c) F is exact; and (d) δ is an isomorphism.

REMARK 1. (1) By the conditions (a) and (b), for each pair of $X \in \text{Ob}(\mathcal{A})$ and

$M \in \text{Ob}(\mathcal{B})$ we have a natural isomorphism

$$\theta_{X,M} : \text{Hom}_{\mathcal{B}}(FX, M) \rightarrow \text{Hom}_{\mathcal{A}}(X, GM), \beta \mapsto G\beta \circ \varepsilon_X$$

with $\theta_{X,M}^{-1}(\alpha) = \delta_M \circ F\alpha$ for $\alpha \in \text{Hom}_{\mathcal{A}}(X, GM)$. Namely, G is a right adjoint of F . In particular, G is left exact.

(2) By the conditions (a), (b) and (d), $G : \mathcal{B} \rightarrow \mathcal{A}$ is fully faithful.

(3) By the conditions (a) and (d), $F\varepsilon : F \rightarrow FGF$ is an isomorphism with $F\varepsilon^{-1} = \delta_F$.

(4) By the conditions (b) and (d), $\varepsilon_G : G \rightarrow GFG$ is an isomorphism with $\varepsilon_G^{-1} = G\delta$.

Though the following lemmas are well known and more or less obvious, we include proofs for completeness.

Lemma 1.1. *Let $X \in \text{Ob}(\mathcal{A})$ be simple with $FX \neq 0$. Then $FX \in \text{Ob}(\mathcal{B})$ is simple.*

Proof. Let $\beta : FX \rightarrow M$ be a nonzero morphism in \mathcal{B} . We claim β monic. Note that $\beta = \delta_M \circ F(G\beta \circ \varepsilon_X)$. Thus $G\beta \circ \varepsilon_X : X \rightarrow GM$ is nonzero and monic, so is $\beta = \delta_M \circ F(G\beta \circ \varepsilon_X)$. \square

Lemma 1.2. *Let $\mu : Y \rightarrow X$ be an essential monomorphism in \mathcal{A} with ε_Y monic. Then $F\mu : FY \rightarrow FX$ is an essential monomorphism in \mathcal{B} .*

Proof. Let $\beta : FX \rightarrow M$ be a morphism in \mathcal{B} with $\beta \circ F\mu$ monic. We claim β monic. Since $(G\beta \circ \varepsilon_X) \circ \mu = G\beta \circ GF\mu \circ \varepsilon_Y = G(\beta \circ F\mu) \circ \varepsilon_Y$ is monic, $G\beta \circ \varepsilon_X$ is monic and so is $\beta = \delta_M \circ F(G\beta \circ \varepsilon_X)$. \square

Lemma 1.3. *Let $X \in \text{Ob}(\mathcal{A})$ be injective with ε_X monic. Then $\varepsilon_X : X \rightarrow GFX$ is an isomorphism and $FX \in \text{Ob}(\mathcal{B})$ is injective.*

Proof. Since $F\varepsilon_X$ is an isomorphism, $F(\text{Cok } \varepsilon_X) \cong \text{Cok } F\varepsilon_X = 0$ and $\text{Hom}_{\mathcal{A}}(\text{Cok } \varepsilon_X, GFX) \cong \text{Hom}_{\mathcal{B}}(F(\text{Cok } \varepsilon_X), FX) = 0$. Thus, since $\varepsilon_X : X \rightarrow GFX$ is a split monomorphism, $\text{Cok } \varepsilon_X = 0$. Hence for each $M \in \text{Ob}(\mathcal{B})$ we have a natural isomorphism

$$\eta_M : \text{Hom}_{\mathcal{B}}(M, FX) \rightarrow \text{Hom}_{\mathcal{A}}(GM, X), \beta \mapsto \varepsilon_X^{-1} \circ G\beta.$$

Let $\nu : N \rightarrow M$ be a monomorphism in \mathcal{B} . Since $G\nu$ is monic, $\text{Hom}_{\mathcal{A}}(G\nu, X)$ is epic and so is $\text{Hom}_{\mathcal{B}}(\nu, FX) = \eta_N^{-1} \circ \text{Hom}_{\mathcal{A}}(G\nu, X) \circ \eta_M$. \square

REMARK 2. (1) An object $M \in \text{Ob}(\mathcal{B})$ is injective if and only if so is $GM \in$

$\text{Ob}(\mathcal{A})$.

(2) The canonical monomorphism $\text{Im} \varepsilon_X \rightarrow GFX$ is an essential monomorphism for every $X \in \text{Ob}(\mathcal{A})$ with $FX \neq 0$.

(3) If $\nu : N \rightarrow M$ is an essential monomorphism in \mathcal{B} , so is $G\nu : GN \rightarrow GM$.

(4) For $X \in \text{Ob}(\mathcal{A})$ with ε_X monic, a monomorphism $\mu : Y \rightarrow X$ in \mathcal{A} is an essential monomorphism if and only if so is $F\mu : FY \rightarrow FX$.

2. Injective pairs

Throughout the rest of this note, A stands for a ring with Jacobson radical J . For an i -pair (eA, Af) in A , we denote by $\mathcal{A}_\ell(eA, Af)$ the lattice of submodules X of ${}_{eAe}eA$ with $\ell_{eA}(r_{Af}(X)) = X$ and by $\mathcal{A}_r(eA, Af)$ the lattice of submodules M of Af_fAf with $r_{Af}(\ell_{eA}(M)) = M$.

REMARK 3. Let (eA, Af) be an i -pair in A . Let X be a submodule of ${}_{eAe}eA$. Then $Xr_{Af}(X) = 0$ implies $X \subset \ell_{eA}(r_{Af}(X))$ and thus $r_{Af}(\ell_{eA}(r_{Af}(X))) \subset r_{Af}(X)$. Also, $\ell_{eA}(r_{Af}(X))r_{Af}(X) = 0$ implies $r_{Af}(X) \subset r_{Af}(\ell_{eA}(r_{Af}(X)))$. Thus $r_{Af}(X) \in \mathcal{A}_r(eA, Af)$. Similarly, $\ell_{eA}(M) \in \mathcal{A}_\ell(eA, Af)$ for every submodule M of Af_fAf . It follows that $\mathcal{A}_\ell(eA, Af)$ is anti-isomorphic to $\mathcal{A}_r(eA, Af)$.

The following lemmas have been established in [5], [3], [1], [8], [6] and so on. However, for the benefit of the reader, we provide direct proofs.

Lemma 2.1. *Let $e, f \in A$ be idempotents and assume $\ell_{eA}(Af) = 0 = r_{Af}(eA)$. Then the following hold.*

- (1) *For a two-sided ideal I of A , $eI = 0$ if and only if $If = 0$.*
- (2) *$\ell_{eA}(I) = \ell_{eA}(If)$ for every right ideal I of A .*
- (3) *$r_{Af}(I) = r_{Af}(eI)$ for every left ideal I of A .*

Proof. (1) Assume $eI = 0$. Then $eAIf = eIf = 0$ and $If \subset r_{Af}(eA) = 0$. By symmetry, $If = 0$ implies $eI = 0$.

(2) Since $If \subset I$, $\ell_{eA}(I) \subset \ell_{eA}(If)$. For any $x \in \ell_{eA}(If)$, since $xIAf = xIf = 0$, $xI \subset \ell_{eA}(Af) = 0$ and $x \in \ell_{eA}(I)$. Thus $\ell_{eA}(If) \subset \ell_{eA}(I)$.

(3) Similar to (2). □

Lemma 2.2. *Let (eA, Af) be an i -pair in A . Then the following hold.*

- (1) *$\ell_{eA}(Af) = 0 = r_{Af}(eA)$.*
- (2) *eAf_fAf and ${}_{eAe}eAf$ have simple essential socles and $\text{soc}(eA_A)f = \text{soc}(eAf_fAf) = \text{soc}({}_{eAe}eAf) = e(\text{soc}({}_A Af))$.*

Proof. (1) For any $0 \neq x \in eA$, since $\text{soc}(eA_A) \subset xA$, $0 \neq \text{soc}(eA_A)f \subset xAf$ and $x \notin \ell_{eA}(Af)$. Thus $\ell_{eA}(Af) = 0$. Similarly $r_{Af}(eA) = 0$.

(2) Since by Lemma 1.1 $\text{soc}(eA_A)f_fAf_f$ and ${}_{eAe}e(\text{soc}({}_A Af_f))$ are simple, and since by Lemma 1.2 $\text{soc}(eA_A)f_fAf_f \subset eAf_fAf_f$ and ${}_{eAe}e(\text{soc}({}_A Af_f)) \subset {}_{eAe}eAf_f$ are essential extensions, the assertion follows. \square

Lemma 2.3. *Let (eA, Af) be an i -pair in A . Then for any $n \geq 1$ $eJ^n = 0$ if and only if $J^n f = 0$, so that eA_A and ${}_A Af$ have the same Loewy length.*

Proof. By Lemmas 2.2(1) and 2.1(1). \square

Lemma 2.4. *Let (eA, Af) be an i -pair in A . Let N, M be submodules of Af_fAf_f with $N \subset M$ and M/N simple. Assume $N \in \mathcal{A}_r(eA, Af)$. Then the following hold.*

- (1) ${}_{eAe}e\ell_{eA}(N)/\ell_{eA}(M)$ is simple.
- (2) $M \in \mathcal{A}_r(eA, Af)$.

Proof. (1) Let $a \in M$ with $a \notin N$. Then $M = N + afAf_f$. Also, since $M \neq N = r_{Af}(\ell_{eA}(N))$, $\ell_{eA}(M) \subset \ell_{eA}(N)$ with $\ell_{eA}(N)/\ell_{eA}(M) \neq 0$. Since $0 \neq \ell_{eA}(N)M = \ell_{eA}(N)afAf_f$ and $\ell_{eA}(N)afJf = 0$, $\ell_{eA}(N)afAf_f = \text{soc}(eAf_fAf_f)$. Thus by Lemma 2.2(2) $\ell_{eA}(N)a = \text{soc}({}_{eAe}eAf_f)$ and, since $\ell_{eA}(M)a = 0$, ${}_{eAe}e\ell_{eA}(N)/\ell_{eA}(M) \cong \text{soc}({}_{eAe}eAf_f)$.

(2) Since $\ell_{eA}(M) \subset \ell_{eA}(N) \subset {}_{eAe}eA$ with $\ell_{eA}(M) \in \mathcal{A}_\ell(eA, Af)$ and $\ell_{eA}(N)/\ell_{eA}(M)$ simple, we can apply the part (1) to conclude that $r_{Af}(\ell_{eA}(M))/r_{Af}(\ell_{eA}(N))$ is simple. Thus $r_{Af}(\ell_{eA}(N)) = N \subset M \subset r_{Af}(\ell_{eA}(M))$ with both $r_{Af}(\ell_{eA}(M))/r_{Af}(\ell_{eA}(N))$ and M/N simple, so that $M = r_{Af}(\ell_{eA}(M))$. \square

Lemma 2.5. *Let (eA, Af) be an i -pair in A . Then $M \in \mathcal{A}_r(eA, Af)$ for every submodule M of Af_fAf_f of finite composition length.*

Proof. Lemma 2.4(2) together with Lemma 2.2(1) enables us to make use of induction on the composition length. \square

Lemma 2.6. *Let (eA, Af) be an i -pair in A . Then ${}_{eAe}eA$ and Af_fAf_f have the same composition length.*

Proof. By symmetry, we may assume Af_fAf_f has finite composition length. Let $0 = M_0 \subset M_1 \subset \dots \subset M_n = Af$ be a composition series of Af_fAf_f . Put $X_i = \ell_{eA}(M_i)$ for $0 \leq i \leq n$. Since by Lemma 2.5 $M_i \in \mathcal{A}_r(eA, Af)$ for all $0 \leq i \leq n$, by Lemmas 2.4(1) and 2.2(1) we have a composition series $0 = X_n \subset \dots \subset X_1 \subset X_0 = eA$ of ${}_{eAe}eA$. \square

Lemma 2.7. *Let (eA, Af) be an i -pair in A . Then the following are equivalent.*

- (1) eA_A is A -simple-injective.
- (2) $\ell_{eA}(M) = \ell_{eA}(N)$ implies $N = M$ for submodules N, M of Af_fAf with $N \subset M$.
- (3) $M \in \mathcal{A}_r(eA, Af)$ for every submodule M of Af_fAf .

Proof. (1) \Rightarrow (2). Let N, M be submodules of Af_fAf with $N \subset M$ and $M/N \neq 0$. Since $(MA/NA)f \cong M/N \neq 0$, there exist submodules K, I of MA_A such that $NA \subset K \subset I$ and $I/K \cong fA/fJ$. Let $\mu : I_A \rightarrow A_A$ denote the inclusion. Since we have $\theta : I_A \rightarrow eA_A$ with $\text{Im } \theta = \text{soc}(eA_A)$ and $\text{Ker } \theta = K$, there exists $\phi : A_A \rightarrow eA_A$ with $\phi \circ \mu = \theta$. Then $\phi(1)I = \phi(I) = \theta(I) \neq 0$ and $\phi(1)K = \phi(K) = \theta(K) = 0$. Thus $\phi(1) \in \ell_{eA}(K)$ and $\phi(1) \notin \ell_{eA}(I)$. Since $\ell_{eA}(M) = \ell_{eA}(MA) \subset \ell_{eA}(I) \subset \ell_{eA}(K) \subset \ell_{eA}(NA) = \ell_{eA}(N)$, $\ell_{eA}(I) \neq \ell_{eA}(K)$ implies $\ell_{eA}(M) \neq \ell_{eA}(N)$.

(2) \Rightarrow (3). Let M be a submodule of Af_fAf and put $L = r_{Af}(\ell_{eA}(M))$. Then $M \subset L$ and $\ell_{eA}(L) = \ell_{eA}(r_{Af}(\ell_{eA}(M))) = \ell_{eA}(M)$. Thus $M = L$.

(3) \Rightarrow (1). Let I be a nonzero right ideal and $\mu : I_A \rightarrow A_A$ the inclusion. Let $\theta : I_A \rightarrow eA_A$ with $\text{Im } \theta = \text{soc}(eA_A)$ and put $K = \text{Ker } \theta$. Then by Lemma 1.1 $I f / K f_f A f \cong (I / K) f_f A f$ is simple, so is ${}_{eAe} \ell_{eA}(K f) / \ell_{eA}(I f)$ by Lemma 2.4(1). Let $a \in I f$ with $a \notin K f$. Then, since $\ell_{eA}(K f) a \neq 0$ and $\ell_{eA}(I f) a = 0$, ${}_{eAe} \ell_{eA}(K f) a$ is simple. Thus by Lemma 2.2(2) $\ell_{eA}(K f) a = \text{soc}(eA_A) f$, so that $\theta(a) = \theta(a f) = \theta(a) f = b a$ with $b \in \ell_{eA}(K f)$. Define $\phi : A_A \rightarrow eA_A$ by $1 \mapsto b$. Then, since by Lemmas 2.2(1) and 2.1(2) $b \in \ell_{eA}(K)$, and since $I = K + aA$, we have $\phi \circ \mu = \theta$. □

Lemma 2.8. *Let (eA, Af) be an i -pair in A . Assume eA_A is injective. Then the canonical homomorphism ${}_{eAe} eA_A \rightarrow {}_{eAe} \text{Hom}_{fAf}(Af, eAf)_A$, $a \mapsto (b \mapsto ab)$, is an isomorphism and eAf_fAf is injective.*

Proof. By Lemmas 2.2(1) and 1.3. □

3. Injective pairs in perfect rings

In this section, we extend results of Baba and Oshiro [1] to left perfect rings. We refer to [2] for perfect rings. We abbreviate the ascending (resp. descending) chain condition as the ACC (resp. DCC).

REMARK 4. (1) Let (eA, Af) be an i -pair in A . Then, since $\mathcal{A}_\ell(eA, Af)$ is anti-isomorphic to $\mathcal{A}_r(eA, Af)$, $\mathcal{A}_\ell(eA, Af)$ satisfies the ACC (resp. DCC) if and only if $\mathcal{A}_r(eA, Af)$ satisfies the DCC (resp. ACC).

(2) Let $e \in A$ be an idempotent. Then, since ${}_{eAe} eAe$ appears as a direct sum-

mand in ${}_{eAe}eA$, ${}_{eAe}eA$ is artinian if and only if it has finite composition length.

(3) Every module L_A with $\text{soc}(L_A) = 0$ is A -simple-injective.

Lemma 3.1 (cf. [1, Proposition 5]). *Let (eA, Af) be an i -pair in A . Assume $\mathcal{A}_r(eA, Af)$ satisfies the ACC and fAf is a left perfect ring. Then Af_fAf is artinian and $M \in \mathcal{A}_r(eA, Af)$ for every submodule M of Af_fAf .*

Proof. It follows by Lemma 2.5 that there exists a maximal element M in the set of submodules of Af_fAf of finite composition length. We claim $M = Af_fAf$. Otherwise, there exists a submodule L of Af_fAf with $M \subset L$ and L/M simple, a contradiction. Thus Af_fAf has finite composition length and again by Lemma 2.5 the last assertion follows. □

Proposition 3.2. *Let (eA, Af) be an i -pair in a left perfect ring A . Then the following are equivalent.*

- (1) ${}_{eAe}eA$ is artinian.
- (2) $\mathcal{A}_\ell(eA, Af)$ satisfies both the ACC and the DCC.
- (3) $\mathcal{A}_\ell(eA, Af)$ satisfies the ACC.

Proof. The implications (1) \Rightarrow (2) \Rightarrow (3) are obvious.

(3) \Rightarrow (1). Since the ascending chain $\ell_{eA}(Af) \subset \ell_{eA}(Jf) \subset \ell_{eA}(J^2f) \subset \dots$ in $\mathcal{A}_\ell(eA, Af)$ terminates, $\ell_{eA}(J^n f) = \ell_{eA}(J^{n+1} f)$ for some $n \geq 0$. We claim $\ell_{eA}(J^n f) = eA$. Suppose otherwise. Then there exists a submodule M of eA_A with $\ell_{eA}(J^n f) \subset M$ and $M/\ell_{eA}(J^n f)$ simple. Since $MJ \subset \ell_{eA}(J^n f)$, $MJ^{n+1}f \subset \ell_{eA}(J^n f)J^n f = 0$ and $M \subset \ell_{eA}(J^{n+1} f) = \ell_{eA}(J^n f)$, a contradiction. Thus $\ell_{eA}(J^n f) = eA$ and by Lemma 2.2(1) $J^n f \subset r_{Af}(\ell_{eA}(J^n f)) = 0$. Then by Lemma 2.3 $eJ^n = 0$ and eAe is a semiprimary ring. Thus by Lemma 3.1 ${}_{eAe}eA$ is artinian. □

Lemma 3.3. *Let $e \in A$ be a local idempotent. Assume eA_A is A -simple-injective and has nonzero socle. Then $\text{soc}(eA_A)$ is simple.*

Proof. Let S be a simple submodule of $\text{soc}(eA_A)_A$. We claim $S = \text{soc}(eA_A)$. Suppose otherwise. Let $\pi : \text{soc}(eA_A) \rightarrow S_A$ be a projection and $\mu : \text{soc}(eA_A) \rightarrow eA_A$, $\nu : S_A \rightarrow eA_A$ inclusions. There exists $\phi : eA_A \rightarrow eA_A$ with $\phi \circ \mu = \nu \circ \pi$. Since π is not monic, ϕ is not an isomorphism. Thus $\phi(e) \in eJe$ and $(e - \phi(e))$ is a unit in eAe . For any $x \in S$, since $\phi(e)x = \phi(x) = \pi(x) = x$, $(e - \phi(e))x = 0$ and thus $x = 0$, a contradiction. □

Lemma 3.4 (cf. [1, Proposition 2]). *Let A be a semiperfect ring and $e \in A$ a local idempotent. Assume eA_A is A -simple-injective and has finite Loewy length. Then*

eA_A is injective.

Proof. Let I be a nonzero right ideal and $\mu : I_A \rightarrow A_A$ the inclusion. Let $\theta : I_A \rightarrow eA_A$. We make use of induction on the Loewy length of $\theta(I)$ to show the existence of $\phi : A_A \rightarrow eA_A$ with $\theta = \phi \circ \mu$. Let $n = \min\{k \geq 0 \mid \theta(I)J^k = 0\}$. We may assume $n > 0$. Since eA_A has nonzero socle, by Lemma 3.3 $\text{soc}(eA_A)$ is simple and $\text{soc}(eA_A) = \theta(I)J^{n-1} = \theta(IJ^{n-1})$. Let μ_1 and θ_1 denote the restrictions of μ and θ to IJ^{n-1} , respectively. Then $\text{Im } \theta_1 = \text{soc}(eA_A)$ and there exists $\phi_1 : A_A \rightarrow eA_A$ with $\phi_1 \circ \mu_1 = \theta_1$. Since $(\theta - \phi_1 \circ \mu)(I)J^{n-1} = 0$, by induction hypothesis there exists $\phi_2 : A_A \rightarrow eA_A$ with $\phi_2 \circ \mu = \theta - \phi_1 \circ \mu$. Then $\theta = (\phi_1 + \phi_2) \circ \mu$. \square

Lemma 3.5 (cf. [1, Proposition 4]). *Let A be a semiperfect ring and $e \in A$ a local idempotent. Assume eA_A is A -simple-injective and has essential socle. Then there exists a local idempotent $f \in A$ such that (eA, Af) is an i -pair in A .*

Proof. By Lemma 3.3 $S_A = \text{soc}(eA_A)$ is simple. Let $f \in A$ be a local idempotent with $Sf \neq 0$. We claim that (eA, Af) is an i -pair in A . Let $0 \neq a \in Sf$. It suffices to show $a \in Ab$ for all $0 \neq b \in Af$. Let $0 \neq b \in Af$. Define $\alpha : fA_A \rightarrow aA_A$ by $x \mapsto ax$ and $\beta : fA_A \rightarrow bA_A$ by $x \mapsto bx$. Since $\text{Ker } \beta = r_{fA}(b) \subset fJ = r_{fA}(a) = \text{Ker } \alpha$, we have $\theta : bA_A \rightarrow aA_A = S_A$ with $\alpha = \theta \circ \beta$. Let $\mu : S_A \rightarrow eA_A$, $\nu : bA_A \rightarrow A_A$ be inclusions. Then there exists $\phi : A_A \rightarrow eA_A$ with $\phi \circ \nu = \mu \circ \theta$ and $a = \alpha(f) = \theta(\beta(f)) = \theta(b) = \phi(b) = \phi(1)b \in Ab$. \square

Theorem 3.6 (cf. [1, Theorem 1]). *Let A be a left perfect ring and $e \in A$ a local idempotent. Then the following are equivalent.*

- (1) eA_A is A -simple-injective.
- (2) There exists a local idempotent $f \in A$ such that (eA, Af) is an i -pair in A and $M \in \mathcal{A}_r(eA, Af)$ for every submodule M of Af_{fAf} .

Proof. By Lemmas 3.5 and 2.7. \square

Theorem 3.7 (cf. [1, Theorem 2]). *Let (eA, Af) be an i -pair in a left perfect ring A . Then the following are equivalent.*

- (1) $eA_e eA$ is artinian.
- (2) Af_{fAf} is artinian.
- (3) Both eA_A and ${}_A Af$ are injective.

Proof. (1) \Leftrightarrow (2). By Lemma 2.6.

(2) \Rightarrow (3). By Lemmas 2.6, 2.5 and 2.7 both eA_A and ${}_A Af$ are A -simple-injective. Also, by Lemma 2.3 both eA_A and ${}_A Af$ have finite Loewy length. Thus by Lemma 3.4 both eA_A and ${}_A Af$ are injective.

(3) \Rightarrow (1). By Lemma 2.8 the canonical homomorphism

$${}_{eAe}eA_A \rightarrow {}_{eAe}\text{Hom}_{fAf}(Af, eAf)_A$$

is an isomorphism and eAf_{fAf} is injective. Similarly, the canonical homomorphism ${}_AAf_{fAf} \rightarrow {}_A\text{Hom}_{eAe}(eA, eAf)_{fAf}$ is an isomorphism and ${}_{eAe}eAf$ is injective. It follows that ${}_{eAe}eAf_{fAf}$ defines a Morita duality. Thus by [7, Theorem 3] eAe is left artinian and ${}_{eAe}eA$ has finite Loewy length. Since the canonical homomorphism ${}_{eAe}eA \rightarrow {}_{eAe}\text{Hom}_{fAf}(\text{Hom}_{eAe}(eA, eAf), eAf)$ is an isomorphism, it follows by [7, Lemma 13] that ${}_{eAe}eA$ has finite composition length. \square

REMARK 5. In Theorem 3.7 the assumption that A is left perfect cannot be replaced by a weaker condition that A is semiperfect (see [7, Example 1]).

References

- [1] Y. Baba and K. Oshiro: *On a theorem of Fuller*, J. Algebra, **154** (1993), 86–94.
- [2] H. Bass: *Finitistic dimension and a homological generalization of semi-primary rings*, Trans. Amer. Math. Soc. **95** (1960), 466–488.
- [3] K.R. Fuller: *On indecomposable injectives over artinian rings*, Pacific J. Math. **29** (1969), 115–135.
- [4] M. Harada: *Note on almost relative projectives and almost relative injectives*, Osaka J. Math. **29** (1992), 435–446.
- [5] T. Kato: *Torsionless modules*, Tôhoku Math. J. **20** (1968), 234–243.
- [6] M. Morimoto and T. Sumioka: *Generalizations of theorems of Fuller*, Osaka J. Math. to appear.
- [7] B.L. Osofsky: *A generalization of quasi-Frobenius rings*, J. Algebra, **4** (1966), 373–387.
- [8] T. Sumioka and S. Tozaki: *On almost QF-rings*, Osaka J. Math. **33** (1996), 649–661.

M. Hoshino
Institute of Mathematics
University of Tsukuba
Ibaraki, 305-0006
Japan

T. Sumioka
Department of Mathematics
Osaka City University
Osaka, 558-8585
Japan