

INFINITE ALGEBRAIC SUBGROUPS OF THE REAL CREMONA GROUP

MARIA FERNANDA ROBAYO and SUSANNA ZIMMERMANN

(Received January 4, 2017, revised April 28, 2017)

Abstract

We give the classification of the maximal infinite algebraic subgroups of the real Cremona group of the plane up to conjugacy and present a parametrisation space of each conjugacy class. Moreover, we show that the real plane Cremona group is not generated by a countable union of its infinite algebraic subgroups.

Contents

| | |
|---|-----|
| 1. Introduction | 681 |
| 2. First steps | 684 |
| 2.1. Algebraic groups | 684 |
| 2.2. Minimal surfaces | 687 |
| 3. Real rational del Pezzo surfaces of degree 6 | 690 |
| 3.1. The surfaces obtained by blowing up the sphere | 690 |
| 3.2. The surfaces obtained by blowing up \mathbb{F}_0 | 694 |
| 4. Pairs of real conic bundles | 697 |
| 4.1. Real conic bundles obtained by blowing up a del Pezzo surface | 698 |
| 4.2. Real conic bundles obtained by blowing up a Hirzebruch surface | 704 |
| 5. The maximal infinite algebraic subgroups | 707 |
| References | 711 |

1. Introduction

It is most natural to study actions of algebraic groups on algebraic varieties. One tends to assume the action to be regular, but this is quite restrictive; any algebraic group acting regularly on the n -dimensional projective space \mathbb{P}^n is a subgroup of $\text{Aut}_k(\mathbb{P}^n) = \text{PGL}_{n+1}(k)$. It is therefore interesting to study rational group actions on \mathbb{P}^n , or, equivalently, algebraic subgroups of $\text{Bir}_k(\mathbb{P}^n)$. Algebraic subgroups of the complex Cremona group have been studied by many mathematicians, we refer to [2] for a historical note where also the complete classification of the maximal algebraic subgroups of the complex Cremona group $\text{Bir}_{\mathbb{C}}(\mathbb{P}^2)$

2010 Mathematics Subject Classification. Primary 14E07; Secondary 14L99, 14P99.

The first author gratefully acknowledges support from the Department of Mathematics and Computer Science, University of Basel. The second author gratefully acknowledges support by the Swiss National Science Foundation via grants PP00P2_153026/1 and P2BSP2_168743.

is presented.

In this paper, we give a classification of the maximal infinite algebraic subgroups of the real Cremona group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. The classification is built on the classification of minimal real smooth projective surfaces given in [7], which is manageable, quite opposed to the case of a general perfect field.

The finite subgroups of the real Cremona group of odd order have already been classified in [24]; they are conjugate to a subgroup of the automorphism group of a real del Pezzo surface. It is also interesting to look at the group of birational transformations of a rational real smooth minimal model X that are well defined on the set of real points. The group is usually called the group of birational diffeomorphisms of X and has been studied for instance in [1, 18, 11, 14, 4, 16]. The birational diffeomorphisms of the sphere of prime order have been classified in [16].

In the classification of the infinite algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, two of the infinite families in the classification of the complex algebraic subgroups of the Cremona group split into two families each: There exist four isomorphism classes of real del Pezzo surfaces of degree 6, and the automorphism group of two of them are infinite maximal algebraic subgroups. The family of conic bundles splits into real conic bundles coming from a del Pezzo surface of degree 6 obtained by blowing up the sphere in a pair of non-real conjugate points and into real conic bundles coming from Hirzebruch surfaces. In both cases, an infinite number of them have automorphism groups that are maximal infinite algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, pairwise non-conjugate. More concretely, the classification is as follows.

By D_6 we denote the dihedral group with twelve elements.

Theorem 1.1. *Every infinite algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is contained in a maximal algebraic subgroup.*

An infinite maximal algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is conjugate to $G = \text{Aut}_{\mathbb{R}}(X)$ where X is a real del Pezzo surface or to $G = \text{Aut}_{\mathbb{R}}(X, \pi)$ where $\pi: X \rightarrow \mathbb{P}^1$ is a real conic bundle, and the pairs (X, G) are described as follows:

- (1) $X \simeq \mathbb{P}^2$ and $G \simeq \text{PGL}_3(\mathbb{R})$,
- (2) $X \simeq \mathcal{Q}_{3,1} \subset \mathbb{P}^3$ is the real rational minimal surface defined by $w^2 = x^2 + y^2 + z^2$ whose real part is diffeomorphic to the 2-sphere \mathbb{S}^2 , and $G \simeq \text{PO}_{\mathbb{R}}(3, 1)$,
- (3) $X \simeq \mathbb{P}^1 \times \mathbb{P}^1$ and $G \simeq (\text{PGL}_2(\mathbb{R}) \times \text{PGL}_2(\mathbb{R})) \rtimes \langle \tau \rangle$, where $\tau: (x, y) \mapsto (y, x)$,
- (4) X is a del Pezzo surface of degree 6 obtained by blowing up a pair of non-real conjugate points on \mathbb{F}_0 , and the action of G on $\text{Pic}(X)$ induces the split exact sequence

$$1 \rightarrow \text{SO}_2(\mathbb{R})^2 \rightarrow G \rightarrow D_6 \rightarrow 1,$$

- (5) X is a del Pezzo surface of degree 6 obtained by blowing up two real points on \mathbb{F}_0 , and the action of G on $\text{Pic}(X)$ induces the split exact sequence

$$1 \rightarrow (\mathbb{R}^*)^2 \rightarrow G \rightarrow D_6 \rightarrow 1,$$

- (6) $X \simeq \mathbb{F}_n$, $n \geq 2$, is the n -th Hirzebruch surface and $G \simeq \mathbb{R}^{n+1} \rtimes \text{GL}_2(\mathbb{R})/\mu_n$, where $\mu_n = \{\pm 1\}$ if n is even and $\mu_n = \{1\}$ if n is odd.
- (7) $\eta: X \rightarrow X_{[2]}$ is a birational morphism of real conic bundles, where $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$ is the conic bundle obtained by blowing up a pair of non-real conjugate points on $\mathcal{Q}_{3,1}$. The morphism η blows up $n \geq 1$ pairs of non-real conjugate points with non-real

fibres on the non-real conjugate disjoint (-1) -curves of $X_{[2]}$ that are the exceptional divisors of $X_{[2]} \rightarrow \mathcal{Q}_{3,1}$. The action of G on \mathbb{P}^1 induces the split exact sequence

$$1 \rightarrow \mathrm{SO}_2(\mathbb{R}) \rtimes \mathbb{Z}/2\mathbb{Z} \rightarrow G \rightarrow H_\Delta \rightarrow 1$$

where $H_\Delta \subset \mathrm{PGL}_2(\mathbb{R})$ is the subgroup preserving the set of images in \mathbb{P}^1 of the points blown up by η and the interval $\pi(X(\mathbb{R})) = \pi_{[2]}(X_{[2]}(\mathbb{R}))$,

- (8) *$\eta: X \rightarrow \mathbb{F}_n$ is a birational morphism of real conic bundles that is the blow-up of $2n \geq 4$ points on the zero section s_n of self-intersection n (see Section 4.2). The action of G on \mathbb{P}^1 induces a split exact sequence*

$$1 \rightarrow (\mathbb{R}^*/\mu_n) \rtimes \mathbb{Z}/2\mathbb{Z} \rightarrow G \rightarrow H_\Delta \rightarrow 1,$$

where $H_\Delta \subset \mathrm{PGL}_2(\mathbb{R})$ is the subgroup preserving the set of images in \mathbb{P}^1 of the points blown up by η , and $\mu_n = \{\pm 1\}$ if n is even and $\mu_n = \{1\}$ if n is odd.

Furthermore, the families (1) – (8) are distinct and pairwise non-conjugate in $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.

We also give the parameter space of the maximal infinite algebraic subgroups of $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.

Theorem 1.2. *The families in Theorem 1.1 are distinct families and the conjugacy classes in each family are parametrised by*

- (1)-(5) *One point.*
- (6) *One point for each $n \geq 2$.*
- (7) *For each $n \geq 1$, the set of n pairs of non-real conjugate points in \mathbb{P}^1 modulo the action of $\mathrm{Aut}_{\mathbb{R}}(\mathbb{P}^1, [0, \infty])$.*
- (8) *For each $n \geq 2$, the set of $2n$ points in \mathbb{P}^1 consisting of real points or pairs of non-real conjugate points, modulo the action of $\mathrm{Aut}_{\mathbb{R}}(\mathbb{P}^1) = \mathrm{PGL}_2(\mathbb{R})$.*

The groups listed in Theorem 1.1 are contained in infinite (complex) algebraic subgroups of $\mathrm{Bir}_{\mathbb{C}}(\mathbb{P}^2)$ (classified in [2, Theorem 2]), and they are all dense in their complex counterpart.

The elements of the group of birational diffeomorphisms of the sphere of prime order are contained in $\mathrm{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1})$, or are automorphisms of real del Pezzo surfaces of degree 2 or 4, or are automorphisms of real conic bundles as in family (7) [16]. The finite subgroups of $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ of odd order are contained in automorphism groups appearing in the families (1), (2), (3), (4) or in automorphism groups of del Pezzo surfaces of degree 5 [24]. The classification in Theorem 1.1 does not list automorphism groups of del Pezzo surfaces of degree 5 because they are finite.

The real Cremona group of the plane is generated by the family of standard quintic transformations and $\mathrm{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ [4], and its abelianisation $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)/\langle\langle \mathrm{Aut}_{\mathbb{R}}(\mathbb{P}^2) \rangle\rangle \simeq \bigoplus_{(0,1)} \mathbb{Z}/2\mathbb{Z}$ is generated by the classes of the standard quintic transformations [25]. The classification of the maximal infinite subgroups of $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ yields the following theorem.

Theorem 1.3. *An infinite algebraic subgroup of $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ with non-trivial image in the abelianisation $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)/\langle\langle \mathrm{Aut}_{\mathbb{R}}(\mathbb{P}^2) \rangle\rangle \simeq \bigoplus_{(0,1)} \mathbb{Z}/2\mathbb{Z}$ has finite image in the abelianisation and is conjugate to a subgroup of an algebraic group in family (7) of Theorem 1.1.*

Furthermore, for each generator of $\bigoplus_{(0,1)} \mathbb{Z}/2\mathbb{Z}$ there is a conjugacy class of infinite algebraic groups in family (7) of Theorem 1.1 which is sent onto the generator.

Corollary 1.4. *The group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is not generated by a countable union of infinite algebraic subgroups.*

Quite the opposite is true for the Cremona group $\text{Bir}_{\mathbb{C}}(\mathbb{P}^2)$ of the complex plane. As the standard quadratic transformation of \mathbb{P}^2 is conjugate to an automorphism of $\mathbb{P}^1 \times \mathbb{P}^1$, it follows from the Noether-Castelnuovo theorem [6] that $\text{Bir}_{\mathbb{C}}(\mathbb{P}^2)$ is generated by the infinite algebraic groups $\text{Aut}_{\mathbb{C}}(\mathbb{P}^2)$ and $\text{Aut}_{\mathbb{C}}(\mathbb{P}^1 \times \mathbb{P}^1)$.

2. First steps

2.1. Algebraic groups. For a projective algebraic variety X defined over k , we denote by $\text{Bir}_k(X)$ its group of birational self-maps defined over k and by $\text{Aut}_k(X) \subset \text{Bir}_k(X)$ the group of k -automorphisms of X defined over k .

Note on the side: the group $\text{Aut}_k(X)$ is the group of k -rational points of a group scheme locally of finite type over k , having at most countably many connected components.

The definition of rational actions on varieties goes back to Weil and Rosenlicht (see for instance [23, 17]).

DEFINITION 2.1. Let X be an algebraic variety over k .

(1) Let G be an algebraic group defined over k . A *k -rational action* ρ of G on X is a k -rational morphism $\rho: G \times X \dashrightarrow X$ such that

- $\rho(e, x) = x$ for all $x \in X$,
- ρ is associative whenever it is defined, i.e. $\rho(g_1, \rho(g_2, x)) = \rho(g_1 g_2, x)$ whenever ρ is defined at (g_2, x) , at $(g_1, \rho(g_2, x))$ and at $\rho(g_1 g_2, x)$,
- there exist open dense subsets $U, V \subset G \times X$ such that the k -rational map $G \times X \dashrightarrow G \times X$, $(g, x) \mapsto (g, gx)$ restricts to an isomorphism $U \rightarrow V$ and the projection of U and V to the first factor is surjective onto G .

For each k -rational point $g \in G$, we get a k -birational map $\rho(g, \cdot): X \dashrightarrow X$, and this induces a group homomorphism $G(k) \rightarrow \text{Bir}_k(X)$.

(2) We say that X is a *G -variety* if ρ is regular. Then the action induces a group homomorphism $G(k) \rightarrow \text{Aut}_k(X)$.

REMARK 2.2. Let G be an algebraic group with a k -rational action on X . For $g \in G$ we obtain a k -birational map $\rho(g, \cdot): X \dashrightarrow X$ if and only if $g \in G(k)$. The induced homomorphism $G(k) \rightarrow \text{Bir}_k(X)$ is injective if the action of G on X is faithful.

In this paper, we classify the infinite (not necessarily connected) maximal algebraic subgroups $G(k)$ up to conjugation inside $\text{Bir}_k(X)$ for $k = \mathbb{R}$. The classification over algebraically closed fields of characteristic zero is done in [2] (in fact, it also classifies the finite maximal ones).

As result of looking at $G(\mathbb{R})$ only, the groups in the classification are real linear algebraic groups, i.e. can be embedded into some $\text{GL}_N(\mathbb{R})$.

DEFINITION 2.3. Let G be an algebraic group with a k -rational faithful action on X . Then the induced group homomorphism $G(k) \rightarrow \text{Bir}_k(X)$ is injective, and we call $G(k)$ an *algebraic subgroup* of $\text{Bir}_k(X)$.

REMARK 2.4. Note that classically, the algebraic group G is called algebraic subgroup of $\text{Bir}_k(X)$, and not the group $G(k)$. However, only the k -rational points of G can be viewed as elements of $\text{Bir}_k(X)$.

For a real variety (X, σ) , one can also define an algebraic subgroup of $\text{Bir}_{\mathbb{R}}(X)$ to be an algebraic subgroup $G \subset \text{Bir}_{\mathbb{C}}(X)$ such that $\sigma(G) = G$. If we further impose that $G(\mathbb{R})$ is dense in G , then the classification of infinite maximal algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is exactly the one in Theorem 1.1. However, without this condition, the classification is longer. The additional examples appear in Section 3.1. Let $G := \text{Aut}_{\mathbb{C}}(X_{[2]}) = (\mathbb{C}^*)^2 \rtimes D_6$. Then

$$G \supset \text{Aut}_{\mathbb{R}}(X_{[2]}) = (\mathbb{R}_{>0} \times \text{SO}_2(\mathbb{R})) \rtimes (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})$$

and

$$G \supset \text{Aut}_{\mathbb{R}}(X_{[3, Q_{3,1}]}) = \text{SO}_2(\mathbb{R}) \rtimes (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})$$

are two maximal algebraic groups ([2, Théorème 2]) whose real locus $G(\mathbb{R}) = \text{Aut}_{\mathbb{R}}(X_{[2]})$ is not dense in G .

REMARK 2.5. Let G be an algebraic group. Recall that a *morphism* $G \rightarrow \text{Bir}_k(X)$ is defined as follows (see for instance [3]): let $\mu: G \times X \dashrightarrow G \times X$ be a k -rational map inducing an isomorphism $U \rightarrow V$, where $U, V \subset G \times X$ are open dense subsets whose projections onto G are surjective. The rational map μ is given by $(g, x) \mapsto (g, p_2(\mu(g, x)))$, where p_2 is the second projection, and for each k -rational point $g \in G$, the birational map $x \mapsto p_2(\mu(g, x))$ corresponds to an element $f_g \in \text{Bir}_k(X)$. The maps $g \mapsto f_g$ represent a map from $G(k)$ to $\text{Bir}_k(X)$, which is called a *morphism* from G to $\text{Bir}_k(X)$.

If the map $G(k) \rightarrow \text{Bir}_k(X)$ is a homomorphism of groups, then its image is an algebraic subgroup of $\text{Bir}_k(X)$.

The group $\text{Bir}_k(X)$ can be endowed with the so-called Zariski topology, introduced by [3, 9, 20], which is compatible with the concept of morphism of varieties into $\text{Bir}_k(X)$; a subset $F \subset \text{Bir}_k(X)$ is closed if for any algebraic variety A and any morphism $A \rightarrow \text{Bir}_k(X)$, the pre-image of F is closed. Endowed with the Zariski topology, $\text{Bir}_k(\mathbb{P}^n)$ is not an ind-variety, algebraic stack or algebraic space if $n \geq 2$ [3, Theorem 1, Remark 3.5]. The following remark gives a sufficient and necessary condition for a subgroup of $\text{Bir}_k(\mathbb{P}^n)$ to be an algebraic subgroup.

REMARK 2.6. Let $G \subset \text{Bir}_k(\mathbb{P}^n)$ be a subgroup. It is an algebraic subgroup if and only if it is closed in the Zariski topology and of bounded degree [3, Remark 2.20]. In that case, G is by definition the group of k -rational points of an algebraic group H acting rationally on \mathbb{P}^n . The group H is affine [3, Remark 2.21], and so G is an affine [22, §4.4, Corollary].

The fact that algebraic subgroups of $\text{Bir}_k(\mathbb{P}^2)$ are linear algebraic groups, makes them approachable with classical tools.

Lemma 2.7. *Let k be a perfect field, G a linear algebraic group and X a smooth G -surface, all defined over k . Then there exists a smooth projective G -surface Y and a G -equivariant birational map $X \dashrightarrow Y$.*

Proof. Forgetting about the group action, we see that X is contained as open set in a complete surface, which can be desingularised [15]. Smooth complete surfaces are projective, hence X is contained as open set in a projective surface and thus is quasi-projective. Let G_0 be the neutral component of G . We can apply [21, Theorem 4.9]; there exists a G_0 -equivariant smooth completion of X . Equivalently, X admits a G_0 -linearisable ample line bundle. [5, Lemma 3.2] implies that X admits a G -linearised ample line bundle and hence a G -equivariant completion Y of X . We may replace Y with a G -equivariant desingularisation. We have found a birational G -equivariant map $X \dashrightarrow Y$ to a smooth projective G -surface. \square

By \bar{k} we denote the algebraic closure of the field k .

DEFINITION 2.8. An algebraic variety X over k is *geometrically rational* if $X_{\bar{k}} := X \times_{\text{Spec}(k)} \text{Spec}(\bar{k})$ is rational, i.e. if it is rational as variety over \bar{k} .

A geometrically rational variety X is *k-rational* if there is a birational map $X \dashrightarrow \mathbb{P}^n$ defined over k .

The following lemma is classical and states a necessary and sufficient condition for a (abstract) subgroup of $\text{Aut}_k(X)$ to have the structure of a linear algebraic group acting regularly on X .

DEFINITION 2.9. For a smooth projective variety X over a perfect field k with $X(k) \neq \emptyset$, we have $\bar{k}[X_{\bar{k}}]^* = (\bar{k})^*$ and then $\text{Pic}(X) = \text{Pic}(X_{\bar{k}})^{\text{Gal}(\bar{k}/k)}$ is the Galois-invariant Picard group [19, Lemma 6.3(iii)]. We denote by $\text{Pic}(X)^G = \text{Pic}(X_{\bar{k}})^{\text{Gal}(\bar{k}/k) \times G}$ the G -invariant Picard group.

The action of $\text{Aut}_k(X)$ on X induces a homomorphism of (abstract) groups

$$\text{Aut}_k(X) \rightarrow \text{Aut}(\text{Pic}(X)).$$

Lemma 2.10. *Let X be a smooth projective variety defined over a perfect field k .*

- (1) *Let D be a very ample divisor on X defined over k and $G \subset \text{Aut}_k(X)$ the group of elements fixing D . Then the k -embedding $X \hookrightarrow \mathbb{P}^n$ given by the linear system of D conjugates G to a closed subgroup of $\text{PGL}_{n+1}(k)$.*
- (2) *The kernel $K = \ker(\text{Aut}_k(X) \rightarrow \text{Aut}(\text{Pic}(X)))$ has the structure of a linear algebraic group acting regularly on X via the given inclusion $K \subset \text{Aut}_k(X)$.*
- (3) *Any subgroup $G \subset \text{Aut}_k(X)$ containing K whose action on $\text{Pic}(X)$ is finite has the structure of a linear algebraic group acting regularly on X via the inclusion $G \subset \text{Aut}_k(X)$.*
- (4) *Suppose that G is a linear algebraic group and X a k -rational G -variety. If G contains K , then G has finite action on $\text{Pic}(X)$.*

Proof. (1): The linear system of D induces a closed embedding $\varphi_D: X \hookrightarrow \mathbb{P}^N$ defined over k . Let $G \subset \text{Aut}_k(X)$ be the subgroup of elements whose image in $\text{Aut}(\text{Pic}(X))$ fix D . Denote by $H \subset \text{PGL}_{N+1}(k)$ the subgroup preserving $\varphi_D(X)$, which is a closed subgroup. It is also a subgroup of $\text{Aut}_k(\varphi_D(X))$, and G is conjugate to H via φ_D .

(2): For any very ample divisor D on X defined over k , the group G from (1) contains K as a closed subgroup, which is therefore a linear algebraic group as well and acts on X regularly.

(3): If $G \subset \text{Aut}_k(X)$ has finite action on $\text{Pic}(X)$ and contains K , then G/K is finite. As K is a linear algebraic group, also G is one.

(4): The group G is a linear algebraic group by assumption and the group K is a normal, closed linear algebraic group by (2), so the group G/K is a linear algebraic group. If X is a k -rational smooth projective variety then $\text{Pic}(X)$ is finitely generated and has no torsion, i.e. $\text{Pic}(X) \simeq \mathbb{Z}^n$ for some $n \in \mathbb{N}$. By assumption, X is a G -variety, so there is an inclusion $G \subset \text{Aut}_k(X)$ of abstract groups. Then $G/K \subset \text{Aut}(\text{Pic}(X)) \subset \text{GL}_n(\mathbb{Z})$. Since $\text{GL}_n(\mathbb{Z})$ is countable, the group G/K is finite. \square

2.2. Minimal surfaces.

DEFINITION 2.11. We denote by (X, G) the pair consisting of a smooth projective surface X defined over k and G a subgroup of $\text{Aut}_k(X)$.

- (1) We say that (X, G) is a *minimal pair* (or X is G -*minimal*) if for any smooth projective surface Y over k any birational G -morphism $X \rightarrow Y$ is an isomorphism.
- (2) Let $\pi: X \rightarrow C$ be a G -equivariant morphism, where C a curve. We say that π is *relatively G -minimal* if for any decomposition $\pi: X \xrightarrow{\eta} Y \xrightarrow{\pi'} C$, where π' is a G -equivariant morphism and η is a birational G -equivariant morphism, η is in fact an isomorphism,

Note that for $G = \{1\}$, a G -minimal surface is just a minimal surface.

DEFINITION 2.12. We say that a smooth projective G -surface X admits a conic bundle structure if there exists $\pi: X \rightarrow C$, where C is a smooth curve and the fibre over closed point t is isomorphic to a reduced conic over the residue field $k(t)$ of t .

REMARK 2.13. A G -surface admitting a conic bundle structure $\pi: X \rightarrow C$ is relatively G -minimal if any k -birational G -equivariant morphism $X \rightarrow Y$ of conic bundles is an isomorphism.

If X is a geometrically rational smooth surface, then $C_{\bar{k}} \simeq \mathbb{P}^1$. If moreover $X(k) \neq \emptyset$, then $C(k) \neq \emptyset$, and so $C \simeq \mathbb{P}^1$ over k .

A real smooth projective surface X can be seen as a pair $(X_{\mathbb{C}}, \sigma)$ consisting of a smooth projective complex variety $X_{\mathbb{C}}$ and an antiholomorphic involution σ .

DEFINITION 2.14. For a real conic bundle $\pi: X \rightarrow \mathbb{P}^1$, we define

$$\text{Aut}_{\mathbb{R}}(X, \pi) := \{f \in \text{Aut}_{\mathbb{R}}(X) \mid \exists \alpha \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^1) : \pi f = \alpha \pi\} \subset \text{Aut}_{\mathbb{R}}(X),$$

the group of automorphisms preserving the conic bundle structure, and

$$\text{Aut}_{\mathbb{R}}(X/\pi) := \{f \in \text{Aut}_{\mathbb{R}}(X, \pi) \mid \pi f = \pi\} \subset \text{Aut}_{\mathbb{R}}(X, \pi),$$

its subgroup acting trivially on \mathbb{P}^1 .

Every Hirzebruch surface \mathbb{F}_n admits a natural real structure with real points: writing

$$\mathbb{F}_n \simeq \{([x_0 : x_1 : x_2], [u : v]) \in \mathbb{P}^2 \times \mathbb{P}^1 \mid x_1 v^n = x_2 u^n\}$$

the standard antiholomorphic involution of $\mathbb{P}^2 \times \mathbb{P}^1$, that is, the standard antiholomorphic involution on either factors, descends to a antiholomorphic involution on \mathbb{F}_n .

DEFINITION 2.15. By $\mathcal{Q}_{3,1} \subset \mathbb{P}^3$, we denote the real surface given by $w^2 = x^2 + y^2 + z^2$ endowed with the standard antiholomorphic involution on \mathbb{P}^3 .

REMARK 2.16. Note that $\mathcal{Q}_{3,1}(\mathbb{R}) = \mathbb{S}^2$ is the 2-dimensional real sphere, $\mathbb{F}_{2n}(\mathbb{R}) = \mathbb{S}^1 \times \mathbb{S}^1$ is the real torus and $\mathbb{F}_{2n+1}(\mathbb{R})$ is the Klein bottle for any $n \geq 0$. The isomorphism of complex surfaces

$$\begin{aligned} \varphi: (\mathcal{Q}_{3,1})_{\mathbb{C}} &\longrightarrow (\mathbb{P}^1 \times \mathbb{P}^1)_{\mathbb{C}} \\ [w : x : y : z] &\mapsto ([w + z : y + \mathbf{i}x], [w + z : y - \mathbf{i}x]) \\ &= ([y - \mathbf{i}x : w - z], [y + \mathbf{i}x : w - z]) \\ \varphi^{-1}: ([x_0 : x_1], [y_0 : y_1]) &\mapsto [x_0y_0 + x_1y_1 : \mathbf{i}(x_0y_1 - x_1y_0) : x_0y_1 + x_1y_0 : x_0y_0 - x_1y_1] \end{aligned}$$

induces an isomorphism of real surfaces $\varphi: \mathcal{Q}_{3,1} \rightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_S)$, where

$$\sigma_S: ([x_0 : x_1], [y_0 : y_1]) \mapsto ([\bar{y}_0 : \bar{y}_1], [\bar{x}_0 : \bar{x}_1])$$

Note that $\mathcal{Q}_{3,1}$ is a del Pezzo surface of degree 8 with $\text{rk}(\text{Pic}(\mathcal{Q}_{3,1})) = 1$, whereas $\text{rk}(\text{Pic}(\mathbb{P}^1 \times \mathbb{P}^1)) = 2$.

Theorem 2.17 ([7]). *Let X be a minimal geometrically rational real surface X with $X(\mathbb{R}) \neq \emptyset$. If X is \mathbb{R} -rational, then X is \mathbb{R} -isomorphic to \mathbb{P}^2 , to the quadric $\mathcal{Q}_{3,1}$ or to a real Hirzebruch surface \mathbb{F}_n , $n \neq 1$.*

DEFINITION 2.18. By $X_{[2]}$ we denote a del Pezzo surface of degree 6 obtained by blowing up $\mathcal{Q}_{3,1}$ in a pair of non-real conjugate points. The notation is motivated by the fact that $\text{rkPic}(X_{[2]}) = 2$.

Lemma 2.19. *Any real surface $X_{[2]}$ is isomorphic to*

$$X_{[2]} \simeq \{([w : x : y : z], [u : v]) \in \mathbb{P}^3 \times \mathbb{P}^1 \mid wz = x^2 + y^2, uz = vw\}$$

endowed with the antiholomorphic involution that is the restriction of the standard antiholomorphic involution on $\mathbb{P}^3 \times \mathbb{P}^1$.

Proof. Consider the surface $S \subset \mathbb{P}^3$ given by $wz = x^2 + y^2$. The isomorphism $[w : x : y : z] \mapsto [w + z : x : y : w - z]$ yields an isomorphism $\mathcal{Q}_{3,1} \simeq S$. Pick a non-real point $p \in S$ and denote by $\eta: X_{[2]} \rightarrow S$ the blow-up of p, \bar{p} . We find an automorphism of S that sends p onto $[0 : 1 : \mathbf{i} : 0]$ [16, Lemma 4.8]. The blow-up of p, \bar{p} on $\mathcal{Q}_{3,1}$ is the restriction of the blow-up of \mathbb{P}^3 along the line l given by $w = z = 0$, as the intersection of l and S is transversal and equal to the set $\{p, \bar{p}\}$. This yields the claim. \square

REMARK 2.20. The projection $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$, $([w : x : y : z], [u : v]) \mapsto [u : v]$ is a real conic bundle morphism with two singular fibres, which lie over 0 and ∞ , and without sections.

Proposition 2.21. *Let G be an infinite algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. Then there exists a G -equivariant real birational map $\mathbb{P}^2 \dashrightarrow X$ to a real smooth G -surface X , which is one of*

the following:

- (1) X is a real del Pezzo surface of degree 6, 8 or 9 such that $\text{rk}(\text{Pic}(X)^G) = 1$.
- (2) X admits a real conic bundle structure $\pi_X: X \rightarrow \mathbb{P}^1$ with $\text{rk}(\text{Pic}(X)^G) = 2$ and $G \subset \text{Aut}_{\mathbb{R}}(X, \pi_X)$.

Furthermore, in (2), there is a birational morphism of conic bundles $\eta: X \rightarrow Y$, where Y is a Hirzebruch surface \mathbb{F}_n , $n \geq 1$, or $Y \simeq X_{[2]}$.

Proof. By Remark 2.6, G is a linear algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. So, there is a real smooth algebraic G -surface X' and a G -equivariant real birational map $\phi: X' \dashrightarrow \mathbb{P}^2$ by [23, Proposition 4] and [17, Theorem 1]. By Lemma 2.7 there exists a smooth real projective G -surface X'' and a G -equivariant real birational map $X \dashrightarrow X''$. After contracting all the sets of disjoint G -invariant real (-1) -curves and all sets of disjoint G -invariant pairs of non-real conjugate (-1) -curves, we obtain a real smooth projective G -variety X that can be one of the following possibilities by [13, Excerise 2.18] and Lemma 2.10 (4):

- (i) X is a del Pezzo surface and $\text{rk}(\text{Pic}(X)^G) = 1$,
- (ii) X admits a real conic bundle structure $X \xrightarrow{\pi_X} \mathbb{P}^1$ and $\text{rk}(\text{Pic}(X)^G) = 2$.

Due to Comessatti [7], the minimal smooth geometrically rational real surfaces are \mathbb{R} -isomorphic to \mathbb{P}^2 , to $\mathcal{Q}_{3,1}$ or to a real Hirzebruch surface \mathbb{F}_n , $n \neq 1$. Thus, forgetting about the action of G and π_X , there is a real birational morphism $X \rightarrow Y$ where Y is one of these minimal surfaces.

In case (i), the situation is as follows: Any real geometrically rational del Pezzo surface is the blow-up of at most 8 \mathbb{C} -points on one of the three real minimal surfaces. The automorphism group of a del Pezzo surface of degree ≤ 5 is finite [10, Section 6]. Any real del Pezzo surface X of degree 7 has three (-1) -curves, one of which is real. Hence X is the blow-up of \mathbb{F}_0 or $\mathcal{Q}_{3,1}$ in one real point p and any automorphism of X preserves its exceptional divisor and hence is the lift of an automorphism from $\mathcal{Q}_{3,1}$ or \mathbb{F}_0 . In particular, $\text{Aut}_{\mathbb{R}}(X)$ is conjugate by the blow-up of p to a subgroup of $\text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$ or $\text{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1})$. This leaves degree 6, 8 and 9.

In case (ii), the situation is as follows: Forgetting the action of G , there is a real birational morphism of real conic bundles $X \rightarrow Y$, which is the contraction of all disjoint real and disjoint pairs of non-real (-1) -curves in the fibres. We obtain a relatively minimal real conic bundle $\pi_Y: Y \rightarrow \mathbb{P}^1$ with no real (-1) -curves and no non-real conjugate singular fibres, i.e. it has at most real singular fibres whose components are non-real conjugate (-1) -curves. Forgetting about π_Y , we obtain a real birational morphism $Y \rightarrow Z$ to a minimal real smooth rational surface, and $Z(\mathbb{R})$ is connected and homeomorphic to the real projective plane, the sphere, the torus or the Klein bottle (see Remark 2.16). It follows that Y has either none or exactly two singular fibres (otherwise $Y(\mathbb{R})$ is not connected and thus not rational). In this case, Y is the blow-up of $\mathcal{Q}_{3,1}$ in a pair of non-real conjugate points, i.e. $Y \simeq X_{[2]}$, or Y has no singular fibres and is isomorphic to a real Hirzebruch surface \mathbb{F}_n , $n \neq 1$. □

Lemma 2.22 (Real version of [2, Proposition 2.2.6]). *Let X be a real smooth projective geometrically rational surface.*

- (1) *If X is a del Pezzo surface, then $\text{Aut}_{\mathbb{R}}(X)$ is a linear algebraic group.*
- (2) *If $\pi: X \rightarrow \mathbb{P}^1$ is a real conic bundle, $\text{Aut}_{\mathbb{R}}(X, \pi)$ is a linear algebraic group.*

Proof. (1): Any element of $\text{Aut}_{\mathbb{R}}(X)$ fixes any multiple of the anti-canonical divisor, so the claim follows from Lemma 2.10 (1).

(2): Since X is smooth projective and geometrically rational, $\text{Pic}(X)$ is generated by K_X and the real classes of the irreducible components singular fibres of π (or if there are none, the general fibre) of which there are finitely many. Furthermore, because X is geometrically rational, $\text{Pic}(X) \simeq \mathbb{Z}^n$. Let $K := \ker(\text{Aut}_{\mathbb{R}}(X) \rightarrow \text{Aut}(\text{Pic}(X)))$. Then $\text{Aut}_{\mathbb{R}}(X)/K$ is a subgroup of $\text{GL}_n(\mathbb{Z})$ and fixes K_X and the class of the general fibre. It therefore corresponds to a subgroup of permutations of the components of the singular fibres, and is thus a finite group. Lemma 2.10 (3) implies that $\text{Aut}_{\mathbb{R}}(X)$ is a linear algebraic group. \square

REMARK 2.23. It follows from Proposition 2.21 that every infinite algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is contained in the automorphism group of one of the surfaces in Proposition 2.21, which are linear algebraic groups by Lemma 2.22. It now suffices to study the pairs $(X, \text{Aut}_{\mathbb{R}}(X))$ and $(X, \text{Aut}_{\mathbb{R}}(X, \pi))$ for the cases stated in Proposition 2.21 and to determine which automorphism groups are maximal algebraic groups up to conjugacy.

3. Real rational del Pezzo surfaces of degree 6

According to Proposition 2.21, the maximal infinite algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ are contained in the automorphism groups of real del Pezzo surfaces of degree 9, 8 or 6 or the automorphism groups of real conic bundles. In this section, we first classify the real del Pezzo surfaces of degree 6 and give their automorphism groups as explicitly as we dare.

Lemma 3.1. *Let X be a real del Pezzo surface of degree 6.*

- (1) *Then X is the blow-up of $\mathbb{Q}_{3,1}$ or \mathbb{F}_0 in two real or a pair of non-real conjugate points and there are four isomorphism classes, represented in Figures 1, 2, 3 and 4.*
- (2) *The rank of their invariant Picard group is 2, 3, 3 and 4, respectively.*

Proof. The complex surface $X_{\mathbb{C}}$ is the blow-up of three points in \mathbb{P}^2 . It has thus exactly six (-1) -curves, which are arranged as a hexagon on X . The antiholomorphic involution σ on X acts on the hexagon as symmetry of order 2. The only possible cases are shown in Figure 1, 2, 3 and 4, the action of σ indicated by arrows. The second claim follows from the first. \square

Let X be a real Del Pezzo surface of degree 6. There is an exact sequence

$$1 \rightarrow K \rightarrow \text{Aut}_{\mathbb{R}}(X) \xrightarrow{\rho} \text{Aut}(\text{Pic}(X))$$

and K is of finite index, because the action of $\text{Aut}_{\mathbb{R}}(X)$ on $\text{Pic}(X)$ is finite by Lemma 2.10 and Lemma 2.22. The image of $\rho(\text{Aut}_{\mathbb{R}}(X))$ is a subgroup of the dihedral group D_6 acting on the hexagon of (-1) -curves.

3.1. The surfaces obtained by blowing up the sphere. Blowing up the sphere in a pair of non-real conjugate points p, \bar{p} , we obtain the del Pezzo surface $X_{[2]}$ with $\text{rk}(\text{Pic}(X_{[2]})) = 2$. The lift of the antiholomorphic involution is indicated in Figure 1.

REMARK 3.2. For a non-real point $p \in \mathbb{Q}_{3,1}$, we denote by $\text{Aut}_{\mathbb{R}}(\mathbb{Q}_{3,1}, p, \bar{p}) \subset \text{Aut}_{\mathbb{R}}(\mathbb{Q}_{3,1})$ the subgroup of $\text{Aut}_{\mathbb{R}}(\mathbb{Q}_{3,1})$ that fixes both points p and \bar{p} . Choosing $p = ([1 : 0], [0 : 1])$,




Fig.1. $\text{rk}(\text{Pic}(X_{[2]})) = 2$




Fig.2. $\text{rk}(\text{Pic}(X_{[3, Q_{3,1}]})) = 3$




Fig.3. $\text{rk}(\text{Pic}(X_{[3, F_0]})) = 3$




Fig.4. $\text{rk}(\text{Pic}(X_{[4]})) = 4$

it is isomorphic to the group $\{(d, \bar{d}) \in \text{PGL}_2(\mathbb{C}) \times \text{PGL}_2(\mathbb{C}) \mid d \text{ diagonal}\}$ [16, Lemma 4.5]. Conjugating with the real birational map $Q_{3,1} \dashrightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_S)$ from Remark 2.16, we obtain that

$$\text{Aut}_{\mathbb{R}}(Q_{3,1}, p, \bar{p}) \simeq \{(d, \bar{d}) \in \text{PGL}_2(\mathbb{C}) \times \text{PGL}_2(\mathbb{C}) \mid d \text{ diagonal}\} \simeq \mathbb{R}_{>0} \times \text{SO}_2(\mathbb{R}).$$

Proposition 3.3.

(1) The surface $X_{[2]}$ is isomorphic to

$$X_{[2]} \simeq \{([w : x : y : z], [u : v]) \in \mathbb{P}^3 \times \mathbb{P}^1 \mid wz = x^2 + y^2, uz = vw\}.$$

(2) There is an exact sequence

$$1 \rightarrow \ker(\rho) \rightarrow \text{Aut}_{\mathbb{R}}(X_{[2]}) \xrightarrow{\rho} \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \rightarrow 1$$

where $\ker(\rho) \simeq \text{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1}, p, \bar{p}) \simeq \mathbb{R}_{>0} \times \text{SO}_2(\mathbb{R})$ and $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \simeq \langle \rho(\alpha_1) \rangle \times \langle \rho(\alpha_2) \rangle$, where

$$\alpha_1: ([w : x : y : z], [u : v]) \mapsto ([z : -x : y : w], [v : u]),$$

$$\alpha_2: ([w : x : y : z], [u : v]) \mapsto ([w : -x : y : z], [u : v]),$$

and $\rho(\alpha_1)$ is a rotation of order 2 and $\rho(\alpha_2)$ is a reflection, both exchanging E_p and $E_{\bar{p}}$.

- (3) The automorphisms α_1, α_2 are lifts of elements of $\text{Aut}(\mathcal{Q}_{3,1}, \{p, \bar{p}\})$. In particular, the pair $(X_{[2]}, \text{Aut}_{\mathbb{R}}(X_{[2]}))$ is not a minimal pair and the contraction morphism $X_{[2]} \rightarrow \mathcal{Q}_{3,1}$ induces an embedding $\text{Aut}_{\mathbb{R}}(X_{[2]}) \hookrightarrow \text{Aut}(\mathcal{Q}_{3,1})$.

Proof. (1) is Lemma 2.19. Let $S \subset \mathbb{P}^3$ be the surface given by $wz = x^2 + z^2$ and $\psi: S \rightarrow \mathcal{Q}_{3,1}$, $\psi: [w : x : y : z] \mapsto [w + z : 2x : 2y : w - z]$. Any automorphism of $X_{[2]}$ preserves the hexagon in Figure 1 and so $\rho(\text{Aut}_{\mathbb{R}}(X_{[2]}))$ is contained in D_6 . The action of the antiholomorphic involution indicated in Figure 1 implies that any element of $\text{Aut}_{\mathbb{R}}(X_{[2]})$ preserves the set $\{E_p, E_{\bar{p}}\}$. The kernel of ρ is contained in the subgroup of $\text{Aut}_{\mathbb{R}}(X_{[2]})$ fixing E_p and $E_{\bar{p}}$. Any such automorphism descends to an automorphism of S fixing both points p, \bar{p} . Any element of $\text{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1}, p, \bar{p})$ also fixes f_p and $f_{\bar{p}}$ and thus lifts to an element of $\ker(\rho)$. It follows that $\ker(\rho) \simeq \text{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1}, p, \bar{p}) \simeq \mathbb{R}_{>0} \times \text{SO}_2(\mathbb{R})$ (see Remark 3.2 for the isomorphisms).

The only non-trivial elements of D_6 preserving $\{E_p, E_{\bar{p}}\}$ and respecting the action of the antiholomorphic involution are the rotation of order 2 and two reflections, one exchanging $E_p, E_{\bar{p}}$ and one fixing them. The automorphisms

$$\alpha_1: ([w : x : y : z], [u : v]) \mapsto ([z : -x : y : w], [v : u])$$

and

$$\alpha_2: ([w : x : y : z], [u : v]) \mapsto ([w : -x : y : z], [u : v])$$

are the lifts of automorphisms of S exchanging p and \bar{p} and hence exchange E_p and $E_{\bar{p}}$. In fact, via the \mathbb{R} -isomorphism $S \xrightarrow{\varphi\psi^{-1}} (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_S)$, where φ is as in Remark 2.16, α_1 and α_2 are conjugate to

$$\varphi\psi^{-1}\alpha_1\psi\varphi^{-1}: ([x_0 : x_1], [y_0 : y_1]) \mapsto ([x_1 : x_0], [y_1 : y_0])$$

$$\varphi\psi^{-1}\alpha_2\psi\varphi^{-1}: ([x_0 : x_1], [y_0 : y_1]) \mapsto ([y_0 : y_1], [x_0 : x_1])$$

and $\varphi\psi^{-1}(p) = ([0 : 1], [1 : 0])$. This description implies that $\rho(\alpha_1)$ is a rotation of order 2 and $\rho(\alpha_2)$ is the reflection exchanging $E_p, E_{\bar{p}}$. We have $\alpha_1^2 = \alpha_2^2 = (\alpha_1\alpha_2)^2 = \text{Id}$ and hence $\rho: \text{Aut}_{\mathbb{R}}(X_{[4]}) \rightarrow \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is surjective and has a section. On the other hand, it follows that every element of $\text{Aut}_{\mathbb{R}}(X_{[2]})$ is the lift of an element of $\text{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1})$, which yields (3). □

Blowing up the sphere in two real points p, q , we obtain the del Pezzo surface $X_{[3, \mathcal{Q}_{3,1}]}$ with $\text{rk}(\text{Pic}(X_{[3, \mathcal{Q}_{3,1}]})) = 3$. The lift of the antiholomorphic involution is indicated in Figure 2.

Contracting two non-real conjugate (-1) -curves and one real (-1) -curve that are pairwise disjoint, we obtain a birational morphism $X_{[3, \mathcal{Q}_{3,1}]} \rightarrow \mathbb{P}^2$ which is the blow-up of a real point and a pair of non-real conjugate points on \mathbb{P}^2 .

Proposition 3.4.

(1) The surface $X_{[3, \mathcal{Q}_{3,1}]}$ is isomorphic to

$$\{([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid x_0y_0 = x_1y_2 + x_2y_1, x_1y_1 = x_2y_2\}.$$

(2) There is a split exact sequence

$$1 \rightarrow \ker(\rho) \rightarrow \text{Aut}_{\mathbb{R}}(X_{[3, \mathcal{Q}_{3,1}]}) \xrightarrow{\rho} \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \rightarrow 1,$$

where $\ker(\rho) \simeq \text{SO}_2(\mathbb{R})$ and $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \simeq \langle \rho(\alpha_1) \rangle \times \langle \rho(\alpha_2) \rangle$, where

$$\alpha_1 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([y_0 : y_1 : y_2], [x_0 : x_1 : x_2]),$$

$$\alpha_2 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([x_0 : x_2 : x_1], [y_0 : y_2 : y_1]),$$

and $\rho(\alpha_1)$ is a rotation of order 2 and $\rho(\alpha_2)$ a reflection fixing E_p, E_q .

(3) The automorphisms α_1, α_2 are lifts of elements of $\text{Aut}(\mathcal{Q}_{3,1})$. In particular, $(X_{[3, \mathcal{Q}_{3,1}]}, \text{Aut}_{\mathbb{R}}(X_{[3, \mathcal{Q}_{3,1]}}))$ is not a minimal pair, and the contraction $X_{[3, \mathcal{Q}_{3,1}]} \rightarrow \mathcal{Q}_{3,1}$ induces an embedding $\text{Aut}_{\mathbb{R}}(X_{[3, \mathcal{Q}_{3,1}]}) \hookrightarrow \text{Aut}(\mathcal{Q}_{3,1})$.




Fig. 5. The surface $X_{[3, \mathcal{Q}_{3,1}]}$ and the blow-ups ε and η .

Proof. Let $X_{[3, \mathcal{Q}_{3,1}]} \rightarrow \mathcal{Q}_{3,1}$ be the blow-up of two real points p, q on $\mathcal{Q}_{3,1}$. On X there are two pairs of non-real conjugate (-1) -curves – they are the strict transforms of the fibres $f_p, \overline{f_p}$ and $f_q, \overline{f_q}$ passing through p and q (see Figure 2), and by abuse of notation we denote them by $f_p, \overline{f_p}$ and $f_q, \overline{f_q}$ as well. The contraction of the disjoint (-1) -curves $E_p, f_q, \overline{f_q}$ yields a real birational morphism $\varepsilon : X_{[3, \mathcal{Q}_{3,1}]} \rightarrow \mathbb{P}^2$. We call the images of the (-1) -curves r, s, \overline{s} respectively. Composing with an automorphism of \mathbb{P}^2 , we may choose $r = [1 : 0 : 0]$ and $s = [0 : 1 : \mathbf{i}]$. The contraction of the disjoint (-1) -curves $E_q, f_p, \overline{f_p}$ yields a real birational

morphism $\eta: X_{[3, \mathbb{Q}_{3,1}]} \rightarrow \mathbb{P}^2$ and we can assume that they are contracted onto r, s, \bar{s} as well. Our choice implies that the pencil of lines through r, s, \bar{s} respectively is sent onto the pencil of lines through r, s, \bar{s} respectively (see Figure 5). In fact, $\eta\varepsilon^{-1}$ is – up to automorphisms of \mathbb{P}^2 fixing the points r, s, \bar{s} – just the birational involution

$$\eta\varepsilon^{-1}: [x_0 : x_1 : x_2] \mapsto [x_1^2 + x_2^2 : x_0x_2 : x_0x_1].$$

The blow-ups ε and η yield an injection $\varepsilon \times \eta: X_{[3, \mathbb{Q}_{3,1}]} \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$ whose image is described in (1).

The kernel of ρ is isomorphic to the subgroup of $\mathrm{PGL}_3(\mathbb{R})$ fixing the points r, s, \bar{s} , which is isomorphic to $\mathrm{SO}_2(\mathbb{R})$.

Any automorphism of $X_{[3, \mathbb{Q}_{3,1}]}$ preserves the hexagon in Figure 2 and is hence a subgroup of the dihedral group D_6 . The action of the antiholomorphic involution indicated in Figure 2 shows that any automorphism of $X_{[3, \mathbb{Q}_{3,1}]}$ preserves the set $\{E_p, E_q\}$, which means that $\rho(\mathrm{Aut}_{\mathbb{R}}(X_{[3, \mathbb{Q}_{3,1}]}))$ contains, besides the identity map, at most a rotation of order 2 (exchanging E_p, E_q) and two reflections, one fixing E_p and E_q and one exchanging them. The automorphism

$$\alpha_1: ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([y_0 : y_1 : y_2], [x_0 : x_1 : x_2])$$

is the lift of the real birational involution $\eta\varepsilon^{-1}$ of \mathbb{P}^2 and exchanges E_p, E_q and f_p, f_q and so $\rho(\alpha_1)$ is a rotation of order 2. The automorphism

$$\alpha_2: ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([x_0 : x_2 : x_1], [y_0 : y_2 : y_1])$$

is the lift of a linear map of \mathbb{P}^2 exchanging s, \bar{s} and fixing r . It therefore fixes E_p and E_q and exchanges f_p, \bar{f}_p , which means that $\rho(\alpha_2)$ is a reflection. Moreover, $\alpha_3 := \alpha_1\alpha_2 = \alpha_2\alpha_1$ is the reflection exchanging E_p and E_q . The relations $\alpha_1^2 = \alpha_2^2 = (\alpha_2\alpha_1)^2 = \mathrm{Id}$ imply that $\mathrm{Aut}_{\mathbb{R}}(X_{[3, \mathbb{Q}_{3,1}]}) \xrightarrow{\rho} \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is surjective and $\rho(\alpha_i) \mapsto \alpha_i$ is a section of ρ . This yields (2).

The automorphisms α_1, α_2 both preserve the set $\{E_p, E_q\}$ and descend via the contractions of E_p and E_q to automorphisms of S that respectively exchange or fix the points p, q . This yields (3). □

3.2. The surfaces obtained by blowing up \mathbb{F}_0 . Blowing up a pair of non-real conjugate points p, \bar{p} on \mathbb{F}_0 , we obtain the del Pezzo surface $X_{[3, \mathbb{F}_0]}$ with $\mathrm{rk}(\mathrm{Pic}(X_{[3, \mathbb{F}_0]})) = 3$. The lift of the action of the antiholomorphic involution is indicated in Figure 3 by arrows. By $\mathrm{Aut}(\mathbb{F}_0, p, \bar{p}, \mathrm{pr}) \subset \mathrm{Aut}_{\mathbb{R}}(\mathbb{F}_0)$ we denote the subgroup fixing p and \bar{p} and preserving the fibrations.

Proposition 3.5.

(1) *The surface $X_{[3, \mathbb{F}_0]}$ is isomorphic to*

$$\{([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) \in \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \mid x_0y_0z_1 + x_0y_1z_0 + x_1y_0z_0 - x_1y_1z_1 = 0\}.$$

(2) *There is a split exact sequence*

$$0 \rightarrow \ker(\rho) \rightarrow \mathrm{Aut}_{\mathbb{R}}(X_{[3, \mathbb{F}_0]}) \xrightarrow{\rho} D_6 \rightarrow 1$$

where $\ker(\rho) \simeq \mathrm{Aut}_{\mathbb{R}}(\mathbb{F}_0, p, \bar{p}, \mathrm{pr}) \simeq \mathrm{SO}_2(\mathbb{R}) \times \mathrm{SO}_2(\mathbb{R})$.

(3) *The group $\rho(\mathrm{Aut}_{\mathbb{R}}(X_{[3, \mathbb{F}_0]})) \simeq D_6$ is generated by the reflection $\rho(\alpha_1)$ fixing E_p and*

$E_{\bar{p}}$, where

$$\alpha_1 : ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) \mapsto ([y_0 : y_1], [x_0 : x_1], [z_0 : z_1]),$$

and the rotation $\rho(\alpha_2)$ of order 6, where

$$\alpha_2 : ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) \mapsto ([z_1 : z_0], [x_0 : -x_1], [y_1 : y_0]).$$

- (4) The pair $(X_{[3, \mathbb{F}_0]}, \text{Aut}_{\mathbb{R}}(X_{[3, \mathbb{F}_0]}))$ is a minimal pair.
- (5) There is exactly one finite $\text{Aut}_{\mathbb{R}}(X_{[3, \mathbb{F}_0]})$ -orbit on $X_{[3, \mathbb{F}_0]}$, namely the one of the six intersection points of the (-1) -curves.




Fig. 6. The surface $X_{[3, \mathbb{F}_0]}$ and the blow-ups ε and η .

Proof. Let $\varepsilon : X_{[3, \mathbb{F}_0]} \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ be the blow-up of two non real conjugate points in $\mathbb{P}^1 \times \mathbb{P}^1$. We may assume that the points are $p = ([1 : \mathbf{i}], [1 : \mathbf{i}])$ and its conjugate. With this choice of the points, there is a birational morphism $\eta : X \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ which corresponds to the contraction of the fibres g_p and \bar{g}_p (see Figure 6). This yields an injection $\varepsilon \times \eta : X \rightarrow (\mathbb{P}^1)^4$. The fibration given by g (meaning the fibres linearly equivalent to g , drawn punctuated in Figure 6) is preserved by the birational map $\eta\varepsilon^{-1}$ from $\mathbb{P}^1 \times \mathbb{P}^1$ to itself. Composing η with an automorphism of the first factor, we obtain that $\eta\varepsilon^{-1}$ is the identity map on the first factor. Furthermore, we calculate that – up to isomorphism of the second factor – the map $\eta\varepsilon^{-1}$ is given by

$$\eta\varepsilon^{-1} : ([x_0 : x_1], [y_0 : y_1]) \mapsto ([x_0 : x_1], [x_0y_0 + x_1y_1 : x_0y_1 - x_1y_0]).$$

The projection of $(\mathbb{P}^1)^4$ dropping the third factor thus yields an injection $\varphi : X \xrightarrow{\varepsilon \times \eta} (\mathbb{P}^1)^4 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and we get (1) after the isomorphism $y_1 \mapsto -y_1$.

For the second part, the map ρ stands for the induced map coming from the action of $\text{Aut}(X)$ on $\text{Pic}(X)$. As the action of any automorphism of X preserves the hexagon in Figure 3, the image of ρ is contained in the dihedral group D_6 . The image contains the reflections $\rho(\alpha_1)$ and $\rho(\alpha_0)$, where

$$\alpha_1 : ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) \mapsto ([y_0 : y_1], [x_0 : x_1], [z_0 : z_1]),$$

which is the lift of an automorphism exchanging the fibrations of \mathbb{F}_0 and whose image by ρ exchanges f_p and g_p and fixes E_p , and

$$\alpha_0 : ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) \mapsto ([x_0 : -x_1], [z_1 : z_0], [y_1 : y_0]),$$

which is the lift of an automorphism of \mathbb{F}_0 (the one on the right side in Figure 6) exchanging g_p and $\overline{g_p}$ and whose image by ρ exchanges $g_p, \overline{g_p}$ and E_p, f_p . Their composition

$$\alpha_2 := \alpha_1 \alpha_0 : ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) \mapsto ([z_1 : z_0], [x_0 : -x_1], [y_1 : y_0])$$

has order 6. The image $\rho(\alpha_2)$ is the composition of the two reflections $\rho(\alpha_1)$ and $\rho(\alpha_0)$ and hence is a rotation of order 6. The elements $\rho(\alpha_1)$ and $\rho(\alpha_2)$ generate D_6 . This yields the exact sequence (2). Moreover, the relations $\alpha_1^2 = \alpha_2^6 = (\alpha_2 \alpha_1)^2 = \text{Id}$ imply that $\rho(\alpha_i) \mapsto \alpha_i$, $i = 1, 2$ is a section of $\rho : \text{Aut}_{\mathbb{R}}(X_{[3,S_1]}) \rightarrow D_6$ and the sequence splits.

Last but not least, $\ker(\rho)$ is the group of automorphisms of \mathbb{F}_0 fixing the points p and \bar{p} and preserving the fibrations of \mathbb{F}_0 , and which is isomorphic to $\text{SO}_2(\mathbb{R}) \times \text{SO}_2(\mathbb{R})$. Its only finite orbits are its fixed points p, \bar{p} and the intersection points of f_p with $\overline{g_p}$ and of $\overline{f_p}$ with g_p . The group D_6 acts transitively on the intersection points of the (-1) -curves of $X_{[3,\mathbb{F}_0]}$, and this yields (5).

Finally, the description of $\text{Aut}_{\mathbb{R}}(X_{[3,\mathbb{F}_0]})$ in (2)–(3) implies that we cannot contract any (-1) curves on $X_{[3,\mathbb{F}_0]}$ $\text{Aut}_{\mathbb{R}}(X_{[3,\mathbb{F}_0]})$ -equivariantly. In particular, the pair $(X_{[3,\mathbb{F}_0]}, \text{Aut}_{\mathbb{R}}(X_{[3,\mathbb{F}_0]}))$ is a minimal pair. □

Blowing up two real points p, q on \mathbb{F}_0 , we obtain a del Pezzo surface $X_{[4]}$ with $\text{rk}(\text{Pic}(X_{[4]})) = 4$. The lift of the antiholomorphic involution is indicated on Figure 4 by arrows. Blowing down three of the six real (-1) -curves on $X_{[4]}$ yields a birational morphism $X_{[4]} \rightarrow \mathbb{P}^2$ which is the blow-up of three real non-collinear points r_1, r_2, r_3 on \mathbb{P}^2 .

Proposition 3.6.

(1) *The surface $X_{[4]}$ is isomorphic to*

$$\{([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid x_0 y_0 = x_1 y_1 = x_2 y_2\}.$$

(2) *There is a split exact sequence*

$$1 \rightarrow \ker(\rho) \rightarrow \text{Aut}_{\mathbb{R}}(X_{[4]}) \xrightarrow{\rho} D_6 \rightarrow 1$$

where $\ker(\rho) \simeq (\mathbb{R}^*)^2$ is the diagonal subgroup of $\text{PGL}_3(\mathbb{R})$.

(3) *The group $\rho(\text{Aut}_{\mathbb{R}}(X_{[4]})) = D_6$ is generated by the rotation $\rho(\alpha_1)$ and the reflection $\rho(\alpha_2)$, where*

$$\alpha_1 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([y_2 : y_0 : y_1], [x_2 : x_0 : x_1])$$

and

$$\alpha_2 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([x_1 : x_0 : x_2], [y_1 : y_0 : y_2]).$$

(4) *The pair $(X_{[4]}, \text{Aut}_{\mathbb{R}}(X_{[4]}))$ is a minimal pair.*

(5) *There is only one finite $\text{Aut}_{\mathbb{R}}(X_{[4]})$ -orbit on $X_{[4]}$, namely the one of the six intersection points of the (-1) -curves.*

Proof. The surface $X_{[4]}$ is the blow-up of three non-collinear real points in \mathbb{P}^2 and hence isomorphic to $\{([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid x_0y_0 = x_1y_1 = x_2y_2\}$, the blow-up of the real points $[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]$.

The kernel of ρ is isomorphic to the subgroup of $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2) = \text{PGL}_3(\mathbb{R})$ fixing each of the three points $[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]$, which is the diagonal subgroup.

The image by ρ of $\text{Aut}_{\mathbb{R}}(X_{[4]})$ is contained in D_6 because any automorphism of $X_{[4]}$ preserves the hexagon in Figure 4. The involution

$$\beta_1 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([y_0 : y_1 : y_2], [x_0 : x_1 : x_2])$$

is the lift of the standard Cremona involution on \mathbb{P}^2 and hence $\rho(\beta_1)$ is a rotation of order 2. The automorphism

$$\beta_2 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([x_2 : x_0 : x_1], [y_2 : y_0 : y_1])$$

of order 3 is the lift of the automorphism of \mathbb{P}^2 that permutes the three points $[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]$ and so $\rho(\beta_2)$ is a rotation of order 3. Their composition

$$\alpha_1 := \beta_2\beta_1 = \beta_1\beta_2 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([y_2 : y_0 : y_1], [x_2 : x_0 : x_1])$$

is of order 6 and $\rho(\alpha_1)$ is a rotation of order 6 by construction. Furthermore, we find that the involution

$$\alpha_2 : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([x_1 : x_0 : x_2], [y_1 : y_0 : y_2])$$

is the lift of the automorphism of \mathbb{P}^2 that exchanges $[1 : 0 : 0]$ and $[0 : 1 : 0]$ and fixes $[0 : 0 : 1]$, hence $\rho(\alpha_2)$ acts as a reflection. It follows that $\rho(\alpha_1)$ and $\rho(\alpha_2)$ generated D_6 , and we get the exact sequence in (2). Furthermore, the relations $\alpha_1^6 = \alpha_2^2 = (\alpha_1\alpha_2)^2 = \text{Id}$ imply that $\rho(\alpha_i) \mapsto \alpha_i$ is a section of $\rho : \text{Aut}_{\mathbb{R}}(X_{[4]}) \rightarrow D_6$ and the sequence splits.

The above description of $\text{Aut}_{\mathbb{R}}(X_{[4]})$ yields that we cannot contract any (-1) -curve on $X_{[4]}$ $\text{Aut}_{\mathbb{R}}(X_{[4]})$ -equivariantly. In particular, $(X_{[4]}, \text{Aut}_{\mathbb{R}}(X_{[4]}))$ is a minimal pair.

The fact that D_6 acts transitively on the intersection points of the six (-1) -curves and that the only fixed points of $\ker(\rho)$ are the points $[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]$ implies (5). □

4. Pairs of real conic bundles

Recall that for a real conic bundle $\pi : X \rightarrow \mathbb{P}^1$ we denote by $\text{Aut}_{\mathbb{R}}(X, \pi) \subset \text{Aut}_{\mathbb{R}}(X)$ the subgroup of automorphisms respecting the conic bundle structure on X . The morphism π induces a homomorphism $\alpha : \text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow \text{Aut}_{\mathbb{R}}(\mathbb{P}^1) = \text{PGL}_2(\mathbb{R})$ whose kernel we denote by $\ker(\alpha) = \text{Aut}_{\mathbb{R}}(X/\pi)$. We get an exact sequence

$$(*) \quad 1 \rightarrow \text{Aut}_{\mathbb{R}}(X/\pi) \rightarrow \text{Aut}_{\mathbb{R}}(X, \pi) \xrightarrow{\alpha} \text{Aut}_{\mathbb{R}}(\mathbb{P}^1).$$

Proposition 2.21 and Lemma 2.22 imply that a minimal pair (X, G) of a \mathbb{R} -rational G -variety is either a del Pezzo surface of degree 9, 8 or 6 and $G = \text{Aut}_{\mathbb{R}}(X)$, or it admits a real conic bundle $\pi : X \rightarrow \mathbb{P}^1$ with a real birational morphism of conic bundles $X \rightarrow Y$, where Y is the sphere blown up in a pair of non-real conjugate points or a Hirzebruch surface, and $G = \text{Aut}_{\mathbb{R}}(X, \pi)$. In this section, we aim at classifying the real conic bundles $\pi : X \rightarrow \mathbb{P}^1$ that

are relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal.

The following lemma is an adaption of [2, Lemma 4.3.5] to our purpose.

Lemma 4.1. *Suppose $\pi: X \rightarrow \mathbb{P}^1$ is a relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal real conic bundle with a morphism $\eta: X \rightarrow Y$ of real conic bundles where Y is as above, and η is not an isomorphism. Let*

$$G := \text{Aut}_{\mathbb{R}}(X/\pi) \cap (\ker(\text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow \text{Aut}(\text{Pic}(X)))$$

If G is non-trivial, there exists $n \geq 1$ and a (perhaps non-real) birational morphism $X \rightarrow \mathbb{F}_n$ of conic bundles defined over \mathbb{C} that blows up $2n$ points in a section s with $s^2 = n$ which is disjoint from the exceptional section of \mathbb{F}_n , and the strict transform of these two sections are exchanged by an element of $\text{Aut}_{\mathbb{R}}(X, \pi)$.

If G is trivial, then $\text{Aut}_{\mathbb{R}}(X/\pi) \simeq (\mathbb{Z}/2\mathbb{Z})^r$ for $r \in \{0, 1, 2\}$.

Proof. By assumption, η blows up at least one point on Y , hence X has at least one singular fibre. All its singular fibres have exactly two irreducible components because $\pi: X \rightarrow \mathbb{P}^1$ is relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal. Note that G is a normal subgroup of $\text{Aut}(X, \pi)$. Forgetting the antiholomorphic involution on X , we contract in each singular fibre one component and obtain a (perhaps non-real) G -equivariant morphism $\eta': X \rightarrow \mathbb{F}_n$ for some $n \geq 0$. By changing the choice of the components we contract, we obtain $n \geq 1$ and further that η' does not blow-up any points on the exceptional section E_n of \mathbb{F}_n . Let $R := \eta'G(\eta')^{-1} \subset \text{Aut}_{\mathbb{R}}(\mathbb{F}_n)$.

Suppose that G is not trivial. The group R fixes the points blown-up by η' and it preserves E_n . Hence G preserves the strict transform \tilde{E}_n of E_n in X . By construction of η' , the curve \tilde{E}_n intersects exactly one component of each singular fibre of X . The morphism $\pi: X \rightarrow \mathbb{P}^1$ is relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal, so there exists $h \in \text{Aut}_{\mathbb{R}}(X, \pi)$ exchanging the components of singular fibres of X . As G is normal in $\text{Aut}_{\mathbb{R}}(X, \pi)$, then we have $hGh^{-1} = G$. Thus $\text{Fix}(G) = \text{Fix}(hGh^{-1}) = h(\text{Fix}(G))$ contains the section $h(\tilde{E}_n)$, and $h(\tilde{E}_n) \neq \tilde{E}_n$. In particular, R preserves the section $s := \eta'(h(\tilde{E}_n)) \neq E_n$. As $R \subset \text{PGL}_2(\mathbb{C}(t))$ is a non-trivial subgroup, it fixes at most two points on each fibre of \mathbb{F}_n , thus $E_n \cup s = \text{Fix}(R)$ and s contains all points blown up by η' . Further, the two curves \tilde{E}_n and $h(\tilde{E}_n)$ have the same self-intersection, which is equal to $-n$ because η' does not blow up any points on E_n . Moreover, since the elements of $R \subset \text{PGL}_2(\mathbb{C}(t))$ fix exactly two points on all but finitely many fibres, they are diagonalisable and hence fix two points on every fibre. It follows that E_n and s are disjoint. Hence $s \sim E_n + nf$ and therefore $s^2 = n$. It follows that η' is the blow-up of $2n$ points on s .

Suppose that G is trivial. Then every non-trivial element of $\text{Aut}_{\mathbb{R}}(X/\pi)$ is an involution. As $\text{Aut}_{\mathbb{R}}(X/\pi) \subset \text{PGL}_2(\mathbb{C}(x))$, it follows that $\text{Aut}_{\mathbb{R}}(X/\pi)$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^r$ for $r \in \{0, 1, 2\}$. □

4.1. Real conic bundles obtained by blowing up a del Pezzo surface. In this subsection, we study the ones with a real birational morphism of conic bundles $\eta: X \rightarrow Y$ to the surface Y obtained by blowing up the sphere in a pair of non-real conjugate points. The surface Y is a del Pezzo surface of degree 6 with $\text{rk}(\text{Pic}(Y)) = 2$ and so, by Lemma 3.1,

$$\text{pr}_1: Y = X_{[2]} \rightarrow \mathcal{Q}_{3,1}$$

is isomorphic to the blow-up of the points $r := [0 : 1 : \mathbf{i} : 0]$ and $\bar{r} = [0 : 1 : -\mathbf{i} : 0]$ (see Proposition 3.3). The generic fibre of the projection

$$\text{pr}: X_{[2]} \rightarrow \mathbb{P}^1, \quad ([w : x : y : z], [u : v]) \mapsto [u : v]$$

is the non-rational conic $C \subset \mathbb{P}_{\mathbb{R}(t)}^2$ given by $x^2 + y^2 - tz^2 = 0$, which makes $\text{pr}: X_{[2]} \rightarrow \mathbb{P}^1$ a conic bundle. However, the description in Lemma 4.2 of the conic bundle $(X_{[2]}, \text{pr})$ will turn out to be more convenient.

As described in Remark 2.16, the isomorphism of complex surfaces $\varphi: \mathcal{Q}_{3,1} \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$,

$$\begin{aligned} \varphi: [w : x : y : z] &\longmapsto ([w + z : y + \mathbf{i}x], [w + z : y - \mathbf{i}x]) \\ &= ([y - \mathbf{i}x : w - z], [y + \mathbf{i}x : w - z]) \end{aligned}$$

$$\varphi^{-1}: ([x_0 : x_1], [y_0 : y_1]) \mapsto [x_0y_0 + x_1y_1 : \mathbf{i}(x_0y_1 - x_1y_0) : x_0y_1 + x_1y_0 : x_0y_0 - x_1y_1]$$

induces an isomorphism of real surfaces $\varphi: \mathcal{Q}_{3,1} \rightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_S)$, where

$$\sigma_S: ([x_0 : x_1], [y_0 : y_1]) \mapsto ([\bar{y}_0 : \bar{y}_1], [\bar{x}_0 : \bar{x}_1])$$

and $p := \varphi(r) = ([0 : 1], [1 : 0])$ and $\bar{p} = \varphi(\bar{r}) = ([1 : 0], [0 : 1])$.

Lemma 4.2.

(1) *The real surface $X_{[2]}$ is isomorphic to*

$$(X_{[2]}, \sigma) \simeq \left(\{([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid x_0y_0 = x_1y_1 = x_2y_2\}, \sigma_{[2]}\right)$$

where $\sigma_{[2]}: ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([\bar{y}_1 : \bar{y}_0 : \bar{y}_2], [\bar{x}_1 : \bar{x}_0 : \bar{x}_2])$ and the conic bundle structure $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$ is given by

$$\pi_{[2]}: ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \longmapsto [x_0 : x_1] = [y_1 : y_0].$$

(2) *The irreducible components of the singular fibres of $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$ are given by*

$$f_p: y_1 = y_2 = 0, \quad \bar{f}_p: x_0 = x_2 = 0, \quad f_{\bar{p}}: x_1 = x_2 = 0, \quad \bar{f}_{\bar{p}}: y_0 = y_2 = 0$$

(see Figure 7) and the pair of non-real conjugate (-1) -“sections” by

$$s: x_0 = x_1 = 0, \quad \bar{s}: y_0 = y_1 = 0.$$

Proof. Over \mathbb{C} , there is only one del Pezzo surface of degree 6 and it is isomorphic to

$$Z := \{([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid x_0y_0 = x_1y_1 = x_2y_2\}$$

(see Proposition 3.6). The abstract birational morphism

$$\begin{aligned} Z &\longrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \\ ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) &\longmapsto ([x_0 : x_2], [x_2 : x_1]) = ([y_2 : y_0], [y_1 : y_2]) \\ ([ru : sv : su], [sv : ru : rv]) &\longleftarrow ([r : s], [u : v]) \end{aligned}$$

contracts the (-1) -curves $s_1 = \{x_0 = x_1 = 0\}$ and $s_2 = \{y_0 = y_1 = 0\}$ onto p and \bar{p} , respectively. In Figure 1, they are therefore denoted by E_p and $E_{\bar{p}}$. The lift of the antiholomorphic involution σ_S onto Z is $\sigma_{[2]}$ and makes $(Z, \sigma_{[2]})$ a real del Pezzo surface isomorphic to $X_{[2]}$ (Lemma 3.1) and $s_2 = \bar{s}_1$. The morphism

$$\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1, \quad ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \longmapsto [x_0 : x_1] = [y_1 : y_0]$$

is the projection onto the (-1) -curves s and \bar{s} and is thus a conic bundle. The antiholomorphic

involution $\sigma_{[2]}$ descends to the standard antiholomorphic involution $[u : v] \mapsto [\bar{u} : \bar{v}]$ on \mathbb{P}^1 , which makes $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$ a real conic bundle. The morphisms are visualised in Figure 7. The equations of the irreducible components of the singular fibres are checked by calculation.

By abuse of notation, we will denote the surface Z by $X_{[2]}$ endowed with $\sigma_{[2]}$. □




Fig. 7. The real conic bundle $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$.

The following lemma gives a necessary condition for a real conic bundle $\pi: X \rightarrow \mathbb{P}^1$ to be relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal. It will turn out in Proposition 4.5 that the condition is also sufficient. A reference picture is drawn in Figure 8.

Lemma 4.3. *Let $\pi: X \rightarrow \mathbb{P}^1$ be a relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal real conic bundle with a birational morphism of real conic bundles $\eta: X \rightarrow X_{[2]}$ that is not an isomorphism. Then $\text{Aut}_{\mathbb{R}}(X, \pi)$ is finite, or $\eta: X \rightarrow X_{[2]}$ is the blow-up of $n \geq 1$ pairs of non-real conjugate points of $X_{[2]}$ contained in $s \cup \bar{s}$ and in non-real fibres.*

Proof. The morphism η blows up at least one point, so X has at least four singular fibres. Let $G := \text{Aut}_{\mathbb{R}}(X/\pi) \cap (\text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow \text{Aut}(\text{Pic}(X)))$.

Suppose that G is non-trivial. By Lemma 4.1, there exists a non-real birational morphism $\eta': X \rightarrow \mathbb{F}_n, n \geq 1$, of conic bundles defined over \mathbb{C} which blows up $2n \geq 2$ points on a section s' disjoint from the exceptional section E_n of \mathbb{F}_n with $(s')^2 = n$.

Denote by \tilde{s}' and \tilde{E}_n the strict transforms of s' and E_n respectively. Note that they are the unique $(-n)$ -curves on X and hence are two real or a pair of non-real conjugate curves. They descend via $\eta: X \rightarrow X_{[2]}$ onto curves c_1 and c_2 on $X_{[2]}$. As $X_{[2]}$ does not have any real sections, we get $c_2 = \bar{c}_1$, and hence $\tilde{s}' = \tilde{E}_n$. In particular, the real morphism η contracts $n - 1$ components of singular fibres only intersecting \tilde{s}' and $n - 1$ components of singular fibres only intersecting \tilde{E}_n . In other words, η blows up $n \geq 1$ pairs of non-real conjugate points contained in $c_1 \cup \bar{c}_2$, no two on the same fibre, so $c_1^2 = -1$. In particular, $c_1 \cup \bar{c}_1 = s \cup \bar{s}$.

Suppose that G is trivial. Lemma 4.1 implies that $\text{Aut}_{\mathbb{R}}(X/\pi)$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^r$ for $r \in \{0, 1, 2\}$. The group $\text{Aut}_{\mathbb{R}}(X, \pi)$ preserves $X(\mathbb{R})$, hence its image H in $\text{PGL}_2(\mathbb{R})$ preserves

$\pi(X(\mathbb{R})) = \pi_{[2]}(X_{[2]}(\mathbb{R})) = [0, \infty]$, i.e.

$$H \subset \text{Aut}_{\mathbb{R}}(\mathbb{P}^1, [0, \infty]) \simeq \mathbb{R}_{>0} \times \mathbb{Z}/2\mathbb{Z}$$

Furthermore, H preserves the set of the images in \mathbb{P}^1 of the singular fibres of X , of which there are at least four. This implies that H is finite. As $\text{Aut}_{\mathbb{R}}(X/\pi)$ and H are both finite, also $\text{Aut}_{\mathbb{R}}(X, \pi)$ is finite. □

DEFINITION 4.4(AND CONSTRUCTION). The abstract birational morphism

$$\begin{aligned} \varepsilon: X_{[2]} &\longrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \\ ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) &\longmapsto ([x_0 : x_1], [x_2 : x_0]) = ([y_1 : y_0], [y_0 : y_2]) \\ ([u_0 v_1 : u_1 v_1 : u_0 v_0], [u_1 v_0 : u_0 v_0 : u_1 v_1]) &\longleftarrow ([u_0 : u_1], [v_0, v_1]) \end{aligned}$$

contracts the components f_p and $f_{\bar{p}}$ of the singular fibres onto the points $([0 : 1], [1 : 0])$ and $([1 : 0], [0 : 1])$, respectively, and the sections s, \bar{s} onto the sections $\varepsilon(s) = \mathbb{P}^1 \times \{[1 : 0]\}$ and $\varepsilon(\bar{s}) = \mathbb{P}^1 \times \{[0 : 1]\}$. The antiholomorphic involution $\sigma_{[2]}$ descends to a rational antiholomorphic involution

$$\sigma_C: ([u_0 : u_1], [v_0 : v_1]) \mapsto ([\overline{u_0} : \overline{u_1}], [\overline{u_1 v_1} : \overline{u_0 v_0}])$$

on $\mathbb{P}^1 \times \mathbb{P}^1$, not defined at $([1 : 0], [0 : 1])$ and $([0 : 1], [1 : 0])$. It makes ε a real birational morphism of conic bundles, i.e. the diagram

$$\begin{array}{ccc} (X_{[2]}, \sigma_{[2]}) & \xrightarrow{\varepsilon} & (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C) \\ \pi_{[2]} \searrow & & \swarrow \text{pr}_1 \\ & \mathbb{P}^1 & \end{array}$$

is commutative. The construction is visualised in Figure 8.




Fig. 8. The real birational morphism $\varepsilon: (X_{[2]}, \sigma_{[2]}) \rightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C)$.

Proposition 4.5. *Let $\eta: X \rightarrow X_{[2]}$ be the blow-up of $n \geq 1$ pairs of non-real conjugate points in $s \cup \bar{s}$ and in non-real fibres. Then $\pi := \pi_{[2]}\eta: X \rightarrow \mathbb{P}^1$ is a relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal real conic bundle.*

Let $\Delta \subset \mathbb{P}^1$ be the image of the $2n + 2$ singular fibres of X and $H_{\Delta} \subset \text{PGL}_2(\mathbb{R})$ be the subgroup preserving Δ and $\pi(X(\mathbb{R})) = [0, \infty]$. Then

(1) *there exists a split exact sequence*

$$1 \rightarrow \text{Aut}(X/\pi) \rightarrow \text{Aut}(X, \pi) \rightarrow H_{\Delta} \rightarrow 1$$

where $\text{Aut}(X/\pi) \simeq \text{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1}, p, \bar{p}) \rtimes \mathbb{Z}/2\mathbb{Z} \simeq \text{SO}_2(\mathbb{R}) \rtimes \mathbb{Z}/2\mathbb{Z}$,

- (2) *an element of $\text{SO}_2(\mathbb{R}) \subset \text{Aut}(X/\pi)$ fixes the two $-(n + 1)$ -sections of X and the generator of $\mathbb{Z}/2\mathbb{Z}$ exchanges them,*
- (3) *an element of $\text{Aut}(X/\pi) \setminus \text{SO}_2(\mathbb{R})$ is an involution fixing an irreducible curve on $X_{[2]}$ which is a double cover of \mathbb{P}^1 ramified at Δ ,*
- (4) *the group $\text{SO}_2(\mathbb{R})$ acts trivially on $\text{Pic}(X)$.*

Proof. Any automorphism of X preserves the set of real points $X(\mathbb{R})$, which is diffeomorphic via η to $X_{[2]}(\mathbb{R})$ and is mapped to the interval $[0, \infty]$ by π . Therefore, the exact sequence (*) yields the exact sequence (1). Any element of H_{Δ} lifts to a real automorphism of $X_{[2]}$ fixing the points blown-up by η and thus lifts to an automorphism of X . The sequence splits.

Over \mathbb{C} , there is a birational morphism $X \xrightarrow{\eta} X_{[2]} \rightarrow \mathbb{F}_0$, hence X has exactly two $-(n + 1)$ -sections [2, Lemma 4.3.1], and they are the strict transforms s', \bar{s}' of the (-1) -curves s, \bar{s} on $X_{[2]}$. So $\text{Aut}_{\mathbb{R}}(X/\pi)$ acts on $\{s', \bar{s}'\}$ and we claim that it acts non-trivially; we now construct a birational involution of $X_{[2]}$ whose lift onto X is an automorphism respecting π and exchanging s', \bar{s}' .

Let $q_1, \dots, q_n \in s$ and $\bar{q}_1, \dots, \bar{q}_n \in \bar{s}$ be the points blown up by η , and define $p_i := \pi_{[2]}(q_i)$. Let $\varepsilon: (X_{[2]}, \sigma_{[2]}) \rightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C)$ be the birational morphism of real conic bundles given in Definition 4.4. Then $\varepsilon(q_i) = (p_i, [1 : 0])$ and $\varepsilon(\bar{q}_i) = (\bar{p}_i, [0 : 1])$. Let $m_1, \dots, m_n \in \mathbb{C}[u_0, u_1]$ be homogenous linear polynomials vanishing on p_1, \dots, p_n respectively and define $P(u_0, u_1) := \prod_{i=1}^n m_i(u_0, u_1)$. The involution $\varphi: \mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$,

$$\varphi: ([u_0 : u_1], [v_0 : v_1]) \dashrightarrow ([u_0 : u_1], [u_1 v_1 \bar{P}(u_0, u_1) : u_0 v_0 P(u_0, u_1)])$$

commutes with the antimeromorphic involution σ_C and is undefined exactly at $\varepsilon(q_1), \dots, \varepsilon(q_n), \varepsilon(\bar{q}_1), \dots, \varepsilon(\bar{q}_n), ([1 : 0], [0 : 1]), ([0 : 1], [1 : 0])$ and exchanges $\varepsilon(s)$ and $\varepsilon(\bar{s})$. The map φ is visualised in Figure 9.

The involution φ thus lifts via ε to a real birational involution of $X_{[2]}$ that exchanges s, \bar{s} and is undefined exactly at $q_1, \dots, q_n, \bar{q}_1, \dots, \bar{q}_n$. So, it lifts to a real automorphism of X that exchanges s', \bar{s}' . Therefore $\text{Aut}_{\mathbb{R}}(X/\pi)$ acts non-trivially on the set $\{s', \bar{s}'\}$, which yields the split exact sequence

$$1 \rightarrow K \rightarrow \text{Aut}_{\mathbb{R}}(X/\pi) \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 0.$$

This also shows that we cannot contract any components of the singular fibres $\text{Aut}_{\mathbb{R}}(X/\pi)$ -equivariantly, and hence also not $\text{Aut}_{\mathbb{R}}(X, \pi)$ -equivariantly. In particular, $\pi: X \rightarrow \mathbb{P}^1$ is relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal.

By definition of K , all of its elements fix s'_1, s'_2 and thus descend to a subgroup of $\text{Aut}_{\mathbb{R}}(X_{[2]}/\pi_{[2]})$ and hence via $\text{pr}_1: X_{[2]} \rightarrow \mathcal{Q}_{3,1}$ to a subgroup of $\text{Aut}(\mathcal{Q}_{3,1}, p, \bar{p})$, the automor-

phism group of $\mathcal{Q}_{3,1}$ fixing p and \bar{p} , which is isomorphic to $\{(A, \bar{A}) \in \mathrm{PGL}_2(\mathbb{C})^2 \mid A \text{ diagonal}\}$ via $\mathcal{Q}_{3,1} \simeq (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_S)$ [16, Lemma 4.5]. On the other hand, any $(\mathrm{diag}(a, 1), \mathrm{diag}(\bar{a}, 1)) \in \mathrm{Aut}(\mathcal{Q}_{3,1}, p, \bar{p})$ lifts to the real automorphism

$$\beta_a : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([a\bar{a}x_0 : x_1 : \bar{a}x_2], [y_0 : a\bar{a}y_1 : ay_2])$$

of $X_{[2]}$ which fixes s and \bar{s} . On them, it acts by $[y_0 : y_1] \mapsto [y_0 : a\bar{a}y_1]$ and $[x_0 : x_1] \mapsto [a\bar{a}x_0 : x_1]$, respectively. Hence, β_a lifts to an automorphism of X if and only if it fixes the points blown up by η (there is at least one), which is equivalent to $a\bar{a} = 1$. The lift of β_a on X then descends via $\pi_{[2]}$ to the identity map on \mathbb{P}^1 . It follows that $(\mathrm{diag}(a, 1), \mathrm{diag}(\bar{a}, 1)) \in \mathrm{Aut}_{\mathbb{R}}(\mathcal{Q}_{3,1}, p, \bar{p})$ is contained in $\mathrm{Aut}(X/\pi)$ if and only if $a\bar{a} = 1$, which implies that $K = \{(\mathrm{diag}(a, 1), \mathrm{diag}(\bar{a}, 1)) \in \mathrm{PGL}_2(\mathbb{C})^2 \mid a\bar{a} = 1\}$. Conjugating K with the real isomorphism $\mathcal{Q}_{3,1} \rightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_S)$ from Remark 2.16 yields $K \simeq \mathrm{SO}_2(\mathbb{R})$. This finishes the proof of (1) and yields (2), (4) and the first half of (3).

The group $\mathrm{SO}_2(\mathbb{R})$ acts via ε on $(\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C)$ by

$$([u_0 : u_1], [v_0 : v_1]) \mapsto ([u_0 : u_1], [v_0 : av_1]).$$

It follows that $(a, \varphi) \in \mathrm{Aut}(X/\pi) \setminus \mathrm{SO}_2(\mathbb{R})$ is an involution on $(\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C)$ fixing the irreducible curve $au_0v_0^2P(u_0, u_1) - u_1v_1^2\bar{P}(u_0, u_1) = 0$, which is an irreducible double cover of \mathbb{P}^1 ramified at Δ and $([1 : 0], [0 : 1]), ([0 : 1], [1 : 0])$. Its strict transform on $X_{[2]}$ is an irreducible double cover over \mathbb{P}^1 ramified over Δ . This yields the second part of (3). \square

REMARK 4.6. Note that the generator of $\mathbb{Z}/2\mathbb{Z}$ in the description of $\mathrm{Aut}(X, \pi)$ in Proposition 4.5 is the composition of elementary links of the real conic bundle $\pi_{[2]} : X_{[2]} \rightarrow \mathbb{P}^1$, each blowing up a pair of non-real points on $s \cup \bar{s}$.




Fig. 9. The real birational map $\varphi : (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C) \rightarrow (\mathbb{P}^1 \times \mathbb{P}^1, \sigma_C)$ that lifts to an automorphism of X generating $\mathbb{Z}/2\mathbb{Z} \subset \mathrm{Aut}_{\mathbb{R}}(X/\pi)$

4.2. Real conic bundles obtained by blowing up a Hirzebruch surface. The n -th Hirzebruch surface is given by

$$\mathbb{F}_n = \{([x_0 : x_1 : x_2], [u : v]) \in \mathbb{P}^2 \times \mathbb{P}^1 \mid u^n x_2 = v^n x_1\} \subset \mathbb{P}^2 \times \mathbb{P}^1.$$

The canonical projection $\text{pr}_n : \mathbb{F}_n \rightarrow \mathbb{P}^1$ onto the second factor makes it a real conic bundle. In this subsection, we study real conic bundles with a birational morphism $X \rightarrow \mathbb{F}_n$ of real conic bundles.

If $n = 0$, then $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$ with the standard antiholomorphic involution on it, and $\text{Aut}_{\mathbb{R}}(\mathbb{F}_0, \text{pr}_0) = \text{PGL}_2(\mathbb{R})^2$.

If $n = 1$, then \mathbb{F}_1 is isomorphic to the blow-up of $[1 : 0 : 0] \in \mathbb{P}^2$, and any automorphism preserves the unique (-1) -curve on it, which yields $\text{Aut}_{\mathbb{R}}(\mathbb{F}_1) = \text{Aut}_{\mathbb{R}}(\mathbb{F}_1, \text{pr}_1) \simeq \text{Aut}_{\mathbb{R}}(\mathbb{P}^2, [1 : 0 : 0])$.

If $n \geq 2$, the automorphism group of \mathbb{F}_n is

$$\text{Aut}_{\mathbb{R}}(\mathbb{F}_n) \simeq \mathbb{R}^{n+1} \rtimes \text{GL}_2(\mathbb{R})/\mu_n,$$

where $\mu_n = \{\mu \text{Id} \mid \mu^n = 1\}$ [2]. An element

$$\left((a_0, \dots, a_n), \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) \in \mathbb{R}^{n+1} \rtimes \text{GL}_2(\mathbb{R})/\mu_n$$

acts on the chart $u \neq 0$ by

$$([xu^n : yu^n : yv^n], [u : v]) \mapsto ([xu^n + y(a_0v^n + a_1uv^{n-1} + \dots + a_nv^n) : y(au + bv)^n : y(cu + dv)^n], [au + bv : cu + dv])$$

and in particular respects the conic bundle structure on \mathbb{F}_n . Multiples of the identity matrix act trivially on the base, and we get

$$\text{Aut}_{\mathbb{R}}(\mathbb{F}_n/\text{pr}_n) \simeq \mathbb{R}^{n+1} \rtimes (\mathbb{R}^*/\mu_n),$$

where we see $\mu_n \subset \mathbb{R}^*$ as $\mu_n = \{\pm 1\}$ if n is even and $\mu_n = \{1\}$ if n is odd.

We denote by $E_n := \{([1 : 0 : 0], [u : v]) \mid [u : v] \in \mathbb{P}^1\} \subset \mathbb{F}_n$ its $(-n)$ -section, by f the general fibre of pr_n and by $s_n \subset \mathbb{F}_n$ the section given by $x_0 = 0$, i.e.

$$s_n := \{([0 : u^n : v^n], [u : v]) \mid [u : v] \in \mathbb{P}^1\}.$$

The conditions $s_n E_n = 0$ and $s_n f = 1$ yield $s_n \sim E_n + n f$ as divisors and hence $s_n^2 = n$.

Let us give a necessary description of minimal pairs $(X, \text{Aut}_{\mathbb{R}}(X, \pi))$ equipped with a birational morphism $\eta : X \rightarrow \mathbb{F}_n$ of real conic bundles.

Lemma 4.7. *Let $\pi : X \rightarrow \mathbb{P}^1$ be a relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal real conic bundle equipped with a brational morphism $\eta : X \rightarrow \mathbb{F}_n$ of real conic bundles that is not an isomorphism.*

If $\text{Aut}_{\mathbb{R}}(X/\pi) \cap \ker(\text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow \text{Aut}(\text{Pic}(X)))$ contains a non-trivial element, then there exists a birational morphism $X \rightarrow \mathbb{F}_N$ of real conic bundles blowing up $2N \geq 2$ points, all contained in s_N with pairwise distinct fibres.

Else, $\text{Aut}_{\mathbb{R}}(X/\pi) \simeq (\mathbb{Z}/2\mathbb{Z})^r$ for $r \in \{0, 1, 2\}$.

Proof. If $G := \text{Aut}_{\mathbb{R}}(X/\pi) \cap \ker(\text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow \text{Aut}(\text{Pic}(X)))$ is trivial, the claim follows from Lemma 4.1. Suppose that G is non-trivial. By Lemma 4.1, there exists a (perhaps non-real) birational morphism $X \rightarrow \mathbb{F}_N$, $N \geq 1$, blowing up $2N$ points on a section s of \mathbb{F}_N disjoint from E_N and of self-intersection N . By c_1 and c_2 we denote the strict transforms of s and E_N in X . The real birational morphism η blows down one component of each fibre. Furthermore, c_1 is sent onto a curve of self-intersection $N - r$, where r is the number of contracted components intersecting $\eta(c_1)$, and $\eta(c_2)$ is a curve of self-intersection $N - (2N - r) = -N + r$. We can assume that $r \leq N$ (else we exchange the indices of c_1 and c_2). Then $\eta(c_2)^2 \leq 0$, hence $\eta(c_2)$ is the exceptional section of \mathbb{F}_n . In particular, it is a real curve, hence also c_2 is a real curve. By Lemma 4.1 there is an element of $\text{Aut}_{\mathbb{R}}(X, \pi)$ exchanging c_1 and c_2 , thus c_1 is a real curve as well. Therefore, only contracting components intersecting c_1 commutes with the antiholomorphic involution of X . It follows that the birational morphism $X \rightarrow \mathbb{F}_N$ from Lemma 4.1 is in fact a real morphism. \square

In the first assertion of the lemma, the cases $N = 0$ and $N = 1$ yield relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal conic bundles, but the groups $\text{Aut}_{\mathbb{R}}(X, \pi)$ are not maximal algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.

Proposition 4.8. *Let $\eta: X \rightarrow \mathbb{F}_n$ be the blow-up of $2n \geq 4$ points contained in s_n . Then $\pi := \text{pr}_n \eta: X \rightarrow \mathbb{P}^1$ is a relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal conic bundle.*

Let $\Delta \subset \mathbb{P}^1$ be the projection of the points blown up by η , $H_{\Delta} \subset \text{PGL}_2(\mathbb{R})$ the subgroup preserving Δ and $\mu_n = \{\pm 1\}$ if n is even and $\mu_n = \{1\}$ if n is odd. Then:

- (1) *There is a split exact sequence*

$$1 \rightarrow \text{Aut}_{\mathbb{R}}(X/\pi) \rightarrow \text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow H_{\Delta} \rightarrow 1$$

where $\text{Aut}_{\mathbb{R}}(X/\pi) \simeq (\mathbb{R}^/\mu_n) \rtimes \mathbb{Z}/2\mathbb{Z}$.*

- (2) *An element of $(\mathbb{R}^*/\mu_n) \subset \text{Aut}(X/\pi)$ fixes the two $(-n)$ -sections of X and the generator of $\mathbb{Z}/2\mathbb{Z}$ exchanges them.*
- (3) *An element of $\text{Aut}(X/\pi) \setminus (\mathbb{R}^*/\mu_n)$ is an involution fixing an irreducible curve on \mathbb{F}_n which is a double cover of \mathbb{P}^1 ramified at Δ .*
- (4) *The group \mathbb{R}^*/μ_n acts trivially on $\text{Pic}(X)$.*

Proof. Denote by $\tilde{s}_n, \tilde{E}_n \subset X$ the strict transforms of s_n and E_n , respectively. Then $\pi := \text{pr} \eta: X \rightarrow \mathbb{P}^1$ is a conic bundle with $2n$ singular fibres, whose components either intersect \tilde{s}_n or \tilde{E}_n . The action of $\text{Aut}_{\mathbb{R}}(X, \pi)$ descends to an action on \mathbb{P}^1 that preserves the set Δ . On the other hand, any element of H_{Δ} lifts to an automorphism of \mathbb{F}_n preserving the set of the points blown-up by η and hence lifts to an automorphism of X that permutes the singular fibres. Thus the sequence splits.

The group $\text{Aut}_{\mathbb{R}}(X/\pi)$ acts on the set $\{\tilde{E}_n, \tilde{s}_n\}$ non-trivially: We can assume that the points blown up by η are in the chart $u = 1$. They are thus of the form $p_i = ([0 : 1 : v_i^n], [1 : v_i]) \in s_n$, $i = 1, \dots, 2n$. The number of non-real points in Δ is even, hence the number of real points in Δ is even as well. We order the points such that p_1, \dots, p_{2k} are real points and $p_{2k+1}, p_{2k+2} := \overline{p_{2k+1}}, \dots, p_{2n-1}, p_{2n} := \overline{p_{2n-1}}$ are pairs of non-real conjugate points. For $i = 1, \dots, 2k$, let $l_i := t - v_i^n \in \mathbb{R}[t]$. For $i = 1, \dots, n - k$, let $m_i := (t - v_{2(k+i)-1}^n)(t - v_{2(k+i)}^n) \in \mathbb{R}[t]$. We define

$$P(r) := \prod_{i=1}^k l_i \prod_{i=1}^{n-k} m_i \in \mathbb{R}[t].$$

Then the rational map $\varphi: \mathbb{F}_n \dashrightarrow \mathbb{F}_n$ given on the chart $u = 1$ by

$$\varphi: ([x_0 : x_1 : x_1 v^n], [1 : v]) \mapsto ([x_1 P(v) : x_0 : x_0 v^n], [1 : v])$$

is a real involution respecting the conic bundle structure of \mathbb{F}_n . It is undefined exactly at the points p_1, \dots, p_{2n} and exchanges E_n and s_n . It furthermore contracts the fibre through p_i onto p_i . The map φ is visualised in Figure 10.

Therefore, φ lifts to an automorphism of the conic bundle X that exchanges \tilde{s}_n and \tilde{E}_n . This induces the split exact sequence

$$1 \rightarrow K \rightarrow \text{Aut}_{\mathbb{R}}(X/\pi) \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 1.$$

It also proves that we cannot contract any components of the singular fibres on X $\text{Aut}_{\mathbb{R}}(X/\pi)$ -equivariantly and thus also not $\text{Aut}_{\mathbb{R}}(X, \pi)$ -equivariantly. In particular, $\pi: X \rightarrow \mathbb{P}^1$ is relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal.

By definition of K , all of its elements fix \tilde{s}_n and \tilde{E}_n pointwise and thus descend to elements of $\text{Aut}_{\mathbb{R}}(\mathbb{F}_n/\text{pr})$. On the other hand, an element $((a_0, \dots, a_n), r) \in \text{Aut}_{\mathbb{R}}(\mathbb{F}_n/\text{pr}_n) \simeq \mathbb{R}^{n+1} \rtimes (\mathbb{R}^* \text{Id})/\mu_n$ acts on the chart $u \neq 0$ by

$$([x_0 : x_1 : v^n x_1], [1 : v]) \mapsto ([x_0 + x_1(a_n v^n + a_{n-1} v^{n-1} + \dots + a_0) : r^n x_1 : r^n v^n x_1], [1 : v])$$

and lifts to an automorphism of X if and only if it preserves s_n (which is given by $x_0 = 0$). It follows that $K = \mathbb{R}^*/\mu_n$. This completes (1) and (2).

Every element $(r, \varphi) \in \text{Aut}_{\mathbb{R}}(X/\pi) \setminus (\mathbb{R}^*/\mu_n)$ fixes the curve $x_1^2 P(v) - r^n x_0^2 = 0$ (given on the chart $u \neq 0$), which is a double cover of \mathbb{P}^1 ramified over Δ . This is (3).

Finally, the action of $\mathbb{R}^*/\mu_n \subset \text{Aut}_{\mathbb{R}}(X/\pi)$ fixes each fibre and it fixes E_n and s_n . Hence it acts trivially on $\text{Pic}(X)$, which is (4). □




Fig. 10. The real birational map $\varphi: \mathbb{F}_n \dashrightarrow \mathbb{F}_n$ that lifts to an automorphism of X .

The following lemma is an adapted version of Lemma [2, Lemma 5.2.1].

Lemma 4.9. *Let $\pi: X \rightarrow \mathbb{P}^1$ be a relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal conic bundle with a birational morphism $\eta: X \rightarrow \mathbb{F}_n$ of real conic bundles. Suppose that $\text{Aut}_{\mathbb{R}}(X/\pi) \cap \ker(\text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow \text{Aut}_{\mathbb{R}}(\text{Pic}(X))) = \{1\}$. Then $\text{Aut}_{\mathbb{R}}(X, \pi)$ is finite or strictly contained in the automorphism group of a real del Pezzo surface.*

Proof. Suppose X has at least three singular fibres. The action of $\text{Aut}_{\mathbb{R}}(X, \pi)$ on \mathbb{P}^1 induces the exact sequence

$$1 \rightarrow \text{Aut}_{\mathbb{R}}(X/\pi) \rightarrow \text{Aut}_{\mathbb{R}}(X, \pi) \rightarrow H,$$

where $H \subset \text{Aut}_{\mathbb{R}}(\mathbb{P}^1)$ is the subgroup fixing the image of the set of the singular fibres. The conic bundle X having three singular fibres implies that H is finite. By Lemma 4.7, $\text{Aut}_{\mathbb{R}}(X/\pi)$ is finite as well, so $\text{Aut}_{\mathbb{R}}(X, \pi)$ is finite.

Suppose X has one or two singular fibres. Let \tilde{E}_n be the strict transform of the $-n$ curve of \mathbb{F}_n . It is of self-intersection $-r \leq -n$. As the pair $(X, \text{Aut}_{\mathbb{R}}(X, \pi))$ is minimal, the singular fibres intersect \tilde{E}_n , have exactly two components and there exists $g \in \text{Aut}_{\mathbb{R}}(X, \pi)$ exchanging the components of each fibre. Then $s := g(\tilde{E}_n) \neq \tilde{E}_n$ is a real section of self-intersection $-r$. Forgetting about the action of $\text{Aut}_{\mathbb{R}}(X, \pi)$, we contract in each fibre the component intersecting s . This is a blow-down $\eta': X \rightarrow \mathbb{F}_r$, and $r \leq \eta'(s)^2 = -r + m$, where $1 \leq m \leq 2$ is the number of points blown-up by η' . It follows that $m = 2$ and $r \in \{0, 1\}$. The case $r = 0$ is not possible, so $r = 1$. Hence η' blows up two points in different fibres of \mathbb{F}_1 , and X is a del Pezzo surface of degree 6. Therefore, $\text{Aut}_{\mathbb{R}}(X, \pi) \subset \text{Aut}_{\mathbb{R}}(X)$. Figures 4 and 3 and Propositions 3.5 and 3.6 imply that the inclusion is strict. \square

5. The maximal infinite algebraic subgroups

This section aims at proving Theorem 1.1 and Theorem 1.2. We first prove that any infinite algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is contained in one of the groups in Theorem 1.1. We then have to prove that all listed groups are in fact maximal.

Proposition 5.1. *Let $G \subset \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ be an infinite algebraic subgroup. Then G is conjugate to a subgroup of one of the groups in Theorem 1.1.*

Proof. Proposition 2.21 states that for an infinite algebraic subgroup G of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ there exists a G -equivariant birational morphism $\mathbb{P}^2 \dashrightarrow X$ where G acts on X regularly and X is one of the following:

- (1) X is a Del Pezzo surface of degree 6, 8 or 9 such that $\text{rk}(\text{Pic}(X)^G) = 1$.
- (2) X admits a real conic bundle structure $\pi_X: X \rightarrow \mathbb{P}^1$ with $\text{rk}(\text{Pic}(X)^G) = 2$ and $G \subset \text{Aut}_{\mathbb{R}}(X, \pi_X)$, and there is a birational morphism of conic bundles $\eta: X \rightarrow Y$, where $Y \simeq X_{[2]}$ is the sphere blown up in a pair of non-real conjugate points or Y is a real Hirzebruch surface $Y = \mathbb{F}_n$, $n \neq 1$.

So G is conjugate to a subgroup of $\text{Aut}_{\mathbb{R}}(X)$ or $\text{Aut}_{\mathbb{R}}(X, \pi)$, where X is as in (1) or (2) respectively. The pairs $(X, \text{Aut}_{\mathbb{R}}(X))$ and $(X, \text{Aut}_{\mathbb{R}}(X, \pi))$ are described as follows.

In the first case, we get:

- If $\text{deg}(X) = 9$, then $X = \mathbb{P}^2$ and $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \simeq \text{PGL}_3(\mathbb{R})$.
- If $\text{deg}(X) = 8$, then $X = \mathbb{F}_1$ and $\text{Aut}_{\mathbb{R}}(X)$ is conjugate to a subgroup of $\text{PGL}_3(\mathbb{R})$, or X is one of the following two by [7]:

$$X = \mathbb{Q}_{3,1} \text{ and } \text{Aut}_{\mathbb{R}}(\mathbb{Q}_{3,1}) \simeq \mathbb{P}\text{O}_{\mathbb{R}}(3, 1).$$

$$X = \mathbb{F}_0 \text{ and } \text{Aut}_{\mathbb{R}}(\mathbb{F}_0) = \text{PGL}_2(\mathbb{R})^2 \rtimes \langle \tau' \rangle, \text{ where } \tau' : (x, y) \mapsto (y, x).$$

- If $\text{deg}(X) = 6$, then Lemma 3.1 and Propositions 3.3, 3.4, 3.5 and 3.6 imply that $X = X_{[2]}$ and $\text{Aut}_{\mathbb{R}}(X_{[2]})$ is conjugate to a subgroup of $\text{Aut}_{\mathbb{R}}(\mathbb{Q}_{3,1})$ (Proposition 3.3).
 $X = X_{[3, \mathbb{Q}_{3,1}]}$ and $\text{Aut}_{\mathbb{R}}(X_{[3, \mathbb{Q}_{3,1}]})$ is conjugate to a subgroup of $\text{Aut}_{\mathbb{R}}(\mathbb{Q}_{3,1})$ (Proposition 3.4),
 $X = X_{[3, \mathbb{F}_0]}$ and by Proposition 3.5 states the action of $\text{Aut}_{\mathbb{R}}(X)$ on $\text{Pic}(X)$ induces the split exact sequence

$$1 \rightarrow \text{SO}_2(\mathbb{R}) \times \text{SO}_2(\mathbb{R}) \rightarrow \text{Aut}_{\mathbb{R}}(X_{[3, \mathbb{F}_0]}) \rightarrow D_6 \rightarrow 1,$$

or $X = X_{[4]}$ and Proposition 3.6 states that the action of $\text{Aut}_{\mathbb{R}}(X)$ induces the split exact sequence

$$1 \rightarrow (\mathbb{R}^*)^2 \text{Aut}_{\mathbb{R}}(X_{[4]}) \rightarrow D_6 \rightarrow 1.$$

In the second case, we look up the results of Section 4:

- Lemma 4.3 implies that $\eta: X \rightarrow Y \simeq X_{[2]}$ is the blow-up of $n \geq 1$ pairs of non-real conjugate points in the exceptional divisors of $X_{[2]} \rightarrow \mathbb{Q}_{3,1}$ and contained in non-real fibres, and $\pi = \pi_{[2]}\eta: X \rightarrow \mathbb{P}^1$ is relatively $\text{Aut}_{\mathbb{R}}(X, \pi)$ -minimal. Proposition 4.5 implies that the action of $\text{Aut}_{\mathbb{R}}(X)$ on \mathbb{P}^1 induces the split exact sequence

$$1 \rightarrow \text{SO}_2(\mathbb{R}) \rtimes \mathbb{Z}/2\mathbb{Z} \rightarrow \text{Aut}_{\mathbb{R}}(X) \rightarrow \rtimes H_{\Delta} \rightarrow,$$

where $H_{\Delta} \subset \text{PGL}_2(\mathbb{R})$ is the subgroup preserving the image in \mathbb{P}^1 of the $2n + 2$ singular fibres of X and the interval $\pi(X(\mathbb{R})) = \pi_{[2]}(X_{[2]}(\mathbb{R})) = [0, \infty]$.

- If $\eta: X \rightarrow Y = \mathbb{F}_n$, then the following possibilities occur:
 - If η is an isomorphism, then $X = \mathbb{F}_n$ and

$$\text{Aut}_{\mathbb{R}}(\mathbb{F}_0, \text{pr}_0) \subseteq \text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$$

$$\text{Aut}_{\mathbb{R}}(\mathbb{F}_1, \text{pr}_1) \subseteq \text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \quad (\text{see beginning of Section 4.2})$$

$$\text{Aut}_{\mathbb{R}}(\mathbb{F}_n, \text{pr}_n) = \text{Aut}_{\mathbb{R}}(\mathbb{F}_n) \simeq \mathbb{R}^{n+1} \rtimes (\text{GL}_2(\mathbb{R})/\{\mu \text{Id} \mid \mu^n = 1\}), \quad n \geq 2$$

– Else, $\text{Aut}_{\mathbb{R}}(X, \pi)$ being infinite and maximal, Lemma 4.7 and Lemma 4.9 imply that there exists a birational morphism $\eta': X \rightarrow \mathbb{F}_N$ that is the blow-up of $2N$ points on s_N (see definition in beginning of Section 4.2).

If $N = 0$, then $\eta': Y \rightarrow \mathbb{F}_0$ is an isomorphism, and we have already listed this case.

If $N = 1$, then Y is a del Pezzo surface of degree 6; if η' blows up two real points, then $Y \simeq X_{[4]}$ and if η' blows up a pair of non-real conjugate points, then $Y \simeq X_{[3, \mathbb{F}_0]}$ (see Section 3). In either case, $\text{Aut}_{\mathbb{R}}(Y, \pi) \subseteq \text{Aut}_{\mathbb{R}}(Y)$.

If $N \geq 2$, then Proposition 4.8 states that $\pi = \pi_N \eta': Y \rightarrow \mathbb{P}^1$ is relatively $\text{Aut}_{\mathbb{R}}(Y, \pi)$ -minimal and the action of $\text{Aut}_{\mathbb{R}}(X)$ on \mathbb{P}^1 induces the split exact sequence


$$1 \rightarrow \mathbb{R}^*/\mu_N \rtimes \mathbb{Z}/2\mathbb{Z} \rightarrow \text{Aut}_{\mathbb{R}}(Y) \rightarrow H_{\Delta} \rightarrow 1,$$

where $H_{\Delta} \subset \text{PGL}_2(\mathbb{R})$ is the subgroup fixing the image in \mathbb{P}^1 of the $2N$ points and $\mu_N \subset \mathbb{R}^*$ the group of N -th roots of unity. □

Proposition 5.2. *The groups in Theorem 1.1 are maximal algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ and the classes are pairwise non-conjugate.*

Proof. Let (X, G) be a pair in the list of Theorem 1.1. To prove that the group G is a maximal algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, we have to check because of Proposition 5.1 that any G -equivariant birational map $f: X \dashrightarrow Y$, where Y is one of the surfaces listed in Theorem 1.1, is in fact a G -equivariant isomorphism. This will also prove that all the classes are distinct.

As X and Y are del Pezzo surfaces or conic bundles, [12, Theorem 2.5] (and [8, Appendix]) implies that any birational map $X \dashrightarrow Y$ is an isomorphism or the composition of elementary links, which are divided into type I–IV, shown by the commutative diagrams below, where the horizontal maps are blow-ups defined over \mathbb{R} , and $T \in \{\mathbb{P}^1, *\}$.



The decomposition into links can be made G -equivariant because G is a linear algebraic group (Lemma 2.22) and its action on the Picard group is finite by Lemma 2.10. The horizontal maps of the G -equivariant links blow up the finite G -orbit of a real point or the finite G -orbit of a pair of non-real points.

Suppose that f is not an isomorphism and let $f = \Phi_n \cdots \Phi_1$ be its decomposition into G -equivariant links of type I–IV. Then:

- If Φ_1 is a link of type I, then [12, Theorem 2.6] implies that X is a del Pezzo surface of degree 9, 8 or 4. The latter case does not appear, and $X = \mathbb{P}^2$ or $X = \mathcal{Q}_{3,1}$ or $X = \mathbb{F}_0$ by [7]. However, there are no finite G -orbits on X , which makes such a link impossible.
- If Φ_1 is a link of type III, then [12, Theorem 2.6] implies that $X \simeq \mathbb{F}_1$, $X \simeq X_{[3,S]}$ or $X \simeq X_{[2]}$ or $X \simeq X_{[4]}$. Only the latter is in our list. [12, Theorem 2.6] says that Φ_1 must be the contraction of an orbit of order 2, which does not exist by Proposition 3.6.
- If $\Phi_1 = \eta_2 \eta_1^{-1}$ is a link of type II, then either X, X_1 are both del Pezzo surfaces and $T = *$ or X, X_1 are both conic bundles and $T = \mathbb{P}^1$. We look at these cases separately:

If X and X_1 are del Pezzo surfaces, then [12, Theorem 2.6] implies that the degree of X is 9, 8, 6, 5, 4, 3 or 2. Only the first three degrees appear in our list. If X has degree 9 or 8, then, again, there are no finite G -orbits on X , so a link of type II is not possible. Suppose that X is of degree 6, i.e. $X \simeq X_{[3,\mathbb{F}_0]}$ or $X \simeq X_{[4]}$ in Lemma 3.1. By [12, Theorem 2.6], η_1 is the blow-up of at most 5 points on X . That is impossible because the only finite G -orbit on X has cardinality six by Propositions 3.5 and 3.6.

Suppose that X and X_1 are real conic bundles. If $X = \mathbb{F}_n$ for some $n \in \mathbb{N}$, then G does not have a finite orbit and hence a link of type II cannot start with \mathbb{F}_n .

If there exists a birational morphism $X \rightarrow X_{[2]}$ of real conic bundles, then Proposition 4.5 states that $\text{Aut}_{\mathbb{R}}(X/\pi)$ contains an element exchanging the two unique $-(n+1)$ -sections of X . If there exists a birational morphism $X \rightarrow \mathbb{F}_n$ of conic bundles that is not an isomorphism, then Proposition 4.8 states that $\text{Aut}_{\mathbb{R}}(X/\pi)$ contains an element exchanging the two $(-n)$ -

sections of X . In either case, a G -orbit on X contains at least two points that are contained in the same fiber, which is not allowed. So there is no link of type II starting with X .

- If Φ_1 is a link of type IV, then [12, Theorem 2.6] implies that $K_X^2 = 8, 4, 2, 1$. So, consulting our list, $K_X^2 = 8$ and, again by [12, Theorem 2.6], $X = X_1 = \mathbb{F}_0$ and Φ_1 exchanges the two fibrations. In particular, $\Phi_1 \in \text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$.

Summarised, one cannot find a decomposition of f into elementary links and hence f is an isomorphism. □

Proof of Theorem 1.1. By Proposition 5.1, any algebraic subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is conjugate to a subgroup of one of the groups in the list. By Proposition 5.2, all of these groups are maximal and pairwise non-conjugate. □

Proof of Theorem 1.2. The claim is clear for families (1)-(3) and (6). The rest of the claim follows from the description of $\text{Aut}_{\mathbb{R}}(X)$ in Propositions 3.5 and 3.6 and of $\text{Aut}_{\mathbb{R}}(X, \pi)$ in Propositions 4.5 and 4.8. □

Let us take a look at which infinite algebraic subgroups survive the abelianisation of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.

REMARK 5.3. (1) The construction of the abelianisation $\varphi: \text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \rightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ in [25, §3.2.1, Proposition 5.3] yields the following: Let g_1 and g_2 be elementary links of the real conic bundle $\pi_{[2]}: X_{[2]} \rightarrow \mathbb{P}^1$ contracting pairs of non-real conjugate fibres f_1, \bar{f}_1 and f_2, \bar{f}_2 respectively. Then g_1 and g_2 have the same image in the quotient if and only if

$$\pi_{[2]}(f_1) \in \mathbb{R} \cdot \pi_{[2]}(f_2) \cup \mathbb{R} \cdot \pi_{[2]}(\bar{f}_2) \quad \text{in } \mathbb{P}^1.$$

(2) Let $\pi: X \xrightarrow{\eta} X_{[2]} \xrightarrow{\pi_{[2]}} \mathbb{P}^1$ be a surface as in Theorem 1.1 (7). We can see elements of $\text{Aut}_{\mathbb{R}}(X, \pi)$ as birational transformations of $X_{[2]}$ preserving the conic bundle structure. Let $(q_1, \bar{q}_1), \dots, (q_n, \bar{q}_n)$ be the pairs of non-real conjugate points blown up by η . Then, by definition of φ in [25, Definition 3.17] and the description of $\text{Aut}_{\mathbb{R}}(X, \pi)$ in the proof of Proposition 4.5, we have

$$\varphi(g) = \sum_{i=1}^n e_{\nu(\pi(q_i))} \text{ if } g \in \text{Aut}_{\mathbb{R}}(X, \pi) \setminus (\text{SO}_2(\mathbb{R}) \cup H_{\Delta}), \quad \varphi(g) = 0 \text{ if } g \in \text{SO}_2(\mathbb{R}) \cup H_{\Delta}$$

where $\nu([a + \mathbf{i}b : 1]) = \nu([a - \mathbf{i}b : 1]) = 1 - \frac{|a|}{a^2+b^2}$ and e_r is the “standard vector” with entry 1 at r and zero everywhere else.

Proof of Theorem 1.3. It suffices to see the image of the maximal infinite subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ have in the quotient $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)/\langle\langle \text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \rangle\rangle \simeq \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$. The quotient is abelian, so it suffices to check the groups listed in Theorem 1.1.

Groups (1)–(3) have trivial image because their elements are conjugate to transformations of \mathbb{P}^2 of degree at most 2.

The generators of group (4) either descend to automorphisms of \mathbb{F}_0 or to birational transformations of \mathbb{F}_0 sending one fibration onto the other (Proposition 3.5). The latter are conjugate to transformations of \mathbb{P}^2 sending a pencil of lines through a real point onto the pencil of lines through another real point. So, also the generators of (4) have trivial image in the quotient.

The generators of the group (5) are conjugate to transformations of \mathbb{P}^2 of degree at most 2 by Proposition 3.6, so they are contained in $\langle\langle \text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \rangle\rangle$ [25, Lemma 6.6].

The groups in families (6) and (8) are conjugate to transformations of \mathbb{P}^2 preserving a pencil of lines through a point. So, they have trivial image in the quotient [25, Proposition 5.3].

By Remark 5.3 there exist real conic bundles $\pi: X \xrightarrow{\eta} X_{[2]} \xrightarrow{\pi_{[2]}} \mathbb{P}^1$ as in family (7) that have non-trivial image in $\bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$. The image is finite and they are mapped onto the generator e_r if and only if η blows up exactly one pair of non-real conjugate points q, \bar{q} such that $v(\pi(p)) = r$. \square

Proof of Corollary 1.4. This is a direct consequence of the fact that the images of the algebraic subgroups are finite by Theorem 1.3. \square

ACKNOWLEDGEMENTS. The authors would like to express their warmest thanks to Jérémy Blanc for many helpful discussions and Michel Brion for his detailed remarks on algebraic groups and equivariant completions. They would also like to thank Egor Yasinsky for interesting discussions about the real Cremona group.

References

- [1] I. Biswas and J. Huisman: *Rational real algebraic models of topological surfaces*, Doc. Math. **12** (2007), 549–567.
- [2] J. Blanc: *Sous-groupes algébriques du groupe de Cremona*, Transformation groups, **14** (2009), 249–285.
- [3] J. Blanc and J.-P. Furter: *Topologies and structures on the Cremona groups*, Ann. of Math. **178** (2013), 1173–1198.
- [4] J. Blanc and F. Mangolte: *Cremona groups of real surfaces*, in Automorphisms in birational and affine geometry, Springer Proc. Math. Stat. **79** (2014), 35–58.
- [5] M. Brion: *On actions of connected algebraic groups*, arXiv:1412.1906v1.
- [6] G. Castelnuovo: *Le trasformazioni generatrici del gruppo cremoniano nel piano*, Atti della R. Accad. delle Scienze di Torino, **36** (1901), 861–874.
- [7] A. Comessatti: *Fondamenti per la geometria sopra le superficie razionali dal punto di vista reale*, Math. Ann. **73** (1912), 1–72.
- [8] A. Corti: *Factoring birational maps of threefolds after Sarkisov*, J. Algebraic. Geom. **4** (1995), 223–254.
- [9] M. Demazure: *Sous-groupes algébriques de rang maximum du groupe de Cremona*, Ann. Sci. École Norm. Sup. (4) **3** (1970), 507–588.
- [10] I. Dolgachev and V. Iskovskikh: *Finite subgroups of the plane Cremona group*, in Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I, Progr. Math. **269**, Birkhäuser Boston, Inc., Boston, MA, (2009), 443–548.
- [11] J. Huisman and F. Mangolte: *The group of automorphisms of a real rational surface is n-transitive*, Bull. Lond. Math. Soc. **41** (2009), 563–568.
- [12] V.A. Iskovskikh: *Factorization of birational mappings of rational surfaces from the viewpoint of Mori theory*, Uspekhi Mat. Nauk **51**, 3–72. English translation: Russian Math. Surveys **51** (1996), 585–652.
- [13] J. Kollár and S. Mori: *Birational geometry of algebraic varieties*, with the collaboration of C.H. Clemens and A. Corti. Translated from the 1998 Japanese original. Cambridge Tracts in Mathematics, **134**, Cambridge University Press, Cambridge, 1998.
- [14] J. Kollár and F. Mangolte: *Cremona transformations and diffeomorphisms of surfaces*, Adv. Math. **222** (2009), 44–61.
- [15] J. Limpan: *Desingularization of two-dimensional schemes*, Ann. Math. (2) **107** (1978), 151–207.

- [16] M.F. Robayo: *Prime order birational diffeomorphisms of the sphere*, Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5) **16** (2016), 909–970.
- [17] M. Rosenlicht: *Some basic theorems on algebraic groups*, Amer. J. Math. **78** (1956), 401–443.
- [18] F. Ronga and T. Vust: *Diffeomorfismi birazionali del piano proiettivo reale*, Comm. Math. Helv. **80** (2005), 517–540.
- [19] J.-J. Sansuc: *Groupe de Brauer et arithmétique des groupes algébriques linéaires sur un corps de nombres*, J. Reine Angew. Math. **327** (1981), 12–80.
- [20] J.-P. Serre: *Le groupe de Cremona et ses sous-groupes finis*, Séminaire Bourbaki. Volume 2008/2009. Astérisque No. **332** (2010), Exp. No. 1000, vii, 75–100.
- [21] H. Sumihiro: *Equivariant completion II*, J. Math. Kyoto Univ. **15** (1975), 573–605.
- [22] W.C. Waterhouse: *Introduction to Affine Group Schemes*, Graduate Texts in Mathematics, **66**. Springer-Verlag, New York-Berlin, 1979.
- [23] A. Weil: *On Algebraic Groups of Transformations*, Amer. J. Math. **77** (1955), 355–391.
- [24] E. Yasinsky: *Subgroups of odd order in the real plane Cremona group*, J. Algebra **461** (2016), 87–120.
- [25] S. Zimmermann: *The abelianisation of the real Cremona group*, Duke Math. J. **167** (2018), 211–267.

Maria Fernanda Robayo
Basel
Switzerland
e-mail: maferobayo@gmail.com

Susanna Zimmermann
LAREMA
Université d'Angers
France
e-mail: zimmermann@math.univ-angers.fr