

## The proportion of cyclic quartic fields with discriminant divisible by a given prime

By Blair K. SPEARMAN<sup>\*)</sup> and Kenneth S. WILLIAMS<sup>\*\*)</sup>

(Communicated by Heisuke HIRONAKA, M. J. A., Sept. 13, 2004)

**Abstract:** An asymptotic formula is given for the number of cyclic quartic fields with discriminant  $\leq x$  and divisible by a given prime.

**Key words:** Discriminant; cyclic quartic field.

**1. Introduction.** It was shown in [1, Theorem, p. 97] that the number  $N(x)$  of cyclic quartic fields  $K$  with discriminant  $d(K) \leq x$  satisfies

$$(1.1) \quad N(x) = \frac{3}{\pi^2} \left\{ \frac{(24 + \sqrt{2})}{24} C - 1 \right\} x^{1/2}, \\ + O(x^{1/3} \log^3 x),$$

as  $x \rightarrow \infty$ , where

$$(1.2) \quad C = \prod_{p \equiv 1 \pmod{4}} \left( 1 + \frac{2}{(p+1)\sqrt{p}} \right).$$

Here and throughout this paper  $p$  denotes a prime. Let  $q$  be a fixed prime. In this paper, which should be viewed as a continuation of [1], we determine an asymptotic formula for the number  $N_q(x)$  of cyclic quartic fields  $K$  with discriminant  $d(K) \leq x$  and  $d(K) \equiv 0 \pmod{q}$ . We prove

**Theorem.** *Let  $q$  be a prime. Then*

$$(1.3) \quad N_q(x) = E_q x^{1/2} + O(x^{1/3} \log^3 x),$$

as  $x \rightarrow \infty$ , where

$$E_2 = \frac{1}{\pi^2} \left( \frac{(8 + \sqrt{2})}{8} C - 1 \right), \\ E_q = \frac{3}{\pi^2(q+1)} \left( \left( \frac{24 + \sqrt{2}}{24} \right) C - 1 \right), \\ \text{if } q \equiv 3 \pmod{4},$$

$$E_q = \frac{3}{\pi^2(q+1)} \left( \left( \frac{24 + \sqrt{2}}{24} \right) \left( \frac{1 + \frac{2}{\sqrt{q}}}{1 + \frac{2}{(q+1)\sqrt{q}}} \right) C - 1 \right), \\ \text{if } q \equiv 1 \pmod{4}.$$

This theorem is proved in Section 3 after some preliminary results are given in Section 2.

The proportion  $d_q$  of cyclic quartic fields with discriminant divisible by the fixed prime  $q$  is

$$d_q = \lim_{x \rightarrow \infty} \frac{N_q(x)}{N(x)} = \frac{E_q}{\frac{3}{\pi^2} \left\{ \frac{(24 + \sqrt{2})}{24} C - 1 \right\}}.$$

Appealing to the values of  $E_q$  given in the Theorem, the proportion  $d_q$  is given by

$$d_q = \frac{(8 + \sqrt{2})C - 8}{(24 + \sqrt{2})C - 24}, \quad \text{if } q = 2, \\ d_q = \frac{1}{q+1}, \quad \text{if } q \equiv 3 \pmod{4}, \\ d_q = \frac{(24 + \sqrt{2}) \left( \frac{1 + \frac{2}{\sqrt{q}}}{1 + \frac{2}{(q+1)\sqrt{q}}} \right) C - 24}{(q+1)((24 + \sqrt{2})C - 24)}, \\ \text{if } q \equiv 1 \pmod{4}.$$

**2. Some Lemmas.** The results of this section are used in Section 3. They are either contained in [1] or [2] or are simple extensions of results there. We use ‘ $n$  sqf’ to indicate that the positive integer  $n$  is required to be squarefree. As usual, for  $n \in \mathbf{N}$ ,  $\phi(n)$  is Euler’s totient function and  $d(n)$  counts the number of positive divisors of  $n$ . The greatest common divisor of the positive integers  $a$  and  $b$  is

2000 Mathematics Subject Classification. Primary 11R16; Secondary 11R21, 11R29.

<sup>\*)</sup> Department of Mathematics and Statistics, Okanagan University College, Kelowna, B.C. Canada V1V 1V7.

<sup>\*\*) School of Mathematics and Statistics, Carleton University, Ottawa, Ontario, Canada K1S 5B6.</sup>

denoted by  $(a, b)$ .

**Lemma 2.1.** *Let  $k \in \mathbf{N}$ . Then*

$$\sum_{\substack{1 \leq n \leq x \\ n \text{ sqf} \\ (n,k)=1}} 1 = x \frac{6}{\pi^2} \frac{\phi(k)}{k} \prod_{p|k} \left(1 - \frac{1}{p^2}\right)^{-1} + O(x^{1/2} d(k)),$$

as  $x \rightarrow \infty$ , where the implied constant is absolute.

*Proof.* See [2, Lemma 3, p. 182].  $\square$

**Lemma 2.2.** *Let  $k \in \mathbf{N}$ . Let  $q$  be an odd prime. Then*

$$\sum_{\substack{1 \leq n \leq x \\ n \text{ sqf} \\ (n,k)=1 \\ q|n}} 1 = \begin{cases} 0, & \text{if } q \mid k, \\ \frac{x}{q+1} \frac{6}{\pi^2} \frac{\phi(k)}{k} \prod_{p|k} \left(1 - \frac{1}{p^2}\right)^{-1} \\ \quad + O(x^{1/2} q^{-1/2} d(k)), & \text{if } q \nmid k, \end{cases}$$

where the implied constant is absolute.

*Proof.* The result is clear for  $q \mid k$ . For  $q \nmid k$  we have

$$\begin{aligned} \sum_{\substack{1 \leq n \leq x \\ n \text{ sqf} \\ (n,k)=1 \\ q|n}} 1 &= \sum_{\substack{1 \leq n \leq x/q \\ n \text{ sqf} \\ q \nmid n \\ (qn,k)=1}} 1 = \sum_{\substack{1 \leq n \leq x/q \\ n \text{ sqf} \\ (n,qk)=1}} 1 \\ &= \frac{x}{q} \frac{6}{\pi^2} \frac{\phi(qk)}{qk} \prod_{p|qk} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad + O\left(\left(\frac{x}{q}\right)^{1/2} d(qk)\right) \\ &= \frac{x}{q+1} \frac{6}{\pi^2} \frac{\phi(k)}{k} \prod_{p|k} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad + O(x^{1/2} q^{-1/2} d(k)), \end{aligned}$$

by Lemma 2.1.  $\square$

Following [1, eq. (3.7), p. 100] we set

$$\wp = \{D \mid D = q_1 \cdots q_r \ (r \geq 1), q_1, \dots, q_r \text{ distinct primes } \equiv 1 \pmod{4}\}.$$

Note that  $1 \notin \wp$ . We set (as in [1, eq. (3.8), p. 100])

$$S(x) := \sum_{\substack{D \leq x^{1/3} \\ D \in \wp}} d(D) \sum_{\substack{1 \leq A \leq \sqrt{xD^{-3}} \\ A \text{ sqf} \\ (A,2D)=1}} 1.$$

**Lemma 2.3.**

$$S(x) = \frac{4}{\pi^2} (C - 1) x^{1/2} + O(x^{1/3} \log^3 x),$$

where the implied constant is absolute.

*Proof.* See [1, p. 103]. Note that we have  $c_0 + 1 = C$ .  $\square$

Let  $q$  be an odd prime. We define

$$S_1(x) := \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q|D}} d(D) \sum_{\substack{1 \leq A \leq \sqrt{xD^{-3}} \\ A \text{ sqf} \\ (A,2D)=1}} 1$$

and

$$S_2(x) := \sum_{\substack{D \leq x^{1/3} \\ D \in \wp}} d(D) \sum_{\substack{1 \leq A \leq \sqrt{xD^{-3}} \\ A \text{ sqf} \\ (A,2D)=1 \\ q|A}} 1.$$

We note that

$$S_1(x) = 0, \quad \text{if } q \equiv 3 \pmod{4}.$$

**Lemma 2.4.** *Let  $q$  be an odd prime. Then*

$$S_2(x) = \frac{4}{\pi^2(q+1)} (C' - 1) x^{1/2} + O(x^{1/3} \log^3 x),$$

where the implied constant depends only on  $q$ , and

$$(2.1) \quad C' = \prod_{\substack{p \equiv 1 \pmod{4} \\ p \neq q}} \left(1 + \frac{2}{(p+1)\sqrt{p}}\right).$$

We note that  $C' = C$  if  $q \equiv 3 \pmod{4}$ , whereas

$$(2.2) \quad C' = \frac{C}{\left(1 + \frac{2}{(q+1)\sqrt{q}}\right)}$$

if  $q \equiv 1 \pmod{4}$ .

*Proof.* We have by Lemma 2.2

$$\begin{aligned} S_2(x) &= \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q \nmid D}} d(D) \left\{ \frac{x^{1/2}}{D^{3/2}} \frac{1}{q+1} \right. \\ &\quad \times \frac{6}{\pi^2} \frac{\phi(2D)}{2D} \prod_{p|2D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad \left. + O(x^{1/4} D^{-3/4} d(D)) \right\} \\ &= \frac{4}{\pi^2} \frac{x^{1/2}}{q+1} \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q \nmid D}} d(D) D^{-5/2} \\ &\quad \times \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \end{aligned}$$

$$+ O(x^{1/4} \sum_{\substack{D \leq x^{1/3} \\ D \in \wp}} d^2(D) D^{-3/4}).$$

It is shown in [1, p. 103] that

$$\sum_{\substack{D \leq x^{1/3} \\ D \in \wp}} d^2(D) D^{-3/4} = O(x^{1/12} \log^3 x).$$

Also

$$\begin{aligned} & \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q \nmid D}} d(D) D^{-5/2} \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &= \sum_{\substack{D=1 \\ D \in \wp \\ q \nmid D}}^{\infty} d(D) D^{-5/2} \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &+ O\left(\sum_{\substack{D > x^{1/3} \\ D \in \wp}} d(D) D^{-5/2} \phi(D)\right), \end{aligned}$$

as

$$\prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} < \frac{\pi^2}{6}.$$

Clearly

$$\begin{aligned} & \sum_{\substack{D=1 \\ D \in \wp \\ q \nmid D}}^{\infty} d(D) D^{-5/2} \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &= \prod_{\substack{p \equiv 1 \pmod{4} \\ p \neq q}} \left(1 + \frac{2}{(p+1)\sqrt{p}}\right) - 1 = C' - 1. \end{aligned}$$

Also

$$\sum_{\substack{D > x^{1/3} \\ D \in \wp}} d(D) D^{-5/2} \phi(D) = O(x^{-1/6} \log x),$$

see [1, p. 103]. Thus

$$\begin{aligned} S(x) &= \frac{4}{\pi^2} \frac{x^{1/2}}{q+1} (C' - 1) + O(x^{1/2-1/6} \log x) \\ &\quad + O(x^{1/4+1/12} \log^3 x), \end{aligned}$$

which gives the asserted result.  $\square$

**Lemma 2.5.** *Let  $q$  be a prime  $\equiv 1 \pmod{4}$ . Then*

$$S_1(x) = \frac{8}{\pi^2} \frac{x^{1/2}}{(q+1)\sqrt{q}} C' + O(x^{1/3} \log^3 x).$$

*Proof.* We have by Lemma 2.1

$$\begin{aligned} S_1(x) &= \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q|D}} d(D) \left\{ \left(\frac{x}{D^3}\right)^{1/2} \frac{6}{\pi^2} \right. \\ &\quad \times \frac{\phi(2D)}{2D} \prod_{p|2D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad \left. + O\left(\left(\frac{x}{D^3}\right)^{1/4} d(D)\right) \right\} \\ &= \frac{4}{\pi^2} x^{1/2} \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q|D}} d(D) D^{-5/2} \\ &\quad \times \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad + O\left(x^{1/4} \sum_{\substack{D \leq x^{1/3} \\ D \in \wp}} d^2(D) D^{-3/4}\right). \end{aligned}$$

As in Lemma 2.4 the error term is

$$O(x^{1/4+1/12} \log^3 x) = O(x^{1/3} \log^3 x).$$

Also

$$\begin{aligned} & \sum_{\substack{D \leq x^{1/3} \\ D \in \wp \\ q|D}} d(D) D^{-5/2} \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &= \sum_{\substack{D=1 \\ D \in \wp \\ q \nmid D}}^{\infty} d(D) D^{-5/2} \phi(D) \prod_{p|D} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad + O\left(\sum_{\substack{D > x^{1/3} \\ D \in \wp}} d(D) D^{-5/2} \phi(D)\right) \\ &= d(q) q^{-5/2} \phi(q) \left(1 - \frac{1}{q^2}\right)^{-1} \\ &\quad \times \prod_{\substack{p \equiv 1 \pmod{4} \\ p \neq q}} \left(1 + \frac{2}{(p+1)\sqrt{p}}\right) \\ &\quad + O(x^{-1/6} \log x) \\ &= \frac{2}{(q+1)\sqrt{q}} C' + O(x^{-1/6} \log x). \end{aligned}$$

Finally

$$\begin{aligned} S_1(x) &= \frac{4x^{1/2}}{\pi^2} \left( \frac{2}{(q+1)\sqrt{q}} C' + O(x^{-1/6} \log x) \right) \\ &\quad + O(x^{1/3} \log^3 x) \\ &= \frac{8}{\pi^2} \frac{x^{1/2}}{(q+1)\sqrt{q}} C' + O(x^{1/3} \log^3 x), \end{aligned}$$

which is the assertion of Lemma 2.5.  $\square$

**3. Proof of Theorem.** From [1, (3.3) and (3.4), p. 100] we see that if  $q = 2$

$$\begin{aligned} N_q(x) &= 2 \sum_{\substack{A \leq (x/2^{11})^{1/2} \\ A \text{ sqf} \\ A \text{ odd}}} 1 + 2S(2^{-11}x) \\ &\quad + S(2^{-6}x) + \frac{1}{2}S(2^{-4}x); \end{aligned}$$

if  $q \equiv 3 \pmod{4}$

$$\begin{aligned} N_q(x) &= 2 \sum_{\substack{A \leq (x/2^{11})^{1/2} \\ A \text{ sqf} \\ A \text{ odd} \\ q|A}} 1 + 2S_2(2^{-11}x) \\ &\quad + S_2(2^{-6}x) + \frac{1}{2}S_2(2^{-4}x) + \frac{1}{2}S_2(x); \end{aligned}$$

and if  $q \equiv 1 \pmod{4}$

$$\begin{aligned} N_q(x) &= 2 \sum_{\substack{A \leq (x/2^{11})^{1/2} \\ A \text{ sqf} \\ A \text{ odd} \\ q|A}} 1 + 2S_2(2^{-11}x) + S_2(2^{-6}x) \\ &\quad + \frac{1}{2}S_2(2^{-4}x) + \frac{1}{2}S_2(x) + 2S_1(2^{-11}x) \\ &\quad + S_1(2^{-6}x) + \frac{1}{2}S_1(2^{-4}x) + \frac{1}{2}S_1(x). \end{aligned}$$

For  $q = 2$  we have by Lemmas 2.1 and 2.3

$$\begin{aligned} N_q(x) &= 2 \left( \left( \frac{x}{2^{11}} \right)^{1/2} \frac{6}{\pi^2} \frac{\phi(2)}{2} \right. \\ &\quad \times \left. \prod_{p|2} \left( 1 - \frac{1}{p^2} \right)^{-1} + O(x^{1/4}) \right) \\ &\quad + \frac{8}{\pi^2} (C-1) \left( \frac{x}{2^{11}} \right)^{1/2} + \frac{4}{\pi^2} (C-1) \left( \frac{x}{2^6} \right)^{1/2} \\ &\quad + \frac{2}{\pi^2} (C-1) \left( \frac{x}{2^4} \right)^{1/2} + O(x^{1/3} \log^3 x) \\ &= \frac{1}{2^{5/2}} \frac{x^{1/2}}{\pi^2} + \frac{2(C-1)}{\pi^2} x^{1/2} \left( \frac{4}{2^{11/2}} + \frac{2}{2^3} + \frac{1}{2^2} \right) \\ &\quad + O(x^{1/3} \log^3 x) \end{aligned}$$

$$\begin{aligned} &= \frac{x^{1/2}}{8\pi^2} (\sqrt{2} + (C-1)(\sqrt{2} + 8)) + O(x^{1/3} \log^3 x) \\ &= \frac{x^{1/2}}{8\pi^2} ((8 + \sqrt{2})C - 8) + O(x^{1/3} \log^3 x) \\ &= \frac{1}{\pi^2} \left( \frac{(8 + \sqrt{2})}{8} C - 1 \right) x^{1/2} + O(x^{1/3} \log^3 x). \end{aligned}$$

For  $q \equiv 3 \pmod{4}$  we have by Lemmas 2.2 and 2.4

$$\begin{aligned} N_q(x) &= 2 \left( \left( \frac{x}{2^{11}} \right)^{1/2} \frac{1}{q+1} \frac{6}{\pi^2} \frac{\phi(2)}{2} \right. \\ &\quad \times \left. \prod_{p|2} \left( 1 - \frac{1}{p^2} \right)^{-1} + O(x^{1/4}) \right) \\ &\quad + \frac{8}{\pi^2} \frac{(C-1)}{(q+1)} \left( \frac{x}{2^{11}} \right)^{1/2} \\ &\quad + \frac{4(C-1)}{\pi^2(q+1)} \left( \frac{x}{2^6} \right)^{1/2} \\ &\quad + \frac{2(C-1)}{\pi^2(q+1)} \left( \frac{x}{2^4} \right)^{1/2} + \frac{2(C-1)}{\pi^2(q+1)} x^{1/2} \\ &\quad + O(x^{1/3} \log^3 x) \\ &= \frac{x^{1/2}}{(q+1)\pi^2} \left( \frac{2^3}{2^{11/2}} \right) + \frac{(C-1)}{(q+1)\pi^2} x^{1/2} \\ &\quad \times \left( \frac{8}{2^{11/2}} + \frac{4}{2^3} + \frac{2}{2^2} + 2 \right) \\ &\quad + O(x^{1/3} \log^3 x) \\ &= \frac{3}{\pi^2} \frac{x^{1/2}}{(q+1)} \left( \left( \frac{24 + \sqrt{2}}{24} \right) C - 1 \right) \\ &\quad + O(x^{1/3} \log^3 x). \end{aligned}$$

For  $q \equiv 1 \pmod{4}$  we have by Lemmas 2.2, 2.4 and 2.5

$$\begin{aligned} N_q(x) &= \frac{3}{\pi^2} \frac{x^{1/2}}{q+1} \left( \left( \frac{24 + \sqrt{2}}{24} \right) C' - 1 \right) \\ &\quad + \frac{8C'x^{1/2}}{\pi^2(q+1)\sqrt{q}} \left\{ \frac{2}{2^{11/2}} + \frac{1}{2^3} + \frac{1}{2^3} + \frac{1}{2} \right\} \\ &\quad + O(x^{1/3} \log^3 x) \\ &= \frac{3}{\pi^2} \frac{x^{1/2}}{(q+1)} \\ &\quad \times \left( \left( \frac{24 + \sqrt{2}}{24} \right) \left( 1 + \frac{2}{\sqrt{q}} \right) C' - 1 \right) \\ &\quad + O(x^{1/3} \log^3 x). \end{aligned}$$

This completes the proof of the Theorem.  $\square$

**Acknowledgement.** Both authors were supported by grants from the Natural Sciences and Engineering Research Council of Canada.

[ 2 ] Spearman, B. K., and Williams, K. S.: Integers which are discriminants of bicyclic or cyclic quartic fields. *JP J. Algebra Number Theory Appl.*, **1**, 179–194 (2001).

### References

- [ 1 ] Ou, Z. M., and Williams, K. S.: On the density of cyclic quartic fields. *Canad. Math. Bull.*, **44**, 97–104 (2001).