

87. Difference Approximation of Evolution Equations and Generation of Nonlinear Semigroups

By Yoshikazu KOBAYASHI

Department of Mathematics, Waseda University

(Comm. by Kinjirō KUNUGI, M. J. A., June 3, 1975)

We consider the following nonlinear evolution equation

$$(DE) \quad (d/dt)u(t) \in Au(t), \quad 0 < t < T,$$

where A is a (multi-valued) quasi-dissipative operator. In this note, we construct the solution of the evolution equation (DE) by the method of difference approximation. In addition, we give a generation theorem of nonlinear semigroups through the difference approximation. We sketch here our results. The details will be treated in [6].

1. Preliminaries. Let X be a real Banach space. For the multi-valued operator A , we use the following notations:

$$D(A) = \{x \in X; Ax \neq \emptyset\}, \quad R(A) = \bigcup_{x \in D(A)} \{y; y \in Ax\},$$

$$\text{and } \|Ax\| = \inf \{\|y\|; y \in Ax\} \quad \text{for } x \in D(A).$$

We identify the multi-valued operator A with its graph, so that we write $[x, y] \in A$ if $y \in Ax$.

Let F be the duality map from X into X^* . Then we set

$$\langle y, x \rangle_i = \inf \{\langle y, f \rangle; f \in F(x)\} \quad \text{for } x, y \in X.$$

Let $A \subset X \times X$. A is said to be *dissipative* if for any $[x_i, y_i] \in A$ ($i=1, 2$),

$$\langle y_1 - y_2, x_1 - x_2 \rangle_i \leq 0.$$

According to Takahashi [9], we introduce the following notion as a generalization of that of dissipative operators.

Definition 1. Let $A \subset X \times X$. A is said to be *quasi-dissipative* if for any $[x_i, y_i] \in A$ ($i=1, 2$),

$$\langle y_1, x_1 - x_2 \rangle_i + \langle y_2, x_2 - x_1 \rangle_i \leq 0.$$

The following example shows that quasi-dissipative operators are not always dissipative.

Example (I. Miyadera). Let $X = R^2$ with the maximum norm. Let $x_1 = (1, 1)$ and $x_2 = (0, 0)$. We set $D(A) = \{x_1, x_2\}$, $Ax_1 = \{(\alpha, \beta); \alpha \leq 0 \text{ or } \beta \leq 0\}$ and $Ax_2 = \{(\alpha, \beta); \alpha \geq 0 \text{ or } \beta \geq 0\}$. Then A is quasi-dissipative in X but $A - \omega$ is not dissipative in X for any real ω . In addition, $R(I - \lambda A) \supset D(A)$ for any $\lambda > 0$.

The following plays a central role in our argument.

Lemma 1. Let $A \subset X \times X$. Then the following are equivalent:

(i) A is quasi-dissipative;

- (ii) for any $[x_i, y_i] \in A$ ($i=1, 2$) and $\lambda, \mu > 0$,
 $(\lambda + \mu) \|x_1 - x_2\| \leq \lambda \|x_1 - x_2 - \mu y_1\| + \mu \|x_2 - x_1 - \lambda y_2\|$;
- (iii) for any $[x_i, y_i] \in A$ ($i=1, 2$) and $\lambda > 0$,
 $2 \|x_1 - x_2\| \leq \|x_1 - x_2 - \lambda y_1\| + \|x_2 - x_1 - \lambda y_2\|$.

We can verify Lemma 1 similarly as Lemma 1.1 in Kato [4].

Let $X_0 \subset X$. A one parameter family $\{T(t); t \geq 0\}$ of operators from X_0 into itself is called (nonlinear) *contraction semigroup* on X_0 if it has the following properties :

- (i) $\|T(t)x - T(t)y\| \leq \|x - y\|$ for $x, y \in X_0$ and $t \geq 0$;
- (ii) $T(0)x = x$ for $x \in X_0$ and $T(t+s) = T(t)T(s)$ for $t, s \geq 0$;
- (iii) for each $x \in X_0$, $T(t)x$ is strongly continuous in $t \geq 0$.

2. Cauchy problems and difference approximation. Let A be a quasi-dissipative operator in X . Let $x_0 \in X$ and $T > 0$. Then we treat the following Cauchy problem for the evolution equation (DE) :

$$(CP; x_0) \quad \begin{cases} (d/dt)u(t) \in Au(t) & \text{for } t \in (0, T), \\ u(0) = x_0. \end{cases}$$

For the Cauchy problem (CP; x_0), we consider the following type of difference approximation :

$$(DS; x_0) \quad \begin{cases} \frac{x_k^n - x_{k-1}^n}{t_k^n - t_{k-1}^n} - \varepsilon_k^n \in Ax_k^n, & k=1, 2, \dots, N_n; n \geq 1, \\ x_0^n = x_0, \end{cases}$$

where for each n , $\{t_k^n\}$ represents the partition of $[0, T]$ such that $0 = t_0^n < t_1^n < \dots < t_{N_n-1}^n < T \leq t_{N_n}^n$ and $\delta_n = \max_{1 \leq k \leq N_n} (t_k^n - t_{k-1}^n) \rightarrow 0$ as $n \rightarrow \infty$. The ε_k^n may be referred as an error which occurs at the k -th step of the n -th approximation of the difference approximation. In this sense (DS; x_0) can be regarded as an approximating difference scheme for (CP; x_0) which permits errors.

Definition 2. Let $u_n(t)$ be a sequence in $L^\infty(0, T; X)$. We say that $u_n(t)$ is a (backward) DS-*approximate solution* of the Cauchy problem (CP; x_0) if there exists a difference approximation (DS; x_0) satisfying the followings :

- (i) $u_n(0) = x_0^n = x_0, n \geq 1$;
- (ii) $u_n(t) = x_k^n$ for $t \in (t_{k-1}^n, t_k^n] \cap (0, T], k=1, 2, \dots, N_n; n \geq 1$,
- (iii) $\sum_{k=1}^{N_n} \|\varepsilon_k^n\| (t_k^n - t_{k-1}^n) \rightarrow 0$ as $n \rightarrow \infty$.

Then we have

Theorem 1. Let $x_0 \in \overline{D(A)}$ and $u_n(t)$ be a DS-*approximate solution* of (CP; x_0) on $[0, T]$. Then there exists a $u(t) \in C([0, T]; X)$ satisfying the followings :

- (i) $u(t) = \lim_{n \rightarrow \infty} u_n(t)$ for $t \in [0, T]$, and the convergence is uniform on $[0, T]$;
- (ii) $u(t) \in \overline{D(A)}$ for $t \in [0, T]$ and $u(0) = x_0$;
- (iii) for any DS-*approximate solution* $\hat{u}_n(t)$ of (CP; x_0),

$$u(t) = \lim_{n \rightarrow \infty} \hat{u}_n(t) \quad \text{for } t \in [0, T].$$

Remarks. 1) Kenmochi-Oharu [5] and Takahashi [9], [10] studied the convergence (i) under the additional condition, which is called the stability condition by them. Our result is an extension of their results.

2) By Bénilan's method [2], we find that the limiting function $u(t)$ is the unique integral solution of the Cauchy problem (CP; x_0).

The proof of Theorem 1 is based on the following.

Lemma 2. *Let (DS; x_0) and (DS; \hat{x}_0) be two difference approximations as above of the Cauchy problems (CP; x_0) and (CP; \hat{x}_0) on $[0, T]$, respectively. Let the notations with the symbol “ \wedge ” represent the difference approximation (DS; \hat{x}_0). Then we have*

$$(1) \quad \begin{aligned} \|x_i^m - \hat{x}_j^n\| \leq & \|x_0 - u\| + \|\hat{x}_0 - u\| \\ & + \{(t_i^m - \hat{t}_j^n)^2 + \delta_m t_i^m + \delta_n \hat{t}_j^n\}^{1/2} \|Au\| \\ & + \sum_{k=1}^i \varepsilon_k^m \|t_k^m - t_{k-1}^m\| + \sum_{k=1}^j \varepsilon_k^n \|\hat{t}_k^n - \hat{t}_{k-1}^n\|, \end{aligned}$$

for $0 \leq i \leq N_m$, $0 \leq j \leq \hat{N}_n$ and $u \in D(A)$.

Lemma 2 is proved by the method of Crandall-Liggett [3], modified by Rasmussen [8] (see also Yosida [13]), by using Lemma 1.

Remark. Let A be a dissipative operator in X such that $R(I - \lambda A) \supset D(A)$ for $\lambda > 0$. Then the estimate (1) gives

$$\|(I - \lambda A)^{-n} x - (I - \mu A)^{-m} x\| \leq \{(n\lambda - m\mu)^2 + n\lambda^2 + m\mu^2\}^{1/2} \|Ax\|$$

for $n, m \geq 1$, $\lambda, \mu > 0$ and $x \in D(A)$. This estimate is similar to but different from that of Crandall-Liggett [3].

By virtue of Theorem 1, we define the following.

Definition 3. Let $u(t) \in C([0, T]; X)$ and $x_0 \in \overline{D(A)}$. We say that $u(t)$ is a (backward) DS-limit solution of the Cauchy problem (CP; x_0) on $[0, T]$ if there exists a (backward) DS-approximate solution $u_n(t)$ of (CP; x_0) on $[0, T]$, such that $u_n(t)$ converges to $u(t)$, uniformly for $t \in [0, T]$.

By Lemma 2, we have also

Corollary. *Let $u(t), \hat{u}(t)$ be two DS-limit solutions of (CP) on $[0, T]$. Then*

$$\|u(t) - \hat{u}(t)\| \leq \|u(0) - \hat{u}(0)\| \quad \text{for } t \in [0, T].$$

3. Generation of semigroups. By Theorem 1 and Corollary, we have a generation theorem of semigroups.

Definition 4. Let A be a quasi-dissipative operator in X . We say that A has the property (\mathfrak{D}) if for any $x \in \overline{D(A)}$ and $T > 0$, there exists a DS-approximate solution of the Cauchy problem (CP; x) on $[0, T]$.

Theorem 2. *Let A be a quasi-dissipative operator in X , having the property (\mathfrak{D}). Then there exists a contraction semigroup $\{T(t); t \geq 0\}$ on $\overline{D(A)}$ such that for each $x \in \overline{D(A)}$ and $T > 0$, $u(t) = T(t)x$ is the unique DS-limit solution of the Cauchy problem (CP; x) on $[0, T]$.*

We give a sufficient condition that a quasi-dissipative operator has the property (\mathfrak{D}) . Let A be a quasi-dissipative operator in X . We add the following condition on A :

for any $x \in \overline{D(A)}$, there exist a sequence $\delta_n \downarrow 0$ and $[x_n, y_n] \in A$ (R_t) ($n \geq 1$) such that

$$\lim_{n \rightarrow \infty} \delta_n^{-1} \|x_n - x - \delta_n y_n\| = 0.$$

Then we have

Theorem 3. *Let A be a quasi-dissipative operator in X , satisfying the condition (R_t) . Then A has the property (\mathfrak{D}) . Thus A generates a contraction semigroup on $\overline{D(A)}$, in the sense of Theorem 2.*

Remarks. 1) This theorem implies the fundamental result of Crandall-Liggett [3]; a part of the results of Martin [7] on ordinary differential equations; and the results of Webb [11] and Barbu [1] on the continuous perturbations of m -dissipative operators.

2) Yorke announces in [12] that he obtained a similar result.

Sketch of the proof of Theorem 3. Let $x_0 \in \overline{D(A)}$ and $\varepsilon_n \downarrow 0$. Let n be fixed. Then for each $x \in \overline{D(A)}$, we define

$$\delta_n(x) = \sup \{ \delta; 0 < \delta \leq \varepsilon_n \text{ and there exists } [x_\delta, y_\delta] \in A \text{ such that } \|x_\delta - x - \delta y_\delta\| \leq \delta \varepsilon_n \}.$$

Then each $\delta_n(x)$ is positive by the assumption. Therefore, inductively, we can choose $h_k^n > 0$ and $[x_k^n, y_k^n] \in A$, for $k=1, 2, \dots$, so that they satisfy the followings:

- (i) $x_0^n = x_0$;
- (ii) $(1/2)\delta_n(x_{k-1}^n) < h_k^n \leq \varepsilon_n$, for $k=1, 2, \dots$,
- (iii) $\|x_k^n - x_{k-1}^n - h_k^n y_k^n\| \leq h_k^n \varepsilon_n$, for $k=1, 2, \dots$.

Then we set $t_i^n = \sum_{k=1}^i h_k^n$. We may show that $t_i^n \rightarrow \infty$ as $i \rightarrow \infty$. For the purpose, we establish the following estimate:

$$(2) \quad \|x_i^n - x_j^n\| \leq (t_i^n - t_j^n) \|y_k^n\| + \varepsilon_n(t_i^n - t_k^n) + \varepsilon_n(t_j^n - t_k^n)$$

for any $i \geq j \geq k \geq 1$. This estimate may be verified by the induction for (i, j) with $i \geq j \geq k$ for each fixed $k \geq 1$, by using Lemma 1.

Acknowledgement. The author would like to express his hearty thanks to Prof. I. Miyadera and Mr. T. Takahashi for their valuable advices.

References

- [1] V. Barbu: Continuous perturbations of nonlinear m -accretive operators in Banach spaces. *Boll. U. M. I.*, **6**, 270–278 (1972).
- [2] Ph. Bénéilan: Equation d'évolution dans un espace Banach quelconque et applications. *Thèse Orsay* (1972).
- [3] M. Crandall and T. Liggett: Generation of semigroups of nonlinear transformations in general Banach spaces. *Amer. J. Math.*, **93**, 265–298 (1971).
- [4] T. Kato: Nonlinear semigroups and evolution equations. *J. Math. Soc. Japan*, **19**, 508–520 (1967).

- [5] N. Kenmochi and S. Oharu: Difference approximation of nonlinear evolution equations. *Publ. Res. Inst. Math. Sci.*, **10**, 265–520 (1974).
- [6] Y. Kobayashi: Difference approximation of Cauchy problems for quasi-dissipative operators and generation of nonlinear semigroups (to appear).
- [7] R. Martin: Differential equations on closed subsets of a Banach space. *Trans. Amer. Math. Soc.*, **179**, 399–414 (1973).
- [8] S. Rasmussen: Nonlinear semigroups, evolution equations and product integral representations. *Various Publication Series, No. 2, Aarhus Universitet* (1971/72).
- [9] T. Takahashi: Difference approximation of Cauchy problems for quasi-dissipative operators and generation of semigroups of nonlinear contractions (to appear).
- [10] —: Convergence of difference approximation of nonlinear evolution equations and generation of semigroups (to appear).
- [11] G. Webb: Continuous nonlinear perturbations of linear accretive operators in Banach spaces. *J. Funct. Anal.*, **10**, 191–203 (1972).
- [12] J. Yorke: Ordinary differential equations and evolutionary equations (preprint).
- [13] K. Yosida: *Functional Analysis (Fourth Edition)*. Springer-Verlag (1974).