

NON-LINEAR DIFFERENTIAL EQUATIONS ON CONES IN BANACH SPACES

CHARLES V. COFFMAN

1. Introduction. Let y denote a vector in n -dimensional Euclidean space E^n , $y \geq 0$ means that each component of y is nonnegative. Let $f(t, y)$ be a function with values in E^n defined on the domain $D = \{(t, y), 0 \leq t < \infty, y \in E^n, y \geq 0\}$. Assume that $f(t, y)$ is continuous on D and that for each $(t, y) \in D$, $-f(t, y) \geq 0$, and that $f(t, 0) \equiv 0$. Let C be a positive constant. It is known, [*, 4], that the differential equation $dy/dt = f(t, y)$ possesses at least one solution $y = y(t)$, defined for $0 \leq t < \infty$, with $y(t) \geq 0$ for $0 \leq t < \infty$, and satisfying $|y(0)| = C$, where $|\dots|$ denotes the Euclidean norm in E^n . (Actually the hypothesis of the theorem referred to involves an additional condition on $f(t, y)$, however the proof given in [4] can be modified so as to avoid the need for this condition.)

It was shown in [2] that the linear case of the result just quoted has a generalization to an infinite dimensional case. Before describing this result it will be necessary to introduce the following terminology and notation. Throughout the remainder of the paper Y will denote a (real) Banach space with norm $\|\dots\|$ and K some fixed, closed convex cone in Y . For $y \in Y$, $y \geq 0$ will mean $y \in K$. A linear operator A on Y will be said to be nonnegative if $Ay \geq 0$ whenever $y \geq 0$. A nonempty set of the form $H = H(y^*) = \{y : y \in K, y^*(y) = 1\}$, where y^* is in the dual space Y^* of Y will be called a cross-section of the cone K . Theorem 1 of [2] concerns the linear differential equation on the Banach space Y ,

$$(1.1) \quad dy/dt = -A(t)y,$$

where for each t , $0 \leq t < \infty$, $A(t)$ is a nonnegative bounded linear operator on Y , and $A(t)$ is strongly continuous on $0 \leq t < \infty$. The result states that if the cone K has a weakly compact cross-section then the differential equation (1.1) has a nontrivial solution $y = y(t)$ satisfying $y(t) \geq 0$ for $0 \leq t < \infty$.

Theorem 3 of [2] is the dual of Theorem 1 and concerns the differential equation on Y^* adjoint to (1.1)

$$(1.2) \quad dy^*/dt = -A^*(t)y^*,$$

where for each t , $0 \leq t < \infty$, $A^*(t)$ is the adjoint of $A(t)$. Let $A(t)$ be as above, this theorem states that if K has an interior point then (1.2) has a nontrivial solution $y^* = y^*(t)$ satisfying $y^*(t) \in K^*$ for $0 \leq t < \infty$. K^* in Y^* is the dual cone of the cone K in Y , consisting of those

elements y^* such that $y^*(y) \geq 0$ for each $y \in K$.

The main result of this note, Theorem 4.1, generalizes the result of [4] mentioned above to an infinite dimensional case and contains both of the results just quoted.

The following standard terminology will be used. A closed linear manifold Γ in the dual space Y^* of Y is called *determining* for Y if, when $y \in Y$, $\gamma(y) = 0$ for each $\gamma \in \Gamma$ implies $y = 0$. Such a manifold defines a topology, referred to as the weak Γ -topology, on Y . The generalized sequence $\{y_\alpha\}$ of elements of Y has limit 0 in the weak Γ -topology if and only if $\lim_\alpha \gamma(y_\alpha) = 0$ for each $\gamma \in \Gamma$.

2. Differential equations on a Banach space. Let Γ be a closed linear manifold in Y^* which is determining for Y . A function $y(t)$ on some interval I on the real line, with values in Y , will be said to have a weak Γ -derivative, or to be weakly Γ -differentiable at $t \in I$ if the limit as $h \rightarrow 0$ of the difference quotient $(y(t+h) - y(t))/h$ exists in the weak Γ -topology. If $y(t)$ is weakly Γ -differentiable on I , its weak Γ -derivative will be denoted by $D_\Gamma y(t)$. Let C be a subset of Y and let I be, as above, an interval on the real line. Consider the differential equation

$$(2.1) \quad dy/dt = f(t, y)$$

where $f(t, y)$ is a function from $I \times C \rightarrow Y$, continuous in the weak Γ -topology. Let $y(t)$ be a function on I with values in C . If $y(t)$ is strongly continuous and strongly differentiable on I with a strongly continuous derivative dy/dt , and if (2.1) holds, $y(t)$ will be said to be a strong solution of (2.1) on I . If $y(t)$ is strongly continuous, weakly absolutely continuous and strongly differentiable a.e. on I , and if (2.1) holds a.e. on I , $y(t)$ will be said to be a strong solution in the extended sense of (2.1) on I . Finally $y(t)$ will be said to be a weak Γ -solution of (2.1) on I if it is weakly Γ -differentiable on I and if $D_\Gamma y(t) = f(t, y(t))$ on I . Obviously if $y(t)$ is a weak Γ -solution of (2.1) then $D_\Gamma y(t)$ is weakly Γ -continuous.

LEMMA 2.1. *If $y(t)$ is a strong solution in the extended sense of (2.1) on I , then it can be expressed as the Bochner integral of its derivative.*

Proof. Since $y(t)$ is strongly continuous, its range lies in a separable subspace of Y . Clearly the values of dy/dt , where it exists will be in the same subspace. Put

$$(2.2) \quad u(t) = f(t, y(t)) ,$$

then $u(t)$ is weakly Γ -continuous and coincides with dy/dt a.e. where the latter is defined. Since $u(t)$ is almost separably valued and weakly Γ -continuous, it follows from Theorem 1.1.7, [3, p. 330], that $u(t)$, hence dy/dt is locally Bochner integrable on I . Since $y(t)$ is weakly absolutely continuous it may be expressed as the indefinite integral of dy/dt , Theorem 3.8.6, [5].

LEMMA 2.2. *Let $y(t)$ be a function defined on I with values in C . In order that $y(t)$ be a strong solution in the extended sense of (2.1), it is necessary and sufficient that $y(t)$ be a weak Γ -solution of (2.1) and that $D_r y(t)$ be almost separably valued.*

REMARK. If $\Gamma = Y^*$ and if $y(t)$ is a weak Γ -solution of (2.1), then $D_r y(t)$ is necessarily separably valued, since it is weakly Γ -continuous; see [5], p. 59.

Proof of Lemma 2.2. Necessity. Let $y(t)$ be a strong solution in the extended sense of (2.1). Let $u(t)$ be defined by (2.2). From the proof of Lemma 2.1, $u(t)$ is almost separably valued and for $t, t' \in I$, $y(t) - y(t') = \int_{t'}^t u(t) dt$. For each $\gamma \in \Gamma$, $\gamma(u(t))$ is continuous, consequently $\gamma(y(t)) \in C^1$ and $d[\gamma(y(t))]/dt = \gamma(u(t))$, since one has $\gamma\left[\int_{t'}^t u(t) dt\right] = \int_{t'}^t \gamma(u(t)) dt$. It follows that $D_r y(t)$ exists everywhere on I and that $D_r y(t) = u(t)$.

Sufficiency. Let $y(t)$ be a weak Γ -solution of (2.1) and let $D_r y(t)$ be almost separably valued. By Theorem 1.1.7, [3], $D_r y(t)$ is locally Bochner integrable on I . For each $\gamma \in \Gamma$, $\gamma(D_r y(t))$ is continuous on I , so for $t, t' \in I$, $\gamma(y(t)) - \gamma(y(t')) = \int_{t'}^t \gamma(D_r y(t)) dt = \gamma\left[\int_{t'}^t D_r y(t) dt\right]$. Since Γ is determining for Y it follows that $y(t)$ can be expressed as the indefinite Bochner integral of $D_r y(t)$. Hence $y(t)$ is in fact strongly absolutely continuous, and, by Corollary 2, p. 88, [5], $y(t)$ is strongly differentiable a.e., and dy/dt coincides a.e., where it is defined with $D_r y(t)$.

3. Convergence of approximate solutions. Let Y, Γ, C, I , and $f(t, y)$ be as in § 2. The following generalization of the notion of an ϵ -approximate solution of a differential equation will be used. Let V be a neighborhood of 0 in the weak Γ -topology. A function $y(t)$ will be called a V -approximate weak Γ -solution of (2.1) on I if

- (i) $y(t)$ is defined on I , has values in C and is weakly Γ -continuous;
- (ii) for some finite set of points S on I , $D_r y(t)$ exists and is weakly Γ -continuous on $I - S$, further $D_r y(t)$ has only simple discontinuities, in the weak Γ -topology, at points of S ;
- (iii) $(D_r y(t) - f(t, y)) \in V$ for $t \in I - S$.

REMARK. In view of the principle of uniform boundedness (ii) implies that $\|D_r y(t)\|$ is locally bounded on I .

LEMMA 3.1. *Let $\{y_n(t)\}$ be a sequence of weakly Γ -continuous functions defined on I with values in C . Let C be compact in the weak Γ -topology. For each neighborhood V of 0, in the weak Γ -topology, and for each compact subinterval I' of I , let there exist an $N = N(V, I')$ such that for all $n \geq N$, $y_n(t)$ is a V -approximate weak Γ -solution of (2.1) on I' . Then there exists a subsequence $\{y_{n_k}(t)\}$ which converges uniformly on every compact subinterval of I to a weak Γ -solution $y(t)$ of (2.1) on I .*

Proof. The existence of a subsequence convergent on each compact subinterval of I will follow from Ascoli's theorem, [1, p. 43], if it is shown that the sequence is equicontinuous in the weak Γ -topology. Let $t_0 \in I$ be arbitrary, it will suffice to show that for each $\gamma \in \Gamma$, and for each $\varepsilon > 0$ there exists a $\delta = \delta(\varepsilon, \gamma, t_0)$ such that for $n = 1, 2, \dots$

$$(3.1) \quad |\gamma(y_n(t) - y_n(t_0))| < \varepsilon$$

whenever $t \in I$, and $|t - t_0| < \delta$. For some $\delta_0 > 0$, there exists, by the principle of uniform boundedness, an M such that $\|f(t, y)\| \leq M$ for $t \in I$, $|t - t_0| \leq \delta_0$ and $y \in C$. For $n = 1, 2, \dots$, one has

$$(3.2) \quad \begin{aligned} \gamma(y_n(t)) - \gamma(y_n(t_0)) &= \int_{t_0}^t \gamma(f(t, y_n(t))) dt \\ &+ \int_{t_0}^t \gamma(D_r y_n(t) - f(t, y_n(t))) dt . \end{aligned}$$

For all sufficiently large n , say $n \geq n_0$, $|\gamma(D_r y_n(t) - f(t, y_n(t)))| < \varepsilon/2 \|\gamma\| \delta_0$, for $t \in I$, $|t - t_0| < \delta_0$. Thus, if $\delta_1 < \min(\delta_0, \varepsilon/2M \|\gamma\| \delta_0)$ then (3.1) holds for $t \in I$, $|t - t_0| < \delta_1$ and $n \geq n_0$. Clearly it is possible to choose a δ , $0 < \delta \leq \delta_1$, such that (3.1) holds for all n when $t \in I$, $|t - t_0| < \delta$.

Let the subsequence $\{y_{n_k}(t)\}$ of the original sequence be uniformly convergent on each compact subinterval of I to $y(t)$. For each $\gamma \in \Gamma$, $\gamma(f(t, y_{n_k}(t))) \rightarrow \gamma(f(t, y(t)))$ and $\gamma(D_r y_{n_k}(t) - f(t, y_{n_k}(t))) \rightarrow 0$ uniformly on each compact subinterval of I . It follows that $\gamma(y(t) - y(t_0)) = \int_{t_0}^t \gamma(f(t, y(t))) dt$, for $t, t_0 \in I$, for each $\gamma \in \Gamma$. Consequently $D_r y(t)$ exists on I and $D_r y(t) = f(t, y(t))$ there.

4. An existence theorem. As a consequence of the results of the last section, one has the following local existence theorem for the problem (2.1).

THEOREM 4.1. *Let the unit sphere in Y be compact in the weak*

Γ -topology. For some real interval I , some point $y_0 \in Y$ and some positive constant k let $f(t, y)$ be defined and weakly Γ -continuous for $t \in I, \|y - y_0\| \leq k$. Let t_0 be in the interior of I , then on some open subinterval I' of I , with $t_0 \in I'$, there exists a weak Γ -solution $y(t)$ of (2.1) satisfying the initial condition

$$(4.1) \quad y(t_0) = y_0.$$

Proof. It can be assumed that I is compact, so that $\|f(t, y)\| \leq M < \infty$ for $t \in I, \|y - y_0\| \leq k$. Let δ be some number on the range $0 < \delta < k/M$, such that the interval $I' = \{t : |t - t_0| < \delta\} \subset I$. For a positive integer n , subdivide I' by the points $t_i = t_0 + l\delta/n, l = 0, \pm 1, \dots, \pm n - 1$. Define a function $y_n(t)$ on I' as follows: let $y_n(t_0) = y_0$, and let $y_n(t) = y_0 + (t - t_0)f(t_0, y_0)$ for $t_0 \leq t \leq t_1$, clearly then $\|y_n(t) - y_0\| \leq M(t_1 - t_0) \leq k/n$ for $t_0 \leq t \leq t_1$. Assume that $y_n(t)$ is defined for $t_0 \leq t \leq t_i, i \leq n - 1$, with $\|y_n(t) - y_0\| \leq lk/n$ for $t_0 \leq t \leq t_i$, and put $y_n(t) = y_n(t_i) + (t - t_i)f(t_i, y_n(t_i))$ for $t_i \leq t \leq t_{i+1}$. Thus $\|y_n(t) - y_0\| \leq (l + 1)k/n$, and the process can be continued to define $y_n(t)$ on $t_0 \leq t < \delta$, with $\|y_n(t) - y_0\| < k$ there. $y_n(t)$ is defined to the left of t_0 in a similar fashion.

Let $y_n(t)$ be so defined for $n = 1, 2, \dots$. The interval I being compact, $f(t, y)$ is uniformly weakly Γ -continuous for $t \in I, \|y - y_0\| \leq k$. It readily follows that, given $V \subset Y$, a neighborhood of zero in the weak Γ -topology, then for all sufficiently large n , $y_n(t)$ is a V -approximate weak Γ -solution of (2.1) on I' .

The existence of a weak Γ -solution of (2.1) on I' satisfying the initial condition (4.1) now follows from Lemma 3.1.

5. Differential equations on cones. The main result of this note is the following.

THEOREM 5.1. *Let K be a closed convex cone in the Banach space Y . Let Γ be as above, and let there exist a $\gamma_0 \in K^* \cap \Gamma$ such that the cross-section $H(\gamma_0)$ of K is compact in the weak Γ -topology on Y . Let $I = [0, \infty)$ and let $f(t, y)$ be defined on $I \times K$, and continuous in the weak Γ -topology, and suppose that*

$$(5.1) \quad -f(t, y) \in K \quad \text{for } (t, y) \in K.$$

assume also that

$$(5.2) \quad f(t, 0) \equiv 0.$$

Then there exists a weak Γ -solution $y(t)$ of (2.1) on I with $y(t) \in K$ for all $t \in I$, and with $y(0) \in H$.

Proof. First it will be shown that for each $T > 0$ there exists a weak Γ -solution $y_T(t)$ of (2.1) on $[0, T]$ with $y_T(t) \in K$ for $0 \leq t \leq T$, and having $y_T(0) \in H$. Let $T > 0$ be given, let n be a positive integer and let η be an arbitrary element of K . Subdivide $[0, T]$ into n subintervals of equal length, and construct a polygonal approximate solution $y_n(t) = y_n(t, \eta)$ as in § 4, beginning at $t = T$ with $y_n(T, \eta) = \eta$ and proceeding to the left. That is, $t_k = k T/n, k = 0, \dots, n$, being the endpoints of the intervals of the subdivision, if $y_n(t, \eta)$ is assumed to be defined on $[t_k, T], k > 0$, the definition is extended to $[t_{k-1}, T]$ by putting

$$(5.3) \quad y_n(t, \eta) = y_n(t_k, \eta) + (t - t_k)f(t_k, y_n(t_k, \eta))$$

on $[t_{k-1}, t_k]$. Notice that because of (5.1) at no point can the polygon leave K .

From the fact that $f(t, y)$ is weakly Γ -continuous it readily follows, using (5.3) and an induction argument that the mapping of $K \rightarrow K$ given by $\eta \rightarrow y_n(0, \eta)$ is continuous in the weak Γ -topology. From (5.2), it follows that $y_n(0, 0) = 0$. Because of (5.1) one has, in view of (5.3) that $y_n(t, \eta) \geq y_n(t', \eta)$ for $0 \leq t < t' \leq T$, and consequently that

$$(5.4) \quad \gamma_0(y_n(t, \eta)) \geq \gamma_0(y_n(t', \eta)) \quad \text{for } 0 \leq t < t' \leq T.$$

The weak Γ -continuity of the mapping $\eta \rightarrow y_n(0, \eta)$ implies therefore, the existence of values $\eta \in K$ for which $y_n(0, \eta) \in H$. For each $n = 1, 2, \dots$, let η_n be chosen so that $y_n(0, \eta_n) \in H$. By (5.4) one has that for $n = 1, 2, \dots y_n(t, \eta_n) \in K_1$ for $0 \leq t \leq T$, where $K_1 = \{y : y \in K, \gamma_0(y) \leq 1\}$. Since K_1 is weak Γ -compact, $f(t, y)$ is uniformly weakly Γ -continuous on $[0, T] \times K_1$, thus it follows, as in the proof of Theorem 4.1, that the sequence $y_n(t, \eta_n)$ satisfies the hypotheses of Lemma 3.1. Consequently that lemma implies that there exists a weak Γ -solution of (2.1) on $[0, T]$ such that $y_T(t) \in K_1$ on $[0, T]$ and $y_T(0) \in H$.

For each $n = 1, 2, \dots$, let $y_n(t)$ be a weakly Γ -continuous function from I to K_1 with $y_n(0) \in H$ and such that on $[0, n], y_n(t)$ is a weak Γ -solution of (2.1). The sequence $\{y_n(t)\}$ satisfies the hypotheses of Lemma 3.1, and the conclusion of the theorem follows by an application of that lemma.

Theorem 3 of [2] follows immediately from Theorem 4.1 above in view of the observation (ii) of [2], namely that the dual cone K^* of a cone K with interior has a weak* compact cross-section. If $\Gamma = Y^*$ then the solution of (2.1) whose existence is asserted in Theorem 4.1 is, by Lemma 2.2 and the remark following the statement of that lemma, a strong solution, in the extended sense, of (2.1) on I . If the mapping $f(t, y)$ of $I \times K$ to $-K$ is continuous with respect to the strong topology on Y it is not difficult to see that a strong solution in the

extended sense, of (2.1) is in fact a strong solution of (2.1). Thus Theorem 1 of [2] also follows from Theorem 5.1.

If Y is finite dimensional, Theorem 5.1 implies a result similar to Theorem (*) of [4] referred to in § 1. In the latter result the initial value $y(0)$ is asserted to lie on the surface of a sphere $\|y\| = C, C > 0$, rather than on some hyperplane. For the finite dimensional case it is clear that the proof given above can be modified to give this different normalization of the initial value.

REFERENCES

1. N. Bourbaki, *Topologie Générale*, Chap. X, Actualities Scientifiques et Industrielles n. 1084, Paris 1949.
2. C. V. Coffman, *Linear differential equations on cones in Banach spaces*, Pacific J. Math., **12** (1962), 69-75.
3. N. Dunford and B. J. Pettis, *Linear operations on summable functions*, Trans. Amer. Math. Soc., **47** (1940), 323-391.
4. P. Hartman and A. Wintner, *On monotone solutions of non-linear differential equations*, Amer. J. Math., **76** (1954), 860-886.
5. E. Hille and R. S. Phillips, *Functional analysis and semi-groups*, Amer. Math. Soc. Colloquium Publ. vol. XXXI, 2nd ed. 1957.

