

WEAK SOLUTIONS TO 3-D CAHN–HILLIARD SYSTEM IN ELASTIC SOLIDS

IRENA PAWŁOW — WOJCIECH M. ZAJĄCZKOWSKI

ABSTRACT. In this paper we prove the existence and some time regularity of weak solutions to a three-dimensional (3-D) Cahn–Hilliard system coupled with nonstationary elasticity. Such nonlinear parabolic-hyperbolic system arises as a model of phase separation in deformable alloys. The regularity result is based on the analysis of time differentiated problem by means of the Faedo–Galerkin method. The obtained regularity provides a first step to the proof of strong solvability of the problem to be presented in a forthcoming paper [22].

1. Introduction

The present paper is concerned with the existence and regularity of weak solutions to a three-dimensional (3-D) Cahn–Hilliard system coupled with nonstationary elasticity. Such system arises as a model of phase separation in a binary deformable alloy quenched below a critical temperature. The problem under consideration has the following form:

$$(1.1) \quad \begin{aligned} \mathbf{u}_{tt} - \nabla \cdot W_{,\varepsilon}(\varepsilon(\mathbf{u}), \chi) &= \mathbf{b} && \text{in } \Omega^T = \Omega \times (0, T), \\ \mathbf{u}|_{t=0} = \mathbf{u}_0, \quad \mathbf{u}_t|_{t=0} &= \mathbf{u}_1 && \text{in } \Omega, \\ \mathbf{u} &= \mathbf{0} && \text{in } S^T = S \times (0, T), \end{aligned}$$

2000 *Mathematics Subject Classification.* 35K50, 35K60, 35L20, 35Q72.

Key words and phrases. Cahn–Hilliard, nonstationary elasticity, phase separation, weak solutions.

The paper partially supported by the Polish Grant No 1/P03A/02130.

$$(1.2) \quad \begin{aligned} \chi_t - \nabla \cdot \mathbf{M}\nabla\mu &= 0 && \text{in } \Omega^T, \\ \chi|_{t=0} &= \chi_0 && \text{in } \Omega, \\ \mathbf{n} \cdot \mathbf{M}\nabla\mu &= 0 && \text{on } S^T, \end{aligned}$$

$$(1.3) \quad \begin{aligned} \mu &= -\nabla \cdot \mathbf{\Gamma}\nabla\chi + \psi'(\chi) + W_{,\chi}(\varepsilon(\mathbf{u}), \chi) && \text{in } \Omega^T, \\ \mathbf{n} \cdot \mathbf{\Gamma}\nabla\chi &= 0 && \text{on } S^T. \end{aligned}$$

Here $\Omega \subset \mathbb{R}^3$ is a bounded domain with a smooth boundary S , occupied by a body in a reference configuration with constant mass density $\rho = 1$; \mathbf{n} is the unit outward normal to S , and $T > 0$ is an arbitrary fixed time. The body is a binary $a - b$ alloy.

The unknowns are the fields \mathbf{u} , χ and μ , where $\mathbf{u}: \Omega^T \rightarrow \mathbb{R}^3$ is the *displacement vector*, $\chi: \Omega^T \rightarrow \mathbb{R}$ is the *order parameter* (phase ratio) and $\mu: \Omega^T \rightarrow \mathbb{R}$ is the chemical potential difference between the components, shortly referred to as the *chemical potential*.

The second order tensor

$$\varepsilon = \varepsilon(\mathbf{u}) = \frac{1}{2}(\nabla\mathbf{u} + (\nabla\mathbf{u})^T)$$

denotes the linearized strain tensor.

In case of a binary $a - b$ alloy the order parameter is related to the volumetric fraction of one of the two phases, characterized by different crystalline structures of the components, e.g. $\chi = -1$ is identified with the phase a and $\chi = 1$ with the phase b .

The function $W(\varepsilon(\mathbf{u}), \chi)$ denotes the elastic energy defined by

$$(1.4) \quad W(\varepsilon(\mathbf{u}), \chi) = \frac{1}{2}(\varepsilon(\mathbf{u}) - \bar{\varepsilon}(\chi)) \cdot \mathbf{A}(\varepsilon(\mathbf{u}) - \bar{\varepsilon}(\chi)).$$

The corresponding derivatives

$$\begin{aligned} W_{,\varepsilon}(\varepsilon(\mathbf{u}), \chi) &= \mathbf{A}(\varepsilon(\mathbf{u}) - \bar{\varepsilon}(\chi)), \\ W_{,\chi}(\varepsilon(\mathbf{u}), \chi) &= -\bar{\varepsilon}'(\chi) \cdot \mathbf{A}(\varepsilon(\mathbf{u}) - \bar{\varepsilon}(\chi)) \end{aligned}$$

represent respectively the stress tensor and the elastic part of the chemical potential.

The fourth order tensor $\mathbf{A} = (A_{ijkl})$ denotes a constant elasticity tensor given by

$$(1.5) \quad \varepsilon(\mathbf{u}) \mapsto \mathbf{A}\varepsilon(\mathbf{u}) = \bar{\lambda} \text{tr}\varepsilon(\mathbf{u})\mathbf{I} + 2\bar{\mu}\varepsilon(\mathbf{u}),$$

where $\mathbf{I} = (\delta_{ij})$ is the identity tensor, and $\bar{\lambda}$, $\bar{\mu}$ are the Lamé constants with values within the elasticity range (see Section 2). The form (1.5) refers to the isotropic, homogeneous medium with the same elastic properties of the phases.

The second order tensor $\bar{\varepsilon}(\chi)$ denotes the eigenstrain, i.e. the stress free strain corresponding to the phase ratio χ , defined by

$$(1.6) \quad \bar{\varepsilon}(\chi) = (1 - z(\chi))\bar{\varepsilon}_a + z(\chi)\bar{\varepsilon}_b,$$

with $\bar{\varepsilon}_a, \bar{\varepsilon}_b$ denoting constant eigenstrains of the phases a, b , and $z: \mathbb{R} \rightarrow [0, 1]$ being a sufficiently smooth interpolation function (called shape function) satisfying

$$(1.7) \quad z(\chi) = 0 \quad \text{for } \chi \leq -1 \quad \text{and} \quad z(\chi) = 1 \quad \text{for } \chi \geq 1.$$

Furthermore, the function $\psi: \mathbb{R} \rightarrow \mathbb{R}$ denotes the chemical energy of the system at zero stress. This function depends on temperature and is convex above a critical temperature and a nonconvex for temperatures less than the critical one. Here we assume it in the simplest double-well form

$$(1.8) \quad \psi(\chi) = \frac{1}{4}(1 - \chi^2)^2$$

with two minima at $\chi = -1$ and $\chi = 1$.

The second order tensors $\mathbf{M} = (M_{ij})$ and $\mathbf{\Gamma} = (\Gamma_{ij})$ represent respectively the mobility matrix and the interfacial energy matrix. For simplicity, we shall confine ourselves to the isotropic, homogeneous situation assuming that

$$(1.9) \quad \mathbf{M} = M\mathbf{I}, \quad \mathbf{\Gamma} = \Gamma\mathbf{I}, \quad M = \Gamma = 1,$$

with positive constants M, Γ normalized to unity.

System (1.1)–(1.3) represents respectively the linear momentum balance, the mass balance and a generalized equation for the chemical potential. In a thermodynamical theory due to Gurtin [10] equation (1.3) is identified with a microforce balance. The free energy density underlying system (1.1)–(1.3) has the Landau–Ginzburg–Cahn–Hilliard form

$$(1.10) \quad f(\varepsilon(\mathbf{u}), \chi, \nabla\chi) = W(\varepsilon(\mathbf{u}), \chi) + \psi(\chi) + \frac{1}{2}\nabla\chi \cdot \mathbf{\Gamma}\nabla\chi,$$

with the three terms on the right-hand side representing respectively the elastic, chemical and interfacial energy.

The remaining quantities in (1.1)–(1.3) have the following meaning: $\mathbf{b}: \Omega^T \rightarrow \mathbb{R}^3$ represents the external body force, and $\mathbf{u}_0, \mathbf{u}_1: \Omega \rightarrow \mathbb{R}^3, \chi_0: \Omega \rightarrow \mathbb{R}$ are the initial conditions respectively for the displacement, the velocity and the order parameter.

The homogeneous boundary conditions in (1.1)–(1.3) are chosen for the sake of simplicity. The condition (1.1)₃ means that the body is fixed at the boundary S , (1.2)₃ reflects the mass isolation at S , and (1.3)₂ is the natural boundary condition for (1.10).

Before discussing the results of the paper let us place our study in the present theory of Cahn–Hilliard systems in elastic solids. In recent years such systems have been the subject of many different modelling, mathematical and numerical studies, we refer e.g. to [18], [19], [1], [21] for up to date references. It is known from the materials science literature that the elastic effects have a pronounced effect on the microstructure evolution of the phase separation process and consequently on the resulting material properties.

The most general setting of the Cahn–Hilliard system coupled with elasticity, accounting for additional anisotropic, heterogeneous and kinetic effects, was derived by Gurtin [10] within the frame of his thermodynamical theory of phase transitions based on a microforce balance. System (1.1)–(1.3) represents a simplified version of Gurtin’s model with neglected anisotropic, heterogeneous and kinetic effects; for more details see [1] where the full Gurtin’s model was studied.

In view of the fact that the mechanical equilibrium is usually attained on a much faster time scale than diffusion in most of the studies a quasi-stationary approximation of (1.1)₁, obtained by neglecting the inertial term \mathbf{u}_{tt} , was assumed. Various variants of the Cahn–Hilliard system with quasi-stationary elasticity were analyzed by Garcke [7]–[9], Bonetti et al. [2], Carrive et al. [4] and [5], Miranville [14]–[17]. We stress on the fact that the results of [7], [8], [2] included mathematically difficult case of nonhomogeneous elasticity with tensor $\mathbf{A} = \mathbf{A}(\chi)$ depending on the order parameter.

The Cahn–Hilliard system with nonstationary elasticity was studied in [15] and [1] where the existence and properties of weak solutions were examined, and in [21] where the classical solvability was proved in 1-D case. It is clear that with the quasi-stationary hypothesis the hyperbolic elasticity system is replaced by the elliptic one and thereby the mathematical analysis becomes qualitatively different.

We point out that the study of the Cahn–Hilliard problem with nonstationary elasticity — apart from the mathematical interest on its own — is of special importance for the initial stages of phase separation at which the formation of the microstructure is on a very fast time scale.

The goal of the present paper is to prove the existence and some regularity of weak solutions to system (1.1)–(1.3). Our ultimate aim is to obtain the existence and uniqueness of a strong solution to (1.1)–(1.3), i.e. such a solution that its all derivatives appearing in the equations are at least in L_2 . The strong solvability theory will be presented separately in [22]. It is based on the regularity results proved in the present paper together with some additional time regularity. More precisely, having sufficiently high time regularity we shall apply the standard elliptic regularity theory to conclude further space regularity, and consequently the strong solvability.

The main results of the present paper were outlined in [23].

As was already mentioned, the strong solvability of system (1.1)–(1.3) in 1-D case was proved in [21]. The strong solvability of the single Cahn–Hilliard equation in 1-D and 3-D cases was analyzed first by Elliott and Zheng [6].

We point out that in three space dimensions the coupled system (1.1)–(1.3) shows features that make its analysis much more difficult than in one-dimensional setting. The arguments used by the authors in the single space dimension in [21], based on the space regularity of the wave equation, do not extend to the 3-D case.

The key idea of the regularity theory presented here and in the forthcoming paper [22] consists in the analysis of time-differentiated versions of problem (1.1)–(1.3) which yield solutions with sufficiently high time derivatives. The analysis is performed with the help of the Faedo–Galerkin approximation. The procedure is straightforward except for some difficulties of technical nature due to many nonlinear terms that appear in the system after differentiation with respect to time variable.

For further analysis it is convenient to introduce a simplified formulation of problem (1.1)–(1.3) which results on account of particular constitutive equations (1.4)–(1.6) and (1.9). Let \mathbf{Q} be the linear elasticity operator defined by

$$(1.11) \quad \mathbf{u} \mapsto \mathbf{Q}\mathbf{u} = \nabla \cdot \mathbf{A}\varepsilon(\mathbf{u}) = \bar{\mu}\Delta\mathbf{u} + (\bar{\lambda} + \bar{\mu})\nabla(\nabla \cdot \mathbf{u}).$$

Moreover, let us denote

$$(1.12) \quad \mathbf{B} = -\mathbf{A}(\bar{\varepsilon}_b - \bar{\varepsilon}_a), \quad D = -\mathbf{B} \cdot (\bar{\varepsilon}_b - \bar{\varepsilon}_a), \quad E = -\mathbf{B} \cdot \bar{\varepsilon}_a,$$

where $\mathbf{B} = (B_{ij})$ is a symmetric second order tensor, and D, E are two scalars. With such a notation we have

$$(1.13) \quad \nabla \cdot W_{,\varepsilon}(\varepsilon(\mathbf{u}), \chi) = \nabla \cdot \mathbf{A}\varepsilon(\mathbf{u}) - \nabla \cdot \mathbf{A}(\bar{\varepsilon}_a + z(\chi)(\bar{\varepsilon}_b - \bar{\varepsilon}_a)) = \mathbf{Q}\mathbf{u} + z'(\chi)\mathbf{B}\nabla\chi,$$

and

$$W_{,\chi}(\varepsilon(\mathbf{u}), \chi) = z'(\chi)(\mathbf{B} \cdot \varepsilon(\mathbf{u}) + Dz(\chi) + E),$$

so that (1.1)–(1.3) simplifies to

$$(1.14) \quad \begin{aligned} \mathbf{u}_{tt} - \mathbf{Q}\mathbf{u} &= z'(\chi)\mathbf{B}\nabla\chi + \mathbf{b} && \text{in } \Omega^T, \\ \mathbf{u}|_{t=0} &= \mathbf{u}_0, \quad \mathbf{u}_t|_{t=0} = \mathbf{u}_1 && \text{in } \Omega, \\ \mathbf{u} &= \mathbf{0} && \text{on } S^T, \end{aligned}$$

$$(1.15) \quad \begin{aligned} \chi_t - \Delta\mu &= 0 && \text{in } \Omega^T, \\ \chi|_{t=0} &= \chi_0 && \text{in } \Omega, \\ \mathbf{n} \cdot \nabla\mu &= 0 && \text{on } S^T, \end{aligned}$$

$$(1.16) \quad \begin{aligned} \mu &= -\Delta\chi + \psi'(\chi) + W_{,\chi}(\varepsilon(\mathbf{u}), \chi) && \text{in } \Omega^T, \\ \mathbf{n} \cdot \nabla\chi &= 0 && \text{on } S^T, \end{aligned}$$

with $W_{,\chi}(\varepsilon(\mathbf{u}), \chi)$ given by (1.13)₂.

Let us note that the combined systems (1.15) and (1.16) yield the following Cahn–Hilliard problem

$$(1.17) \quad \begin{aligned} \chi_t + \Delta^2\chi &= \Delta[\psi'(\chi) + z'(\chi)(\mathbf{B} \cdot \varepsilon(\mathbf{u}) + Dz(\chi) + E)] && \text{in } \Omega^T, \\ \chi|_{t=0} &= \chi_0 && \text{in } \Omega, \\ \mathbf{n} \cdot \nabla\chi &= 0 && \text{on } S^T, \\ \mathbf{n} \cdot \nabla\Delta\chi &= z'(\chi)\mathbf{n} \cdot \nabla(\mathbf{B} \cdot \varepsilon(\mathbf{u})) && \text{on } S^T, \end{aligned}$$

coupled with the elasticity system (1.14). It is seen that the problems are coupled not only through the right-hand sides but also through the boundary conditions. Moreover, by definition (1.7) of the shape function z , the problems decouple for $\chi \leq -1$ and $\chi \geq 1$. We point out that the boundary coupling is characteristic for the multidimensional problem and does not appear in its one-dimensional setting. In fact, in 1-D case assuming that $b = 0$ on S^T , it follows from (1.14)₁, (1.14)₃ and (1.17)₃ that $u_{xx} = 0$ on S^T , and consequently condition (1.17)₄ yields $\chi_{xxx} = 0$ on S^T . This fact was used in [21] in the analysis of the 1-D version of problem (1.1)–(1.3).

The plan of the paper is as follows: In Section 2 we present our main assumptions and results stated in Theorems 2.1 and 2.2. Theorem 2.1 asserts the existence of a weak solution to problem (1.1)–(1.3). Theorem 2.2 provides a time-regularity result obtained by differentiating (1.1)–(1.3) with respect to time variable. In Section 3 we introduce a Faedo–Galerkin approximation of (1.1)–(1.3). We derive primary energy estimates with constants independent of approximation and time, and investigate their implications. In Section 4 we consider a time-differentiated version of the approximate problem and establish the first regularity estimates with constants uniform in approximation but depending on time. The subsequent sections 5 and 6 provide the existence proofs of Theorems 2.1 and 2.2 respectively. The proofs are based on the previously established uniform a priori estimates which, by standard arguments, allow us to pass to the limit in the corresponding versions of the approximate problems.

We remark that having in mind a future examination of the long time behaviour of solutions we record time-dependence of various constants. The obtained regularity estimates turn out to depend exponentially on time, thus in the present form are not useful for the long time analysis. We point out that in long-time analysis the crucial point is to show property that if $\chi(0) \in [-1, 1]$ then $\chi(t) \in [-1, 1]$ for all $t > 0$, in other words that the order parameter attains physically meaningful values for all times. This question is left open in the present paper.

We use the following notation:

$$\begin{aligned} \mathbf{x} &= (x_i)_{i=1,2,3} \in \Omega \subset \mathbb{R}^3 \quad \text{the material point,} \\ f_{,i} &= \frac{\partial f}{\partial x_i}, \quad f_t = \frac{df}{dt} \quad \text{the material space and time derivatives,} \\ \varepsilon &= (\varepsilon_{ij})_{i,j=1,2,3}, \quad W_{,\varepsilon}(\varepsilon, \chi) = \left(\frac{\partial W(\varepsilon, \chi)}{\partial \varepsilon_{ij}} \right)_{i,j=1,2,3}, \\ W_{,\chi}(\varepsilon, \chi) &= \frac{\partial W(\varepsilon, \chi)}{\partial \chi}, \quad \psi'(\chi) = \frac{d\psi(\chi)}{d\chi}. \end{aligned}$$

For simplicity, whenever there is no danger of confusion, we omit the arguments (ε, χ) . The specification of tensor indices is omitted as well.

Vector- and tensor-valued mappings are denoted by bold letters.

The summation convention over repeated indices is used, as well as the notation: for vectors $\mathbf{a} = (a_i)$, $\tilde{\mathbf{a}} = (a_i)$ and tensors $\mathbf{B} = (B_{ij})$, $\tilde{\mathbf{B}} = (\tilde{B}_{ij})$, $\mathbf{A} = (A_{ijkl})$, we write

$$\begin{aligned} \mathbf{a} \cdot \tilde{\mathbf{a}} &= a_i \tilde{a}_i, & \mathbf{B} \cdot \tilde{\mathbf{B}} &= B_{ij} \tilde{B}_{ij}, \\ \mathbf{AB} &= (A_{ijkl} B_{kl}), & \mathbf{BA} &= (B_{ij} A_{ijkl}), \\ |\mathbf{a}| &= (a_i a_i)^{1/2}, & |\mathbf{B}| &= (B_{ij} B_{ij})^{1/2}. \end{aligned}$$

The symbols ∇ and $\nabla \cdot$ denote the gradient and the divergence operators with respect to the material point \mathbf{x} . For the divergence of a tensor field we use the convention of the contraction over the last index, e.g. $\nabla \cdot \varepsilon(\mathbf{x}) = (\varepsilon_{ij,j}(\mathbf{x}))$.

We use the standard Sobolev spaces notation $H^m(\Omega) = W_2^m(\Omega)$ for $m \in \mathbb{N}$. Besides,

$$H_0^1(\Omega) = \{v \in H^1(\Omega) : v = 0 \text{ on } S\}, \quad H_N^2(\Omega) = \{v \in H^2(\Omega) : \mathbf{n} \cdot \nabla v = 0 \text{ on } S\},$$

where \mathbf{n} is the outward unit normal to $S = \partial\Omega$, denote the subspaces respectively of $H^1(\Omega)$ and $H^2(\Omega)$, with the standard norms of $H^1(\Omega)$ and $H^2(\Omega)$.

We denote by bold face letters the spaces of vector- or tensor-valued functions, e.g.

$$\mathbf{L}_2(\Omega) = (L_2(\Omega))^n, \quad \mathbf{H}^1(\Omega) = (H^1(\Omega))^n, \quad n \in \mathbb{N};$$

if there is no confusion we do not specify dimension n .

Moreover, we write

$$\|\mathbf{a}\|_{\mathbf{L}_2(\Omega)} = \|\mathbf{a}\|_{L_2(\Omega)}, \quad \|\mathbf{a}\|_{\mathbf{H}^1(\Omega)} = \|\mathbf{a}\|_{L_2(\Omega)} + \|\nabla \mathbf{a}\|_{L_2(\Omega)}$$

for the corresponding norms of a vector-valued function $\mathbf{a}(\mathbf{x}) = (a_i(\mathbf{x}))$; similarly for tensor-valued functions.

As common, the symbol (\cdot, \cdot) denotes the scalar product in $L_2(\Omega)$. For simplicity, we use the same symbol to denote scalar products in $\mathbf{L}_2(\Omega) = (L_2(\Omega))^n$,

e.g. we write

$$(a, \tilde{a}) = \int_{\Omega} a(\mathbf{x}) \tilde{a}(\mathbf{x}) dx, \quad (\mathbf{a}, \tilde{\mathbf{a}}) = \int_{\Omega} a_i(\mathbf{x}) \tilde{a}_i(\mathbf{x}) dx,$$

$$(\mathbf{B}, \tilde{\mathbf{B}}) = \int_{\Omega} B_{ij}(\mathbf{x}) \tilde{B}_{ij}(\mathbf{x}) dx.$$

The dual of the space V is denoted by V' , and $\langle \cdot, \cdot \rangle_{V', V}$ stands for the duality pairing between V' and V .

By c and $c(T)$ we denote generic positive constants different in various instances, depending on the data of the problem and domain Ω ; whenever it is of interest their dependence on parameters is specified. The argument T indicates the time horizon dependence. Moreover, δ denotes a generic, sufficiently small positive constant.

2. Assumptions and main results

System (1.1)–(1.3) (in simplified form (1.14)–(1.16)) is studied under the following assumptions:

(A1) $\Omega \subset \mathbb{R}^3$ is a bounded domain with the boundary S of class at least C^2 ; $T > 0$ is an arbitrary final time.

(A2) The coefficients of the elasticity operator \mathbf{Q} , defined by (1.11), satisfy

$$(2.1) \quad \bar{\mu} > 0, \quad 3\bar{\lambda} + 2\bar{\mu} > 0 \quad (\text{elasticity range}).$$

These two conditions assure the following:

(i) Coercivity and boundedness of the operator \mathbf{A}

$$(2.2) \quad \underline{c}|\varepsilon|^2 \leq \varepsilon \cdot \mathbf{A}\varepsilon \leq \bar{c}|\varepsilon|^2 \quad \text{for all } \varepsilon \in \mathbf{S}^2,$$

where \mathbf{S}^2 denotes the set of symmetric second order tensors in \mathbb{R}^3 , and

$$\underline{c} = \min\{3\bar{\lambda} + 2\bar{\mu}, 2\bar{\mu}\}, \quad \bar{c} = \max\{3\bar{\lambda} + 2\bar{\mu}, 2\bar{\mu}\};$$

(ii) Strong ellipticity of the operator \mathbf{Q} (property holding true under weaker assumption $\bar{\mu} > 0$, $\bar{\lambda} + 2\bar{\mu} > 0$, see [24, Section 7]). Thanks to this property the following estimate holds true (see [20, Lemma 3.2]):

$$(2.3) \quad c\|\mathbf{u}\|_{\mathbf{H}^2(\Omega)} \leq \|\mathbf{Q}\mathbf{u}\|_{L_2(\Omega)} \quad \text{for } \mathbf{u} \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$$

with constant c depending on Ω .

Hence, since clearly $\|\mathbf{Q}\mathbf{u}\|_{L_2(\Omega)} \leq \bar{c}\|\mathbf{u}\|_{\mathbf{H}^2(\Omega)}$, it follows that the norms $\|\mathbf{Q}\mathbf{u}\|_{L_2(\Omega)}$ and $\|\mathbf{u}\|_{\mathbf{H}^2(\Omega)}$ are equivalent on $\mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$.

The next two assumptions concern the ingredients of the free energy (see (1.10) with $\mathbf{\Gamma} = \mathbf{I}$)

$$(2.4) \quad f(\varepsilon(\mathbf{u}), \chi, \nabla\chi) = W(\varepsilon(\mathbf{u}), \chi) + \psi(\chi) + \frac{1}{2}|\nabla\chi|^2.$$

(A3) The elastic energy $W(\varepsilon(\mathbf{u}), \chi)$ is given by (1.4)–(1.6). The interpolation function $z: \mathbb{R} \rightarrow [0, 1]$ in definition (1.6) of $\bar{\varepsilon}(\chi)$ is at least of class C^1 with the property (1.7). Hence,

$$(2.5) \quad 0 \leq z(\chi) \leq 1 \quad \text{and} \quad |z'(\chi)| \leq c \quad \text{for all } \chi \in \mathbb{R}.$$

(A4) The chemical energy $\psi(\chi)$ has the form of the standard double-well potential (1.8), so

$$(2.6) \quad \psi'(\chi) = \chi^3 - \chi, \quad \psi''(\chi) = 3\chi^2 - 1, \quad \psi'''(\chi) = 6\chi.$$

Moreover, for simplicity it is assumed that

(A5) The mobility tensor \mathbf{M} and the interfacial tensor $\mathbf{\Gamma}$ are the identity matrices $\mathbf{M} = \mathbf{I}$, $\mathbf{\Gamma} = \mathbf{I}$.

The second order symmetric tensor \mathbf{B} and the scalars D , E are defined in (1.12).

We note that assumptions (A3) and (A4) imply the following bounds for all $\varepsilon \in \mathbf{S}^2$ and $\chi \in \mathbb{R}$:

$$(2.7) \quad \begin{aligned} |\bar{\varepsilon}(\chi)| &\leq |\bar{\varepsilon}_a| + |\bar{\varepsilon}_b| \leq c, \\ |\bar{\varepsilon}'(\chi)| &= |z'(\chi)(\bar{\varepsilon}_b - \bar{\varepsilon}_a)| \leq c, \\ |W(\varepsilon, \chi)| &\leq \frac{1}{2}c|\varepsilon - \bar{\varepsilon}(\chi)|^2 \leq c(|\varepsilon|^2 + 1), \\ |W_{,\varepsilon}(\varepsilon, \chi)| + |W_{,\chi}(\varepsilon, \chi)| &\leq c(|\varepsilon| + 1), \\ |\psi(\chi)| &\leq c(\chi^4 + 1), \quad |\psi'(\chi)| \leq c(|\chi|^3 + 1) \end{aligned}$$

with some positive constant c . Moreover, by the Young inequality, we have

$$(2.8) \quad W(\varepsilon, \chi) \geq \frac{1}{2}c|\varepsilon - \bar{\varepsilon}(\chi)|^2 \geq \frac{1}{4}c|\varepsilon|^2 - \frac{1}{2}c|\bar{\varepsilon}(\chi)|^2 \geq \frac{1}{4}c|\varepsilon|^2 - c(|\bar{\varepsilon}_a|^2 + |\bar{\varepsilon}_b|^2),$$

and

$$\psi(\chi) \geq \frac{1}{8}\chi^4 - \frac{1}{4}.$$

This shows that free energy (2.4) satisfies the following structure condition

$$(2.9) \quad \begin{aligned} f(\varepsilon, \chi, \nabla\chi) &\geq \frac{1}{4}c|\varepsilon|^2 + \frac{1}{8}\chi^4 + \frac{1}{2}|\nabla\chi|^2 - c(|\bar{\varepsilon}_a|^2 + |\bar{\varepsilon}_b|^2) - \frac{1}{4} \\ &\geq c_f(|\varepsilon|^2 + \chi^4 + |\nabla\chi|^2) - c'_f \end{aligned}$$

with constants $c_f > 0$ and c'_f given by

$$c_f = \min \left\{ \frac{1}{4}c, \frac{1}{8} \right\}, \quad c'_f = c(|\bar{\varepsilon}_a|^2 + |\bar{\varepsilon}_b|^2) - \frac{1}{4}.$$

This bound plays the key role in the derivation of energy estimates for problem (1.1)–(1.3) (see Section 4).

For further purposes we recall here the following two additional properties of the operator \mathbf{Q} :

(2.10) \mathbf{Q} is selfadjoint on $\mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$, i.e.

$$(\mathbf{Q}\mathbf{u}, \mathbf{v}) = -\bar{\mu}(\nabla\mathbf{u}, \nabla\mathbf{v}) - (\bar{\lambda} + \bar{\mu})(\nabla \cdot \mathbf{u}, \nabla \cdot \mathbf{v}) = (\mathbf{u}, \mathbf{Q}\mathbf{v})$$

for $\mathbf{u}, \mathbf{v} \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$,

(2.11) $-\mathbf{Q}$ is positive on $\mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$, i.e.

$$(-\mathbf{Q}\mathbf{u}, \mathbf{u}) = \bar{\mu}\|\nabla\mathbf{u}\|_{L_2(\Omega)}^2 + (\bar{\lambda} + \bar{\mu})\|\nabla \cdot \mathbf{u}\|_{L_2(\Omega)}^2 \geq 0$$

for $\mathbf{u} \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$.

We state now the main results of the paper.

THEOREM 2.1 (Weak solutions). *Let assumptions (A1)–(A5) hold true. Moreover, let the data satisfy*

$$(2.12) \quad \mathbf{b} \in L_2(\Omega^T), \quad \mathbf{u}_0 \in \mathbf{H}_0^1(\Omega), \quad \mathbf{u}_1 \in L_2(\Omega), \quad \chi_0 \in H^1(\Omega).$$

Then there exist functions (\mathbf{u}, χ, μ) such that

$$(2.13) \quad \begin{aligned} \mathbf{u} &\in L_\infty(0, T; \mathbf{H}_0^1(\Omega)), \quad \mathbf{u}_t \in L_\infty(0, T; L_2(\Omega)), \\ \mathbf{u}_{tt} &\in L_2(0, T; (\mathbf{H}_0^1(\Omega))'), \\ \chi &\in L_\infty(0, T; H^1(\Omega)) \cap L_2(0, T; H_N^2(\Omega)), \quad \chi_t \in L_2(0, T; (H^1(\Omega))'), \\ \mu &\in L_2(0, T; H^1(\Omega)), \\ \mathbf{u}(0) &= \mathbf{u}_0, \quad \mathbf{u}_t(0) = \mathbf{u}_1, \quad \chi(0) = \chi_0, \end{aligned}$$

which satisfy problem (1.14)–(1.16) in the following weak sense

$$\int_0^T \langle \mathbf{u}_{tt}, \boldsymbol{\eta} \rangle_{(\mathbf{H}_0^1(\Omega))', \mathbf{H}_0^1(\Omega)} dt + \int_0^T (\mathbf{A}\varepsilon(\mathbf{u}), \varepsilon(\boldsymbol{\eta})) dt = \int_0^T (z'(\chi)\mathbf{B}\nabla\chi + \mathbf{b}, \boldsymbol{\eta}) dt$$

for all $\boldsymbol{\eta} \in L_2(0, T; \mathbf{H}_0^1(\Omega))$,

$$(2.14) \quad \int_0^T \langle \chi_t, \xi \rangle_{(H^1(\Omega))', H^1(\Omega)} dt + \int_0^T (\nabla\mu, \nabla\xi) dt = 0$$

for all $\xi \in L_2(0, T; H^1(\Omega))$,

$$\int_0^T (\mu, \zeta) dt = - \int_0^T (\Delta\chi, \zeta) dt + \int_0^T (\psi'(\chi) + W_{,\chi}(\varepsilon(\mathbf{u}), \chi), \zeta) dt$$

for all $\zeta \in L_2(0, T; L_2(\Omega))$.

Moreover, (\mathbf{u}, χ, μ) satisfy a priori estimates:

$$(2.15) \quad \begin{aligned} &\|\mathbf{u}_t\|_{L_\infty(0, T; L_2(\Omega))} + \|\varepsilon(\mathbf{u})\|_{L_\infty(0, T; L_2(\Omega))} + \|\chi\|_{L_\infty(0, T; L_4(\Omega))} \\ &+ \|\nabla\chi\|_{L_\infty(0, T; L_2(\Omega))} + \|\nabla\mu\|_{L_2(\Omega^T)} + \|\chi_t\|_{L_2(0, T; (H^1(\Omega))')} \leq c_0, \end{aligned}$$

$$(2.16) \quad \|\mathbf{u}\|_{L_\infty(0,T;\mathbf{H}_0^1(\Omega))} + \|\chi\|_{L_\infty(0,T;H^1(\Omega))} \leq c_1,$$

$$(2.17) \quad \|\chi\|_{L_2(0,T;H_N^2(\Omega))} + \|\mu\|_{L_2(0,T;H^1(\Omega))} \leq c_2(T),$$

$$(2.18) \quad \|\mathbf{u}_{tt}\|_{L_2(0,T;(\mathbf{H}_0^1(\Omega))')} \leq c_3(T),$$

with positive constants

$$\begin{aligned} c_0 &= c(\|\mathbf{u}_0\|_{\mathbf{H}^1(\Omega)}, \|\mathbf{u}_1\|_{L_2(\Omega)}, \|\chi_0\|_{H^1(\Omega)}, \|\mathbf{b}\|_{L_1(0,T;L_2(\Omega))}, c_f, c'_f), \\ c_1 &= c(c_0, \Omega), \quad c_2(T) = c(c_1)T^{1/2}, \quad c_3(T) = c(c_0, \|\mathbf{b}\|_{L_2(\Omega^T)})T^{1/2}. \end{aligned}$$

The second theorem states a time-regularity result which is concluded from a time-differentiated version of problem (1.14)–(1.16).

In compatibility with equations (1.14)₁, (1.15)₁ and (1.16)₁, we define the following initial conditions corresponding respectively to $\mathbf{u}_{tt}(0)$, $\chi_t(0)$ and $\mu(0)$:

$$(2.19) \quad \begin{aligned} \mathbf{u}_2 &:= \mathbf{Q}\mathbf{u}_0 + z'(\chi_0)B\nabla\chi_0 + \mathbf{b}(0), \\ \chi_1 &:= \Delta\mu_0, \\ \mu_0 &:= -\Delta\chi_0 + \psi'(\chi_0) + z'(\chi_0)(\mathbf{B} \cdot \varepsilon(\mathbf{u}_0) + Dz(\chi_0) + E). \end{aligned}$$

THEOREM 2.2 (Time regularity). *Let (A1)–(A5) hold, the boundary S of domain Ω be of class C^4 , and*

$$(2.20) \quad z: \mathbb{R} \rightarrow [0, 1] \text{ be of class } C^2 \text{ with } |z'(\chi)| + |z''(\chi)| \leq c \text{ for all } \chi \in \mathbb{R}.$$

Moreover, let the data satisfy

$$(2.21) \quad \begin{aligned} &\mathbf{b} \in H^1(0, T; L_2(\Omega)), \\ \mathbf{u}_0 &\in \mathbf{H}^3(\Omega) \cap \mathbf{H}_0^1(\Omega), \quad \mathbf{u}_1 \in \mathbf{H}_0^1(\Omega), \quad \chi_0 \in H^4(\Omega) \cap H_N^2(\Omega), \\ &\mathbf{u}_2 \in L_2(\Omega), \quad \chi_1 \in L_2(\Omega), \quad \mu_0 \in H_N^2(\Omega). \end{aligned}$$

Then there exist functions (\mathbf{u}, χ, μ) such that

$$(2.22) \quad \begin{aligned} \mathbf{u} &\in L_\infty(0, T; \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)), \quad \mathbf{u}_t \in L_\infty(0, T; \mathbf{H}_0^1(\Omega)), \\ \mathbf{u}_{tt} &\in L_\infty(0, T; L_2(\Omega)), \quad \mathbf{u}_{ttt} \in L_2(0, T; (\mathbf{H}_0^1(\Omega))'), \\ \chi &\in C^{1/2}([0, T]; H_N^2(\Omega)), \quad \chi_t \in L_\infty(0, T; L_2(\Omega)) \cap L_2(0, T; H_N^2(\Omega)), \\ \chi_{tt} &\in L_2(0, T; (H_N^2(\Omega))'), \\ \mu &\in L_\infty(0, T; H_N^2(\Omega)), \quad \mu_t \in L_2(\Omega^T), \\ \mathbf{u}(0) &= \mathbf{u}_0, \quad \mathbf{u}_t(0) = \mathbf{u}_1, \quad \mathbf{u}_{tt}(0) = \mathbf{u}_2, \\ \chi(0) &= \chi_0, \quad \chi_t(0) = \chi_1, \quad \mu(0) = \mu_0, \end{aligned}$$

which satisfy problem (1.14)–(1.16) in the sense of the identities:

$$(2.23) \quad \begin{aligned} &\int_0^T \langle \mathbf{u}_{ttt}, \boldsymbol{\eta} \rangle_{(\mathbf{H}_0^1(\Omega))', \mathbf{H}_0^1(\Omega)} dt + \int_0^T (\mathbf{A}\varepsilon(\mathbf{u}_t), \varepsilon(\boldsymbol{\eta})) dt \\ &= \int_0^T ([z'(\chi)B\nabla\chi]_t + \mathbf{b}_t, \boldsymbol{\eta}) dt \quad \text{for all } \boldsymbol{\eta} \in H^1(0, T; \mathbf{H}_0^1(\Omega)), \quad \boldsymbol{\eta}(T) = 0, \end{aligned}$$

$$(2.24) \quad \int_0^T \langle \chi_{tt}, \xi \rangle_{(H_N^2(\Omega))', H_N^2(\Omega)} dt = \int_0^T (\mu_t, \Delta \xi) dt$$

for all $\xi \in H^1(0, T; H_N^2(\Omega))$, $\xi(T) = 0$,

$$(2.25) \quad \int_0^T (\mu_t, \zeta) dt = - \int_0^T (\Delta \chi_t, \zeta) dt + \int_0^T ([\psi'(\chi) + W_{,\chi}(\varepsilon(\mathbf{u}), \chi)]_{,t}, \zeta) dt$$

for all $\zeta \in H^1(0, T; L_2(\Omega))$, $\zeta(T) = 0$,

where

$$(2.26) \quad [\psi'(\chi) \mathbf{B} \nabla \chi]_{,t} = z''(\chi) \chi_t \mathbf{B} \nabla \chi + z'(\chi) \mathbf{B} \nabla \chi_t,$$

$$[\psi'(\chi) + W_{,\chi}(\varepsilon(\mathbf{u}), \chi)]_{,t} = \psi''(\chi) \chi_t + z''(\chi) \chi_t (\mathbf{B} \cdot \varepsilon(\mathbf{u}) + D z(\chi) + E)$$

$$+ z'(\chi) (\mathbf{B} \cdot \varepsilon(\mathbf{u}_t) + D z'(\chi) \chi_t).$$

In addition, (\mathbf{u}, χ, μ) satisfy estimates (2.15)–(2.18) and

$$(2.27) \quad \|\mathbf{u}\|_{L_\infty(0, T; \mathbf{H}^2(\Omega))} + \|\mathbf{u}_t\|_{L_\infty(0, T; \mathbf{H}^1(\Omega))} + \|\mathbf{u}_{tt}\|_{L_\infty(0, T; L_2(\Omega))} \leq c_5(T),$$

$$\|\chi\|_{C^{1/2}([0, T]; H_N^2(\Omega))} + \|\chi_t\|_{L_\infty(0, T; L_2(\Omega))}$$

$$+ \|\chi_t\|_{L_2(0, T; H_N^2(\Omega))} + \|\mu\|_{L_\infty(0, T; H_N^2(\Omega))} \leq c_4(T),$$

$$\|\mathbf{u}_{ttt}\|_{L_2(0, T; (\mathbf{H}_0^1(\Omega))')} + \|\chi_{tt}\|_{L_2(0, T; (H_N^2(\Omega))')} + \|\mu_t\|_{L_2(\Omega^T)} \leq T^{1/2} c_5(T),$$

with constants

$$c_4(T) = c(T^{1/2} E_1(T) + \|\chi_1\|_{L_2(\Omega)}) [\exp a(T)]^{1/2}, \quad c_5(T) = T^{1/2} c_4(T),$$

where

$$E_1(T) = T^{1/2} \|\mathbf{b}_t\|_{L_2(\Omega^T)} + \|\mathbf{u}_2\|_{L_2(\Omega)} + \|\varepsilon(\mathbf{u}_1)\|_{L_2(\Omega)}, \quad a(T) = c(c_0) T^8 \exp(cT).$$

REMARK 2.3. Integrating by parts with respect to time in identities (2.23)–(2.25) and using compatibility conditions (2.19) we obtain the standard weak formulation of (1.14)–(1.16).

3. The Faedo–Galerkin approximation

In this section we introduce a Faedo–Galerkin approximation of problem (1.1)–(1.3) (in simplified form (1.14)–(1.16)) and derive basic energy estimates. These estimates are used to prove the existence of weak solutions in Theorem 2.1. Throughout this section we assume that the domain Ω has the boundary S at least of class C^2 .

3.1. Approximation. Let us consider the following two eigenvalue problems

$$(3.1) \quad -\mathbf{Q} \mathbf{v}_j = \lambda_j \mathbf{v}_j \quad \text{in } \Omega, \quad \mathbf{v}_j = \mathbf{0} \quad \text{on } S, \quad j \in \mathbb{N},$$

where \mathbf{Q} is the elliptic operator defined by (1.11), and

$$(3.2) \quad -\Delta w_j = \lambda'_j w_j \quad \text{in } \Omega, \quad \mathbf{n} \cdot \nabla w_j = 0 \quad \text{on } S, \quad j \in \mathbb{N}.$$

We recall that, by virtue of the elliptic regularity theory, if the domain Ω has the boundary of class C^l , $l \in \mathbb{N}$, then the solutions of (3.1) and (3.2) satisfy

$$\mathbf{v}_j \in \mathbf{H}^l(\Omega), \quad w_j \in H^l(\Omega).$$

We take the family $\{\mathbf{v}_j\}_{j \in \mathbb{N}}$ as a basis of the space $\mathbf{H}_0^1(\Omega)$ and the family $\{w_j\}_{j \in \mathbb{N}}$ as a basis of the space $H_N^2(\Omega) = \{w \in H^2(\Omega) : \mathbf{n} \cdot \nabla w = 0 \text{ on } S\}$. Such choice is possible thanks to the following properties of (3.1) and (3.2). On account of (2.10) we have

$$(3.3) \quad \begin{aligned} \lambda_i(\mathbf{v}_i, \mathbf{v}_j) &= (-\mathbf{Q}\mathbf{v}_i, \mathbf{v}_j) = \bar{\mu}(\nabla \mathbf{v}_i, \nabla \mathbf{v}_j) + (\bar{\lambda} + \bar{\mu})(\nabla \cdot \mathbf{v}_i, \nabla \cdot \mathbf{v}_j) \\ &= (\mathbf{v}_i, -\mathbf{Q}\mathbf{v}_j) = \lambda_j(\mathbf{v}_i, \mathbf{v}_j) \end{aligned}$$

for $i, j \in \mathbb{N}$. Identities (3.3) show, by the Poincaré–Friedrichs inequality, that the family $\{\mathbf{v}_j\}_{j \in \mathbb{N}}$ is orthogonal in $\mathbf{H}^1(\Omega)$ and $L_2(\Omega)$ scalar products.

We shall assume that \mathbf{v}_j are normalized so that $(\mathbf{v}_i, \mathbf{v}_i) = 1$. Thereby the basis $\{\mathbf{v}_j\}_{j \in \mathbb{N}}$ becomes orthonormal in $L_2(\Omega)$ and orthogonal in $\mathbf{H}^1(\Omega)$ scalar products.

Similarly, the family $\{w_j\}_{j \in \mathbb{N}}$ satisfies

$$(3.4) \quad \begin{aligned} \lambda_i(w_i, w_j) &= (-\Delta w_i, w_j) = (\nabla w_i, \nabla w_j) = (w_i, -\Delta w_j) = \lambda_j(w_i, w_j), \\ \lambda_i \lambda_j(w_i, w_j) &= (\Delta w_i, \Delta w_j) \quad \text{for } i, j \in \mathbb{N}. \end{aligned}$$

Hence, by the Poincaré inequality, it follows that the family $\{w_j\}_{j \in \mathbb{N}}$ is orthogonal in $H^2(\Omega)$, $H^1(\Omega)$ and $L_2(\Omega)$ scalar products. We normalize w_j so that $(w_i, w_i) = 1$. Then the basis $\{w_j\}_{j \in \mathbb{N}}$ becomes orthonormal in $L_2(\Omega)$ and orthogonal in $H^1(\Omega)$ and $H^2(\Omega)$ scalar products. Furthermore, we assume without loss of generality that $w_1 = 1$.

For $m \in \mathbb{N}$ we denote by $\mathbf{V}_{0m} = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$, $V_m = \text{span}\{w_1, \dots, w_m\}$ the finite dimensional subspaces, respectively of $\mathbf{H}_0^1(\Omega)$ and $H_N^2(\Omega)$, spanned by $\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$ and $\{w_1, \dots, w_m\}$.

Now, let us introduce the following approximation of problem (1.14)–(1.16): For any $m \in \mathbb{N}$ find a triple of functions $(\mathbf{u}^m, \chi^m, \mu^m)$ of the form

$$(3.5) \quad \begin{aligned} \mathbf{u}^m(\mathbf{x}, t) &= \sum_{i=1}^m e_i^m(t) \mathbf{v}_i(\mathbf{x}), \\ \chi^m(\mathbf{x}, t) &= \sum_{i=1}^m c_i^m(t) w_i(\mathbf{x}), \\ \mu^m(\mathbf{x}, t) &= \sum_{i=1}^m d_i^m(t) w_i(\mathbf{x}), \end{aligned}$$

with $e_i^m(t)$, $c_i^m(t)$, $d_i^m(t)$ being determined so that

$$\begin{aligned}
 (\mathbf{u}_{tt}^m, \mathbf{v}_j) + (\mathbf{A}\varepsilon(\mathbf{u}^m), \varepsilon(\mathbf{v}_j)) &= (z'(\chi^m)\mathbf{B}\nabla\chi^m + \mathbf{b}, \mathbf{v}_j), \\
 (\chi_t^m, w_j) + (\nabla\mu^m, \nabla w_j) &= 0, \\
 (\mu^m, w_j) &= -(\Delta\chi^m, w_j) + (\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m), w_j), \\
 &\text{for } j = 1, \dots, m, \\
 \mathbf{u}^m(0) &= \mathbf{u}_0^m, \quad \mathbf{u}_t^m(0) = \mathbf{u}_1^m, \quad \chi^m(0) = \chi_0^m,
 \end{aligned}
 \tag{3.6}$$

where $\mathbf{u}_0^m, \mathbf{u}_1^m \in \mathbf{V}_{0m}$ and $\chi_0^m \in V_m$ are the projections of $\mathbf{u}_0, \mathbf{u}_1$ respectively, and χ_0 satisfying for $m \rightarrow \infty$

$$\begin{aligned}
 \mathbf{u}_0^m &\rightarrow \mathbf{u}_0 \quad \text{strongly in } \mathbf{H}_0^1(\Omega), \\
 \mathbf{u}_1^m &\rightarrow \mathbf{u}_1 \quad \text{strongly in } \mathbf{L}_2(\Omega), \\
 \chi_0^m &\rightarrow \chi_0 \quad \text{strongly in } H^1(\Omega).
 \end{aligned}
 \tag{3.7}$$

Clearly, (3.6) can be expressed as a system of first order ordinary differential equations for the coefficients (e_1^m, \dots, e_m^m) , $(e_{1,t}^m, \dots, e_{m,t}^m)$, (c_1^m, \dots, c_m^m) , with the right-hand sides being by assumptions continuous functions of their arguments. Thus (3.6) has a local in time solution on an interval $[0, T_m]$, $T_m > 0$. The uniform in m a priori estimates proved in lemmas below show that $T_m = T$, i.e. (3.6) has a solution on the interval $[0, T]$.

3.2. Energy estimates.

LEMMA 3.1. *Let (A1)–(A5) hold and the data satisfy*

$$\mathbf{u}_0 \in \mathbf{H}_0^1(\Omega), \quad \mathbf{u}_1 \in \mathbf{L}_2(\Omega), \quad \chi_0 \in H^1(\Omega), \quad \mathbf{b} \in L_1(0, T; \mathbf{L}_2(\Omega)).
 \tag{3.8}$$

Then a solution $(\mathbf{u}^m, \chi^m, \mu^m)$ to problem (3.6) satisfies the following uniform estimate

$$\begin{aligned}
 \|\mathbf{u}_t^m\|_{L_\infty(0, T; \mathbf{L}_2(\Omega))} + \|\varepsilon(\mathbf{u}^m)\|_{L_\infty(0, T; \mathbf{L}_2(\Omega))} + \|\chi^m\|_{L_\infty(0, T; L_4(\Omega))} \\
 + \|\nabla\chi^m\|_{L_\infty(0, T; \mathbf{L}_2(\Omega))} + \|\nabla\mu^m\|_{L_2(\Omega^T)} \leq c_0,
 \end{aligned}
 \tag{3.9}$$

with the constant

$$c_0 = c_0(\|\mathbf{u}_0\|_{\mathbf{H}^1(\Omega)}, \|\mathbf{u}_1\|_{\mathbf{L}_2(\Omega)}, \|\chi_0\|_{H^1(\Omega)}, \|\mathbf{b}\|_{L_1(0, T; \mathbf{L}_2(\Omega))}, c_f, c'_f).$$

PROOF. We derive the energy identity for system (3.6). First, let us note that, according to (1.13),

$$z'(\chi)\mathbf{B}\nabla\chi = -\nabla \cdot \mathbf{A}(\bar{\varepsilon}_a + z(\chi)(\bar{\varepsilon}_b - \bar{\varepsilon}_a)),$$

thus an equivalent form of (3.6)₁ is

$$(\mathbf{u}_{tt}^m, \mathbf{v}_j) + (W_{,\varepsilon}(\varepsilon(\mathbf{u}^m), \chi^m), \varepsilon(\mathbf{v}_j)) = (\mathbf{b}, \mathbf{v}_j).
 \tag{3.10}$$

Testing (3.10) by $\mathbf{u}_t^m(t)$ (i.e. multiplying by $e_j^m(t)$ and summing over j from $j = 1$ to $j = m$) gives

$$(3.11) \quad \frac{1}{2} \frac{d}{dt} \|\mathbf{u}_t^m\|_{\mathbf{L}_2(\Omega)}^2 + (W_{,\varepsilon}(\varepsilon(\mathbf{u}^m), \chi^m), \varepsilon(\mathbf{u}_t^m)) = (\mathbf{b}, \mathbf{u}_t^m).$$

Further, testing (3.6)₂ by μ^m yields

$$(3.12) \quad (\chi_t^m, \mu^m) + \|\nabla \mu^m\|_{\mathbf{L}_2(\Omega)}^2 = 0.$$

Finally, testing (3.6)₃ by $-\chi_t^m(t)$ and integrating by parts, leads to

$$(3.13) \quad -(\mu^m, \chi_t^m) + \frac{1}{2} \frac{d}{dt} \|\nabla \chi^m\|_{\mathbf{L}_2(\Omega)}^2 + (\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m), \chi_t^m) = 0.$$

Summing up (3.11)–(3.13) we arrive at the following energy identity

$$(3.14) \quad \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\mathbf{u}_t^m|^2 dx + \frac{d}{dt} \int_{\Omega} \left[W(\varepsilon(\mathbf{u}^m), \chi^m) + \psi(\chi^m) + \frac{1}{2} |\nabla \chi^m|^2 \right] dx \\ + \int_{\Omega} |\nabla \mu^m|^2 dx = \int_{\Omega} \mathbf{b} \cdot \mathbf{u}_t^m dx.$$

Integration of (3.14) over $(0, t)$ gives

$$(3.15) \quad \frac{1}{2} \int_{\Omega} |\mathbf{u}_t^m|^2 dx + \int_{\Omega} f(\varepsilon(\mathbf{u}^m), \chi^m, \nabla \chi^m) dx + \int_{\Omega^t} |\nabla \mu^m|^2 dx dt' \\ = \frac{1}{2} \int_{\Omega} |\mathbf{u}_1^m|^2 dx + \int_{\Omega} f(\varepsilon(\mathbf{u}_0^m), \chi_0^m, \nabla \chi_0^m) dx + \int_{\Omega^t} \mathbf{b} \cdot \mathbf{u}_{t'}^m dx dt',$$

with $f(\varepsilon, \chi, \nabla \chi)$ defined by (2.4). Now, bearing in mind that $f(\varepsilon, \chi, \nabla \chi)$ satisfies the structure condition (2.9), we can estimate the left-hand side of (3.15) from below by

$$\frac{1}{2} \|\mathbf{u}_t^m\|_{\mathbf{L}_2(\Omega)}^2 + c_f (\|\varepsilon(\mathbf{u}^m)\|_{\mathbf{L}_2(\Omega)}^2 + \|\chi^m\|_{L_4(\Omega)}^4 + \|\nabla \chi^m\|_{\mathbf{L}_2(\Omega)}^2) + \|\nabla \mu^m\|_{\mathbf{L}_2(\Omega^t)}^2 - c'_f.$$

Further, in view of growth conditions (2.7) and the convergences (3.7), we have

$$\int_{\Omega} f(\varepsilon(\mathbf{u}_0^m), \chi_0^m, \nabla \chi_0^m) dx \leq c (\|\varepsilon(\mathbf{u}_0)\|_{\mathbf{L}_2(\Omega)}^2 + \|\chi_0\|_{L_4(\Omega)}^4 + \|\nabla \chi_0\|_{\mathbf{L}_2(\Omega)}^2 + 1).$$

Thus the sum of the first two terms on the right-hand side of (3.15) is bounded from above by a constant depending on $\|\chi_0\|_{H^1(\Omega)}$, $\|\mathbf{u}_0\|_{\mathbf{H}^1(\Omega)}$ and $\|\mathbf{u}_1\|_{\mathbf{L}_2(\Omega)}$.

Finally, estimating the third term on the right-hand side of (3.15) by

$$\left| \int_{\Omega^t} \mathbf{b} \cdot \mathbf{u}_{t'}^m dx dt' \right| \leq \|\mathbf{u}_{t'}^m\|_{L_{\infty}(0,t;\mathbf{L}_2(\Omega))} \|\mathbf{b}\|_{L_1(0,t;\mathbf{L}_2(\Omega))} \\ \leq \frac{1}{4} \|\mathbf{u}_{t'}^m\|_{L_{\infty}(0,t;\mathbf{L}_2(\Omega))}^2 + \|\mathbf{b}\|_{L_1(0,t;\mathbf{L}_2(\Omega))}^2,$$

we arrive at the following uniform in m estimate

$$(3.16) \quad \frac{1}{4} \|\mathbf{u}_t^m\|_{\mathbf{L}_2(\Omega)}^2 + c_f (\|\varepsilon(\mathbf{u}^m)\|_{\mathbf{L}_2(\Omega)}^2 + \|\chi^m\|_{L_4(\Omega)}^4 + \|\nabla \chi^m\|_{\mathbf{L}_2(\Omega)}^2) \\ + \|\nabla \mu^m\|_{\mathbf{L}_2(\Omega^t)}^2 \leq c$$

for $t \in (0, T]$, with constant c depending only on $\|\mathbf{u}_0\|_{\mathbf{H}^1(\Omega)}$, $\|\mathbf{u}_1\|_{\mathbf{L}_2(\Omega)}$, $\|\chi_0\|_{H^1(\Omega)}$, $\|\mathbf{b}\|_{L_1(0,t;L_2(\Omega))}$ and c'_f . This proves the assertion. \square

3.3. Further estimates. Clearly, (3.9) implies that

$$(3.17) \quad \|\chi^m\|_{L_\infty(0,T;H^1(\Omega))} \leq c_1$$

with constant $c_1 = c(c_0, \Omega)$. Hence, by the Sobolev imbedding,

$$(3.18) \quad \|\chi^m\|_{L_\infty(0,T;L_6(\Omega))} \leq c_1.$$

Further, since $\mathbf{u}^m = \mathbf{0}$ on S^T , it follows from (3.9) by Korn's inequality that

$$(3.19) \quad \|\mathbf{u}^m\|_{L_\infty(0,T;\mathbf{H}^1(\Omega))} \leq c_1.$$

We note that setting $w_j = 1$ in (3.6)₂ (admissible by assumption) yields

$$(3.20) \quad \frac{d}{dt} \int_{\Omega} \chi^m dx = 0,$$

which shows that the mean value of χ^m is preserved

$$\int_{\Omega} \chi^m(t) dx = \int_{\Omega} \chi_0^m dx \quad \text{for } t \in [0, T].$$

This property will be used in later analysis. We remark also that thanks to (3.20) the structure condition (2.9) on $f(\varepsilon, \chi, \nabla \chi)$ could be in fact replaced by a weaker one

$$f(\varepsilon, \chi, \nabla \chi) \geq c_f(|\varepsilon|^2 + |\nabla \chi|^2) - c'_f.$$

In such a case, by the Poincaré inequality, estimate $\|\nabla \chi\|_{L_\infty(0,T;L_2(\Omega))} \leq c_0$ would still guarantee bounds (3.17) and (3.18) with a constant

$$c = c\left(c_0, \Omega, \int_{\Omega} \chi_0 dx\right).$$

On the basis of (3.9) and (3.18) we derive an additional estimate on μ^m .

LEMMA 3.2. *Let assumptions of Lemma 3.1 be satisfied. Then, for $t \in (0, T]$,*

$$(3.21) \quad \|\mu^m\|_{L_2(0,t;H^1(\Omega))} \leq c_2(t)$$

with constant $c_2(t) = c(c_1)t^{1/2}$.

PROOF. Setting $w_j = 1$ in (3.6)₃ (admissible by assumption) it follows that

$$\int_{\Omega} \mu^m dx = \int_{\Omega} [\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)] dx.$$

Hence, using growth conditions (2.7) and estimates (3.9), (3.18) we obtain

$$(3.22) \quad \left| \int_{\Omega} \mu^m dx \right| \leq c \int_{\Omega} (|\chi|^3 + |\varepsilon(\mathbf{u})| + 1) dx \leq c(c_1)$$

for almost all $t \in (0, T]$. Consequently, by the Poincaré inequality, estimates (3.9) and (3.22) imply that

$$\begin{aligned} \|\mu^m\|_{L_2(\Omega^t)} &\leq c\|\nabla\mu^m\|_{L_2(\Omega^t)} + c\left[\int_0^t \left(\int_{\Omega} \mu^m dx\right)^2 dt'\right]^{1/2} \\ &\leq cc_0 + c(c_1)t^{1/2} \leq c_2(t). \end{aligned}$$

This shows (3.21). \square

By virtue of Lemma 3.2 we can deduce further estimates on χ^m .

LEMMA 3.3. *Let assumptions of Lemma 3.1 hold. Then, for $t \in (0, T]$,*

$$(3.23) \quad \|\chi^m\|_{L_2(0,t;H^2_{\chi}(\Omega))} \leq c_2(t).$$

PROOF. In view of (3.2), identity (3.6)₃ implies that

$$(\mu^m, \Delta w_j) = (-\Delta\chi^m, \Delta w_j) + (\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m), \Delta w_j).$$

Testing the above equality by $\chi^m(t)$ and integrating with respect to t yields

$$\begin{aligned} \int_0^t \int_{\Omega} (\Delta\chi^m)^2 dx dt' &= - \int_0^t \int_{\Omega} \mu^m \Delta\chi^m dx dt' \\ &\quad + \int_0^t \int_{\Omega} [\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)] \Delta\chi^m dx dt'. \end{aligned}$$

Now, using the Cauchy–Schwarz inequality, and then growth conditions (2.7) and estimates (3.9), (3.18), (3.21), we obtain

$$\begin{aligned} (3.24) \quad \|\Delta\chi^m\|_{L_2(\Omega^t)} &\leq \|\mu^m\|_{L_2(\Omega^t)} + \|\psi'(\chi^m)\|_{L_2(\Omega^t)} \\ &\quad + \|W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)\|_{L_2(\Omega^t)} \\ &\leq \|\mu^m\|_{L_2(\Omega^t)} + ct^{1/2}(\|\chi^m\|_{L_{\infty}(0,t;L_6(\Omega))}^3 \\ &\quad + \|\varepsilon(\mathbf{u}^m)\|_{L_{\infty}(0,t;L_2(\Omega))} + 1) \leq c_2(t). \end{aligned}$$

On account of the ellipticity property of the Laplace operator (see e.g. [11, Chapter III.8]) we have

$$(3.25) \quad \|\chi^m\|_{H^2(\Omega)} \leq c\left(\|\Delta\chi^m\|_{L_2(\Omega)} + \left|\int_{\Omega} \chi^m dx\right|\right).$$

Hence, by (3.24), (3.20) and the convergences (3.7)₃, we conclude (3.23). \square

Using standard duality arguments we shall estimate also time derivatives \mathbf{u}_{tt}^m and χ_t^m .

LEMMA 3.4. *Let assumptions of Lemma 3.1 hold, and $\mathbf{b} \in \mathbf{L}_2(\Omega^T)$. Then, for $t \in (0, T]$,*

$$(3.26) \quad \|\mathbf{u}_{t't'}^m\|_{L_2(0,t;(\mathbf{H}_0^1(\Omega))')} \leq c_3(t), \quad \|\chi_{t'}^m\|_{L_2(0,t;(H^1(\Omega))')} \leq c_0,$$

where $c_3(t) = c(c_0, \|\mathbf{b}\|_{L_2(\Omega^t)})t^{1/2}$.

PROOF. For $\boldsymbol{\eta} \in L_2(0, T; \mathbf{H}_0^1(\Omega))$, we test (3.6)₁ by $\boldsymbol{\eta}^m = \mathbf{P}^m \boldsymbol{\eta}$, where \mathbf{P}^m denotes the projection defined by

$$(3.27) \quad \mathbf{P}^m \boldsymbol{\eta} = \sum_{i=1}^m (\boldsymbol{\eta}, \mathbf{v}_i) \mathbf{v}_i.$$

Then, using the Cauchy–Schwarz inequality, and recalling estimate (3.9), we obtain

$$\begin{aligned} \left| \int_0^t (\mathbf{u}_{t't'}^m, \boldsymbol{\eta}) dt' \right| &= \left| \int_0^t (\mathbf{u}_{t't'}^m, \mathbf{P}^m \boldsymbol{\eta}) dt' \right| \\ &= \left| \int_0^t [-(\mathbf{A}\varepsilon(\mathbf{u}^m), \varepsilon(\mathbf{P}^m \boldsymbol{\eta})) + (z'(\chi^m) \mathbf{B} \nabla \chi^m + \mathbf{b}, \mathbf{P}^m \boldsymbol{\eta})] dt' \right| \\ &\leq c[\|\varepsilon(\mathbf{u}^m)\|_{L_2(\Omega^t)} \|\nabla \mathbf{P}^m \boldsymbol{\eta}\|_{L_2(\Omega^t)} + (\|\nabla \chi^m\|_{L_2(\Omega^t)} + \|\mathbf{b}\|_{L_2(\Omega^t)}) \|\mathbf{P}^m \boldsymbol{\eta}\|_{L_2(\Omega^t)}] \\ &\leq c(c_0 t^{1/2} + \|\mathbf{b}\|_{L_2(\Omega^t)}) \|\mathbf{P}^m \boldsymbol{\eta}\|_{L_2(0,t;H^1(\Omega))} \leq c_3(t) \|\boldsymbol{\eta}\|_{L_2(0,t;H^1(\Omega))} \end{aligned}$$

for all $\boldsymbol{\eta} \in L_2(0, t; \mathbf{H}_0^1(\Omega))$. This shows (3.26)₁. Similarly, for $\xi \in L_2(0, T; H^1(\Omega))$, we test (3.6)₂ by

$$(3.28) \quad \xi^m = P^m \xi = \sum_{i=1}^m (\xi, w_i) w_i,$$

to obtain

$$\begin{aligned} \left| \int_0^t (\chi_{t'}^m, \xi) dt' \right| &= \left| \int_0^t (\chi_{t'}^m, P^m \xi) dt' \right| = \left| \int_0^t (\nabla \mu^m, \nabla P^m \xi) dt' \right| \\ &\leq \|\nabla \mu^m\|_{L_2(\Omega^t)} \|\nabla P^m \xi\|_{L_2(\Omega^t)} \leq c_0 \|\xi\|_{L_2(0,t;H^1(\Omega))}, \end{aligned}$$

where in the last inequality we used (3.9). This implies (3.26)₂ and completes the proof. \square

4. Time regularity estimates

In this section we derive uniform in m time-regularity estimates for solutions of approximate system (3.6). These estimates result from time-differentiated version of (3.6) and lead to the existence result of Theorem 2.2. Throughout this section we assume that the domain Ω has the boundary S of class C^4 .

Let us differentiate system (3.6)₁–(3.6)₃ with respect to t and rewrite it in the following form:

$$\begin{aligned}
 (\mathbf{u}_{ttt}^m, \mathbf{v}_j) + (\mathbf{A}(\varepsilon(\mathbf{u}_t^m), \varepsilon(\mathbf{v}_j))) &= ([z'(\chi^m)\mathbf{B}\nabla\chi^m]_{,t} + \mathbf{b}_t, \mathbf{v}_j), \\
 (\chi_{tt}^m, w_j) - (\mu_t^m, \Delta w_j) &= 0, \\
 (\mu_t^m, w_j) &= -(\Delta\chi_t^m, w_j) + ([\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)]_{,t}, w_j),
 \end{aligned}
 \tag{4.1}$$

for $j = 1, \dots, m$, where the explicit expressions for $[z'(\chi)\mathbf{B}\nabla\chi]_{,t}$ and $[\psi'(\chi) + W_{,\chi}(\varepsilon(\mathbf{u}), \chi)]_{,t}$ are given in (2.26). The above system is considered with the initial conditions

$$\begin{aligned}
 \mathbf{u}^m(0) &= \mathbf{u}_0^m, & \mathbf{u}_t^m(0) &= \mathbf{u}_1^m, & \chi^m(0) &= \chi_0, \\
 \mathbf{u}_{tt}^m(0) &= \mathbf{u}_2^m, & \chi_t^m(0) &= \chi_1^m, & \mu^m(0) &= \mu_0^m,
 \end{aligned}
 \tag{4.2}$$

where $\mathbf{u}_2^m \in \mathbf{V}_{0m}$ and $\chi_1^m, \mu_0^m \in V_m$ are the projections of the data \mathbf{u}_2, χ_1 and μ_0 , respectively defined in (2.19).

We assume that the following convergences in the strong sense are satisfied:

$$\begin{aligned}
 \mathbf{u}_0^m &\rightarrow \mathbf{u}_0 & \text{in } \mathbf{H}^3(\Omega) \cap \mathbf{H}_0^1(\Omega), & & \mathbf{u}_1^m &\rightarrow \mathbf{u}_1 & \text{in } \mathbf{H}_0^1(\Omega), \\
 \chi_0^m &\rightarrow \chi_0 & \text{in } H^4(\Omega) \cap H_N^2(\Omega), & & \mathbf{u}_2^m &\rightarrow \mathbf{u}_2 & \text{in } \mathbf{L}_2(\Omega), \\
 \chi_1^m &\rightarrow \chi_1 & \text{in } L_2(\Omega), & & \mu_0^m &\rightarrow \mu_0 & \text{in } H_N^2(\Omega).
 \end{aligned}
 \tag{4.3}$$

4.1. The basic estimate.

LEMMA 4.1. *Let (A1)–(A5) hold, the boundary of the domain Ω be of class C^4 , the function $z: \mathbb{R} \rightarrow [0, 1]$ be of class C^2 with property (2.20) and the data satisfy (2.21). Then a solution $(\mathbf{u}^m, \chi^m, \mu^m)$ of approximate problem (3.6) satisfies system (4.1) with the initial conditions (4.2). Moreover, $(\mathbf{u}^m, \chi^m, \mu^m)$ satisfy the estimates in Lemmas 3.1–3.4, and*

$$\begin{aligned}
 \|\mathbf{u}_{t't'}^m\|_{L_\infty(0,t;\mathbf{L}_2(\Omega))} + \|\mathbf{u}_{t'}^m\|_{L_\infty(0,t;\mathbf{H}^1(\Omega))} &\leq c_5(t), \\
 \|\chi_{t't'}^m\|_{L_\infty(0,t;L_2(\Omega))} + \|\chi_{t'}^m\|_{L_2(0,t;H_N^2(\Omega))} &\leq c_4(t)
 \end{aligned}
 \tag{4.4}$$

for $t \in (0, T]$, with constants $c_4(t), c_5(t)$ (independent of m) given by

$$c_4(t) = c(t^{1/2}E_1(t) + \|\chi_1\|_{L_2(\Omega)})[\exp a(t)]^{1/2}, \quad c_5(t) = t^{1/2}c_4(t),$$

where

$$\begin{aligned}
 E_1(t) &= t^{1/2}\|\mathbf{b}_{t'}\|_{L_2(\Omega^t)} + \|\mathbf{u}_2\|_{L_2(\Omega)} + \|\varepsilon(\mathbf{u}_1)\|_{L_2(\Omega)}, \\
 a(t) &= c(c_0)t^8 \exp(ct).
 \end{aligned}$$

PROOF. In the first step we estimate \mathbf{u}_t^m in terms of the $L_2(0, t; H^2(\Omega))$ -norm of χ_t^m . Testing (4.1)₁ by $\mathbf{u}_{tt}^m(t)$, we obtain

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} (|\mathbf{u}_{tt}^m|^2 + \varepsilon(\mathbf{u}_t^m) \cdot \mathbf{A}\varepsilon(\mathbf{u}_t^m)) dx = \int_{\Omega} ([z'(\chi^m)\mathbf{B}\nabla\chi^m]_{,t} + \mathbf{b}_t) \cdot \mathbf{u}_{tt}^m dx.$$

Hence, by the Cauchy–Schwarz inequality,

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left[\left(\int_{\Omega} (|\mathbf{u}_{tt}^m|^2 + \varepsilon(\mathbf{u}_t^m) \cdot \mathbf{A}\varepsilon(\mathbf{u}_t^m)) dx \right)^{1/2} \right]^2 \\ & \leq c \left(\int_{\Omega} ((\chi_t^m)^2 |\nabla \chi^m|^2 + |\nabla \chi_t^m|^2 + |\mathbf{b}_t|^2) dx \right)^{1/2} \\ & \quad \cdot \left(\int_{\Omega} (|\mathbf{u}_{tt}^m|^2 + \varepsilon(\mathbf{u}_t^m) \cdot \mathbf{A}\varepsilon(\mathbf{u}_t^m)) dx \right)^{1/2}, \end{aligned}$$

so

$$(4.5) \quad \frac{d}{dt} \left(\int_{\Omega} (|\mathbf{u}_{tt}^m|^2 + \varepsilon(\mathbf{u}_t^m) \cdot \mathbf{A}\varepsilon(\mathbf{u}_t^m)) dx \right)^{1/2} \leq c \left(\int_{\Omega} ((\chi_t^m)^2 |\nabla \chi^m|^2 + |\nabla \chi_t^m|^2 + |\mathbf{b}_t|^2) dx \right)^{1/2}.$$

Integrating (4.5) with respect to t and using the coercivity of \mathbf{A} (see (2.2)), it follows that

$$(4.6) \quad \begin{aligned} & \int_{\Omega} (|\mathbf{u}_{tt}^m|^2 + |\varepsilon(\mathbf{u}_t^m)|^2) dx \\ & \leq c \left[\int_0^t \left(\int_{\Omega} ((\chi_{t'}^m)^2 |\nabla \chi^m|^2 + |\nabla \chi_{t'}^m|^2 + |\mathbf{b}_{t'}|^2) dx \right)^{1/2} dt' \right]^2 \\ & \quad + \int_{\Omega} (|\mathbf{u}_{tt}^m(0)|^2 + |\varepsilon(\mathbf{u}_t^m(0))|^2) dx \\ & \leq ct \int_{\Omega^t} ((\chi_{t'}^m)^2 |\nabla \chi^m|^2 + |\nabla \chi_{t'}^m|^2) dx dt' + c(E_1^m(t))^2, \end{aligned}$$

where $E_1^m(t) := t^{1/2} \|\mathbf{b}_t\|_{L_2(\Omega^t)} + \|\mathbf{u}_2^m\|_{L_2(\Omega)} + \|\varepsilon(\mathbf{u}_1^m)\|_{L_2(\Omega)}$. Clearly, due to convergences (4.3), $E_1^m(t) \leq cE_1(t)$.

We shall estimate the first two terms on the right-hand side of (4.6). On account of Lemma 3.1, we obtain

$$(4.7) \quad \begin{aligned} \int_{\Omega^t} (\chi_{t'}^m)^2 |\nabla \chi^m|^2 dx dt' & \leq \sup_{t'} \int_{\Omega} |\nabla \chi^m|^2 dx \int_0^t \|\chi_{t'}^m\|_{L_{\infty}(\Omega)}^2 dt' \\ & \leq c_0 \int_0^t \|\chi_{t'}^m\|_{L_{\infty}(\Omega)}^2 dt' = I_1. \end{aligned}$$

Now, applying the interpolation inequality (see e.g. [3, Chapter III, Section 10])

$$\|\chi_t^m\|_{L_{\infty}(\Omega)} \leq \varepsilon^{1-\varkappa_1} \|\nabla^2 \chi_t^m\|_{L_2(\Omega)} + c\varepsilon^{-\varkappa_1} \|\chi_t^m\|_{L_2(\Omega)}$$

with $\varkappa_1 = 3/4$, $\varepsilon > 0$, and setting $\delta_1 = \varepsilon^{1/4}$, yields

$$(4.8) \quad I_1 \leq \delta_1 \int_0^t \|\nabla^2 \chi_{t'}^m\|_{L_2(\Omega)}^2 dt' + c(c_1)\delta_1^{-3} \int_0^t \|\chi_{t'}^m\|_{L_2(\Omega)}^2 dt'.$$

Similarly,

$$(4.9) \quad \int_{\Omega^t} |\nabla \chi_{t'}^m|^2 dx dt' \leq \delta_2 \int_0^t \|\nabla^2 \chi_{t'}^m\|_{L_2(\Omega)}^2 dt' + c\delta_2^{-1} \int_0^t \|\chi_{t'}^m\|_{L_2(\Omega)}^2 dt',$$

where the interpolation inequality

$$\|\nabla \chi_t^m\|_{L_2(\Omega)} \leq \varepsilon^{1-\varkappa_2} \|\nabla^2 \chi_t^m\|_{L_2(\Omega)} + c\varepsilon^{-\varkappa_2} \|\chi_t^m\|_{L_2(\Omega)}$$

with $\varkappa_2 = 1/2$, $\varepsilon > 0$ and $\delta_2 = \varepsilon^{1/2}$, was applied. Using (4.7)–(4.9) in (4.6) yields

$$\begin{aligned} \int_{\Omega} (|\mathbf{u}_{tt}^m|^2 + |\varepsilon(\mathbf{u}_t^m)|^2) dx &\leq ct(\delta_1 + \delta_2) \int_0^t \|\nabla^2 \chi_{t'}^m\|_{L_2(\Omega)}^2 dt' \\ &\quad + c(c_0)t(\delta_1^{-3} + \delta_2^{-1}) \int_0^t \|\chi_{t'}^m\|_{L_2(\Omega)}^2 dt' + c(E_1(t))^2. \end{aligned}$$

Hence, assuming $\delta_1 = \delta_2$ and choosing $\delta = ct\delta_1$ we arrive at the estimate

$$(4.10) \quad \begin{aligned} \|\mathbf{u}_{tt}^m\|_{L_2(\Omega)}^2 + \|\varepsilon(\mathbf{u}_t^m)\|_{L_2(\Omega)}^2 \\ \leq \delta \int_0^t \|\nabla^2 \chi_{t'}^m\|_{L_2(\Omega)}^2 dt' + c(1/\delta, c_0, t) \int_0^t \|\chi_{t'}^m\|_{L_2(\Omega)}^2 dt' + c(E_1(t))^2 \end{aligned}$$

for $t \in (0, T]$, where $c(1/\delta, c_0, t) = c(c_0)\delta^{-3}t^4$, $\delta > 0$ (arbitrary).

In the second step we consider system (4.1)₂, (4.1)₃ which on account of (3.2) can be rewritten in the form of the following equation:

$$(4.11) \quad (\chi_{tt}^m, w_j) = -(\Delta \chi_t^m, \Delta w_j) + ([\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)],_t, \Delta w_j).$$

Testing (4.11) by $\chi_t^m(t)$ gives

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} (\chi_t^m)^2 dx + \int_{\Omega} (\Delta \chi_t^m)^2 dx = \int_{\Omega} [\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)],_t \Delta \chi_t^m dx.$$

Hence, by the Young inequality, it follows that

$$(4.12) \quad \begin{aligned} \frac{d}{dt} \int_{\Omega} (\chi_t^m)^2 dx + \int_{\Omega} (\Delta \chi_t^m)^2 dx &\leq \int_{\Omega} [\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)],_t^2 dx \\ &\leq c \int_{\Omega} [(\chi^m)^4 (\chi_t^m)^2 + (\chi_t^m)^2 |\varepsilon(\mathbf{u}^m)|^2 + (\chi_t^m)^2 + |\varepsilon(\mathbf{u}_t^m)|^2] dx, \end{aligned}$$

where in the last inequality we used identity (2.26)₂ and the assumptions on ψ and z .

Let us examine the first two terms on the right-hand side of (4.12). Using the Hölder inequality and then recalling estimate (3.18), we obtain

$$(4.13) \quad \begin{aligned} \int_{\Omega} (\chi^m)^4 (\chi_t^m)^2 dx &\leq \sup_t \|\chi^m\|_{L_6(\Omega)}^4 \|\chi_t^m\|_{L_6(\Omega)}^2 \\ &\leq c_1 \|\chi_t^m\|_{L_6(\Omega)}^2 \leq \delta_3 \|\nabla^2 \chi_t^m\|_{L_2(\Omega)}^2 + c(1/\delta_3) \|\chi_t^m\|_{L_2(\Omega)}^2, \end{aligned}$$

for $\delta_3 > 0$, where in the last line the interpolation inequality was used.

Similarly, an application of the Hölder inequality, estimate (3.9) and the interpolation inequality to the second term on the right-hand side of (4.12) yields

$$(4.14) \quad \int_{\Omega} (\chi_t^m)^2 |\varepsilon(\mathbf{u}^m)|^2 dx \leq \sup_t \|\varepsilon(\mathbf{u}^m)\|_{L_2(\Omega)}^2 \|\chi_t^m\|_{L_{\infty}(\Omega)}^2 \\ \leq c_0 \|\chi_t^m\|_{L_{\infty}(\Omega)}^2 \leq \delta_4 \|\nabla^2 \chi_t^m\|_{L_2(\Omega)}^2 + c(1/\delta_4) \|\chi_t^m\|_{L_2(\Omega)}^2,$$

for $\delta_4 > 0$. Now, using the inequality

$$(4.15) \quad \|\chi_t^m\|_{H_N^2(\Omega)} \leq c \|\Delta \chi_t^m\|_{L_2(\Omega)},$$

which holds true because $\int_{\Omega} \chi_t^m dx = 0$ (see (3.20)), applying (4.13) and (4.14) in (4.12), and choosing δ_3, δ_4 sufficiently small, we conclude that

$$(4.16) \quad \frac{d}{dt} \|\chi_t^m\|_{L_2(\Omega)}^2 + \|\chi_t^m\|_{H_N^2(\Omega)}^2 \leq c(\|\chi_t^m\|_{L_2(\Omega)}^2 + \|\varepsilon(\mathbf{u}^m)\|_{L_2(\Omega)}^2).$$

At this point we apply estimate (4.10) to the second term on the right-hand side of (4.16). This leads to

$$(4.17) \quad \frac{d}{dt} \|\chi_t^m\|_{L_2(\Omega)}^2 + \|\chi_t^m\|_{H_N^2(\Omega)}^2 \leq c \|\chi_t^m\|_{L_2(\Omega)}^2 + \delta \int_0^t \|\nabla^2 \chi_{t'}^m\|_{L_2(\Omega)}^2 dt' \\ + c(c_0) \delta^{-3} t^4 \int_0^t \|\chi_{t'}^m\|_{L_2(\Omega)}^2 dt' + c(E_1(t))^2$$

for $t \in (0, T]$. Multiplying (4.17) by e^{-ct} , integrating with respect to t , and using that, on account of (4.3), $\|\chi_1^m\|_{L_2(\Omega)} \leq c \|\chi_1\|_{L_2(\Omega)}$, we arrive at

$$e^{-ct} \|\chi_t^m(t)\|_{L_2(\Omega)}^2 + \int_0^t e^{-ct'} \|\chi_{t'}^m(t')\|_{H_N^2(\Omega)}^2 dt' \\ \leq \delta \int_0^t e^{-ct'} \int_0^{t'} \|\nabla^2 \chi_{t''}^m(t'')\|_{L_2(\Omega)}^2 dt'' dt' \\ + c(c_0) \delta^{-3} \int_0^t (t')^4 e^{-ct'} \int_0^{t'} \|\chi_{t''}^m(t'')\|_{L_2(\Omega)}^2 dt'' dt' \\ + c \int_0^t e^{-ct'} (E_1(t'))^2 dt' + \|\chi_1\|_{L_2(\Omega)}^2.$$

Hence,

$$(4.18) \quad \|\chi_t^m(t)\|_{L_2(\Omega)}^2 + \int_0^t \|\chi_{t'}^m(t')\|_{H_N^2(\Omega)}^2 dt' \leq \delta t e^{ct} \int_0^t \|\chi_{t'}^m(t')\|_{H_N^2(\Omega)}^2 dt' \\ + c(c_0) \delta^{-3} t^5 e^{ct} \int_0^t \|\chi_{t'}^m(t')\|_{L_2(\Omega)}^2 dt' + c t e^{ct} (E_1(t))^2 + e^{ct} \|\chi_1\|_{L_2(\Omega)}^2.$$

Choosing $\delta = t^{-1}e^{-ct}/2$, we obtain

$$\begin{aligned} \|\chi_t^m(t)\|_{L_2(\Omega)}^2 + \int_0^t \|\chi_{t'}^m(t')\|_{H_N^2(\Omega)}^2 dt' \\ \leq c(c_0)t^8 e^{ct} \int_0^t \|\chi_{t'}^m(t')\|_{L_2(\Omega)}^2 dt' + cte^{ct}(E_1(t))^2 + e^{ct}\|\chi_1\|_{L_2(\Omega)}^2. \end{aligned}$$

Now, the application of the Gronwall lemma yields

$$(4.19) \quad \begin{aligned} \|\chi_t^m(t)\|_{L_2(\Omega)}^2 + \int_0^t \|\chi_{t'}^m(t')\|_{H_N^2(\Omega)}^2 dt' \\ \leq c(t(E_1(t))^2 + \|\chi_1\|_{L_2(\Omega)}) \exp(ct + c(c_0)t^8 \exp(ct)) \leq c_4^2(t) \end{aligned}$$

for almost all $t \in (0, T]$. This proves estimate (4.4)₂. Applying (4.19) in (4.10) and setting there $\delta = t$ we conclude, by virtue of Korn’s inequality, estimate (4.4)₁. Thereby the proof is completed. \square

4.2. Further estimates. Firstly, we note that in view of the inequality

$$|\chi_{xx}(t) - \chi_{xx}(t')| \leq |t - t'|^{1/2} \left(\int_{t'}^t \chi_{t''xx}^2 dt'' \right)^{1/2},$$

estimate (4.4)₂ implies that $\chi^m \in C^{1/2}([0, T]; H_N^2(\Omega))$, and

$$(4.20) \quad \|\chi^m\|_{C^{1/2}([0,t]; H_N^2(\Omega))} \leq c_4(t).$$

Next, we prove the following

LEMMA 4.2. *Let assumptions of Lemma 4.1 hold. Then, for $t \in (0, T]$,*

$$(4.21) \quad \|\mathbf{u}^m\|_{L_\infty(0,t; H^2(\Omega))} \leq c_5(t).$$

PROOF. Using (3.1) we rewrite (3.6)₁ in the form

$$(\mathbf{u}_{tt}^m, \mathbf{Q}\mathbf{v}_j) - (\mathbf{Q}\mathbf{u}^m, \mathbf{Q}\mathbf{v}_j) = (z'(\chi^m)\mathbf{B}\nabla\chi^m + \mathbf{b}, \mathbf{Q}\mathbf{v}_j).$$

Testing this equality by $\mathbf{u}^m(t)$ gives

$$\|\mathbf{Q}\mathbf{u}^m\|_{L_2(\Omega)}^2 = (\mathbf{u}_{tt}^m - z'(\chi^m)\mathbf{B}\nabla\chi^m + \mathbf{b}, \mathbf{Q}\mathbf{u}^m) \quad \text{for a.a. } t \in (0, T].$$

Hence, using the Cauchy–Schwarz inequality and then recalling estimates (3.9) and (4.4)₁, we obtain

$$\begin{aligned} \|\mathbf{Q}\mathbf{u}^m\|_{L_\infty(0,t; L_2(\Omega))} \\ \leq c(\|\mathbf{u}_{t't'}^m\|_{L_\infty(0,t; L_2(\Omega))} + \|\nabla\chi^m\|_{L_\infty(0,t; L_2(\Omega))} + \|\mathbf{b}\|_{L_\infty(0,t; L_2(\Omega))}) \\ \leq c(c_5(t) + c_0 + 1) \leq c_5(t). \end{aligned}$$

This, by the ellipticity property of \mathbf{Q} (see (2.3)), implies (4.21). \square

The next result provides an additional regularity estimate for μ^m .

LEMMA 4.3. *Under assumptions of Lemma 4.1,*

$$(4.22) \quad \|\mu^m\|_{L_\infty(0,t;H_N^2(\Omega))} \leq c_4(t), \quad \|\mu_{t'}^m\|_{L_2(\Omega^t)} \leq t^{1/2}c_5(t), \quad t \in (0, T].$$

PROOF. Using (3.2) we rewrite identity (3.6)₂ in the form

$$(\chi_t^m, \Delta w_j) = (\Delta \mu^m, \Delta w_j).$$

Testing this equality by $\mu^m(t)$ and using the Cauchy–Schwarz inequality yields

$$\|\Delta \mu^m\|_{L_2(\Omega)} \leq \|\chi_t^m\|_{L_2(\Omega)} \quad \text{for a.a. } t \in (0, T].$$

Consequently, due to estimate (4.4)₂,

$$(4.23) \quad \|\Delta \mu^m\|_{L_\infty(0,t;L_2(\Omega))} \leq \|\chi_{t'}^m\|_{L_\infty(0,t;L_2(\Omega))} \leq c_4(t).$$

Thus, recalling bound (3.22) on the mean value of μ^m , estimate (4.22)₁ follows from (4.23) on account of the ellipticity property of the Laplace operator.

To show (4.22)₂ we test identity (4.1)₃ by $\mu_t^m(t)$. Then, with the help of the Cauchy–Schwarz inequality, it follows that

$$(4.24) \quad \|\mu_t^m\|_{L_2(\Omega)} \leq \|\Delta \chi_t^m\|_{L_2(\Omega)} + \|[\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)]_{,t}\|_{L_2(\Omega)}.$$

The second term on the right-hand side of (4.24) can be estimated with the help of bounds (4.12)–(4.14) (with $\delta_3 = \delta_4 = 1$) in the proof of Lemma 4.1. Consequently, we obtain

$$\begin{aligned} \|\mu_t^m\|_{L_2(\Omega^t)} &\leq c(\|\Delta \chi_t^m\|_{L_2(\Omega^t)} + \|\nabla^2 \chi_t^m\|_{L_2(\Omega^t)} + \|\chi_t^m\|_{L_2(\Omega^t)} + \|\varepsilon(\mathbf{u}_t^m)\|_{L_2(\Omega^t)}) \\ &\leq c(c_4(t) + t^{1/2}c_5(t)) \leq t^{1/2}c_5(t), \end{aligned}$$

where in the last line Lemma 4.1 was used. This completes the proof. \square

Finally, we estimate time derivatives \mathbf{u}_{ttt}^m and χ_{tt}^m .

LEMMA 4.4. *Under assumptions of Lemma 4.1,*

$$(4.25) \quad \begin{aligned} \|\mathbf{u}_{t't't'}^m\|_{L_2(0,t;(\mathbf{H}_0^1(\Omega))')} &\leq t^{1/2}c_5(t), \\ \|\chi_{t't'}^m\|_{L_2(0,t;(H_N^2(\Omega))')} &\leq t^{1/2}c_5(t), \quad t \in (0, T]. \end{aligned}$$

PROOF. We proceed similarly as in Lemma 3.4. For $\boldsymbol{\eta} \in L_2(0, T; \mathbf{H}_0^1(\Omega))$ we test (4.1)₁ by $\boldsymbol{\eta}^m = \mathbf{P}^m \boldsymbol{\eta}$, where the projectin \mathbf{P}^m is defined by (3.27). Then

$$\begin{aligned} &\left| \int_0^t (\mathbf{u}_{t't't'}^m, \boldsymbol{\eta}) dt' \right| \\ &= \left| \int_0^t \{-(\mathbf{A}\varepsilon(\mathbf{u}_{t'}^m), \varepsilon(\mathbf{P}^m \boldsymbol{\eta})) + ([z'(\chi^m)\mathbf{B}\nabla \chi^m + \mathbf{b}]_{,t'}, \mathbf{P}^m \boldsymbol{\eta})\} dt' \right| \\ &\leq c[\|\varepsilon(\mathbf{u}_{t'}^m)\|_{L_2(\Omega^t)} \|\nabla \mathbf{P}^m \boldsymbol{\eta}\|_{L_2(\Omega^t)} \\ &\quad + (\|\chi_{t'}^m \nabla \chi^m\|_{L_2(\Omega^t)} + \|\nabla \chi_{t'}^m\|_{L_2(\Omega^t)} + \|\mathbf{b}_{t'}\|_{L_2(\Omega^t)}) \|\mathbf{P}^m \boldsymbol{\eta}\|_{L_2(\Omega^t)}]. \end{aligned}$$

Hence, recalling Lemmas 3.1, 4.1, and the estimate

$$\|\chi_{t'}^m \nabla \chi^m\|_{L_2(\Omega^t)} \leq \|\chi_{t'}^m\|_{L_2(0,t;L_\infty(\Omega))} \|\nabla \chi^m\|_{L_\infty(0,t;L_2(\Omega))} \leq c_0 c_4(t),$$

it follows that

$$\begin{aligned} \left| \int_0^t (\mathbf{u}_{t't'}^m, \boldsymbol{\eta}) dt' \right| &\leq c(t^{1/2} c_5(t) + c_4(t) + \|\mathbf{b}_{t'}\|_{L_2(\Omega^t)}) \|\mathbf{P}^m \boldsymbol{\eta}\|_{L_2(0,t;H^1(\Omega))} \\ &\leq t^{1/2} c_5(t) \|\boldsymbol{\eta}\|_{L_2(0,t;H^1(\Omega))} \end{aligned}$$

for all $\boldsymbol{\eta} \in L_2(0,t;H_0^1(\Omega))$. This shows (4.25)₁.

Similarly, for any $\xi \in L_2(0,T;H_N^2(\Omega))$, testing (4.1)₂ by $\xi^m = P^m \xi$, where the projection P^m is defined by (3.28), we obtain

$$\begin{aligned} \left| \int_0^t (\chi_{t't'}^m, \xi) dt' \right| &= \left| \int_0^t (\mu_{t'}^m, \Delta P^m \xi) dt' \right| \\ &\leq \|\mu_{t'}^m\|_{L_2(\Omega^t)} \|\Delta P^m \xi\|_{L_2(\Omega^t)} \leq t^{1/2} c_5(t) \|\xi\|_{L_2(0,t;H_N^2(\Omega))}, \end{aligned}$$

where in the last inequality Lemma 4.3 was applied. This shows (4.25)₂. \square

5. Proof of Theorem 2.1

From Lemmas 3.2–3.4 it follows that there exists a triple (\mathbf{u}, χ, μ) with

$$\begin{aligned} \mathbf{u} &\in L_\infty(0,T;H_0^1(\Omega)), \quad \mathbf{u}_t \in L_\infty(0,T;L_2(\Omega)), \quad \mathbf{u}_{tt} \in L_2(0,T;(H_0^1(\Omega))'), \\ (5.1) \quad \chi &\in L_\infty(0,T;H^1(\Omega)) \cap L_2(0,T;H_N^2(\Omega)), \quad \chi_t \in L_2(0,T;(H^1(\Omega))'), \\ &\mu \in L_2(0,T;H^1(\Omega)), \end{aligned}$$

and a subsequence of solutions $(\mathbf{u}^m, \chi^m, \mu^m)$ to problem (3.6) (which we still denote by the same indices) such that as $m \rightarrow \infty$:

$$\begin{aligned} (5.2) \quad \mathbf{u}^m &\rightarrow \mathbf{u} \quad \text{weakly-}^* \text{ in } L_\infty(0,T;H_0^1(\Omega)), \\ \mathbf{u}_t^m &\rightarrow \mathbf{u}_t \quad \text{weakly-}^* \text{ in } L_\infty(0,T;L_2(\Omega)), \\ \mathbf{u}_{tt}^m &\rightarrow \mathbf{u}_{tt} \quad \text{weakly in } L_2(0,T;(H_0^1(\Omega))'), \\ \chi^m &\rightarrow \chi \quad \text{weakly-}^* \text{ in } L_\infty(0,T;H^1(\Omega)) \text{ and} \\ &\quad \text{weakly in } L_2(0,T;H_N^2(\Omega)), \\ \chi_t^m &\rightarrow \chi_t \quad \text{weakly in } L_2(0,T;(H^1(\Omega))'), \\ \mu^m &\rightarrow \mu \quad \text{weakly in } L_2(0,T;H^1(\Omega)). \end{aligned}$$

Using the compactness results (see e.g. Lions [12], Simon [25, Section 8]) it follows that for a subsequence (still denoted by the same indices)

$$\begin{aligned}
 \mathbf{u}^m &\rightarrow \mathbf{u} && \text{strongly in } L_2(0, T; \mathbf{L}_q(\Omega)) \cap C([0, T]; \mathbf{L}_q(\Omega)), \quad q < 6, \\
 &&& \text{and a.e. in } \Omega^T, \\
 (5.3) \quad \mathbf{u}_t^m &\rightarrow \mathbf{u}_t && \text{strongly in } C([0, T]; (\mathbf{H}_0^1(\Omega))'), \\
 \chi^m &\rightarrow \chi && \text{strongly in } L_2(0, T; H^1(\Omega)) \cap C([0, T]; L_2(\Omega)) \\
 &&& \text{and a.e. in } \Omega^T.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 \mathbf{u}^m(0) = \mathbf{u}_0^m &\rightarrow \mathbf{u}(0) && \text{strongly in } \mathbf{L}_q(\Omega), \quad q < 6, \\
 \mathbf{u}_t^m(0) = \mathbf{u}_1^m &\rightarrow \mathbf{u}_t(0) && \text{strongly in } (\mathbf{H}_0^1(\Omega))', \\
 \chi^m(0) = \chi_0^m &\rightarrow \chi(0) && \text{strongly in } L_2(\Omega),
 \end{aligned}$$

what together with convergences (3.7) implies that

$$(5.4) \quad \mathbf{u}(0) = \mathbf{u}_0, \quad \mathbf{u}_t(0) = \mathbf{u}_1, \quad \chi(0) = \chi_0.$$

The relations (5.1) and (5.4) imply assertion (2.13) of the theorem.

Now, let us introduce the following weak formulation of (3.6):

$$\begin{aligned}
 (5.5) \quad &\int_0^T \langle \mathbf{u}_{tt}^m, \boldsymbol{\eta} \rangle_{(\mathbf{H}_0^1(\Omega))', \mathbf{H}_0^1(\Omega)} dt + \int_0^T (\mathbf{A}\boldsymbol{\varepsilon}(\mathbf{u}^m), \boldsymbol{\varepsilon}(\boldsymbol{\eta})) dt \\
 &= \int_0^T (z'(\chi^m) \mathbf{B}\nabla\chi^m + \mathbf{b}, \boldsymbol{\eta}) dt \quad \text{for all } \boldsymbol{\eta} \in L_2(0, T; \mathbf{V}_{0m}),
 \end{aligned}$$

$$\begin{aligned}
 (5.6) \quad &\int_0^T \langle \chi_t^m, \xi \rangle_{(H^1(\Omega))', H^1(\Omega)} dt + \int_0^T (\nabla\mu^m, \nabla\xi) dt = 0 \\
 &\text{for all } \xi \in L_2(0, T; V_m),
 \end{aligned}$$

$$\begin{aligned}
 (5.7) \quad &\int_0^T (\mu^m, \zeta) dt = - \int_0^T (\Delta\chi^m, \zeta) dt + \int_0^T (\psi'(\chi^m) + W_{,\chi}(\boldsymbol{\varepsilon}(\mathbf{u}^m), \chi^m), \zeta) dt \\
 &\text{for all } \zeta \in L_2(0, T; V_m).
 \end{aligned}$$

To pass to the limit $m \rightarrow \infty$ in identities (5.5)–(5.7) we follow the standard procedure (see e.g. Lions–Magenes [13]). Namely, we fix $m = m_0 \in \mathbb{N}$ in the spaces of test functions $\boldsymbol{\eta}$, ξ , ζ and take subsequences (5.2) with $m \geq m_0$. Clearly, by virtue of the weak convergences (5.2), the linear terms in (5.5)–(5.7) converge to the corresponding limits. Thus, it remains to examine the convergence of the nonlinear terms $z'(\chi^m) \mathbf{B}\nabla\chi^m$ and $\psi'(\chi^m) + W_{,\chi}(\boldsymbol{\varepsilon}(\mathbf{u}^m), \chi^m)$.

Recalling the growth conditions (2.7), and using the energy bounds (3.9), (3.18), it follows that

$$\begin{aligned} \|z'(\chi^m)\mathbf{B}\nabla\chi^m\|_{L_\infty(0,T;\mathbf{L}_2(\Omega))} &\leq c\|\nabla\chi^m\|_{L_\infty(0,T;\mathbf{L}_2(\Omega))} \leq cc_0, \\ \|\psi'(\chi^m)\|_{L_\infty(0,T;L_2(\Omega))} &\leq c(\|\chi^m\|_{L_\infty(0,T;L_6(\Omega))}^3 + 1) \leq c(c_1), \\ \|W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)\|_{L_\infty(0,T;L_2(\Omega))} &\leq c(\|\varepsilon(\mathbf{u}^m)\|_{L_\infty(0,T;\mathbf{L}_2(\Omega))} + 1) \leq c(c_0). \end{aligned}$$

Thanks to these uniform in m estimates and the pointwise convergences (5.3) we can apply the standard nonlinear convergence lemma (see Lions [12, Chapter 1, Lemma 1.3]) to conclude that

$$\begin{aligned} z'(\chi^m)\mathbf{B}\nabla\chi^m &\rightharpoonup z(\chi)\mathbf{B}\nabla\chi && \text{weakly-}^* \text{ in } L_\infty(0,T;\mathbf{L}_2(\Omega)), \\ \psi'(\chi^m) &= (\chi^m)^3 - \chi^m \rightharpoonup \chi^3 - \chi = \psi(\chi) && \text{weakly-}^* \text{ in } L_\infty(0,T;L_2(\Omega)), \end{aligned}$$

$$\begin{aligned} W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m) &= z'(\chi^m)(\mathbf{B} \cdot \varepsilon(\mathbf{u}^m) + Dz(\chi^m) + E) \rightharpoonup \\ &= z'(\chi)(\mathbf{B} \cdot \varepsilon(\mathbf{u}) + Dz(\chi) + E) = W_{,\chi}(\varepsilon(\mathbf{u}), \chi) \\ &\text{weakly-}^* \text{ in } L_\infty(0,T;L_2(\Omega)). \end{aligned}$$

Consequently, passing to the limit in (5.5)–(5.7) for a subsequence $m_0 \leq m \rightarrow \infty$, we conclude that the identities in Theorem 2.1 are satisfied for all test functions $\boldsymbol{\eta} \in L_2(0,T;\mathbf{V}_{0m_0})$, $\xi \in L_2(0,T;V_{m_0})$ and $\zeta \in L_2(0,T;V_{m_0})$. Next, passing to the limit $m_0 \rightarrow \infty$, we arrive by density arguments at identities (2.14). Clearly, a priori estimates (2.15)–(2.18) are the consequences of the uniform estimates in Lemmas 3.1–3.4 and the weak convergences (5.2). This proves the theorem. \square

6. Proof of Theorem 2.2

From Lemmas 3.1–3.4 and 4.1–4.4 it follows that there exists a triple (\mathbf{u}, χ, μ) with

$$(6.1) \quad \begin{aligned} \mathbf{u} &\in L_\infty(0,T;\mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)), \quad \mathbf{u}_t \in L_\infty(0,T;\mathbf{H}^1(\Omega)), \\ \mathbf{u}_{tt} &\in L_\infty(0,T;\mathbf{L}_2(\Omega)), \quad \mathbf{u}_{ttt} \in L_2(0,T;(\mathbf{H}_0^1(\Omega))'), \end{aligned}$$

$$(6.2) \quad \begin{aligned} \chi &\in C^{1/2}([0,T];H_N^2(\Omega)), \quad \chi_t \in L_\infty(0,T;L_2(\Omega)) \cap L_2(0,T;H_N^2(\Omega)), \\ \chi_{tt} &\in L_2(0,T;(H_N^2(\Omega))'), \\ \mu &\in L_\infty(0,T;H_N^2(\Omega)), \quad \mu_t \in L_2(\Omega^T), \end{aligned}$$

and a subsequence of solutions $(\mathbf{u}^m, \chi^m, \mu^m)$ to problem (3.6) (which we still denote by the same indices) such that as $m \rightarrow \infty$:

$$\begin{aligned}
(6.3) \quad & \mathbf{u}^m \rightarrow \mathbf{u} \quad \text{weakly-}^* \text{ in } L_\infty(0, T; \mathbf{H}^2(\Omega)), \\
& \mathbf{u}_t^m \rightarrow \mathbf{u}_t \quad \text{weakly-}^* \text{ in } L_\infty(0, T; \mathbf{H}^1(\Omega)), \\
& \mathbf{u}_{tt}^m \rightarrow \mathbf{u}_{tt} \quad \text{weakly-}^* \text{ in } L_\infty(0, T; \mathbf{L}_2(\Omega)), \\
& \mathbf{u}_{ttt}^m \rightarrow \mathbf{u}_{ttt} \quad \text{weakly in } L_2(0, T; (\mathbf{H}_0^1(\Omega))'), \\
& \chi^m \rightarrow \chi \quad \text{weakly-}^* \text{ in } L_\infty(0, T; H_N^2(\Omega)), \\
& \chi_t^m \rightarrow \chi_t \quad \text{weakly-}^* \text{ in } L_\infty(0, T; L_2(\Omega)) \text{ and} \\
& \quad \quad \quad \text{weakly in } L_2(0, T; H_N^2(\Omega)), \\
& \chi_{tt}^m \rightarrow \chi_{tt} \quad \text{weakly in } L_2(0, T; (H_N^2(\Omega))'), \\
& \mu^m \rightarrow \mu \quad \text{weakly-}^* \text{ in } L_\infty(0, T; H_N^2(\Omega)), \\
& \mu_t^m \rightarrow \mu_t \quad \text{weakly in } L_2(\Omega).
\end{aligned}$$

Using the compactness results [25] it follows that for a subsequence (still denoted by the same indices)

$$\begin{aligned}
(6.4) \quad & \mathbf{u}^m \rightarrow \mathbf{u} \quad \text{strongly in } L_2(0, T; \mathbf{H}_0^1(\Omega)) \cap C([0, T]; \mathbf{L}_2(\Omega)) \\
& \quad \quad \quad \text{and a.e. in } \Omega^T, \\
& \mathbf{u}_t^m \rightarrow \mathbf{u}_t \quad \text{strongly in } \mathbf{L}_2(\Omega^T) \cap C([0, T]; \mathbf{L}_2(\Omega)) \\
& \quad \quad \quad \text{and a.e. in } \Omega^T, \\
& \mathbf{u}_{tt}^m \rightarrow \mathbf{u}_{tt} \quad \text{strongly in } C([0, T]; (\mathbf{H}_0^1(\Omega))'), \\
& \chi^m \rightarrow \chi \quad \text{strongly in } L_2(0, T; H^1(\Omega)) \cap C([0, T]; H^1(\Omega)) \\
& \quad \quad \quad \text{and a.e. in } \Omega^T, \\
& \chi_t^m \rightarrow \chi_t \quad \text{strongly in } L_2(0, T; H^1(\Omega)) \cap C([0, T]; (H_N^2(\Omega))') \\
& \quad \quad \quad \text{and a.e. in } \Omega^T, \\
& \mu^m \rightarrow \mu \quad \text{strongly in } L_2(0, T; H^1(\Omega)) \cap C([0, T]; H^1(\Omega)) \\
& \quad \quad \quad \text{and a.e. in } \Omega^T.
\end{aligned}$$

Hence,

$$\begin{aligned}
& \mathbf{u}^m(0) = \mathbf{u}_0^m \rightarrow \mathbf{u}(0) \quad \text{strongly in } \mathbf{H}_0^1(\Omega), \\
& \mathbf{u}_t^m(0) = \mathbf{u}_1^m \rightarrow \mathbf{u}_t(0) \quad \text{strongly in } \mathbf{L}_2(\Omega), \\
& \mathbf{u}_{tt}^m(0) = \mathbf{u}_2^m \rightarrow \mathbf{u}_{tt}(0) \quad \text{strongly in } (\mathbf{H}_0^1(\Omega))', \\
& \chi^m(0) = \chi_0^m \rightarrow \chi(0) \quad \text{strongly in } H^1(\Omega), \\
& \chi_t^m(0) = \chi_1^m \rightarrow \chi_t(0) \quad \text{strongly in } (H_N^2(\Omega))', \\
& \mu^m(0) = \mu_0^m \rightarrow \mu(0) \quad \text{strongly in } H^1(\Omega),
\end{aligned}$$

what together with convergences (4.3) implies that

$$\begin{aligned}
(6.5) \quad & \mathbf{u}(0) = \mathbf{u}_0, \quad \mathbf{u}_t(0) = \mathbf{u}_1, \quad \mathbf{u}_{tt}(0) = \mathbf{u}_2, \\
& \chi(0) = \chi_0, \quad \chi_t(0) = \chi_1, \quad \mu(0) = \mu_0.
\end{aligned}$$

The relations (6.1)–(6.2) and (6.5) imply assertion (2.22) of the theorem.

We introduce now the following weak formulation of system (4.1):

$$\begin{aligned}
 & \int_0^T \langle \mathbf{u}_{tt}^m, \boldsymbol{\eta} \rangle_{(\mathbf{H}_0^1(\Omega))', \mathbf{H}_0^1(\Omega)} dt + \int_0^T (\mathbf{A}\varepsilon(\mathbf{u}_t^m), \varepsilon(\boldsymbol{\eta})) dt \\
 &= \int_0^T ([z'(\chi^m) \mathbf{B}\nabla\chi^m]_{,t} + \mathbf{b}_t, \boldsymbol{\eta}) dt \quad \text{for all } \boldsymbol{\eta} \in H^1(0, T; \mathbf{V}_{0m}), \boldsymbol{\eta}(T) = 0, \\
 (6.6) \quad & \int_0^T \langle \chi_{tt}^m, \xi \rangle_{(H_N^2(\Omega))', H_N^2(\Omega)} = \int_0^T (\mu_t, \Delta\xi) dt \\
 & \quad \text{for all } \xi \in H^1(0, T; V_m), \xi(T) = 0, \\
 & \int_0^T (\mu_t^m, \zeta) dt = - \int_0^T (\Delta\chi_t^m, \zeta) dt + \int_0^T ([\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)]_{,t}, \zeta) dt \\
 & \quad \text{for all } \zeta \in H^1(0, T; V_m), \zeta(T) = 0.
 \end{aligned}$$

We pass to the limit $m \rightarrow \infty$ in a similar fashion as in the proof of Theorem 2.1. Clearly, due to the weak convergences (6.3), all linear terms in identities (6.6) converge to the corresponding limits. It remains to examine the convergence of the nonlinear terms $[z'(\chi^m) \mathbf{B}\nabla\chi^m]_{,t}$ and $[\psi'(\chi^m) + W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)]_{,t}$ whose explicit expressions are given in (2.26).

Recalling assumptions on z, ψ and using Lemmas 3.1, 4.1 we obtain the following bounds (these bounds can be also directly concluded from the proofs of Lemmas 4.3, 4.4):

$$\begin{aligned}
 & \| [z'(\chi^m) \mathbf{B}\nabla\chi^m]_{,t} \|_{L_2(\Omega^T)} \leq c(\|\chi_t^m \nabla\chi^m\|_{L_2(\Omega^T)} + \|\nabla\chi_t^m\|_{L_2(\Omega^T)}) \\
 & \leq c(\|\chi_t^m\|_{L_2(0, T; L_\infty(\Omega))} \|\nabla\chi^m\|_{L_\infty(0, T; L_2(\Omega))} + \|\nabla\chi_t^m\|_{L_2(\Omega^T)}) \\
 & \leq c(c_0 + 1)c_4(T), \\
 & \| [\psi'(\chi^m)]_{,t} \|_{L_2(\Omega^T)} \leq c\|(\chi^m)^2 + 1\|_{L_2(\Omega^T)} \|\chi_t^m\|_{L_2(\Omega^T)} \\
 & \leq c(\|\chi^m\|_{L_\infty(0, T; L_4(\Omega))}^2 \|\chi_t^m\|_{L_2(0, T; L_\infty(\Omega))} + \|\chi_t^m\|_{L_2(\Omega^T)}) \\
 & \leq c(c_1^2 + 1)c_4(T), \\
 & \| [W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)]_{,t} \|_{L_2(\Omega^T)} \\
 & \leq c(\|\chi_t^m \varepsilon(\mathbf{u}^m)\|_{L_2(\Omega^T)} + \|\chi_t^m\|_{L_2(\Omega^T)} + \|\varepsilon(\mathbf{u}_t^m)\|_{L_2(\Omega^T)}) \\
 & \leq c(\|\chi_t^m\|_{L_2(0, T; L_\infty(\Omega))} \|\varepsilon(\mathbf{u}^m)\|_{L_\infty(0, T; L_2(\Omega))} + \|\chi_t^m\|_{L_2(\Omega^T)} \\
 & \quad + \|\varepsilon(\mathbf{u}_t^m)\|_{L_2(\Omega^T)}) \leq c(c_0 + 1)c_4(T) + c_5(T).
 \end{aligned}$$

Thanks to these uniform in m bounds and the pointwise convergences (6.4) we can apply the nonlinear convergence lemma (see [12, Chapter 1, Lemma 1.3]) to conclude that

$$\begin{aligned}
 (6.7) \quad & [z'(\chi^m) \mathbf{B}\nabla\chi^m]_{,t} = z''(\chi^m) \chi_t^m \mathbf{B}\nabla\chi^m + z'(\chi^m) \mathbf{B}\nabla\chi_t^m \\
 & \rightarrow z''(\chi) \chi_t \mathbf{B}\nabla\chi + z'(\chi) \mathbf{B}\nabla\chi_t = [z'(\chi) \mathbf{B}\nabla\chi]_{,t} \quad \text{weakly in } L_2(\Omega^T),
 \end{aligned}$$

$$(6.8) \quad [\psi'(\chi^m)]_{,t} = (3(\chi^m)^2 - 1)\chi_t^m \rightarrow (3\chi^2 - 1)\chi_t = [\psi'(\chi)]_{,t}$$

weakly in $L_2(\Omega^T)$,

$$(6.9) \quad [W_{,\chi}(\varepsilon(\mathbf{u}^m), \chi^m)]_{,t}$$

$$= z''(\chi^m)\chi_t^m(\mathbf{B} \cdot \varepsilon(\mathbf{u}^m) + Dz(\chi^m) + E) + z'(\chi^m)(\mathbf{B} \cdot \varepsilon(\mathbf{u}_t^m) + Dz'(\chi^m)\chi_t^m)$$

$$\rightarrow z''(\chi)\chi_t(\mathbf{B} \cdot \varepsilon(\mathbf{u}) + Dz(\chi) + E) + z'(\chi)(\mathbf{B} \cdot \varepsilon(\mathbf{u}_t) + Dz'(\chi)\chi_t)$$

$$= [W_{,\chi}(\varepsilon(\mathbf{u}), \chi)]_{,t} \quad \text{weakly in } L_2(\Omega^T).$$

In view of (6.7)–(6.9), passing to the limit $m \rightarrow \infty$ in identities (6.6) we conclude (2.23)–(2.25). We also note that a priori estimates (2.27) result immediately from the estimates in Lemmas 3.1–4.3, 4.1–4.4 and the weak convergences (6.3). This completes the proof of the theorem. \square

REFERENCES

- [1] Ł. BARTKOWIAK, I. PAWŁOW, *The Cahn–Hilliard–Gurtin system coupled with elasticity*, Control Cybernet. **34** (2005), no. 4, 1005–1043.
- [2] E. BONETTI, P. COLLI, W. DREYER, G. GILIARDI, G. SCHIMPERNA AND J. SPREKELS, *On a model for phase separation in binary alloys driven by mechanical effects*, Physica D **165** (2002), 48–65.
- [3] O. V. BESOV, V. P. IL'IN AND S. M. NIKOLSKI, *Integral Representations of Functions and Imbedding Theorems*, Nauka, Moscow, 1996. (in Russian)
- [4] M. CARRIVE, A. MIRANVILLE AND A. PIÉTRUS, *The Cahn–Hilliard equation for deformable elastic continua*, Adv. Math. Sci. Appl. **10** (2000), no. 2, 539–569.
- [5] M. CARRIVE, A. MIRANVILLE, A. PIÉTRUS AND J. M. RAKOTOSON, *The Cahn–Hilliard equation for an isotropic deformable continuum*, Appl. Math. Letters **12** (1999), 23–28.
- [6] CH. M. ELLIOTT AND S. ZHENG, *On the Cahn–Hilliard equation*, Arch. Rational Mech. Anal. **96** (1986), no. 4, 339–357.
- [7] H. GARCKE, *On mathematical models for phase separation in elastically stressed solids*, Habilitation Thesis (2000), University of Bonn.
- [8] H. GARCKE, *On Cahn–Hilliard systems with elasticity*, Proc. Roy. Soc. Edinburgh Sect. A **133** (2003), 307–331.
- [9] ———, *On a Cahn–Hilliard model for phase separation with elastic misfit*, Inst. H. Poincaré Anal. Non Linéaire **22** (2005), 165–185.
- [10] M. E. GURTIN, *Generalized Ginzburg–Landau and Cahn–Hilliard equations based on a microforce balance*, Physica D **92** (1996), 178–192.
- [11] O. A. LADYZHENSKAYA AND N. N. URAL'TSEVA, *Linear and Quasilinear Equations of Elliptic type*, Nauka, Moscow, 1973. (in Russian)
- [12] J. L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, 1969.
- [13] J. L. LIONS AND E. MAGENES, *Non-Homogeneous Boundary Value Problems and Applications*, vol. I, II, Springer, Berlin, 1972.
- [14] A. MIRANVILLE, *Some generalizations of the Cahn–Hilliard equation*, Asymptotic Anal. **22** (2000), 235–259.
- [15] ———, *Long-time behavior of some models of Cahn–Hilliard equations in deformable continua*, Nonlinear Analysis: Real World Applications **2** (2001), 273–304.

- [16] ———, *Consistent models of Cahn–Hilliard–Gurtin equations with Neumann boundary conditions*, *Physica D* **158** (2001), no. 1–4, 233–257.
- [17] ———, *Generalized Cahn–Hilliard equations based on a microforce balance*, *J. Appl. Math.* **4** (2003), 165–185.
- [18] A. MIRANVILLE AND G. SCHIMPERNA, *Nonisothermal phase separation based on a microforce balance*, *Discrete Contin. Dynam. Systems Ser. B* **5** (2005), 753–768.
- [19] ———, *Generalized Cahn–Hilliard equations for multicomponent alloys*, Preprint (2005).
- [20] J. NEČAS, *Les Methodés Directes en Théorie des Equations Elliptiques*, Mason, Paris, 1967.
- [21] I. PAWŁOW, W. M. ZAJĄCZKOWSKI, *Classical solvability of 1-D Cahn–Hilliard equation coupled with elasticity*, *Math. Methods Appl. Sci.* **29** (2006), 853–876.
- [22] ———, *Strong solvability of 3-D Cahn–Hilliard system in elastic solids*, *Math. Methods Appl. Sci.*, in press.
- [23] ———, *Regular weak solutions to 3-D Cahn–Hilliard in elastic solids*, *Discrete Contin. Dynam. Systems, Supplement Volume* (2007), (in press).
- [24] I. PAWŁOW AND A. ŻOCHOWSKI, *Existence and uniqueness for a three-dimensional thermoelastic system*, *Dissertationes Math.* **406** (2002), 46 pp.
- [25] J. SIMON, *Compact sets in the space $L^p(0, T; B)$* , *Ann. Mat. Pura Appl.* **146** (1987), 65–97.

Manuscript received November 10, 2006

IRENA PAWŁOW
 Systems Research Institute
 Polish Academy of Sciences
 Newelska 6
 01-447 Warsaw, POLAND
 and
 Institute of Mathematics and Cryptology
 Cybernetics Faculty
 Military University of Technology
 Kaliskiego 2
 00-908 Warsaw, POLAND
E-mail address: pawlow@ibspan.waw.pl

WOJCIECH M. ZAJĄCZKOWSKI
 Institute of Mathematics
 Polish Academy of Sciences
 Śniadeckich 8
 00-950 Warsaw, POLAND
 and
 Institute of Mathematics and Cryptology
 Cybernetics Faculty
 Military University of Technology
 Kaliskiego 2
 00-908 Warsaw, POLAND
E-mail address: wz@impan.gov.pl