

Tunisian Journal of Mathematics

an international publication organized by the Tunisian Mathematical Society

On the mod-2 cohomology of $SL_3(\mathbb{Z}[\frac{1}{2}, i])$

Hans-Werner Henn

2019 vol. 1 no. 4



On the mod-2 cohomology of $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$

Hans-Werner Henn

Let $\Gamma = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$, let X be any mod-2 acyclic Γ -CW complex on which Γ acts with finite stabilizers and let X_s be the 2-singular locus of X . We calculate the mod-2 cohomology of the Borel construction of X_s with respect to the action of Γ . This cohomology coincides with the mod-2 cohomology of Γ in cohomological degrees bigger than 8 and the result is compatible with a conjecture of Quillen which predicts the structure of the cohomology ring $H^*(\Gamma; \mathbb{F}_2)$.

1. Introduction

The main motivation for this paper comes from a conjecture of Quillen [1971, Conjecture 14.7] which concerns the structure of the mod- p cohomology ring of the group $\mathrm{GL}_n(\Lambda)$ of invertible matrices of rank n with coefficients in a ring Λ of S -integers in a number field; the assumption on Λ is that p is invertible in Λ and Λ contains a primitive p -th root of unity. The conjecture stipulates that under these assumptions $H^*(\mathrm{GL}_n(\Lambda); \mathbb{Z}/p)$ is a free module over the polynomial algebra $\mathbb{Z}/p[c_1, \dots, c_n]$ where the c_i are the mod- p Chern classes associated to an embedding of Λ into the complex numbers. In the sequel we will denote this conjecture by $C(n, \Lambda, p)$.

For $p = 2$ the simplest ring for which the assumptions of Quillen's conjecture hold is the ring $\mathbb{Z}[\frac{1}{2}]$. Let $\mathbb{Z}[\frac{1}{2}, i]$ be the ring obtained from the Gaussian integers $\mathbb{Z}[i]$ by inverting 2.

Conjecture $C(n, \mathbb{Z}[\frac{1}{2}], 2)$ is trivially true for $n = 1$ and known to be true for $n = 2$ by [Mitchell 1992] and $n = 3$ by [Henn 1999]; it is known to be false for $n = 32$ by [Dwyer 1998] and even for $n \geq 14$ (Henn and Lannes, unpublished). The positive results have been established by direct calculation and while a direct calculation is perhaps still doable for $n = 4$, it would be extremely involved. For larger n a complete calculation does not look realistic. In fact, the negative results have been obtained by very indirect methods which depend on étale approximations for the homotopy type of the 2-completion of $\mathrm{BGL}_n(\mathbb{Z}[\frac{1}{2}])$. These étale approximations can also be used to show that if $C(2n, \mathbb{Z}[\frac{1}{2}], 2)$ holds then $C(n, \mathbb{Z}[\frac{1}{2}, i], 2)$ holds

MSC2010: primary 20G10; secondary 55R40.

Keywords: cohomology of special and general linear groups, Quillen conjecture.

as well (Henn and Lannes, unpublished). This gives particular motivation to study conjecture $C(n, \mathbb{Z}[\frac{1}{2}, i], 2)$.

We will show in [Theorem 5.1](#) that $C(n, \mathbb{Z}[\frac{1}{2}, i], 2)$ holds if and only if there is an isomorphism

$$H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \cong \mathbb{F}_2[c_1, \dots, c_n] \otimes E(e_1, e'_1, \dots, e_{2n-1}, e'_{2n-1})$$

where the classes c_i are the Chern classes of the tautological n -dimensional complex representation of $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$, E denotes an exterior algebra and the classes e_{2i-1}, e'_{2i-1} are of cohomological degree $2i - 1$ for $i = 1, \dots, n$. These exterior classes are closely related to Quillen's exterior classes in the mod-2 cohomology of $\mathrm{GL}_n(\mathbb{F}_p)$ if p is a prime such that $p \equiv 1 \pmod{4}$ (see [\(5-1\)](#) for more details).

Conjecture $C(n, \mathbb{Z}[\frac{1}{2}, i], 2)$ is again trivially true for $n = 1$ and has been verified by direct calculation for $n = 2$ in [\[Weiss 2006\]](#). Dwyer's method [\[1998\]](#) using étale approximations X_n for the homotopy type of the 2-completion of $\mathrm{BGL}_n(\mathbb{Z}[\frac{1}{2}])$ and comparing the set of homotopy classes of $[\mathrm{BP}, X_n]$ with that of $[\mathrm{BP}, \mathrm{BGL}_n(\mathbb{Z}[\frac{1}{2}])]$ for suitable cyclic groups P of order 2^n can be adapted to disprove $C(16, \mathbb{Z}[\frac{1}{2}, i], 2)$. However, we will not dwell on this in this paper.

This paper embarks on a study of conjecture $C(3, \mathbb{Z}[\frac{1}{2}, i], 2)$ which is more accessible than conjecture $C(4, \mathbb{Z}[\frac{1}{2}], 2)$. In order to calculate $H^*(\mathrm{GL}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ we first try to calculate $H^*(\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$. For this we propose the same strategy as the one which was used in the case of $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}])$ and which finally led to a verification of conjecture $C(3, \mathbb{Z}[\frac{1}{2}], 2)$. In a first step one uses a centralizer spectral sequence introduced in [\[Henn 1997\]](#) in order to calculate the mod-2 Borel cohomology $H_G^*(X_s; \mathbb{F}_2)$ where X is any mod-2 acyclic G -CW complex on which a given discrete group G acts with finite stabilizers and X_s is the 2-singular locus of X , i.e., the subcomplex consisting of all points for which the isotropy group of the action of G is of even order. For $G = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}])$ this step was carried out in [\[Henn 1997\]](#) and for $G = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ it is carried out in this paper. The precise form of X does not really matter in this step.

The second step involves a very laborious analysis of the relative mod-2 Borel cohomology $H_G^*(X, X_s; \mathbb{F}_2)$ and of the connecting homomorphism for the Borel cohomology of the pair (X, X_s) . In the case of $G = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}])$ this was carried out by hand in [\[Henn 1999\]](#). A by hand calculation looks forbidding in the case of $G = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ and this paper makes no attempt on such a calculation. However, we do make some comments on what is likely to be involved in such an attempt.

Here are the main results of this paper. In these results the elements b_2 and b_3 are of degree 4 and 6, respectively. They are given as Chern classes of the tautological 3-dimensional complex representation of $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$. The indices of the other elements give their cohomological degrees. These elements come from Quillen's exterior cohomology classes in the cohomology of $\mathrm{GL}_3(\mathbb{F}_p)$ for suitable primes p ,

for example $p = 5$ (see [Section 3.2](#) for more details). Furthermore Σ^n denotes n -fold suspension so that $\Sigma^4\mathbb{F}_2$ is a one dimensional \mathbb{F}_2 -vector space concentrated in degree 4.

Theorem 1.1. *Let $\Gamma = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ and let X be any mod-2 acyclic Γ -CW complex such that the isotropy group of each cell is finite. Then the centralizer spectral sequence of [\[Henn 1997\]](#) collapses at E_2 and gives a short exact sequence*

$$0 \rightarrow \Sigma^4\mathbb{F}_2 \oplus \Sigma^4\mathbb{F}_2 \oplus \Sigma^7\mathbb{F}_2 \rightarrow H_\Gamma^*(X_s; \mathbb{F}_2) \rightarrow \mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5) \rightarrow 0$$

in which the epimorphism is a map of graded commutative algebras with unit.

Next let

$$\psi : H^*(\Gamma; \mathbb{F}_2) = H_\Gamma^*(X; \mathbb{F}_2) \rightarrow H_\Gamma^*(X_s; \mathbb{F}_2) \rightarrow \mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5)$$

be the composition of the map induced by the inclusion $X_s \subset X$ and the epimorphism of [Theorem 1.1](#).

Theorem 1.2. *Let $\Gamma = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ and X be as in the previous theorem.*

- (a) *If $\mathrm{SD}_3(\mathbb{Z}[\frac{1}{2}, i])$ denotes the subgroup of diagonal matrices of Γ then the target of ψ can be identified with a subalgebra of $H^*(\mathrm{SD}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ in such a way that ψ is induced by the restriction homomorphism*

$$H^*(\Gamma; \mathbb{F}_2) \rightarrow H^*(\mathrm{SD}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2).$$

- (b) *The homomorphism ψ admits a multiplicative section*

$$\varphi : \mathbb{F}_2[c_2, c_3] \otimes E(e_3, e'_3, e_5, e'_5) \rightarrow H^*(\Gamma; \mathbb{F}_2)$$

that sends c_i to b_i for $i = 2, 3$ and sends e_i and e'_i respectively to d_i and d'_i for $i = 3, 5$.

- (c) *The homomorphism ψ is surjective in all degrees, an isomorphism in degrees $* > 8$ and its kernel is finite-dimensional in degrees $* \leq 8$.*

Remark 1.3. Conjecture $C(3, \mathbb{Z}[\frac{1}{2}, i], 2)$ would hold if the maps ψ and φ of part (b) of [Theorem 1.2](#) turned out to be isomorphisms (see [Proposition 5.5](#)).

The following result is an immediate consequence of [Theorem 1.2](#).

Corollary 1.4. *Let $\Gamma = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ and X be as in [Theorem 1.1](#). Then the following conditions are equivalent:*

- (a) *The restriction homomorphism $H^*(\Gamma; \mathbb{F}_2) \rightarrow H^*(\mathrm{SD}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ is injective and $H^*(\Gamma; \mathbb{F}_2)$ is isomorphic as a graded \mathbb{F}_2 -algebra to $\mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5)$.*

(b) *There is an isomorphism*

$$H_{\Gamma}^*(X, X_s; \mathbb{F}_2) \cong \Sigma^5 \mathbb{F}_2 \oplus \Sigma^5 \mathbb{F}_2 \oplus \Sigma^8 \mathbb{F}_2$$

and the connecting homomorphism $H_{\Gamma}^*(X_s; \mathbb{F}_2) \rightarrow H_{\Gamma}^{*+1}(X, X_s; \mathbb{F}_2)$ is surjective. \square

The paper is organized as follows. In [Section 2](#) we recall the centralizer spectral sequence and in [Section 3](#) we prove [Theorems 1.1 and 1.2](#). In [Section 4](#) we make some comments on step 2 of the program of a complete calculation of $H^*(\Gamma; \mathbb{F}_2)$. Finally in [Section 5](#) we establish [Theorem 5.1](#) and discuss the relation between [Theorem 1.2](#) and conjecture $C(3, \mathbb{Z}[\frac{1}{2}, i], 2)$.

2. The centralizer spectral sequence

We recall the centralizer spectral sequence introduced in [\[Henn 1997\]](#).

Let G be a discrete group and let p be a fixed prime. Let $\mathcal{A}(G)$ be the category whose objects are the elementary abelian p -subgroups E of G , i.e., subgroups which are isomorphic to $(\mathbb{Z}/p)^k$ for some integer k ; if E_1 and E_2 are elementary abelian p -subgroups of G , then the set of morphisms from E_1 to E_2 in $\mathcal{A}(G)$ consists precisely of those group homomorphisms $\alpha: E_1 \rightarrow E_2$ for which there exists an element $g \in G$ with $\alpha(e) = geg^{-1}$ for all $e \in E_1$. Let $\mathcal{A}_*(G)$ be the full subcategory of $\mathcal{A}(G)$ whose objects are the nontrivial elementary abelian p -subgroups.

For an elementary abelian p -subgroup E we denote its centralizer in G by $C_G(E)$. Then the assignment $E \mapsto H^*(C_G(E); \mathbb{F}_p)$ determines a functor from $\mathcal{A}_*(G)$ to the category \mathcal{E} of graded \mathbb{F}_p -vector spaces. The inverse limit functor is a left exact functor from the functor category $\mathcal{E}^{\mathcal{A}_*(G)}$ to \mathcal{E} . Its right derived functors are denoted by \lim^s . The p -rank $r_p(G)$ of a group G is defined as the supremum of all k such that G contains a subgroup isomorphic to $(\mathbb{Z}/p)^k$.

For a G -space X and a fixed prime p we denote by X_s the p -singular locus, i.e., the subspace of X consisting of points whose isotropy group contains an element of order p . Let EG be the total space of the universal principal G -bundle. The mod- p cohomology of the Borel construction $EG \times_G X$ of a G space X will be denoted $H_G^*(X; \mathbb{F}_p)$. The following result is a special case of part (a) of [Corollary 0.4 of \[Henn 1997\]](#).

Theorem 2.1. *Let G be a discrete group and assume there exists a finite-dimensional mod- p acyclic G -CW complex X such that the isotropy group of each cell is finite. Then there exists a cohomological second quadrant spectral sequence*

$$E_2^{s,t} = \lim_{\mathcal{A}_*(G)}^s H^t(C_G(E); \mathbb{F}_p) \Rightarrow H_G^{s+t}(X_s; \mathbb{F}_p)$$

with $E_2^{s,t} = 0$ if $s \geq r_p(G)$ and $t \geq 0$.

Remark 2.2. The edge homomorphism in this spectral sequence is a map of algebras

$$H_G^*(X_s; \mathbb{F}_p) \rightarrow \lim_{\mathcal{A}_*(G)} H^*(C_G(E); \mathbb{F}_p),$$

which is given as follows.

Let X^E be the fixed points for the action of E on X . The G -action on X restricts to an action of the centralizer $C_G(E)$ on X^E and the G -equivariant maps

$$G \times_{C_G(E)} X^E \rightarrow X_s, \quad (g, x) \mapsto gx.$$

for $E \in \mathcal{A}_*(G)$ induce compatible maps in Borel cohomology

$$H_G^*(X_s; \mathbb{F}_2) \rightarrow H_G^*(G \times_{C_G(E)} X^E; \mathbb{F}_2) \cong H_{C_G(E)}^*(X^E; \mathbb{F}_2) \cong H^*(C_G(E); \mathbb{F}_2)$$

which assemble to give the map to the inverse limit. Here we have used that by classical Smith theory X^E is mod p -acyclic if X is mod- p acyclic and hence we get canonical isomorphisms $H_{C_G(E)}^*(X^E; \mathbb{F}_2) \cong H_{C_G(E)}^*(\ast; \mathbb{F}_2) \cong H^*(C_G(E); \mathbb{F}_2)$.

Furthermore the composition

$$H^*(G; \mathbb{F}_p) = H_G^*(X; \mathbb{F}_p) \rightarrow H_G^*(X_s; \mathbb{F}_p) \rightarrow H^*(C_G(E); \mathbb{F}_p) \quad (2-1)$$

is induced by the inclusions $C_G(E) \rightarrow G$ as E varies through $\mathcal{A}_*(G)$.

In [Henn 1997] we have used this spectral sequence in the case $p = 2$ and $G = \mathrm{SL}_3(\mathbb{Z})$. Here we will use it in the case $p = 2$ and $G = \mathrm{SL}(3, \mathbb{Z}[\frac{1}{2}, i])$. In both cases we have $r_2(G) = 2$ and hence the spectral sequence collapses at E_2 and degenerates into a short exact sequence

$$0 \rightarrow \lim_{\mathcal{A}_*(G)}^1 H^*(C_G(E); \mathbb{F}_2) \rightarrow H_G^{*+1}(X_s; \mathbb{F}_2) \rightarrow \lim_{\mathcal{A}_*(G)} H^{*+1}(C_G(E); \mathbb{F}_2) \rightarrow 0. \quad (2-2)$$

3. The centralizer spectral sequence for $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$

3.1. The Quillen category. Let K be any number field, let \mathcal{O}_K be its ring of integers and consider the ring of S -integers $\mathcal{O}_K[\frac{1}{2}]$. Then, up to equivalence, the Quillen category of $G := \mathrm{SL}_3(\mathcal{O}_K[\frac{1}{2}])$ for the prime 2 is independent of K . In fact, because 2 is invertible every elementary abelian 2-subgroup is conjugate to a diagonal subgroup, and hence $\mathcal{A}_*(G)$ has a skeleton, say \mathcal{A} , with exactly two objects, say E_1 and E_2 of rank 1 and 2, respectively. We take E_1 to be the subgroup generated by the diagonal matrix whose first two diagonal entries are -1 and whose third diagonal entry is 1, and E_2 to be the subgroup of all diagonal matrices with diagonal entries 1 or -1 and determinant 1.

The automorphism group of E_1 is trivial, of course, while $\mathrm{Aut}_{\mathcal{A}}(E_2)$ is isomorphic to the group of all abstract automorphisms of E_2 which we can identify

with \mathfrak{S}_3 , the symmetric group on three elements. There are three morphisms from E_1 to E_2 and $\text{Aut}_{\mathcal{A}}(E_2)$ acts transitively on them.

3.2. The centralizers and their cohomology. For centralizers in $H := \text{GL}_3(\mathcal{O}_K[\frac{1}{2}])$ we find $C_H(E_1) = \text{GL}_2(\mathcal{O}_K[\frac{1}{2}]) \times \text{GL}_1(\mathcal{O}_K[\frac{1}{2}])$ and $C_H(E_2) = D_3(\mathcal{O}_K[\frac{1}{2}])$ if $D_n(\mathcal{O}_K[\frac{1}{2}])$ denotes the subgroup of diagonal matrices in $\text{GL}_n(\mathcal{O}_K[\frac{1}{2}])$. This implies

$$\begin{aligned} C_G(E_1) &\cong \text{GL}_2(\mathcal{O}_K[\frac{1}{2}]), \\ C_G(E_2) &= \text{SD}_3(\mathcal{O}_K[\frac{1}{2}]) \cong D_2(\mathcal{O}_K[\frac{1}{2}]) \cong \mathcal{O}_K[\frac{1}{2}]^\times \times \mathcal{O}_K[\frac{1}{2}]^\times, \end{aligned}$$

where as before $\text{SD}_3(\mathcal{O}_K[\frac{1}{2}])$ denotes special diagonal matrices with coefficients in $\mathcal{O}_K[\frac{1}{2}]$.

From now on we specialize to the case $K = \mathbb{Q}[i]$ where we have $\mathcal{O}_K[\frac{1}{2}] = \mathbb{Z}[\frac{1}{2}, i]$. In this case the cohomology of the centralizers is explicitly known. In the sequel we abbreviate $\text{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ by Γ .

3.2.1. The cohomology of $C_\Gamma(E_2)$. There is an isomorphism of groups

$$\mathbb{Z}/4 \times \mathbb{Z} \cong \mathbb{Z}[\frac{1}{2}, i]^\times, \quad (n, m) \mapsto i^n(1+i)^m$$

and therefore we get an isomorphism

$$H^*(C_\Gamma(E_2); \mathbb{F}_2) \cong H^*(\mathbb{Z}[\frac{1}{2}, i]^\times \times \mathbb{Z}[\frac{1}{2}, i]^\times; \mathbb{F}_2) \cong \mathbb{F}_2[y_1, y_2] \otimes E(x_1, x'_1, x_2, x'_2) \quad (3-1)$$

with y_1 and y_2 in degree 2 and the other generators in degree 1. We agree to choose the generators so that y_1, x_1 and x'_1 come from the first factor with x_1 and x'_1 being the dual basis to the basis of

$$H_1(\mathbb{Z}[\frac{1}{2}, i]^\times; \mathbb{F}_2) \cong \mathbb{Z}[\frac{1}{2}, i]^\times / (\mathbb{Z}[\frac{1}{2}, i]^\times)^2 \cong \mathbb{Z}/2 \times \mathbb{Z}/2$$

given by the image of i and $(1+i)$ in the mod-2 reduction of the abelian group $\mathbb{Z}[\frac{1}{2}, i]^\times$ and y_1 coming from $H^2(\mathbb{Z}/4; \mathbb{F}_2)$; likewise with y_2, x_2 and x'_2 coming from the second factor.

3.2.2. The cohomology of $C_\Gamma(E_1)$. This cohomology has been calculated in [Weiss 2006]. In fact, from Theorem 1 of [Weiss 2006] we know

$$H^*(C_\Gamma(E_1); \mathbb{F}_2) \cong H^*(\text{GL}_2(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \cong \mathbb{F}_2[c_1, c_2] \otimes E(e_1, e'_1, e_3, e'_3). \quad (3-2)$$

Here we give a short summary of this calculation. The classes e_1, e'_1, e_3 and e'_3 are pulled back from Quillen's exterior classes q_1 and q_3 [1972] in

$$H^*(\text{GL}_2(\mathbb{F}_5); \mathbb{F}_2) \cong \mathbb{F}_2[c_1, c_2] \otimes E(q_1, q_3) \quad (3-3)$$

via two ring homomorphisms

$$\pi : \mathbb{Z}[\frac{1}{2}, i] \rightarrow \mathbb{F}_5, \quad \pi' : \mathbb{Z}[\frac{1}{2}, i] \rightarrow \mathbb{F}_5. \quad (3-4)$$

We choose π such that i is sent to 3 and π' such that i is sent to 2.

Now consider the two commutative diagrams (with horizontal arrows induced by inclusion and vertical arrows induced by π and, respectively, π')

$$\begin{array}{ccc} D_2(\mathbb{Z}[\frac{1}{2}]) & \longrightarrow & GL_2(\mathbb{Z}[\frac{1}{2}]) \\ \downarrow & & \downarrow \\ D_2(\mathbb{F}_5) & \longrightarrow & GL_2(\mathbb{F}_5). \end{array} \quad (3-5)$$

By abuse of notation we can write

$$H^*(D_2(\mathbb{F}_5); \mathbb{F}_2) \cong H^*(\mathbb{F}_5^\times \times \mathbb{F}_5^\times; \mathbb{F}_2) \cong \mathbb{F}_2[y_1, y_2] \otimes E(x_1, x_2) \quad (3-6)$$

with $y_1 \in H^2(\mathbb{F}_5^\times; \mathbb{F}_2)$ and $x_1 \in H^2(\mathbb{F}_5^\times; \mathbb{F}_2)$ coming from the first factor and likewise with y_2 and x_2 coming from the second factor. Then π and π' induce two homomorphisms

$$\pi^*, \pi'^* : H^*(D_2(\mathbb{F}_5); \mathbb{F}_2) \rightarrow H^*(D_2(\mathbb{Z}[\frac{1}{2}]); \mathbb{F}_2)$$

which in terms of the isomorphisms (3-6) and (3-1) are explicitly given by

$$\pi^*(y_i) = y_i = \pi'^*(y_i), \quad \pi^*(x_i) = x_i, \quad \pi'^*(x_i) = x_i + x'_i \quad \text{for } i = 1, 2. \quad (3-7)$$

The cohomology of $GL_2(\mathbb{F}_5)$ is detected by restriction to the cohomology of diagonal matrices and restriction is given explicitly as follows:

$$c_1 \mapsto y_1 + y_2, \quad c_2 \mapsto y_1 y_2, \quad q_1 \mapsto x_1 + x_2, \quad q_3 \mapsto y_1 x_2 + y_2 x_1. \quad (3-8)$$

Then e_1, e'_1, e_3, e'_3 are defined via

$$e_1 = \pi^*(q_1), \quad e_3 = \pi^*(q_3), \quad e'_1 = \pi'^*(q_1), \quad e'_3 = \pi'^*(q_3). \quad (3-9)$$

If c_1 and c_2 are the Chern classes of the tautological 2-dimensional complex representation of $GL_2(\mathbb{Z}[\frac{1}{2}], i)$, then the restriction homomorphism which sends $H^*(GL_2(\mathbb{Z}[\frac{1}{2}], i); \mathbb{F}_2)$ to the cohomology of the subgroup of diagonal matrices is injective and by using (3-5) and (3-8) we see that it is explicitly given by

$$\begin{array}{ll} c_1 \mapsto y_1 + y_2, & c_2 \mapsto y_1 y_2, \\ e_1 \mapsto x_1 + x_2, & e_3 \mapsto y_1 x_2 + y_2 x_1, \\ e'_1 \mapsto x_1 + x'_1 + x_2 + x'_2, & e'_3 \mapsto y_1(x_2 + x'_2) + y_2(x_1 + x'_1). \end{array} \quad (3-10)$$

3.2.3. Functoriality. We note that together with the isomorphisms (3-1) and (3-2) the restriction (3-10) also describes the map

$$\alpha_* : H^*(C_\Gamma(E_1); \mathbb{F}_2) \rightarrow H^*(C_\Gamma(E_2); \mathbb{F}_2)$$

induced from the standard inclusion of E_1 into E_2 .

To finish the description of $H^*(C_\Gamma(-); \mathbb{F}_2)$ as a functor on \mathcal{A} it remains to describe the action of the symmetric group $\text{Aut}_{\mathcal{A}}(E_2) \cong \mathfrak{S}_3$ of rank 3 on

$$H^*(C_\Gamma(E_2); \mathbb{F}_2) \cong \mathbb{F}_2[y_1, y_2] \otimes \Lambda(x_1, x'_1, x_2, x'_2).$$

Because of the multiplicative structure we need it only on the generators.

If $\tau \in \text{Aut}_{\mathcal{A}}(E_2)$ corresponds to permuting the factors in $C_\Gamma(E_2) \cong \text{GL}_1(\mathbb{Z}[\frac{1}{2}, i]) \times \text{GL}_1(\mathbb{Z}[\frac{1}{2}, i])$ then

$$\begin{aligned} \tau_*(y_1) &= y_2, & \tau_*(x_1) &= x_2, & \tau_*(x'_1) &= x'_2, \\ \tau_*(y_2) &= y_1, & \tau_*(x_2) &= x_1, & \tau_*(x'_2) &= x'_1, \end{aligned} \quad (3-11)$$

and if $\sigma \in \text{Aut}_{\mathcal{A}}(E_2)$ corresponds to the cyclic permutation of the diagonal entries (in suitable order) then

$$\begin{aligned} \sigma_*(y_1) &= y_2, & \sigma_*(x_1) &= x_2, & \sigma_*(x'_1) &= x'_2, \\ \sigma_*(y_2) &= y_1 + y_2, & \sigma_*(x_2) &= x_1 + x_2, & \sigma_*(x'_2) &= x'_1 + x'_2. \end{aligned} \quad (3-12)$$

3.3. Calculating the limit and its derived functors. In Proposition 4.3 of [Henn 1997] we showed that for any functor F from \mathcal{A} to $\mathbb{Z}_{(2)}$ -modules there is an exact sequence

$$0 \rightarrow \lim_{\mathcal{A}} F \rightarrow F(E_1) \xrightarrow{\varphi} \text{Hom}_{\mathbb{Z}[\mathfrak{S}_3]}(\text{St}_{\mathbb{Z}}, F(E_2)) \rightarrow \lim_{\mathcal{A}}^1 F \rightarrow 0 \quad (3-13)$$

where $\text{St}_{\mathbb{Z}}$ is the $\mathbb{Z}[\mathfrak{S}_3]$ -module given by the kernel of the augmentation map $\mathbb{Z}[\mathfrak{S}_3/\mathfrak{S}_2] \rightarrow \mathbb{Z}$, and if a and b are chosen to give an integral basis of $\text{St}_{\mathbb{Z}}$ on which τ and σ act via

$$\begin{aligned} \tau_*(a) &= b, & \tau_*(b) &= a, \\ \sigma_*(a) &= -b, & \sigma_*(b) &= a - b, \end{aligned} \quad (3-14)$$

then $\varphi(x)(a) = \alpha_*(x) - (\sigma_*)^2 \alpha_*(x)$ and $\varphi(x)(b) = \alpha_*(x) - \sigma_* \alpha_*(x)$ if $x \in F(E_1)$.

Because in our case the functor takes values in \mathbb{F}_2 -vector spaces we can replace $\text{Hom}_{\mathbb{Z}[\mathfrak{S}_3]}$ by $\text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}$ and $\text{St}_{\mathbb{Z}}$ by its mod-2 reduction. The following elementary lemma is needed in the analysis of the third term in the exact sequence (3-13).

Lemma 3.1. (a) *Let St be the $\mathbb{F}_2[\mathfrak{S}_3]$ -module given as the kernel of the augmentation $\mathbb{F}_2[\mathfrak{S}_3/\mathfrak{S}_2] \rightarrow \mathbb{F}_2$. The tensor product $\text{St} \otimes \text{St}$ decomposes as $\mathbb{F}_2[\mathfrak{S}_3]$ -module canonically as*

$$\text{St} \otimes \text{St} \cong \mathbb{F}_2[\mathfrak{S}_3/A_3] \oplus \text{St}$$

where A_3 denotes the alternating group on three letters. In fact, the decomposition is given by

$$\text{St} \otimes \text{St} \cong \text{Im}(\text{id} + \sigma_* + \sigma_*^2) \oplus \text{Ker}(\text{id} + \sigma_* + \sigma_*^2)$$

and the first summand is isomorphic to $\mathbb{F}_2[\mathfrak{S}_3/A_3]$ while the second summand is isomorphic to St .

(b) The tensor product $\mathbb{F}_2[\mathfrak{S}_3/A_3] \otimes St$ is isomorphic to $St \oplus St$.

Proof.

(a) It is well-known that St is a projective $\mathbb{F}_2[\mathfrak{S}_3]$ -module, hence $St \otimes St$ is also projective. It is also well-known that every projective indecomposable $\mathbb{F}_2[\mathfrak{S}_3]$ -module is isomorphic to either St or $\mathbb{F}_2[\mathfrak{S}_3/A_3]$. The two modules can be distinguished by the fact that $e := id + \sigma_* + \sigma_*$ acts trivially on St and as the identity on $\mathbb{F}_2[\mathfrak{S}_3/A_3]$.

Furthermore e is a central idempotent in $\mathbb{F}_2[\mathfrak{S}_3]$ and hence each $\mathbb{F}_2[\mathfrak{S}_3]$ -module M decomposes as direct sum of $\mathbb{F}_2[\mathfrak{S}_3]$ -modules

$$M \cong \text{Im}(e : M \rightarrow M) \oplus \text{Ker}(e : M \rightarrow M).$$

An easy calculation shows that in the case of $St \otimes St$ both submodules are nontrivial and this together with the fact these submodules must be projective proves the claim.

(b) Again each of the factors in the tensor product is a projective $\mathbb{F}_2[\mathfrak{S}_3]$ -module, hence the tensor product is a projective $\mathbb{F}_2[\mathfrak{S}_3]$ -module. Because σ acts as the identity on $\mathbb{F}_2[\mathfrak{S}_3/A_3]$ we see that the idempotent e acts trivially on the tensor product and this forces the tensor product to be isomorphic to $St \oplus St$. \square

Lemma 3.2. *The Poincaré series χ_2 of $\text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(St, \mathbb{F}_2[y_1, y_2] \otimes E(x_1, x'_1, x_2, x'_2))$ is given by*

$$\chi_2 = \frac{2t^2(1+3t^2+3t^4+t^6) + 2t(1+2t^2+2t^4+2t^6+t^8)}{(1-t^4)(1-t^6)}.$$

Proof. The isomorphism of (3-1) is an isomorphism of $\mathbb{F}_2[\mathfrak{S}_3]$ -modules where the action of \mathfrak{S}_3 is given by equations (3-11) and (3-12). In particular we see that $H^1(\text{GL}_1(\mathbb{Z}[\frac{1}{2}, i]) \times \text{GL}_1(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ is isomorphic to $St \oplus St$ generated by x_1, x'_1, x_2, x'_2 . The exterior powers of H^1 are given as

$$E^k(x_1, x_2, x'_1, x'_2) \cong E^k(St \oplus St) \cong \bigoplus_{j=0}^k E^j St \otimes E^{k-j} St$$

and, because $E^k(St)$ is isomorphic to $\Sigma^k \mathbb{F}_2$ if $k = 0, 2$, isomorphic to ΣSt if $k = 1$, and trivially otherwise, we obtain

$$E^k(x_1, x_2, x'_1, x'_2) \cong \begin{cases} \Sigma^k \mathbb{F}_2 & \text{if } k = 0, 4, \\ \Sigma^k(St \oplus St) & \text{if } k = 1, 3, \\ \Sigma^2 \mathbb{F}_2 \oplus \Sigma^2(St \otimes St) \oplus \Sigma^2 \mathbb{F}_2 & \text{if } k = 2, \\ 0 & \text{if } k \neq 0, 1, 2, 3, 4, \end{cases}$$

where \mathbb{F}_2 denotes the trivial $\mathbb{F}_2[\mathfrak{S}_3]$ -module whose additive structure is that of \mathbb{F}_2 .

Therefore the Poincaré series χ_2 of $\text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St}, H^*(C_\Gamma(E_2); \mathbb{F}_2))$ decomposes according to the decomposition of $\Lambda(x_1, x'_2, x'_1, x'_2)$ as the sum

$$\chi_2 := (1 + 2t^2 + t^4)\chi_{2,0} + t^2\chi_{2,1} + 2(t + t^3)\chi_{2,2} \quad (3-15)$$

where here we denote by $\chi_{2,0}$ the Poincaré series of $\text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St}, \mathbb{F}_2[y_1, y_2])$, by $\chi_{2,1}$ the Poincaré series of $\text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St}, \text{St} \otimes \text{St} \otimes \mathbb{F}_2[y_1, y_2])$ and by $\chi_{2,2}$ that of $\text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St}, \text{St} \otimes \mathbb{F}_2[y_1, y_2])$.

It is well-known (and elementary to verify) that there is an isomorphism of $\mathbb{F}_2[\mathfrak{S}_3]$ -modules $\text{St} \oplus \text{St} \otimes \mathbb{F}_2[\mathfrak{S}_3/A_3] \cong \mathbb{F}_2[\mathfrak{S}_3]$ and therefore an isomorphism

$$\begin{aligned} \mathbb{F}_2[y_1, y_2] &\cong \text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St} \oplus \text{St} \otimes \mathbb{F}_2[\mathfrak{S}_3/A_3], \mathbb{F}_2[y_1, y_2]) \\ &\cong \text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St}, \mathbb{F}_2[y_1, y_2])^{\oplus 2} \oplus \mathbb{F}_2[y_1, y_2]^{A_3}. \end{aligned}$$

Together with the elementary fact that the A_3 -invariants $\mathbb{F}_2[y_1, y_2]^{A_3}$ form a free module over $\mathbb{F}_2[y_1, y_2]^{\mathfrak{S}_3} \cong \mathbb{F}_2[c_2, c_3]$ on the two generators 1 and $y_1^3 + y_1y_2^2 + y_2^3$ of degree 0 and 6, respectively, this implies

$$2\chi_{2,0} + \frac{1+t^6}{(1-t^4)(1-t^6)} = \frac{1}{(1-t^2)^2}$$

and hence

$$\chi_{2,0} = \frac{t^2}{(1-t^2)(1-t^6)}. \quad (3-16)$$

It is elementary to check that St and $\mathbb{F}_2[\mathfrak{S}_3/A_3]$ are both self-dual $\mathbb{F}_2[\mathfrak{S}_3]$ -modules and hence [Lemma 3.1](#) gives

$$\text{St} \otimes \text{St}^* \cong \mathbb{F}_2[\mathfrak{S}_3/A_3] \oplus \text{St}$$

and

$$\begin{aligned} \text{St} \otimes \text{St}^* \otimes \text{St}^* &\cong (\mathbb{F}_2[\mathfrak{S}_3/A_3] \oplus \text{St}) \otimes \text{St}^* \\ &\cong (\mathbb{F}_2[\mathfrak{S}_3/A_3] \otimes \text{St}) \oplus (\text{St} \otimes \text{St}) \\ &\cong \mathbb{F}_2[\mathfrak{S}_3/A_3] \oplus \text{St} \oplus \text{St} \oplus \text{St}. \end{aligned}$$

Therefore, if $\chi_{\mathbb{F}_2[y_1, y_2]^{A_3}}$ denotes the Poincaré series of the A_3 -invariants then

$$\begin{aligned} \chi_{2,1} &= \chi_{\mathbb{F}_2[y_1, y_2]^{A_3}} + 3\chi_{2,0} \\ &= \frac{1+t^6}{(1-t^4)(1-t^6)} + \frac{3t^2}{(1-t^2)(1-t^6)} = \frac{1+3t^2+3t^4+t^6}{(1-t^4)(1-t^6)}, \end{aligned} \quad (3-17)$$

$$\begin{aligned} \chi_{2,2} &= \chi_{2,0} + \chi_{\mathbb{F}_2[y_1, y_2]^{A_3}} \\ &= \frac{t^2}{(1-t^2)(1-t^6)} + \frac{1+t^6}{(1-t^4)(1-t^6)} = \frac{1+t^2+t^4+t^6}{(1-t^4)(1-t^6)}. \end{aligned} \quad (3-18)$$

Finally (3-15), (3-16), (3-17) and (3-18) give

$$\begin{aligned}\chi_2 &= \frac{(1+2t^2+t^4)t^2(1+t^2)+t^2(1+3t^2+3t^4+t^6)+2(t+t^3)(1+t^2+t^4+t^6)}{(1-t^4)(1-t^6)} \\ &= \frac{2t^2(1+3t^2+3t^4+t^6)+2t(1+2t^2+2t^4+2t^6+t^8)}{(1-t^4)(1-t^6)},\end{aligned}$$

and this finishes the proof. \square

Theorem 1.1 is now an immediate consequence of **Theorem 2.1** and the following result.

Proposition 3.3. *Let $p = 2$ and $\Gamma = SL_3(\mathbb{Z}[\frac{1}{2}, i])$.*

(a) *There is an isomorphism of graded \mathbb{F}_2 -algebras*

$$\lim_{\mathcal{A}} H^*(C_\Gamma(E); \mathbb{F}_2) \cong \mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5).$$

Furthermore, if we identify this limit with a subalgebra of $H^(C_\Gamma(E_1); \mathbb{F}_2) \cong \mathbb{F}_2[c_1, c_2] \otimes E(e_1, e'_1, e_3, e'_3)$ then*

$$\begin{aligned}b_2 &= c_1^2 + c_2, & b_3 &= c_1 c_2, \\ d_3 &= e_3, & d_5 &= c_1 e_3 + c_2 e_1, \\ d'_3 &= e'_3, & d'_5 &= c_1 e'_3 + c_2 e'_1.\end{aligned}$$

(b) *There is an isomorphism of graded \mathbb{F}_2 -vector spaces*

$$\lim_{\mathcal{A}}^1 H^*(C_\Gamma(E); \mathbb{F}_2) \cong \Sigma^3 \mathbb{F}_2 \oplus \Sigma^3 \mathbb{F}_2 \oplus \Sigma^6 \mathbb{F}_2.$$

(c) *For any $s > 1$*

$$\lim_{\mathcal{A}}^s H^*(C_\Gamma(E); \mathbb{F}_2) = 0.$$

Proof. (a) It is easy to check that the subalgebra of $\mathbb{F}_2[c_1, c_2] \otimes E(e_1, e'_1, e_3, e'_3)$ generated by the elements $c_1^2 + c_2$, $c_1 c_2$, e_3 , e'_3 , $c_1 e_3 + c_2 e_1$, and $c_1 e'_3 + c_2 e'_1$ is isomorphic to the tensor product of a polynomial algebra on two generators b_2 and b_3 of degrees 4 and 6 and an exterior algebra on 4 generators d_3 , d'_3 , d_5 and d'_5 of degrees 3, 3, 5 and 5. In fact, it is clear that $c_1^2 + c_2$ and $c_1 c_2$ are algebraically independent and the elements e_3 , e'_3 , $c_1 e_3 + c_2 e_1$, and $c_1 e'_3 + c_2 e'_1$ are exterior classes; their product is given as $c_2^2 e_3 e'_3 e_1 e'_1 \neq 0$, and this implies easily that the exterior monomials in these elements are linearly independent over the polynomial algebra generated by $c_1^2 + c_2$ and $c_1 c_2$. From now on we identify b_2 , b_3 , d_3 , d'_3 , d_5 and d'_5 with $c_1^2 + c_2$, $c_1 c_2$, e_3 , e'_3 , $c_1 e_3 + c_2 e_1$ and $c_1 e'_3 + c_2 e'_1$.

Now we use the exact sequence (3-13) and the description of φ to determine the inverse limit. Because α_* is injective, we see that if we identify $H^*(C_\Gamma(E_1); \mathbb{F}_2)$ with its image in $H^*(C_\Gamma(E_2); \mathbb{F}_2)$ then the inverse limit can be identified with the intersection of the image of α_* with the invariants in $\mathbb{F}_2[y_1, y_2] \otimes E(x_1, x'_1, x_2, x'_2)$

with respect to the action of the cyclic group of order 3 of $\text{Aut}_{\mathcal{A}}(E_2) \cong \mathfrak{S}_3$ generated by σ . This action has been described in (3-12) and with these formulas it is straightforward to check that the elements

$$\begin{aligned}
 b_2 &= y_1^2 + y_1 y_2 + y_2^2, \\
 b_3 &= y_1 y_2 (y_1 + y_2), \\
 d_3 &= y_1 x_2 + y_2 x_1, \\
 d_5 &= (y_1 + y_2)(y_1 x_2 + y_2 x_1) + y_1 y_2 (x_1 + x_2) = y_1^2 x_2 + y_2^2 x_1, \\
 d'_3 &= y_1 (x_2 + x'_2) + y_2 (x_1 + x'_1), \\
 d'_5 &= (y_1 + y_2)(y_1 (x_2 + x'_2) + y_2 (x_1 + x'_1)) + y_1 y_2 (x_1 + x'_1 + x_2 + x'_2) \\
 &= y_1^2 (x_2 + x'_2) + y_2^2 (x_1 + x'_1)
 \end{aligned} \tag{3-19}$$

all belong to the inverse limit.

Now consider the Poincaré series

$$\begin{aligned}
 \chi_0 &:= \sum_{n \geq 0} \dim_{\mathbb{F}_2} (\mathbb{F}_2[b_2, b_3] \otimes E(e_3, e'_3, e_5, e'_5)^n) t^n = \frac{(1+t^3)^2(1+t^5)^2}{(1-t^4)(1-t^6)}, \\
 \chi_1 &:= \sum_{n \geq 0} \dim_{\mathbb{F}_2} H^n(C_\Gamma(E_1); \mathbb{F}_2) t^n = \frac{(1+t)^2(1+t^3)^2}{(1-t^2)(1-t^4)}, \\
 \chi_2 &:= \frac{2t^2(1+3t^2+3t^4+t^6) + 2t(1+2t^2+2t^4+2t^6+t^8)}{(1-t^4)(1-t^6)}.
 \end{aligned}$$

Then we have the identity

$$\chi_0 + \chi_2 - \chi_1 = \frac{p}{(1-t^4)(1-t^6)}$$

with

$$\begin{aligned}
 p &= (1+t^3)^2(1+t^5)^2 + 2t^2(1+3t^2+3t^4+t^6) \\
 &\quad + 2t(1+2t^2+2t^4+2t^6+t^8) - (1+t)^2(1+t^3)^2(1+t^2+t^4) \\
 &= 2t^3 + t^6 - 2t^7 - 2t^9 - t^{10} - t^{12} + 2t^{13} + t^{16} \\
 &= (2t^3 + t^6)(1-t^4)(1-t^6)
 \end{aligned}$$

and therefore

$$\chi_0 + \chi_2 = \chi_1 + (2t^3 + t^6). \tag{3-20}$$

This, together with the fact that $\lim_{\mathcal{A}} H^*(C_\Gamma(E); \mathbb{F}_2)$ contains a subalgebra isomorphic to $\mathbb{F}_2[b_2, b_3] \otimes \Lambda(d_3, d'_3, d_5, d'_5)$, already implies that the sequence

$$\begin{aligned}
 0 \rightarrow \mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5) &\rightarrow H^*(C_\Gamma(E_1); \mathbb{F}_2) \\
 &\xrightarrow{\varphi} \text{Hom}_{\mathbb{F}_2[\mathfrak{S}_3]}(\text{St}, H^*(C_\Gamma(E_1); \mathbb{F}_2)) \rightarrow 0
 \end{aligned}$$

in which the left-hand arrow is given by inclusion is exact except possibly in dimensions 3 and 6.

In order to complete the proof of (a) it is now enough to verify that in degrees 3 and 6 the inverse limit is not bigger than $\mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5)$. We leave this straightforward verification to the reader.

Then (b) follows immediately from (a) together with (3-20) and the exact sequence (3-13), and (c) follows from Theorem 2.1 and the fact that $r_2(G) = 2$. \square

We can now give the proof of Theorem 1.2.

Proof.

(a) The exact sequence of Theorem 1.1 is obtained from the exact sequence (2-2) via Proposition 3.3. Therefore the epimorphism of Theorem 1.1 is the edge homomorphism of the centralizer spectral sequence. The result then follows from (2-1) by observing that we have identified the target of the edge homomorphism with the subalgebra $\mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5)$ of $H^*(C_\Gamma(E_1); \mathbb{F}_2)$ and by recalling that $C_\Gamma(E_1)$ is equal to the subgroup of special diagonal matrices $\mathrm{SD}_3(\mathbb{Z}[\frac{1}{2}, i])$.

(b) The two ring homomorphisms $\pi, \pi' : \mathbb{Z}[\frac{1}{2}, i] \rightarrow \mathbb{F}_5$ of (3-4) determine homomorphisms $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i]) \subset \mathrm{GL}_3(\mathbb{Z}[\frac{1}{2}, i]) \rightarrow \mathrm{GL}_3(\mathbb{F}_5)$. By [Quillen 1972] we have

$$H^* \mathrm{GL}_3(\mathbb{F}_5); \mathbb{F}_2 \cong \mathbb{F}_3[c_1, c_2, c_3] \otimes E(q_1, q_3, q_5).$$

We get a well-defined homomorphism of \mathbb{F}_2 -graded algebras

$$\varphi : \mathbb{F}_2[c_2, c_3] \otimes E(e_3, e'_3, e_5, e'_5) \rightarrow H^*(\Gamma; \mathbb{F}_2)$$

by sending c_i to the i -th Chern class of the tautological 3-dimensional representation of Γ and by declaring $\varphi(e_i) = \pi^*(q_i)$ and $\varphi(e'_i) = \pi'^*(q'_i)$ for $i = 3, 5$. The classes q_1, q_3 and q_5 are the symmetrizations of x_1, y_1x_2 and $y_1y_2x_3$, respectively, with respect to the natural action of \mathfrak{S}_3 on

$$H^*(\mathrm{GL}_3(\mathbb{F}_5); \mathbb{F}_2) \cong \mathbb{F}_2[y_1, y_2, y_3] \otimes E(x_1, x_2, x_3).$$

Compare (5-1) below.

Next we determine the composition $\psi\varphi$. The universal Chern classes c_i are the elementary symmetric polynomials in variables, say y_i , and the inclusion $\mathrm{GL}_2(\mathbb{C}) \subset \mathrm{SL}_3(\mathbb{C}) \subset \mathrm{GL}_3(\mathbb{C})$ imposes the relation $y_1 + y_2 + y_3 = 0$. This implies that the behavior of ψ on Chern classes is given by

$$c_1 \mapsto 0, \quad c_2 \mapsto c_1^2 + c_2 = y_1^2 + y_1y_2 + y_2^2 = b_2, \quad c_3 \mapsto c_1c_2 = y_1y_2(y_1 + y_2) = b_3.$$

In these equations we have identified $H^*(\mathrm{GL}_2(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$, as in the proof of Proposition 3.3, via restriction with a subalgebra of $\mathbb{F}_2[y_1, y_2] \otimes E(x_1, x'_1, x_3, x'_3)$.

In order to determine the composition $\psi\phi$ on the classes e_3, e'_3, e_5 and e'_5 we calculate at the level of \mathbb{F}_5 and use naturality with respect to the homomorphisms induced by π and π' , i.e., we consider the maps induced in cohomology by the following commutative diagram in which the horizontal maps are induced by inclusion and the vertical maps are induced by π and, respectively, π' :

$$\begin{array}{ccccc} \mathrm{GL}_2(\mathbb{Z}[\frac{1}{2}]) & \longrightarrow & \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}]) & \longrightarrow & \mathrm{GL}_3(\mathbb{Z}[\frac{1}{2}]) \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{GL}_2(\mathbb{F}_5) & \longrightarrow & \mathrm{SL}_3(\mathbb{F}_5) & \longrightarrow & \mathrm{GL}_3(\mathbb{F}_5). \end{array}$$

On the level of \mathbb{F}_5 the composition induces in cohomology a map

$$\mathbb{F}_3[c_1, c_2, c_3] \otimes E(q_1, q_3, q_5) \rightarrow \mathbb{F}_2[c_1, c_2] \otimes E(e_1, e_3) \subset \mathbb{F}_2[y_1, y_2] \otimes E(q_1, q_3)$$

which is easily determined from (5-1) below by imposing the relations $y_1 + y_2 + y_3 = 0$ and $x_1 + x_2 + x_3 = 0$ on the symmetrization of the classes y_1x_2 and $y_1y_2x_3$ with respect to the natural action of \mathfrak{S}_3 on the cohomology of diagonal matrices $H^*(D_3(\mathbb{F}_5); \mathbb{F}_2) \cong \mathbb{F}_2[y_1, y_2, y_3] \otimes E(x_1, x_2, x_3)$. Explicitly we get

$$\begin{aligned} c_1 &\mapsto 0, & c_2 &\mapsto y_1^2 + y_1y_2 + y_2^2, & c_3 &\mapsto y_1y_2(y_1 + y_2), \\ q_1 &\mapsto 0, & q_3 &\mapsto y_1x_2 + y_2x_1, & q_5 &\mapsto y_1^2x_2 + y_2^2x_1 \end{aligned}$$

and by using (3-7) and (3-19) we see that the composition $\psi\phi$ maps the elements e_3, e_5, e'_3 , and e'_5 as follows:

$$\begin{aligned} e_3 &\mapsto \pi^*(y_1x_2 + y_2x_1) = d_3, \\ e_5 &\mapsto \pi^*(y_1^2x_2 + y_2^2x_1) = d_5, \\ e'_3 &\mapsto \pi'^*(y_1x_2 + y_2x_1) = d'_3, \\ e'_5 &\mapsto \pi'^*(y_1^2x_2 + y_2^2x_1) = d'_5. \end{aligned}$$

Here we have identified the target of ψ with a subalgebra of $H^*(\mathrm{GL}_2(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ and the latter via restriction with a subalgebra of $\mathbb{F}_2[y_1, y_2] \otimes E(x_1, x'_1, x_3, x'_3)$.

(c) The space X can be taken to be the product of symmetric space

$$X_\infty := \mathrm{SL}_3(\mathbb{C}) / \mathrm{SU}(3)$$

and the Bruhat–Tits building X_2 for $\mathrm{SL}_3(\mathbb{Q}_2[i])$. Now $\mathrm{SL}_3(\mathbb{Q}_2[i]) \backslash X_2$ is a 2-simplex [Brown 1989] and the projection map $X \rightarrow X_2$ induces a map

$$\mathrm{SL}_3(\mathbb{Q}_2[i]) \backslash X \rightarrow \mathrm{SL}_3(\mathbb{Q}_2[i]) \backslash X_2$$

whose fibers have the homotopy type of a 6-dimensional $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ -invariant deformation retract (see Section 4). Therefore we get $H_G^n(X, X_s; \mathbb{F}_2) = 0$ if $n > 8$

and the inclusion $X_s \subset X$ induces an isomorphism $H_G^n(X; \mathbb{F}_2) \cong H_G^n(X_s; \mathbb{F}_2)$ if $n > 8$. Then part (c) simply follows from (a) except for the finiteness statement for the kernel for which we refer to (4-1) and (4-2) below. \square

4. Comments on step 2

The situation for $p = 2$ and $G = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ is analogous to the situation for $p = 2$ and $G = \mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}])$ for which step 2 was carried out in [Henn 1999] via a detailed study of the relative cohomology $H_G^*(X, X_s; \mathbb{F}_2)$ for X equal to the product of the symmetric space $X_\infty := \mathrm{SL}_3(\mathbb{R})/\mathrm{SO}(3)$ with the Bruhat–Tits building X_2 for $\mathrm{SL}_3(\mathbb{Q}_2)$; the spaces involved had a few hundred cells and the calculation was painful. In the case of $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ with X the product of $\mathrm{SL}_3(\mathbb{C})/\mathrm{SU}(3)$ with the Bruhat–Tits building for $\mathrm{SL}_3(\mathbb{Q}_2[i])$ the calculational complexity of the second step is much more involved and an explicit calculation by hand does not look feasible. However, in recent years there have been a lot of machine aided calculations of the cohomology of various arithmetic groups (for example [Dutour Sikirić et al. 2016; Bui et al. 2016]) and a machine aided calculation seems to be within reach.

The natural strategy for undertaking this second step is to follow the same path as in [Henn 1999]. The equivariant cohomology $H_\Gamma^*(X, X_s; \mathbb{F}_2)$ can be studied via the spectral sequence of the projection map

$$p : X = X_\infty \times X_2 \rightarrow X_2.$$

This gives a spectral sequence with

$$E_1^{p,q} \cong \bigoplus_{\sigma \in \Lambda_p} H_{\Gamma_\sigma}^q(X_\infty, X_{\infty,s}; \mathbb{F}_2) \Rightarrow H_\Gamma^{p+q}(X, X_s; \mathbb{F}_2). \quad (4-1)$$

Here Λ_p indexes the p -dimensional cells in the orbit space of X_2 with respect to the action of Γ . The orbit space is a 2-simplex, i.e., Λ_0 and Λ_1 contain 3 elements and Λ_2 is a singleton. Furthermore Γ_σ is the isotropy group of a chosen representative in X_2 of the cell σ in the quotient space. For fixed p all p -dimensional cells have isomorphic isotropy groups because the Γ -action on the Bruhat–Tits building is the restriction of a natural action of $\mathrm{GL}_3(\mathbb{Z}[\frac{1}{2}, i])$ on X_2 and this action is transitive on the set of p -dimensional cells [Brown 1989].

Therefore all isotropy subgroups for the action on X_2 are, up to isomorphism, subgroups of $\mathrm{SL}_3(\mathbb{Z}[i])$ which itself appears as isotropy group of a 0-dimensional cell in X_2 . The isotropy groups of 1-dimensional and 2-dimensional cells are isomorphic to well-known congruence subgroups of $\mathrm{SL}_3(\mathbb{Z}[i])$. By the Soulé–Lannes method the fiber X_∞ of the projection map p admits a 6-dimensional $\mathrm{SL}_3(\mathbb{Z}[i])$ -equivariant deformation retract (the space of “well-rounded hermitian forms” modulo arithmetic equivalence) with compact quotient [Ash 1984] and

therefore we have

$$E_1^{s,t} = 0 \text{ unless } s = 0, 1, 2, 0 \leq t \leq 6, \quad \text{and} \quad \dim_{\mathbb{F}_2} E_1^{s,t} < \infty \text{ for all } (s, t). \quad (4-2)$$

The E_1 -term of this spectral sequence should be accessible to machine calculation. The spectral sequence will necessarily degenerate at E_3 and the calculation of the differentials is likely to need human intervention, as in the case of $\mathrm{SL}(3, \mathbb{Z}[\frac{1}{2}])$ (compare Section 3.4 of [Henn 1999]). Likewise the calculation of the connecting homomorphism for the mod-2 Borel cohomology of the pair (X, X_s) is likely to require human intervention.

5. On Quillen's conjecture for $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$

The next result gives two reformulations of the conjecture of Quillen briefly discussed in the introduction. The classes e_{2k-1} and e'_{2k-1} in part (c) will be introduced in (5-1) below.

Theorem 5.1. *Suppose $n \geq 2$. The following statements are equivalent:*

- (a) *Conjecture $C(n, \mathbb{Z}[\frac{1}{2}, i], 2)$ holds, i.e., $H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ is a free module over $\mathbb{Z}/2[c_1, \dots, c_n]$ where the c_i are the mod-2 Chern classes of the tautological n -dimensional complex representation of $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$.*
- (b) *The restriction homomorphism*

$$H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \rightarrow H^*(D_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$$

is injective, where $D_n(\mathbb{Z}[\frac{1}{2}, i])$ is the subgroup of diagonal matrices in $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$.

- (c) *There are isomorphisms*

$$H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \cong \mathbb{F}_2[c_1, \dots, c_n] \otimes E(e_1, e'_1, \dots, e_{2n-1}, e'_{2n-1})$$

where the classes c_k are the Chern classes of the tautological n -dimensional complex representation of $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$ and the classes e_{2k-1}, e'_{2k-1} are of cohomological degree $2k - 1$ for $k = 1, \dots, n$.

Proof. It is trivial that (c) implies (a).

In order to show that (a) implies (b) we observe that $D_n(\mathbb{Z}[\frac{1}{2}, i])$ is the centralizer of the unique, up to conjugacy, maximal elementary abelian 2-subgroup E_n of $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$ given by the subgroup of diagonal matrices of order 2. Now consider the top Dickson invariant ω in $H^*(\mathrm{BGL}_n(\mathbb{C}); \mathbb{F}_2)$, i.e., the class whose restriction to $H^*(B(\prod_{i=1}^n \mathrm{GL}_1(\mathbb{C})); \mathbb{F}_2)$ is the product of all nontrivial classes of degree 2. The image of ω in $H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ restricts trivially to the cohomology of all elementary abelian 2-subgroups E of $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i])$ of rank less than n . If (a) holds then the image of ω is not a zero divisor in $H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ and hence

Corollary I.5.8 of [Henn et al. 1995] implies that the restriction to the centralizer of E_n is injective.

The implication (b) \Rightarrow (c) follows from Proposition 5.3 below. \square

Before we go on we introduce the classes e_{2k-1} and e'_{2k-1} . As in the case of GL_2 they are obtained from Quillen's classes [1972] $q_{2k-1} \in H^{2k-1}(\mathrm{GL}_n(\mathbb{F}_5); \mathbb{F}_2)$ which restrict in the cohomology of diagonal matrices in \mathbb{F}_5 to the symmetrization of the classes $y_1 \cdots y_{k-1} x_k$ where y_k is of cohomological degree 2 corresponding to the k -th factor in the product $\prod_{k=1}^n \mathbb{F}_5^\times$ and x_k is of cohomological degree 1 of the same factor. We define

$$e_{2k-1} := \pi^*(q_{2k-1}), \quad e'_{2k-1} := \pi'^*(q_{2k-1}) \quad (5-1)$$

where π and π' are the two ring homomorphisms $\mathbb{Z}[\frac{1}{2}, i] \rightarrow \mathbb{F}_5$ with π sending i to 3 and π' sending i to 2 which we considered earlier in Section 3. We identify the mod-2 cohomology $H^*(D_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ with $\mathbb{F}_2[y_1, \dots, y_n] \otimes E(x_1, x'_1, \dots, x_n, x'_n)$ with $y_k, k = 1, \dots, n$ of degree 2 and $x_k, x'_k, k = 1, \dots, n$ of degree 1 where as before we choose x_k and x'_k to be the basis which is dual to the basis of the k -th factor in

$$D_n(\mathbb{Z}[\frac{1}{2}, i])/D_n(\mathbb{Z}[\frac{1}{2}, i])^2 \cong (\mathbb{Z}[\frac{1}{2}, i]^\times / (\mathbb{Z}[\frac{1}{2}, i]^\times)^2)^n$$

given by the classes of i and $1 + i$. Then we get the following lemma which generalizes (3-10) and whose straightforward proof we leave to the reader.

Lemma 5.2. *The class e_{2k-1} restricts in the cohomology of the subgroup of diagonal matrices $H^*(D_n(\mathbb{Z}[\frac{1}{2}, i]; \mathbb{F}_2))$ to the symmetrization of $y_1 \cdots y_{k-1} x_k$ and the class e'_{2k-1} restricts to the symmetrization of $y_1 \cdots y_{k-1} (x_k + x'_k)$. \square*

The following result determines the image of the restriction homomorphism and shows that (b) implies (c) in Theorem 5.1. It resembles results of Mitchell [1992] for $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}])$ for $p = 2$ and of Anton [1999] for $\mathrm{GL}_n(\mathbb{Z}[\frac{1}{3}, \zeta_3])$ for $p = 3$. Its proof uses crucially condition (5-3) below, which also plays a central role in [Anton 2003].

Proposition 5.3. *Let $n \geq 1$ be an integer. The image of the restriction map*

$$\begin{aligned} i^* : H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \\ \rightarrow H^*(D_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \cong \mathbb{F}_2[y_1, \dots, y_n] \otimes E(x_1, x'_1, \dots, x_n, x'_n) \end{aligned}$$

is isomorphic to

$$\mathbb{F}_2[c_1, \dots, c_n] \otimes E(e_1, e'_1, \dots, e_{2n-1}, e'_{2n-1}).$$

Here we have identified the Chern classes c_i and the classes e_{2i-1} and e'_{2i-1} with their image via i^* . The images of the elements c_i are, of course, the elementary symmetric polynomials in the y_i and the images of the classes e_{2i-1} and e'_{2i-1} have been determined in [Lemma 5.2](#). We remark that even though i^* need not be injective, it is injective on the subalgebra of $H^*(\mathrm{GL}_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ generated by the classes c_i , e_{2i-1} and e'_{2i-1} , $1 \leq i \leq n$.

Proof. In this proof we denote the subalgebra

$$\mathbb{F}_2[c_1, \dots, c_n] \otimes E(e_1, e'_1, \dots, e_{2n-1}, e'_{2n-1}).$$

of $H^*(D_n(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2)$ by C_n and the image of the restriction map by B_n . We need to show that $B_n = C_n$. This is trivial if $n = 1$ and for $n = 2$ this follows from Theorem 1 of [\[Weiss 2006\]](#) (compare (3-2), (3-10) and [Lemma 5.2](#)).

The classes c_1, \dots, c_n are in B_n as images of the Chern classes with the same name and the classes $e_1, \dots, e_{2n-1}, e'_1, \dots, e'_{2n-1}$ are in B_n by [Lemma 5.2](#). Therefore we have $C_n \subset B_n$. We will show $B_n \subset C_n$ for $n \geq 2$ by induction on n . This will be done in three steps:

1. From the inclusions

$$\begin{aligned} \mathrm{GL}_{n-2}(\mathbb{Z}[\tfrac{1}{2}, i]) \times \mathrm{GL}_2(\mathbb{Z}[\tfrac{1}{2}, i]) &\subset \mathrm{GL}_n(\mathbb{Z}[\tfrac{1}{2}, i]) \\ \mathrm{GL}_{n-1}(\mathbb{Z}[\tfrac{1}{2}, i]) \times \mathrm{GL}_1(\mathbb{Z}[\tfrac{1}{2}, i]) &\subset \mathrm{GL}_n(\mathbb{Z}[\tfrac{1}{2}, i]) \end{aligned}$$

given by matrix block sum and the identifications of $D_{n-2}(\mathbb{Z}[\frac{1}{2}, i]) \times D_2(\mathbb{Z}[\frac{1}{2}, i])$ and of $D_{n-1}(\mathbb{Z}[\frac{1}{2}, i]) \times D_1(\mathbb{Z}[\frac{1}{2}, i])$ with $D_n(\mathbb{Z}[\frac{1}{2}, i])$ we see that

$$B_n \subset B_{n-1} \otimes B_1 \cap B_{n-2} \otimes B_2$$

and by the induction hypothesis the latter subalgebra is equal to

$$C_{n-1} \otimes C_1 \cap C_{n-2} \otimes C_2,$$

in particular we have

$$B_n \subset C_{n-1} \otimes C_1 \cap C_{n-2} \otimes C_2. \quad (5-2)$$

2. The monomial basis in

$$H^*(D_n(\mathbb{Z}[\tfrac{1}{2}, i]); \mathbb{F}_2) \cong \mathbb{F}_2[y_1, \dots, y_n] \otimes E(x_1, \dots, x_n, x'_1, \dots, x'_n)$$

is in bijection with the set $S(n)$ of sequences

$$I = (a_1, \varepsilon_{1,1}, \varepsilon_{2,1}, \dots, a_n, \varepsilon_{1,n}, \varepsilon_{2,n})$$

where the a_i are integers ≥ 0 and $\varepsilon_{i,j} \in \{0, 1\}$ for $i = 1, 2$ and $1 \leq j \leq n$. More precisely to I we associate the monomial

$$y^I := y_1^{a_1} \cdots y_n^{a_n} x_1^{\varepsilon_{1,1}} \cdots x_n^{\varepsilon_{1,n}} x_1'^{\varepsilon_{2,1}} \cdots x_n'^{\varepsilon_{2,n}}.$$

We equip $S(n)$ with the lexicographical order and denote it by $<_n$. This order has the property that for each $1 \leq k < n$ it agrees with the lexicographical order on $S(k) \times S(n-k)$ if $S(k)$ and $S(n-k)$ are equipped with the orders $<_k$ and $<_{n-k}$ and $S(n)$ is identified with $S(k) \times S(n-k)$ via concatenation of sequences.

In what follows we replace the symmetrizations of the elements $y_1 \cdots y_{i-1}(x_i + x_i')$, $i = 1, \dots, n$, by the symmetrization of $y_1 \cdots y_{i-1}x_i'$ and by abuse of notation we continue to denote them by e'_{2i-1} . This does not change the subalgebra C_n . This subalgebra

$$\begin{aligned} \mathbb{F}_2[c_1, \dots, c_n] \otimes E(e_1, e'_1, \dots, e_{2n-1}, e'_{2n-1}) \\ \subset \mathbb{F}_2[y_1, \dots, y_n] \otimes E(x_1, \dots, x_n, x'_1, \dots, x'_n) \end{aligned}$$

has a monomial basis which is in bijection with the set $T(n)$ of sequences

$$K = (k_1, \dots, k_n; \phi_{1,1}, \dots, \phi_{1,n}; \phi_{2,1}, \dots, \phi_{2,n})$$

where the k_i are integers ≥ 0 and $\phi_{i,j} \in \{0, 1\}$ for $i = 1, 2$ and $1 \leq j \leq n$. More precisely to K we associate the monomial

$$c^K := c_1^{k_1} \cdots c_n^{k_n} e_1^{\phi_{1,1}} \cdots e_n^{\phi_{1,n}} e'_1{}^{\phi_{2,1}} \cdots e'_n{}^{\phi_{2,n}}.$$

We define a map

$$\alpha : T(n) \rightarrow S(n)$$

by associating to $K \in T(n)$ the largest monomial in $S(n)$ which occurs in the decomposition of c^K as linear combination of elements x^I with $I \in S(n)$. The proof of the following result is elementary and is left to the reader.

Lemma 5.4. *The map α is explicitly given by*

$$\alpha((k_1, \dots, k_n; \phi_{1,1}, \dots, \phi_{1,n}; \phi_{2,1}, \dots, \phi_{2,n})) = (a_1, \varepsilon_{1,1}, \varepsilon_{2,1}, \dots, a_n, \varepsilon_{1,n}, \varepsilon_{2,n})$$

with

$$a_j = k_j + \sum_{i=j+1}^n (k_i + \phi_{1,i} + \phi_{2,i}), \quad 1 \leq j < n,$$

$$a_n = k_n,$$

$$\varepsilon_{i,j} = \phi_{i,j}, \quad 1 \leq j \leq n, \quad i = 1, 2.$$

□

From this lemma it is obvious that α is injective and a sequence

$$I = (a_1, \varepsilon_{1,1}, \varepsilon_{2,1}, \dots, a_n, \varepsilon_{1,n}, \varepsilon_{2,n}) \in S(n)$$

is in the image of α if and only if we have

$$a_j - a_{j+1} \geq \varepsilon_{1,j+1} + \varepsilon_{2,j+1} \quad \text{for all } 1 \leq j < n. \quad (5-3)$$

In particular, if an element x is in C_n then the maximal sequence which appears in the decomposition of x as a linear combination of the monomials x^I with $I \in S(n)$ satisfies (5-3) for all $1 \leq j < n$. Likewise, if x is in $C_i \otimes C_{n-i}$ then this maximal sequence is equal to the maximal sequence which appears in the decomposition of x as a linear combination of the monomials x^I with $I \in S(k) \times S(n-k)$ and hence it satisfies (5-3) for all $1 \leq j < i$ and $i+1 \leq j < n$.

3. Now let x be a homogeneous element of B_n and let I_0 be the maximal sequence in $S(n)$ appearing in the decomposition of x as a linear combination of the monomials x^I with $I \in S(n)$. By (5-2) we have $x \in C_{n-1} \otimes C_1$ and $x \in C_{n-2} \otimes C_2$, and I_0 remains the maximal sequence in $S(n-1) \times S(1)$ and $S(n-2) \times S(2)$, respectively, appearing in the decomposition of x as a linear combination of the monomials x^I with, respectively, $I \in S(n-1) \times S(1)$ and $I \in S(n-2) \times S(2)$. Hence I_0 satisfies conditions (5-3) for $1 \leq j < n-1$ and, respectively, $1 \leq j < n-2$ and $j = n-1$. In particular condition (5-3) holds for all $1 \leq j < n$ and therefore there exists $K_0 \in T(n)$ such that $\alpha(K_0) = I_0$. Then $x - c^{K_0}$ is still in B_n and the maximal sequence appearing in the decomposition of $x - c^{K_0}$ is smaller than that of x . By iterating this procedure we see that x belongs to C_n . \square

Finally we relate $C(3, \mathbb{Z}[\frac{1}{2}, i], 2)$ to the behavior of the restriction homomorphism

$$H^*(\Gamma; \mathbb{F}_2) \rightarrow H^*(C_\Gamma(E_2); \mathbb{F}_2).$$

For this we observe that the subgroups $\Gamma = \text{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$ and the center $Z \cong \mathbb{Z}[\frac{1}{2}, i]^\times$ of $\text{GL}_3(\mathbb{Z}[\frac{1}{2}, i])$ have trivial intersection and their product is the kernel of the homomorphism

$$\text{GL}_3(\mathbb{Z}[\frac{1}{2}, i]) \rightarrow \mathbb{Z}[\frac{1}{2}, i]^\times \rightarrow \mathbb{Z}[\frac{1}{2}, i]^\times / (\mathbb{Z}[\frac{1}{2}, i]^\times)^3 \cong \mathbb{Z}/3$$

given as the composition of the determinant with the natural quotient map. Therefore the spectral sequence of the extension

$$1 \rightarrow \text{SL}_3(\mathbb{Z}[\frac{1}{2}, i]) \times Z \rightarrow \text{GL}_3(\mathbb{Z}[\frac{1}{2}, i]) \rightarrow \mathbb{Z}/3 \rightarrow 1$$

gives an isomorphism

$$H^*(\text{GL}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \cong (H^*(\text{SL}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \otimes H^*(Z; \mathbb{F}_2))^{\mathbb{Z}/3}. \quad (5-4)$$

Proposition 5.5. *Conjecture $C(3, \mathbb{Z}[\frac{1}{2}, i], 2)$ holds if and only if either*

- (a) $H^*(\text{SL}_3(\mathbb{Z}[\frac{1}{2}, i]); \mathbb{F}_2) \cong \mathbb{F}_2[b_2, b_3] \otimes E(d_3, d'_3, d_5, d'_5)$ or

(b) *the kernel of the map ψ of Theorem 1.2 is a finite-dimensional vector space for which the action of $\mathbb{Z}/3 \cong \mathbb{Z}[\frac{1}{2}, i]^\times / (\mathbb{Z}[\frac{1}{2}, i]^\times)^3$ has trivial invariants.*

Proof. Clearly $\mathbb{Z}/3 \cong \mathbb{Z}[\frac{1}{2}, i]^\times / (\mathbb{Z}[\frac{1}{2}, i]^\times)^3$ acts trivially on $H^*(Z; \mathbb{F}_2)$ and on the image of the homomorphism φ of Theorem 1.2. Hence, the corollary follows immediately from (5-4) and Theorem 1.2. \square

Acknowledgement

The author is happy to acknowledge numerous enlightening conversations with Jean Lannes over many years on the topics discussed in this paper.

References

- [Anton 1999] M. F. Anton, “On a conjecture of Quillen at the prime 3”, *J. Pure Appl. Algebra* **144**:1 (1999), 1–20. [MR](#) [Zbl](#)
- [Anton 2003] M. F. Anton, “An elementary invariant problem and general linear group cohomology restricted to the diagonal subgroup”, *Trans. Amer. Math. Soc.* **355**:6 (2003), 2327–2340. [MR](#) [Zbl](#)
- [Ash 1984] A. Ash, “Small-dimensional classifying spaces for arithmetic subgroups of general linear groups”, *Duke Math. J.* **51**:2 (1984), 459–468. [MR](#) [Zbl](#)
- [Brown 1989] K. S. Brown, *Buildings*, Springer, 1989. [MR](#) [Zbl](#)
- [Bui et al. 2016] A. T. Bui, A. Rahm, and M. Wendt, “The Farrell–Tate and Bredon homology for $\mathrm{PSL}_4(\mathbb{Z})$ and other arithmetic groups”, preprint, 2016. [arXiv](#)
- [Dutour Sikirić et al. 2016] M. Dutour Sikirić, H. Gangl, P. E. Gunnells, J. Hanke, A. Schürmann, and D. Yasaki, “On the cohomology of linear groups over imaginary quadratic fields”, *J. Pure Appl. Algebra* **220**:7 (2016), 2564–2589. [MR](#) [Zbl](#)
- [Dwyer 1998] W. G. Dwyer, “Exotic cohomology for $\mathrm{GL}_n(\mathbb{Z}[1/2])$ ”, *Proc. Amer. Math. Soc.* **126**:7 (1998), 2159–2167. [MR](#) [Zbl](#)
- [Henn 1997] H.-W. Henn, “Centralizers of elementary abelian p -subgroups, the Borel construction of the singular locus and applications to the cohomology of discrete groups”, *Topology* **36**:1 (1997), 271–286. [MR](#) [Zbl](#)
- [Henn 1999] H.-W. Henn, “The cohomology of $\mathrm{SL}(3, \mathbb{Z}[1/2])$ ”, *K-Theory* **16**:4 (1999), 299–359. [MR](#) [Zbl](#)
- [Henn et al. 1995] H.-W. Henn, J. Lannes, and L. Schwartz, “Localizations of unstable A -modules and equivariant mod p cohomology”, *Math. Ann.* **301**:1 (1995), 23–68. [MR](#) [Zbl](#)
- [Mitchell 1992] S. A. Mitchell, “On the plus construction for $\mathrm{BGL}_2(\mathbb{Z}[\frac{1}{2}])$ at the prime 2”, *Math. Z.* **209**:2 (1992), 205–222. [MR](#) [Zbl](#)
- [Quillen 1971] D. Quillen, “The spectrum of an equivariant cohomology ring, I, II”, *Ann. of Math.* (2) **94** (1971), 549–572; *ibid.* (2) **94**, 573–602. [MR](#) [Zbl](#)
- [Quillen 1972] D. Quillen, “On the cohomology and K -theory of the general linear groups over a finite field”, *Ann. of Math.* (2) **96** (1972), 552–586. [MR](#) [Zbl](#)
- [Weiss 2006] N. Weiss, *Cohomologie de $\mathrm{GL}_2(\mathbb{Z}[i, 1/2])$ à coefficients dans \mathbb{F}_2* , thèse, Université Louis Pasteur, Strasbourg, 2006, <https://hal.archives-ouvertes.fr/tel-00174888>.

HANS-WERNER HENN:

henn@math.unistra.fr

Institut de Recherche Mathématique Avancée, C.N.R.S., Université de Strasbourg, France

Tunisian Journal of Mathematics

msp.org/tunis

EDITORS-IN-CHIEF

Ahmed Abbes CNRS & IHES, France
abbes@ihes.fr
Ali Baklouti Faculté des Sciences de Sfax, Tunisia
ali.baklouti@fss.usf.tn

EDITORIAL BOARD

Hajer Bahouri CNRS & LAMA, Université Paris-Est Créteil, France
hajer.bahouri@u-pec.fr
Arnaud Beauville Laboratoire J. A. Dieudonné, Université Côte d'Azur, France
beauville@unice.fr
Bassam Fayad CNRS & Institut de Mathématiques de Jussieu-Paris Rive Gauche, Paris, France
bassam.fayad@imj-prg.fr
Benoit Fresse Université Lille 1, France
benoit.fresse@math.univ-lille1.fr
Dennis Gaitsgory Harvard University, United States
gaitsgde@gmail.com
Emmanuel Hebey Université de Cergy-Pontoise, France
emmanuel.hebey@math.u-cergy.fr
Mohamed Ali Jendoubi Université de Carthage, Tunisia
ma.jendoubi@gmail.com
Sadok Kallel Université de Lille 1, France & American University of Sharjah, UAE
sadok.kallel@math.univ-lille1.fr
Minhyong Kim Oxford University, UK & Korea Institute for Advanced Study, Seoul, Korea
minhyong.kim@maths.ox.ac.uk
Toshiyuki Kobayashi The University of Tokyo & Kavli IPMU, Japan
toshi@kurims.kyoto-u.ac.jp
Yanyan Li Rutgers University, United States
yyli@math.rutgers.edu
Nader Masmoudi Courant Institute, New York University, United States
masmoudi@cims.nyu.edu
Haynes R. Miller Massachusetts Institute of Technology, United States
hrm@math.mit.edu
Nordine Mir Texas A&M University at Qatar & Université de Rouen Normandie, France
nordine.mir@qatar.tamu.edu
Detlef Müller Christian-Albrechts-Universität zu Kiel, Germany
mueller@math.uni-kiel.de
Mohamed Sifi Université Tunis El Manar, Tunisia
mohamed.sifi@fst.utm.tn
Daniel Tataru University of California, Berkeley, United States
tataru@math.berkeley.edu
Sundaram Thangavelu Indian Institute of Science, Bangalore, India
veluma@math.iisc.ernet.in
Saïd Zarati Université Tunis El Manar, Tunisia
said.zarati@fst.utm.tn

PRODUCTION

Silvio Levy (Scientific Editor)
production@msp.org

The Tunisian Journal of Mathematics is an international publication organized by the Tunisian Mathematical Society (<http://www.tms.rnu.tn>) and published in electronic and print formats by MSP in Berkeley.

See inside back cover or msp.org/tunis for submission instructions.

The subscription price for 2019 is US \$315/year for the electronic version, and \$370/year (+\$20, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Tunisian Journal of Mathematics (ISSN 2576-7666 electronic, 2576-7658 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

TJM peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Tunisian Journal of Mathematics

2019

vol. 1

no. 4

Grothendieck–Messing deformation theory for varieties for K3 type	455
ANDREAS LANGER and THOMAS ZINK	
Purity of crystalline strata	519
JINGHAO LI and ADRIAN VASIU	
On the mod-2 cohomology of $\mathrm{SL}_3(\mathbb{Z}[\frac{1}{2}, i])$	539
HANS-WERNER HENN	
Geometric origin and some properties of the arctangential heat equation	561
YANN BRENIER	
Horn’s problem and Fourier analysis	585
JACQUES FARAUT	