

GLOBAL EXISTENCE AND ENERGY DECAY OF SOLUTIONS TO THE CAUCHY PROBLEM FOR A WAVE EQUATION WITH A WEAKLY NONLINEAR DISSIPATION

ABBÈS BENAÏSSA AND SOUFIANE MOKEDDEM

Received 21 November 2003

We prove the global existence and study decay properties of the solutions to the wave equation with a weak nonlinear dissipative term by constructing a stable set in $H^1(\mathbb{R}^n)$.

1. Introduction

We consider the Cauchy problem for the nonlinear wave equation with a weak nonlinear dissipation and source terms of the type

$$\begin{aligned}u'' - \Delta_x u + \lambda^2(x)u + \sigma(t)g(u') &= |u|^{p-1}u \quad \text{in } \mathbb{R}^n \times [0, +\infty[, \\u(x, 0) = u_0(x), \quad u'(x, 0) &= u_1(x) \quad \text{in } \mathbb{R}^n,\end{aligned}\tag{1.1}$$

where $g : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous nondecreasing function and λ and σ are positive functions.

When we have a bounded domain instead of \mathbb{R}^n , and for the case $g(x) = \delta x$ ($\delta > 0$) (without the term $\lambda^2(x)u$), Ikehata and Suzuki [8] investigated the dynamics, they have shown that for sufficiently small initial data (u_0, u_1) , the trajectory $(u(t), u'(t))$ tends to $(0, 0)$ in $H_0^1(\Omega) \times L^2(\Omega)$ as $t \rightarrow +\infty$. When $g(x) = \delta|x|^{m-1}x$ ($m \geq 1, \lambda \equiv 0, \sigma \equiv 1$), Georgiev and Todorova [4] introduced a new method and determined suitable relations between m and p , for which there is global existence or alternatively finite-time blow up. Precisely they showed that the solutions continue to exist globally in time if $m \geq p$ and blow up in finite time if $m < p$ and the initial energy is sufficiently negative. This result was later generalized to an abstract setting by Levine and Serrin [12] and Levine et al. [11]. In these papers, the authors showed that no solution with negative initial energy can be extended on $[0, \infty[$, if the source term dominates over the damping term ($p > m$). This generalization allowed them also to apply their result to quasilinear wave equations (see [1, 17]). Quite recently, Ikehata [7] proved that a global solution exists with no relation between p and m by the use of a stable set method due to Sattinger [18].

For the Cauchy problem (1.1) with $\lambda \equiv 1$ and $\sigma \equiv 1$, when $g(x) = \delta|x|^{m-1}x$ ($m \geq 1$) Todorova [21] (see [16]) proved that the energy decay rate is $E(t) \leq (1+t)^{-(2-n(m-1))/(m-1)}$

for $t \geq 0$. She used a general method introduced by Nakao [14] on condition that the data have compact support. Unfortunately, this method does not seem to be applicable in the case of more general functions λ and σ .

Our purpose in this paper is to give a global solvability in the class H^1 and energy decay estimates of the solutions to the Cauchy problem (1.1) for a weak linear perturbation and a weak nonlinear dissipation.

We use a new method recently introduced by Martinez [13] (see also [2]) to study the decay rate of solutions to the wave equation $u'' - \Delta_x u + g(u') = 0$ in $\Omega \times \mathbb{R}^+$, where Ω is a bounded domain of \mathbb{R}^n . This method is based on a new integral inequality that generalizes a result of Haraux [6]. So we proceed with the argument combining the method in [13] with the concept of modified stable set on $H^1(\mathbb{R}^n)$. Here the modified stable set is the extended \mathbb{R}^n version of Sattinger's stable set.

2. Preliminaries and main results

$\lambda(x)$, $\sigma(t)$, and g satisfy the following hypotheses.

(i) $\lambda(x)$ is a locally bounded measurable function defined on \mathbb{R}^n and satisfies

$$\lambda(x) \geq d(|x|), \tag{2.1}$$

where d is a decreasing function such that $\lim_{y \rightarrow \infty} d(y) = 0$.

(ii) $\sigma : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a nonincreasing function of class C^1 on \mathbb{R}_+ .

Consider $g : \mathbb{R} \rightarrow \mathbb{R}$ a nondecreasing C^0 function and suppose that there exist $C_i > 0$, $i = 1, 2, 3, 4$, such that

$$c'_3 |v|^m \leq |g(v)| \leq c'_4 |v|^{1/m} \quad \text{if } |v| \leq 1, \tag{2.2}$$

$$c_1 |v| \leq |g(v)| \leq c_2 |v|^r \quad \forall |v| \geq 1, \tag{2.3}$$

where $m \geq 1$ and $1 \leq r \leq (n + 2)/(n - 2)^+$.

We first state two well-known lemmas, and then we state and prove two other lemmas that will be needed later.

LEMMA 2.1. *Let q be a number with $2 \leq q < +\infty$ ($n = 1, 2$) or $2 \leq q \leq 2n/(n - 2)$ ($n \geq 3$). Then there is a constant $c_* = c(q)$ such that*

$$\|u\|_q \leq c_* \|u\|_{H^1(\mathbb{R}^n)} \quad \text{for } u \in H^1(\mathbb{R}^n). \tag{2.4}$$

LEMMA 2.2 (Gagliardo-Nirenberg). *Let $1 \leq r < q \leq +\infty$ and $p \geq 2$. Then, the inequality*

$$\|u\|_p \leq C \|\nabla_x^m u\|_2^\theta \|u\|_r^{1-\theta} \quad \text{for } u \in \mathcal{D}((-\Delta)^{m/2})L^r \tag{2.5}$$

holds with some constant $C > 0$ and

$$\theta = \left(\frac{1}{r} - \frac{1}{p}\right) \left(\frac{m}{n} + \frac{1}{r} - \frac{1}{2}\right)^{-1} \tag{2.6}$$

provided that $0 < \theta \leq 1$ (assuming that $0 < \theta < 1$ if $m - n/2$ is a nonnegative integer).

LEMMA 2.3 [10]. Let $E : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a nonincreasing function and assume that there are two constants $p \geq 1$ and $A > 0$ such that

$$\int_S^{+\infty} E^{(p+1)/2}(t) dt \leq AE(S), \quad 0 \leq S < +\infty, \tag{2.7}$$

then

$$\begin{aligned} E(t) &\leq cE(0)(1+t)^{-2/(p-1)} \quad \forall t \geq 0, \text{ if } p > 1, \\ E(t) &\leq cE(0)e^{-\omega t} \quad \forall t \geq 0, \text{ if } p = 1, \end{aligned} \tag{2.8}$$

where c and ω are positive constants independent of the initial energy $E(0)$.

LEMMA 2.4 [13]. Let $E : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a nonincreasing function and $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ an increasing C^2 function such that

$$\phi(0) = 0, \quad \phi(t) \rightarrow +\infty \quad \text{as } t \rightarrow +\infty. \tag{2.9}$$

Assume that there exist $p \geq 1$ and $A > 0$ such that

$$\int_S^{+\infty} E(t)^{(p+1)/2}(t)\phi'(t) dt \leq AE(S), \quad 0 \leq S < +\infty, \tag{2.10}$$

then

$$\begin{aligned} E(t) &\leq cE(0)(1+\phi(t))^{-2/(p-1)} \quad \forall t \geq 0, \text{ if } p > 1, \\ E(t) &\leq cE(0)e^{-\omega\phi(t)} \quad \forall t \geq 0, \text{ if } p = 1, \end{aligned} \tag{2.11}$$

where c and ω are positive constants independent of the initial energy $E(0)$.

Proof of Lemma 2.4. Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be defined by $f(x) := E(\phi^{-1}(x))$, (we remark that ϕ^{-1} has a sense by the hypotheses assumed on ϕ). f is nonincreasing, $f(0) = E(0)$, and if we set $x := \phi(t)$, we obtain

$$\begin{aligned} \int_{\phi(S)}^{\phi(T)} f(x)^{(p+1)/2} dx &= \int_{\phi(S)}^{\phi(T)} E(\phi^{-1}(x))^{(p+1)/2} dx = \int_S^T E(t)^{(p+1)/2} \phi'(t) dt \\ &\leq AE(S) = Af(\phi(S)), \quad 0 \leq S < T < +\infty. \end{aligned} \tag{2.12}$$

Setting $s := \phi(S)$ and letting $T \rightarrow +\infty$, we deduce that

$$\int_s^{+\infty} f(x)^{(p+1)/2} dx \leq Af(s), \quad 0 \leq s < +\infty. \tag{2.13}$$

Thanks to Lemma 2.3, we deduce the desired results. □

Before stating the global existence theorem and decay property of problem (1.1), we will introduce the notion of the modified stable set. Let

$$\begin{aligned} K(u) &= \|\nabla_x u\|_2^2 + \|u\|_2^2 - \|u\|_{p+1}^{p+1} \quad \text{if } \lambda \equiv 1, \\ I(u) &= \|\nabla_x u\|_2^2 - \|u\|_{p+1}^{p+1} \quad \text{if } \lambda \neq \text{const}, \end{aligned} \tag{2.14}$$

for $u \in H^1(\mathbb{R}^n)$. Then we define the modified stable set \widetilde{W}^* and \widetilde{W}^{**} by

$$\begin{aligned} \widetilde{W}^* &\equiv \{u \in H^1(\mathbb{R}^n) \setminus K(u) > 0\} \cup \{0\} \quad \text{if } \lambda \equiv 1, \\ \widetilde{W}^{**} &\equiv \{u \in H^1(\mathbb{R}^n) \setminus I(u) > 0\} \cup \{0\} \quad \text{if } \lambda \neq \text{const}. \end{aligned} \tag{2.15}$$

Next, let $J(u)$ and $E(t)$ be the potential and energy associated with problem (1.1), respectively:

$$\begin{aligned} J(u) &= \frac{1}{2} \|\nabla_x u\|_2^2 + \frac{1}{2} \|\lambda(x)u\|_2^2 - \frac{1}{p+1} \|u\|_{p+1}^{p+1} \quad \text{for } u \in H^1(\mathbb{R}^n), \\ E(t) &= \frac{1}{2} \|u'\|_2^2 + J(u). \end{aligned} \tag{2.16}$$

We get the local existence solution.

THEOREM 2.5. *Let $1 < p \leq (n+2)/(n-2)$ ($1 < p < \infty$ if $n = 1, 2$) and assume that $(u_0, u_1) \in H^1(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ and u_0 belong to the modified stable set \widetilde{W}^* . Then there exists $T > 0$ such that the Cauchy problem (1.1) has a unique solution $u(t)$ on $\mathbb{R}^n \times [0, T)$ in the class*

$$u(t, x) \in C([0, T); H^1(\mathbb{R}^n)) \cap C^1([0, T); L^2(\mathbb{R}^n)), \tag{2.17}$$

satisfying

$$u(t) \in \widetilde{W}^*, \tag{2.18}$$

and this solution can be continued in time as long as $u(t) \in \widetilde{W}^*$.

When $\lambda \neq \text{const}$, we use the following theorem of local existence in the space $H^2 \times H^1$, and the decay property of the energy $E(t)$ is necessarily required for the local solution to remain in \widetilde{W}^{**} as $t \rightarrow \infty$; this fact of course guarantees the global existence in $H^2 \times H^1$ and by approximation, we obtain global existence in $H^1 \times L^1$.

THEOREM 2.6 [15]. *Let $(u_0, u_1) \in H^2 \times H^1$. Suppose that*

$$1 \leq p \leq \frac{n}{n-4} \quad (1 \leq \infty \text{ if } n \leq 4). \tag{2.19}$$

Then under the hypotheses (2.1), (2.2), and (2.3), problem (1.1) admits a unique local solution $u(t)$ on some interval $[0, T[$, $T \equiv T(u_0, u_1) > 0$, in the class $W^{2,\infty}([0, T[; L^2) \cap W^{1,\infty}([0, T[; H^1) \cap L^\infty([0, T[; H^2)$, satisfying the finite propagation property.

Proof of Theorem 2.5 (see [15, 18]). Since the argument is standard, we only sketch the main idea of the proof. Let $(u_0, u_1) \in H^1 \times L^2$ and $u_0 \in \widetilde{W}^*$. Then we have a unique local solution $u(t)$ for some $T > 0$. Indeed, taking suitable approximate functions f_j such that (see [20])

$$f_j(u) = f(u) \quad \text{if } |u| \leq j, \quad |f_j(u)| \leq |f(u)|, \quad |f_j(u)| \leq c_j |u|, \tag{2.20}$$

problem (1.1) with $f(u) \equiv |u|^{p-1}u$ replaced by $f_j(u)$ admits a unique solution $u_j(t) \in C([0, T); H^1(\mathbb{R}^n)) \cap C^1([0, T); L^2(\mathbb{R}^n))$. Further, we can prove that $u_j(t) \in \widetilde{W}^*$, $0 < t < T$,

for sufficiently large j , and there exists a subsequence of $\{u_j(t)\}$ which converges to a function $\tilde{u}(t)$ in certain senses. $\tilde{u}(t)$ is, in fact, a weak solution in $C([0, T]; H^1(\mathbb{R}^n)) \cap C^1([0, T]; L^2(\mathbb{R}^n))$ (see [19, 20]) and such a solution is unique by Ginibre and Velo [5] and Brenner [3]. We can also construct such a solution which meets moreover the finite propagation property, if we assume that the initial data $u_0(x)$ and $u_1(x)$ are of compact support:

$$\text{supp } u_0 \cup \text{supp } u_1 \subset \{x \in \mathbb{R}^n, |x| < L\}, \quad \text{for some } L > 0. \tag{2.21}$$

Applying [9, Appendix 1] of John, then the solution is also of compact support: $\text{supp } u_j(t) \subset \{x \in \mathbb{R}^n, |x| < L + t\}$. So, we have $\text{supp } \tilde{u}(t) \subset \{x \in \mathbb{R}^n, |x| < L + t\}$. \square

We denote the life span of the solution $u(t, x)$ of the Cauchy problem (1.1) by T_{\max} . First we consider the case $\lambda(x) \equiv \text{const}$ ($\lambda(x) \equiv 1$ without loss of generality). And construct a stable set in $H^1(\mathbb{R}^n)$.

Setting

$$C_0 \equiv K \left\{ \frac{2(p+1)}{(p-1)} \right\}^{(p-1)/2}, \tag{2.22}$$

$$\int_0^\infty \sigma(\tau) d\tau = +\infty \quad \text{if } m = 1, \tag{2.23}$$

$$\int_0^\infty (1+\tau)^{-n(m-1)/2} \sigma(\tau) d\tau = +\infty \quad \text{if } m > 1. \tag{2.24}$$

THEOREM 2.7. *Let $u(t, x)$ be a local solution of problem (1.1) on $[0, T_{\max})$ with initial data $u_0 \in \widetilde{W}^*$, $u_1 \in L^2(\mathbb{R}^n)$ with sufficiently small initial energy $E(0)$ so that*

$$C_0 E(0)^{(p-1)/2} < 1. \tag{2.25}$$

Then $T_{\max} = \infty$. Furthermore, the global solution of the Cauchy problem (1.1) has the following energy decay property. Under (2.22), (2.3), and (2.23),

$$E(t) \leq E(0) \exp \left(1 - \omega \int_0^t \sigma(\tau) d\tau \right) \quad \forall t > 0. \tag{2.26}$$

Under (2.2), (2.3), and (2.24),

$$E(t) \leq \left(\frac{C(E(0))}{\int_0^t (1+\tau)^{-n(m-1)/2} \sigma(\tau) d\tau} \right)^{2/(m-1)} \quad \forall t > 0. \tag{2.27}$$

Secondly, we consider the case $\lambda(x) \not\equiv \text{const}$ and we assume that

$$\frac{n+4}{n} \leq p \leq \frac{n}{n-2}. \tag{2.28}$$

(1) If $\sigma(t) = \mathcal{O}(\bar{d}(t))$, where $\bar{d}(t) = d(L+t)$.

If $m = 1$, we suppose that

$$\int_0^\infty \sigma(\tau) d\tau = +\infty \tag{2.29}$$

with

$$\begin{aligned} (\tilde{d}(t))^{-4-(n-2)(p-1)/2} \exp\left(1 - \omega \int_0^t \sigma(\tau) d\tau\right)^{(p-1)/2} < \infty, \\ (\tilde{d}(t))^{-1} \exp\left(\frac{1}{2} - \frac{\omega}{2} \int_0^t \sigma(\tau) d\tau\right) < \infty. \end{aligned} \tag{2.30}$$

If $m > 1$, we suppose that

$$\int_0^\infty (1 + \tau)^{-n(m-1)/2} \sigma(\tau) d\tau = \infty \tag{2.31}$$

with

$$\begin{aligned} \frac{(\tilde{d}(t))^{-4-(n-2)(p-1)/2}}{\left(\int_0^t (1 + \tau)^{-n(m-1)/2} \sigma(\tau) d\tau\right)^{(p-1)/(m-1)}} < \infty, \\ \frac{(\tilde{d}(t))^{-1}}{\left(\int_0^t (1 + \tau)^{-n(m-1)/2} \sigma(\tau) d\tau\right)^{1/(m-1)}} < \infty. \end{aligned} \tag{2.32}$$

(2) If $\tilde{d}(t) = \mathcal{O}(\sigma(t))$.

If $m = 1$, we suppose that for some $0 \leq \alpha < 1$,

$$\int_0^\infty \frac{\tilde{d}^2(\tau)}{\sigma^\alpha(\tau)} d\tau = +\infty \tag{2.33}$$

with

$$\begin{aligned} (\tilde{d}(t))^{-4-(n-2)(p-1)/2} \exp\left(1 - \omega \int_0^t \frac{\tilde{d}^2(\tau)}{\sigma^\alpha(\tau)} d\tau\right)^{(p-1)/2} < \infty, \\ (\tilde{d}(t))^{-1} \exp\left(\frac{1}{2} - \frac{\omega}{2} \int_0^t \frac{\tilde{d}^2(\tau)}{\sigma^\alpha(\tau)} d\tau\right) < \infty. \end{aligned} \tag{2.34}$$

If $m > 1$, we suppose that for some $0 \leq \alpha < 1$,

$$\int_0^\infty (1 + \tau)^{-n(m-1)/2} \sigma^{-((1+\alpha)(1+m)-2)/2}(\tau) \tilde{d}^{m+1}(\tau) d\tau = \infty \tag{2.35}$$

with

$$\frac{(\tilde{d}(t))^{-4-(n-2)(p-1)/2}}{\left(\int_0^t (1+\tau)^{-n(m-1)/2} \sigma^{-((1+\alpha)(1+m)-2)/2}(\tau) \tilde{d}^{m+1}(\tau) d\tau\right)^{(p-1)/(m-1)}} < \infty,$$

$$\frac{(\tilde{d}(t))^{-1}}{\left(\int_0^t (1+\tau)^{-n(m-1)/2} \sigma^{-((1+\alpha)(1+m)-2)/2}(\tau) \tilde{d}^{m+1}(\tau) d\tau\right)^{1/(m-1)}} < \infty. \tag{2.36}$$

We have the following theorem.

THEOREM 2.8. *Let $(u_0, u_1) \in H^1 \times L^2$, $u_0 \in \widetilde{W}^{**}$, and let the initial energy $E(0)$ be sufficiently small. The following cases are considered.*

(i) $\sigma(t) = \mathbb{O}(\tilde{d}(t))$.

Suppose (2.2), (2.3), (2.29), and (2.30) or (2.2), (2.3), (2.31), and (2.32). Then problem (1.1) admits a unique solution $u(t) \in C([0, \infty); H^1) \cap C^1([0, \infty); L^2)$ and has the same decay property as Theorem 2.7.

(ii) $\tilde{d}(t) = \mathbb{O}(\sigma(t))$.

Suppose (2.2), (2.3), (2.33), and (2.34) or (2.2), (2.3), (2.35), and (2.36). Then problem (1.1) admits a unique solution $u(t) \in C([0, \infty); H^1) \cap C^1([0, \infty); L^2)$. Furthermore, the global solution of the Cauchy problem (1.1) has the following energy decay property:

$$E(t) \leq E(0) \exp\left(1 - \omega \int_0^t \frac{\tilde{d}^2(\tau)}{\sigma^\alpha(\tau)} d\tau\right) \quad \forall t > 0 \text{ if } m = 1, \tag{2.37}$$

$$E(t) \leq \left(\frac{C(E(0))}{\int_0^t (1+\tau)^{-n(m-1)/2} \sigma^{-((1+\alpha)(1+m)-2)/2}(\tau) \tilde{d}^{m+1}(\tau) d\tau}\right)^{2/(m-1)} \quad \forall t > 0 \text{ if } m > 1. \tag{2.38}$$

Remark 2.9. In Theorem 2.7, the global existence and energy decay are independent, but in Theorem 2.8, we need the estimation of the energy decay for a local solution to prove global existence.

Examples 2.10. (1) If $\sigma(t) = 1/t^\theta$, by applying Theorem 2.7 we obtain

$$E(t) \leq E(0)e^{1-\omega t^{1-\theta}} \quad \text{if } m = 1,$$

$$E(t) \leq C(E(0))(1+t)^{-(2-n(m-1)-2\theta)/(m-1)} \quad \text{if } 1 < m < 1 + \frac{2-2\theta}{n}, 0 < \theta < 1, \tag{2.39}$$

$$E(t) \leq C(E(0))(\ln t)^{-2/(m-1)} \quad \text{if } m = 1 + \frac{2-2\theta}{n}, 0 < \theta < 1.$$

(2) If $\sigma(t) = 1/t^\theta \ln t \ln_2 t \cdots \ln_p t$, by applying Theorem 2.7, we obtain

$$E(t) \leq E(0)(\ln_p t)^{-\omega} \quad \text{if } m = 1, \theta = 1. \tag{2.40}$$

For example, if $n(m-1)/2 + \theta = 1$, that is, $1 < m < 1 + 2/n$,

$$E(t) \leq C(E(0))(\ln_p t)^{-2/(m-1)}. \tag{2.41}$$

(3) If $\sigma(t) = 1/t^\theta$ and $d(r) = 1/r^\gamma$ with $\theta \geq \gamma$ by applying [Theorem 2.8](#), we obtain

$$E(t) \leq C(E(0))(1+t)^{-(2-n(m-1)-2\theta)/(m-1)} \quad \text{if } 1 < m < 1 + \frac{2-2\theta}{2\gamma+n}, \quad 0 < \theta < 1, \tag{2.42}$$

$$E(t) \leq C(E(0))(\ln t)^{-2/(m-1)} \quad \text{if } m = 1 + \frac{2-2\theta}{2\gamma+n}, \quad 0 < \theta < 1.$$

In order to show the global existence, it suffices to obtain the a priori estimates for $E(t)$ and $\|u(t)\|_2$ in the interval of existence.

To prove [Theorem 2.7](#) we first have the following energy identity to problem (1.1).

LEMMA 2.11 (energy identity). *Let $u(t, x)$ be a local solution to problem (1.1) on $[0, T_{\max})$ as in [Theorem 2.5](#). Then*

$$E(t) + \int_{\mathbb{R}^n} \int_0^t \sigma(s)u'(s)g(u'(s)) ds dx = E(0) \tag{2.43}$$

for all $t \in [0, T_{\max})$.

Next we state several facts about the modified stable set \widetilde{W}^* .

LEMMA 2.12. *Suppose that*

$$1 < p \leq \frac{n+2}{n-2}. \tag{2.44}$$

Then

- (i) \widetilde{W}^* is a neighborhood of 0 in $H^1(\mathbb{R}^n)$,
- (ii) for $u \in \widetilde{W}^*$,

$$J(u) \geq \frac{p-1}{2(p+1)} (\|\nabla_x u\|_2^2 + \|u\|_2^2). \tag{2.45}$$

Proof of Lemma 2.12. (i) From [Lemma 2.1](#) we have

$$\|u\|_{p+1}^{p+1} \leq K \|u\|_{H^1}^{p+1} \leq K \|u\|_{H^1}^{p-1} (\|u\|_2^2 + \|\nabla_x u\|_2^2). \tag{2.46}$$

Let

$$U(0) \equiv \left\{ u \in H^1(\mathbb{R}^N) \setminus \|u\|_{H^1}^{p-1} < \frac{1}{K} \right\}. \tag{2.47}$$

Then, for any $u \in U(0) \setminus \{0\}$, we deduce from (2.46) that

$$\|u\|_{p+1}^{p+1} < \|u\|_2^2 + \|\nabla_x u\|_2^2, \tag{2.48}$$

that is, $K(u) > 0$. This implies $U(0) \subset \widetilde{W}^*$.

(ii) By the definition of $K(u)$ and $J(u)$, we have the identity

$$(p+1)J(u) = K(u) + \frac{(p-1)}{2} (\|\nabla_x u\|_2^2 + \|u\|_2^2). \tag{2.49}$$

Since $u \in \widetilde{W}^*$, we have $K(u) \geq 0$. Therefore from (2.44) we get the desired in-equality (2.45). \square

LEMMA 2.13. Let $u(t)$ be a solution to problem (1.1) on $[0, T_{\max})$. Suppose (2.44) holds. If $u_0 \in \widetilde{W}^*$ and $u_1 \in L^2(\mathbb{R}^n)$ satisfy

$$C_0 E(0)^{(p-1)/2} < 1, \tag{2.50}$$

then

- (i) $u(t) \in \widetilde{W}^*$ on $[0, T_{\max})$,
- (ii) $\|u(t)\|_2 \leq I_0$ on $[0, T_{\max})$.

Proof of Lemma 2.13. Suppose that there exists a number $t^* \in [0, T_{\max}[$ such that $u(t) \in \widetilde{W}^*$ on $[0, t^*[$ and $u(t^*) \notin \widetilde{W}^*$. Then we have

$$K(u(t^*)) = 0, \quad u(t^*) \neq 0. \tag{2.51}$$

Since $u(t) \in \widetilde{W}^*$ on $[0, t^*[$, it holds that

$$\frac{p-1}{2(p+1)} (\|\nabla_x u\|_2^2 + \|u\|_2^2) \leq J(u) \leq E(t); \tag{2.52}$$

it follows from the nonincreasing of the energy that

$$\|\nabla_x u\|_2^2 + \|u\|_2^2 \leq \frac{2(p+1)}{p-1} E(0) \equiv I_0^2. \tag{2.53}$$

Hence, we obtain

$$\|u\|_2^2 \leq \frac{2(p+1)}{p-1} E(0) \equiv I_0^2 \quad \text{on } [0, t^*]. \tag{2.54}$$

Next, from Lemma 2.1 and (2.54) we have

$$\begin{aligned} \|u\|_{p+1}^{p+1} &\leq K \|u(t)\|_{H^1(\mathbb{R}^n)}^{p+1} \\ &\leq K \|u(t)\|_{H^1(\mathbb{R}^n)}^{p-1} (\|\nabla_x u\|_2^2 + \|u\|_2^2) \\ &\leq K I_0^{p-1} (\|\nabla_x u\|_2^2 + \|u\|_2^2) \\ &\leq C_0 E(0)^{(p-1)/2} (\|u(t)\|_2^2 + \|\nabla_x u(t)\|_2^2) \end{aligned} \tag{2.55}$$

for all $t \in [0, t^*]$, where C_0 is the constant defined by (2.22). Note that from (2.55) and our hypothesis

$$\eta_0 \equiv C_0 E(0)^{(p-1)/2} < 1, \tag{2.56}$$

it follows that

$$\|u(t)\|_{p+1}^{p+1} \leq (1 - \eta_0) (\|u(t)\|_2^2 + \|\nabla_x u(t)\|_2^2). \tag{2.57}$$

Therefore, we obtain

$$K(u(t^*)) \geq \eta_0(\|u(t^*)\|_2^2 + \|\nabla_x u(t^*)\|_2^2) \tag{2.58}$$

which contradicts (2.51). Thus, we conclude that $u(t) \in \widetilde{\mathcal{W}}^*$ on $[0, T_{\max}[$. The assertion (ii) can be obtained by the same argument as for (2.54). This completes the proof of Lemma 2.13. \square

LEMMA 2.14. *Under the same assumptions as in Lemma 2.13, there exists a constant M_2 depending on $\|u_0\|_{H^1}$ and $\|u_1\|_2$ such that*

$$\|u(t)\|_{H^1}^2 + \|u'(t)\|_2^2 \leq M_2^2 \tag{2.59}$$

for all $t \in [0, T_{\max}[$.

Proof of Lemma 2.14. It follows from Lemma 2.13 that $u(t) \in \widetilde{\mathcal{W}}^*$ on $[0, T_{\max}[$. So Lemma 2.12(ii) implies that

$$J(u) \geq \frac{p-1}{2(p+1)}(\|u(t)\|_2^2 + \|\nabla_x u(t)\|_2^2) \quad \text{on } [0, T_{\max}[. \tag{2.60}$$

Hence, from Lemma 2.11 and (2.60) we get

$$\frac{1}{2}\|u'(t)\|_2^2 + \frac{p-1}{2(p+1)}(\|u\|_2^2 + \|\nabla_x u(t)\|_2^2) \leq E(t) \leq E(0). \tag{2.61}$$

So we get

$$\|u(t)\|_{H^1}^2 + \|u'(t)\|_2^2 \leq M_2^2, \tag{2.62}$$

for some $M_2 > 0$.

The above inequality and the continuation principle lead to the global existence of the solution, that is, $T_{\max} = \infty$. \square

Proof of the energy decay. From now on, we denote by c various positive constants which may be different at different occurrences. We multiply the first equation of (1.1) by $E^q \phi' u$, where ϕ is a function satisfying all the hypotheses of Lemma 2.4. We obtain

$$\begin{aligned} 0 &= \int_S^T E^q \phi' \int_{\mathbb{R}^n} u(u'' - \Delta u + u + \sigma(t)g(u') - |u|^{p-1}u) \, dx \, dt \\ &= \left[E^q \phi' \int_{\mathbb{R}^n} uu' \, dx \right]_S^T - \int_S^T (qE'E^{q-1}\phi' + E^q\phi'') \int_{\mathbb{R}^n} uu' \, dx \, dt - 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 \, dx \, dt \\ &\quad + \int_S^T E^q \phi' \int_{\mathbb{R}^n} \left(u'^2 + |u|^2 + |\nabla u|^2 - \frac{2}{p+1}|u|^{p+1} \right) \, dx \, dt + \int_S^T E^q \phi' \int_{\mathbb{R}^n} \sigma(t)ug(u') \, dx \, dt \\ &\quad \quad \quad \int_S^T E^q \phi' \int_{\mathbb{R}^n} \left(\frac{2}{p+1} - 1 \right) |u|^{p+1} \, dx \, dt. \end{aligned} \tag{2.63}$$

Since

$$\begin{aligned} \left(1 - \frac{2}{p+1}\right) \int_{\mathbb{R}^n} |u|^{p+1} dx &\leq (1 - \eta_0) \frac{p-1}{p+1} \|u(t)\|_{H^1(\mathbb{R}^n)}^2 dx \\ &\leq (1 - \eta_0) \frac{p-1}{p+1} \frac{2(p+1)}{p-1} E(t) \\ &= 2(1 - \eta_0)E(t), \end{aligned} \tag{2.64}$$

we deduce that

$$\begin{aligned} 2\eta_0 \int_S^T E^{q+1} \phi' dt &\leq - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T + \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt \\ &\quad + 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt - \int_S^T E^q \phi' \int_{\mathbb{R}^n} \sigma(t)ug(u') dx dt \\ &\leq - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T + \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt \tag{2.65} \\ &\quad + 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt + c(\varepsilon) \int_S^T E^q \phi' \int_{|u'| \leq 1} g(u')^2 dx dt \\ &\quad + \varepsilon \int_S^T E^q \phi' \int_{\mathbb{R}^n} u^2 dx dt + \int_S^T E^q \phi' \int_{|u'| \geq 1} \sigma(t)ug(u') dx dt \end{aligned}$$

for every $\varepsilon > 0$. Also, applying Hölder’s and Young’s inequalities, we have

$$\begin{aligned} &\int_S^T E^q \phi' \int_{|u'| > 1} \sigma(t)ug(u') dx dt \\ &\leq \int_S^T E^q \phi' \sigma(t) \left(\int_{\Omega} |u|^{r+1} dx \right)^{1/(r+1)} \left(\int_{|u'| > 1} |g(u')|^{(r+1)/r} dx \right)^{r/(r+1)} dt \\ &\leq c \int_S^T E^{(2q+1)/2} \phi' \sigma^{1/(r+1)}(t) \left(\int_{|u'| > 1} \sigma(t)u'g(u') dx \right)^{r/(r+1)} dt \\ &\leq \int_S^T \phi' \sigma^{1/(r+1)}(t) E^{(2q+1)/2} (-E')^{r/(r+1)} dt \tag{2.66} \\ &\leq c \int_S^T \phi' \left(\sigma^{1/(r+1)}(t) E^{(2q+1)/2 - r/(r+1)} \right) \left((-E')^{r/(r+1)} E^{r/(r+1)} \right) dt \\ &\leq c(\varepsilon') \int_S^T \phi' (-E'E) dt + \varepsilon' \int_S^T \phi' \sigma(t) E^{(r+1)((2q+1)/2 - r/(r+1))} dt \\ &\leq c(\varepsilon') E(S)^2 + \varepsilon' \sigma(0) E(0)^{(2rq-r-1)/2} \int_S^T \phi' E^{q+1} dt \end{aligned}$$

for every $\varepsilon' > 0$. Choosing ε and ε' small enough, we obtain

$$\begin{aligned} \int_S^T E^{q+1} \phi' dt &\leq - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T + \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt \\ &\quad + \int_{|u'| \geq 1} \sigma(t) u g(u') dx dt + c \int_S^T E \phi' \int_{\mathbb{R}^n} u'^2 dx dt \\ &\leq cE(S) + c \int_S^T E \phi' \int_{\mathbb{R}^n} u'^2 dx dt. \end{aligned} \tag{2.67}$$

Since $xg(x) \geq 0$ for all $x \in \mathbb{R}$, it follows that the energy is nonincreasing, locally absolutely continuous and $E'(t) = - \int_{\mathbb{R}^n} \sigma(t) u' g(u') dx$ a.c. in \mathbb{R}_+ . \square

Proof of (2.26). We consider the case $m = 1$, that is,

$$c_3 |v| \leq |g(v)| \leq c_4 |v| \quad \text{for all } |v| \leq 1. \tag{2.68}$$

Then we have

$$u'^2 \leq \frac{c_{13}}{\sigma(t)} u' \rho(t, u') \quad \forall t \in \mathbb{R}, \forall x \in \mathbb{R}^n, \tag{2.69}$$

where $\rho(t, s) = \sigma(t)g(s)$ for all $s \in \mathbb{R}$. Therefore we deduce from (2.67) (applied with $q = 0$) that

$$\int_S^T E(t) \phi'(t) dt \leq CE(S) + 2C \int_S^T \phi'(t) \int_{\mathbb{R}^n} \frac{1}{\sigma(t)} u' \rho(t, u') dx dt. \tag{2.70}$$

Define

$$\phi(t) = \int_0^t \sigma(\tau) d\tau. \tag{2.71}$$

It is clear that ϕ is a nondecreasing function of class C^2 on \mathbb{R}_+ . The hypothesis (2.23) ensures that

$$\phi(t) \longrightarrow +\infty \quad \text{as } t \longrightarrow +\infty. \tag{2.72}$$

Then we deduce from (2.70) that

$$\int_S^T E(t) \phi'(t) dt \leq CE(S) + 2C \int_S^T \int_{\mathbb{R}^n} u' \rho(t, u') dx dt \leq 3CE(S), \tag{2.73}$$

and thanks to Lemma 2.4 we obtain

$$E(t) \leq E(0) e^{(1-\phi(t))/(3C)}. \tag{2.74}$$

\square

Proof of (2.27). Now we assume that $m > 1$ in (2.2). Define ϕ by (2.71). We apply Lemma 2.4 with $q = (m - 1)/2$.

We need to estimate

$$\int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt. \tag{2.75}$$

For $t \geq 0$, consider

$$\Omega_1 = \{x \in \mathbb{R}^n, |u'| \leq 1\}, \quad \Omega_2 = \{x \in \mathbb{R}^n, |u'| > 1\}. \tag{2.76}$$

First we note that for every $t \geq 0$,

$$\Omega_1 \cup \Omega_2 = \mathbb{R}^n. \tag{2.77}$$

Next we deduce from (2.2) and (2.3) that for every $t \geq 0$,

- (i) if $x \in \Omega_1$, then $u'^2 \leq ((1/\sigma(t))u'\rho(t, u'))^{2/(m+1)}$,
- (ii) if $x \in \Omega_2$, then $u'^2 \leq (1/\sigma(t))u'\rho(t, u')$.

Hence, using Hölder’s inequality, we get that

$$\begin{aligned} & \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt \\ & \leq 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} \frac{1}{\sigma(t)} u' \rho(t, u') dx dt + 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} \left(\frac{1}{\sigma(t)} u' \rho(t, u') \right)^{2/(m+1)} dx dt \\ & \leq 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} \frac{1}{\sigma(t)} u' \rho(t, u') dx dt + 2 \int_S^T E^q \phi' \int_{\{|x| \leq (L+t)\}} (u' g(u'))^{2/(m+1)} dx dt \\ & \leq 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} \frac{1}{\sigma(t)} u' \rho(t, u') dx dt \\ & \quad + 2 \int_S^T E^q \phi' (1+t)^{n(m-1)/(m+1)} \left(\int_{\mathbb{R}^n} u' g(u') dx \right)^{2/(m+1)} dt \\ & \leq cE(S)^{1+q} + c' \int_S^T E^q \phi' (1+t)^{n(m-1)/(m+1)} \left(\frac{-E'}{\sigma(t)} \right)^{2/(m+1)} dt \\ & \leq cE(S)^{1+q} + c' \int_S^T E^q \phi' (1+t)^{n(m-1)/(m+1)} \sigma^{-2/(m+1)}(t) (-E')^{2/(m+1)} dt. \end{aligned} \tag{2.78}$$

Set $\varepsilon > 0$; thanks to Young’s inequality and to our definitions of p and ϕ , we obtain

$$\begin{aligned} & \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt \\ & \leq cE(S)^{1+q} + 2 \frac{m-1}{m+1} \varepsilon^{(m+1)/(m-1)} \int_S^T E^{1+q} (\phi')^{(m+1)/(m-1)} (1+t)^n \sigma^{-2/(m-1)} dt \\ & \quad + \frac{4}{m+1} \frac{1}{\varepsilon^{(m+1)/2}} E(S). \end{aligned} \tag{2.79}$$

We choose ϕ' such that

$$\phi'^{2/(m-1)} (1+t)^n \sigma^{-2/(m-1)} = 1, \tag{2.80}$$

so

$$\phi(t) = \int_0^t (1+s)^{-n(m-1)/2} \sigma(s) ds. \tag{2.81}$$

Then we deduce from (2.79) that

$$\int_S^T E^{1+q} \phi' dt \leq 2CE(S), \tag{2.82}$$

and thanks to Lemma 2.4 (applied with $c = 0$) we obtain

$$E(t) \leq \frac{C}{\phi(t)^{2/(m-1)}}. \tag{2.83}$$

□

Proof of Theorem 2.8. First, we see that if $u \in \widetilde{\mathcal{W}}^{**}$, then

$$\|\nabla_x u\|_2^2 + 2 \int_{\mathbb{R}^n} F(u(t)) dx \geq \|\nabla_x u\|_2^2 - \frac{2}{p+1} \|u(t)\|_{p+1}^{p+1} \geq \frac{p-1}{p+1} \|\nabla_x u\|_2^2. \tag{2.84}$$

In the proof, we often use the following inequality:

$$\|u(t)\|_2 \leq \frac{1}{\tilde{d}(t)} \|\lambda(x)u(t)\|_2. \tag{2.85}$$

Now, we assume that $I(u_0) > (1/2)\|\nabla_x u_0\|_2^2$. Then

$$I(u(t)) \geq \frac{1}{2} \|\nabla_x u(t)\|_2^2 \tag{2.86}$$

for some interval near $t = 0$. As long as (2.86) holds, we have $J(t) \equiv I(u(t))$. Thus

$$\begin{aligned} 2\eta_0 \int_S^T E^{q+1} \phi' dt &\leq - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T + \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt \\ &\quad + 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt - \int_S^T E^q \phi' \int_{\mathbb{R}^n} \sigma(t)ug(u') dx dt \\ &\leq - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T + \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt \\ &\quad + 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} u'^2 dx dt + c(\varepsilon) \int_S^T E^q \phi' \int_{|u'| \leq 1} \left(\frac{\sigma(t)}{\lambda(x)} \right)^2 g(u')^2 dx dt \\ &\quad + \varepsilon \int_S^T E^q \phi' \int_{\mathbb{R}^n} \lambda^2(x)u^2 dx dt + \int_S^T E^q \phi' \int_{|u'| \geq 1} \sigma(t)ug(u') dx dt. \end{aligned} \tag{2.87}$$

If $\sigma(t) = \mathcal{O}(\tilde{d}(t))$, that is, $\sigma(t) \rightarrow 0$ as $t \rightarrow \infty$ more rapidly than $\tilde{d}(t)$, we find the same results of asymptotic behaviour as in Theorem 2.7.

If $\tilde{d}(t) = \mathcal{O}(\sigma(t))$, so, we obtain

$$\begin{aligned} \int_S^T E^{q+1} \phi' dt &\leq - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T + \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt \\ &\int_S^T E^q \phi' \int_{|u'| \geq 1} u'^2 dx dt \int_S^T E^q \phi' \int_{|u'| \geq 1} \sigma(t) u g(u') dx dt \\ &+ c(\varepsilon) \int_S^T E^q \phi' \int_{|u'| \leq 1} \left(\frac{\sigma(t)}{\lambda(x)} \right)^2 |u'|^2 dx dt. \end{aligned} \tag{2.88}$$

We consider the case $m = 1$. Thus under (2.2) and (2.3), we have

$$\int_S^T E^q \phi' \int_{\mathbb{R}^n} \left(\frac{\sigma(t)}{\lambda(x)} \right)^2 u'^2 dx dt \leq C \int_S^T E^q \phi' \int_{\mathbb{R}^n} \left(\frac{\sigma^\alpha(t)}{\tilde{d}^2(t)} \right) \sigma(t) u' g(u') dx dt \tag{2.89}$$

for all $0 \leq \alpha < 1$. We choose

$$\phi(t) = \int_0^t \frac{\tilde{d}^2(s)}{\sigma^\alpha(s)} ds. \tag{2.90}$$

It is clear that ϕ is a nondecreasing function of class C^2 on \mathbb{R}_+ . Hypothesis (2.33) ensures that

$$\phi(t) \longrightarrow +\infty \quad \text{as } t \longrightarrow +\infty. \tag{2.91}$$

By (2.85), the definition of E , and the Cauchy-Schwartz inequality, we have

$$\begin{aligned} - \left[E^q \phi' \int_{\mathbb{R}^n} uu' dx \right]_S^T &= E^q(S) \phi'(S) \int_{\mathbb{R}^n} u(S) u'(S) dx - E^q(T) \phi'(T) \int_{\mathbb{R}^n} u(T) u'(T) dx \\ &\leq C E^{q+1}(S) \left[\frac{\phi'(S)}{\tilde{d}(S)} + \frac{\phi'(T)}{\tilde{d}(T)} \right] \leq C E^{q+1}(S), \\ \int_S^T (qE' E^{q-1} \phi' + E^q \phi'') \int_{\mathbb{R}^n} uu' dx dt &\leq \int_S^T q |E'| E^q \frac{\phi'(t)}{\tilde{d}(t)} dt + \int_S^T E^{q+1} \frac{|\phi''(t)|}{\tilde{d}(t)} dt \end{aligned} \tag{2.92}$$

when we have (in the case $m = 1$)

$$\phi' = \frac{\tilde{d}^2(t)}{\sigma^\alpha(t)}, \quad \phi''(t) = \frac{2\tilde{d}(t)\tilde{d}'(t)}{\sigma^\alpha(t)} - \alpha \frac{\sigma'(t)\tilde{d}^2(t)}{\sigma^{\alpha+1}(t)}. \tag{2.93}$$

So

$$\left| \frac{\phi''(t)}{\tilde{d}(t)} \right| \leq - \frac{2\tilde{d}'(t)\tilde{d}^\alpha(t)}{\tilde{d}^\alpha(t)\sigma^\alpha(t)} - \alpha \frac{\tilde{d}(t)\sigma'(t)}{\sigma(t)\sigma^\alpha(t)}, \tag{2.94}$$

$\tilde{d}(t)/\sigma(t)$ is bounded, so we obtain

$$\begin{aligned} \int_S^T E^{q+1} \frac{|\phi''(t)|}{\tilde{d}(t)} dt &\leq -E^{q+1}(S) [\tilde{d}^{1-\alpha}(t) + \sigma^{1-\alpha}(t)]_S^T \\ &\leq E^{q+1}(S) (\tilde{d}^{1-\alpha}(t) + \sigma^{1-\alpha}(t)) \\ &\leq CE^{q+1}(S). \end{aligned} \tag{2.95}$$

Then we deduce from (2.88) that

$$\int_S^T E\phi' dt \leq CE(S) + 2C \int_S^T \int_{\mathbb{R}^n} \sigma(t)u'g(u') dx dt \leq 3CE(S), \tag{2.96}$$

and thanks to Lemma 2.4 we obtain

$$E(t) \leq E(0)e^{(1-\phi(t))/3C}. \tag{2.97}$$

Using the condition that $\tilde{d}(t) = \mathbb{O}(\sigma(t))$ and using Hölder’s inequality, we get that

$$\begin{aligned} &\int_S^T E^q \phi' \int_{|u'| \leq 1} \frac{\sigma^2(t)}{\lambda^2(x)} u'^2 dx dt \\ &\leq 2 \int_S^T E^q \phi' \int_{\mathbb{R}^n} \frac{\sigma^2(t)}{\lambda^2(x)} \left(\frac{1}{\sigma(t)} \sigma(t)u'g(u') \right)^{2/(m+1)} dx dt \\ &\leq 2 \int_S^T E^q \phi' \int_{\{|x| \leq L+t\}} \frac{\sigma^2(t)}{\lambda^2(x)} \left(\frac{1}{\sigma(t)} \sigma(t)u'g(u') \right)^{2/(m+1)} dx dt \\ &\leq 2 \int_S^T E^q \phi' \left(\frac{\sigma(t)}{\tilde{d}(t)} \right)^2 (1+t)^{n(m-1)/(m+1)} \left(\int_{\mathbb{R}^n} u'g(u') dx \right)^{2/(m+1)} dt \\ &\leq c' \int_S^T E^q \phi' \left(\frac{\sigma(t)}{\tilde{d}(t)} \right)^2 (1+t)^{n(m-1)/(m+1)} \left(\frac{-E'}{\sigma(t)} \right)^{2/(m+1)} dt \\ &\leq c' \int_S^T E^q \phi' \left(\frac{\sigma(t)}{\tilde{d}(t)} \right)^2 (1+t)^{n(m-1)/(m+1)} \sigma^{-2/(m+1)}(t) (-E')^{2/(m+1)} dt \\ &\leq c' \int_S^T E^q \phi' \frac{\sigma^{\alpha+1}(t)}{\tilde{d}^2(t)} (1+t)^{n(m-1)/(m+1)} \sigma^{-2/(m+1)}(t) (-E')^{2/(m+1)} dt. \end{aligned} \tag{2.98}$$

Set $\varepsilon > 0$; thanks to Young’s inequality and to our definitions of p and ϕ , we obtain

$$\begin{aligned} &\int_S^T E^q \phi' \int_{\mathbb{R}^n} \left(\frac{\sigma(t)}{\lambda(x)} \right)^2 u'^2 dx dt \\ &\leq cE(S)^{1+q} + 2 \frac{m-1}{m+1} \varepsilon^{(m+1)/(m-1)} \\ &\quad \times \int_S^T E^{1+q} (\phi')^{(m+1)/(m-1)} \frac{\sigma^{(1+\alpha)(m+1)/(m-1)}(t)}{\tilde{d}^{2(m+1)/(m-1)}(t)} (1+t)^n \sigma^{-2/(m-1)}(t) dt + \frac{4}{m+1} \frac{1}{\varepsilon^{(m+1)/2}} E(S). \end{aligned} \tag{2.99}$$

We choose ϕ such that

$$\phi(t) = \int_0^t (1 + \tau)^{-n(m-1)/2} \sigma^{-((1+\alpha)(m+1)-2)/2}(\tau) \tilde{d}^{m+1}(\tau) d\tau. \tag{2.100}$$

It is clear that ϕ is a nondecreasing function of class C^2 on \mathbb{R}_+ . The hypothesis (2.35) ensures that $\phi(t) \rightarrow +\infty$ as $t \rightarrow +\infty$. By (2.85), the definition of E , and the Cauchy-Schwartz inequality we have (2.92) when we have (the case $m > 1$)

$$\begin{aligned} \phi' &= (1+t)^{-n(m-1)/2} \sigma^{-((1+\alpha)(m+1)-2)/2}(t) \tilde{d}^{m+1}(t), \\ \phi''(t) &= -\frac{n(m-1)}{2} (1+t)^{-n(m-1)/2-1} \sigma^{-((1+\alpha)(m+1)-2)/2}(t) \tilde{d}^{m+1}(t) \\ &\quad + (1+t)^{-n(m-1)/2} \left(-\frac{(1+\alpha)(m+1)-2}{2} \sigma^{-((1+\alpha)(m+1)-2)/2}(t) \sigma'(t) \tilde{d}^{m+1}(t) \right. \\ &\quad \left. + (m+1) \tilde{d}^m(t) \tilde{d}'(t) \sigma^{-((1+\alpha)(m+1)-2)/2}(t) \right). \end{aligned} \tag{2.101}$$

Thus

$$\begin{aligned} \left| \frac{\phi''(t)}{\tilde{d}(t)} \right| &\leq C \frac{\tilde{d}^m(t)}{\sigma^m(t)} \sigma^{(1-\alpha)(m+1)/2}(t) - C' \frac{\tilde{d}^m(t)}{\sigma^m(t)} \frac{\sigma'(t)}{\sigma^{((1+\alpha)(m+1)-2m)/2}(t)} \\ &\quad - C'' \frac{\tilde{d}^{((1+\alpha)(m+1)-2)/2}(t)}{\sigma^{((1+\alpha)(m+1)-2)/2}(t)} \frac{\tilde{d}'(t)}{\tilde{d}^{((1+\alpha)(m+1)-2m)/2}(t)}, \end{aligned} \tag{2.102}$$

$\tilde{d}(t)/\sigma(t)$ is bounded, so we obtain

$$\begin{aligned} \int_S^T E^{q+1} \frac{|\phi''(t)|}{\tilde{d}(t)} dt &\leq -E^{q+1}(S) \left[\tilde{d}^{(1-\alpha)(m+1)/2}(t) + \sigma^{(1-\alpha)(m+1)/2}(t) \right]_S^T \\ &\leq E^{q+1}(S) (\tilde{d}^{(1-\alpha)(m+1)/2}(S) + \sigma^{(1-\alpha)(m+1)/2}(S)) \\ &\leq CE^{q+1}(S). \end{aligned} \tag{2.103}$$

We deduce from this choice

$$\int_S^T E^{1+q} \phi' dt \leq 2CE(S), \tag{2.104}$$

and thanks to Lemma 2.4 (applied with $c = 0$), we obtain

$$E(t) \leq \frac{C(E(0))}{\phi^{2/(m-1)}}. \tag{2.105}$$

Since $u_0 \in \widetilde{W}^{**}$ and \widetilde{W}^{**} is an open set, putting

$$T_1 = \sup \{t \in [0, +\infty) : u(s) \in \widetilde{W}^{**} \text{ for } 0 \leq s \leq t\}, \tag{2.106}$$

we see that $T_1 > 0$ and $u(t) \in \widetilde{W}^{**}$ for $0 \leq t < T_1$. If $T_1 < T_{\max} < \infty$, where T_{\max} is the lifespan of the solution, then $u(T_1) \in \partial \widetilde{W}^{**}$; that is,

$$I(u(T_1)) = 0, \quad u(T_1) \neq 0. \tag{2.107}$$

We see from Lemma 2.2 and (2.85) that

$$\begin{aligned} \|u(t)\|_{p+1}^{p+1} &\leq C \|u(t)\|_2^{(4-(n-2)(p-1))/2} \|\nabla_x u(t)\|_2^{n(p-1)/2} \\ &\leq (\tilde{d}(t))^{-(4-(n-2)(p-1))/2} E^{(p-1)/2} \|\nabla_x u(t)\|_2^2 \\ &\leq B(t) \|\nabla_x u(t)\|_2^2 \end{aligned} \tag{2.108}$$

for $0 \leq t \leq T_1$, where we have used the assumption $p \geq (n+4)/n$ and

$$B(t) = \frac{C(E(0))\tilde{d}^{-(4-(n-2)(p-1))/2}(t)}{\left(\int_0^t (1+\tau)^{-n(m-1)/2} \sigma^{-((1+\alpha)(m+1)-2)/2}(\tau) \tilde{d}^{m+1}(\tau) d\tau\right)^{(p-1)/(m-1)}}. \tag{2.109}$$

Next, we put

$$T_2 \equiv \sup \left\{ t \in [0, +\infty) : B(s) < \frac{1}{2} \text{ for } 0 \leq s < t \right\}, \tag{2.110}$$

and then we see that $T_2 > 0$ and $T_2 = T_1$ because $B(t) < 1/2$ by the condition that $E(0)$ is small. Then

$$I(u(t)) \geq \|\nabla_x u(t)\|_2^2 - B(t) \|\nabla_x u(t)\|_2^2 \geq \frac{1}{2} \|\nabla_x u(t)\|_2^2 \tag{2.111}$$

for $0 \leq t \leq T_1$. Moreover, (2.107) and (2.111) imply

$$K(u(T_1)) \geq \frac{1}{2} \|\nabla_x u(T_1)\|_2^2 > 0, \tag{2.112}$$

which is a contradiction, and hence, it might be $T_1 = T_{\max}$. Therefore, (2.105) holds true for $0 \leq T \leq T_{\max}$. To prove global existence in $H^2 \cap H^1$, we need to derive the estimates for second derivatives of $u(t)$ on the basis of the energy estimate of $E(t)$, we utilize the differentiated equation

$$u_{ttt} - \Delta_x u' + \lambda^2(x)u' + \sigma'(t)g(u') + \sigma(t)g(u')u'' + f'(u)u' = 0, \tag{2.113}$$

where $f(u) = |u|^{p-1}u$. Multiplying (2.113) by u'' , we have

$$\begin{aligned} \frac{d}{dt} E_2(t) + 2\sigma(t) \int_{\mathbb{R}^n} g'(u') |u''(t)|^2 dx \\ \leq 2 \int_{\mathbb{R}^n} |f'(u)| |u'(t)| |u''(t)| dx + 2 |\sigma'(t)| \int_{\mathbb{R}^n} |g(u')| |u''(t)| dx, \end{aligned} \tag{2.114}$$

where we set

$$E_2(t) = \|u''(t)\|_2^2 + \|\nabla_x u(t)\|_2^2 + \|\lambda u'(t)\|_2^2. \tag{2.115}$$

By (2.2) and (2.3), we have

$$\begin{aligned} \int_{\mathbb{R}^n} |g(u')|^2 dx &\leq C \int_{|u'| \leq 1} |u'|^{2/m} dx + C' \int_{|u'| \geq 1} |u'|^{2r} dx \\ &\leq C(L+t)^{n(m-1)/m} E^{1/m} + C' E^{2-(n-2)(r-1)/2} E_2^{n(r-1)/2} \end{aligned} \tag{2.116}$$

and by [Lemma 2.2](#)

$$\begin{aligned}
 2 \int_{\mathbb{R}^n} |f'(u)| |u'(t)| |u''(t)| \, dx &\leq C \|u(t)\|_{n(p-1)}^{p-1} \|u'(t)\|_{2n/(n-2)} \|u''(t)\|_2 \\
 &\leq c \|u(t)\|_{n(p-1)}^{p-1} \|\nabla_x u'(t)\|_2 \|u''(t)\|_2 \\
 &\leq c \|u(t)\|_{n(p-1)}^{p-1} E_2(t).
 \end{aligned}
 \tag{2.117}$$

Since $(n + 2)/n \leq p \leq n/(n - 2)$, then

$$\|u(t)\|_{n(p-1)}^{p-1} \leq \tilde{d}^{-(2-(n-2)(p-1))/2} E^{(p-1)/2}.
 \tag{2.118}$$

Thus, we have

$$\begin{aligned}
 \frac{d}{dt} E_2 &\leq \tilde{d}^{-(2-(n-2)(p-1))/2} E^{(p-1)/2} E_2(t) \\
 &\quad + 2 |\sigma'(t)| ((L + t)^{n(m-1)/m} E^{1/m} + CE^{(2-(n-2)(r-1))/2} E_2^{n(r-1)/2})^{1/2} E_2^{1/2}(t).
 \end{aligned}
 \tag{2.119}$$

We have also, applying Young inequality,

$$E^{(2-(n-2)(r-1))/2} E_2^{n(r-1)/2} \leq CE^{(2-(n-2)(r-1))/(2-n(r-1))}(t) + E_2(t),
 \tag{2.120}$$

hence, we deduce that

$$\frac{d}{dt} E_2(t) \leq C(t)(1 + E_2(t)).
 \tag{2.121}$$

So, we obtain

$$E_2(t) \leq C' e^{\int_0^t C(s) ds}.
 \tag{2.122}$$

From [\(2.122\)](#) and the first equation of problem [\(1.1\)](#), we also prove easily that

$$\|\Delta_x u(t)\|_2 \leq C'(t) < \infty
 \tag{2.123}$$

for all $t \geq 0$. Indeed, we have

$$\|\Delta_x u(t)\|_2 \leq \|u''(t)\|_2 + \|\lambda^2 u(t)\|_2 + \sigma(t) \|g(u'(t))\|_2 + \|f(u)\|_2
 \tag{2.124}$$

and also we have

$$\|f(u)\|_2^2 \leq C \|u(t)\|_{2p}^{2p} \leq C \|u(t)\|_{n(p-1)/2}^{2(p-1)} \|\Delta_x u(t)\|_2^2.
 \tag{2.125}$$

Here, to check the last inequality of [\(2.125\)](#), we note that if $(n + 4)/n \leq p \leq (n + 2)/(n - 2)$, then

$$\begin{aligned}
 C \|u(t)\|_{n(p-1)/2}^{2(p-1)} &\leq C \|u(t)\|_2^{4-(n-2)(p-1)} \|\nabla_x u(t)\|_2^{n(p-1)-4} \\
 &\leq (\tilde{d}(t))^{4-(n-2)(p-1)} E^{p-1}(t) < \frac{1}{2}.
 \end{aligned}
 \tag{2.126}$$

Thus the solution in the sense of [Theorem 2.6](#) exists globally in time t under the assumption $(u_0, u_1) \in H^2 \times H^1$.

When $(u_0, u_1) \in H^1 \times L^2$ we approximate (u_0, u_1) by $(u_0^k, u_1^k) \in H^2 \times H^1$, $k = 1, 2, \dots$, in the topologies of $H^1 \times L^2$, which satisfy

$$\text{supp } u_0^k \cap \text{supp } u_1^k \subset \{x \in \mathbb{R}^n \mid |x| \leq L\}. \quad (2.127)$$

Since $\lim_{k \rightarrow \infty} (u_0^k, u_1^k) = (u_0, u_1)$ in $H^1 \times L^2$, then for these initial data problem (1.1) has global solutions $u^k \in W_{\text{loc}}^{2, \infty}([0, \infty); L^2) \cap W_{\text{loc}}^{1, \infty}([0, \infty); H^1) \cap L_{\text{loc}}^\infty([0, \infty); H^2)$, which satisfy (2.122) and (2.123). We can easily see that $\{u^k(t)\}$ converges uniformly on each compact interval $[0, T]$, $T > 0$. Uniqueness follow from a standard argument. The proof of [Theorem 2.8](#) is now completed. \square

References

- [1] A. Benaissa and S. A. Messaoudi, *Blow-up of solutions for the Kirchhoff equation of q -Laplacian type with nonlinear dissipation*, Colloq. Math. **94** (2002), no. 1, 103–109.
- [2] A. Benaissa and L. Rahmani, *Global existence and energy decay of solutions for Kirchhoff-Carrier equations with weakly nonlinear dissipation*, Bull. Belg. Math. Soc. Simon Stevin **11** (2004), no. 4, 547–574.
- [3] P. Brenner, *On space-time means and strong global solutions of nonlinear hyperbolic equations*, Math. Z. **201** (1989), no. 1, 45–55.
- [4] V. Georgiev and G. Todorova, *Existence of a solution of the wave equation with nonlinear damping and source terms*, J. Differential Equations **109** (1994), no. 2, 295–308.
- [5] J. Ginibre and G. Velo, *The global Cauchy problem for the nonlinear Klein-Gordon equation*, Math. Z. **189** (1985), no. 4, 487–505.
- [6] A. Haraux, *Two remarks on hyperbolic dissipative problems*, Nonlinear Partial Differential Equations and Their Applications. Collège de France Seminar, Vol. VII (Paris, 1983–1984), Res. Notes in Math., vol. 122, Pitman, Massachusetts, 1985, pp. 161–179.
- [7] R. Ikehata, *Some remarks on the wave equations with nonlinear damping and source terms*, Nonlinear Anal. **27** (1996), no. 10, 1165–1175.
- [8] R. Ikehata and T. Suzuki, *Stable and unstable sets for evolution equations of parabolic and hyperbolic type*, Hiroshima Math. J. **26** (1996), no. 3, 475–491.
- [9] F. John, *Nonlinear Wave Equations, Formation of Singularities*, University Lecture Series, vol. 2, American Mathematical Society, Rhode Island, 1990.
- [10] V. Komornik, *Exact Controllability and Stabilization*, RAM: Research in Applied Mathematics, Masson, Paris, 1994.
- [11] H. A. Levine, P. Pucci, and J. Serrin, *Some remarks on global nonexistence for nonautonomous abstract evolution equations*, Harmonic Analysis and Nonlinear Differential Equations (Riverside, Calif, 1995), Contemp. Math., vol. 208, American Mathematical Society, Rhode Island, 1997, pp. 253–263.
- [12] H. A. Levine and J. Serrin, *Global nonexistence theorems for quasilinear evolution equations with dissipation*, Arch. Rational Mech. Anal. **137** (1997), no. 4, 341–361.
- [13] P. Martinez, *A new method to obtain decay rate estimates for dissipative systems*, ESAIM Control Optim. Calc. Var. **4** (1999), 419–444.
- [14] M. Nakao, *A difference inequality and its application to nonlinear evolution equations*, J. Math. Soc. Japan **30** (1978), no. 4, 747–762.
- [15] M. Nakao and K. Ono, *Existence of global solutions to the Cauchy problem for the semilinear dissipative wave equations*, Math. Z. **214** (1993), no. 2, 325–342.

- [16] ———, *Global existence to the Cauchy problem of the semilinear wave equation with a nonlinear dissipation*, Funkcial. Ekvac. **38** (1995), no. 3, 417–431.
- [17] K. Ono, *On global solutions and blow-up solutions of nonlinear Kirchhoff strings with nonlinear dissipation*, J. Math. Anal. Appl. **216** (1997), no. 1, 321–342.
- [18] D. H. Sattinger, *On global solution of nonlinear hyperbolic equations*, Arch. Rational Mech. Anal. **30** (1968), 148–172.
- [19] W. A. Strauss, *On continuity of functions with values in various Banach spaces*, Pacific J. Math. **19** (1966), 543–551.
- [20] ———, *Nonlinear Wave Equations*, CBMS Regional Conference Series in Mathematics, vol. 73, Conference Board of the Mathematical Sciences, Washington, D.C., 1989.
- [21] G. Todorova, *Stable and unstable sets for the Cauchy problem for a nonlinear wave equation with nonlinear damping and source terms*, C. R. Acad. Sci. Paris Sér. I Math. **328** (1999), no. 2, 117–122.

Abbès Benaïssa: Département de Mathématiques, Faculté des Sciences, Université Djillali Liabès,
B. P. 89, Sidi Bel Abbès 22000, Algeria
E-mail address: benaïssa_abbes@yahoo.com

Soufiane Mokeddem: Département de Mathématiques, Faculté des Sciences, Université Djillali
Liabès, B. P. 89, Sidi Bel Abbès 22000, Algeria
E-mail address: mok5_so5@yahoo.fr