

SIMULATION OF A SPACE–TIME BOUNDED DIFFUSION

BY G. N. MILSTEIN AND M. V. TRETYAKOV¹

Weierstraß-Institut and Ural State University

Mean-square approximations, which ensure boundedness of both time and space increments, are constructed for stochastic differential equations in a bounded domain. The proposed algorithms are based on a space–time discretization using a random walk over boundaries of small space–time parallelepipeds. To realize the algorithms, exact distributions for exit points of the space–time Brownian motion from a space–time parallelepiped are given. Convergence theorems are stated for the proposed algorithms. A method of approximate searching for exit points of the space–time diffusion from the bounded domain is constructed. Results of several numerical tests are presented.

1. Introduction. The paper is devoted to mean-square approximations for a system of stochastic differential equations (SDE),

$$(1.1) \quad dX = \chi_{\tau_{t,x} > s} b(s, X) ds + \chi_{\tau_{t,x} > s} \sigma(s, X) dw(s), \quad X(t) = X_{t,x}(t) = x,$$

in a space–time bounded domain $Q = [t_0, t_1] \times G \subset R^{d+1}$. Here X and b are d -dimensional vectors, σ is a $d \times d$ -matrix, $(w(s), \mathcal{F}_s)$, $s \geq t_0$, is a d -dimensional standard Wiener process defined on a probability space (Ω, \mathcal{F}, P) , G is a bounded open domain in R^d , and the Markov moment $\tau_{t,x}$ is the first-passage time of the process $(s, X_{t,x}(s))$, $s \geq t$, to $\Gamma = \overline{Q} \setminus Q$. The set Γ is a part of the boundary ∂Q consisting of the lateral surface and the upper base of the cylinder \overline{Q} . We put $X_{t,x}(s) = X_{t,x}(\tau_{t,x})$ under $s \geq \tau_{t,x}$, and thus, the process $(s, X_{t,x}(s))$ is defined for all $t \leq s < t_1$. The coefficients $b^i(s, x)$ and $\sigma^{ij}(s, x)$, $(s, x) \in \overline{Q}$, and the boundary ∂G are assumed to be sufficiently smooth, while the strict ellipticity condition is imposed on the matrix $a(s, x) := \sigma(s, x)\sigma^T(s, x)$.

The first numerical method concerning simulation of a diffusion process in a bounded domain is constructed in [25]. The method is based on a random walk over touching spheres and applied to solving the Dirichlet problem for elliptic equations with constant coefficients by a Monte Carlo technique.

Probabilistic methods for solving boundary value problems, which involve the weak sense numerical integration of ordinary SDE, are the main subject of the works [19, 20, 23]. These methods ensure that the proposed weak approximations belong to the bounded domain associated with a considered boundary value problem. Some other probabilistic approaches are also available in [5, 8, 16, 29].

Received September 1997; revised May 1998.

¹Supported in part by the Alexander von Humboldt Foundation with a research fellowship.

AMS 1991 subject classifications. Primary 60H10; secondary 60J15, 65U05.

Key words and phrases. Space–time Brownian motion, random walk, mean-square approximation, the Dirichlet problem for equations of parabolic and elliptic type.

A mean-square approximation for simulation of an autonomous diffusion process in a space bounded domain is considered in [22, 24]. The algorithm is based on a space discretization (quantization) using a random walk over small spheres. It gives the points which are close in the mean-square sense to the points of the real phase trajectory for SDE in the space bounded domain. To realize the algorithm, the exit point of the Wiener process from a d -dimensional ball has to be constructed at each step. Due to independence of the first exit time and the first exit point of the Wiener process from the ball, it is possible to simulate them separately. It is known, that the exit point is distributed uniformly on the sphere, but simulation of the exit time is a fairly laborious problem. Consequently, the algorithm gives only the phase component of the approximate trajectory without modelling the corresponding time component like the algorithm over touching spheres [25]. The space-time point lies on the d -dimensional lateral surface of a semicylinder with sphere base in the $(d + 1)$ -dimensional semispace $[t_0, \infty) \times R^d$. The algorithm ensures smallness of the phase increments at each step, but the nonsimulated time increments can take arbitrary large values with some probability.

As is well known, “ordinary” mean-square methods (see, e.g., [14, 18, 26]), intended to solve SDE on a finite time interval, are based on a time discretization (sampling). The space-time point, corresponding to an “ordinary” one-step approximation constructed at a time point t_k , lies on the d -dimensional plane $t = t_k$, which belongs to the $(d + 1)$ -dimensional semispace $[t_0, \infty) \times R^d$. The “ordinary” mean-square methods give both time and phase components of the approximate trajectory. They ensure smallness of time increments at each step, but space increments can take arbitrary large values with some probability.

The mean-square approximations, which are the subject of the present paper, control boundedness of both space increments and time increments at each step. In addition, they give approximate values for both phase and time components of the space-time diffusion in the space-time bounded domain Q . It is possible to solve this problem in a constructive manner by the implementation of a space-time discretization by a random walk over boundaries of small space-time parallelepipeds. It turns out that the first exit point $(\bar{\theta}, w(\bar{\theta}))$ of the space-time Brownian motion $(s, w(s))$, $s > 0$, from the space-time parallelepiped $\Pi_r = [0, lr^2) \times C_r$, $C_r \subset R^d$ is a cube with center at the origin and edge length equal to $2r$, can be simulated in a sufficiently easy way (some aspects of the space-time Brownian motion under $d = 1$ are considered in [11]). To construct a one-step approximation, we introduce the system with frozen coefficients (both t, x fixed),

$$(1.2) \quad d\bar{X} = b(t, x) ds + \sigma(t, x) dw(s), \quad \bar{X}(t) = x.$$

As an approximation of the point $(t + \bar{\theta}, X_{t,x}(t + \bar{\theta}))$ of the space-time diffusion $(s, X_{t,x}(s))$, $s \geq t$, we take the point $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$, where $\bar{X}_{t,x}(t + \bar{\theta})$ is a solution of (1.2):

$$(1.3) \quad \bar{X}_{t,x}(t + \bar{\theta}) = x + b(t, x)\bar{\theta} + \sigma(t, x)(w(t + \bar{\theta}) - w(t)),$$

and $(\bar{\theta}, w(t + \bar{\theta}) - w(t))$ is the exit point of the space–time Brownian motion $(s - t, w(s) - w(t))$, $s > t$, from the space–time parallelepiped Π_r .

The point $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$ lies on the lateral surface or on the upper base of a certain parallelepiped obtained from Π_r by a linear transformation, that is, it is constructed on a bounded d -dimensional manifold in contrast to the “ordinary” mean-square approximations and to the approximations of [22, 24], which are constructed on the d -dimensional unbounded manifolds.

On the basis of the one-step approximation (1.3), we form a Markov chain $(\bar{\vartheta}_k, \bar{X}_k)$ which belongs to Q at each step and approximates the points $(\bar{\vartheta}_k, X(\bar{\vartheta}_k))$ of the trajectory $(s, X_{t,x}(s))$, $s \geq t$, in the mean-square sense.

In the paper we present some constructive methods of simulating the Markov chain $(\bar{\vartheta}_k, \bar{X}_k)$, state some convergence theorems, propose a method of approximate searching for exit points of the space–time diffusion from a bounded domain and indicate possible applications.

2. Auxiliary knowledge. Let G be a bounded domain in R^d , $Q = [t_0, t_1] \times G$ be a cylinder in R^{d+1} , $\Gamma = \bar{Q} \setminus Q$. The set Γ is a part of the boundary of the cylinder Q consisting of the upper base and the lateral surface.

Consider the first boundary value problem for the equation of parabolic type

$$(2.1) \quad \begin{aligned} \frac{\partial u}{\partial t} + \frac{1}{2} \sum_{i,j=1}^d a^{ij}(t, x) \frac{\partial^2 u}{\partial x^i \partial x^j} \\ + \sum_{i=1}^d b^i(t, x) \frac{\partial u}{\partial x^i} + c(t, x)u + e(t, x) = 0, \quad (t, x) \in Q, \end{aligned}$$

with the initial condition on the upper base

$$(2.2) \quad u(t_1, x) = f(x), \quad x \in \bar{G},$$

and the boundary condition on the lateral surface

$$(2.3) \quad u(t, x) = g(t, x), \quad t_0 \leq t \leq t_1, \quad x \in \partial G.$$

Introduce the function φ defined on Γ such that it is equal to $f(x)$ on the upper base and it is equal to $g(t, x)$ on the lateral surface. Then the conditions (2.2) and (2.3) may be rewritten shortly as

$$(2.4) \quad u|_{\Gamma} = \varphi.$$

The coefficients $a^{ij} = a^{ji}$ are assumed to satisfy the property of strong ellipticity in \bar{Q} , that is,

$$\lambda_1^2 = \min_{(t,x) \in \bar{Q}} \min_{1 \leq i \leq d} \lambda_i^2(t, x) > 0,$$

where $\lambda_1^2(t, x) \leq \lambda_2^2(t, x) \leq \dots \leq \lambda_d^2(t, x)$ are eigenvalues of the matrix $a(t, x) = \{a^{ij}(t, x)\}$.

Let $\lambda_d^2 = \max_{(t,x) \in \bar{Q}} \lambda_d^2(t, x)$. Then for any $(t, x) \in \bar{Q}$ and $y \in \mathbf{R}^d$ the following inequality:

$$(2.5) \quad \lambda_1^2 \sum_{i=1}^d y^{i^2} \leq \sum_{i,j=1}^d a^{ij}(t, x) y^i y^j \leq \lambda_d^2 \sum_{i=1}^d y^{i^2}$$

holds.

The solution to the problem (2.1), (2.4) has the following probabilistic representation ([6], [9], page 299):

$$(2.6) \quad u(t, x) = E[\varphi(\tau, X_{t,x}(\tau))Y_{t,x,1}(\tau) + Z_{t,x,1,0}(\tau)],$$

where $X_{t,x}(s)$, $Y_{t,x,y}(s)$, $Z_{t,x,y,z}(s)$, $s \geq t$, is the solution of the Cauchy problem to the following system of stochastic differential equations:

$$(2.7) \quad \begin{aligned} dX &= b(s, X) ds + \sigma(s, X) dw(s), & X(t) &= x, \\ dY &= c(s, X) ds, & Y(t) &= y, \\ dZ &= e(s, X) Y ds, & Z(t) &= z. \end{aligned}$$

Here the point (t, x) belongs to Q , $\tau = \tau_{t,x}$ is the first-passage time of the trajectory $(s, X_{t,x}(s))$ to the boundary Γ . In the system (2.7), Y and Z are scalars, $w(s) = (w^1(s), \dots, w^d(s))^T$ is a d -dimensional standard Wiener process, $b(s, x)$ is a column vector of dimension d compounded from the coefficients $b^i(s, x)$, $\sigma(s, x)$ is a matrix of dimension $d \times d$ which is received from the equation

$$(2.8) \quad \sigma(s, x)\sigma^T(s, x) = a(s, x), \quad a(s, x) = \{a^{ij}(s, x)\}.$$

Setting in (2.1), (2.7)

$$(2.9) \quad c = 0, \quad e = 0, \quad f = 0, \quad g = \chi_{(\partial G)_0}(x),$$

where $(\partial G)_0 \subseteq \partial G$, we get the following formula:

$$(2.10) \quad u(t, x) = P(\tau_{t,x} < t_1, X_{t,x}(\tau_{t,x}) \in (\partial G)_0), \quad t_0 \leq t < t_1,$$

where the time $\tau_{t,x}$ is the first-passage time of the trajectory $X_{t,x}(s)$ to the boundary ∂G .

In particular, if

$$(2.11) \quad c = 0, \quad e = 0, \quad f = 0, \quad g = 1,$$

then

$$(2.12) \quad u(t, x) = P(\tau_{t,x} < t_1), \quad t_0 \leq t < t_1.$$

Setting in (2.1), (2.7),

$$(2.13) \quad c = 0, \quad e = 0, \quad f = \chi_{G_0}(x), \quad g = 0,$$

where $G_0 \subset G$, we get the following formula:

$$(2.14) \quad u(t, x) = P(\tau_{t,x} \geq t_1, X_{t,x}(t_1) \in G_0).$$

In autonomous cases (i.e., a^{ij}, b^i, c, e, g do not depend on t) we shall consider the first boundary value problem for parabolic equations in the following form:

$$(2.15) \quad \frac{\partial u}{\partial t} = \frac{1}{2} \sum_{i,j=1}^d a^{ij}(x) \frac{\partial^2 u}{\partial x^i \partial x^j} + \sum_{i=1}^d b^i(x) \frac{\partial u}{\partial x^i} + c(x)u + e(x), \quad t > 0, \quad x \in G,$$

$$(2.16) \quad u(0, x) = f(x), \quad x \in \overline{G},$$

$$(2.17) \quad u(t, x) = g(x), \quad t > 0, \quad x \in \partial G.$$

Using (2.9) and (2.10) and (2.13) and (2.14), it is not difficult to obtain that the function

$$(2.18) \quad u(t, x) = P(\tau_{0,x} < t, X_{0,x}(\tau_{0,x}) \in (\partial G)_0), \quad t > 0,$$

is the solution of the problem (2.15)–(2.17) under (2.9); the function

$$(2.19) \quad u(t, x) = P(\tau_{0,x} < t), \quad t > 0,$$

is the solution of the problem (2.15)–(2.17) under (2.11); the function

$$(2.20) \quad u(t, x) = P(\tau_{0,x} \geq t, X_{0,x}(t) \in G_0)$$

is the solution of the problem (2.15)–(2.17) under (2.13).

Here $X_{0,x}(s)$ is the solution to the Cauchy problem,

$$(2.21) \quad dX = b(X) ds + \sigma(X) dw(s), \quad X(0) = x,$$

and $\tau_{0,x}$ is the first-passage time of the trajectory $X_{0,x}(s)$ to the boundary ∂G .

3. Some distributions for a one-dimensional Wiener process. A part of distributions for the Wiener process which we give in this paper (see Sections 3, 4, 9) may be found in the literature. For instance, in [4, 7, 11] some distributions for the one-dimensional Wiener process are written down in a certain form. But we do not know whether all the distributions needed for our goals are available in the literature. Moreover, we need various analytical forms of one and the same distribution due to computational aspects. That is why, for completeness of the exposition, we derive all the distributions here and give them in the forms, which are suitable for practical realization.

Introduce the first-passage time $\tau_x := \tau_{0,x}$ of the one-dimensional Wiener process $x + W(t)$, $-1 \leq x \leq 1$, $t > 0$, to the boundary of the interval $[-1, 1]$. Derive the formulas for

$$u(t, x) = P(\tau_x < t).$$

From (2.15)–(2.17) under (2.11), we obtain that the function [see (2.19)]

$$v(t, x) = u(t, x) - 1 = P(\tau_x < t) - 1$$

satisfies the following boundary value problem:

$$(3.1) \quad \frac{\partial v}{\partial t} = \frac{1}{2} \frac{\partial^2 v}{\partial x^2}, \quad t > 0, \quad -1 < x < 1,$$

$$(3.2) \quad v(0, x) = -1, \quad v(t, -1) = v(t, 1) = 0.$$

By the method of separation of variables, we get the following formula:

$$(3.3) \quad P(\tau_x < t) = 1 - \frac{4}{\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} \cos \frac{\pi(2k+1)x}{2} \exp\left(-\frac{1}{8} \pi^2(2k+1)^2 t\right).$$

Further, extending the initial data in (3.1) and (3.2) by the odd way on the whole axis and solving the obtained Cauchy problem, we get another form for the same distribution

$$(3.4) \quad P(\tau_x < t) = 1 - \int_{-1}^1 G(t, x, y) dy,$$

where

$$(3.5) \quad G(t, x, y) = \frac{1}{\sqrt{2\pi t}} \sum_{k=-\infty}^{\infty} \left(\exp\left(-\frac{1}{2t}(x-4k-y)^2\right) - \exp\left(-\frac{1}{2t}(x-(4k+2)+y)^2\right) \right).$$

We shall use (3.3) and (3.4) under $x = 0$. Denote $\tau = \tau_0$,

$$\mathcal{P}(t) := P(\tau < t),$$

and introduce the density $\mathcal{P}'(t)$. From (3.3) and (3.4) one can obtain the following lemma.

LEMMA 3.1. *Let τ be the first-passage time of the one-dimensional standard Wiener process $W(t)$ to the boundary of the interval $[-1, 1]$. Then the following formulas for its distribution and density take place:*

$$(3.6) \quad \mathcal{P}(t) = 1 - \frac{4}{\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} \exp\left(-\frac{1}{8} \pi^2(2k+1)^2 t\right), \quad t > 0$$

and

$$(3.7) \quad \mathcal{P}(t) = 2 \sum_{k=0}^{\infty} (-1)^k \operatorname{erfc} \frac{2k+1}{\sqrt{2t}}, \quad t > 0,$$

$$(3.8) \quad \mathcal{P}'(t) = \frac{\pi}{2} \sum_{k=0}^{\infty} (-1)^k (2k+1) \exp\left(-\frac{1}{8} \pi^2(2k+1)^2 t\right), \quad t > 0$$

and

$$(3.9) \quad \mathcal{P}'(t) = \frac{2}{\sqrt{2\pi t^3}} \sum_{k=0}^{\infty} (-1)^k (2k+1) \exp\left(-\frac{1}{2t}(2k+1)^2\right), \quad t > 0.$$

Remember

$$\operatorname{erfc} x = \frac{2}{\sqrt{\pi}} \int_x^\infty \exp(-s^2) ds, \quad \operatorname{erfc} 0 = 1.$$

Formulas (3.6) and (3.8) are suitable for calculations under large t , and (3.7) and (3.9) are suitable under small t . The remainders of the series (3.8) and (3.9) are evaluated by the quantities

$$r_k(t) = \frac{\pi}{2}(2k + 3) \exp\left(-\frac{1}{8} \pi^2(2k + 3)^2 t\right)$$

and

$$\rho_k(t) = \frac{2}{\sqrt{2\pi t^3}}(2k + 3) \exp\left(-\frac{1}{2t}(2k + 3)^2\right)$$

correspondingly.

These quantities coincide under $t = 2/\pi$ and

$$\begin{aligned} r_k(t) &< r_k\left(\frac{2}{\pi}\right), & t > \frac{2}{\pi}, \\ \rho_k(t) &< r_k\left(\frac{2}{\pi}\right), & t < \frac{2}{\pi}. \end{aligned}$$

If we take k , for example, equal to 2, then

$$r_2\left(\frac{2}{\pi}\right) = \frac{7\pi}{2} \exp(-49\pi/4) < 2.13 \cdot 10^{-16},$$

and consequently,

$$\bar{\mathcal{P}}'(t) = \begin{cases} \frac{2}{\sqrt{2\pi t^3}}(\exp(-1/2t) - 3 \exp(-9/2t) + 5 \exp(-25/2t)), & 0 < t < \frac{2}{\pi}, \\ \frac{\pi}{2}(\exp(-\pi^2 t/8) - 3 \exp(-9\pi^2 t/8) + 5 \exp(-25\pi^2 t/8)), & t > \frac{2}{\pi}, \end{cases}$$

differs from $\bar{\mathcal{P}}'(t)$ by a quantity of $2.13 \cdot 10^{-16}$ on the whole interval $[0, \infty)$.

It is not difficult to evaluate that

$$\bar{\mathcal{P}}(t) = \int_0^t \bar{\mathcal{P}}'(s) ds$$

differs from $\mathcal{P}(t)$ on the whole interval $[0, \infty)$ by $(8/7\pi) \exp(-49\pi/4) < 7.04 \cdot 10^{-18}$. Such an exactness is quite sufficient for practical calculations. See the curves of the distribution $\mathcal{P}(t)$ and its density $\mathcal{P}'(t)$ in Figure 1.




FIG. 1. Distribution function $\mathcal{P}(t)$ and density $\mathcal{P}'(t)$.

Denote the inverse function to \mathcal{P} by \mathcal{P}^{-1} , and let γ be a random variable uniformly distributed on $[0, 1]$. Then the random variable

$$\tau = \mathcal{P}^{-1}(\gamma)$$

is distributed by the law $\mathcal{P}(t)$.

To simulate this law in practice, we have to solve the following equation:

$$(3.10) \quad \bar{\mathcal{P}}(t) = \gamma.$$

Let us note that because of the analytical simplicity of the function $\bar{\mathcal{P}}(t)$, it is natural to use the Newton method for solving (3.10).

LEMMA 3.2. For the conditional probability

$$\mathcal{Q}(\beta; t) := P(W(t) < \beta / |W(s)| < 1, 0 < s < t),$$

where $-1 < \beta \leq 1$, the following equalities hold:

$$(3.11) \quad \begin{aligned} \mathcal{Q}(\beta; t) &= \frac{P(W(t) < \beta, \tau \geq t)}{P(\tau \geq t)} \\ &= \frac{1}{1 - \mathcal{P}(t)} \frac{2}{\pi} \sum_{k=0}^{\infty} \frac{1}{2k+1} \left((-1)^k + \sin \frac{\pi(2k+1)\beta}{2} \right) \\ &\quad \times \exp\left(-\frac{1}{8} \pi^2 (2k+1)^2 t\right), \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{D}(\beta; t) &= \frac{1}{1 - \mathcal{P}(t)} \\
 (3.12) \quad &\times \sum_{k=0}^{\infty} \frac{(-1)^k}{2} \left(\operatorname{erfc} \frac{2k-1}{\sqrt{2t}} - \operatorname{erfc} \frac{2k+\beta}{\sqrt{2t}} \right. \\
 &\quad \left. - \operatorname{erfc} \frac{2k+2-\beta}{\sqrt{2t}} + \operatorname{erfc} \frac{2k+3}{\sqrt{2t}} \right).
 \end{aligned}$$

PROOF. The first equality in (3.11) flows out of equivalence of the events $(|W(s)| < 1, 0 < s < t)$ and $(\tau \geq t)$. Let us prove the second one. To this end consider the probability

$$u(t, x) = P(\tau_x \geq t, \alpha \leq x + W(t) < \beta),$$

where $\alpha \geq -1$.

Due to (2.15)–(2.17), (2.20) under (2.13), this probability is the solution of the following boundary value problem:

$$(3.13) \quad \frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}, \quad t > 0, \quad -1 < x < 1,$$

$$(3.14) \quad u(0, x) = \chi_{[\alpha, \beta]}(x), \quad u(t, -1) = u(t, 1) = 0, \quad t > 0.$$

Solving this problem, we get

$$\begin{aligned}
 u(t, x) &= \frac{2}{\pi} \sum_{k=1}^{\infty} \frac{1}{k} \sin \frac{\pi k(\alpha + \beta)}{2} \sin \frac{\pi k(\beta - \alpha)}{2} \sin \pi kx \exp\left(-\frac{1}{2} \pi^2 k^2 t\right) \\
 &\quad + \frac{4}{\pi} \sum_{k=0}^{\infty} \frac{1}{2k+1} \sin \frac{\pi(2k+1)(\beta - \alpha)}{4} \cos \frac{\pi(2k+1)(\beta + \alpha)}{4} \\
 &\quad \times \cos \frac{\pi(2k+1)x}{2} \exp\left(-\frac{1}{8} \pi^2 (2k+1)^2 t\right).
 \end{aligned}$$

As $P(W(t) < \beta, \tau \geq t) = u(t, 0)$ under $\alpha = -1, x = 0$, we arrive at (3.11) from here. Equality (3.12) follows from

$$u(t, x) = \frac{1}{\sqrt{2\pi t}} \int_{\alpha}^{\beta} G(t, x, y) dy$$

obtained analogously to (3.4). \square

Let us note that the series (3.11) and (3.12) are of the Leibniz type, (3.11) is convenient for calculations under large t , and (3.12) is convenient under small t . We draw attention to the denominator $(1 - \mathcal{P}(t))$ in (3.11), which is close to zero for $t \gg 1$. But it is not difficult to transform (3.11) to the form proper for calculations. See the curves of the distribution $\mathcal{D}(\beta; t)$ for some values of t in Figure 2.




FIG. 2. The distribution function $Q(\beta; \cdot)$; under $t \geq 0.5$ the curves coincide visually.

Let the function $\mathcal{D}^{-1}(\cdot; t)$ be the inverse function to $\mathcal{D}(\cdot; t)$ for every fixed t . Then the random variable

$$\xi = \mathcal{D}^{-1}(\gamma; t)$$

has $\mathcal{D}(\beta; t)$ as its distribution function.

4. Simulation of exit time and exit point of a Wiener process from a cube. Let $C \subset R^d$ be a d -dimensional cube with center at the origin and with edge length equal to 2. We suppose all the edges of the cube to be parallel to the coordinate axes, that is, $C = \{x = (x^1, \dots, x^d): |x^i| < 1, i = 1, \dots, d\}$. Let $W(s) = (W^1(s), \dots, W^d(s))^T$ be a d -dimensional standard Wiener process, τ be the first-passage time of $W(s)$ to the boundary ∂C of the cube C .

Let us give the following evident result in the form of a lemma.

LEMMA 4.1. The distribution function $\mathcal{P}_d(t)$ for τ is equal to

$$(4.1) \quad \mathcal{P}_d(t) = P(\tau < t) = 1 - (1 - \mathcal{D}(t))^d$$

and the random variable

$$(4.2) \quad \tau = \mathcal{D}^{-1}(1 - \gamma^{1/d})$$

is distributed by the law $\mathcal{P}_d(t)$.

Our nearest goal is to construct an algorithm for simulation of the point $(\tau, W(\tau))$. To this end, let us obtain some distributions connected with the d -dimensional Wiener process.

LEMMA 4.2. *Let τ^j be the first-passage time of the component $W^j(t)$ to the boundary of the interval $[-1, 1]$. Then*

$$(4.3) \quad \begin{aligned} P\left(\bigcap_{i \neq j} (W^i(\tau^j) < \beta^i, |W^i(s)| < 1, 0 < s < \tau^j)/\tau^j\right) \\ = (1 - \mathcal{P}(\tau^j))^{d-1} \prod_{i \neq j} \mathcal{Q}(\beta^i; \tau^j). \end{aligned}$$

PROOF. We shall use an assertion of the following kind: if $\zeta \geq 0$ is $\tilde{\mathcal{F}}$ -measurable (where $\tilde{\mathcal{F}}$ is a σ -subalgebra of a general σ -algebra \mathcal{F}), a random variable $\varphi(t, \omega)$ under every $t \geq 0$ does not depend on $\tilde{\mathcal{F}}$ [$\varphi(t, \omega)$ is supposed to be measurable on t], and $E\varphi(t, \omega) = h(t)$, then $E(\varphi(\zeta, \omega)/\tilde{\mathcal{F}}) = h(\zeta)$ (see [9], page 67, [15], page 158).

Due to Lemma 3.2 and independence of the processes $W^i(s)$, we get for any $t \geq 0$,

$$P\left(\bigcap_{i \neq j} (W^i(t) < \beta^i, |W^i(s)| < 1, 0 < s < t)\right) = (1 - \mathcal{P}(t))^{d-1} \prod_{i \neq j} \mathcal{Q}(\beta^i; t).$$

This equality implies (4.3) in accordance with the above-mentioned assertion because the processes $W^i(s)$, $i \neq j$, do not depend on the process $W^j(s)$. \square

Introduce the random variable \varkappa which takes the value j for $\omega \in \{\omega: W^j(\tau) = \pm 1\}$. This variable is defined uniquely with probability 1, and $P(\varkappa = j) = 1/d$. Let $\nu := W^\varkappa(\tau)$. Clearly, the distribution law for ν is given by $P(\nu = -1) = P(\nu = 1) = \frac{1}{2}$.

LEMMA 4.3. *The following equality takes place:*

$$(4.4) \quad \begin{aligned} P\left(\varkappa = j, \tau < \theta, \bigcap_{i \neq j} (W^i(\tau) < \beta^i)\right) \\ = \int_0^\theta (1 - \mathcal{P}(\vartheta))^{d-1} \prod_{i \neq j} \mathcal{Q}(\beta^i; \vartheta) \mathcal{P}'(\vartheta) d\vartheta. \end{aligned}$$

PROOF. We have

$$\begin{aligned} P\left(\varkappa = j, \tau < \theta, \bigcap_{i \neq j} (W^i(\tau) < \beta^i)\right) \\ = P\left(\bigcap_{i \neq j} (W^i(\tau^j) < \beta^i, |W^i(s)| < 1, 0 < s < \tau^j), \tau^j < \theta\right) \\ = \int_0^\theta P\left(\bigcap_{i \neq j} (W^i(\tau^j) < \beta^i, |W^i(s)| < 1, 0 < s < \tau^j)/\tau^j = \vartheta\right) d\mathcal{P}_{\tau^j}(\vartheta), \end{aligned}$$

where $\mathcal{P}_{\tau^j}(\vartheta)$ is the distribution function for τ^j . Clearly $\mathcal{P}_{\tau^j}(\vartheta) = \mathcal{P}(\vartheta)$. Now the assertion (4.4) arises from Lemma 4.2. \square

LEMMA 4.4. *The following equality takes place:*

$$(4.5) \quad P\left(\bigcap_{i \neq j} (W^i(\tau) < \beta^i) / \varkappa = j, \tau = \theta\right) = \prod_{i \neq j} \mathcal{P}(\beta^i; \theta).$$

PROOF. The random variables \varkappa and τ are independent. Indeed, $P(\varkappa = 1, \tau < \theta) = \dots = P(\varkappa = d, \tau < \theta)$ on the strength of symmetry. Hence $P(\varkappa = i, \tau < \theta) = (1/d)P(\tau < \theta) = P(\varkappa = i)P(\tau < \theta)$. Further [see (4.1)],

$$dP(\varkappa = j, \tau < \theta) = \frac{1}{d}d\mathcal{P}_d(\theta) = (1 - \mathcal{P}(\theta))^{d-1}\mathcal{P}'(\theta)d\theta.$$

From here we get

$$(4.6) \quad \begin{aligned} & P\left(\varkappa = j, \tau < \theta, \bigcap_{i \neq j} (W^i(\tau) < \beta^i)\right) \\ &= \int_0^\theta P\left(\bigcap_{i \neq j} (W^i(\tau) < \beta^i) / \varkappa = j, \tau = \vartheta\right) (1 - \mathcal{P}(\vartheta))^{d-1} \mathcal{P}'(\vartheta) d\vartheta. \end{aligned}$$

Comparing (4.4) with (4.6), we obtain (4.5). \square

Let us note that the point $(\tau, W(\tau)) \in [0, \infty) \times \partial C$, that is, this point belongs to the lateral surface of the unbounded semicylinder $[0, \infty) \times C$ with cubic base in $(d + 1)$ -dimensional space of variables (t, x^1, \dots, x^d) .

THEOREM 4.1 (Algorithm for simulating exit point to lateral surface of a cylinder with cubic base). *Let $\varkappa, \nu, \gamma, \gamma^1, \dots, \gamma^{d-1}$ be independent random variables. Let \varkappa and ν be simulated by the laws $P(\varkappa = j) = 1/d, j = 1, \dots, d; P(\nu = \pm 1) = \frac{1}{2}$, and let $\gamma, \gamma^1, \dots, \gamma^{d-1}$ be uniformly distributed on $[0, 1]$. Then the point $(\tau, \xi) = (\tau, \xi^1, \dots, \xi^d)$ with*

$$(4.7) \quad \begin{aligned} \tau &= \mathcal{D}^{-1}(1 - \gamma^{1/d}), & \xi^1 &= \mathcal{D}^{-1}(\gamma^1; \tau), \dots, \xi^{\varkappa-1} = \mathcal{D}^{-1}(\gamma^{\varkappa-1}; \tau), \\ \xi^\varkappa &= \nu, & \xi^{\varkappa+1} &= \mathcal{D}^{-1}(\gamma^\varkappa; \tau), \dots, \xi^d = \mathcal{D}^{-1}(\gamma^{d-1}; \tau) \end{aligned}$$

has the same distribution as $(\tau, W(\tau))$.

This theorem is a simple consequence of Lemmas 4.1 and 4.4.

COROLLARY 4.1. *Let $C_r = \{x = (x^1, \dots, x^d): |x^i| < r, i = 1, \dots, d\} \subset R^d$ be a d -dimensional cube with center at the origin and with edge length equal to $2r$. Let $\bar{\theta}$ be the first-passage time of the d -dimensional standard Wiener process $w(s)$ to the boundary ∂C_r of the cube C_r . Then the point*

$$(\bar{\theta}, \bar{w}) = (r^2\tau, r\xi),$$

where (τ, ξ) is simulated by the algorithm for simulating exit point to lateral surface of cylinder with the cubic base C , has the same distribution as $(\bar{\theta}, w(\bar{\theta}))$.

The proof easily follows from the fact that if $W(t)$ is a Wiener process, then $w(t) = rW(t/r^2)$ is a Wiener process as well.

REMARK 4.1. The algorithm for simulating exit point to lateral surface of a cylinder with parallelepiped base is more complicated because of the dependence of κ and τ . This algorithm will be adduced later as a consequence of some following results.

5. Simulation of exit point of the space–time Brownian motion from a space–time parallelepiped with cubic base. Now let us consider the space–time parallelepiped $\Pi = [0, l] \times C \subset R^{d+1}$, where the cube $C \subset R^d$ is defined as above, and construct an algorithm for simulating the exit point $(\tau(l), W(\tau(l)))$ from the parallelepiped Π . The random variable $\tau(l)$ is found as $\min(\tau, l)$, where τ is the first-passage time of $W(s)$ to the boundary ∂C as above. The distribution function of $\tau(l)$ is equal to

$$(5.1) \quad P(\tau(l) < t) = \begin{cases} 1 - (1 - \mathcal{P}(t))^d, & t \leq l, \\ 1, & t > l. \end{cases}$$

THEOREM 5.1 (Algorithm for simulating exit point from a space–time parallelepiped with cubic base). *Let $\iota, \kappa, \nu, \gamma, \gamma^1, \dots, \gamma^{d-1}$ be independent random variables. Let ι be simulated by the law*

$$P(\iota = -1) = 1 - (1 - \mathcal{P}(l))^d, \quad P(\iota = 1) = (1 - \mathcal{P}(l))^d,$$

and the random variables $\kappa, \nu, \gamma, \gamma^1, \dots, \gamma^{d-1}$ be simulated as in Theorem 4.1.

Then a random point $(\tau(l), \xi)$, distributed as the exit point $(\tau(l), W(\tau(l)))$, is simulated by the following algorithm.

If the simulated value of ι is equal to -1 , then the point $(\tau(l), \xi)$ belongs to the lateral surface of Π , and

$$\begin{aligned} \tau(l) &= \mathcal{P}^{-1}(1 - [1 - \gamma(1 - (1 - \mathcal{P}(l))^d)]^{1/d}), \\ \xi^1 &= \mathcal{D}^{-1}(\gamma^1; \tau(l)), \dots, \xi^{\kappa-1} = \mathcal{D}^{-1}(\gamma^{\kappa-1}; \tau(l)), \quad \xi^\kappa = \nu, \\ \xi^{\kappa+1} &= \mathcal{D}^{-1}(\gamma^\kappa; \tau(l)), \dots, \xi^d = \mathcal{D}^{-1}(\gamma^{d-1}; \tau(l)); \end{aligned}$$

otherwise, when $\iota = 1$, the point $(\tau(l), \xi)$ belongs to the upper base of Π , and

$$\begin{aligned} \tau(l) &= l, \\ \xi^1 &= \mathcal{D}^{-1}(\gamma; l), \quad \xi^2 = \mathcal{D}^{-1}(\gamma^1; l), \dots, \xi^d = \mathcal{D}^{-1}(\gamma^{d-1}; l). \end{aligned}$$

PROOF. Using Lemma 4.1, we have

$$(5.2) \quad \begin{aligned} P(\tau(l) < l) &= P(\tau < l) = 1 - (1 - \mathcal{P}(l))^d, \\ P(\tau(l) = l) &= P(\tau \geq l) = (1 - \mathcal{P}(l))^d. \end{aligned}$$

The conditional probability $P(\tau(l) < t/\tau(l) < l)$ is equal to

$$P(\tau(l) < t/\tau(l) < l) = \frac{P((\tau(l) < t) \cap (\tau(l) < l))}{P(\tau(l) < l)} = \chi_{[l, \infty)}(t) + \chi_{[0, l)}(t) \frac{P(\tau < t)}{P(\tau < l)},$$

and the random variable $\mathcal{D}^{-1}(1 - [1 - \gamma(1 - (1 - \mathcal{D}(l))^d)]^{1/d})$ is distributed by the law $P(\tau(l) < t/\tau(l) < l)$.

Carrying out reasoning similar to Lemmas 4.2, 4.3 and 4.4 we obtain

$$(5.3) \quad P\left(\bigcap_{i \neq j} (W^i(\tau(l)) < \beta^i) / \kappa = j, \tau(l) = \theta < l\right) = \chi_{[0, l)}(\theta) \prod_{i \neq j} \mathcal{D}(\beta^i; \theta).$$

Further, the equality

$$(5.4) \quad \begin{aligned} &P\left(\bigcap_{i=1}^d (W^i(\tau(l)) < \beta^i) / \tau(l) = l\right) \\ &= P\left(\bigcap_{i=1}^d (W^i(l) < \beta^i) / \tau \geq l\right) \\ &= \frac{1}{P(\tau \geq l)} P\left(\bigcap_{i=1}^d (W^i(l) < \beta^i, |W^i(s)| < 1, 0 < s < l)\right) \\ &= \frac{1}{\prod_{i=1}^d P(\tau^i \geq l)} P\left(\bigcap_{i=1}^d (W^i(l) < \beta^i, \tau^i \geq l)\right) = \prod_{i=1}^d \mathcal{D}(\beta^i; l) \end{aligned}$$

holds due to the mutual independence of the components $W^i, i = 1, \dots, d$, and Lemma 3.2.

Now the statement of the theorem easily follows from (5.2)–(5.4). \square

The following corollary has the same proof as Corollary 4.1.

COROLLARY 5.1. *Let $\Pi_r = [0, lr^2) \times C_r = \{(t, x) = (t, x^1, \dots, x^d) : 0 \leq t < lr^2, |x^i| < r, i = 1, \dots, d\} \subset R^{d+1}$ be a space-time parallelepiped. Let $\bar{\theta}$ be the first-passage time of the process $(s, w(s)), s > 0$, to the boundary $\partial\Pi_r$. Then the point*

$$(\bar{\theta}, \bar{w}) = (r^2\tau(l), r\xi),$$

where $(\tau(l), \xi)$ is simulated by the algorithm for simulating exit point from the space-time parallelepiped Π , has the same distribution as $(\bar{\theta}, w(\bar{\theta}))$.

6. Theorem on local mean-square approximation. Let us return to the problem (1.1). Here and in the next sections we assume that the coefficients of (1.1) belong to the class $C^{1,2}([t_0, t_1] \times \bar{G})$, the boundary ∂G of the domain G is twice continuously differentiable, and the strict ellipticity condition is imposed on the matrix $a(s, x) = \sigma(s, x)\sigma^T(s, x)$ [see (2.5)].

Introduce the space–time parallelepiped $U_r^{\sigma(t,x)}(x)$,

$$U_r^{\sigma(t,x)}(x) = \bigcup_{0 \leq s < lr^2} \{t + s\} \times C_r^{\sigma(t,x)}(x + b(t,x)s),$$

where $(t, x) \in Q$ and $C_r^{\sigma(t,x)}(x + b(t,x)s)$ is the space parallelepiped in R^d obtained from the open cube C_r by the linear transformation $\sigma(t, x)$ and the shift $x + b(t, x)s$, and as in the previous section, C_r is the cube with center at the origin and with edges of length $2r$ which are parallel to the coordinate axes. The time size of $U_r^{\sigma(t,x)}(x)$ is taken lr^2 as the characteristic exit time of a diffusion process from a space cube of linear size r is proportional to r^2 .

Let Γ_δ be an intersection of a δ -neighborhood of the set Γ with the domain Q . Remember that the set Γ is a part of the boundary ∂Q consisting of the lateral surface and the upper base of the cylinder \bar{Q} . The size δ of the layer Γ_δ may depend on r . The condition of strict ellipticity ensures for any $\beta > 0$ the existence of a constant $\alpha > 0$ such that under all sufficiently small r for every point $(t, x) \in Q \setminus \Gamma_{\alpha r}$ the following relations take place:

$$(6.1) \quad U_r^{\sigma(t,x)}(x) \subset Q, \quad \min_{0 \leq s \leq lr^2} \rho(\partial C_r^{\sigma(t,x)}(x + b(t,x)s), \partial G) \geq \beta r.$$

Indeed, due to the property of strong ellipticity, we get

$$\max_{0 \leq s \leq lr^2} \rho(x, \partial C_r^{\sigma(t,x)}(x + b(t,x)s)) \leq lr^2 \max_{(s,y) \in \bar{Q}} |b(s,y)| + 2r\sqrt{d}\lambda_d.$$

It is easy to see that if we take

$$\alpha = lr \max_{(s,y) \in \bar{Q}} |b(s,y)| + 2\sqrt{d}\lambda_d + \beta,$$

then under a sufficiently small r the relations (6.1) are fulfilled.

The values β , α and r used below are assumed to ensure (6.1).

To construct a one-step approximation for the system (1.1), we consider the system with frozen coefficients,

$$(6.2) \quad d\bar{X} = b(t, x) ds + \sigma(t, x) dw(s), \quad \bar{X}(t) = x, \quad (t, x) \in Q \setminus \Gamma_{\alpha r}.$$

Let $\bar{\theta}$ be the first-passage time of the process $(s - t, w(s) - w(t))$, $s > t$, to the boundary $\partial \Pi_r$ of the space–time parallelepiped $\Pi_r = [0, lr^2] \times C_r \subset R^{d+1}$. Clearly, $\bar{\theta} \leq lr^2$. The point $(\bar{\theta}, w(t + \bar{\theta}) - w(t))$ is simulated in accordance with Corollary 5.1.

Let us take the point $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$ with $\bar{X}_{t,x}(t + \bar{\theta})$ calculated by

$$(6.3) \quad \bar{X}_{t,x}(t + \bar{\theta}) = x + b(t, x)\bar{\theta} + \sigma(t, x)(w(t + \bar{\theta}) - w(t))$$

as an approximation of the point $(t + \bar{\theta}, X_{t,x}(t + \bar{\theta}))$, $(t, x) \in Q \setminus \Gamma_{\alpha r}$, where $X_{t,x}(s)$ is a solution of the system (1.1). Remember that if $t + \bar{\theta} \geq \tau_{t,x}$, then $X_{t,x}(t + \bar{\theta}) = X_{t,x}(\tau_{t,x})$.

The point $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$ belongs to the lateral surface or to the upper base of the space–time parallelepiped $U_r^{\sigma(t,x)}(x) \subset Q$.

It follows from (6.1) that

$$(6.4) \quad \rho(\bar{X}_{t,x}(t+s), \partial G) \geq \beta r, \quad 0 \leq s \leq lr^2.$$

THEOREM 6.1. *For every natural m there exists a constant $K > 0$ such that for any sufficiently small r and for any point $(t, x) \in Q \setminus \Gamma_{cr}$ the inequality*

$$(6.5) \quad E|X_{t,x}(t+\bar{\theta}) - \bar{X}_{t,x}(t+\bar{\theta})|^{2m} \leq K r^{4m}$$

holds.

PROOF. Below we use the same letter K without any index for various constants, which depend on the system (1.1) only and do not depend on (t, x) , r and so on. Thereby, we write K instead of, for example, $K + K$, $2K$, K^2 , and so on.

We have [see (1.1)] that $\tau_{t,x} \leq t_1$, $X_{t,x}(s) \in G$ under $s \in [t, \tau_{t,x})$, and $X_{t,x}(s) = X_{t,x}(\tau_{t,x})$ under $s \geq \tau_{t,x}$.

Let us rewrite the local error in the form

$$(6.6) \quad \begin{aligned} & E|X_{t,x}(t+\bar{\theta}) - \bar{X}_{t,x}(t+\bar{\theta})|^{2m} \\ &= E \left| \int_t^{t+\bar{\theta}} (\chi_{\tau_{t,x}>s} b(s, X_{t,x}(s)) - b(t, x)) ds \right. \\ & \quad \left. + \int_t^{t+\bar{\theta}} (\chi_{\tau_{t,x}>s} \sigma(s, X_{t,x}(s)) - \sigma(t, x)) dw(s) \right|^{2m} \\ &\leq K E \left| \int_t^{t+\bar{\theta}} (\chi_{\tau_{t,x}>s} b(s, X_{t,x}(s)) - b(t, x)) ds \right|^{2m} \\ & \quad + K E \left| \int_t^{(t+\bar{\theta}) \wedge \tau_{t,x}} (\sigma(s, X_{t,x}(s)) - \sigma(t, x)) dw(s) \right|^{2m} \\ & \quad + K E \left| \int_{(t+\bar{\theta}) \wedge \tau_{t,x}}^{t+\bar{\theta}} \sigma(t, x) dw(s) \right|^{2m}. \end{aligned}$$

We obtain for the first term in (6.6),

$$(6.7) \quad K E \left| \int_t^{t+\bar{\theta}} (\chi_{\tau_{t,x}>s} b(s, X_{t,x}(s)) - b(t, x)) ds \right|^{2m} \leq K E \bar{\theta}^{2m} \leq K r^{4m}$$

because of the boundedness of $b(s, x)$, $(s, x) \in \bar{Q}$ and $\bar{\theta} \leq lr^2$.

Below we need the following inequality for Itô integrals in the case of the scalar Wiener process (see, e.g., [9], page 26):

$$(6.8) \quad \begin{aligned} & E \left(\int_t^{t+T} \varphi(s) dw(s) \right)^{2m} \\ & \leq (m(2m-1))^{m-1} T^{m-1} \int_t^{t+T} E \varphi^{2m}(s) ds, \quad m = 1, 2, \dots \end{aligned}$$

Clearly, in the case of the d -dimensional Wiener process, the inequality (6.8) implies

$$(6.9) \quad \begin{aligned} & E \left| \int_t^{t+T} \varphi(s) dw(s) \right|^{2m} \\ & \leq KT^{m-1} \int_t^{t+T} E \sum_{i,j=1}^d (\varphi^{ij}(s))^{2m} ds, \quad m = 1, 2, \dots, \end{aligned}$$

where the constant K depends on m , of course.

If φ is bounded, we also have

$$(6.10) \quad E \left| \int_t^{t+T} \varphi(s) dw(s) \right|^{2m} \leq KT^m, \quad m = 1, 2, \dots$$

Due to inequality (6.9), smoothness of $\sigma(s, x)$, $(s, x) \in \bar{Q}$ and $(t + \bar{\theta}) \wedge \tau_{t,x} \leq t + lr^2$, we obtain for the second term of (6.6)

$$(6.11) \quad \begin{aligned} & KE \left| \int_t^{(t+\bar{\theta}) \wedge \tau_{t,x}} (\sigma(s, X(s)) - \sigma(t, x)) dw(s) \right|^{2m} \\ & = KE \left| \int_t^{t+lr^2} \chi_{(t+\bar{\theta}) \wedge \tau_{t,x} > s} (\sigma(s, X_{t,x}(s)) - \sigma(t, x)) dw(s) \right|^{2m} \\ & \leq Kr^{2m-2} \int_t^{t+lr^2} E \left(\chi_{(t+\bar{\theta}) \wedge \tau_{t,x} > s} \sum_{i,j=1}^d |\sigma^{ij}(s, X_{t,x}(s)) - \sigma^{ij}(t, x)|^{2m} \right) ds \\ & \leq Kr^{2m-2} \int_t^{t+lr^2} (E \chi_{\tau_{t,x} > s} |X_{t,x}(s) - x|^{2m} + (s-t)^{2m}) ds \\ & \leq Kr^{2m-2} \int_t^{t+lr^2} E \chi_{\tau_{t,x} > s} |X_{t,x}(s) - x|^{2m} ds + Kr^{6m}. \end{aligned}$$

Further,

$$\begin{aligned} & E \chi_{\tau_{t,x} > s} |X_{t,x}(s) - x|^{2m} \\ & = E \chi_{\tau_{t,x} > s} \left| \int_t^s b(s', X_{t,x}(s')) ds' + \int_t^s \sigma(s', X_{t,x}(s')) dw(s') \right|^{2m}, \end{aligned}$$

whence, due to (6.10),

$$E \chi_{\tau_{t,x} > s} |X_{t,x}(s) - x|^{2m} \leq K(s-t)^{2m} + K(s-t)^m.$$

Substituting this inequality in (6.11), we obtain

$$(6.12) \quad KE \left| \int_t^{(t+\bar{\theta}) \wedge \tau_{t,x}} (\sigma(s, X_{t,x}(s)) - \sigma(t, x)) dw(s) \right|^{2m} \leq Kr^{4m}.$$

It follows from inequalities (6.7) and (6.12) that

$$(6.13) \quad E |X_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta})) - \bar{X}_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta}))|^{2m} \leq Kr^{4m}.$$

Now let us estimate the third term in (6.6). We have, due to (6.9),

$$\begin{aligned}
 (6.14) \quad & KE \left| \int_{(t+\bar{\theta}) \wedge \tau_{t,x}}^{t+\bar{\theta}} \sigma(t, x) dw(s) \right|^{2m} \\
 &= KE \left| \int_t^{t+lr^2} (\chi_{(t+\bar{\theta})>s} - \chi_{(t+\bar{\theta}) \wedge \tau_{t,x}>s}) \sigma(t, x) dw(s) \right|^{2m} \\
 &\leq Kr^{2m-2} \int_t^{t+lr^2} E(\chi_{(t+\bar{\theta})>s} - \chi_{(t+\bar{\theta}) \wedge \tau_{t,x}>s}) ds \\
 &= Kr^{2m-2} E\chi_{\tau_{t,x} < (t+\bar{\theta})} ((t + \bar{\theta}) - (t + \bar{\theta}) \wedge \tau_{t,x}) \\
 &\leq Kr^{2m} P(\tau_{t,x} < t + \bar{\theta}).
 \end{aligned}$$

Evaluate the probability $P(\tau_{t,x} < t + \bar{\theta})$ using the reception from [24]. If $\tau_{t,x} < t + \bar{\theta}$, then $\tau_{t,x} < t_1$ and, consequently, $\bar{X}_{t,x}(\tau_{t,x}) \in \partial G$. At the same time, due to (6.4),

$$\rho(\bar{X}_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta})), \partial G) \geq \beta r.$$

Therefore,

$$\begin{aligned}
 & E(\chi_{\tau_{t,x} < t+\bar{\theta}} | X_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta})) - \bar{X}_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta}))|^m) \\
 & \geq P(\tau_{t,x} < t + \bar{\theta})(\beta r)^m, \quad m = 1, 2, \dots
 \end{aligned}$$

On the other hand, due to (6.13), we have

$$\begin{aligned}
 & P(\tau_{t,x} < t + \bar{\theta})(\beta r)^m \\
 & \leq E(\chi_{\tau_{t,x} < t+\bar{\theta}} | X_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta})) - \bar{X}_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta}))|^m) \\
 & \leq \sqrt{P(\tau_{t,x} < t + \bar{\theta}) [E|X_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta})) - \bar{X}_{t,x}(\tau_{t,x} \wedge (t + \bar{\theta}))|^{2m}]^{1/2}} \\
 & \leq Kr^{2m} \sqrt{P(\tau_{t,x} < t + \bar{\theta})}.
 \end{aligned}$$

Consequently,

$$(6.15) \quad P(\tau_{t,x} < t + \bar{\theta}) \leq Kr^{2m}, \quad m = 1, 2, \dots$$

Now inequality (6.6) together with (6.7), (6.12) and (6.14) gives (6.5). \square

7. Global algorithm and convergence theorems. Let us construct a random walk over small space-time parallelepipeds based on the one-step approximation (6.3) of the previous section. Let $(\bar{\theta}_1, w(t + \bar{\theta}_1) - w(t))$ be the first exit point of the process $(s - t, w(s) - w(t))$, $s > t$, from the parallelepiped Π simulated in accordance with Corollary 5.1, $(\bar{\theta}_2, w(t + \bar{\theta}_1 + \bar{\theta}_2) - w(t + \bar{\theta}_1))$ be the exit point of the process $(s - t - \bar{\theta}_1, w(s) - w(t + \bar{\theta}_1))$, $s > t + \bar{\theta}_1$, from the parallelepiped Π , and so on.

Suppose that $(t, x) \in Q \setminus \Gamma_{\alpha r}$. Then, we construct the recurrence sequence $(\bar{\vartheta}_k, \bar{X}_k), k = 0, 1, \dots, \bar{\nu}$,

$$\begin{aligned} \bar{\vartheta}_0 &= t, & \bar{X}_0 &= x, \\ \bar{\vartheta}_k &= \bar{\vartheta}_{k-1} + \bar{\theta}_k, \\ \bar{X}_k &= \bar{X}_{k-1} + b(\bar{\vartheta}_{k-1}, \bar{X}_{k-1})\bar{\theta}_k \\ &\quad + \sigma(\bar{\vartheta}_{k-1}, \bar{X}_{k-1})(w(\bar{\vartheta}_k) - w(\bar{\vartheta}_{k-1})), \quad k = 1, \dots, \bar{\nu}, \end{aligned}$$

where the number $\bar{\nu} = \bar{\nu}_{t,x}$ is the first one for which $(\bar{\vartheta}_k, \bar{X}_k) \in \Gamma_{\alpha r}$.

If $(t, x) \in \Gamma_{\alpha r}$, we put $\bar{\nu} = 0$.

Let $(\bar{\vartheta}_k, \bar{X}_k) = (\bar{\vartheta}_{\bar{\nu}}, \bar{X}_{\bar{\nu}})$ under $k > \bar{\nu}$. The obtained sequence $(\bar{\vartheta}_k, \bar{X}_k), k = 0, 1, \dots$, is a Markov chain stopping at the Markov moment $\bar{\nu}$. It is clear that the random number of steps $\bar{\nu}$ depends on the domain $Q \setminus \Gamma_{\alpha r}$. That is why the more rigorous notation for $\bar{\nu}$ is $\bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r})$.

Following the technique proposed in [19, 22, 23], we get the theorems on average characteristics of $\bar{\nu}_{t,x} = \bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r})$.

THEOREM 7.1. *The mean number of steps $\bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r})$ is estimated as*

$$(7.1) \quad E\bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r}) \leq \frac{K}{r^2},$$

where the positive constant K does not depend on r .

THEOREM 7.2. *For every $L > 0$, the inequality*

$$(7.2) \quad \begin{aligned} &P\left\{\bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r}) \geq \frac{L}{r^2}\right\} \\ &\leq (1 + t_1 - t_0) \exp\left(-c_r \frac{\gamma}{1 + t_1 - t_0} L\right), \quad c_r \rightarrow 1 \text{ as } r \rightarrow 0, \end{aligned}$$

is valid.

Further, we need two auxiliary lemmas.

LEMMA 7.1. *There exists a constant K such that, for all r small enough and all $(t, x) \in Q \setminus \Gamma_{\alpha r}$, the inequality*

$$(7.3) \quad |E(X_{t,x}(t + \bar{\theta}_1) - \bar{X}_{t,x}(t + \bar{\theta}_1))| \leq Kr^4$$

is valid.

PROOF. By the Itô formula, smoothness of $b(s, x)$ and the inequality (6.15) under $m = 1$, we obtain

$$\begin{aligned}
 & \left| E(X_{t,x}(t + \bar{\theta}_1) - \bar{X}_{t,x}(t + \bar{\theta}_1)) \right| \\
 &= \left| E \int_t^{t+\bar{\theta}_1} (\chi_{\tau_{t,x} > s} b(s, X_{t,x}(s)) - b(t, x)) ds \right. \\
 & \quad \left. + E \int_t^{t+\bar{\theta}_1} (\chi_{\tau_{t,x} > s} \sigma(s, X_{t,x}(s)) - \sigma(t, x)) dw(s) \right| \\
 &= \left| E \int_t^{t+\bar{\theta}_1} (\chi_{\tau_{t,x} > s} b(s, X_{t,x}(s)) - b(t, x)) ds \right| \\
 &= \left| E \left(\int_t^{(t+\bar{\theta}_1) \wedge \tau_{t,x}} (b(s, X_{t,x}(s)) - b(t, x)) ds \right) \right. \\
 & \quad \left. - b(t, x) E((t + \bar{\theta}_1) - \tau_{t,x} \wedge (t + \bar{\theta}_1)) \right| \\
 &\leq \left| E \left(\int_t^{(t+\bar{\theta}_1) \wedge \tau_{t,x}} \int_t^s Lb(s', X_{t,x}(s')) ds' ds \right) \right| \\
 & \quad + KE((t + \bar{\theta}_1) - \tau_{t,x} \wedge (t + \bar{\theta}_1)) \\
 &\leq KE\bar{\theta}_1^2 + Kr^2 P(\tau_{t,x} < t + \bar{\theta}_1) \leq Kr^4,
 \end{aligned}$$

where

$$L = \frac{\partial}{\partial s} + \frac{1}{2} \sum_{i,j=1}^d \alpha^{ij} \frac{\partial^2}{\partial x^i \partial x^j} + \sum_{i=1}^d b^i \frac{\partial}{\partial x^i}. \quad \square$$

LEMMA 7.2. Let the random variable Z be defined by the relation

$$X_{t,x}(t + \bar{\theta}_1) - X_{t,y}(t + \bar{\theta}_1) = x - y + Z.$$

Then for every natural m there exists a positive constant K such that for any r small enough and all $(t, x), (t, y) \in \mathcal{Q} \setminus \Gamma_{\alpha r}$ the inequalities

$$(7.4) \quad E|Z|^m \leq K r^m (|x - y|^m + r^m),$$

$$(7.5) \quad |EZ| \leq K r^2 (|x - y| + r^2)$$

hold.

PROOF. We have, due to $(t, x), (t, y) \in \mathcal{Q} \setminus \Gamma_{\alpha r}$ and (6.3),

$$\bar{X}_{t,x}(t + \bar{\theta}_1) = x + b(t, x)\bar{\theta}_1 + \sigma(t, x)(w(t + \bar{\theta}_1) - w(t))$$

and

$$\bar{X}_{t,y}(t + \bar{\theta}_1) = y + b(t, y)\bar{\theta}_1 + \sigma(t, y)(w(t + \bar{\theta}_1) - w(t)).$$

Then

$$\begin{aligned} Z &= X_{t,x}(t + \bar{\theta}_1) - X_{t,y}(t + \bar{\theta}_1) - (x - y) \\ &= (X_{t,x}(t + \bar{\theta}_1) - \bar{X}_{t,x}(t + \bar{\theta}_1)) - (X_{t,y}(t + \bar{\theta}_1) - \bar{X}_{t,y}(t + \bar{\theta}_1)) \\ &\quad + (b(t, x) - b(t, y))\bar{\theta}_1 + (\sigma(t, x) - \sigma(t, y))(w(t + \bar{\theta}_1) - w(t)). \end{aligned}$$

By Lemma 7.1 and the smoothness of $b(s, x)$, $(s, x) \in \bar{Q}$, we get

$$\begin{aligned} |EZ| &\leq |E(X_{t,x}(t + \bar{\theta}_1) - \bar{X}_{t,x}(t + \bar{\theta}_1))| \\ &\quad + |E(X_{t,y}(t + \bar{\theta}_1) - \bar{X}_{t,y}(t + \bar{\theta}_1))| \\ &\quad + |b(t, x) - b(t, y)|E\bar{\theta}_1 \\ &\leq Kr^4 + K|x - y|r^2, \end{aligned}$$

that gives (7.5).

Now consider the $2n$ th moments of Z . Using Theorem 6.1, the property (6.9), boundedness of $b(s, x)$, $(s, x) \in \bar{Q}$ and smoothness of $\sigma(s, x)$, $(s, x) \in \bar{Q}$, we obtain

$$\begin{aligned} E|Z|^{2n} &\leq K E|X_{t,x}(t + \bar{\theta}_1) - \bar{X}_{t,x}(t + \bar{\theta}_1)|^{2n} \\ &\quad + K E|X_{t,y}(t + \bar{\theta}_1) - \bar{X}_{t,y}(t + \bar{\theta}_1)|^{2n} \\ &\quad + K|b(t, x) - b(t, y)|^{2n} E\bar{\theta}_1^{2n} \\ &\quad + E \left| \int_t^{t+lr^2} \chi_{t+\bar{\theta}_1 > s} (\sigma(t, x) - \sigma(t, y)) dw(s) \right|^{2n} \\ &\leq Kr^{4n} + Kr^{2n}|x - y|^{2n} \end{aligned}$$

that gives (7.4) in the case of the even m .

In the case of the odd m , we come to (7.4) using the Cauchy–Bunyakovskii inequality,

$$E|Z|^m \leq (E|Z|^{2m})^{1/2} \leq (Kr^{2m}(|x - y|^{2m} + r^{2m}))^{1/2} \leq Kr^m(|x - y|^m + r^m). \quad \square$$

For every $\varepsilon \in (0, 1]$ and any $\beta > 0$ it is possible to introduce the layer $\Gamma_{\alpha r^{1-\varepsilon}}$ with a constant α such that under a sufficiently small r and for every $(t, x) \in Q \setminus \Gamma_{\alpha r^{1-\varepsilon}}$ the following relations together with the relations (6.1) take place:

$$U_r^{\sigma(t, x)}(x) \subset Q, \quad \min_{0 \leq s \leq lr^2} \rho(\partial C_r^{\sigma(t, x)}(x + b(t, x)s), \partial G) \geq \beta r^{1-\varepsilon}.$$

Clearly, $\Gamma_{\alpha r} \subset \Gamma_{\alpha r^{1-\varepsilon}}$.

The Markov moment $\bar{v}_{t,x}(Q \setminus \Gamma_{\alpha r^{1-\varepsilon}})$, when the chain $(\bar{\vartheta}_k, \bar{X}_k)$ leaves the domain $Q \setminus \Gamma_{\alpha r^{1-\varepsilon}}$, satisfies the inequality

$$\bar{v}_{t,x}(Q \setminus \Gamma_{\alpha r^{1-\varepsilon}}) \leq \bar{v}_{t,x}(Q \setminus \Gamma_{\alpha r}).$$

We shall use the old notation $(\bar{\vartheta}_k, \bar{X}_k)$ for the new Markov chain, which is constructed by the same rules as above but stops in the layer $\Gamma_{\alpha r^{1-\varepsilon}}$ at the new Markov moment $\bar{\nu} = \bar{\nu}_{t,x}(\mathcal{Q} \setminus \Gamma_{\alpha r^{1-\varepsilon}})$. We believe that the use of the same notation $(\bar{\vartheta}_k, \bar{X}_k)$ for various Markov chains and $\bar{\nu}$ for various stopping moments will cause no confusion below.

Consider the sequence $(\bar{\vartheta}_k, \bar{X}_k)$, $k = 0, 1, \dots$, with X_k ,

$$\begin{aligned} X_0 &= x, \\ X_1 &= X_{t,x}(\bar{\vartheta}_1), \\ &\dots \\ X_k &= X_{t,x}(\bar{\vartheta}_k) = X_{\bar{\vartheta}_{k-1}, X_{k-1}}(\bar{\vartheta}_k), \\ &\dots \end{aligned}$$

connected with the system (1.1).

The sequence $(\bar{\vartheta}_k, \bar{X}_k)$ is a Markov chain, which stops at the random moment $\bar{\nu}$ due to $\bar{\vartheta}_k = \bar{\vartheta}_{\bar{\nu}}$ under $k > \bar{\nu}$.

The following theorem states the closeness of X_k and \bar{X}_k for $N = L/r^2$ steps.

THEOREM 7.3. *Let $\bar{\nu} = \bar{\nu}_{t,x}(\mathcal{Q} \setminus \Gamma_{\alpha r^{1-\varepsilon}})$, $0 < \varepsilon \leq 1$, be the first exit moment of the Markov chain $(\bar{\vartheta}_i, \bar{X}_i)$, $i = 1, 2, \dots$, from the domain $\mathcal{Q} \setminus \Gamma_{\alpha r^{1-\varepsilon}}$. Then there exist constants $K > 0$ and $\gamma > 0$ such that for all r small enough, the inequality*

$$(E|X_{N \wedge \bar{\nu}} - \bar{X}_{N \wedge \bar{\nu}}|^2)^{1/2} = (E|X_N - \bar{X}_N|^2)^{1/2} \leq K e^{\gamma L} r$$

holds.

PROOF. Below we use the same letter K for various constants (see the note in Theorem 6.1).

Remember that $\bar{\vartheta}_k = \bar{\vartheta}_{k \wedge \bar{\nu}}$, $\bar{X}_k = \bar{X}_{k \wedge \bar{\nu}}$ and $X_k = X_{k \wedge \bar{\nu}} = X(\bar{\vartheta}_{k \wedge \bar{\nu}} \wedge \tau_{t,x})$.

Here we follow the proof of the corresponding theorem in [24].

Let ν be the first number at which $X_\nu \in \Gamma_{cr}$,

$$\nu = \begin{cases} \min\{k: X_k \in \Gamma_{cr}, k \leq \bar{\nu}\}, \\ \infty, X_k \notin \Gamma_{cr}, k \leq \bar{\nu}, \end{cases}$$

where $c \leq (\beta/2)r^{-\varepsilon}$ (here β relates to $\Gamma_{\alpha r^{1-\varepsilon}}$).

Then under $\nu \leq \bar{\nu}$,

$$(7.6) \quad |X_\nu - \bar{X}_\nu| \geq \frac{\beta}{2} r^{1-\varepsilon}.$$

We rewrite the global error in the form (l is a diameter of G):

$$\begin{aligned} E|X_N - \bar{X}_N|^2 &= E\chi_{\nu \geq N \wedge \bar{\nu}} |X_N - \bar{X}_N|^2 + E\chi_{\nu < N \wedge \bar{\nu}} |X_N - \bar{X}_N|^2 \\ (7.7) \quad &\leq E\chi_{\nu \geq N \wedge \bar{\nu}} |X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^2 + l^2 P(\nu < N \wedge \bar{\nu}) \\ &\leq E|X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^2 + l^2 P(\nu < N \wedge \bar{\nu}). \end{aligned}$$

Due to (7.6), we have

$$E\chi_{\nu < N \wedge \bar{\nu}} |X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^n \geq P(\nu < N \wedge \bar{\nu}) \left(\frac{\beta}{2}\right)^n r^{n-\varepsilon n}, \quad n = 1, 2, \dots$$

On the other hand,

$$E\chi_{\nu < N \wedge \bar{\nu}} |X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^n \leq \sqrt{P(\nu < N \wedge \bar{\nu})} \cdot [E|X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^{2n}]^{1/2}.$$

Consequently,

$$(7.8) \quad P(\nu < N \wedge \bar{\nu}) \leq K r^{-2n+2\varepsilon n} E|X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^{2n}.$$

To prove the theorem, we need to find bounds for $E|d_k|^{2n}$, $k = 0, 1, \dots, N$, where $d_k := X_{k \wedge \nu} - \bar{X}_{k \wedge \nu}$. Note that the first term in (7.7) is equal to $E|d_N|^2$.

We have

$$\begin{aligned} d_k &= X_{k \wedge \nu} - \bar{X}_{k \wedge \nu} = (X_{\bar{\nu}_{(k-1) \wedge \nu}, X_{(k-1) \wedge \nu}}(\bar{\vartheta}_{k \wedge \nu}) - X_{\bar{\nu}_{(k-1) \wedge \nu}, \bar{X}_{(k-1) \wedge \nu}}(\bar{\vartheta}_{k \wedge \nu})) \\ &\quad + (X_{\bar{\nu}_{(k-1) \wedge \nu}, \bar{X}_{(k-1) \wedge \nu}}(\bar{\vartheta}_{k \wedge \nu}) - \bar{X}_{k \wedge \nu}). \end{aligned}$$

Denote the second term by ρ_k and define Z_k similarly to Z in Lemma 7.2,

$$\begin{aligned} &X_{\bar{\nu}_{(k-1) \wedge \nu}, X_{(k-1) \wedge \nu}}(\bar{\vartheta}_{k \wedge \nu}) - X_{\bar{\nu}_{(k-1) \wedge \nu}, \bar{X}_{(k-1) \wedge \nu}}(\bar{\vartheta}_{k \wedge \nu}) \\ &= X_{(k-1) \wedge \nu} - \bar{X}_{(k-1) \wedge \nu} + \chi_{\nu \wedge \bar{\nu} > k-1} Z_k. \end{aligned}$$

Then

$$\begin{aligned} d_k &= X_{(k-1) \wedge \nu} - \bar{X}_{(k-1) \wedge \nu} + \chi_{\nu \wedge \bar{\nu} > k-1} Z_k + \chi_{\nu \wedge \bar{\nu} > k-1} \rho_k \\ &= d_{k-1} + \chi_{\nu \wedge \bar{\nu} > k-1} (Z_k + \rho_k). \end{aligned}$$

We have

$$\begin{aligned} E|d_k|^{2n} &= E|d_{k-1} + \chi_{\nu \wedge \bar{\nu} > k-1} (Z_k + \rho_k)|^{2n} \\ &= E[(d_{k-1}, d_{k-1}) + 2(d_{k-1}, \chi_{\nu \wedge \bar{\nu} > k-1} (Z_k + \rho_k)) \\ &\quad + \chi_{\nu \wedge \bar{\nu} > k-1} (Z_k + \rho_k, Z_k + \rho_k)]^n \\ &\leq E|d_{k-1}|^{2n} + 2n E|d_{k-1}|^{2n-2} (d_{k-1}, \chi_{\nu \wedge \bar{\nu} > k-1} (Z_k + \rho_k)) \\ &\quad + K \sum_{m=2}^{2n} E|d_{k-1}|^{2n-m} \chi_{\nu \wedge \bar{\nu} > k-1} |Z_k + \rho_k|^m. \end{aligned}$$

Due to \mathcal{F}_{k-1} -measurability (we denote $\mathcal{F}_m = \mathcal{F}_{\bar{\nu}_m}$) of d_{k-1} and $\chi_{\nu \wedge \bar{\nu} > k-1}$ and due to the conditional variants of (7.5) and (7.3), we get

$$\begin{aligned} &E|d_{k-1}|^{2n-2} (d_{k-1}, \chi_{\nu \wedge \bar{\nu} > k-1} (Z_k + \rho_k)) \\ &= E[|d_{k-1}|^{2n-2} (d_{k-1}, \chi_{\nu \wedge \bar{\nu} > k-1} E((Z_k + \rho_k)/\mathcal{F}_{k-1}))] \\ &\leq E[|d_{k-1}|^{2n-1} \chi_{\nu \wedge \bar{\nu} > k-1} (|E(Z_k/\mathcal{F}_{k-1})| + |E(\rho_k/\mathcal{F}_{k-1})|)] \\ &\leq K r^2 E|d_{k-1}|^{2n} + K r^4 E|d_{k-1}|^{2n-1}. \end{aligned}$$

Now consider $E|d_{k-1}|^{2n-m} \chi_{\nu \wedge \bar{\nu} > k-1} |Z_k + \rho_k|^m$. Using \mathcal{F}_{k-1} -measurability of d_{k-1} and $\chi_{\nu \wedge \bar{\nu} > k-1}$ and the conditional variants of (7.4) and (6.5), we obtain for $2 \leq m \leq 2n$,

$$\begin{aligned} & E|d_{k-1}|^{2n-m} \chi_{\nu \wedge \bar{\nu} > k-1} |Z_k + \rho_k|^m \\ &= E[|d_{k-1}|^{2n-m} \chi_{\nu \wedge \bar{\nu} > k-1} E(|Z_k + \rho_k|^m / \mathcal{F}_{k-1})] \\ &\leq KE[|d_{k-1}|^{2n-m} \chi_{\nu \wedge \bar{\nu} > k-1} E(|Z_k|^m / \mathcal{F}_{k-1})] \\ &\quad + KE[|d_{k-1}|^{2n-m} \chi_{\nu \wedge \bar{\nu} > k-1} E(|\rho_k|^m / \mathcal{F}_{k-1})] \\ &\leq KE(|d_{k-1}|^{2n} r^m + |d_{k-1}|^{2n-m} r^{2m}). \end{aligned}$$

Then

$$\begin{aligned} E|d_k|^{2n} &\leq E|d_{k-1}|^{2n} + Kr^2 E|d_{k-1}|^{2n} + Kr^4 E|d_{k-1}|^{2n-1} \\ &\quad + Kr^2 \sum_{m=2}^{2n} E|d_{k-1}|^{2n-m} r^{2m-2} \\ &\leq E|d_{k-1}|^{2n} + Kr^2 E \sum_{m=0}^{2n} |d_{k-1}|^{2n-m} r^m. \end{aligned}$$

Using the elementary inequality $ab \leq a^p/p + b^q/q$, $a, b > 0$, $p, q > 1$, $1/p + 1/q = 1$, we get

$$|d_{k-1}|^{2n-m} r^m \leq \frac{|d_{k-1}|^{2n}}{2n/(2n-m)} + \frac{r^{2n}}{2n/m}, \quad 1 \leq m < 2n.$$

Hence

$$E|d_k|^{2n} \leq E|d_{k-1}|^{2n} + Kr^2 E|d_{k-1}|^{2n} + Kr^{2n+2}, \quad d_0 = 0,$$

and we obtain for $N = L/r^2$,

$$(7.9) \quad E|d_N|^{2n} = E|X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^{2n} \leq Ke^{2\gamma L} r^{2n}.$$

Taking $n \geq 1/\varepsilon$ and substituting (7.9) in (7.8), we get

$$(7.10) \quad P(\nu < N \wedge \bar{\nu}) \leq Ke^{2\gamma L} r^2.$$

Note that K and γ depend on ε .

The inequality (7.7) together with (7.9) under $n = 1$ and (7.10) gives the statement of the theorem. \square

THEOREM 7.4. *Let $\bar{\nu} = \bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r^{1-\varepsilon}})$, $0 < \varepsilon \leq 1$, be the first exit moment of the Markov chain $(\bar{\nu}_i, \bar{X}_i)$, $i = 1, 2, \dots$, from the domain $Q \setminus \Gamma_{\alpha r^{1-\varepsilon}}$. Then, there exist constants $K > 0$ and $\gamma > 0$ such that for all r small enough the inequality,*

$$(E|X_{\bar{\nu}} - \bar{X}_{\bar{\nu}}|^2)^{1/2} \leq K (\exp(\gamma L) r + \exp(-c_r \gamma L/2))$$

holds.

PROOF. Introduce two sets $\mathcal{C} = \{\bar{\nu} \leq L/r^2\}$ and $\Omega \setminus \mathcal{C} = \{\bar{\nu} > L/r^2\}$. Let l be a diameter of G . Using Theorems 7.2 and 7.3, we obtain

$$\begin{aligned} E|X_{\bar{\nu}} - \bar{X}_{\bar{\nu}}|^2 &= E(|X_{\bar{\nu}} - \bar{X}_{\bar{\nu}}|^2; \mathcal{C}) + E(|X_{\bar{\nu}} - \bar{X}_{\bar{\nu}}|^2; \Omega \setminus \mathcal{C}) \\ &= E(|X_{N \wedge \bar{\nu}} - \bar{X}_{N \wedge \bar{\nu}}|^2; \mathcal{C}) + E(|X_{\bar{\nu}} - \bar{X}_{\bar{\nu}}|^2; \Omega \setminus \mathcal{C}) \\ &\leq E|X_N - \bar{X}_N|^2 + l^2 P(\Omega \setminus \mathcal{C}) \\ &\leq K \exp(2\gamma L) r^2 + K \exp(-c_r \gamma L). \end{aligned} \quad \square$$

REMARK 7.1. Let $\bar{\nu} = \bar{\nu}_{t,x}(Q \setminus \Gamma_{\alpha r^{1-\varepsilon}})$, $0 < \varepsilon \leq 1$, be the first exit moment of the Markov chain $(\bar{\vartheta}_i, \bar{X}_i)$, $i = 1, 2, \dots$, from the domain $Q \setminus \Gamma_{\alpha r^{1-\varepsilon}}$. Then for every natural m there exist constants $K > 0$ and $\gamma > 0$ such that for all r small enough the following inequalities:

$$(7.11) \quad E|X_N - \bar{X}_N|^{2m} \leq K \exp(2\gamma L) r^{2m},$$

$$(7.12) \quad E|X_{\bar{\nu}} - \bar{X}_{\bar{\nu}}|^{2m} \leq K(\exp(2\gamma L) r^{2m} + \exp(-c_r \gamma L))$$

and

$$(7.13) \quad P(\tau_{t,x} < \bar{\vartheta}_N) \leq K \exp(2\gamma L) r^{2n}, \quad n = 1, 2, \dots,$$

hold.

Indeed, taking $n \geq m/\varepsilon$ in (7.9) and in (7.8), we get

$$(7.14) \quad P(\nu < N \wedge \bar{\nu}) \leq K e^{2\gamma L} r^{2m}$$

instead of (7.10).

Similarly to (7.7), we have

$$E|X_N - \bar{X}_N|^{2m} \leq E|X_{N \wedge \nu} - \bar{X}_{N \wedge \nu}|^{2m} + l^{2m} P(\nu < N \wedge \bar{\nu}).$$

Now inequality (7.11) can be easily obtained from (7.9) under $n = m$.

Inequality (7.12) follows by the arguments as in the proof of Theorem 7.4.

Let us prove inequality (7.13). Remember that $X_N = X_{t,x}(\tau_{t,x} \wedge \bar{\vartheta}_N) = X_{t,x}(\tau_{t,x})$ under $\tau_{t,x} < \bar{\vartheta}_N$, and $\rho(\bar{X}_N, \partial G) \geq \beta r^{1-\varepsilon}$. Therefore

$$\begin{aligned} E\chi_{\tau_{t,x} < \bar{\vartheta}_N} |X_{t,x}(\tau_{t,x} \wedge \bar{\vartheta}_N) - \bar{X}_N|^m \\ \geq P(\tau_{t,x} < \bar{\vartheta}_N) \cdot \beta^m \cdot r^{m-\varepsilon m}, \quad m = 1, 2, \dots \end{aligned}$$

On the other hand,

$$E\chi_{\tau_{t,x} < \bar{\vartheta}_N} |X_{t,x}(\tau_{t,x} \wedge \bar{\vartheta}_N) - \bar{X}_N|^m \leq \sqrt{P(\tau_{t,x} < \bar{\vartheta}_N) [E|X_N - \bar{X}_N|^{2m}]^{1/2}}.$$

Consequently, we get

$$P(\tau_{t,x} < \bar{\vartheta}_N) \leq K r^{2\varepsilon m - 2m} E|X_N - \bar{X}_N|^{2m}.$$

Using (7.11) under $m \geq n/\varepsilon$, we come to (7.13).

8. Approximation of exit point $(\tau, X(\tau))$. Here we are interested in an approximation of the exit point $(\tau_{t,x}, X_{t,x}(\tau_{t,x}))$ of the space-time diffusion $(s, X_{t,x}(s))$, $s \geq t$, from the space-time domain Q . For the sake of simplicity in proofs we restrict ourselves to the case of the convex domain G in this section.

We have $(\bar{\vartheta}_N, \bar{X}_N) = (\bar{\vartheta}_{\bar{\nu}}, \bar{X}_{\bar{\nu}}) \in \Gamma_{\alpha r^{1-\varepsilon}}$ on the set $\mathcal{C} = \{\bar{\nu} \leq L/r^2\}$. Let $(\bar{\tau}_{t,x}, \bar{\xi}_{t,x})(\omega)$, $\omega \in \mathcal{C}$, be a point on Γ defined as if $\bar{\vartheta}_{\bar{\nu}} \geq t_1 - \alpha r^{1-\varepsilon}$ then $\bar{\tau}_{t,x} = t_1$ and $\bar{\xi}_{t,x} = \bar{X}_{\bar{\nu}} \in G$, otherwise (i.e., when $\rho(\bar{X}_{\bar{\nu}}, \partial G) \leq \alpha r^{1-\varepsilon}$) $\bar{\tau}_{t,x} = \bar{\vartheta}_{\bar{\nu}}$ and a point $\bar{\xi}_{t,x} \in \partial G$ is such that

$$(8.1) \quad |\bar{X}_{\bar{\nu}} - \bar{\xi}_{t,x}| \leq \alpha r^{1-\varepsilon}, \quad \omega \in \mathcal{C}.$$

To complete the definition of $(\bar{\tau}_{t,x}, \bar{\xi}_{t,x})(\omega)$ on the set $\Omega \setminus \mathcal{C}$, we put $\bar{\tau}_{t,x}$ be equal to $\bar{\vartheta}_N$ and $\bar{\xi}_{t,x}$ be a point on ∂G nearest to \bar{X}_N .

It is natural to take the point $(\bar{\tau}_{t,x}, \bar{\xi}_{t,x})$ as an approximate one to the exit point $(\tau_{t,x}, X_{t,x}(\tau_{t,x}))$.

Below we need the following lemma (it is analogous to the corresponding lemma from [22]).

LEMMA 8.1. *There exists a constant $K > 0$ such that for all $(t, x) \in \bar{Q}$ and $y \in \partial G$ the inequalities*

$$(8.2) \quad E(\tau_{t,x} - t) \leq K|x - y|,$$

$$(8.3) \quad E(X_{t,x}(\tau_{t,x}) - y)^2 \leq K|x - y|$$

are valid.

PROOF. Without loss of generality, one can suppose that the domain G belongs to the band, bounded by the plains $x^1 = h$, $x^1 = H$ with $0 < h < H$, and the point y has the coordinates: $y^1 = H$, $y^2 = \dots = y^d = 0$. Introduce the function (see [13])

$$V(x) := H^{2n} - (x^1)^{2n}, \quad n \geq 1.$$

Clearly,

$$0 \leq V(x) = (y^1)^{2n} - (x^1)^{2n} \leq K|y - x|,$$

where a constant $K > 0$ depends on the domain G only.

We have

$$\begin{aligned} LV &:= \frac{\partial V}{\partial t} + \frac{1}{2} \sum_{i,j=1}^d a^{ij}(t,x) \frac{\partial^2 V}{\partial x^i \partial x^j} + \sum_{i=1}^d b^i(t,x) \frac{\partial V}{\partial x^i} \\ &= -\frac{1}{2} a^{11}(t,x) 2n(2n-1)(x^1)^{2n-2} - b^1(t,x) 2n(x^1)^{2n-1} := g_n(t,x). \end{aligned}$$

As $x^1 \geq h$, $a^{11}(t,x) \geq \lambda_1^2$, and $b(t,x)$ is uniformly bounded in \bar{Q} , one can reach the inequality $g_n(t,x) \geq \gamma > 0$ on account of a choice of a sufficiently big n .

Consider the Dirichlet problem

$$(8.4) \quad Lu = -g(t, x), \quad (t, x) \in Q,$$

$$(8.5) \quad u(t, x)|_{\Gamma} = V(x).$$

Under $g = g_n$ the solution to this problem is the function $V(x)$. Therefore

$$K|y - x| \geq V(x) = EV(X_{t,x}(\tau_{t,x})) + E \int_t^{\tau_{t,x}} g_n(s, X_{t,x}(s)) ds \geq \gamma E(\tau_{t,x} - t),$$

whence inequality (8.2) follows.

Now introduce the function

$$g(t, x) := -Lv,$$

where $v(x) = (x - y)^2$ (remember that y is fixed).

Clearly, $|g(t, x)| \leq K$, $(t, x) \in \bar{Q}$. The function $u(t, x) = (x - y)^2$ is the solution of equation (8.4) with the condition

$$u(t, x)|_{\Gamma} = (x - y)^2.$$

Therefore

$$\begin{aligned} |x - y|^2 = u(t, x) &= E(X_{t,x}(\tau_{t,x}) - y)^2 + E \int_t^{\tau_{t,x}} g(s, X_{t,x}(s)) ds \\ &\geq E(X_{t,x}(\tau_{t,x}) - y)^2 - KE(\tau_{t,x} - t) \end{aligned}$$

and due to inequality (8.2) we get (8.3). \square

THEOREM 8.1. *Let $\bar{v} = \bar{v}_{t,x}(Q \setminus \Gamma_{\alpha r^{1-\varepsilon}})$, $0 < \varepsilon \leq 1$, be the first exit moment of the Markov chain $(\bar{\vartheta}_i, \bar{X}_i)$, $i = 1, 2, \dots$, from the domain $Q \setminus \Gamma_{\alpha r^{1-\varepsilon}}$. Then, there exist positive constants K and γ such that for all r small enough the inequalities*

$$(8.6) \quad [E(|X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{E})]^{1/2} \leq K r^{(1-\varepsilon)/2}$$

and

$$(8.7) \quad [E|X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2]^{1/2} \leq K(r^{(1-\varepsilon)/2} + \exp(-c_r \gamma L/2))$$

hold.

PROOF. Consider the distance between $X_{t,x}(\tau_{t,x})$ and $\xi_{t,x}$ on \mathcal{E} :

$$(8.8) \quad \begin{aligned} E(|X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{E}) &= E(\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{E}) \\ &\quad + E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{E}). \end{aligned}$$

We get for the first term of (8.8):

$$(8.9) \quad \begin{aligned} &E(\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{E}) \\ &= E\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \bar{X}_N|^2 \\ &\leq 2E\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - X_N|^2 + 2E|X_N - \bar{X}_N|^2. \end{aligned}$$

Due to Theorem 7.3, the second term of (8.9) is estimated by $Ke^{2\gamma L} r^2$, and we have for the first term of (8.9),

$$\begin{aligned}
 & E\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - X_N|^2 \\
 &= E|\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}}(X_{t,x}(\tau_{t,x}) - X_N)|^2 \\
 &\leq 2E\left|\int_{\bar{\vartheta}_N \wedge \tau_{t,x}}^{\tau_{t,x}} \chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} b(s, X_{t,x}(s)) ds\right|^2 \\
 &\quad + 2E\left|\int_{\bar{\vartheta}_N \wedge \tau_{t,x}}^{\tau_{t,x}} \chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} \sigma(s, X_{t,x}(s)) dw(s)\right|^2 \\
 &\leq KE\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} (\tau_{t,x} - \tau_{t,x} \wedge \bar{\vartheta}_N)^2 \\
 &\quad + KE\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} (\tau_{t,x} - \tau_{t,x} \wedge \bar{\vartheta}_N) \\
 &\leq KE\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} (t_1 - \bar{\vartheta}_N)^2 \\
 &\quad + KE\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} (t_1 - \bar{\vartheta}_N) \leq Kr^{1-\varepsilon},
 \end{aligned}$$

whence it follows that

$$(8.10) \quad E(\chi_{\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{C}) \leq Kr^{1-\varepsilon}.$$

Consider the second term of (8.8). Due to its definition, the point $\xi_{t,x}(\omega)$, $\omega \in \mathcal{C}$, belongs to ∂G if $\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}$. Then by the conditional version of Lemma 8.1, we get (note that $\xi_{t,x}$ is measurable with respect to \mathcal{F}_N)

$$\begin{aligned}
 & E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{C}) \\
 &= E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} E(|X_{\bar{\vartheta}_N, X_N}(\tau_{\bar{\vartheta}_N, X_N}) - \xi_{t,x}|^2 / \mathcal{F}_N); \mathcal{C}) \\
 &\leq KE(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |X_N - \xi_{t,x}|; \mathcal{C}).
 \end{aligned}$$

Theorem 7.3 and inequality (8.1) imply

$$\begin{aligned}
 & E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |X_N - \xi_{t,x}|; \mathcal{C}) \\
 (8.11) \quad & \leq [E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |X_N - \xi_{t,x}|^2; \mathcal{C})]^{1/2} \\
 & \leq [2E|X_N - \bar{X}_N|^2 + 2(E\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |\bar{X}_N - \xi_{t,x}|^2; \mathcal{C})]^{1/2} \\
 & \leq Ke^{\gamma L} r + 2\alpha r^{1-\varepsilon} \leq Kr^{1-\varepsilon}.
 \end{aligned}$$

Thus,

$$E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} |X_{t,x}(\tau_{t,x}) - \xi_{t,x}|^2; \mathcal{C}) \leq Kr^{1-\varepsilon}.$$

Substituting this inequality and inequality (8.10) in (8.8), we get (8.6).

Inequality (8.7) is obtained by Theorem 7.2 analogously to the proof of Theorem 7.4. \square

THEOREM 8.2. *Under the assumptions of Theorem 8.1, the inequalities*

$$(8.12) \quad E(|\tau_{t,x} - \bar{\tau}_{t,x}|; \mathcal{E}) \leq K r^{1-\varepsilon},$$

$$(8.13) \quad E|\tau_{t,x} - \bar{\tau}_{t,x}| \leq K(r^{1-\varepsilon} + \exp(-\alpha_r \gamma L))$$

hold.

PROOF. Remember that $\tau_{t,x} \leq t_1$, $\bar{\vartheta}_N \leq t_1$. Further, $\bar{\tau}_{t,x} = t_1$ under $\bar{\vartheta}_N \geq t_1 - \alpha r^{1-\varepsilon}$ and $\bar{\tau}_{t,x} = \bar{\vartheta}_N$ otherwise. Consequently, $\bar{\tau}_{t,x} \geq \bar{\vartheta}_N$. Let below $\tau := \tau_{t,x}$, $\bar{\tau} := \bar{\tau}_{t,x}$.

Consider the difference $|\tau - \bar{\tau}|$ on the set \mathcal{E} . We have

$$(8.14) \quad E(|\tau - \bar{\tau}|; \mathcal{E}) = E((\bar{\tau} - \tau \wedge \bar{\tau}); \mathcal{E}) + E((\tau - \tau \wedge \bar{\tau}); \mathcal{E}).$$

We get for the first term

$$\begin{aligned} E((\bar{\tau} - \tau \wedge \bar{\tau}); \mathcal{E}) &\leq E(\bar{\tau} - \tau \wedge \bar{\tau}) = E\chi_{\tau < \bar{\tau}}(\bar{\tau} - \tau \wedge \bar{\tau}) \\ &= E\chi_{\tau < \bar{\vartheta}_N}(\bar{\tau} - \tau \wedge \bar{\tau}) + E\chi_{\bar{\vartheta}_N \leq \tau < \bar{\tau}}(\bar{\tau} - \tau \wedge \bar{\tau}) \\ &\leq (t_1 - t_0)P(\tau < \bar{\vartheta}_N) + E\chi_{t_1 - \alpha r^{1-\varepsilon} \leq \tau < t_1}(t_1 - \tau). \end{aligned}$$

Then using (7.13) under $n = 1$, we obtain

$$(8.15) \quad E((\bar{\tau} - \tau \wedge \bar{\tau}); \mathcal{E}) \leq Ke^{2\gamma L} r^2 + \alpha r^{1-\varepsilon} \leq Kr^{1-\varepsilon}.$$

Consider the second term of (8.14). Due to $\xi_{t,x} \in \partial G$ under $\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}$, Lemma 8.1 and inequality (8.11), we get

$$\begin{aligned} E((\tau - \tau \wedge \bar{\tau}); \mathcal{E}) &= E(\chi_{\bar{\tau} < \tau}(\tau - \tau \wedge \bar{\tau}); \mathcal{E}) \\ &= E(\chi_{\bar{\vartheta}_N < \tau} \chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}}(\tau - \tau \wedge \bar{\vartheta}_N); \mathcal{E}) \\ &= E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}}(\tau_{\bar{\vartheta}_N, X_N} - \bar{\vartheta}_N); \mathcal{E}) \\ &= E(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} E(\tau_{\bar{\vartheta}_N, X_N} - \bar{\vartheta}_N / \mathcal{F}_N); \mathcal{E}) \\ &\leq KE(\chi_{\bar{\vartheta}_N < t_1 - \alpha r^{1-\varepsilon}} | X_N - \xi_{t,x}|; \mathcal{E}) \leq Kr^{1-\varepsilon}. \end{aligned}$$

Substituting this inequality and inequality (8.15) in (8.14), we get (8.12).

Inequality (8.13) is obtained by Theorem 7.2 analogously to the proof of Theorem 7.4. \square

9. Simulation of Brownian motion with drift provided bounded space increment. Sections 6–8 are connected with the one-step approximation $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$, $(t, x) \in Q \setminus \Gamma_{\alpha r}$ [see (6.3)], which is based on the simulation of the exit point $(\bar{\theta}, w(t + \bar{\theta}) - w(t))$ of the process $(s - t, w(s) - w(t))$, $s > t$, from the space–time parallelepiped $\Pi_r = [0, lr^2] \times C_r$ with the cubic base C_r .

It is possible to derive other constructive one-step approximations. Let us consider a one-step approximation based on a simulation of exit points for the Brownian motion with drift $W_\mu(s)$,

$$W_\mu(s) = \mu s + W(s), \quad W_\mu(0) = 0,$$

where μ is a d -dimensional fixed vector and $W(s)$ is a d -dimensional standard Wiener process.

If $(\bar{\theta}, w_\mu(t + \bar{\theta}) - w_\mu(t))$ is the first exit point of the process $(s - t, w_\mu(s) - w_\mu(t))$, $s > t$, under $\mu = \sigma^{-1}(t, x)b(t, x)$, $(t, x) \in Q \setminus \Gamma_{\alpha r}$, from the space-time parallelepiped $[0, l) \times C_r$, $l \leq t_1 - t$, then it is easy to see that the approximation

$$(9.1) \quad \bar{X}_{t,x}(t + \bar{\theta}) = x + \sigma(t, x)(w_\mu(t + \bar{\theta}) - w_\mu(t))$$

belongs to the space parallelepiped $\bar{C}_r^\sigma(x)$ even not under small l .

Then we are able to ensure that $\bar{X}_{t,x}(t + \bar{\theta})$ belongs to G , and, consequently, $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$ to Q .

The approximation (9.1) is more universal than the approximation (6.3). However, the approximation (6.3) is simpler in a computational sense than (9.1) and is quite appropriate for the majority of problems.

In this section we give algorithms on simulating exit points for Brownian motion with drift $W_\mu(s)$. The theorems on local error and global convergence connected with the one-step approximation 9.1 can be done analogously to the corresponding theorems of Sections 6–8.

9.1. Some distributions for one-dimensional Brownian motion with drift.

LEMMA 9.1. *Let τ be the first-passage time of the one-dimensional Brownian motion with drift $W_\mu(s) = \mu s + W(s)$, $W_\mu(0) = 0$, to the boundary of the interval $[-1, 1]$. Then its distribution $\mathcal{P}(t; \mu) = P(\tau < t)$ is equal to*

$$(9.2) \quad \begin{aligned} \mathcal{P}(t; \mu) &= 1 - 2\pi \exp\left(-\frac{1}{2}\mu^2 t\right)(e^\mu + e^{-\mu}) \\ &\times \sum_{k=0}^{\infty} (-1)^k \frac{(2k+1)}{\pi^2(2k+1)^2 + 4\mu^2} \exp\left(-\frac{1}{8}\pi^2(2k+1)^2 t\right) \end{aligned}$$

or

$$(9.3) \quad \begin{aligned} \mathcal{P}(t; \mu) &= 1 - \frac{1}{2} \sum_{k=0}^{\infty} (-1)^k e^{2\mu k} \left(\operatorname{erfc} \frac{2k-1+\mu t}{\sqrt{2t}} - \operatorname{erfc} \frac{2k+1+\mu t}{\sqrt{2t}} \right) \\ &- \frac{1}{2} \sum_{k=1}^{\infty} (-1)^k e^{-2\mu k} \left(\operatorname{erfc} \frac{2k-1-\mu t}{\sqrt{2t}} - \operatorname{erfc} \frac{2k+1-\mu t}{\sqrt{2t}} \right). \end{aligned}$$

PROOF. Due to (2.19), the distribution $P(\tau_x < t)$ is equal to $1 - v(t, x)$, where τ_x is the first exit time of $x + W_\mu(s) = x + \mu s + W(s)$, $-1 \leq x \leq 1$, to

the boundary of the interval $[-1, 1]$, and $v(t, x)$ obeys the following boundary value problem:

$$(9.4) \quad \frac{\partial v}{\partial t} = \frac{1}{2} \frac{\partial^2 v}{\partial x^2} + \mu \frac{\partial v}{\partial x}, \quad t > 0, \quad -1 < x < 1,$$

$$(9.5) \quad v(0, x) = 1, \quad v(t, -1) = v(t, 1) = 0.$$

The function

$$u(t, x) = \exp\left(\frac{1}{2}\mu^2 t + \mu x\right)v(t, x)$$

satisfies the boundary value problem

$$(9.6) \quad \frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}, \quad t > 0, \quad -1 < x < 1,$$

$$(9.7) \quad u(0, x) = e^{\mu x}, \quad u(t, -1) = u(t, 1) = 0.$$

Solving this problem analogously to (3.1) and (3.2) and (3.13) and (3.14), we get two expressions for $v(t, x)$,

$$\begin{aligned} v(t, x) = & \exp\left(-\frac{1}{2}\mu^2 t - \mu x\right) \sum_{k=1}^{\infty} \frac{(-1)^k \pi k}{(\pi k)^2 + 4\mu^2} (e^{-\mu} - e^{\mu}) \\ & \times \sin \pi k x \exp\left(-\frac{1}{2}\pi^2 k^2 t\right) \\ & + \exp\left(-\frac{1}{2}\mu^2 t - \mu x\right) \sum_{k=0}^{\infty} \frac{(-1)^k 2\pi(2k+1)}{\pi^2(2k+1)^2 + 4\mu^2} (e^{-\mu} + e^{\mu}) \\ & \times \cos \frac{\pi(2k+1)x}{2} \exp\left(-\frac{1}{8}\pi^2(2k+1)^2 t\right) \end{aligned}$$

and

$$v(t, x) = \exp\left(-\frac{1}{2}\mu^2 t - \mu x\right) \int_{-1}^1 G(t, x, y) \exp(\mu y) dy.$$

The equality $\mathcal{P}(t; \mu) = P(\tau < t) = 1 - v(t, 0)$ gives (9.2) and (9.3). \square

REMARK 9.1. As earlier, (9.2) is convenient for calculations under large t , and (9.3) is convenient under small t . It should be pointed out that if one of $2k \pm 1 \pm \mu t$ takes a negative value, then the corresponding $\operatorname{erfc}(2k \pm 1 \pm \mu t/\sqrt{2t}) > 1$. Therefore, it may be necessary to calculate more terms of the series in (9.3) in comparison with (3.7). However, the number of needed terms is not too large in practice due to very fast convergence of the series under small t .

REMARK 9.2. Using the Laplace transform, it is possible to derive one more expression for $\mathcal{P}(t; \mu)$:

$$\mathcal{P}(t; \mu) = \frac{2(e^{-\mu} + e^{\mu})}{\sqrt{\pi}} \sum_{k=0}^{\infty} (-1)^k \int_{2k+1/\sqrt{2t}}^{\infty} \exp\left(-\frac{\mu^2(2k+1)^2}{4z^2} - z^2\right) dz.$$

It is clear that this expression is convenient for calculations under small t .

LEMMA 9.2. *Let τ be the first-passage time of the one-dimensional Brownian motion with drift $W_\mu(s) = \mu s + W(s)$, $W_\mu(0) = 0$, to the boundary of the interval $[-1, 1]$. Then the probabilities*

$$\mathcal{P}(t; -1; \mu) := P(\tau < t, W_\mu(\tau) = -1), \quad \mathcal{P}(t; 1; \mu) := P(\tau < t, W_\mu(\tau) = 1)$$

are equal to

$$(9.8) \quad \mathcal{P}(t; -1; \mu) = \frac{1}{e^{2\mu} + 1} \mathcal{P}(t; \mu), \quad \mathcal{P}(t; 1; \mu) = \frac{e^{2\mu}}{e^{2\mu} + 1} \mathcal{P}(t; \mu).$$

PROOF. The probability $\mathcal{P}(t; -1; \mu)$ is equal to $v(t, 0)$, where $v(t, x)$ is the solution of the equation (9.4) with the initial and boundary conditions: $v(0, x) = 0$, $v(t, -1) = 1$, $v(t, 1) = 0$ [see the problem (2.15)–(2.17) under (2.9) and its solution (2.18)]. The following change of variables:

$$(9.9) \quad u(t, x) = \exp\left(\frac{1}{2}\mu^2 t + \mu x\right) \left(v(t, x) + \frac{e^{-2\mu} - e^{-2\mu x}}{e^{2\mu} - e^{-2\mu}}\right)$$

leads to the problem

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{1}{2} \frac{\partial^2 u}{\partial x^2}, & t > 0, \quad -1 < x < 1, \\ u(0, x) &= \frac{\exp(-2\mu + \mu x) - \exp(-\mu x)}{\exp(2\mu) - \exp(-2\mu)}, & u(t, -1) = u(t, 1) = 0. \end{aligned}$$

Solving this problem, we get [we restrict ourselves to writing $u(t, 0)$ only]

$$u(t, 0) = 2\pi e^{-\mu} \sum_{k=0}^{\infty} (-1)^{k+1} \frac{(2k+1)}{\pi^2(2k+1)^2 + 4\mu^2} \exp(-\pi^2(2k+1)^2 t/8)$$

or

$$u(t, 0) = \frac{e^{-2\mu}}{e^{2\mu} - e^{-2\mu}} \int_{-1}^1 G(t, 0, y) e^{\mu y} dy - \frac{1}{e^{2\mu} - e^{-2\mu}} \int_{-1}^1 G(t, 0, y) e^{-\mu y} dy.$$

Using (9.9) we obtain (9.8) for $\mathcal{P}(t; -1; \mu)$. The second formula in (9.8) is obtained analogously. \square

REMARK 9.3. Lemma 9.2 is a consequence of Reuter's theorem (see [28], page 84), which asserts that τ and $W_\mu(\tau) = \mu\tau + W(\tau)$ are independent random variables (it is not difficult to show that $P(W_\mu(\tau) = -1) = 1/(e^{2\mu} + 1)$, $P(W_\mu(\tau) = 1) = e^{2\mu}/(e^{2\mu} + 1)$). But the given proof has an independent interest because it can be used for evaluation of some other probabilities, for example, $P(\tau_x < t, W_\mu(\tau) = -1)$.

LEMMA 9.3. *For the conditional probability*

$$\mathcal{D}(\beta; t, \mu) := P(W_\mu(t) < \beta / |W_\mu(s)| < 1, \quad 0 < s < t), \quad 1 < \beta \leq 1$$

the following equalities

$$\begin{aligned} \mathcal{D}(\beta; t, \mu) &= \frac{4}{1 - \mathcal{P}(t; \mu)} \exp(-\mu^2 t/2) \sum_{k=0}^{\infty} \frac{1}{\pi^2(2k+1)^2 + 4\mu^2} \\ &\times \left((-1)^k \frac{\pi(2k+1)}{2} \exp(-\mu) \right. \\ (9.10) \quad &+ \left. \exp(\mu\beta) \left[\mu \cos \frac{\pi(2k+1)\beta}{2} + \frac{\pi(2k+1)}{2} \sin \frac{\pi(2k+1)\beta}{2} \right] \right) \\ &\times \exp\left(-\frac{1}{8} \pi^2(2k+1)^2 t\right), \end{aligned}$$

$$\begin{aligned} \mathcal{D}(\beta; t, \mu) &= \frac{1}{2(1 - \mathcal{P}(t; \mu))} \\ (9.11) \quad &\times \sum_{k=-\infty}^{\infty} \left(\exp(4\mu k) \left[\operatorname{erfc} \frac{4k - \beta + \mu t}{\sqrt{2t}} - \operatorname{erfc} \frac{4k + 1 + \mu t}{\sqrt{2t}} \right] \right. \\ &+ \left. \exp(\mu(4k + 2)) \left[\operatorname{erfc} \frac{4k + 3 + \mu t}{\sqrt{2t}} - \operatorname{erfc} \frac{4k + 2 - \beta + \mu t}{\sqrt{2t}} \right] \right) \end{aligned}$$

hold.

PROOF. We have (as in Lemma 3.2)

$$\mathcal{D}(\beta; t, \mu) = \frac{P(W_\mu(t) < \beta, \tau \geq t)}{P(\tau \geq t)}.$$

Then to prove the lemma, we need expressions for the probability $P(W_\mu(t) < \beta, \tau \geq t)$. This probability is equal to $v(t, 0)$, where $v(t, x)$ is the solution of (9.4) with initial and boundary conditions $v(0, x) = \chi_{[-1, \beta)}(x)$, $v(t, -1) = v(t, 1) = 0, t > 0$ [see the function (2.20), which is the solution of the problem (2.15)–(2.17) under (2.13)]. The following change of variables:

$$(9.12) \quad u(t, x) = \exp\left(\frac{1}{2}\mu^2 t + \mu x\right)v(t, x)$$

leads to the problem

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}, \quad t > 0, \quad -1 < x < 1,$$

$$u(0, x) = \exp(\mu x) \chi_{[-1, \beta)}(x), \quad u(t, -1) = u(t, 1) = 0.$$

Solving this problem analogously to (3.13) and (3.14) and then using (9.12), we get the statement of the lemma. \square

9.2. *Simulation of exit time and exit point of Brownian motion with drift from cube.* Let us consider a d -dimensional Brownian motion with drift $W_\mu(s)$ in the d -dimensional cube $C = \{x = (x^1, \dots, x^d): |x^i| < 1, i = 1, \dots, d\} \subset R^d$, and let τ be the first-passage time of $W_\mu(s)$, $W_\mu(0) = 0$, to the boundary ∂C of the cube C .

LEMMA 9.4. *The distribution function $\mathcal{P}_d(t; \mu)$ for τ is equal to*

$$(9.13) \quad \mathcal{P}_d(t; \mu) = P(\tau < t) = 1 - \prod_{i=1}^d (1 - \mathcal{P}(t; \mu^i)),$$

where μ^i , $i = 1, \dots, d$, are components of the vector μ .

The proof is evident.

Introduce the random variable \varkappa , which takes the value j for $\omega \in \{\omega: W_\mu^j(\tau) = \pm 1\}$.

LEMMA 9.5. *The conditional probability $P(\varkappa = j/\tau = \theta)$ is equal to*

$$(9.14) \quad \begin{aligned} P(\varkappa = j/\tau = \theta) \\ = \frac{\mathcal{P}'(\theta; \mu^j) \prod_{i \neq j} (1 - \mathcal{P}(\theta; \mu^i))}{\sum_{i=1}^d \mathcal{P}'(\theta; \mu^i) \prod_{l \neq i} (1 - \mathcal{P}(\theta; \mu^l))}, \quad j = 1, \dots, d. \end{aligned}$$

PROOF. To prove the lemma, we use two expressions for $P(\varkappa = j, \tau < \theta)$,

$$(9.15) \quad \begin{aligned} P(\varkappa = j, \tau < \theta) &= \int_0^\theta P(\varkappa = j/\tau = \vartheta) dP_\tau(\vartheta) \\ &= \int_0^\theta P(\varkappa = j/\tau = \vartheta) \sum_{i=1}^d \mathcal{P}'(\vartheta; \mu^i) \\ &\quad \times \prod_{l \neq i} (1 - \mathcal{P}(\vartheta; \mu^l)) d\vartheta, \end{aligned}$$

and

$$(9.16) \quad \begin{aligned} P(\varkappa = j, \tau < \theta) &= P\left(\bigcap_{i \neq j} (|W_\mu^i(s)| < 1, 0 < s < \tau^j), \tau^j < \theta\right) \\ &= \int_0^\theta P\left(\bigcap_{i \neq j} (|W_\mu^i(s)| < 1, 0 < s < \tau^j)/\tau^j = \vartheta\right) dP_{\tau^j}(\vartheta) \\ &= \int_0^\theta \prod_{i \neq j} (1 - \mathcal{P}(\vartheta; \mu^i)) \mathcal{P}'(\vartheta; \mu^j) d\vartheta. \end{aligned}$$

The equality $P(\bigcap_{i \neq j} (|W_\mu^i(s)| < 1, 0 < s < \tau^j)/\tau^j = \vartheta) = \prod_{i \neq j} (1 - \mathcal{P}(\vartheta; \mu^i))$ in (9.16) is proved similarly to Lemma 4.2. The expressions (9.15) and (9.16) imply (9.14). \square

LEMMA 9.6. *The following equalities:*

$$(9.17) \quad P(W_\mu^j(\tau) = -1/\varkappa = j, \tau = \theta) = \frac{1}{\exp(2\mu^j) + 1},$$

$$(9.18) \quad P(W_\mu^j(\tau) = 1/\varkappa = j, \tau = \theta) = \frac{\exp(2\mu^j)}{\exp(2\mu^j) + 1},$$

are true.

PROOF. Due to Lemma 9.2 and Remark 9.3, which state the independence of τ^j and $W_\mu^j(\tau^j)$, we get

$$\begin{aligned} P(W_\mu^j(\tau) = -1/\varkappa = j, \tau = \theta) &= P(W_\mu^j(\tau^j) = -1/\tau^j = \theta) \\ &= P(W_\mu^j(\tau^j) = -1) = \frac{1}{e^{2\mu^j} + 1}. \end{aligned}$$

The formula (9.18) is obtained analogously. \square

LEMMA 9.7. *The following equality,*

$$(9.19) \quad P\left(\bigcap_{i \neq j} (W_\mu^i(\tau) < \beta^i) / \varkappa = j, \tau = \theta\right) = \prod_{i \neq j} \mathcal{D}(\beta^i; \theta, \mu^i),$$

is valid. In particular, the relation (9.19) means that provided \varkappa and τ be known, $W_\mu^i(\tau)$, $i \neq j$, are independent.

PROOF. Carrying out reasoning as in Lemma 4.2, we get

$$\begin{aligned} P\left(\bigcap_{i \neq j} (W_\mu^i(\tau^j) < \beta^i, |W_\mu^i(s)| < 1, 0 < s < \tau^j) / \tau^j\right) \\ = \prod_{i \neq j} [(1 - \mathcal{P}(\tau^j; \mu^i)) \mathcal{D}(\beta^i; \tau^j, \mu^i)], \end{aligned}$$

whence, doing as in Lemma 4.3, we obtain

$$(9.20) \quad \begin{aligned} P\left(\varkappa = j, \tau < \theta, \bigcap_{i \neq j} (W_\mu^i(\tau) < \beta^i)\right) \\ = \int_0^\theta \prod_{i \neq j} [(1 - \mathcal{P}(\vartheta; \mu^i)) \mathcal{D}(\beta^i; \vartheta, \mu^i)] \mathcal{P}'(\vartheta; \mu^j) d\vartheta. \end{aligned}$$

We have from (9.16)

$$dP(\varkappa = j, \tau < \theta) = \prod_{i \neq j} (1 - \mathcal{P}(\theta; \mu^i)) \mathcal{P}'(\theta; \mu^j) d\theta.$$

Then

$$\begin{aligned}
 &P\left(\kappa = j, \tau < \theta, \bigcap_{i \neq j} (W_\mu^i(\tau) < \beta^i)\right) \\
 (9.21) \quad &= \int_0^\theta P\left(\bigcap_{i \neq j} (W_\mu^i(\tau) < \beta^i) / \kappa = j, \tau = \vartheta\right) \\
 &\quad \times \prod_{i \neq j} (1 - \mathcal{P}(\vartheta; \mu^i)) \mathcal{P}'(\vartheta; \mu^j) d\vartheta.
 \end{aligned}$$

Comparing (9.20) and (9.21), we come to (9.19). \square

Let us note that the point $(\tau, W_\mu(\tau))$ belongs to the lateral surface of the unbounded semicylinder $[0, \infty) \times C \subset R^{d+1}$ with the cubic base C .

THEOREM 9.1 (Algorithm for simulating exit point of space-time Brownian motion with drift to lateral surface of a cylinder with cubic base). *Let $\bar{\kappa}, \bar{\nu}, \gamma, \gamma^1, \dots, \gamma^{d-1}$ be independent, uniformly distributed on $[0, 1]$ random variables. A random point (τ, ξ_μ) , distributed as the first exit point $(\tau, W_\mu(\tau))$ of the process $(s, W_\mu(s))$ to the lateral surface of the cubic semicylinder, is simulated by the following algorithm:*

$$\tau = \mathcal{P}_d^{-1}(\gamma; \mu),$$

where $\mathcal{P}_d^{-1}(\cdot; \mu)$ is the inverse function to $\mathcal{P}_d(t; \mu)$ with respect to t ; κ is found as

$$\kappa = j \quad \text{if } \bar{\kappa} \in [\alpha_{j-1}, \alpha_j), \quad j = 1, \dots, d,$$

where

$$\alpha_0 = 0, \quad \alpha_j = \alpha_{j-1} + \frac{\mathcal{P}'(\tau; \mu^j) \prod_{i \neq j} (1 - \mathcal{P}(\tau; \mu^i))}{\sum_{i=1}^d \mathcal{P}'(\tau; \mu^i) \prod_{l \neq i} (1 - \mathcal{P}(\tau; \mu^l))};$$

ν is found as

$$\nu = \begin{cases} -1, & \bar{\nu} \in \left[0, \frac{1}{\exp(2\mu^\kappa) + 1}\right), \\ 1, & \bar{\nu} \in \left[\frac{1}{\exp(2\mu^\kappa) + 1}, 1\right]; \end{cases}$$

and then the components $\xi_\mu^i, i = 1, \dots, d$, of ξ_μ are simulated as

$$\begin{aligned}
 \xi_\mu^1 &= \mathcal{D}^{-1}(\gamma^1; \tau, \mu^1), \dots, \xi_\mu^{\kappa-1} = \mathcal{D}^{-1}(\gamma^{\kappa-1}; \tau, \mu^{\kappa-1}), & \xi_\mu^\kappa &= \nu, \\
 \xi_\mu^{\kappa+1} &= \mathcal{D}^{-1}(\gamma^\kappa; \tau, \mu^{\kappa+1}), \dots, \xi_\mu^d = \mathcal{D}^{-1}(\gamma^{d-1}; \tau, \mu^d).
 \end{aligned}$$

The statement of the theorem follows from Lemmas 9.4–9.7.

COROLLARY 9.1. Let $C_r = \{x = (x^1, \dots, x^d): |x^i| < r, i = 1, \dots, d\} \subset R^d$ be the d -dimensional cube with center at the origin and with edge length equal to $2r$. Let $\bar{\theta}$ be the first-passage time for the d -dimensional Brownian motion with drift $w_\mu(s) = \mu s + w(s)$ to the boundary ∂C_r of the cube C_r . Then the point

$$(\bar{\theta}, \bar{w}_\mu) = (r^2\tau, r\xi_{r\mu}),$$

where $(\tau, \xi_{r\mu})$ is simulated by the algorithm for simulating the exit point to lateral surface of cylinder with the cubic base C , has the same distribution as $(\bar{\theta}, w_\mu(\bar{\theta}))$.

PROOF. We have

$$W_{r\mu}\left(\frac{t}{r^2}\right) = r\mu\frac{t}{r^2} + W\left(\frac{t}{r^2}\right).$$

Because if $W(t)$ is a Wiener process, then $w(t) = rW(t/r^2)$ is also a Wiener process, we get

$$w_\mu(t) = \mu t + w(t) = rW_{r\mu}\left(\frac{t}{r^2}\right).$$

Evidently, the point $w_\mu(\bar{\theta})$ belongs to the boundary ∂C_r of the cube C_r and $w_\mu(s) \in C_r$ under $s \in [0, \bar{\theta})$. \square

REMARK 9.4. Consider an application of Theorem 9.1 in the case, when the domain G is bounded, $t_1 = \infty$, and the system (1.1) is autonomous. Then by Corollary 9.1, we are able to construct the following one-step approximation:

$$\bar{X}_{t,x}(t + \bar{\theta}) = x + \sigma(x)(w_\mu(t + \bar{\theta}) - w_\mu(t)),$$

where $\bar{\theta}$ is the first passage time of the Brownian motion with drift $w_\mu(s) - w_\mu(t)$, $s \geq t$, $\mu = \sigma^{-1}(x)b(x)$, to the boundary of the cube $C_r \subset R^d$.

The approximation $\bar{X}_{t,x}(t + \bar{\theta})$ satisfies the equation with frozen coefficients (6.2). The point $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$ belongs to the lateral surface of the semi-cylinder $[t_0, \infty) \times C_r^{\sigma(x)}(x) \subset R^{d+1}$, where the space parallelepiped $C_r^{\sigma(x)}(x)$ is obtained from the cube C_r by the linear transformation $\sigma(x)$ and the shift x . Note that $\bar{\theta}$ can take arbitrary large values with some probability.

The point $(t + \bar{\theta}, \bar{X}_{t,x}(t + \bar{\theta}))$ approximates in the mean-square sense the point $(t + \bar{\theta}, X_{t,x}(t + \bar{\theta}))$. Let us emphasize that we are able to simulate both the time and the space components of this point. Theorems on the local mean-square error, global convergence, and on an approximation of the exit point of the autonomous diffusion process $X(s)$ from the domain G can be stated and proved analogously to the corresponding theorems of [24].

9.3. *Simulation of exit point of the space-time Brownian motion with drift from a space-time parallelepiped with cubic base.* Analogously to Section 5, let us construct an algorithm for simulating the exit point $(\tau(l), W_\mu(\tau(l)))$ of the process $(s, W_\mu(s))$ from the space-time parallelepiped $\Pi = [0, l) \times C \subset R^{d+1}$. The random variable $\tau(l)$ is found as $\min(\tau, l)$, where τ is the first-passage time of $W_\mu(s)$ to the boundary ∂C as above.

THEOREM 9.2. (Algorithm for simulating exit point of space-time Brownian motion with drift from a space-time parallelepiped with cubic base). *Let $\iota, \bar{x}, \bar{v}, \gamma, \gamma^1, \dots, \gamma^{d-1}$ be independent random variables. Let ι be simulated by the law*

$$P(\iota = -1) = \mathcal{P}_d(l; \mu), \quad P(\iota = 1) = 1 - \mathcal{P}_d(l; \mu)$$

and the other random variables be uniformly distributed on $[0, 1]$.

Then a random point $(\tau(l), \xi_\mu)$, simulated by the algorithm given below, is distributed as the exit point $(\tau(l), W_\mu(\tau(l)))$.

If the simulated value of ι is equal to -1 , then the point $(\tau(l), \xi_\mu)$ belongs to the lateral surface of Π , and

$$\tau(l) = \mathcal{P}_d^{-1}(\gamma \mathcal{P}_d(l; \mu); \mu);$$

x is found as

$$x = j \quad \text{if } \bar{x} \in [\alpha_{j-1}, \alpha_j), \quad j = 1, \dots, d,$$

where

$$\alpha_0 = 0, \quad \alpha_j = \alpha_{j-1} + \frac{\mathcal{P}'(\tau(l); \mu^j) \prod_{i \neq j} (1 - \mathcal{P}(\tau(l); \mu^i))}{\sum_{i=1}^d \mathcal{P}'(\tau(l); \mu^i) \prod_{l \neq i} (1 - \mathcal{P}(\tau(l); \mu^l))};$$

v is found as

$$v = \begin{cases} -1, & \bar{v} \in \left[0, \frac{1}{\exp(2\mu^x) + 1}\right) \\ 1, & \bar{v} \in \left[\frac{1}{\exp(2\mu^x) + 1}, 1\right]; \end{cases}$$

and the components $\xi_\mu^i, i = 1, \dots, d$, of ξ_μ are simulated as

$$\xi_\mu^1 = \mathcal{Q}^{-1}(\gamma^1; \tau(l), \mu^1), \dots, \xi_\mu^{x-1} = \mathcal{Q}^{-1}(\gamma^{x-1}; \tau(l), \mu^{x-1}), \quad \xi_\mu^x = v, \\ \xi_\mu^{x+1} = \mathcal{Q}^{-1}(\gamma^x; \tau(l), \mu^{x+1}), \dots, \xi_\mu^d = \mathcal{Q}^{-1}(\gamma^{d-1}; \tau(l), \mu^d);$$

otherwise, when $\iota = 1$, the point $(\tau(l), \xi_\mu)$ belongs to the upper base of Π , and

$$\tau(l) = l, \\ \xi_\mu^1 = \mathcal{Q}^{-1}(\gamma; l, \mu^1), \quad \xi_\mu^2 = \mathcal{Q}^{-1}(\gamma^1; l, \mu^2), \dots, \xi_\mu^d = \mathcal{Q}^{-1}(\gamma^{d-1}; l, \mu^d).$$

The statement of the theorem follows from Lemmas 9.4–9.7 and reasoning similar to that done in the proof of Theorem 5.1.

The following corollary is proved as Corollary 9.1.

COROLLARY 9.2. *Let $\Pi_r = [0, lr^2) \times C_r = \{(t, x) = (t, x^1, \dots, x^d): 0 \leq t < lr^2, |x^i| < r, i = 1, \dots, d\} \subset R^{d+1}$ be a space–time parallelepiped. Let $\bar{\theta}$ be the first-passage time of the process $(s, w_\mu(s)), s > 0$, to the boundary $\partial\Pi_r$. Then the point*

$$(\bar{\theta}, \bar{w}_\mu) = (r^2\tau(l), r\xi_{r\mu}),$$

where $(\tau(l), \xi_{r\mu})$ is simulated by the algorithm for simulating the exit point from the space–time parallelepiped Π , has the same distribution as $(\bar{\theta}, w_\mu(\bar{\theta}))$.

REMARK 9.5. Let α be a d -dimensional vector, $C_\alpha = \{x = (x^1, \dots, x^d): |x^i| < \alpha^i, i = 1, \dots, d\} \subset R^d$ be the d -dimensional parallelepiped and $\Pi_\alpha = [0, l) \times C_\alpha \subset R^{d+1}$ be the corresponding space–time parallelepiped. By the results of Section 3 and the reasoning of this section (see also Remark 4.1), we can prove lemmas which are similar to Lemmas 9.4, 9.5 and 9.7 in the case when τ is the exit time of the d -dimensional Wiener process $W(s), W(0) = 0$, from the parallelepiped C_α . Then it is not difficult to state the corresponding theorems on algorithms for simulating the exit points $(\tau, W(\tau))$ of the process $(s, W(s)), s > 0$, both to the lateral surface of the cylinder $[0, \infty) \times C_\alpha$ with parallelepiped base C_α and to the boundary of the space–time parallelepiped Π_α . Using these theorems, the corresponding one-step approximation can be constructed. Note that we are also able to write down the distributions for the exit points in the case when $W(0) = x, x \neq 0, x \in C_\alpha$.

10. Numerical examples. The numerical methods proposed in this paper are widely applicable. As mentioned in the Introduction, these methods are the first ones which can constructively approximate space–time trajectories of a space–time diffusion process. They can also be applied to solving boundary value problems through a Monte Carlo technique on a level with weak methods. Let us underline that the proposed methods give an estimator for a solution to the Dirichlet problem for parabolic and elliptic equations with constant coefficients which do not contain the error of numerical integration in comparison with weak methods [19, 23].

Here we give three numerical examples. The first and the second examples deal with solving boundary value problems. In Example 2 an elliptic problem is considered, nevertheless, we need the simulation of the space–time exit points. The third example concerns the stability analysis of linear autonomous system of SDE and uses simulation of space–time trajectories essentially.

EXAMPLE 1. Let us consider an application of random walks over touching space–time parallelepipeds to the Dirichlet problem for parabolic equations (2.1)–(2.3) in the case when the coefficients are constant. This problem has the probabilistic representation (2.6)–(2.7), which we use for the Monte Carlo procedure here.

Let $(\bar{\vartheta}_k, \bar{X}_k)$ be a Markov chain which is formed analogously to the one of Section 7 but wandering is realized over touching space-time parallelepipeds (instead of small space-time parallelepipeds in Section 7) and is finished in the layer Γ_δ at a random step $\bar{\nu}$, where $\delta > 0$ is a sufficiently small constant. The equation with frozen coefficients (6.3), which we are able to simulate exactly, coincides with (2.7) when its coefficients are constant. Consequently, the chain $(\bar{\vartheta}_k, \bar{X}_k)$ coincides with the chain $(\bar{\vartheta}_k, X_k)$. In the case considered, the solution $u(t, x)$ to the Dirichlet problem (2.1)–(2.3) under $c = 0$ and $e = 0$ is simulated as [see (2.6)]

$$u(t, x) \doteq \bar{u}(t, x) = \frac{1}{M} \sum_{m=1}^M \varphi(\bar{X}_{\bar{\nu}}^{(m)}) \pm 2[\bar{D}/M]^{1/2},$$

where

$$\varphi(\bar{X}_{\bar{\nu}}^{(m)}) = \begin{cases} f(\bar{X}_{\bar{\nu}}^{(m)}), & \bar{\vartheta}_{\bar{\nu}}^{(m)} \in (t_1 - \delta, t_1], \\ g(\bar{X}_{\bar{\nu}}^{(m)}), & \bar{\vartheta}_{\bar{\nu}}^{(m)} \notin (t_1 - \delta, t_1], \end{cases}$$

$$\bar{D} = \frac{1}{M} \sum_{m=1}^M [\varphi(\bar{X}_{\bar{\nu}}^{(m)})]^2 - \left[\frac{1}{M} \sum_{m=1}^M \varphi(\bar{X}_{\bar{\nu}}^{(m)}) \right]^2,$$

and M is a number of independent Markov chains $(\bar{\vartheta}_k^{(m)}, \bar{X}_k^{(m)})$, $m = 1, \dots, M$.

Because the simulated values $(\bar{\vartheta}_k, \bar{X}_k)$ coincide with the points of exact solution $(\bar{\vartheta}_k, X_k)$ here, the estimator $\bar{u}(t, x)$ does not contain the error of numerical integration (naturally, there is Monte Carlo error depending on M and the error due to approximation of the boundary conditions depending on δ).

The mean number of steps of the random walk over touching spheres up to the boundary of space domain G is estimated by $C \ln(l/2\delta)$ (see, e.g., [8, 29] and also [23]), if G is a convex and l is its diameter. In our case the value of $\bar{\nu}$ is also estimated by $C \ln(l/2\delta)$.

Another Monte Carlo approach, whereby a random walk is made on a maximum square and the differential Laplace operator is approximated by a difference one, was proposed in [10].

As an illustration, we take the following parabolic equation in the domain $Q = [0, t_1] \times G$, $G = \{x = (x_1, x_2): |x_1| < 2, |x_2| < 1\}$ (this example is similar to one in [10]):

$$(10.1) \quad \frac{\partial u}{\partial t} = \frac{1}{2} \Delta u, \quad t > 0, |x_1| < 2, |x_2| < 1,$$

with the initial and boundary conditions

$$(10.2) \quad u(0, x) = 2,$$

$$(10.3) \quad u(t, x)|_{\partial G} = 0, \quad t > 0.$$

TABLE 1
 Test results for the boundary value problem (10.1)–(10.3).
 The exact solution $u(1, 0.7, 0.4) = 0.4796$ ($\delta = 0.00001$)

M	$\bar{u}(1, 0.7, 0.4) \pm 2[\bar{D}/M]^{1/2}$	$E\bar{v}$
1000	0.4460 ± 0.0527	3.142
4000	0.4780 ± 0.0270	3.257
100000	0.4782 ± 0.0054	3.272

By changing time $t = t_1 - s$ in (10.1)–(10.3), we obtain the corresponding boundary value problem [like (2.1)–(2.3)] with the initial condition on the upper base.

The results of numerical tests are presented in Table 1.

Throughout our tests we use a generator of uniform random numbers from [27].

EXAMPLE 2. Consider the boundary value problem for the biharmonic equation

$$(10.4) \quad L^2u + c_1(x)Lu + c_2(x)u = f(x), \quad x \in G \subset R^d,$$

$$(10.5) \quad u|_{\partial G} = \varphi(x), \quad Lu|_{\partial G} = \psi(x),$$

where L is an operator of elliptic type,

$$L = \frac{1}{2} \sum_{i,j=1}^d a^{ij}(x) \frac{\partial^2}{\partial x^i \partial x^j} + \sum_{i=1}^d b^i(x) \frac{\partial}{\partial x^i},$$

and $c_1(x)$, $c_2(x)$, $f(x)$, $\varphi(x)$ and $\psi(x)$ are some known functions.

Introducing the function $v = Lu$, we obtain the system of elliptic equations

$$(10.6) \quad Lu - v = 0, \quad x \in G, \quad u|_{\partial G} = \varphi(x),$$

$$(10.7) \quad Lv + c_1(x)v + c_2(x)u = f(x), \quad x \in G, \quad v|_{\partial G} = \psi(x).$$

Let us give a probabilistic representation of the solution to the problem (10.6)–(10.7) [the first probabilistic representation for the problem (10.6)–(10.7) in the case of constant c_1 and c_2 is obtained in [12]]. To this end introduce the system of SDE,

$$(10.8) \quad dX = b(X) ds + \sigma(X) dw(s),$$

$$\frac{dY_1}{ds} = c_2(X)Y_2,$$

$$(10.9) \quad \frac{dY_2}{ds} = -Y_1 + c_1(X)Y_2,$$

where $w(s)$ is a standard d -dimensional Wiener process, $b(x)$ is the d -dimensional vector with the components $b^i(x)$ introduced above, Y_1 and Y_2 are scalars, and $\sigma(x)$ is a matrix that is obtained from the equality

$$a(x) = \sigma(x)\sigma^T(x), \quad a(x) = \{a^{ij}(x)\}.$$

Under some conditions on the coefficients of the problem (10.6)–(10.7), its solution $(u(x), v(x))$ has the following form (see [17]):

$$\begin{aligned} (10.10) \quad u(x) &= E[\varphi(X_x(\tau))Y_1^{(1)}(\tau) + \psi(X_x(\tau))Y_2^{(1)}(\tau)] \\ &\quad - E \int_0^\tau f(X_x(s))Y_2^{(1)}(s) ds, \\ v(x) &= E[\varphi(X_x(\tau))Y_1^{(2)}(\tau) + \psi(X_x(\tau))Y_2^{(2)}(\tau)] \\ &\quad - E \int_0^\tau f(X_x(s))Y_2^{(2)}(s) ds, \end{aligned}$$

where τ is the first exit time of the process $X_x(s)$, $X(0) = x$, from the domain G , and $(Y_1^{(1)}, Y_2^{(1)})$ is the solution of the system (10.9) with initial data $Y_1^{(1)}(0) = 1$, $Y_2^{(1)}(0) = 0$, and $(Y_1^{(2)}, Y_2^{(2)})$ has initial data $Y_1^{(2)}(0) = 0$, $Y_2^{(2)}(0) = 1$.

The probabilistic representation (10.8)–(10.10) for the boundary value problem (10.4)–(10.5) can be used for solving the problem (10.4)–(10.5) by implementation of the random walk over small space-time parallelepipeds through the Monte Carlo technique. If the coefficients of the elliptic operator L and the scalars c_1 , c_2 , f are constant, we can use the random walk over touching space parallelepipeds that gives an estimator which is free from the error of numerical integration. Note that in this case the sufficient condition, under which the representation (10.10) is valid, consists of $c_1 \leq 0$, $c_2 \geq 0$.

As an illustration, consider the following two-dimensional problem in the square $G = \{x = (x_1, x_2): |x_1| < 1, |x_2| < 1\}$:

$$(10.11) \quad \frac{1}{4}\Delta^2 u = 1, \quad x \in G,$$

$$u|_{\partial G} = \varphi(x), \quad \varphi(x_1, \pm 1) = \frac{1 + x_1^4}{12}, \quad \varphi(\pm 1, x_2) = \frac{1 + x_2^4}{12},$$

$$(10.12) \quad \frac{1}{2}\Delta u|_{\partial G} = \psi(x), \quad \psi(x_1, \pm 1) = \frac{1 + x_1^2}{2}, \quad \psi(\pm 1, x_2) = \frac{1 + x_2^2}{2}.$$

Introducing the function $v = \frac{1}{2}\Delta u$ as above, we obtain the system of elliptic equations,

$$(10.13) \quad \frac{1}{2}\Delta u - v = 0, \quad x \in G, \quad u|_{\partial G} = \varphi(x),$$

$$(10.14) \quad \frac{1}{2}\Delta v = 1, \quad x \in G, \quad v|_{\partial G} = \psi(x).$$

Its exact solution is

$$u(x) = \frac{x_1^4 + x_2^4}{12}, \quad v(x) = \frac{x_1^2 + x_2^2}{2}.$$

Of course, one can solve the problem (10.13)–(10.14) sequentially: first find the function v from problem (10.14) and then u from (10.13). But such an approach requires knowledge of the function v in the whole domain G even if one needs the solution (u, v) only at individual points of the domain G . In the last case, the Monte Carlo approach is preferable.

For the system (10.13)–(10.14), (10.8)–(10.10) acquire the form

$$\begin{aligned} u(x) &= E\varphi(x + w(\tau)) - E[\tau\psi(x + w(\tau))] + \frac{1}{2}E\tau^2, \\ v(x) &= E\psi(x + w(\tau)) - E\tau, \end{aligned}$$

where τ is the first exit time of the process $x + w(s)$ from the domain G .

To simulate the point $(\tau, x + w(\tau))$, we use the random walk over touching space squares, which is finished in a δ -neighborhood of the boundary ∂G belonging to G . Remember that we are able to simulate both the exit point and the exit time of the Wiener process from a square exactly in accordance with Theorem 4.1. Then for the same reasons as in Example 1, the corresponding estimator (\bar{u}, \bar{v}) does not contain the error of numerical integration. The notice on the mean number of steps $E\bar{v}$ from Example 1 is also valid here. Let us underline that the usual method of random walk over touching spheres in the space domain G cannot be applied to this problem, because we essentially use the simulation of both the exit point $x + w(\tau)$ and the exit time τ .

The results of numerical tests are given in Table 2.

EXAMPLE 3. Let us recall some needed facts concerning the stability analysis of stochastic equations. Consider the second-order linear autonomous Itô system of SDE,

$$(10.15) \quad dX = AX dt + \sum_{i=1}^2 B_i X dw_i(t),$$

TABLE 2
Test results for the boundary value problem (10.11) and (10.12) ($\delta = 0.00001$)

M	x_1	x_2	$u(x_1, x_2)$	$\bar{u}(x_1, x_2)$	$v(x_1, x_2)$	$\bar{v}(x_1, x_2)$	$E\bar{v}$
10000	0.3	0.5	0.00588	0.0065 ± 0.0038	0.17000	0.1700 ± 0.0082	4.01
100000				0.0058 ± 0.0012		0.1700 ± 0.0026	3.99
1000000				0.00586 ± 0.00039		0.17005 ± 0.00082	4.00
10000	0.7	0.8	0.05414	0.0531 ± 0.0020	0.56500	0.5637 ± 0.0061	3.98
100000				0.05378 ± 0.00061		0.5651 ± 0.0019	4.03
1000000				0.05419 ± 0.00020		0.56536 ± 0.00062	4.00
10000	0.9	0.9	0.10935	0.1088 ± 0.0010	0.81000	0.8070 ± 0.0038	3.05
100000				0.10918 ± 0.00033		0.8096 ± 0.0012	3.01

where X is a two-dimensional vector, A and B_i , $i = 1, 2$, are constant 2×2 matrices, $w_i(t)$, $i = 1, 2$, are independent standard Wiener processes.

Various characteristics describing asymptotic behavior of solutions of the system (10.15), such as the Lyapunov exponent, moment Lyapunov exponents, the stability index and some others, are considered in [1, 2, 13] (see also references therein). The Lyapunov exponent λ^* of system (10.15) (cf. [13]) is defined as

$$(10.16) \quad \lambda^* := \lim_{t \rightarrow \infty} \frac{1}{t} E \ln |X_x(t)| = \lim_{t \rightarrow \infty} \frac{1}{t} \ln |X_x(t)| \quad \text{a.s.},$$

and the moment Lyapunov exponent $g(p)$ is defined as

$$(10.17) \quad g(p) := \lim_{t \rightarrow \infty} \frac{1}{t} E \ln |X_x(t)|^p, \quad p \in \mathbf{R},$$

where $X_x(t)$, $t \geq 0$, is a nontrivial solution to system (10.15).

The limits λ^* and $g(p)$ exist, and they are independent of x , $x \neq 0$, in the ergodic case. The limit $g(p)$ is a convex analytic function of $p \in \mathbf{R}$, $g(0) = 0$, $g(p)/p$ increases with growing p and

$$(10.18) \quad g'(0) = \lambda^*.$$

If $\lambda^* < 0$ then the trivial solution to system (10.15) is a.s. asymptotically stable. It is well known and follows from (10.18) that in this case $g(p)$ is negative for all sufficiently small $p > 0$, that is, the solution $X = 0$ of (10.15) is p -stable for such p . If $g(p) \rightarrow +\infty$ as $p \rightarrow +\infty$, then the equation

$$(10.19) \quad g(p) = 0$$

has the unique root $\gamma^* > 0$, which is known as the stability index.

It is clear that the solution $X = 0$ of (10.15) is p -stable for $0 < p < \gamma^*$ and p -unstable for $p > \gamma^*$. The stability index γ^* is connected with the asymptotic behavior of the probability $V_\delta(x)$ of the exit of $X_x(t)$ from the ball $|x| < \delta$ (see [3]): $V_\delta(x) := P\{\sup_{t \geq 0} |X_x(t)| > \delta\}$, $|x|/\delta \rightarrow 0$. It turns out that there exists a constant $K > 0$ such that for all $\delta > 0$ and $|x| < \delta$ the following inequality takes place:

$$(10.20) \quad \frac{1}{K} (|x|/\delta)^{\gamma^*} \leq V_\delta(x) \leq K (|x|/\delta)^{\gamma^*}.$$

The unstable case, when (10.19) has a negative root γ^* , is considered analogously [3].

The stability properties of the system (10.15) can also be characterized by the exit time τ of $X_x(t)$ from a certain neighborhood of the origin. In [16] the value of $Ee^{-\mu\tau}$, $\mu > 0$, is simulated. By the algorithms proposed in the present paper, we are able to evaluate the distribution function $P(\tau < t)$, which may

be a good characteristic for description of transient behavior related to the system (10.15). Naturally, we are also able to evaluate functionals on τ , for example, $Ee^{-\mu\tau}$.

We take the following particular case of the two-dimensional system (10.15) for our numerical tests:

$$\begin{aligned}
 dX_1 &= (aX_1 + cX_2) ds + b_1X_1 dw_1(s) + b_2X_2 dw_2(s), \\
 (10.21) \quad dX_2 &= (-cX_1 + aX_2) ds + b_1X_2 dw_1(s) - b_2X_1 dw_2(s), \\
 X(0) &= X_x(0) = x.
 \end{aligned}$$

The function $g(p)$, the Lyapunov exponent λ^* and the stability index γ^* for this system are equal to [21],

$$\begin{aligned}
 (10.22) \quad g(p) &= p \left(a + \frac{1}{2}(b_2^2 - b_1^2) \right) + \frac{1}{2}p^2b_1^2, \\
 \lambda^* &= g'(0) = a + \frac{1}{2}(b_2^2 - b_1^2), \\
 \gamma^* &= -\frac{2a + (b_2^2 - b_1^2)}{b_1^2}.
 \end{aligned}$$

Here we evaluate the distribution function $P(\tau < t)$, where τ is the first exit time of $X_x(s)$ under $X(0) = (1, 1)^\top$ from the square $G = \{(x_1, x_2): |x_i| < 3, i = 1, 2\}$. To simulate the system (10.21), we use the random walk over boundaries of small space–time parallelepipeds constructed in Section 7. The algorithm allows finding $\bar{\tau}$ (see Section 8), which is close to τ . The sampling distribution function $\bar{P}_M(t)$ is calculated as

$$\bar{P}_M(t) = \begin{cases} 0, & t \leq \bar{\tau}_1^{(M)}, \\ m/M, & \bar{\tau}_m^{(M)} < t \leq \bar{\tau}_{m+1}^{(M)}, \\ 1, & t > \bar{\tau}_M^{(M)}, \end{cases}$$

where $\{\bar{\tau}_1^{(M)}, \dots, \bar{\tau}_M^{(M)}\}$ is a sample point of size M sorting in the ascending order, it corresponds to the random variable $\bar{\tau}$.

The sampling function $\bar{P}_M(t)$ is close to the distribution function $\bar{P}(t) = P(\bar{\tau} < t)$ under a sufficiently big M , and $\bar{P}(t)$ is close to $P(\tau < t)$ under a sufficiently small r . We control the accuracy of our simulations by increasing M and decreasing r . We select M and r such that the curves $\bar{P}_M(t)$ are visually almost identical under larger values of M and smaller values of r .

Figure 3 presents the behavior $\bar{P}(t) \doteq \bar{P}_M(t)$ under fixed a, c, b_2 , and various b_1 . Increasing of b_1 leads to stabilization [see (10.22)]. It is interesting to note (see Figure 3) that the probability of the exit of $X_x(s)$ from G at small times t under $\lambda^* > 0$ (unstable case) is lower than the corresponding probability under $\lambda^* < 0$ (stable case). It may be explained in the following way. The




FIG. 3. The distribution function $\bar{P}(t)$ for $a = -1$, $c = 1$, $b_2 = 2$, $X(0) = (1, 1)^T$, $r = 0.02$, $M = 5000$ and for various b_1 : (1) $b_1 = 0.1$ ($\lambda^* = 0.995$, $\gamma^* = -199$); (2) $b_1 = 0.6$ ($\lambda^* = 0.82$, $\gamma^* = -4.556$); (3) $b_1 = \sqrt{5}$ ($\lambda^* = -1.5$, $\gamma^* = 0.6$) and (4) $b_1 = 3$ ($\lambda^* = -3.5$, $\gamma^* = 0.778$).

radius $\rho(s) = \sqrt{X_1^2(s) + X_2^2(s)}$ satisfies the following equation:

$$(10.23) \quad d\rho = \left(a + \frac{b_2^2}{2} \right) \rho ds + b_1 \rho dw_1(s).$$

Due to the selection of the parameters, the Lyapunov exponent λ^* is positive (unstable case) under relatively small b_1 and large b_2 . In this case, the first term of (10.23) plays the main role and the influence of noise is relatively small. So there is a lag time before the trajectory $X_x(s)$ leaves the domain G . In the stable case our parameters are such that b_1 is large and the second term of (10.23) plays an essential role. Then the trajectory $X_x(s)$ can leave the domain G during a small time interval with a rather large probability.

Figure 4 illustrates the behavior of $\bar{P}(t)$ under fixed a , c , and $\lambda^* = a + (b_2^2 - b_1^2)/2$ for various values of the stability index γ^* [see (10.22)]. One can see that the probability of the exit of the trajectory $X_x(s)$ from G decreases with increasing of γ^* in accordance with (10.20).

Figures 3 and 4 also demonstrate that in the unstable case the trajectory leaves the neighborhood of the origin during a finite time interval with the probability equal to 1 (see curves 1 and 2 on Figure 3). However, in the stable case, the probability $P(\tau < \infty)$ of leaving the neighborhood of the origin by the trajectory is less than 1. This probability decreases with decreasing of the Lyapunov exponent λ^* (see curves 3 and 4 on Figure 3) and with increasing of the stability index γ^* (see Figure 4).




FIG. 4. The distribution function $\bar{P}(t)$ for $a = -1$, $c = 1$, $X(0) = (1, 1)^T$, $\lambda^* = -1.5$, $M = 5000$ and for various γ^* : (1) $\gamma^* = 1/3$ ($b_1 = 3$, $b_2 = 2.828$, $r = 0.02$); (2) $\gamma^* = 0.6$ ($b_1 = \sqrt{5}$, $b_2 = 2$, $r = 0.02$) and (3) $\gamma^* = 2.479$ ($b_1 = 1.1$, $b_2 = 0.4683$, $r = 0.05$).

REFERENCES

- [1] ARNOLD, L., CRAUEL, H. and ECKMANN, J.-P., eds. (1991). *Lyapunov Exponents. Lecture Notes in Math.* **1486**. Springer, Berlin.
- [2] ARNOLD, L. and WIHSTUTZ, V., eds. (1986). *Lyapunov Exponents. Lecture Notes in Math.* **1186**. Springer, Berlin.
- [3] BAXENDALE, P. (1987). Moment stability and large deviations for linear stochastic differential equations. In *Proceedings Taniguchi Symposium on Probability Methods in Mathematical Physics* (N. Ikeda, ed.) 31–54. Kinokuniya, Tokyo.
- [4] BORODIN, A. N. and SALMINEN, P. (1996). *Handbook of Brownian Motion. Facts and Formulae*. Birkhäuser, Basel.
- [5] BOULEAU, N. and LÉPINGLE, D. (1994). *Numerical Methods for Stochastic Processes*. Wiley, New York.
- [6] DYNKIN, E. B. (1965). *Markov Processes*. Springer, Berlin.
- [7] DYNKIN, E. B. and YUSHKEVICH, A. A. (1969). *Markov Processes: Theorems and Problems*. Plenum, New York.
- [8] ERMAKOV, S. M., NEKRUTKIN, V. V. and SIPIN, A. S. (1989). *Random Processes for Classical Equations of Mathematical Physics*. Kluwer, Dordrecht.
- [9] GICHMAN, I. I. and SKOROCHOD, A. V. (1968). *Stochastic Differential Equations*. Naukova Dumka, Kiev.
- [10] HOSHINO, S. and ICHIDA, K. (1971). Solution of partial differential equations by a modified random walk. *Numer. Math.* **18** 61–72.
- [11] ITÔ, K. and MCKEAN, H. P. (1965). *Diffusion Processes and Their Sample Paths*. Springer, Berlin.
- [12] KHASHMINSKII, R. Z. (1960). Probabilistic representation of solutions of some differential equations. In *Trydi VI Vses. Sovechaniya on Theory of Probability and Mathematical Statistics* 177–183. Vilnius, Gospolitnauchizdat.
- [13] KHASHMINSKII, R. Z. (1969). *Stability of Differential Equations under Random Perturbations of Their Parameters*. Nauka, Moscow.

- [14] KLOEDEN, P. E. and PLATEN, E. (1992). *Numerical Solution of Stochastic Differential Equations*. Springer, Berlin.
- [15] KRYLOV, N. V. (1977). *Controllable Processes of Diffusion Type*. Nauka, Moscow.
- [16] KUSHNER, H. J. (1977). *Probability Methods for Approximations in Stochastic Control and for Elliptic Equations*. Academic Press, New York.
- [17] MILSTEIN, G. N. (1978). On a probabilistic solution of linear systems of elliptic and parabolic equations. *Theory Probab. Appl.* **23** 852–855.
- [18] MILSTEIN, G. N. (1995). *Numerical Integration of Stochastic Differential Equations*. Kluwer, Dordrecht.
- [19] MILSTEIN, G. N. (1995). Solving the first boundary value problem of parabolic type by numerical integration of stochastic differential equations. *Theory Probab. Appl.* **40** 657–665.
- [20] MILSTEIN, G. N. (1996). Application of numerical integration of stochastic equations for solving boundary value problems with the Neumann boundary conditions. *Theory Probab. Appl.* **41** 210–218.
- [21] MILSTEIN, G. N. (1996). Evaluation of moment Lyapunov exponents for second order stochastic systems. *Random Comput. Dynam.* **4** 301–315.
- [22] MILSTEIN, G. N. (1996). The simulation of phase trajectories of a diffusion process in a bounded domain. *Stochastics Stochastics Rep.* **56** 103–126.
- [23] MILSTEIN, G. N. (1997). Weak approximation of a diffusion process in a bounded domain. *Stochastics Stochastics Rep.* **62** 147–200.
- [24] MILSTEIN, G. N. (1998). On the mean-square approximation of a diffusion process in a bounded domain. *Stochastics Stochastics Rep.* **64** 211–233.
- [25] MÜLLER, M. E. (1956). Some continuous Monte Carlo methods for the Dirichlet problem. *Ann. Math. Statist.* **27** 569–589.
- [26] PARDOUX, E. and TALAY, D. (1985). Discretization and simulation of stochastic differential equations. *Acta Appl. Math.* **3** 23–47.
- [27] PRESS, W. H., TEUKOLSKY, S. A., VETTERLING, W. T. and FLANNERY, B. P. (1992). *Numerical Recipes in C: The Art of Scientific Computing*. Cambridge Univ. Press.
- [28] ROGERS, L. C. G. and WILLIAMS, D. (1987). *Diffusions, Markov Processes, and Martingales 2: Itô Calculus*. Wiley, New York.
- [29] SABELFELD, K. K. (1991). *Monte Carlo Methods in Boundary Value Problems*. Springer, Berlin.

WEIERSTRASS INSTITUTE
MOHRENSTR. 39
D-10117 BERLIN
GERMANY
E-MAIL: milstein@wias-berlin.de

URAL STATE UNIVERSITY
LENIN STR. 51
620083 EKATERINBURG
RUSSIA
E-MAIL: michael.tretyakov@usu.ru