

## EXTINCTION IN LOWER HESSENBURG BRANCHING PROCESSES WITH COUNTABLY MANY TYPES

BY PETER BRAUNSTEINS<sup>\*,1</sup> AND SOPHIE HAUTPHENNE<sup>\*,†,2</sup>

*University of Melbourne<sup>\*</sup> and Swiss Federal Institute of Technology Lausanne<sup>†</sup>*

We consider a class of branching processes with countably many types which we refer to as *Lower Hesseburg branching processes*. These are multitype Galton–Watson processes with typeset  $\mathcal{X} = \{0, 1, 2, \dots\}$ , in which individuals of type  $i$  may give birth to offspring of type  $j \leq i + 1$  only. For this class of processes, we study the set  $S$  of fixed points of the progeny generating function. In particular, we highlight the existence of a continuum of fixed points whose minimum is the global extinction probability vector  $\mathbf{q}$  and whose maximum is the partial extinction probability vector  $\tilde{\mathbf{q}}$ . In the case where  $\tilde{\mathbf{q}} = \mathbf{1}$ , we derive a global extinction criterion which holds under second moment conditions, and when  $\tilde{\mathbf{q}} < \mathbf{1}$  we develop necessary and sufficient conditions for  $\mathbf{q} = \tilde{\mathbf{q}}$ . We also correct a result in the literature on a sequence of finite extinction probability vectors that converge to the infinite vector  $\tilde{\mathbf{q}}$ .

**1. Introduction.** Multitype Galton–Watson branching processes (MGWBPs) describe the evolution of a population of independent individuals who live for a single generation and, at death, randomly give birth to offspring that may be of various types. Classical reference books on MGWBPs include Harris [17], Mode [26], Athreya and Ney [2] and Jagers [20]. MGWBPs have been used to model populations in several fields, including in molecular biology, ecology, epidemiology and evolutionary theory, as well as in particle physics, chemistry and computer science. Recent books with a special emphasis on applications are Axelrod and Kimmel [22] and Haccou, Jagers and Vatutin [16]. Branching processes with an infinite number of types have been used to model the dynamics of escape mutants [31] and the spread of parasites through a host population [3, 4]; see also [22], Chapter 7, for other biological applications of infinite-type branching processes.

One of the main quantities of interest in a MGWBP is the probability that the population eventually becomes empty or *extinct*. Let the vector  $\mathbf{Z}_n = (Z_{n,\ell})_{\ell \in \mathcal{X}}$  record the number of type- $\ell$  individuals alive in generation  $n$  of a population whose

---

Received July 2017; revised January 2019.

<sup>1</sup>Supported by the Australian Research Council (ARC) through the Centre of Excellence for the Mathematical and Statistical Frontiers (ACEMS) and the Discovery Project DP150101459.

<sup>2</sup>Supported by the Discovery Early Career Researcher Award DE150101044.

*MSC2010 subject classifications.* Primary 60J80; secondary 60J05, 60J22.

*Key words and phrases.* Infinite-type branching process, extinction probability, extinction criterion, fixed point, varying environment.

members take types that belong to the countable set  $\mathcal{X}$ . We let

$$(1.1) \quad q_i = \mathbf{P} \left[ \lim_{n \rightarrow \infty} \sum_{\ell \in \mathcal{X}} Z_{n,\ell} = 0 \mid \varphi_0 = i \right]$$

be the probability of *global extinction* given that the population begins with a single individual of type  $\varphi_0 = i$ , and we refer to  $\mathbf{q} := (q_i)_{i \in \mathcal{X}}$  as the *global extinction probability vector*. When the set  $\mathcal{X}$  contains only finitely many types, many of the fundamental questions concerning  $\mathbf{q}$  have been resolved. In particular, it is well known that (i)  $\mathbf{q}$  is the minimal nonnegative solution of the fixed-point equation  $\mathbf{s} = \mathbf{G}(\mathbf{s})$ , where  $\mathbf{G}(\mathbf{s}) := (G_i(\mathbf{s}))_{i \in \mathcal{X}}$ , defined in (2.2), records the probability generating function associated with the reproduction law of each type, and that (ii) if the process is irreducible, then the set of fixed-point solutions

$$(1.2) \quad S = \{ \mathbf{s} \in [0, 1]^{\mathcal{X}} : \mathbf{s} = \mathbf{G}(\mathbf{s}) \}$$

contains at most two elements,  $\mathbf{q}$  and  $\mathbf{1}$ . In addition, there is a well-established extinction criterion, namely  $\mathbf{q} = \mathbf{1}$  if and only if the Perron–Frobenius eigenvalue of the mean progeny matrix (defined in (2.3)) is less than or equal to one.

If we allow  $\mathcal{X}$  to contain countably infinitely many types, then this complicates matters considerably. Indeed, even the definition of extinction is no longer unambiguous. We let

$$(1.3) \quad \tilde{q}_i = \mathbf{P} \left[ \lim_{n \rightarrow \infty} Z_{n,\ell} = 0, \forall \ell \in \mathcal{X} \mid \varphi_0 = i \right],$$

be the probability of *partial extinction* given that the population begins with a single individual of type  $i$ , and we refer to  $\tilde{\mathbf{q}} = (\tilde{q}_i)_{i \in \mathcal{X}}$  as the *partial extinction probability vector*. While global extinction implies partial extinction, there may be a positive chance that every type eventually disappears from the population while the total population size grows without bound; it is then possible that  $\mathbf{q} < \tilde{\mathbf{q}}$  (see [18], Section 5.1, for an example).

At least partly due to these challenges, the set  $S$  is yet to be fully characterised in the infinite-type setting. There is, however, a number of papers that make progress towards this goal: Moyal [27] gives general conditions for  $S$  to contain at most a single solution  $\mathbf{s}$  such that  $\sup_{i \in \mathcal{X}} s_i < 1$ ; Spataru [32] gives a stronger results by stating that  $S$  contains at most two elements,  $\mathbf{q}$  and  $\mathbf{1}$ ; however, Bertacchi and Zucca [6, 7] prove the inaccuracy of the latter by providing an irreducible example where  $S$  contains uncountably many elements such that  $\sup_{i \in \mathcal{X}} s_i = 1$ . Both  $\mathbf{q}$  and  $\tilde{\mathbf{q}}$  are elements of the set  $S$ . It is well known that  $\mathbf{q}$  is the minimal element, but as yet, there has been no attempt to identify the precise location of  $\tilde{\mathbf{q}}$ . We observe that due to the existence of irreducible MGWBPs with  $\mathbf{q} < \tilde{\mathbf{q}} < \mathbf{1}$ , the partial extinction probability vector  $\tilde{\mathbf{q}}$  may be neither the minimal element of  $S$ , which is  $\mathbf{q}$ , nor the maximal element of  $S$ , which is  $\mathbf{1}$ .

Extending the extinction criterion established in the finite-type case to the infinite-type setting has also proven difficult. To resolve the problem in the infinite-type setting, we should give both a partial and a global extinction criterion. A number of authors have progressed in this direction [6, 10, 17, 18, 27, 32, 33]. In the infinite-type case, the analogue of the Perron–Frobenius eigenvalue is the *convergence norm*  $\nu(M)$  of  $M$  defined in (2.4), which gives a partial extinction criterion:  $\tilde{q} = \mathbf{1}$  if and only if  $\nu(M) \leq 1$ ; see [33], Theorem 4.1. However, when partial extinction is almost sure we are still lacking general necessary and sufficient conditions for  $q = \mathbf{1}$ . It turns out that there can be no global extinction criterion based solely upon  $M$ , as highlighted through [33], Example 4.4, but as pointed out by the author, other moment conditions have not been clearly identified. In addition, when  $\tilde{q} < \mathbf{1}$ , following the terminology in [6], the process can exhibit *strong local survival*  $q = \tilde{q} < \mathbf{1}$ , or *nonstrong local survival*  $q < \tilde{q} < \mathbf{1}$ . It is again challenging to derive a general criterion separating the two cases.

The main contribution of this paper is to use a unified probabilistic approach to characterise the set  $S$  and to derive a global extinction criterion applicable when  $\tilde{q} = \mathbf{1}$  for a class of branching processes with countably infinitely many types called *lower Hessenberg branching processes* (LHBPs). In these processes, which have the typeset  $\mathcal{X} = \{0, 1, 2, \dots\}$ , the primary constraint is that *type- $i$  individuals can produce offspring of type no larger than  $i + 1$* ; as a consequence, their (infinite) mean progeny matrices have a lower Hessenberg form. The probabilistic approach we employ relies on a single pathwise argument: we reduce the study of the LHBP to that of a much simpler Galton–Watson process in a varying environment (GWPVE), embedded in the LHBP. GWPVEs are single-type Galton–Watson processes whose offspring distributions vary deterministically with the generation. In our context, the embedded GWPVE is *explosive*, in the sense that individuals may have an infinite number of offspring. In particular, we show the equivalence between global extinction of the LHBP and extinction of the embedded GWPVE, and between partial extinction of the LHBP and the event that all generations of the embedded GWPVE are finite. Based on this relationship, we obtain several results for LHBPs:

(i) We prove that there is a continuum of fixed-points solutions  $s \in S$ , whose componentwise minimum and maximum are the global and partial extinction probability vectors  $q$  and  $\tilde{q}$ , respectively (Theorem 4.1).

(ii) We establish a connection between the growth rates of the embedded GWPVE and the convergence rate of  $s_i$  to 1 as  $i \rightarrow \infty$  for any  $s \in S \setminus \{q, \mathbf{1}\}$ ; this yields a physical interpretation for the fixed points lying in between  $q$  and  $\tilde{q}$  (Theorem 4.4).

(iii) In the nontrivial case where  $\tilde{q} = \mathbf{1}$ , we provide a necessary and sufficient condition for global extinction which holds under some second moment conditions (Theorem 5.1). This is the first extinction criterion for irreducible processes that also applies to cases exhibiting nonexponential growth. We illustrate the broad applicability of the criterion through some examples.

(iv) Finally, under additional assumptions, we build on the global extinction criterion to derive necessary and sufficient conditions for strong local survival (Theorem 7.1).

While there is a vast literature on GWPVEs, the explosive case, which has already been studied for standard Galton–Watson processes [28, 29], is yet to be considered in the context of varying environment. In order to prove our main theorems, we both apply known results on GWPVEs and develop new ones. On the way to studying properties of the embedded GWPVE, we also derive a new partial extinction criterion for LHBP which is computationally more efficient than other existing criteria.

The paper is organised as follows. In Section 2, we define LHBP and introduce the tools we use to study them. In Section 3, we construct the embedded GWPVE and derive relationships between it and its corresponding LHBP. In Section 4, we develop (i) and (ii). In Section 5, we deal with (iii). In Section 6, we illustrate the results of Section 5 through two examples. In Section 7, we address (iv). Finally, in Section 8 we discuss possible extensions of our results.

A number of our results use a sequence of truncated branching processes, labelled  $\{\tilde{\mathbf{Z}}_n^{(k)}\}_{n \geq 0}$ , which are formed by taking the original process  $\{\mathbf{Z}_n\}$  and deleting the descendants of each individual whose type is larger than  $k$ . In particular, these results require the extinction probability vectors of  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ , labelled  $\tilde{\mathbf{q}}^{(k)}$ , to converge componentwise to  $\tilde{\mathbf{q}}$  as  $k \rightarrow \infty$ . This convergence was established in Lemma 3.2 of [18] under general conditions, however, it turns out that this lemma is incorrect as it stands. In Appendix A, we point out where the lemma and its proof break down, and in Theorem A.1 we recover its assertion under stricter conditions by using an alternative approach to prove it.

In this paper, we let  $\mathbf{1}$  and  $\mathbf{0}$  denote the column vectors of 1's and 0's, respectively, and we let  $\mathbf{e}_i$  represent the vector with all entries equal to zero, except entry  $i$  which is equal to 1, the size of these vectors being defined by the context. For any vectors  $\mathbf{x}$  and  $\mathbf{y}$ , we write  $\mathbf{x} \leq \mathbf{y}$  if  $x_i \leq y_i$  for all  $i$ , and  $\mathbf{x} < \mathbf{y}$  if  $\mathbf{x} \leq \mathbf{y}$  with  $x_i < y_i$  for at least one entry  $i$ .

**2. Preliminaries.** Consider a MGWBP with the type set  $\mathcal{X} = \mathbb{N}_0 := \{0, 1, 2, \dots\}$ . We assume that the process initially contains a single individual whose type is denoted by  $\varphi_0$ . The process then evolves according to the following rules:

- (i) each individual lives for a single generation, and
- (ii) at death individuals of type  $i$  give birth to  $\mathbf{r} = (r_\ell)_{\ell \in \{0, 1, \dots, i+1\}}$  offspring, that is,  $r_0$  individuals of type 0,  $r_1$  individuals of type 1,  $\dots$ , and  $r_{i+1}$  individuals of type  $i + 1$ , where the vector  $\mathbf{r}$  is chosen independently of that of all other individuals according to a probability distribution,  $p_i(\cdot)$ , specific to the parental type  $i \in \mathcal{X}$ .

We refer to this as a *lower Hessemberg branching process* (LHBP).

We construct the LHBP on the Ulam–Harris space [17], Chapter VI, labelled  $(\Omega, \mathcal{F}, \mathbf{P})$ , as follows. Let  $\mathcal{J} = \bigcup_{n \geq 0} \mathcal{J}_n$  where  $\mathcal{J}_n$  describes the virtual  $n$ th generation. That is,  $\mathcal{J}_0 = \mathcal{X}$ , where  $\varphi_0 \in \mathcal{J}_0$  specifies the type of the root, and for  $n \geq 1$ ,  $\mathcal{J}_n = \mathcal{X} \times (\mathbb{N} \times \mathcal{X} \times \mathbb{N})^n$ , where  $(\varphi_0; i_1, j_1, y_1; \dots; i_n, j_n, y_n)$  denotes the  $i_n$ th child of type  $j_n$  born to  $(\varphi_0; i_1, j_1, y_1; \dots; i_{n-1}, j_{n-1}, y_{n-1})$  and  $y_n$  denotes the individual’s unique identification number. The identification number is not required to define the branching process with countably many types; however, it is necessary to formally define the embedded branching processes considered in the sequel (see [11], Footnote 2, for an illustration). Each virtual individual  $I \in \mathcal{J}$  is assigned a random offspring vector  $N(I) = (N_\ell(I))_{\ell \in \mathcal{X}}$  that takes values in  $R_j := \{\mathbf{r} \in (\mathbb{N}_0)^\mathcal{X} : r_\ell = 0 \ \forall \ell > j + 1\}$  when  $I$  is of type  $j$  and has distribution  $p_j(\cdot)$ , independently of all other individuals. The random set of individuals who appear in the population,  $X = \bigcup_{n \geq 0} X_n$ , is then defined recursively from the values of  $N(I)$  as follows:

$$(2.1) \quad X_0 = \{\varphi_0\}, \quad X_n = \{x = (\tilde{x}; i_n, j_n, n) \in \mathcal{J}_n : \tilde{x} \in X_{n-1}, i_n \leq N_{j_n}(\tilde{x})\}.$$

The population in generation  $n$  is described by the vector  $\mathbf{Z}_n$  with entries

$$\mathbf{Z}_{n,j} = \sum_{I \in \mathcal{J}_n} \mathbb{1}(I \in X_n, j_n = j), \quad j \in \mathcal{X}.$$

We will often refer to branching processes by their sequence of population vectors  $\{\mathbf{Z}_n\}_{n \geq 0}$ .

We define the *progeny generating vector*  $\mathbf{G}(\cdot) : [0, 1]^\mathcal{X} \rightarrow [0, 1]^\mathcal{X}$ , where

$$(2.2) \quad G_i(\mathbf{s}) = G_i(s_0, s_1, \dots, s_{i+1}) = \sum_{\mathbf{r} \in R_i} p_i(\mathbf{r}) \mathbf{s}^\mathbf{r} = \sum_{\mathbf{r} \in R_i} p_i(\mathbf{r}) \prod_{k=0}^{i+1} s_k^{r_k},$$

and the *mean progeny matrix*  $M = (M_{i,j})_{i,j \in \mathcal{X}}$ , where

$$(2.3) \quad M_{i,j} = \left( \frac{\partial G_i(\mathbf{s})}{\partial s_j} \right) \Big|_{\mathbf{s}=\mathbf{1}}$$

is the expected number of type- $j$  children born to a parent of type  $i$ . By assumption,  $M$  is an infinite lower Hessenberg matrix. To avoid trivialities, we assume that  $M_{i,i+1} > 0$  for all  $i \in \mathcal{X}$ . To  $M$ , we associate a weighted directed graph, referred to as the *mean progeny representation graph*. This graph has vertex set  $\mathcal{X}$  and contains an edge from  $i$  to  $j$  of weight  $M_{i,j}$  if and only if  $M_{i,j} > 0$ . The branching process is said to be *irreducible* if there is a path between any two vertices in the mean progeny representation graph on  $\mathcal{X}$ . We define the *convergence norm* of  $M$ ,

$$(2.4) \quad \nu(M) = \limsup_n \sqrt[n]{(M^n)_{ij}},$$

which, when the process is irreducible, is independent of  $i$  and  $j$ .

The global and partial extinction probability vectors  $\mathbf{q}$  and  $\tilde{\mathbf{q}}$ , defined in (1.1) and (1.3), are both solutions to the fixed-point equation  $s = \mathbf{G}(s)$ , and are thus elements of the set  $S$  defined in (1.2). This can be seen by conditioning on the children of the initial individual and then observing that the process becomes partially (globally) extinct if and only if the daughter processes of these children become partially (globally) extinct. Moreover, following the standard arguments, we can prove that  $\mathbf{q}$  is the componentwise minimal element of  $S$  (see [27], Theorem 3.1). By the lower Hessenberg assumption,  $s = \mathbf{G}(s)$  can be written as  $s_i = G_i(s_0, \dots, s_i, s_{i+1})$  for all  $i \geq 0$ . Thus, by the monotonicity of  $G_i(\cdot)$ , each entry  $s_i$  of any  $s \in S$  is uniquely determined by  $s_0$ . It is then natural to consider the one-dimensional projection sets of  $S$ ,

$$S_i = \{x \in [0, 1] : \exists s \in S, \text{ such that } s_i = x\}, \quad i \in \mathcal{X}.$$

We define two sequences of finite-type branching processes on  $(\Omega, \mathcal{F}, \mathbf{P})$ . The first,  $\{\tilde{\mathbf{Z}}_n^{(k)}\}_{n \geq 0, k \geq -1}$ , is such that the random offspring vector of any virtual individual  $I \in \mathcal{J}$  is given by

$$\tilde{N}^{(k)}(\omega, I) = \begin{cases} N(\omega, I), & t(I) \leq k, \\ \mathbf{0}, & t(I) > k, \end{cases}$$

for any  $\omega \in \Omega$ , where  $t(I)$  is the type of virtual individual  $I$ . For any  $k \geq -1$ , outcomes of  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$  are thus constructed by taking the corresponding outcome of  $\{\mathbf{Z}_n\}$  and removing the descendants of all individuals of type  $i > k$ . These types are said to be *sterile*. The second,  $\{\mathbf{Z}_n^{(k)}\}_{n \geq 0, k \geq -1}$ , is such that the random offspring vector of any virtual individual  $I \in \mathcal{J}$  is given by

$$N^{(k)}(\omega, I) = \begin{cases} N(\omega, I), & t(I) \leq k, \\ \mathbf{e}_{t(I)}, & t(I) > k, \end{cases}$$

for any  $\omega \in \Omega$ . For any  $k \geq -1$ , outcomes of  $\{\mathbf{Z}_n^{(k)}\}$  are thus constructed by taking the corresponding outcome of  $\{\mathbf{Z}_n\}$  and replacing the descendants of all individuals of type  $i > k$  with an infinite string of type- $i$  descendants. These types are said to be *immortal*. An illustration of  $\{\mathbf{Z}_n\}$ ,  $\{\tilde{\mathbf{Z}}_n^{(1)}\}$  and  $\{\mathbf{Z}_n^{(1)}\}$  for a specific  $\omega \in \Omega$  is given in Figure 1. By construction, for all  $\omega \in \Omega$ , (i) for each fixed value of  $k$ , if




FIG. 1. The processes  $\{\mathbf{Z}_n\}$ ,  $\{\tilde{\mathbf{Z}}_n^{(1)}\}$  and  $\{\mathbf{Z}_n^{(1)}\}$  for a specific  $\omega \in \Omega$ .

$\varphi_0 \leq k + 1$  then the sterile and immortal individuals are necessarily of type  $k + 1$ ,  
(ii)

$$(2.5) \quad Z_{n,\ell}^{(k)}(\omega) = \tilde{Z}_{n,\ell}^{(k)}(\omega) \quad \text{for all } n \geq 0 \text{ and } 0 \leq \ell \leq k,$$

and (iii)

$$(2.6) \quad \lim_{n \rightarrow \infty} Z_{n,k+1}^{(k)}(\omega) = \sum_{n=0}^{\infty} \tilde{Z}_{n,k+1}^{(k)}(\omega).$$

We denote the progeny generating vector of  $\{Z_n^{(k)}\}$  by  $G^{(k)}(s)$ , which has entries

$$(2.7) \quad G_i^{(k)}(s) = \begin{cases} G_i(s), & 0 \leq i \leq k, \\ s_{k+1}, & i = k + 1. \end{cases}$$

By equation (2.5), the global extinction probability vectors of  $\{Z_n^{(k)}\}$  and  $\{\tilde{Z}_n^{(k)}\}$ , denoted by  $\tilde{q}^{(k)}$  and  $q^{(k)}$ , are given by

$$\tilde{q}^{(k)} = \lim_{n \rightarrow \infty} G^{(k,n)}(0, \dots, 0, 1) \quad \text{and} \quad q^{(k)} = \lim_{n \rightarrow \infty} G^{(k,n)}(0, \dots, 0, 0),$$

where  $G^{(k,n)}(\cdot)$  is the  $n$ -fold composition of  $G^{(k)}(\cdot)$ . As demonstrated in [18], Lemma 3.1, the sequence  $\{q^{(k)}\}_{k \geq -1}$  increases componentwise to  $q$ . In addition, if  $\{Z_n\}$  is irreducible and nonsingular (i.e., there exists  $i \in \mathcal{X}$  such that  $\sum_{v:|v|=1} p_i(v) < 1$ ), then the sequence  $\{\tilde{q}^{(k)}\}_{k \geq -1}$  decreases componentwise to  $\tilde{q}$  (see Theorem A.1 in Appendix A); recall that [18], Lemma 3.2, is an incorrect version of the same result. Unless stated otherwise, we assume that  $\{Z_n\}$  is nonsingular and irreducible.

**3. An embedded GWPVE with explosions.** We construct the embedded GWPVE  $\{Y_k\}$  on  $(\Omega, \mathcal{F}, \mathbf{P})$  from the paths of  $\{Z_n\}$  by selecting all individuals whose type is strictly larger than that of all their ancestors, and connecting each selected individual to their nearest (in generation) selected ancestor (see Figure 2). More formally, we define a function  $f(\cdot) : \mathcal{J} \rightarrow \mathcal{J}$  that takes a line of descent  $(\varphi_0; i_1, j_1, y_1; \dots; i_n, j_n, y_n)$  and deletes each triple  $(i_k, j_k, y_k)$  whose type is not strictly larger than all its ancestors. For each  $\omega \in \Omega$ , the family tree of  $\{Y_k\}$  is then given by  $f(X(\omega))$ , where  $X(\omega)$  is defined in (2.1). Variants of  $\{Y_k\}$  (which do not permit explosion) can be found in [10] and [15].

We take the convention that  $\{Y_k\}$  starts at the generation number corresponding to the initial type  $\varphi_0$  in  $\{Z_n\}$ . By construction, for any  $\omega \in \Omega$  we then have

$$(3.1) \quad Y_k(\omega) = \sum_{n=0}^{\infty} \tilde{Z}_{n,k}^{(k-1)}(\omega) = \lim_{n \rightarrow \infty} Z_{n,k}^{(k-1)}(\omega),$$

that is, the  $k$ th generation of  $\{Y_k\}$  is made up every sterile (type- $k$ ) individual produced over the lifetime of  $\{\tilde{Z}_n^{(k-1)}\}_{n \geq 0}$ . By the lower Hessenberg assumption, each




FIG. 2. An outcome of  $\{Z_n\}$  and  $\{Y_k\}$  for a specific  $\omega \in \Omega$ . The highlighted type- $k$  individuals represent the sterile individuals in the corresponding realisation of  $\{\tilde{Z}_n^{(k-1)}\}$ .

sterile type- $k$  individual that appears in  $\{\tilde{Z}_n^{(k-1)}\}_{n \geq 0}$  is a descendant of a sterile type- $(k - 1)$  individual that appears in  $\{\tilde{Z}_n^{(k-2)}\}_{n \geq 0}$ . Thus, because the daughter processes of these type- $(k - 1)$  individuals in  $\{\tilde{Z}_n^{(k-1)}\}$  are i.i.d.,  $Y_k$  satisfies the branching process equation

$$(3.2) \quad Y_k \stackrel{d}{=} \sum_{i=1}^{Y_{k-1}} \xi_{k,i},$$

where  $\{\xi_{k,i}\}_{i \geq 1}$  is a sequence of i.i.d. random variables such that,  $\xi_{k,i} \stackrel{d}{=} \sum_{n=0}^{\infty} \tilde{Z}_{n,k}^{(k-1)}$  conditional on  $\varphi_0 = k - 1$ . This means  $\{Y_k\}$  is indeed a GWPVE; it is, however, not a classical one because  $\{Y_k\}$  may have a positive chance of *explosion*, that is, individuals in  $\{Y_k\}$  may give birth to an infinite number of offspring with positive probability. The next lemma states that  $\{Y_k\}$  explodes by generation  $k$  if and only if  $\{\tilde{Z}_n^{(k-1)}\}_{n \geq 0}$  survives globally, and  $\{Y_k\}$  becomes extinct by generation  $k$  if and only if  $\{Z_n^{(k-1)}\}_{n \geq 0}$  becomes globally extinct.

LEMMA 1. For any  $k \geq \varphi_0$ ,

$$(3.3) \quad \{\omega \in \Omega : Y_k(\omega) < \infty\} \stackrel{\text{a.s.}}{=} \left\{ \omega \in \Omega : \lim_{n \rightarrow \infty} \tilde{Z}_n^{(k-1)}(\omega) = \mathbf{0} \right\},$$

and

$$(3.4) \quad \{\omega \in \Omega : Y_k(\omega) = 0\} \stackrel{\text{a.s.}}{=} \left\{ \omega \in \Omega : \lim_{n \rightarrow \infty} Z_n^{(k-1)}(\omega) = \mathbf{0} \right\}.$$

PROOF. To prove (3.3), first suppose that  $\omega \in \{\lim_n \tilde{Z}_n^{(k-1)} = \mathbf{0}\}$ . Then there exists a generation  $N < \infty$  such that  $\tilde{Z}_n^{(k-1)}(\omega) = \mathbf{0}$  for all  $n \geq N$ . This means  $\sum_{n=0}^{\infty} \tilde{Z}_{n,k}^{(k-1)}(\omega) = \sum_{n=0}^N \tilde{Z}_{n,k}^{(k-1)}(\omega) < \infty$ , which implies  $\omega \in \{Y_k < \infty\}$ . It then remains to prove  $\mathbf{P}(Y_k < \infty, \liminf_n |\tilde{Z}_n^{(k-1)}| > 0) = 0$ . Because  $M_{i,i+1} > 0$  for



any  $i \leq k - 1$ , we have  $\mathbf{P}_i(\sum_{n=0}^k \tilde{Z}_{n,k}^{(k-1)} = 0) \leq 1 - \varepsilon_i$ , for some  $\varepsilon_i > 0$ . Thus, for any  $\tilde{z}_0^{(k-1)} \in (\mathbb{N}_0)^{k+1}$ , we have

$$\mathbf{P}\left(\sum_{n=0}^k \tilde{Z}_{n,k}^{(k-1)} = 0, |\tilde{Z}_k^{(k-1)}| > 0 \mid \tilde{Z}_0^{(k-1)} = \tilde{z}_0^{(k-1)}\right) \leq 1 - \varepsilon,$$

where  $\varepsilon := \min_{0 \leq i \leq k} \{\varepsilon_i\} > 0$ . By the Markov property, we then have

$$\begin{aligned} & \mathbf{P}\left(Y_k < \infty, \liminf_{n \rightarrow \infty} |\tilde{Z}_n^{(k-1)}| > 0\right) \\ & \leq \mathbf{P}\left(\exists N \geq 0 : \sum_{m=N}^{\infty} \tilde{Z}_{m,k}^{(k-1)} = 0, |\tilde{Z}_n^{(k-1)}| > 0 \forall n\right) \\ & = \mathbf{P}\left(\exists N \geq 0 : \bigcap_{\ell=0}^{\infty} \left(\sum_{m=N+\ell k}^{N+(\ell+1)k} \tilde{Z}_{m,k}^{(k-1)} = 0, |\tilde{Z}_{N+(\ell+1)k}^{(k-1)}| > 0\right)\right) \\ & \leq \sum_{N=0}^{\infty} \prod_{\ell \geq 0} (1 - \varepsilon) = 0, \end{aligned}$$

leading to (3.3). The same arguments lead to (3.4).  $\square$

Since

$$\left\{\omega \in \Omega : \liminf_n \tilde{Z}_n^{(k-1)}(\omega) > \mathbf{0}\right\} \subseteq \left\{\omega \in \Omega : \liminf_n \tilde{Z}_n^{(k)}(\omega) > \mathbf{0}\right\},$$

and

$$\left\{\omega \in \Omega : \lim_n Z_n^{(k-1)}(\omega) = \mathbf{0}\right\} \subseteq \left\{\omega \in \Omega : \lim_n Z_n^{(k)}(\omega) = \mathbf{0}\right\},$$

the process  $\{Y_k\}$  has two absorbing states, 0 and  $\infty$ . The next corollary formalises the equivalence between the following events:

$\{Z_n\}$ experiences:	$\{Y_k\}$ reaches:
both partial and global extinction	$\equiv$ the absorbing state 0
neither partial nor global extinction	$\equiv$ the absorbing state $\infty$
partial extinction but not global extinction	$\equiv$ neither 0 nor $\infty$ .

**COROLLARY 1.** *The global extinction event  $\mathcal{E}_g \stackrel{\text{a.s.}}{=} \{\omega \in \Omega : \lim_{k \rightarrow \infty} Y_k(\omega) = 0\}$ , and the partial extinction event  $\mathcal{E}_p \stackrel{\text{a.s.}}{=} \{\omega \in \Omega : Y_k(\omega) < \infty, \forall k \geq \varphi_0\}$ .*

**PROOF.** The result follows from Lemma 1 and the arguments in the proofs of [18], Lemma 3.1, and Theorem A.1, respectively.  $\square$

By Corollary 1, we can express any question about the extinction probability vectors  $\mathbf{q}$  and  $\tilde{\mathbf{q}}$  in terms of the process  $\{Y_k\}$ . In the sequel we use the shorthand notation  $\mathbf{P}_i(\cdot)$  for  $\mathbf{P}(\cdot | Y_i = 1)$  and  $\mathbf{E}_i(\cdot)$  for  $\mathbf{E}(\cdot | Y_i = 1)$ .

COROLLARY 2. For any  $k \geq 0$  and  $0 \leq i \leq k$ ,

$$q_i^{(k)} = \mathbf{P}_i(Y_{k+1} = 0) \quad \text{and} \quad \tilde{q}_i^{(k)} = \mathbf{P}_i(Y_{k+1} < \infty),$$

and for any  $i \geq 0$ ,

$$q_i = \mathbf{P}_i\left(\lim_{k \rightarrow \infty} Y_k = 0\right) \quad \text{and} \quad \tilde{q}_i = \mathbf{P}_i(\forall k \geq i, Y_k < \infty).$$

PROOF. The results are immediate consequences of Lemma 1 and Corollary 1. □

To take advantage of Corollary 2, we require the progeny generating function of each generation of the embedded GWPVE. For  $k \geq 0$ , we let

$$(3.5) \quad g_k(s) := \mathbf{E}_k(s^{Y_{k+1}} \mathbb{1}_{\{Y_{k+1} < \infty\}}) = \sum_{\ell \geq 0} \mathbf{P}\left(\sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)} = \ell \mid \varphi_0 = k\right) s^\ell,$$

where  $s \in [0, 1]$ . Due to the possibility of explosion, we may have  $1 > g_k(1) = \mathbf{P}_k(Y_{k+1} < \infty) = \tilde{q}_k^{(k)}$ . By (3.2), the generating function of  $Y_{k+1}$ , conditional on  $Y_i = 1$  for  $i \leq k$ , is given by

$$g_{i \rightarrow k}(s) := g_i \circ g_{i+1} \circ \dots \circ g_k(s), \quad s \in [0, 1].$$

Consequently, by Corollary 2, we have  $q_i^{(k)} = g_{i \rightarrow k}(0)$ ,  $\tilde{q}_i^{(k)} = g_{i \rightarrow k}(1)$ ,  $q_i = \lim_{k \rightarrow \infty} g_{i \rightarrow k}(0)$ , and  $\tilde{q}_i = \lim_{k \rightarrow \infty} g_{i \rightarrow k}(1)$ .

The next two lemmas provide respectively an explicit and an implicit relation between the sequence of progeny generating functions  $\{g_k(\cdot)\}$  and the progeny generating vector  $\mathbf{G}(\cdot)$ . The first requires the following technical assumption.

ASSUMPTION 1. For all  $k \geq 0$ ,

$$(3.6) \quad \mathbf{P}_k\left(\lim_{n \rightarrow \infty} \sum_{i=0}^k Z_{n,i}^{(k)} \rightarrow 0\right) + \mathbf{P}_k\left(\lim_{n \rightarrow \infty} \sum_{i=0}^k Z_{n,i}^{(k)} \rightarrow \infty\right) = 1.$$

LEMMA 2. If Assumption 1 holds, then for all  $k \geq \varphi_0$ , the progeny generating function of  $\{Y_k\}$  at generation  $k$  is given by

$$g_k(s) = \lim_{n \rightarrow \infty} G_k^{(k,n)}(s_0, s_1, \dots, s_k, s), \quad s \in [0, 1],$$

where  $(s_0, s_1, \dots, s_k) \in [0, 1]^{k+1}$ .

PROOF. By (3.1) and (3.5),

$$(3.7) \quad g_k(s) = \mathbf{E}_k\left(s^{\lim_{n \rightarrow \infty} Z_{n,k+1}^{(k)}} \mathbb{1}_{\left\{\lim_{n \rightarrow \infty} Z_{n,k+1}^{(k)} < \infty\right\}}\right).$$

By Assumption 1 and the fact that  $(s_0, \dots, s_k) \in [0, 1)^{k+1}$ ,

$$\mathbf{P}_k \left( \lim_{n \rightarrow \infty} \prod_{i=0}^k s_i^{Z_{n,i}^{(k)}} = 0 \right) + \mathbf{P}_k \left( \lim_{n \rightarrow \infty} \prod_{i=0}^k s_i^{Z_{n,i}^{(k)}} = 1 \right) = 1,$$

that is,  $\lim_{n \rightarrow \infty} \prod_{i=0}^k s_i^{Z_{n,i}^{(k)}}$  is an indicator function. In addition, Lemma 1 implies  $\{\lim_{n \rightarrow \infty} Z_{n,k+1}^{(k)} < \infty\} \stackrel{\text{a.s.}}{=} \{\lim_{n \rightarrow \infty} \tilde{\mathbf{Z}}_n^{(k)} = \mathbf{0}\} = \{\lim_{n \rightarrow \infty} \prod_{i=0}^k s_i^{\tilde{Z}_{n,i}^{(k)}} = 1\} = \{\lim_{n \rightarrow \infty} \prod_{i=0}^k s_i^{Z_{n,i}^{(k)}} = 1\}$ . Thus, (3.7) can be rewritten as

$$g_k(s) = \mathbf{E}_k \left( \lim_{n \rightarrow \infty} s^{Z_{n,k+1}^{(k)}} \prod_{i=0}^k s_i^{Z_{n,i}^{(k)}} \right) = \lim_{n \rightarrow \infty} G_k^{(k,n)}(s_0, s_1, \dots, s_k, s),$$

where the last equality follows from the dominated convergence theorem.  $\square$

LEMMA 3. For any  $k \geq 0$ , the progeny generating function  $g_k(\cdot)$  satisfies

$$(3.8) \quad g_k(s) = G_k(g_{0 \rightarrow k}(s), g_{1 \rightarrow k}(s), \dots, g_k(s), s).$$

PROOF. By conditioning on the offspring of a type- $k$  individual in  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ ,

$$\begin{aligned} g_k(s) &= \mathbf{E} \left[ s^{\sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)}} \mathbb{1} \left\{ \sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)} < \infty \right\} \mid \varphi_0 = k \right] \\ &= \sum_{z \geq 0} \mathbf{E} \left[ s^{\sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)}} \mathbb{1} \left\{ \sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)} < \infty \right\} \mid \varphi_0 = k, \tilde{\mathbf{Z}}_1^{(k)} = z \right] \\ &\quad \times \mathbf{P}[\tilde{\mathbf{Z}}_1^{(k)} = z \mid \varphi_0 = k]. \end{aligned}$$

Then, by the Markov property and the independence between the daughter processes of individuals from the same generation,

$$\begin{aligned} &\mathbf{E} \left[ s^{\sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)}} \mathbb{1} \left\{ \sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)} < \infty \right\} \mid \varphi_0 = k, \tilde{\mathbf{Z}}_1^{(k)} = (z_0, \dots, z_k, z_{k+1}) \right] \\ (3.9) \quad &= s^{z_{k+1}} \prod_{i=0}^k \mathbf{E} \left[ s^{\sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)}} \mathbb{1} \left\{ \sum_{n=1}^{\infty} \tilde{Z}_{n,k+1}^{(k)} < \infty \right\} \mid \varphi_0 = i \right]^{z_i} \\ &= s^{z_{k+1}} \prod_{i=0}^k g_{i \rightarrow k}(s)^{z_i}, \end{aligned}$$

where (3.9) follows from (3.2). This leads to

$$g_k(s) = \sum_{z \geq 0} \prod_{i=0}^k g_{i \rightarrow k}(s)^{z_i} s^{z_{k+1}} \mathbf{P}[\tilde{Z}_1^{(k)} = z \mid \varphi_0 = k],$$

which completes the proof.  $\square$

**4. Fixed points and extinction probabilities.** We now characterise the set  $S$  defined in (1.2). The main results in this section rely on the relation between  $S$  and the set

$$S^{[e]} = \{s \in [0, 1]^{\mathcal{X}} : s_k = g_k(s_{k+1}) \forall k \geq 0\},$$

which corresponds to the set of fixed points of the embedded GWPVE. Because each  $g_k(\cdot)$  is a monotone increasing function, like  $S$ , the set  $S^{[e]}$  is one-dimensional. In this section, we assume that Assumption 1 holds. For any vector  $s \in S$ , we write  $\bar{s}^{(k)} := (s_0, s_1, \dots, s_k)$  for the restriction of  $s$  to its first  $k + 1$  entries.

The next lemma establishes a relationship between  $S$  and  $S^{[e]}$ .

LEMMA 4.  $S = S^{[e]} \cup \{\mathbf{1}\}$ .

PROOF. Suppose  $s \in S$  and  $s \neq \mathbf{1}$ . For any  $k, n \geq 0$ ,  $\bar{s}^{(k+1)}$  satisfies  $\bar{s}^{(k+1)} = G^{(k,n)}(\bar{s}^{(k+1)})$ . Because  $\{Z_n\}$  is irreducible and  $s \neq \mathbf{1}$  we have  $s_i < 1$  for all  $i \in \mathcal{X}$  (see [32], Theorem 2). Thus, using Lemma 2,  $g_k(s_{k+1}) = \lim_{n \rightarrow \infty} G_k^{(k,n)}(\bar{s}^{(k+1)}) = s_k$  for all  $k \geq 0$ , leading to  $s \in S^{[e]}$ . Now suppose  $s \in S^{[e]}$ . Then, by Lemma 3, for all  $k \geq 0$ ,

$$s_k = g_k(s_{k+1}) = G_k(g_{0 \rightarrow k}(s_{k+1}), g_{1 \rightarrow k}(s_{k+1}), \dots, g_k(s_{k+1}), s_{k+1}) = G_k(s),$$

therefore,  $s \in S$ .  $\square$

We now characterise the one-dimensional projection sets  $S_i$  and identify which elements of  $S$  correspond to the global and partial extinction probability vectors.

THEOREM 4.1. *If  $S = \{\mathbf{1}\}$ , then  $q = \tilde{q} = \mathbf{1}$ ; otherwise,*

$$q = \min S \quad \text{and} \quad \tilde{q} = \sup S \setminus \{\mathbf{1}\}.$$

*In particular,  $S_i = [q_i, \tilde{q}_i] \cup 1$  for all  $i \geq 0$ .*

PROOF. We show that

$$(4.1) \quad q = \min S^{[e]} \quad \text{and} \quad \tilde{q} = \max S^{[e]},$$

and for any  $i \geq 0$ ,  $S_i^{[e]} = [q_i, \tilde{q}_i]$ , where

$$S_i^{[e]} = \{x \in [0, 1] : \exists s \in S^{[e]}, \text{ such that } s_i = x\}.$$




FIG. 3. A visual representation of possible sets  $S_i$  in the irreducible case.

These results follow from the fact that  $g_i(\cdot)$  and  $g_i^{-1}(\cdot)$  are monotone increasing functions, and therefore so are  $g_{i \rightarrow j}(\cdot)$  and  $g_{i \rightarrow j}^{-1}(\cdot) := g_{j-1}^{-1} \circ \dots \circ g_i^{-1}(\cdot)$  for  $j > i$ . Let  $s \in S^{[e]}$ , then for all  $0 \leq i < k$ ,

$$q_i^{(k-1)} = g_{i \rightarrow k-1}(0) \leq s_i = g_{i \rightarrow k-1}(s_k) \leq g_{i \rightarrow k-1}(1) = \tilde{q}_i^{(k-1)}.$$

Taking the limit as  $k \rightarrow \infty$  we obtain  $q_i \leq s_i \leq \tilde{q}_i$  for all  $i \geq 0$ , which shows (4.1). Now suppose  $q_i \leq s_i \leq \tilde{q}_i$ . For any  $j < i$ , define  $s_j := g_{j \rightarrow i-1}(s_i)$ ; then

$$q_j = g_{j \rightarrow i-1}(q_i) \leq s_j \leq g_{j \rightarrow i-1}(\tilde{q}_i) = \tilde{q}_j.$$

Similarly, for any  $j > i$ , define  $s_j := g_{i \rightarrow j-1}^{-1}(s_i)$ ; then

$$q_j = g_{i \rightarrow j-1}^{-1}(q_i) \leq s_j \leq g_{i \rightarrow j-1}^{-1}(\tilde{q}_i) = \tilde{q}_j.$$

This shows that for any  $i \geq 0$  and for any  $s_i \in [q_i, \tilde{q}_i]$ , it is possible to construct a vector  $s$  belonging to  $S^{[e]}$ .  $\square$

Theorem 4.1 implies that  $S$  contains one, two, or uncountably many elements. More specifically, it shows that  $\mathbf{q}$  is the minimal element of  $S$  which is the beginning of a continuum of elements whose supremum is  $\tilde{\mathbf{q}}$ , as illustrated in Figure 3.

REMARK 1. In the reducible case, there may be an additional countable number of fixed points  $s$  such that  $\tilde{\mathbf{q}} \leq s \leq \mathbf{1}$ . We refer to [9], Section 4.4, for the details.

With the goal of giving a probabilistic interpretation to the intermediate fixed points  $s \in S$  such that  $\mathbf{q} < s < \tilde{\mathbf{q}}$ , we now derive properties of the infinite-dimensional set  $S$ . We begin by deriving a sufficient condition for

$$(4.2) \quad \lim_{i \rightarrow \infty} s_i = 1 \quad \text{for all } s \in S \setminus \{\mathbf{q}\},$$

that is, for  $S$  to contain *at most* a single element (corresponding to  $\mathbf{q}$ ) whose entries do not converge to 1. In a more general setting, sufficient conditions for (4.2) can

be found in Moyal ([27], Lemmas 3.3 and 3.4), the most notable being ‘ $\inf q_i > 0$ ’. The same author also conjectures a more general condition:

$$\sup_i p_i^{(1)} < 1 \quad \text{where } p_i^{(1)} := \sum_{\mathbf{v}:|\mathbf{v}|=1} p_i(\mathbf{v}).$$

In the case of LHBP, we now provide a stronger result.

**THEOREM 4.2.** *If*

$$(4.3) \quad \sum_{i=0}^{\infty} (1 - p_i^{(1)}) = \infty,$$

*then (4.2) holds.*

The proof of Theorem 4.2 uses the following lemma which we state separately because, for LHBP, it generalises the conditions of [10], Theorem 1.

**LEMMA 5.** *If (4.3) holds, then  $\mathbf{P}(Y_k \rightarrow 0) + \mathbf{P}(Y_k \rightarrow \infty) = 1$ .*

**PROOF.** Following Lindvall [23], we have  $\mathbf{P}(Y_k \rightarrow 0) + \mathbf{P}(Y_k \rightarrow \infty) = 1$  if and only if

$$(4.4) \quad \sum_{k=0}^{\infty} (1 - g'_k(0)) = \infty.$$

Suppose (4.3) holds without (4.4), that is, assume (4.3) and

$$(4.5) \quad \sum_{k=0}^{\infty} (1 - g'_k(0)) < \infty.$$

In this case, there can be only finitely many  $k$  such that  $g'_k(0) = 0$ . Thus, (4.5) holds if and only if there exists  $\ell \geq 0$  such that

$$(4.6) \quad \prod_{k=\ell}^{\infty} g'_k(0) \equiv \mathbf{P}_{\ell}(Y_k = 1, \forall k \geq \ell) > 0.$$

In addition, because  $M_{i,i+1} > 0$  for all  $i \geq 0$ , in every generation of the embedded process (including any for which  $g'_k(0) = 0$ ) individuals have a positive chance of giving birth to at least one offspring. In combination with (4.6), this implies that there exists  $c > 0$  such that for any  $l \geq 0$ ,  $\mathbf{P}_l(Y_k \geq 1, \forall k \geq l) \geq c$ . Recall that each individual in  $\{Y_k\}$  corresponds to an individual in  $\{Z_n\}$ . If the corresponding individual in  $\{Z_n\}$  has no offspring then neither does the individual in  $\{Y_k\}$ , whereas if the corresponding individual in  $\{Z_n\}$  has two or more offspring then the individual in  $\{Y_k\}$  must have at least two offspring with probability greater than or equal to  $c^2$ . Thus, for all  $k \geq 0$ ,  $1 - g'_k(0) \geq c^2(1 - p_k^{(1)})$ , which implies

$\sum_{k=0}^{\infty} (1 - g'_k(0)) \geq c^2 \sum_{k=0}^{\infty} (1 - p_k^{(1)}) = \infty$ , contradicting (4.5). Therefore, if (4.3) holds, we must have (4.4).  $\square$

PROOF OF THEOREM 4.2. By Lemma 4, we may assume  $s \in S^{[e]}$ . Thus, for all  $k \geq 0$ ,

$$(4.7) \quad s_0 = g_{0 \rightarrow k-1}(s_k) = \mathbf{E}_0(s_k^{Y_k} \mathbb{1}\{Y_k < \infty\}) = q_0^{(k-1)} + \mathbf{E}_0(s_k^{Y_k} \mathbb{1}\{0 < Y_k < \infty\}).$$

Suppose  $\liminf_k s_k < 1$ . In this case, there exists an infinite sequence  $\{k_i\}_{i \geq 1}$  such that  $s_{k_i} < 1 - \varepsilon$  for all  $i \geq 1$  and some  $\varepsilon > 0$ . For each  $i \geq 1$  and  $K \geq 1$ ,

$$\mathbf{E}_0(s_{k_i}^{Y_{k_i}} \mathbb{1}\{0 < Y_{k_i} < \infty\}) \leq \mathbf{P}_0(0 < Y_{k_i} < K) + (1 - \varepsilon)^K.$$

By Lemma 5, for any  $K \geq 1$ , we have  $\mathbf{P}_0(0 < Y_{k_i} < K) \rightarrow 0$  as  $i \rightarrow \infty$ . Letting  $K$  be arbitrarily large, we obtain  $\liminf_k \mathbf{E}_0(s_k^{Y_k} \mathbb{1}\{0 < Y_k < \infty\}) = 0$ . Because  $q_0^{(k)} \rightarrow q_0$  as  $k \rightarrow \infty$ , from (4.7) we then obtain  $s_0 = q_0$ . The only element  $s \in S^{[e]}$  such that  $\liminf_k s_k < 1$  is therefore  $s = \mathbf{q}$ .  $\square$

Now that we have general sufficient conditions for  $1 - s_i \rightarrow 0$ , we investigate properties of this convergence. The next two theorems use the following lemma.

LEMMA 6. *If  $\{a_n\}_{n \geq 0}$  and  $\{b_n\}_{n \geq 0}$  are sequences of nonnegative real numbers such that  $a_n \in (0, 1)$  for all  $n \geq 0$ , and  $b_n \rightarrow \infty$ , then*

$$(4.8) \quad \limsup_n a_n^{b_n} = \exp\left\{-\liminf_n b_n(1 - a_n)\right\},$$

$$(4.9) \quad \liminf_n a_n^{b_n} = \exp\left\{-\limsup_n b_n(1 - a_n)\right\}.$$

PROOF. For any  $n \geq 0$ , we have

$$a_n^{b_n} = \left(1 - \frac{b_n(1 - a_n)}{b_n}\right)^{b_n}.$$

The result then follows from  $\lim_{n \rightarrow \infty} (1 - c/n)^n = e^{-c}$  for any  $c \in \mathbb{R}$ .  $\square$

The next result shows that if the entries of  $\mathbf{q}$  converge to 1, then they converge slower than those of any other  $s \in S \setminus \{\mathbf{q}\}$ , whereas the entries of  $\tilde{\mathbf{q}}$  converge to 1 faster than those of any other  $s \in S \setminus \{\tilde{\mathbf{q}}, \mathbf{1}\}$ .

THEOREM 4.3. *If (4.3) holds then, for any  $s \in S \setminus \{\mathbf{q}, \tilde{\mathbf{q}}, \mathbf{1}\}$ ,*

$$\lim_{k \rightarrow \infty} \frac{1 - q_i}{1 - s_i} = \infty \quad \text{and} \quad \lim_{k \rightarrow \infty} \frac{1 - \tilde{q}_i}{1 - s_i} = 0.$$

PROOF. Suppose  $s \in S \setminus \{q, \tilde{q}, \mathbf{1}\}$ . In that case, by Theorem 4.1,  $q < s < \tilde{q}$ . In addition, by Lemma 4, for all  $k \geq 0$ ,

$$\begin{aligned}
 s_0 &= \mathbf{E}_0(s_k^{Y_k} \mathbb{1}\{Y_k < \infty\}) \\
 (4.10) \quad &= q_0^{(k-1)} + \mathbf{E}_0(s_k^{Y_k} \mathbb{1}\{0 < Y_k < \infty\})
 \end{aligned}$$

$$(4.11) \quad = \tilde{q}_0^{(k-1)} + \mathbf{E}_0((s_k^{Y_k} - 1) \mathbb{1}\{0 < Y_k < \infty\}).$$

Without loss of generality, we assume that  $q < \tilde{q}$ , which by Corollary 2 is equivalent to  $\mathbf{P}_0(0 < Y_k < \infty, \forall k \geq 0, ) > 0$ . In this case, by (4.10) and the fact that  $q_0^{(k)} \rightarrow q_0$ , we have

$$s_0 = q_0 \iff \lim_{k \rightarrow \infty} \mathbf{E}_0(s_k^{Y_k} \mid 0 < Y_k < \infty) = 0.$$

Because  $s_k^{Y_k}$  is nonnegative and uniformly bounded by 1, we can write

$$s_0 = q_0 \iff \mathbf{P}_0(s_k^{Y_k} \rightarrow 0 \mid \forall k \geq 0, 0 < Y_k < \infty) = 1.$$

By Lemma 5, we may then apply Lemma 6 to obtain

$$s_0 = q_0 \iff \mathbf{P}_0(Y_k(1 - s_k) \rightarrow \infty \mid \forall k \geq 0, 0 < Y_k < \infty) = 1,$$

hence  $\lim_{k \rightarrow \infty} (1 - q_i)/(1 - s_i) = \infty$ . Using (4.11), a similar argument yields

$$s_0 = \tilde{q}_0 \iff \mathbf{P}_0(Y_k(1 - s_k) \rightarrow 0 \mid \forall k \geq 0, 0 < Y_k < \infty) = 1,$$

and  $\lim_{k \rightarrow \infty} (1 - \tilde{q}_i)/(1 - s_i) = 0$ .  $\square$

The next theorem demonstrates that the rate at which  $1 - s_i$  decays is closely linked to the asymptotic growth of  $\{Y_k\}$ . In this context, we define a *growth rate* to be a sequence of real numbers  $\{C_k\}_{k \geq 0}$  such that

$$\lim_{k \rightarrow \infty} \frac{Y_k}{C_k} = W(\{C_k\}) \text{ exists a.s.,}$$

where  $W(\{C_k\})$  is a nonnegative, potentially defective, random variable with  $\mathbf{P}(0 < W(\{C_k\}) < \infty) > 0$ . We let

$$g_{W(\{C_k\})}(z) = \mathbf{E}_0(z^{W(\{C_k\})} \mathbb{1}\{W(\{C_k\}) < \infty\}).$$

Growth rates of nondefective GWPVEs ( $\tilde{q} = \mathbf{1}$ ) have been studied by a number of authors. Although it is natural to assume that  $\{\mathbf{E}_0[Y_{k+1}]\}_{k \geq 0}$  is a growth rate, it may not always be the case. Sufficient conditions for  $\{\mathbf{E}_0[Y_{k+1}]\}$  to be a growth rate are given in [19], and conditions for it to be the only distinct growth rate are discussed in [12, 13, 21]. Examples of GWPVEs with multiple growth rates can be found in [14, 24].



**THEOREM 4.4.** *Suppose (4.3) holds. If  $s \in S \setminus \{1\}$  and there exists some growth rate  $\{C_k\}$  such that  $g_{W(\{C_k\})}(0) < s_0 < g_{W(\{C_k\})}(1)$ , then*

$$\lim_{k \rightarrow \infty} (1 - s_k)C_k = c \in (0, \infty),$$

where  $c$  is such that  $s_0 = g_{W(\{C_k\})}(e^{-c})$ .

**PROOF.** By the arguments in the proof of Lemma 4, any  $s \in S \setminus \{1\}$  is such that  $s_i < 1$  for all  $i \in \mathcal{X}$ , and  $s \in S^{[e]}$ . Therefore, for all  $k \geq 1$ ,

$$(4.12) \quad s_0 = \mathbf{E}_0(s_k^{Y_k} \mathbb{1}\{Y_k < \infty\}) = \mathbf{E}_0(s_k^{Y_k}),$$

which can be rewritten as

$$(4.13) \quad \begin{aligned} s_0 &= \mathbf{E}_0((s_k^{C_k})^{Y_k/C_k} \mathbb{1}\{W(\{C_k\}) = 0\}) \\ &+ \mathbf{E}_0((s_k^{C_k})^{Y_k/C_k} \mathbb{1}\{0 < W(\{C_k\}) < \infty\}) \\ &+ \mathbf{E}_0((s_k^{C_k})^{Y_k/C_k} \mathbb{1}\{W(\{C_k\}) = \infty\}). \end{aligned}$$

By assumption, we have

$$(4.14) \quad \begin{aligned} g_{W(\{C_k\})}(0) &= \mathbf{P}_0(W(\{C_k\}) = 0) < s_0 < g_{W(\{C_k\})}(1) \\ &= \mathbf{P}_0(W(\{C_k\}) < \infty). \end{aligned}$$

If  $\liminf_k (s_k)^{C_k} = 0$ , then taking  $\liminf_k$  in (4.13) gives  $s_0 \leq \mathbf{P}_0(W(\{C_k\}) = 0)$ , which contradicts (4.14). A similar argument applies to the limit superior, leading to

$$(4.15) \quad 0 < \liminf_k s_k^{C_k} \leq \limsup_k s_k^{C_k} < 1.$$

By (4.12), we then have

$$(4.16) \quad \begin{aligned} s_0 &= \limsup_{k \rightarrow \infty} \mathbf{E}_0((s_k^{C_k})^{Y_k/C_k}) \\ &= \mathbf{E}_0\left(\limsup_{k \rightarrow \infty} (s_k^{C_k})^{Y_k/C_k}\right) \end{aligned}$$

$$(4.17) \quad = \mathbf{E}_0\left(\left(\limsup_{k \rightarrow \infty} s_k^{C_k}\right)^{W(\{C_k\})}\right),$$

where (4.16) follows from the dominated convergence theorem, and (4.17) requires (4.15). If we repeat the same argument with  $\limsup$  replaced by  $\liminf$ , we finally obtain

$$s_0 = g_{W(\{C_k\})}\left(\limsup_k s_k^{C_k}\right) = g_{W(\{C_k\})}\left(\liminf_k s_k^{C_k}\right).$$

By (4.3) and Theorem 4.2, we have  $s_k \rightarrow 1$ , and thus through (4.15) we obtain  $C_k \rightarrow \infty$ . Lemma 6 then gives  $\lim_{k \rightarrow \infty} (s_k)^{C_k} = e^{-c}$ , where  $c = \lim_{k \rightarrow \infty} (1 - s_k)C_k$ . This means  $s_0 = \mathbf{E}_0(e^{-cW(\{C_k\})}) = g_{W(\{C_k\})}(e^{-c})$ .  $\square$

We conclude this section with a summary of our findings on the set  $S$ . The set  $S$  is made up of a continuum of elements whose minimum is  $\mathbf{q}$  and whose maximum is  $\tilde{\mathbf{q}}$ , with the additional fixed point  $\mathbf{1}$ . Under Condition (4.3), for any  $s \in S$ , with the possible exception of  $\mathbf{q}$ , we have  $1 - s_i \rightarrow 0$  as  $i \rightarrow \infty$ . The decay rates of  $1 - q_i$  and  $1 - \tilde{q}_i$  are unique, whereas the intermediate elements  $\mathbf{q} < s < \tilde{\mathbf{q}}$  may share one or several decay rates, which have a one-to-one correspondence with the growth rates of  $\{Y_k\}$ . Furthermore, these intermediate elements completely specify the generating functions  $g_{W(\{C_k\})}(\cdot)$  and thereby the distributions of  $W(\{C_k\})$ . This gives a physical meaning to the intermediate elements: in short, they describe the evolution of  $\{Y_k\}$  when there is partial extinction without global extinction. While this physical interpretation is in terms of the growth of  $\{Y_k\}$ , we expect that it is closely related to the growth of  $\{\mathbf{Z}_n\}$ .

**5. Extinction criteria.** While there exist several well-established partial extinction criteria, determining a global extinction criterion when  $\tilde{\mathbf{q}} = \mathbf{1}$  remains an open question. When  $\tilde{\mathbf{q}} = \mathbf{1}$ , the embedded GWPVE  $\{Y_k\}$  is nonexplosive, and we can directly apply known extinction criteria for GWPVEs. These criteria are generally expressed in terms of the first and second factorial moments

$$\mu_k := g'_k(1) \quad \text{and} \quad a_k := g''_k(1), \quad k \geq 0.$$

The next lemma provides recursive expressions for these moments in terms of those of the offspring distributions of  $\{\mathbf{Z}_n\}$ . We let

$$\begin{aligned}
 m_{i \rightarrow k} &:= \mathbf{E}_i[Y_{k+1}] = g'_{i \rightarrow k}(1) = \prod_{j=i}^k \mu_j, \\
 (5.1) \quad G'_{k,i}(s) &:= \left. \frac{\partial G_k(\mathbf{u})}{\partial u_i} \right|_{\mathbf{u}=s}, \quad G''_{k,ij}(s) := \left. \frac{\partial^2 G_k(\mathbf{u})}{\partial u_i \partial u_j} \right|_{\mathbf{u}=s}, \\
 A_{k,ij} &:= G''_{k,ij}(\mathbf{1}),
 \end{aligned}$$

and we take the convention that  $\prod_{i=k}^{k-1} \cdot = 1$  and  $g_{k+1 \rightarrow k}(s) = s$ .

LEMMA 7. *Suppose  $\tilde{\mathbf{q}} = \mathbf{1}$ , then*

$$(5.2) \quad \mu_0 = \frac{M_{0,1}}{1 - M_{0,0}} \quad \text{and} \quad a_0 = \frac{\mu_0^2 A_{0,00} + A_{0,11} + 2\mu_0 A_{0,01}}{1 - M_{0,0}},$$

and for  $k \geq 1$ ,

$$(5.3) \quad \mu_k = \frac{M_{k,k+1}}{1 - \sum_{i=0}^k M_{k,i} m_{i \rightarrow k-1}},$$

and

$$(5.4) \quad a_k = \frac{\sum_{i=0}^k M_{k,i} \sum_{j=i}^{k-1} a_j m_{i \rightarrow j-1} (\prod_{\ell=j+1}^k \mu_\ell^2) + \sum_{i=0}^{k+1} \sum_{j=0}^{k+1} m_{i \rightarrow k} m_{j \rightarrow k} A_{k,ij}}{1 - \sum_{i=0}^k M_{k,i} m_{i \rightarrow k-1}}.$$

PROOF. By Lemma 3, for any  $k \geq 0$ ,

$$(5.5) \quad \begin{aligned} g'_k(s) &= \frac{d}{ds} [G_k(g_{0 \rightarrow k}(s), \dots, g_{k+1 \rightarrow k}(s))] \\ &= \sum_{i=0}^{k+1} g'_{i \rightarrow k}(s) G'_{k,i}(g_{0 \rightarrow k}(s), \dots, g_{k+1 \rightarrow k}(s)), \end{aligned}$$

where  $g'_{i \rightarrow k}(s) = \prod_{j=i}^k g'_j(g_{j+1 \rightarrow k}(s))$ . The assumption  $\tilde{\mathbf{q}} = \mathbf{1}$  implies  $g_{i \rightarrow k}(1) = 1$  for all  $i, k$  and, therefore,  $\mu_k = g'_k(1) = \sum_{i=0}^k M_{k,i} m_{i \rightarrow k} + M_{k,k+1}$ , which leads to the expression for  $\mu_0$  and the recursive equation (5.3).

Next, by differentiating (5.5) with respect to  $s$ , we obtain

$$\begin{aligned} g''_k(s) &= \sum_{i=0}^{k+1} g''_{i \rightarrow k}(s) G'_{k,i}(g_{0 \rightarrow k}(s), \dots, g_{k+1 \rightarrow k}(s)) \\ &\quad + \sum_{i=0}^{k+1} g'_{i \rightarrow k}(s) \sum_{j=0}^{k+1} g'_{j \rightarrow k}(s) G''_{k,ij}(g_{0 \rightarrow k}(s), \dots, g_{k+1 \rightarrow k}(s)), \end{aligned}$$

where, for  $0 \leq i \leq k$ ,

$$g''_{i \rightarrow k}(s) = \sum_{j=i}^k \left( \prod_{\ell=i}^{j-1} g'_\ell(g_{\ell+1 \rightarrow k}(s)) \right) g''_j(g_{j+1 \rightarrow k}(s)) \left( \prod_{\ell=j+1}^k g'_\ell(g_{\ell+1 \rightarrow k}(s)) \right)^2.$$

This implies

$$a_k = g''_k(1) = \sum_{i=0}^k M_{k,i} \sum_{j=i}^k a_j m_{i \rightarrow j-1} \left( \prod_{\ell=j+1}^k \mu_\ell^2 \right) + \sum_{i=0}^{k+1} \sum_{j=0}^{k+1} m_{i \rightarrow k} m_{j \rightarrow k} A_{k,ij},$$

which gives

$$\begin{aligned} &a_k \left( 1 - \sum_{i=0}^k M_{k,i} m_{i \rightarrow k-1} \right) \\ &= \sum_{i=0}^k M_{k,i} \sum_{j=i}^{k-1} a_j m_{i \rightarrow j-1} \left( \prod_{\ell=j+1}^k \mu_\ell^2 \right) + \sum_{i=0}^{k+1} \sum_{j=0}^{k+1} m_{i \rightarrow k} m_{j \rightarrow k} A_{k,ij}, \end{aligned}$$

leading to the expression for  $a_0$  and the recursive equation (5.4).  $\square$

When  $\tilde{\mathbf{q}} = \mathbf{1}$  is not assumed, the recursive expressions (5.2)–(5.4) can still be used to compute two sequences, which may not correspond to the first and second factorial moments of the progeny distributions of  $\{Y_k\}$ , but which we shall even so denote by  $\{\mu_k\}$  and  $\{a_k\}$ . For these sequences to correspond to well-defined moments, their elements must be nonnegative and finite, that is, the denominator

common to (5.3) and (5.4) must be strictly greater than 0 for all  $k \geq 0$ . Thus, if we let

$$x_k := \sum_{i=0}^k M_{k,i} m_{i \rightarrow k-1},$$

we require

$$(5.6) \quad 0 \leq x_k < 1 \quad \text{for all } k \geq 0.$$

By giving a physical interpretation to  $x_k$ , we now show that, in the irreducible case, (5.6) holds if and only if  $\tilde{q} = \mathbf{1}$ . Note that, if there exists  $k$  such that  $x_k = 1$ , then  $\tilde{q} < \mathbf{1}$ , as justified in the proof of the lemma.

LEMMA 8. *If  $\{\mathbf{Z}_n\}$  is irreducible, then  $\tilde{q} = \mathbf{1}$  if and only if  $0 \leq x_k < 1$  for all  $k \geq 0$ .*

PROOF. For any  $k \geq 0$ , we embed a process  $\{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})\}$  in  $\{\tilde{\mathbf{Z}}_n^{(k)} : \varphi_0 = k\}$  by taking all type- $k$  individuals that appear in  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$  and defining the direct descendants of these individuals as the closest (in generation) type- $k$  descendants in  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ ; the process  $\{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})\}$  evolves as a single-type Galton–Watson process that becomes extinct if and only if type  $k$  becomes extinct in  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ . Because  $M_{i,i+1} > 0$  for all  $i \geq 0$ , the extinction of type  $k$  in  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$  is almost surely equivalent to the extinction of the whole process  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ . Hence, for any  $k \geq 0$ ,

$$\tilde{q}^{(k)} < \mathbf{1} \quad \text{if and only if} \quad m_{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})} > 1,$$

where  $m_{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})}$  is the mean number of offspring born to an individual in  $\{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})\}$ . The value of  $m_{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})}$  is obtained by taking the weighted sum of all first return paths to  $k$  in the mean progeny representation graph of  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ . By conditioning on the progeny of an individual of type  $k$  in  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ , the lower-Hessenberg structure then leads to

$$m_{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})} = M_{k,0}m_{0 \rightarrow k-1} + M_{k,1}m_{1 \rightarrow k-1} + \dots + M_{k,k} = x_k.$$

Thus, if  $0 < x_k < 1$  for all  $k \geq 0$  then  $\tilde{q}^{(k)} = \mathbf{1}$  for all  $k$  and, therefore, by Theorem A.1,  $\tilde{q} = \lim_{k \rightarrow \infty} \tilde{q}^{(k)} = \mathbf{1}$ . Similarly, if there exists  $k$  such that  $x_k > 1$ , then  $\tilde{q} \leq \tilde{q}^{(k)} < \mathbf{1}$ . Now suppose there exists  $k$  such that  $x_k = 1$ . Then by the irreducibility of  $\{\mathbf{Z}_n\}$  there exists  $k^* > k$  such that there is a first return path with strictly positive weight of the form  $k \rightarrow k + 1 \rightarrow \dots \rightarrow k^* \rightarrow \dots \rightarrow k$  in the mean progeny representation graph of  $\{\tilde{\mathbf{Z}}_n^{(k^*)}\}$ . This implies

$$m_{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k^*)})} > m_{E_n^{(k)}(\tilde{\mathbf{Z}}_n^{(k)})} = 1,$$

and hence  $\tilde{q} \leq \tilde{q}^{(k^*)} < \mathbf{1}$ .  $\square$

Combining Lemmas 7 and 8 with [21], Theorem 1, which to the authors' knowledge is the most general extinction criterion currently available for GWPVEs, we obtain for an irreducible LHBP.

**THEOREM 5.1.** *If  $\{\mu_k\}$  and  $\{a_k\}$  are given by (5.2)–(5.4), then*

$$(5.7) \quad 0 < \mu_k < \infty \quad \forall k \geq 0 \quad \Leftrightarrow \quad \tilde{\mathbf{q}} = \mathbf{1},$$

*and when  $\tilde{\mathbf{q}} = \mathbf{1}$ , if  $\sup_k a_k/\mu_k < \infty$  and  $\inf_k \sum_{\{v:v_{k+1} \geq 2\}} p_k(\mathbf{v}) > 0$ , then*

$$(5.8) \quad \sum_{k=0}^{\infty} \frac{1}{m_{0 \rightarrow k}} = \infty \quad \Leftrightarrow \quad \mathbf{q} = \mathbf{1},$$

*where  $m_{0 \rightarrow k}$  is defined in (5.1).*

**PROOF.** The global extinction criterion (5.8) follows from (i)  $\Leftrightarrow$  (iv) in [21], Theorem 1. Indeed, our assumptions imply Condition (A) of that theorem, as well as  $\inf_k a_k/\mu_k > 0$ .  $\square$

**REMARK 2.** Theorem 5.1 demonstrates that by computing the sequence  $\{\mu_k\}$  required for (5.8) we are implementing a partial extinction criterion. We note that it is more efficient to compute  $\{\mu_k\}$  through Lemma 7 than to evaluate the convergence norm of  $M$  as the limit of the sequence of spectral radii of the north-west truncations of the mean progeny matrix  $M$  (see [30], Theorem 6.8).

**REMARK 3.** If  $\liminf_k m_{0 \rightarrow k} = 0$  then, through the Markov inequality, we obtain  $\mathbf{q} = \mathbf{1}$ . Thus, in this case the conditions of Theorem 5.1 do not need to be verified.

When the conditions of Theorem 5.1 do not hold, one may still be able to apply [21], Theorem 1, directly. Condition (A) in that theorem holds under an assumption on the third factorial moments  $g_k'''(1)$  ([21], Condition (C)), which can also be shown to satisfy recursive equations. Alternatively, it may be possible to apply the next theorem, which corresponds to [1], Theorem 1 (see, e.g., the proof of Proposition 1).

**THEOREM 5.2.** *If  $\tilde{\mathbf{q}} = \mathbf{1}$  and  $A_{k,i,j} < \infty$  for all  $k, i, j \geq 0$ , then for any  $1 \leq i < k$ ,*

$$\begin{aligned} & 1 - \left[ \frac{1}{m_{i \rightarrow (k-1)}} + \frac{1}{2} \sum_{j=i}^{k-1} \frac{g_j''(0)}{\mu_j m_{i \rightarrow j}} \right]^{-1} \\ & \leq q_i^{(k)} \leq 1 - \left[ \frac{1}{m_{i \rightarrow (k-1)}} + \sum_{j=i}^{k-1} \frac{a_j}{\mu_j m_{i \rightarrow j}} \right]^{-1}. \end{aligned}$$

Roughly speaking, Theorem 5.1 states that the boundary between almost sure global extinction and potential global survival is the expected linear growth of  $\{Y_k\}$ , that is,  $\mathbf{E}_0(Y_k) = m_{0 \rightarrow k-1} = Ck$ , for some constant  $C > 0$ . It is, however, not immediately clear how to interpret this criteria in terms of the expected growth of the original LHBP  $\{Z_n\}$ . The next theorem develops a link between the expected growth of  $\{Y_k\}$  and the exponential growth rate of the mean total population size in  $\{Z_n\}$ ,

$$(5.9) \quad \xi(M) := \liminf_n \sqrt[n]{\mathbf{E}_i |Z_n|} = \liminf_n \sqrt[n]{(M^n \mathbf{1})_i},$$

which, when  $M$  is irreducible, is independent of  $i$ . We note that in an irreducible MGWBP with finitely many types  $\xi(M) = \nu(M)$ , whereas when there are infinitely many types it is possible that  $\nu(M) < \xi(M)$ .

**THEOREM 5.3.** *Assume  $\nu(M) \leq 1$ . If  $\xi(M) > 1$ , then*

$$(5.10) \quad \limsup_n \sqrt[n]{m_{0 \rightarrow n}} \geq \xi(M),$$

and if  $\xi(M) < 1$ , then

$$(5.11) \quad \liminf_n \sqrt[n]{m_{0 \rightarrow n}} \leq \limsup_n (\mathbf{E}_0 |Z_n|)^{1/n}.$$

**PROOF.** We have  $m_{0 \rightarrow (n-1)} = \sum_{k=0}^n \mathbf{E}_0(Z_{n,k})m_{k \rightarrow (n-1)}$ , where  $m_{n \rightarrow (n-1)} := 1$ , which gives

$$(5.12) \quad \mathbf{E}_0 |Z_n| \inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} \leq m_{0 \rightarrow (n-1)} \leq \mathbf{E}_0 |Z_n| \sup_{0 \leq k \leq n} m_{k \rightarrow (n-1)}.$$

Now suppose  $\xi(M) > 1$ . In order to prove (5.10) we need to show that

$$(5.13) \quad \nexists n_0 < \infty \text{ such that } \inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} < 1 \quad \forall n > n_0.$$

Indeed, if (5.13) holds, because  $m_{n \rightarrow (n-1)} := 1$  we have

$$\limsup_n \inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} = 1,$$

and thus by (5.12),

$$\begin{aligned} \limsup_n \sqrt[n]{m_{0 \rightarrow (n-1)}} &\geq \limsup_n \left( \mathbf{E}_0 |Z_n| \inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} \right)^{1/n} \\ &\geq \liminf_n (\mathbf{E}_0 |Z_n|)^{1/n} \limsup_n \left( \inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} \right)^{1/n} \\ &= \xi(M). \end{aligned}$$

To show (5.13), assume there exists  $n_0 := \sup\{n : \inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} = 1\} < \infty$ , and observe that for any  $n \geq 0$  the recursion

$$\inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} = \min \left\{ \left( \inf_{0 \leq k \leq n-1} m_{k \rightarrow (n-2)} \right) \mu_{n-1}, \mu_{n-1}, 1 \right\}$$

holds. This implies that for all  $n > n_0$ ,

$$\inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} = \left( \inf_{0 \leq k \leq n-1} m_{k \rightarrow (n-2)} \right) \mu_{n-1},$$

and  $\inf_{0 \leq k \leq n} m_{k \rightarrow (n-1)} = m_{n_0 \rightarrow (n-1)}$ , which gives

$$m_{0 \rightarrow n} \left( \inf_{0 \leq k \leq n} m_{k \rightarrow n} \right)^{-1} = m_{0 \rightarrow (n_0-1)} \quad \text{for all } n > n_0.$$

By equation (5.12), we then have  $\mathbf{E}_0|\mathbf{Z}_n| \leq m_{0 \rightarrow (n_0-1)}$ , for all  $n > n_0$ , which contradicts the fact that  $\xi(M) > 1$  and shows (5.13). When  $\xi(M) < 1$ , a similar argument can be used to obtain (5.11).  $\square$

By Theorem 5.3, if both  $\lim_n \sqrt[n]{m_{0 \rightarrow n}}$  and  $\lim_n (\mathbf{E}_0|\mathbf{Z}_n|)^{1/n}$  exist (which is the case in our illustrative examples), then by the *root test for convergence*,

$$\xi(M) > 1 \quad \Rightarrow \quad \sum_{j=0}^{\infty} \frac{1}{m_{0 \rightarrow j}} < \infty \quad \text{and} \quad \xi(M) < 1 \quad \Rightarrow \quad \sum_{j=0}^{\infty} \frac{1}{m_{0 \rightarrow j}} = \infty.$$

Thus, if  $\xi(M) \neq 1$  then in Theorem 5.1  $\sum_{j=0}^{\infty} 1/m_{0 \rightarrow j} = \infty$  may be replaced by  $\xi(M) < 1$ . One contribution of Theorem 5.1, which is motivated by the examples in [5], is to provide an extinction criterion applicable even when  $\xi(M) = 1$ , as we demonstrate in Example 2.

**6. Illustrative examples.** We now illustrate the results of the previous section through two examples. Example 1 demonstrates that the mean progeny matrix  $M$  is not sufficient to determine whether  $\mathbf{q} < \mathbf{1}$  or  $\mathbf{q} = \mathbf{1}$ . This fact was highlighted in [33], Example 4.4, however, in that example, the process behaves asymptotically as a GWPVE because  $\sum_{j \neq i+1} M_{i,j} \rightarrow 0$  as  $i \rightarrow \infty$ . In addition, the proof relies on an explicit expression of the progeny generating vector. Through Example 1, we provide a streamlined proof which applies to a significantly broader class of branching processes.

In Example 2, we apply Theorem 5.1 to a LHBP with  $\xi(M) = 1$ . This example also motivates Section 7 on strong and nonstrong local survival.

The proofs related to the examples are collected in Appendix B.

EXAMPLE 1. Consider a LHBP  $\{\mathbf{Z}_n\}$  with mean progeny matrix

$$(6.1) \quad M = \begin{bmatrix} b & c & 0 & 0 & 0 & \dots \\ a & b & c & 0 & 0 & \\ 0 & a & b & c & 0 & \\ 0 & 0 & a & b & c & \\ \vdots & & & \ddots & \ddots & \ddots \end{bmatrix},$$

and progeny generating vector  $\mathbf{G}(\cdot)$ . We assume that  $a, c > 0$  and that there exists a constant  $B < \infty$  such that

$$(6.2) \quad A_{k,ij} = \left. \frac{\partial^2 G_k(s)}{\partial s_i \partial s_j} \right|_{s=1} \leq B \quad \text{for all } k, i, j \geq 0.$$

Apart from these assumptions, we impose no other condition on  $\{\mathbf{Z}_n\}$ . We now consider a modification of  $\{\mathbf{Z}_n\}$ , which we denote by  $\{\mathbf{Z}_n^{(u)}\}$  for some parameter  $u \geq 1$ , whose progeny generating vector,  $\mathbf{G}^{(u)}(s)$ , is given by

$$(6.3) \quad \begin{aligned} G_i^{(u)}(s_{i-1}, s_i, s_{i+1}) &= \frac{1}{\lceil u^i \rceil} G_i(s_{i-1}, s_i, s_{i+1}^{\lceil u^i \rceil}) \\ &+ \left(1 - \frac{1}{\lceil u^i \rceil}\right) G_i(s_{i-1}, s_i, 1), \quad i \geq 0. \end{aligned}$$

This modification decreases the probability that a type- $i$  individual has any type- $(i + 1)$  offspring by a factor of  $1/\lceil u^i \rceil$ , but when the type- $i$  individual does have type- $(i + 1)$  offspring, their number is increased by a factor of  $\lceil u^i \rceil$ , which causes the mean progeny matrix to remain unchanged. Before providing results on the extinction of  $\{\mathbf{Z}_n^{(u)}\}$ , we require the following lemma on branching processes with the tridiagonal mean progeny matrix (6.1).

LEMMA 9. *Suppose  $\{\mathbf{Z}_n\}$  has a mean progeny matrix given by (6.1), then  $\tilde{\mathbf{q}} = \mathbf{1}$  if and only if*

$$(6.4) \quad b < 1 \quad \text{and} \quad (1 - b)^2 - 4ac \geq 0,$$

and when (6.4) holds

$$(6.5) \quad \mu_k \nearrow \mu := \frac{1 - b - \sqrt{(1 - b)^2 - 4ac}}{2a} \quad \text{as } k \rightarrow \infty.$$

Note that the partial extinction criterion (6.4) was given previously in [18] and is implied by [8], Theorem 1. We are now in a position to characterise the global extinction probability of  $\{\mathbf{Z}_n^{(u)}\}$ .

PROPOSITION 1. *Consider the branching processes  $\{\mathbf{Z}_n^{(u)}\}$  defined in Example 1, and suppose  $b < 1$  and  $(1 - b)^2 - 4ac > 0$ . If  $\mu < 1$  then  $\mathbf{q} = \mathbf{1}$ , whereas if  $\mu \geq 1$ , then*

$$u > \mu \quad \Rightarrow \quad \mathbf{q} = \mathbf{1} \quad \text{and} \quad u < \mu \quad \Rightarrow \quad \mathbf{q} < \mathbf{1},$$

where  $\mu$  is given in (6.5).

An important sub-case of Example 1 is  $u = 1$ , the set of unmodified branching processes. Note that this is the only case where the second moments of the offspring distributions are uniformly bounded. For this sub-class of processes, when combined with Lemma 5, Proposition 1 yields the following.






FIG. 4. The mean progeny representation graph corresponding to Example 2.

COROLLARY 3. If  $u = 1$  and (4.3) holds then  $\mathbf{q} = \mathbf{1}$  if and only if  $\mu \leq 1$ .

EXAMPLE 2. Let  $\{\mathbf{Z}_n\}$  have a mean progeny matrix  $M$  such that  $M_{0,1} = 1$ , and for  $i \geq 1$ ,

$$(6.6) \quad M_{i,i-1} = \gamma \frac{i+1}{i} \quad \text{and} \quad M_{i,i+1} = (1-\gamma) \frac{i+1}{i}, \quad 0 \leq \gamma \leq 1,$$

with all remaining entries being 0. The mean progeny representation graph corresponding to this process is illustrated in Figure 4. We assume that there exists  $B < \infty$  such that  $A_{k,i,j} \leq B$  for all  $i, j, k \geq 0$  and that  $\inf_k \sum_{v:v_{k+1} \geq 2} p_k(\mathbf{v}) > 0$ .

For this example, it is not difficult to show that  $\xi(M) = 1$  if and only if  $\nu(M) \leq 1$ , which is the case for a range of values of  $\gamma$ , as we shall see.

PROPOSITION 2. For the set of branching processes described in Example 2,  $\mathbf{q} = \mathbf{1}$  if and only if  $\gamma = 0$ .

Proposition 2 states that the process experiences almost sure global extinction if and only if type- $i$  individuals can only have type- $(i + 1)$  offspring, that is, if it coincides exactly with the embedded GWPVE.

We choose

$$G_k(\mathbf{s}) = \begin{cases} \frac{1}{4}s_1^4 + \frac{1}{4}, & k = 0, \\ \frac{k+1}{4k}(\gamma s_{k-1} + (1-\gamma)s_{k+1})^4 + \frac{3k-1}{4k}, & k \geq 1, \end{cases}$$

which satisfies (6.6), and in Figure 5 we plot  $q_0^{(8000)} \approx q_0$  and  $\tilde{q}_0^{(8000)} \approx \tilde{q}_0$  for  $\gamma \in [0, 1]$ . Although we proved that  $q_0 = 1$  when  $\gamma = 0$ , we observe that  $q_0^{(8000)} \approx 0.95$  for this value of  $\gamma$ . This is because, when  $\gamma = 0$ , Theorem 5.2 implies

$$q_0 - q_0^{(k)} \sim \left( \sum_{\ell=0}^k \frac{1}{\ell} \right)^{-1} \sim \log^{-1}(k),$$

so the convergence of  $q^{(k)}$  to  $\mathbf{q} = \mathbf{1}$  is slow. For GWPVEs with  $\mathbf{q} < \mathbf{1}$ , little attention has been paid to this convergence rate in the literature, so for this example not




FIG. 5. The extinction probabilities  $q_0^{(8000)}$  (solid) and  $\tilde{q}_0^{(8000)}$  (dashed) for  $\gamma \in [0, 1]$ .

much can be said when  $\gamma > 0$ . Using Lemmas 7 and 8, we numerically determine that  $\tilde{\mathbf{q}} = \mathbf{1}$  if and only if  $\gamma \leq \gamma^*$  where

$$(6.7) \quad \gamma^* = \max\{\gamma : 0 < \mu_k < \infty \forall k \geq 0\} \approx 0.1625.$$

Note that in this particular example a sufficient condition for  $\tilde{\mathbf{q}} = \mathbf{1}$  is the existence of some  $k$  such that  $\mu_k < \mu_{k-1}$  (see the proof of Proposition 2). Thus,  $\gamma^*$  can be evaluated particularly efficiently. Given  $q_0^{(8000)} \leq q_0 \leq \tilde{q}_0 \leq \tilde{q}^{(8000)}$ , by visual inspection, the curves of partial and global extinction seem to merge from some value of  $\gamma$ ; however, the cut-off is not clear and further analysis is required to pinpoint the precise value. We are also interested in understanding whether this value depends only on the mean progeny matrix or whether other offspring distributions lead to different values. We address these questions in the next section.

**7. Strong local survival.** Each irreducible infinite-type branching process falls into one of the four categories  $\mathbf{q} = \tilde{\mathbf{q}} = \mathbf{1}$ ,  $\mathbf{q} < \tilde{\mathbf{q}} = \mathbf{1}$ ,  $\mathbf{q} < \tilde{\mathbf{q}} < \mathbf{1}$  or  $\mathbf{q} = \tilde{\mathbf{q}} < \mathbf{1}$ . The results in the previous section deal with the classification of LHBP with  $\tilde{\mathbf{q}} = \mathbf{1}$ . In the present section, we build on these results to establish a method for determining whether LHBP with  $\tilde{\mathbf{q}} < \mathbf{1}$  experience strong local survival ( $\mathbf{q} = \tilde{\mathbf{q}} < \mathbf{1}$ ), or nonstrong local survival ( $\mathbf{q} < \tilde{\mathbf{q}} < \mathbf{1}$ ). Other attempts at distinguishing between these two cases can be found, for instance, in [6, 7] and [25].

For any  $k \geq 0$ , we partition  $M$  into four components,

$$M = \begin{bmatrix} \tilde{M}^{(k)} & \tilde{M}_{12} \\ \tilde{M}_{21} & {}^{(k)}\tilde{M} \end{bmatrix},$$

where  $\tilde{M}^{(k)}$  is of dimension  $(k + 1) \times (k + 1)$  and the other three submatrices are infinite. We then construct a LHBP branching processes on  $(\Omega, \mathcal{F}, \mathbf{P})$ , denoted as  $\{^{(k)}\tilde{\mathbf{Z}}_n\}$ , with mean progeny matrix  ${}^{(k)}\tilde{M}$ , and global and partial extinction probability vectors  ${}^{(k)}\mathbf{q}$  and  ${}^{(k)}\tilde{\mathbf{q}}$ , respectively. Sample paths of  $\{^{(k)}\tilde{\mathbf{Z}}_n\}$  are constructed

from those of  $\{\mathbf{Z}_n\}$  by immediately killing all offspring of type  $i \leq k$ , and relabelling the types so that type  $i \geq k + 1$  becomes  $i - k - 1$ . We now use  $\{^{(k)}\tilde{\mathbf{Z}}_n\}$  to derive a criterion for strong local survival. In the next theorem, we let  $\text{sp}(\cdot)$  denote the spectral radius.

**THEOREM 7.1.** *Assume that  $\tilde{\mathbf{q}} > \mathbf{0}$ , that there exists  $k \in \mathbb{N}$  such that*

$$(7.1) \quad \text{sp}(\tilde{M}^{(k)}) > 1 \quad \text{and} \quad \nu(^{(k)}\tilde{M}) \leq 1,$$

*and that  $\tilde{M}_{21}$  contains a finite number of strictly positive entries. Then there is strong local survival in  $\{\mathbf{Z}_n\}$  if and only if  $\{^{(k)}\tilde{\mathbf{Z}}_n\}$  becomes globally extinct, that is,*

$$(7.2) \quad \mathbf{q} = \tilde{\mathbf{q}} < \mathbf{1} \quad \text{if and only if} \quad ^{(k)}\mathbf{q} = ^{(k)}\tilde{\mathbf{q}} = \mathbf{1}.$$

**PROOF.** We use [7], Theorem 4.2, which we restate using our notation: let  $\{\mathbf{Z}_n^{(G)}\}$  and  $\{\mathbf{Z}_n^{(G^*)}\}$  be two branching processes on the countable type set  $\mathcal{X}$  with respective probability generating functions  $\mathbf{G}(\cdot)$  and  $\mathbf{G}^*(\cdot)$ , and global extinction probability vectors  $\mathbf{q}$  and  $\mathbf{q}^*$ . Let  $A \subseteq \mathcal{X}$  be a nonempty subset of types and denote by  $\mathbf{q}(A)$  and  $\mathbf{q}^*(A)$  the respective vectors of probability of local extinction in  $A$ . If  $\{\mathbf{Z}_n^{(G)}\}$  and  $\{\mathbf{Z}_n^{(G^*)}\}$  differ on  $A$  only, that is, if  $G_i(s) = G_i^*(s)$  for all  $i \in \mathcal{X} \setminus A$  and  $G_i(s) \neq G_i^*(s)$  for all  $i \in A$ , then

$$(7.3) \quad \mathbf{q} = \mathbf{q}(A) \quad \Leftrightarrow \quad \mathbf{q}^* = \mathbf{q}^*(A).$$

We apply this result with  $A = \{0, 1, \dots, k\}$ ,  $\{\mathbf{Z}_n^{(G)}\} = \{\mathbf{Z}_n\}$ , and  $\{\mathbf{Z}_n^{(G^*)}\}$  being such that  $G_i^*(s) = \mathbf{1}$  for all  $i \in A$ , that is, all types in  $A$  are sterile. We need to show that (7.3) is equivalent to (7.2).

We first observe that  $\mathbf{q}(A) = \tilde{\mathbf{q}}$  since, by (7.1), in  $\{\mathbf{Z}_n\}$ , types in  $\mathcal{X} \setminus A$  are only able to survive through the presence of types in  $A$ . Next, since types in  $A$  are sterile in  $\{\mathbf{Z}_n^{(G^*)}\}$ ,  $\mathbf{q}^* = \mathbf{q}^*(A)$  if and only if  $q_i^* = q_i^*(A)$  for all  $i \geq k + 1$ . It is clear that  $(q_{k+1}^*, q_{k+2}^*, \dots) = ^{(k)}\mathbf{q}$  by construction. It remains to show that  $(q_{k+1}^*(A), q_{k+2}^*(A), \dots) = \mathbf{1}$ . We couple the process  $\{^{(k)}\tilde{\mathbf{Z}}_n : \varphi_0 = \ell - 1\}$  and the process  $\{\mathbf{Z}_n^{(G^*)} : \varphi_0 = k + \ell\}$ ,  $\ell \geq 1$ . Let  $k + \bar{\ell}$  be the largest type in  $\{\mathbf{Z}_n^{(G^*)}\}$  able to generate offspring in  $A$ . Then since  $\nu(^{(k)}\tilde{M}) \leq 1$ , with probability one there exists a generation  $N$  such that  $^{(k)}\tilde{Z}_{n,0} + \dots + ^{(k)}\tilde{Z}_{n,\bar{\ell}} = 0$  for all  $n \geq N$ . This implies that with probability one,  $Z_{n,k+1}^{(G^*)} + \dots + Z_{n,k+\bar{\ell}}^{(G^*)} = 0$  for all  $n \geq N$ , which shows  $(q_{k+1}^*(A), q_{k+2}^*(A), \dots) = \mathbf{1}$ .  $\square$

When  $\nu(^{(k)}\tilde{M}) \leq 1$ , we may apply Theorem 5.1 to determine whether  $^{(k)}\mathbf{q} = \mathbf{1}$ . We are now in a position to answer the questions posed at the end of the previous section. The next result is proved in Appendix B.

PROPOSITION 3. For the branching processes described in Example 2,

$$\begin{aligned}\gamma = 0 &\Rightarrow \mathbf{q} = \tilde{\mathbf{q}} = \mathbf{1}, \\ \gamma \in (0, \gamma^*] &\Rightarrow \mathbf{q} < \tilde{\mathbf{q}} = \mathbf{1}, \\ \gamma \in (\gamma^*, 1/2) &\Rightarrow \mathbf{q} < \tilde{\mathbf{q}} < \mathbf{1}, \\ \gamma \in (1/2, 1] &\Rightarrow \mathbf{q} = \tilde{\mathbf{q}} < \mathbf{1},\end{aligned}$$

where  $\gamma^*$  is given in (6.7).

Proposition 3 demonstrates that the curves for partial and global extinction represented in Figure 5 merge at  $\gamma = 1/2$  and that this value is independent of the particular offspring distributions. At the critical value  $\gamma = 1/2$ , there exists no  $k$  satisfying (7.1), causing this case to remain untreated.

**8. Conclusion.** Besides exploring the set of fixed points for LHBPs, we have introduced a method of classifying LHBPs into one of the categories  $\mathbf{q} = \tilde{\mathbf{q}} = \mathbf{1}$ ,  $\mathbf{q} < \tilde{\mathbf{q}} = \mathbf{1}$ ,  $\mathbf{q} < \tilde{\mathbf{q}} < \mathbf{1}$  or  $\mathbf{q} = \tilde{\mathbf{q}} < \mathbf{1}$ . Through Examples 1 and 2, we showed that our results can be used to rigorously determine which category the process falls in; however, in practical situations where rigorous proofs may not be possible, our results can still be applied computationally as a first step in classifying the process.

The inherent assumption in LHBPs is the constraint that individuals of type  $i$  cannot give birth to offspring whose type is larger than  $i + m$  for  $m = 1$ . The approach of embedding a GWPVE in the original LHBP can be extended to the case where  $m$  takes any finite integer value. The resulting embedded GWPVE then becomes multitype with  $m$  types. Results of Section 3 then naturally generalise, but those of Section 4 rely on the characterisation of the  $m$ -dimensional projection sets of  $S$ , which is more difficult in this case. The global extinction criterion discussed in Section 5 would now build upon extinction criteria for multi-type GWPVE, which are less developed in the literature. These questions are the topic of a subsequent paper [11].

#### APPENDIX A: PARTIAL EXTINCTION PROBABILITY

In this Appendix, we point out an error in Lemma 3.2 of [18] and reprove its assertion under additional assumptions (Theorem A.1). Our arguments hold not only for LHBPs but for more general MGWBPs with countably many types. Recall that a process is *singular* if and only if  $\sum_{\mathbf{v}:|\mathbf{v}|=1} p_i(\mathbf{v}) = 1$  for all  $i \in \mathcal{X}$ .

**THEOREM A.1.** *If  $\{\mathbf{Z}_n\}$  is irreducible and nonsingular then  $\tilde{\mathbf{q}}^{(k)} \searrow \tilde{\mathbf{q}}$  componentwise as  $k \rightarrow \infty$ .*

Before proving Theorem A.1, we briefly review Lemma 3.2 of [18]. This lemma states that  $\tilde{\mathbf{q}}^{(k)} \searrow \tilde{\mathbf{q}}$  without the additional assumptions that the process is irreducible and nonsingular. The key step in the proof of [18], Lemma 3.2, is the assertion that

$$(A.1) \quad \left\{ \omega \in \Omega : \lim_{n \rightarrow \infty} \tilde{\mathbf{Z}}_n^{(k)}(\omega) = \mathbf{0} \right\} \searrow \mathcal{E}_p \quad \text{as } k \rightarrow \infty,$$

where  $\mathcal{E}_p$  denotes the partial extinction event. If true, then by the monotone convergence theorem, we may conclude that  $\tilde{\mathbf{q}}^{(k)} \searrow \tilde{\mathbf{q}}$ . To understand why equation (A.1) is incorrect, consider an outcome  $\omega$  where a single infinite line of descent appears in the population<sup>3</sup> (for instance, the total population size is 1 in all generations) whose type  $j_n$  in successive generations  $n \geq 0$  follows the sequence 0, 0, 1, 0, 1, 2, 0, 1, 2, 3, 0, 1, 2, 3, 4, 0, . . . . For this outcome, the single infinite line of descent returns to type 0 infinitely often so  $\omega \notin \mathcal{E}_p$ , and its largest type is unbounded so  $\omega \in \{ \omega \in \Omega : \lim_{n \rightarrow \infty} \tilde{\mathbf{Z}}_n^{(k)}(\omega) = \mathbf{0} \}$  for all  $k \in \mathbb{N}$ . Thus

$$\lim_{k \rightarrow \infty} \left\{ \omega \in \Omega : \lim_{n \rightarrow \infty} \tilde{\mathbf{Z}}_n^{(k)}(\omega) = \mathbf{0} \right\} \neq \mathcal{E}_p.$$

Instead, the left-hand side of equation (A.1) converges down to

$$A := \left\{ \omega \in \Omega : \nexists (\varphi_0; i_1, j_1, y_1; \dots) \in X(\omega) \text{ such that } \limsup_n j_n < \infty \right\},$$

which is the set of outcomes such that no infinite line of descent with a finite maximum type appears in the population. If  $\omega \in \mathcal{E}_p$ , then  $\omega \in A$ , however, due to outcomes such as the one described above, the converse is not true, therefore,

$$\tilde{\mathbf{q}}_i^{(k)} \searrow \mathbf{P}_i(A) = \tilde{\mathbf{q}}_i + \mathbf{P}_i(A \setminus \mathcal{E}_p).$$

Thus  $\tilde{\mathbf{q}}^{(k)}$  converges to  $\tilde{\mathbf{q}}$  if and only if the measure of the (nonempty) set  $A \setminus \mathcal{E}_p$  is zero for all  $\mathbf{P}_i, i \in \mathcal{X}$ . One subclass of infinite-type processes where this does not hold is the class of singular branching processes that correspond to recurrent irreducible Markov chains with state space  $\mathcal{X}$ . Indeed, as these processes are recurrent,  $\mathbf{P}(\mathcal{E}_p) = 0$ . As they are also irreducible, with probability one, every state  $i \in \mathcal{X}$  is visited infinitely often; this means that with probability one, the single infinite line of descent which appears in the population does not have finite maximum type, leading to  $\mathbf{P}(A) = 1$ . We thus have  $\mathbf{P}(A \setminus \mathcal{E}_p) = 1$ .

Theorem A.1 implies that the *only* irreducible branching processes for which  $\mathbf{P}(A \setminus \mathcal{E}_p) > 0$  are those that correspond to Markov chains on  $\mathcal{X}$  (singular branching processes). Below we do not make use of the set  $A$  defined above, and instead prove Theorem A.1 via a different method.

---

<sup>3</sup>We say that the infinite line of descent  $x = (\varphi_0; i_1, j_1, y_1; i_2, j_2, y_2; \dots)$  appears in the population in outcome  $\omega$  (i.e.  $x \in X(\omega)$ ) if and only if  $(\varphi_0; i_1, j_1, y_1; \dots; i_n, j_n, y_n) \in X(\omega)$  for every  $n \in \mathbb{N}$ .

PROOF. Fix some initial type  $i \in \mathcal{X}$ . By construction, for every  $n \geq 0$  and  $\omega \in \Omega$ ,  $\tilde{\mathbf{Z}}_n^{(k)}(\omega)$  is increasing in  $k$ , which implies  $\tilde{q}_i^{(k)}$  is decreasing in  $k$ . Similarly, if  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$  survives globally, then at least one type  $j \in \{1, \dots, k\}$  must survive in  $\{\mathbf{Z}_n\}$ , which implies  $\tilde{q}_i^{(k)} \geq \tilde{q}_i$  for all  $k$ . We may then assume  $\tilde{q}_i < 1$ .

Because  $\{\mathbf{Z}_n\}$  is irreducible, [11], Corollary 1, implies that  $\tilde{q}_i$  is equal to the probability that type  $i$  eventually disappears from the population. We define a function  $f^{(i)} : \mathcal{J} \rightarrow \mathcal{J}$  that takes lines of descent  $(\varphi_0; i_1, j_1, y_1; \dots; i_n, j_n, y_n)$  and deletes each triple whose type  $j_{(\cdot)}$  is not equal to  $i$ , and we define the processes  $\{V_\ell^{(i)}\}_{\ell \geq 0}$  and  $\{\tilde{V}_\ell^{(i,k)}\}_{\ell \geq 0}$ , whose family trees are given by  $f^{(i)}(X)$  and  $f^{(i)}(\tilde{X}^{(k)})$ , respectively. These are single-type Galton–Watson processes that become extinct if and only if type  $i$  becomes extinct in  $\{\mathbf{Z}_n\}$  and  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$ . Thus, given  $\varphi_0 = i$  the probability that  $\{V_\ell^{(i)}\}$  becomes extinct is  $\tilde{q}_i$ , and the probability that  $\{\tilde{V}_\ell^{(i,k)}\}$  becomes extinct is greater than or equal to  $\tilde{q}_i^{(k)}$  (if  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$  is reducible there may be a positive chance type  $i$  dies out but  $\{\tilde{\mathbf{Z}}_n^{(k)}\}$  survives globally). Note that  $\{V_\ell^{(i)}\}_{\ell \geq 0}$  and  $\{\tilde{V}_\ell^{(i,k)}\}_{\ell \geq 0}$  are subject to explosion, however this makes no difference in the sequel.

Because  $\{\mathbf{Z}_n\}$  is irreducible and nonsingular,  $\{V_\ell^{(i)}\}$  is nonsingular, that is, there is positive chance that individuals in  $\{V_\ell^{(i)}\}$  have a total number of offspring different from 1. Thus, with probability 1,  $\{V_\ell^{(i)}\}$  experiences extinction or unbounded growth ([17], Chapter I, Theorem 6.2). For any  $K > 0$ , we then have

$$(A.2) \quad \lim_{\ell \rightarrow \infty} \mathbf{P}_i(V_\ell^{(i)} \geq K) = 1 - \tilde{q}_i.$$

Observe that, for any fixed  $h \in \mathbb{N}$  and  $K > 0$ ,

$$\begin{aligned} \{\omega \in \Omega : \tilde{V}_h^{(i,1)}(\omega) \geq K\} &\subseteq \{\omega \in \Omega : \tilde{V}_h^{(i,2)}(\omega) \geq K\} \\ &\subseteq \{\omega \in \Omega : \tilde{V}_h^{(i,3)}(\omega) \geq K\} \subseteq \dots \end{aligned}$$

and

$$(A.3) \quad \lim_{k \rightarrow \infty} \{\omega \in \Omega : \tilde{V}_h^{(i,k)}(\omega) \geq K\} = \{\omega \in \Omega : V_h^{(i)}(\omega) \geq K\}.$$

To understand (A.3) observe that if  $\omega \in \{\omega \in \Omega : V_h^{(i)}(\omega) \geq K\}$ , then there exists at least  $K$  lines of descent  $(\varphi_0; i_1, j_1, y_1; \dots; i_n, j_n, y_n) \in X(\omega)$  such that the type  $j_n = i$  is the  $h$ th return to  $i$  (where  $n$  is not necessarily the same for each of these  $K$  lines of descent). By construction, the maximum type on each of these lines of descent is finite. Thus letting  $k_m$  denote the maximum of the maximum type on  $K$  arbitrarily selected such lines of descent, we see that  $\omega \in \{\omega \in \Omega : \tilde{V}_h^{(i,k_m)}(\omega) \geq K\}$ . We may now apply the monotone convergence theorem to obtain, for any  $h \in \mathbb{N}$ ,

$$(A.4) \quad \lim_{k \rightarrow \infty} \mathbf{P}_i(\tilde{V}_h^{(i,k)} \geq K) = \mathbf{P}_i(V_h^{(i)} \geq K).$$

The probability that  $\{\tilde{V}_\ell^{(i,k)}\}_{\ell \geq 0}$  becomes extinct is equal to that of  $\{\tilde{V}_{h\ell}^{(i,k)}\}_{\ell \geq 0}$ , which is less than or equal the probability of extinction  $\tilde{q}_i^{(k,h,K)}$  of the Galton–Watson branching process with progeny generating function

$$G(s) = 1 - \mathbf{P}_i(\tilde{V}_h^{(i,k)} \geq K) + s^K \mathbf{P}_i(\tilde{V}_h^{(i,k)} \geq K).$$

Therefore,  $\tilde{q}_i^{(k)} \leq \tilde{q}_i^{(k,h,K)}$ . Observe that for any fixed  $p \in (0, 1]$  the probability of extinction in a Galton–Watson process with progeny generating function  $G(s) = 1 - p + ps^K$  converges monotonically to  $1 - p$  as  $K \rightarrow \infty$  (to see why note that for any  $0 < \eta < p$  there exists  $K$  large enough to ensure  $G(1 - p + \eta) = (1 - p) + p((1 - p) + \eta)^K \leq 1 - p + \eta$ ).

We are now in a position to show that for any  $\varepsilon > 0$  there exists  $k_i$  such that  $\tilde{q}_i^{(k_i)} < \tilde{q}_i + \varepsilon$ . Given a process with partial extinction probability  $\tilde{q}_i < 1$  and some  $0 < \varepsilon < 1 - \tilde{q}_i$ , we select  $K$  by setting it large enough to ensure that a Galton–Watson branching process with progeny generating function  $G(s) = (\tilde{q}_i + \varepsilon/2) + (1 - (\tilde{q}_i + \varepsilon/2))s^K$  has extinction probability less than  $\tilde{q}_i + \varepsilon$ . By (A.2), for this value of  $K$ , we may select  $h$  large enough to ensure  $|\mathbf{P}_i(V_h^{(i)} \geq K) - (1 - \tilde{q}_i)| < \varepsilon/4$ . By (A.4), for these values of  $K$  and  $h$  we may select  $k_i$  large enough to ensure  $|\mathbf{P}_i(V_h^{(i)} \geq K) - \mathbf{P}_i(\tilde{V}_h^{(i,k_i)} \geq K)| < \varepsilon/4$ . By the triangle inequality and the preceding discussion, for these values of  $k_i, h$  and  $K$ , we have  $\tilde{q}_i \leq \tilde{q}_i^{(k_i)} \leq \tilde{q}_i^{(k_i,h,K)} < \tilde{q}_i + \varepsilon$ . The result then follows from the fact  $\tilde{q}_i^{(k)}$  is decreasing in  $k$ .  $\square$

### APPENDIX B: PROOFS RELATED TO THE EXAMPLES

PROOF OF LEMMA 9. Because (6.4) holds, Lemma 7 gives

$$(B.1) \quad \mu_0 = \frac{c}{1-b} \quad \text{and} \quad \mu_k = \frac{c}{1-b-a\mu_{k-1}} \quad \text{for all } k \geq 0.$$

Because  $\mu_1 > \mu_0$  and

$$\mu_k - \mu_{k-1} = \frac{c}{1-b-a\mu_{k-1}} - \frac{c}{1-b-a\mu_{k-2}},$$

by induction the sequence  $\{\mu_k\}_{k \geq 0}$  is strictly positive and increasing. Therefore, since  $a > 0, \tilde{\mathbf{q}} = \mathbf{1}$  implies that  $\{\mu_k\}$  converges to a finite limit  $\mu$ , where  $\mu$  satisfies the equation  $ax^2 - (1-b)x + c = 0$ , which has real solutions

$$x_\pm = \frac{1-b \pm \sqrt{(1-b)^2 - 4ac}}{2a},$$

since (6.4) holds. When (6.4) holds, we have  $\mu_0 \leq x_-$  which, combined with (B.1) and the fact that  $x_- = c/(1-b-ax_-)$ , implies  $\mu_k \leq x_-$  for all  $k \geq 0$ , hence  $\mu_k \nearrow \mu = x_-$ .  $\square$

PROOF OF PROPOSITION 1. Let  $\Delta = (1 - b)^2 - 4ac > 0$ . First, suppose  $u > \mu$ . In this case, we have

$$1 - q_0^{(k)} = \frac{\mathbf{E}_0(Y_k)}{\mathbf{E}_0(Y_k | Y_k > 0)} \leq \frac{\mu^k}{u^{k-1}},$$

where  $\mathbf{E}_0(Y_k) \leq \mu^k$  follows from Lemma 9 and  $\mathbf{E}_0(Y_k | Y_k > 0) \geq u^{k-1}$  follows from the fact that the minimum number of type- $k$  offspring born to a type- $(k - 1)$  parent is  $\lceil u^{k-1} \rceil$ . This then implies

$$1 - q_0 = 1 - \lim_{k \rightarrow \infty} q_0^{(k)} \leq \lim_{k \rightarrow \infty} \frac{\mu^k}{u^{k-1}} = 0$$

and, therefore,  $\mathbf{q} = \mathbf{1}$  by irreducibility.

Now suppose  $1 \leq u < \mu$ . Note that  $A_{k,i,j}^{(u)} = A_{k,i,j}$  for all  $i, j$  with the exception of  $A_{k,(k+1)(k+1)}^{(u)} = \lceil u^k \rceil A_{k,(k+1)(k+1)} + c(\lceil u^k \rceil - 1)$ . Then, by Lemma 7,

$$\begin{aligned} a_k &= \frac{a\mu_k^2 a_{k-1} + \lceil u^k \rceil A_{k,(k+1)(k+1)} + O(1)}{1 - b - a\mu_{k-1}} \\ &\leq \frac{a\mu^2 a_{k-1} + B\lceil u^k \rceil + O(1)}{1 - b - a\mu} \\ &= a_{k-1}\mu \frac{a \frac{1-b-\Delta^{1/2}}{2a}}{1 - b - a \frac{1-b-\Delta^{1/2}}{2a}} + B^*u^k + O(1) \\ &= a_{k-1}\mu \frac{1 - b - \Delta^{1/2}}{1 - b + \Delta^{1/2}} + B^*u^k + O(1), \end{aligned}$$

for all  $k \geq 0$  and some  $B^* < \infty$ , which implies

$$a_k = O(\lceil \max\{u, \mu(1 - b - \Delta^{1/2})/(1 - b + \Delta^{1/2})\} \rceil^k).$$

By assumption,  $\Delta > 0$  and  $u < \mu$ , thus  $\max\{u, \mu \frac{1-b-\Delta^{1/2}}{1-b+\Delta^{1/2}}\} < \mu$ . Using the fact that  $\mu_k \nearrow \mu$  and the root test, we then obtain

$$\sum_{k=0}^{\infty} \frac{a_k}{\mu_k m_{0 \rightarrow k}} < \infty,$$

which, by the upper bound in Theorem 5.2, gives  $q_0 < 1$ .  $\square$

PROOF OF COROLLARY 3. It remains to show  $\mathbf{q} = \mathbf{1}$  when  $\mu = 1$ . Lemma 5 implies  $\mathbf{P}_0(Y_k \rightarrow 0) + \mathbf{P}_0(Y_k \rightarrow \infty) = 1$  and Lemma 9 implies  $\mathbf{E}_0(Y_k) = \prod_{i=0}^{k-1} \mu_i \leq 1$  for all  $k$ , leading to  $\mathbf{P}_0(Y_k \rightarrow \infty) = 0$  and the result.  $\square$



PROOF OF PROPOSITION 2. If  $\gamma = 0$ , then  $\mu_0 = M_{0,1} = 1$ , and for  $k \geq 1$ ,  $\mu_k = (k + 1)/k$ . This gives  $m_{0 \rightarrow k} = k + 1$  and, therefore,

$$\sum_{k=0}^{\infty} \frac{1}{m_{0 \rightarrow k}} = \sum_{k=1}^{\infty} \frac{1}{k} = \infty.$$

By assumption, when  $\gamma = 0$  the conditions Theorem 5.1 are satisfied, which then implies  $\mathbf{q} = \mathbf{1}$ .

Now suppose  $\gamma > 0$ . By Lemma 7, for  $k \geq 1$ ,

$$(B.2) \quad \mu_k = f_k(\mu_{k-1}) := \frac{\frac{k+1}{k}(1-\gamma)}{1 - \frac{k+1}{k}\gamma\mu_{k-1}}.$$

If there exists  $k$  such that  $\mu_k > 1/\gamma$ , then by Lemma 8 we have  $\mathbf{q} \leq \tilde{\mathbf{q}} < \mathbf{1}$ . Assume from now on that  $\tilde{\mathbf{q}} = \mathbf{1}$ , which implies that  $0 \leq \mu_k \leq 1/\gamma < \infty$  for all  $k$ , and  $\gamma < 1/2$ . Since  $\mu_0 = M_{0,1} = 1$ , using equation (B.2) we can inductively show that  $\mu_k \geq 1$  for all  $k \geq 0$ . We then have, for any  $k \geq 1$ ,

$$(B.3) \quad \mu_k \geq \frac{\frac{k+1}{k}(1-\gamma)}{1 - \frac{k+1}{k}\gamma} \geq 1 + \frac{1}{k(1-\gamma)}.$$

The Raabe–Duhamel test for convergence ensures that  $\sum_{k=0}^{\infty} (1/m_{0 \rightarrow k}) < \infty$ , since for  $k \geq 1$ ,

$$k \left( \frac{(1/m_{0 \rightarrow (k-1)})}{(1/m_{0 \rightarrow k})} - 1 \right) = k(\mu_k - 1) \geq \frac{1}{1-\gamma} > 1.$$

To complete the proof, it remains to show that the condition  $\sup_k a_k/\mu_k < \infty$  in Theorem 5.1 holds. By Lemma 7, for all  $k \geq 1$ ,

$$a_k = \frac{a_{k-1}\gamma\frac{k+1}{k}\mu_k^2}{1 - \gamma\frac{k+1}{k}\mu_{k-1}} + \frac{A_{k,(k-1)(k-1)}\mu_{k-1}^2\mu_k^2 + 2A_{k,(k-1)(k+1)}\mu_{k-1}\mu_k + A_{k,(k+1)(k+1)}}{1 - \gamma\frac{k+1}{k}\mu_{k-1}}.$$

Since  $\tilde{\mathbf{q}} = \mathbf{1}$ , the denominator is uniformly bounded away from 0; in addition, by assumption,  $A_{k,ij} \leq B < \infty$  for all  $i, j, k \geq 0$ , therefore, there exists some constant  $K < \infty$  independent of  $k$  such that

$$a_k \leq \frac{a_{k-1}\gamma\frac{k+1}{k}\mu_k^2}{1 - \frac{k+1}{k}\gamma\mu_{k-1}} + K.$$

If  $\mu_k \rightarrow 1$  (which we show below), then for large  $k$ ,

$$a_k \leq \left( \frac{\gamma}{1-\gamma} + o(1) \right) a_{k-1} + K.$$

Since  $\gamma < 1/2$ , we have  $\gamma/(1 - \gamma) < 1$ , which means that  $\{a_k\}$  is a uniformly bounded sequence. Combining this with the fact that  $\mu_k \geq 1$  for all  $k$  implies  $\sup a_k/\mu_k < \infty$ , and  $\mathbf{q} < \mathbf{1}$  by Theorem 5.1.

Finally, we prove that  $\mu_k \rightarrow 1$ . Observe that (B.2) implies that if  $\mu_k < \mu_{k-1}$  for some  $k$ , then  $\mu_{k+1} < \mu_k$ , and thus  $\mu = \lim_{k \rightarrow \infty} \mu_k$  exists since  $1 \leq \mu_k \leq 1/\gamma$  for all  $k$ . Taking  $k \rightarrow \infty$  in (B.2), we obtain that  $\mu$  satisfies

$$\mu = \frac{1 - \gamma}{1 - \gamma\mu} := f(\mu),$$

which means  $\mu$  is either 1 or  $(1 - \gamma)/\gamma > 1$ . The function  $f(x)$  is convex, thus  $f(x) > x$  for all  $x > (1 - \gamma)/\gamma$ ; in addition, by (B.2),  $\mu_{k+1} > f(\mu_k)$  for all  $k \geq 0$ . These imply that if  $\mu_k > (1 - \gamma)/\gamma$  for some  $k$ , then  $\mu_{k+\ell}$  becomes negative for some  $\ell > 1$ , which is a contradiction. So the sequence  $\{\mu_k\}$  lives in the open interval  $(1, (1 - \gamma)/\gamma)$ . Let

$$v_{\pm}^{(k)} = \frac{1}{2\gamma} \left[ \frac{k}{k+1} \pm \sqrt{\left(\frac{k}{k+1}\right)^2 - 4\gamma(1 - \gamma)} \right]$$

be the solutions of the equation  $x = f_k(x)$ . By the convexity of  $f_k(x)$  for all  $k$ , if there exists  $K \geq 1$  such that  $v_-^{(K+1)} < \mu_K < v_+^{(K+1)}$  then  $\{\mu_k\}_{k \geq K}$  is a decreasing sequence which converges to 1. Suppose  $\mu = (1 - \gamma)/\gamma$ . Then  $\mu_K \geq v_+^{(K+1)}$  for some  $K$ . We can then construct a LHBP,  $\{\mathbf{Z}_n^*\}$ , stochastically smaller than  $\{\mathbf{Z}_n\}$  by selecting a sufficiently large type  $K$  and independently killing each type- $(K + 1)$  child born to a type- $K$  parent with a probability carefully chosen to ensure  $v_-^{*(K+1)} < \mu_K^* < v_+^{*(K+1)}$ . For this modified process, we have  $\mu^* = 1$ , and repeating previous arguments, we obtain  $\mathbf{q} < \mathbf{q}^* < \mathbf{1}$ .  $\square$

**PROOF OF PROPOSITION 3.** Given Proposition 2 and Lemmas 7 and 8, it remains to show that  $\mathbf{q} < \tilde{\mathbf{q}}$  for  $\gamma \in (\gamma^*, 1/2)$  and  $\mathbf{q} = \tilde{\mathbf{q}}$  for  $\gamma \in (1/2, 1]$ . Note that, in either case, since  $\tilde{\mathbf{q}} < \mathbf{1}$ ,  $\exists K_1$  such that  $\text{sp}(\tilde{M}^{(k)}) > 1 \forall k \geq K_1$ . In addition,

$$\forall x > 1, \exists K(x) \text{ s.t. } M_{k,k+1} < x(1 - \gamma) \text{ and } M_{k,k-1} < x\gamma \quad \forall k \geq K(x).$$

Since  $\gamma \neq 1/2$ , we may choose  $\bar{x} > 1$  small enough so that  $1 - 4\bar{x}^2(1 - \gamma)\gamma > 0$ . By Lemma 9, this implies that  $v^{(k)}\tilde{M} < 1$  for all  $k \geq \bar{K} := K(\bar{x})$ , and

$${}^{(\bar{K})}\mu_k \leq \frac{1 - \sqrt{1 - 4\bar{x}^2(1 - \gamma)\gamma}}{2\bar{x}\gamma}$$

for all  $k \geq 0$ , where  $\{{}^{(\bar{K})}\mu_k\}_{k \geq 0}$  is computed using  ${}^{(\bar{K})}\tilde{M}$ .

Assume first that  $\gamma \in (1/2, 1]$ . Then  $(1 - \sqrt{1 - 4(1 - \gamma)\gamma})/(2\gamma) < 1$ , so we may choose  $x^* \leq \bar{x}$  small enough, corresponding to  $K^* := K(x^*) \geq \bar{K}$ , so that  ${}^{(K^*)}\mu_k < 1 - \varepsilon$  for all  $k \geq 0$  and some  $\varepsilon > 0$ . Hence there exists  $K = \max\{K_1, K^*\} < \infty$  satisfying the conditions of Theorem 7.1 with  ${}^{(K)}\mathbf{q} = {}^{(K)}\tilde{\mathbf{q}} = \mathbf{1}$ .

Now suppose  $\gamma \in (c, 1/2)$ . If for any  $K > \bar{K}$ , there exists  $k_1 \geq 0$  such that  ${}^{(K)}\mu_{k_1} \geq 1$ , then by the recursion (B.2) we have  ${}^{(K)}\mu_k \geq 1 + 1/(k(1 - \gamma))$ , for all  $k > k_1$ , and the result is derived by repeating the steps that follow equation (B.3) in the proof of Proposition 2. Suppose instead that there exists  $K > \bar{K}$  such that  ${}^{(K)}\mu_k < 1$  for all  $k \geq 0$ . Then by equation (B.2), we have  ${}^{(K)}\mu_{k-1} < 1 - 1/(\gamma(k + 1))$ , which implies

$$(B.4) \quad \prod_{i=0}^{\infty} {}^{(K)}\mu_i = 0.$$

To show that this leads to a contradiction, we compare  $M$  to a matrix  $M^*$  with strictly smaller entries than  $M$ :  $M^*$  is such that  $M_{0,1}^* = 1 - \gamma$ , and for all  $k \geq 1$ ,  $M_{k,k-1}^* = \gamma$  and  $M_{k,k+1}^* = 1 - \gamma$ , with all other entries 0. The value of  $\prod_{i=0}^{\infty} \mu_i^*$ , with  $\{\mu_i^*\}_{i \geq 0}$  computed using  $M^*$ , then has a probabilistic interpretation: it is the probability that a simple random walk on the integers, with transition probabilities  $p_+ = 1 - \gamma > p_- = \gamma$ , whose initial value is 0, never hits  $-1$ . When  $1 - \gamma > \gamma$ , it is well known that this value is nonzero. By the fact that  ${}^{(K)}M > M^*$ , we then have  ${}^{(K)}\mu_i > \mu_i^*$  for all  $i \geq 0$ , which implies  $\prod_{i=0}^{\infty} {}^{(K)}\mu_i > \prod_{i=0}^{\infty} \mu_i^* > 0$ , contradicting (B.4).  $\square$

**Acknowledgements.** The authors are grateful to two anonymous referees and an Associate Editor for their constructive comments which helped us to improve the manuscript. In particular, they thank the referee who pointed out the error in [18], Lemma 3.2.

## REFERENCES

- [1] AGRESTI, A. (1975). On the extinction times of varying and random environment branching processes. *J. Appl. Probab.* **12** 39–46. [MR0365733](#)
- [2] ATHREYA, K. B. and NEY, P. E. (1972). *Branching Processes. Die Grundlehren der Mathematischen Wissenschaften* **196**. Springer, New York. [MR0373040](#)
- [3] BARBOUR, A. D. (1994). Threshold phenomena in epidemic theory. In *Probability, Statistics and Optimisation. Wiley Ser. Probab. Math. Statist. Probab. Math. Statist.* 101–116. Wiley, Chichester. [MR1320745](#)
- [4] BARBOUR, A. D. and KAFETZAKI, M. (1993). A host-parasite model yielding heterogeneous parasite loads. *J. Math. Biol.* **31** 157–176. [MR1205055](#)
- [5] BERTACCHI, D. and ZUCCA, F. (2009). Characterization of critical values of branching random walks on weighted graphs through infinite-type branching processes. *J. Stat. Phys.* **134** 53–65. [MR2489494](#)
- [6] BERTACCHI, D. and ZUCCA, F. (2014). Strong local survival of branching random walks is not monotone. *Adv. in Appl. Probab.* **46** 400–421. [MR3215539](#)
- [7] BERTACCHI, D. and ZUCCA, F. (2017). A generating function approach to branching random walks. *Braz. J. Probab. Stat.* **31** 229–253. [MR3635904](#)
- [8] BIGGINS, J. D., LUBACHEVSKY, B. D., SHWARTZ, A. and WEISS, A. (1991). A branching random walk with a barrier. *Ann. Appl. Probab.* **1** 573–581. [MR1129775](#)
- [9] BRAUNSTEINS, P. (2018). Extinction in branching processes with countably many types. Ph.D. dissertation.

- [10] BRAUNSTEINS, P., DECROUEZ, G. and HAUTPHENNE, S. (2019). A pathwise approach to the extinction of branching processes with countably many types. *Stochastic Process. Appl.* **129** 713–739. [MR3913265](#)
- [11] BRAUNSTEINS, P. and HAUTPHENNE, S. (2018). The probabilities of extinction in a branching random walk on a strip. Preprint. Available at [arXiv:1805.07634](#).
- [12] D’SOUZA, J. C. (1994). The rates of growth of the Galton–Watson process in varying environments. *Adv. in Appl. Probab.* **26** 698–714. [MR1285455](#)
- [13] D’SOUZA, J. C. and BIGGINS, J. D. (1992). The supercritical Galton–Watson process in varying environments. *Stochastic Process. Appl.* **42** 39–47. [MR1172506](#)
- [14] FOSTER, J. H. and GOETTGE, R. T. (1976). The rates of growth of the Galton–Watson process in varying environment. *J. Appl. Probab.* **13** 144–147. [MR0394910](#)
- [15] GANTERT, N., MÜLLER, S., POPOV, S. and VACHKOVSKAIA, M. (2010). Survival of branching random walks in random environment. *J. Theoret. Probab.* **23** 1002–1014. [MR2735734](#)
- [16] HACCOU, P., JAGERS, P. and VATUTIN, V. A. (2007). *Branching Processes: Variation, Growth, and Extinction of Populations. Cambridge Studies in Adaptive Dynamics* **5**. Cambridge Univ. Press, Cambridge. [MR2429372](#)
- [17] HARRIS, T. E. (1963). *The Theory of Branching Processes. Die Grundlehren der Mathematischen Wissenschaften* **119**. Springer, Berlin. [MR0163361](#)
- [18] HAUTPHENNE, S., LATOUCHE, G. and NGUYEN, G. (2013). Extinction probabilities of branching processes with countably infinitely many types. *Adv. in Appl. Probab.* **45** 1068–1082. [MR3161297](#)
- [19] JAGERS, P. (1974). Galton–Watson processes in varying environments. *J. Appl. Probab.* **11** 174–178. [MR0368197](#)
- [20] JAGERS, P. (1975). *Branching Processes with Biological Applications. Wiley Series in Probability and Mathematical Statistics—Applied Probability and Statistics*. Wiley Interscience, London. [MR0488341](#)
- [21] KERSTING, G. (2017). A unifying approach to branching processes in varying environments. Preprint. Available at [arXiv:1703.01960](#).
- [22] KIMMEL, M. and AXELROD, D. E. (2002). *Branching Processes in Biology. Interdisciplinary Applied Mathematics* **19**. Springer, New York. [MR1903571](#)
- [23] LINDVALL, T. (1974). Almost sure convergence of branching processes in varying and random environments. *Ann. Probab.* **2** 344–346. [MR0378130](#)
- [24] MACPHEE, I. M. and SCHUH, H.-J. (1983). A Galton–Watson branching process in varying environments with essentially constant offspring means and two rates of growth. *Aust. J. Stat.* **25** 329–338. [MR0725212](#)
- [25] MENSHIKOV, M. V. and VOLKOV, S. E. (1997). Branching Markov chains: Qualitative characteristics. *Markov Process. Related Fields* **3** 225–241. [MR1468175](#)
- [26] MODE, C. J. (1971). *Multitype Branching Processes*. Elsevier, New York.
- [27] MOYAL, J. E. (1962). Multiplicative population chains. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **266** 518–526. [MR0138132](#)
- [28] SAGITOV, S. and LINDO, A. (2016). A special family of Galton–Watson processes with explosions. In *Branching Processes and Their Applications. Lect. Notes Stat.* **219** 237–254. Springer, Cham. [MR3587990](#)
- [29] SAGITOV, S. and MINUESA, C. (2017). Defective Galton–Watson processes. *Stoch. Models* **33** 451–472. [MR3765109](#)
- [30] SENETA, E. (2006). *Non-negative Matrices and Markov Chains. Springer Series in Statistics*. Springer, New York. [MR2209438](#)
- [31] SERRA, M. C. and HACCOU, P. (2007). Dynamics of escape mutants. *Theor. Popul. Biol.* **72** 167–178.

- [32] SPĂȚARU, A. (1989). Properties of branching processes with denumerably many types. *Rev. Roumaine Math. Pures Appl.* **34** 747–759. [MR1031234](#)
- [33] ZUCCA, F. (2011). Survival, extinction and approximation of discrete-time branching random walks. *J. Stat. Phys.* **142** 726–753. [MR2773785](#)

SCHOOL OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF MELBOURNE  
PARKVILLE, 3010, MELBOURNE  
AUSTRALIA  
E-MAIL: [braunsteins.p@unimelb.edu.au](mailto:braunsteins.p@unimelb.edu.au)  
[sophiemh@unimelb.edu.au](mailto:sophiemh@unimelb.edu.au)