

# QUENCHED INVARIANCE PRINCIPLES FOR THE MAXIMAL PARTICLE IN BRANCHING RANDOM WALK IN RANDOM ENVIRONMENT AND THE PARABOLIC ANDERSON MODEL

BY JIŘÍ ČERNÝ<sup>1</sup> AND ALEXANDER DREWITZ<sup>2</sup>

<sup>1</sup>*Department of Mathematics and Computer Science, Universität Basel, [jiri.cerny@unibas.ch](mailto:jiri.cerny@unibas.ch)*

<sup>2</sup>*Mathematisches Institut, Universität zu Köln, [drewitz@math.uni-koeln.de](mailto:drewitz@math.uni-koeln.de)*

We consider branching random walk in spatial random branching environment (BRWRE) in dimension one, as well as related differential equations: the Fisher–KPP equation with random branching and its linearized version, the parabolic Anderson model (PAM). When the random environment is bounded, we show that after recentering and scaling, the position of the maximal particle of the BRWRE, the front of the solution of the PAM, as well as the front of the solution of the randomized Fisher–KPP equation fulfill quenched invariance principles. In addition, we prove that at time  $t$  the distance between the median of the maximal particle of the BRWRE and the front of the solution of the PAM is in  $O(\ln t)$ . This partially transfers classical results of Bramson (*Comm. Pure Appl. Math.* **31** (1978) 531–581) to the setting of BRWRE.

## CONTENTS

1. Introduction . . . . .	95
2. Definition of the model and main results . . . . .	96
2.1. Behavior of the maximally displaced particle . . . . .	98
2.2. Implications for the PAM and randomized Fisher–KPP equation . . . . .	99
3. Strategy of the proof . . . . .	102
4. Expected number of particles of given velocity . . . . .	104
4.1. Tilted random walk measures . . . . .	104
4.2. An invariance principle for the empirical Legendre transforms . . . . .	108
4.3. An auxiliary invariance principle . . . . .	111
4.4. The walk lingers in the bulk . . . . .	113
4.5. Initial condition stability . . . . .	114
4.6. Proof of Theorem 2.8 (functional CLT for the PAM) . . . . .	115
5. Breakpoint behavior . . . . .	116
5.1. Perturbation estimates . . . . .	116
5.2. Proof of Theorem 2.6 (functional CLT for the breakpoint) . . . . .	119
5.3. Invariance principle for the breakpoint inverse . . . . .	120
6. The breakpoint approximates the maximum . . . . .	122
6.1. Leading particles . . . . .	122
6.1.1. First moment for the leading particles . . . . .	123
6.1.2. Second moment for the leading particles . . . . .	131
6.2. Proof of Theorem 2.4 and Proposition 2.3 . . . . .	135
7. BRWRE and the randomized Fisher–KPP . . . . .	137
8. Open questions . . . . .	138
Appendix: Auxiliary results . . . . .	138
A.1. Properties of logarithmic moment generating functions . . . . .	138
A.2. Basic properties of the Lyapunov exponent . . . . .	140
A.3. Hoeffding-type inequality for mixing sequences . . . . .	143

Received January 2018; revised January 2019.

*MSC2010 subject classifications.* 60J80, 60G70, 82B44.

*Key words and phrases.* Branching random walk, random environment, parabolic Anderson model, invariance principles.

Acknowledgments . . . . . 144  
 References . . . . . 144

**1. Introduction.** Branching random walk as well as branching Brownian motion, and in particular the position of their maximally displaced particles, have been the subject of highly intensive research during the last couple of decades; see the monographs [8, 63] as well as the references in these sources.

Indeed, in [32, 42] and [7] it has successively been shown that under suitable assumptions the position of the maximal or rightmost particle  $M(n)$  of the branching random walk at time  $n$  satisfies a law of large numbers; that is, almost surely

$$(1.1) \quad \lim_{n \rightarrow \infty} n^{-1}M(n) = \tilde{v}_0$$

for some nonrandom  $\tilde{v}_0 \in \mathbb{R}$ . Subsequently, concentration results for  $M(n)$  around its median  $m(n)$  (cf. [15, 20, 53, 61]), as well as corresponding results on the distributional convergence have been obtained; see [2, 4, 12, 13]. In particular, in [1, 35] the law of large numbers of (1.1) has been improved in that for a wide class of branching random walks the position of the maximal particle  $M(n)$  at time  $n$  satisfies

$$(1.2) \quad M(n) = \tilde{v}_0 n - \frac{3}{2}c \ln n + O(1),$$

where  $c > 0$  is a parameter depending on the specifics of the branching and displacement mechanisms.

In the continuum setting of branching Brownian motion (BBM) with binary branching, replacing  $n$  by  $t$  in a suggestive way for the respective quantities, even more precise asymptotics, namely

$$(1.3) \quad m(t) = \sqrt{2}t - \frac{3}{2\sqrt{2}} \ln t + o(1),$$

has been proved much earlier in seminal works by Bramson [11, 14] already. Bramson made use of the fact that the function  $w^{\text{BBM}}(t, x) := \mathbb{P}(M(t) \geq x)$ ,  $t \geq 0$ ,  $x \in \mathbb{R}$ , solves the Fisher–KPP equation

$$(1.4) \quad \frac{\partial w^{\text{BBM}}}{\partial t} = \frac{1}{2} \Delta w^{\text{BBM}} + w^{\text{BBM}}(1 - w^{\text{BBM}}),$$

with the initial condition  $w^{\text{BBM}}(0, \cdot) = \mathbf{1}_{(-\infty, 0]}$ . He then investigated the solution to this equation through an impressively refined analysis of its Feynman–Kac representation.

While the above results for branching random walk have been derived in the context of homogeneous branching mechanisms, there has recently been an increased activity in the investigation of branching random walk with nonhomogeneous branching rates that depend on either time or space in special *deterministic* ways; see [5, 9, 10, 21, 22, 45, 46, 49, 50, 59]. Among other things, as an interesting consequence of the inhomogeneous branching rates, in these sources second-order terms that differ from the logarithmic correction of [14] and (1.2) have been obtained.

While the above sources focus on the case of deterministic branching environments, there are very compelling reasons for trying to achieve a better understanding of the case of spatially random branching environments. On the one hand, this is already interesting from a purely mathematical point of view. On the other hand, when it comes to modeling real world applications, though branching environments are not random, they oftentimes are locally irregular but exhibit certain spatial averaging properties. One natural approach is then to model

the environment as random and try to understand the evolution of the process either conditionally on a realization of the branching environment or averaged over all such environments. In this setting, notable research has been conducted over the past decades on a variety of aspects such as survival and growth properties, transience versus recurrence, diffusivity, as well as localization properties (see, e.g., [16, 17, 24, 29, 34, 36, 48, 56, 60] for a nonexhaustive list).

To the best of our knowledge, the only source that in some sense focuses on the maximal particle is Comets and Popov [17]. They prove a shape theorem for a BRWRE on  $\mathbb{Z}^d$ ,  $d \geq 1$ , from which, as a corollary, one can infer that the maximal particle has an asymptotic velocity, that is, (1.1) holds.

Finally, branching random walk in an environment that is changing randomly in *time* was studied in [37, 52] recently. Among other results, Huang and Liu [37] proved a law of large numbers for the maximal particle. Mallein and Miłoś [52] considered the backlog of the maximal particle behind what can be interpreted as the breakpoint in their setting (cf. (2.5) below) and proved that it is strictly larger than in the setting of constant branching rates. As a corollary, their displaced particle. It should be noted here that the time-dependent random environment seems to be easier to handle since certain techniques of the theory of multitype branching processes apply in this case. We were not able to use them for the model considered in this paper.

**2. Definition of the model and main results.** Let us now introduce the model of branching random walk in random (branching) environment considered in this paper. The random environment is given by a family  $\xi = (\xi(x))_{x \in \mathbb{Z}}$  of random variables defined on some probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . We assume that the environment is i.i.d. and bounded:

$$\begin{aligned} \text{(POT)} \quad & (\xi(x))_{x \in \mathbb{Z}} \text{ are i.i.d. under } \mathbb{P}, \\ & 0 < \text{ei} := \text{ess inf } \xi(0) < \text{ess sup } \xi(0) =: \text{es} < \infty. \end{aligned}$$

We presume that the i.i.d. property can be relaxed with some additional technical effort, but we prefer to work in this context for the sake of simplicity. The same holds true for the condition  $\text{ei} > 0$  which could be relaxed to  $\text{ei} \geq c \in \mathbb{R}$ , with negative branching rates being interpreted as killing rates. On the other hand, some form of boundedness of  $\xi(x)$  from above is essential for our investigations.

We furthermore assume, again for reasons of simplicity, that the initial configuration  $u_0 : \mathbb{Z} \rightarrow \mathbb{N}_0$  is such that

$$\text{(INI)} \quad C \mathbf{1}_{-\mathbb{N}_0} \geq u_0 \geq \mathbf{1}_{\{0\}} \quad \text{for some } C \in [1, \infty).$$

In particular,  $u_0 = \mathbf{1}_{\{0\}}$  and  $u_0 = \mathbf{1}_{-\mathbb{N}_0}$  fulfill (INI). Later, as a consequence of Lemmas 4.15 and 5.1 below, we show that any initial configuration satisfying (INI) is comparable for our purposes to  $u_0 = \mathbf{1}_{\{0\}}$  in the results that follow. Hence, the reader may assume  $u_0 = \mathbf{1}_{\{0\}}$  from now onward without loss of generality.

Let us now describe the dynamics of the BRWRE in detail. Given a realization of  $\xi$  and an initial condition  $u_0 : \mathbb{Z} \rightarrow \mathbb{N}_0$ , at each  $x \in \mathbb{Z}$  we place  $u_0(x)$  particles at time 0. As time evolves, all particles move independently according to continuous time simple random walk with jump rate 1. In addition, and independently of everything else, while at a site  $x$ , a particle splits into two at rate  $\xi(x)$ , and if it does so, the two new particles evolve independently according to the same diffusion mechanism as the remaining particles. This defines *branching random walk in the branching environment*  $\xi$  with binary branching, where again the latter is for simplicity but not essential. Given a realization of  $\xi$ , we write  $\mathbb{P}_{u_0}^\xi$  for the quenched law of the process conditional on starting with a particle configuration  $u_0$  at time 0, and  $\mathbb{E}_{u_0}^\xi$  for

the corresponding expectation. We use  $\mathbb{P} \times \mathbb{P}_{u_0}^\xi$  to denote the averaged law of the process. To simplify notation, we abbreviate  $\mathbb{P}_x^\xi = \mathbb{P}_{\{x\}}^\xi$ .

We use  $N(t)$  to denote the set of particles alive at time  $t$  in this BRWRE. For any particle  $Y \in N(t)$ , we denote by  $(Y_s)_{s \in [0,t]}$  the trajectory of itself and its ancestors up to time  $t$ . We will also call  $(Y_s)_{s \in [0,t]}$  the *genealogy* of  $Y$ . For  $t \geq 0$  and  $x \in \mathbb{Z}$ , we define

$$(2.1) \quad \begin{aligned} N(t, x) &:= |\{Y \in N(t) : Y_t = x\}| \quad \text{and} \\ N^{\geq}(t, x) &:= |\{Y \in N(t) : Y_t \geq x\}| = \sum_{y \geq x} N(t, y) \end{aligned}$$

as the number of particles in the process at time  $t$  which are located at or to the right of  $x$ .

To state our last assumption, we recall that it is well known from the studies on the parabolic Anderson model (cf. Section 2.2 below) that there is a deterministic function  $\lambda : \mathbb{R} \rightarrow \mathbb{R}$ , the *Lyapunov exponent*, such that for a.e. realization of  $\xi$  the quenched expectation of  $N(t, x)$  satisfies

$$(2.2) \quad \lambda(v) = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \mathbb{E}_0^\xi [N(t, \lfloor tv \rfloor)], \quad v \in \mathbb{R}.$$

Under (POT), one can show that  $\lambda$  is even, concave everywhere and strictly concave exactly on  $(v_c, \infty)$  for some nontrivial critical velocity  $v_c \in (0, \infty)$ ; see Figure 1 for the illustration and Proposition A.3 in the Appendix for the proof. Furthermore, the asymptotic velocity of the maximally displaced particle (cf. (1.1)) is given by the unique  $v_0 \in (0, \infty)$  such that

$$(2.3) \quad \lambda(v_0) = 0.$$

Throughout the paper, we will assume that the maximally displaced particle is faster than  $v_c$ , that is,

$$(VEL) \quad v_0 > v_c.$$

This assumption will ensure that there is certain tilted Gibbs measure related to BRWRE (cf. (4.4) and below) under which the particles have the speed  $v_0$ ; the existence of such a measure is crucial for the techniques employed in this paper. While condition (VEL) is not easy to check in general, in Lemma A.4 of the Appendix we show that it is satisfied for a rich family of random environments. Moreover, so far we have found no examples where (VEL) fails to hold, but a proof that (VEL) is always fulfilled eludes us so far. Figure 1 covers possible shapes of the Lyapunov exponent in terms of convexity and the locations of  $v_0$  and  $v_c$ .




FIG. 1. Qualitative illustration of the behavior of the Lyapunov exponent  $\lambda(v)$  with (VEL) satisfied (left) or not (right). In particular, the Lyapunov exponent is a linear function on two nondegenerate symmetric intervals adjacent to the origin, and strictly convex otherwise; see Proposition A.3 for details.

2.1. *Behavior of the maximally displaced particle.* From a probabilistic point of view, in this article we are mainly interested in the behavior of the position of the maximally displaced particle at time  $t$ ,

$$M(t) := \max\{Y_t : Y \in N(t)\},$$

for which we prove the following functional central limit theorem.

**THEOREM 2.1.** *Assume (POT), (INI) and (VEL). Then there is  $\bar{\sigma}_{v_0} \in (0, \infty)$  given explicitly in (5.20) below, such that the sequence of processes*

$$[0, \infty) \ni t \mapsto \frac{M(nt) - v_0 nt}{\bar{\sigma}_{v_0} \sqrt{n}}, \quad n \in \mathbb{N},$$

converges as  $n \rightarrow \infty$  in  $\mathbb{P} \times \mathbb{P}_{u_0}^{\xi}$ -distribution to standard Brownian motion.

**REMARK 2.2.** Without further mentioning, in the functional central limit theorems we prove, we consider the space of càdlàg functions endowed with the Skorokhod topology as the underlying space.

Theorem 2.1 will directly follow from three intermediate results (Proposition 2.3 and Theorems 2.4, 2.6 below) which are of independent interest. To state these results, we define  $m(t)$  as the quenched median of the distribution of  $M(t)$ ,

$$(2.4) \quad m(t) := \sup\{x \in \mathbb{Z} : \mathbb{P}_{u_0}^{\xi}(M(t) \geq x) \geq 1/2\}.$$

Note here that  $m(t)$  is a random variable on  $(\Omega, \mathcal{F}, \mathbb{P})$ .

We further introduce a quantity  $\bar{m}(t)$  which is sometimes referred to as the *breakpoint* in the case of homogeneous branching rates; incidentally, we already remark at this point that in our setting it is also instructive to interpret it as the front of the solution to the parabolic Anderson model; cf. Section 2.2 below. It is defined as

$$(2.5) \quad \bar{m}(t) := \sup\{x \in \mathbb{Z} : \mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, x)] \geq 1/2\}.$$

As the first ingredient of the proof of Theorem 2.1, we show that  $M(t)$  is sufficiently close to its median so that, for the sake of the functional central limit theorem,  $M(t)$  can effectively be replaced by  $m(t)$ .

**PROPOSITION 2.3.** *Under assumptions (POT), (INI) and (VEL), there is a constant  $C \in (0, \infty)$  such that  $\mathbb{P} \times \mathbb{P}_{u_0}^{\xi}$ -a.s.,*

$$\limsup_{t \rightarrow \infty} \frac{|M(t) - m(t)|}{\ln t} \leq C.$$

The second substantial step to show Theorem 2.1 is the following approximation result. It is one of the main results of this article and it is interesting in its own right.

**THEOREM 2.4.** *Assume (POT), (INI) and (VEL) to hold. Then  $m(t) \leq \bar{m}(t)$ , and there exists a constant  $C \in (0, \infty)$  such that for  $\mathbb{P}$ -a.e. realization of  $\xi$ ,*

$$(2.6) \quad \limsup_{t \rightarrow \infty} \frac{\bar{m}(t) - m(t)}{\ln t} \leq C.$$

REMARK 2.5. This result should be compared to the classical results of Bramson [11, 14] for homogeneous BBM (and to corresponding results for branching random walk [1, 35]). In the case of BBM, the breakpoint satisfies  $\bar{m}(t) = \sqrt{2}t - \frac{1}{2\sqrt{2}} \ln t + o(1)$  which can be proved easily using Gaussian tail estimates. Together with (1.3), this yields that for BBM,

$$\lim_{t \rightarrow \infty} \frac{\bar{m}(t) - m(t)}{\ln t} = \frac{1}{\sqrt{2}}.$$

Our result thus shows that in the case of random branching rates we can recover an upper bound whose order matches that of the homogeneous branching setting. The question of whether there is a limit in (2.6) remains open.

The third and last ingredient of the proof of Theorem 2.1 is the functional central limit theorem for a suitably rescaled and centered version of the process  $\bar{m}(t)$ . In fact, we prove a slightly more general statement: As a generalization to (2.5), we define

$$(2.7) \quad \bar{m}_v(t) := \sup \left\{ x \in \mathbb{N} : \mathbb{E}_{u_0}^\xi [N^{\geq}(t, x)] \geq \frac{1}{2} e^{t\lambda(v)} \right\}, \quad v > 0, t > 0,$$

where  $\lambda$  is the Lyapunov exponent defined in (2.2). Note that due to the definition (2.3) of  $v_0$  we have  $\bar{m}(t) = \bar{m}_{v_0}(t)$ .

THEOREM 2.6. *Under assumptions (POT) and (INI), for every  $v > v_c$ , the sequence of processes*

$$[0, \infty) \ni t \mapsto \frac{\bar{m}_v(nt) - vnt}{\bar{\sigma}_v \sqrt{n}}, \quad n \in \mathbb{N},$$

*converges as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to standard Brownian motion. The value of  $\bar{\sigma}_v \in (0, \infty)$  is given in (5.20) below.*

Theorem 2.1 follows directly from Proposition 2.3 and Theorems 2.4, 2.6. Combining these results, we also immediately obtain a functional limit theorem for the median  $m(t)$ .

COROLLARY 2.7. *Assuming (POT), (INI) and (VEL), with  $\bar{\sigma}_{v_0}$  as in Theorem 2.6, the sequence of processes*

$$[0, \infty) \ni t \mapsto \frac{m(nt) - v_0nt}{\bar{\sigma}_{v_0} \sqrt{n}}, \quad n \in \mathbb{N},$$

*converges as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to standard Brownian motion.*

2.2. *Implications for the PAM and randomized Fisher–KPP equation.* As we have touched upon previously in the Introduction, there is a close connection between certain partial differential equations and branching processes: In the case of BBM, it is easy to see that the density  $u^{\text{BBM}}(t, x)$  of the expected number of particles satisfies

$$(2.8) \quad \frac{\partial}{\partial t} u^{\text{BBM}} = \frac{1}{2} \Delta u^{\text{BBM}} + u^{\text{BBM}}.$$

As this equation is essentially the heat equation (write  $u^{\text{BBM}} = e^t \tilde{u}$ ), this allows to estimate the corresponding breakpoint  $\sup\{x > 0 : u(t, x) \geq 1/2\}$  with high accuracy using Gaussian tail estimates. Moreover, as we already mentioned,  $w^{\text{BBM}}(t, x) = \mathbb{P}(M(t) \geq x)$  satisfies the Fisher–KPP equation (1.4). In particular, the front of the solution to (1.4), defined as  $\sup\{x \in \mathbb{R} : w^{\text{BBM}}(t, x) \geq 1/2\}$ , coincides with the median  $m(t)$  of the distribution of the maximal

particle of the BBM. Therefore, Bramson's result (1.3) immediately gives equally precise information on the position of the front of the solution to (1.4) as well.

In our setting of inhomogeneous branching rates, the situation is both more complicated but also more interesting. The breakpoint in the case of heterogeneous branching rates corresponds to the front of the solution to the *parabolic Anderson model* (PAM), a discrete randomized version of (2.8),

$$(2.9) \quad \begin{aligned} \frac{\partial u}{\partial t}(t, x) &= \Delta_d u(t, x) + \xi(x)u(t, x), \quad t \geq 0, x \in \mathbb{Z}, \\ u(0, x) &= u_0(x), \quad x \in \mathbb{Z}. \end{aligned}$$

Here,  $\Delta_d f(x) = \frac{1}{2}(f(x+1) + f(x-1) - 2f(x))$  stands for the discrete Laplace operator.

It is well known that conditionally on  $\xi$ , the expected number of particles at time  $t$  and position  $x$

$$(2.10) \quad u(t, x) := \mathbb{E}_{u_0}^\xi [N(t, x)]$$

solves (2.9) (cf. the original source [27] as well as [26] and [44] for more recent surveys). Hence, due to (2.5) and (2.10), the process  $\bar{m}(t)$  can be viewed as the *front* of the solution to the PAM, which according to Theorem 2.6, fulfills a corresponding functional central limit theorem.

This functional central limit theorem can be supplied with another one, for the logarithm of the function  $u(t, x)$  itself: Since statement (2.2) can be read as a law of large numbers for  $t^{-1} \ln u(t, \lfloor tv \rfloor)$ , it is natural to inquire about the fluctuations. Our investigations lead to a corresponding invariance principle which is of independent interest.

**THEOREM 2.8.** *Under assumptions (POT) and (INI), for every  $v > v_c$  there exists  $\sigma_v \in (0, \infty)$  given explicitly in (4.25) below, such that the sequence of processes*

$$[0, \infty) \ni t \mapsto \frac{1}{\sigma_v \sqrt{vn}} (\ln u(nt, \lfloor vnt \rfloor) - nt\lambda(v)), \quad n \in \mathbb{N},$$

*converges as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to standard Brownian motion.*

While this result for the front of the solution to the PAM is interesting in its own right, the question naturally arises of what one can say about the front of the solution to its nonlinear version, the randomized discrete Fisher–KPP equation

$$(2.11) \quad \frac{\partial w}{\partial t}(t, x) = \Delta_d w(t, x) + \xi(x)w(t, x)(1 - w(t, x)), \quad t \geq 0, x \in \mathbb{Z},$$

Previous results (in continuum space) on the front of the solution to (2.11) have been obtained in [25] (see also [23]), [58] and [31]. First, under suitable regularity and mixing assumptions, and a Heaviside-type initial condition,  $w(0, \cdot) = \mathbf{1}_{-\mathbb{N}_0}$ , as in (1.4), the existence of the speed of the front

$$(2.12) \quad \hat{m}(t) := \sup\{x \in \mathbb{R} : w(t, x) = 1/2\}$$

of the solution to the randomized Fisher–KPP equation (2.11) is known [23], Theorem 7.6.1: For  $\mathbb{P}$ -a.e. realization of  $\xi$ ,

$$(2.13) \quad \lim_{t \rightarrow \infty} t^{-1} \hat{m}(t) = \hat{v}_0,$$

where  $\hat{v}_0$  is nonrandom and corresponds to the speed of the front of the linearized equation, which is a “continuum space PAM.” Here, as in Bramson's work [11], a precise analysis of the Feynman–Kac formula plays an important role in the proofs.



In the case of  $\xi$  periodic instead of random, in [31] it has been shown that there is a logarithmic correction term between  $m(t)$  and  $\widehat{m}(t)$ , and the authors were able to characterize the constant in front of the logarithmic correction as a certain minimizer.

To the best of our knowledge, nothing is known about the fluctuations of  $\widehat{m}(t)$  for the Heaviside-type initial conditions in the case of random branching rates. For a different and due to technical reasons restricted set of initial conditions, Nolen [58] has derived a central limit theorem for the position of the front of the solution to (2.11) by analytic means. To put our results into context, let us describe the assumptions of [58] more precisely: The initial condition  $w_0(x, \xi)$  of [58] is required to depend on the randomness of the environment. It should satisfy  $\lim_{x \rightarrow -\infty} w_0(x, \xi) = 1$  (which roughly corresponds to our assumption (INI)), and more importantly,

$$(2.14) \quad c(\xi)\tilde{w}(x, \xi, \gamma) \leq w_0(x, \xi) \leq C(\xi)\tilde{w}(x, \xi, \gamma) \quad \text{for all } x > 0.$$

Here,  $\tilde{w} = \tilde{w}(x, \xi, \gamma)$ ,  $t \geq 0$ ,  $x \geq 0$ , is a nonnegative solution to the ordinary differential equation  $\frac{1}{2}\Delta\tilde{w} = (\xi - \gamma)\tilde{w}$  satisfying  $\tilde{w}(0, \xi, \gamma) = 1$  and which decays to 0 as  $x \rightarrow \infty$ . It was known previously that  $\tilde{w}$  exists whenever  $\gamma$  is larger than a certain  $\bar{\gamma}$ . In addition, there is another  $\gamma^* > \bar{\gamma}$  such that whenever  $\gamma \geq \gamma^*$  and the initial condition satisfies (2.14), then the law of large numbers for the velocity of the traveling wave, that is, (2.13) holds with the same speed  $\hat{v}_0$ . In order to prove his central limit theorem, Nolen needs to assume that  $\gamma \in (\bar{\gamma}, \gamma^*)$ , which leads to traveling waves with a larger velocity  $v(\gamma) > \hat{v}_0$ . The initial conditions corresponding to such  $\gamma$  decay to 0 exponentially as  $x \rightarrow \infty$ , but the rate of decay is slow.

It is worthwhile to remark that such a distinction between the waves with the minimal (or “critical”) velocity and the waves with strictly larger velocity is present already in the paper of Bramson [11]. Already there, it turns out that the “supercritical” is easier to handle.

One of our main motivations for writing this paper was to understand the behavior of the front of the traveling wave solution to randomized Fisher–KPP equation in the “critical” case, in particular for initial conditions of the form  $w_0 = \mathbf{1}_{-\mathbb{N}_0}$ , that are, from the point of view of the BRWRE as well as of the PAM, more natural.

**THEOREM 2.9.** *Let  $\widehat{m}(t)$  be the front of the solution to discrete randomized Fisher–KPP equation (2.11) with initial condition  $w_0 = \mathbf{1}_{-\mathbb{N}_0}$  defined similarly as in (2.12) by*

$$(2.15) \quad \widehat{m}(t) := \sup\{x \in \mathbb{Z} : w(t, x) \geq 1/2\}.$$

*Then, assuming that (POT) and (VEL) hold true,  $(\widehat{m}(t) - v_0t)/(\bar{\sigma}_{v_0}\sqrt{t})$  converges as  $t \rightarrow \infty$  in  $\mathbb{P}$ -distribution to a standard normal random variable.*

The previous theorem is a *nonfunctional* central limit theorem only, which might look surprising in view of our previous results. The reason for this is the fact that the connection between the BRWRE and the corresponding randomized Fisher–KPP equation is slightly more complicated than in the homogeneous case, due to the fact that the BRWRE is not translation and reflection invariant (given  $\xi$ ): We will prove in Proposition 7.1 that

$$w(t, x) = \mathbb{P}_x^\xi(M(t) \geq 0)$$

solves the randomized Fisher–KPP equation (2.11) with initial condition  $w(0, \cdot) = \mathbf{1}_{\mathbb{N}_0}$ . This should be contrasted with the definition of  $w^{\text{BBM}}(t, x) = \mathbb{P}(M(t) \geq x)$  used in (1.4).



**3. Strategy of the proof.** We now roughly explain the strategy of the proof of our main results, and describe the organization of the paper. As it is common in the branching random walk literature, a first moment method is used to provide an upper bound on the maximum of the BRWRE; a complementary truncated second moment computation gives a lower bound.

Luckily, similar to the homogeneous case, the moments of the number of particles in the BRWRE (possibly satisfying certain additional restrictions) have an explicit representation. This representation, in terms of expectations of certain functionals of simple random walk, is called the Feynman–Kac formula, “many-to-one lemma” or “many-to-few lemma,” depending on the source and context. To introduce it, for  $x \in \mathbb{Z}$ , let  $P_x$  denote the law of the continuous-time simple random walk  $(X_t)_{t \geq 0}$  on  $\mathbb{Z}$  with jump rate 1 and denote by  $E_x$  the corresponding expectation. The following proposition, which is an adaptation of Section 4.2 of [33] or Theorem 2.1 of [30], gives the representation for first and second moments. Its proof is an easy modification of the proofs of these results, and it is therefore omitted.

**PROPOSITION 3.1 (Feynman–Kac formula).** *Let  $\varphi_1, \varphi_2$  be càdlàg functions from  $[0, t]$  to  $[-\infty, \infty]$  satisfying  $\varphi_1 \leq \varphi_2$ . Then the first and second moments of the number of particles in  $N(t)$  whose genealogy stays between  $\varphi_1$  and  $\varphi_2$  are given by*

$$(3.1) \quad \begin{aligned} & E_0^{\xi}[\{|Y \in N(t) : \varphi_1(r) \leq Y_r \leq \varphi_2(r) \ \forall r \in [0, t]\}|] \\ &= E_0 \left[ \exp \left\{ \int_0^t \xi(X_r) \, dr \right\}; \varphi_1(r) \leq X_r \leq \varphi_2(r) \ \forall r \in [0, t] \right], \end{aligned}$$

$$(3.2) \quad \begin{aligned} & E_0^{\xi}[\{|Y \in N(t) : \varphi_1(r) \leq Y_r \leq \varphi_2(r) \ \forall r \in [0, t]\}^2] \\ &= E_0 \left[ \exp \left\{ \int_0^t \xi(X_r) \, dr \right\}; \varphi_1(r) \leq X_r \leq \varphi_2(r) \ \forall r \in [0, t] \right] \\ &+ 2 \int_0^t E_0 \left[ \exp \left\{ \int_0^s \xi(X_r) \, dr \right\} \xi(X_s) \mathbf{1}_{\varphi_1(r) \leq X_r \leq \varphi_2(r) \ \forall r \in [0, s]} \right. \\ &\quad \left. \times \left( E_{X_s} \left[ \exp \left\{ \int_0^{t-s} \xi(X_r) \, dr \right\} \mathbf{1}_{\varphi_1(r+s) \leq X_r \leq \varphi_2(r+s) \ \forall r \in [0, t-s]} \right] \right)^2 \right] ds. \end{aligned}$$

In particular, (3.1) implies that

$$(3.3) \quad E_{u_0}^{\xi}[N^{\geq}(t, n)] = \sum_{i \in \mathbb{Z}} u_0(i) E_i \left[ \exp \left\{ \int_0^t \xi(X_r) \, dr \right\}; X_t \geq n \right].$$

In the first principal step of the proof, we analyze the first moment formula (3.3) for  $n = vt$  with  $v > 0$ . To understand this analysis, it is useful to recall the corresponding representation from the homogeneous case (cf. [1]). In that setting, it is almost trivial that  $E_0^{\xi \equiv 1}[N^{\geq}(t, vt)] = e^t P_0(X_t \geq vt)$ . The probability on the right-hand side can then be analyzed using exact large deviation results (see, e.g., [18], Theorem 3.7.4) to obtain a precise asymptotic formula.

While (3.3) has a different structure, its asymptotics can be understood, at least at a heuristic level, by the same ingredients that are usually used in the proof of exact large deviation theorems: a tilting and a local central limit theorem. Slightly more in detail, by introducing a tilting parameter  $\eta$ , using (3.3), we can write

$$(3.4) \quad E_0^{\xi}[N(t, vt)] = e^{-t\eta} E_0 \left[ \exp \left\{ \int_0^t (\xi(X_r) + \eta) \, dr \right\}; X_t = vt \right].$$

This suggests to introduce new “Gibbs measures” on the space of random walk trajectories, whose density with respect to simple random walk is the exponential factor in (3.4) (cf. Section 4.1). We then adjust  $\eta$  so that the event  $X_t = vt$  is typical under such a Gibbs measure.

Next, using a suitable local central limit theorem, the right-hand side of (3.4) can be approximated by (cf. Proposition 4.10)

$$\sim \frac{c}{\sqrt{t}} e^{-t\eta} E_0 \left[ \exp \left\{ \int_0^t (\xi(X_r) + \eta) dr \right\}; H_{vt} \leq t \right],$$

where  $H_x$  stands for the hitting time of  $x$  by the simple random walk  $X$ ; see (4.1) below. This can further be rewritten as

$$\sim \frac{c}{\sqrt{t}} e^{-t\eta} E_0 \left[ \prod_{x=1}^{vt} \exp \left\{ \int_{H_{x-1}}^{H_x} (\xi(X_r) + \eta) dr \right\} \times \exp \left\{ \int_{H_{vt}}^t (\xi(X_r) + \eta) dr \right\} \right].$$

If one ignores the last factor in the expectation, which can be justified using the concentration of the hitting times of the random walk under the Gibbs measure (see Section 4.4), then by the Markov property

$$\begin{aligned} &\sim \frac{c}{\sqrt{t}} e^{-t\eta} \prod_{x=1}^{vt} E_{x-1} \left[ \exp \left\{ \int_0^{H_x} (\xi(X_r) + \eta) dr \right\} \right] \\ &= \frac{c}{\sqrt{t}} e^{-t\eta} \exp \left\{ \sum_{x=1}^{vt} \ln E_{x-1} \left[ \exp \left\{ \int_0^{H_x} (\xi(X_r) + \eta) dr \right\} \right] \right\}. \end{aligned}$$

The application of a suitable central limit theorem to the above sum then suggests the central limit theorem behavior of the PAM, Theorem 2.8.

Making the above heuristics rigorous requires a nonnegligible effort. In particular, it turns out that the tilting parameter  $\eta$  making the event  $X_t = vt$  typical under the Gibbs measure is *random* (i.e.,  $\xi$ -dependent). This disallows a straightforward application of a central limit theorem in the last formula above. Section 4.3 deals with this problem, building on a preparatory Section 4.2. Other approximations appearing in the previous heuristic computation are treated in Section 4.4; Section 4.5 controls the influence of the initial conditions. The functional central limit theorem for the PAM, Theorem 2.8, then follows easily; cf. Section 4.6.

In order to show the functional central limit theorem for the breakpoint, Theorem 2.6, we essentially need to find the largest root of the function  $x \mapsto \ln E_0^\xi [N^\geq(t, x)]$ , which requires, in a certain sense, to invert the functional central limit theorem for the PAM; cf. Section 5.2. In order to perform this inversion, we study how sensitive  $E_0^\xi [N^\geq(t, x)]$  is to perturbations in the space and time direction; cf. Section 5.1.

Let us now comment on the second moment computation required to prove the remaining main results of this paper. Similar to the homogeneous case, the second moment of  $N^\geq(t, vt)$  explodes too quickly to yield any useful estimates. This explosion is, essentially, due to particles that are much faster than the breakpoint at times in the bulk of the interval  $[0, t]$ . In the case of homogeneous branching rates, this is solved by a truncation which involves considering only so-called *leading* particles, that is, the particles that are slower than the breakpoint,  $X_s \leq v_0 s$  for all  $s \in [0, t]$  (here,  $v_0 t$  is a first-order breakpoint asymptotics). The principal ingredient for the computation of the moments for the leading particles is then a “ballot theorem” for the random walk bridge, which gives the probability that a random walk bridge from  $(0, 0)$  to  $(t, 0)$  stays positive for all intermediate times.

Following the above strategy in the case of BRWRE suggests to call a particle  $Y \in N(t)$  *leading at time  $t$*  if (a)  $Y_t$  is close to the breakpoint  $\bar{m}(t)$ , and (b)  $Y$  is slower than breakpoint at intermediate times,  $Y_s \leq \bar{m}(s)$  for all  $s \in [0, t]$ .<sup>1</sup> Since  $\bar{m}(t)$  satisfies a functional central limit theorem itself, it naturally leads to a ballot estimate of the following form: Let  $B, W$

<sup>1</sup>The actual definition of leading particles in Section 6 is slightly different for technical reasons.

be two independent Brownian motions (or centered random walks, possibly not identically distributed). What is the behavior of

$$\mathbb{P}(B(t) \geq W(t), B(s) \leq 1 + W(s) \forall s \in [0, t] \mid \sigma(W))?$$

Observe that the process  $W$  is “quenched” in this computation as we condition on the  $\sigma$ -field  $\sigma(W)$  generated by  $W$ . This modified ballot problem was recently studied by Mallein and Miłoś [51]. We were however not able to use their results directly due to the lack of the independence that we encounter in our model.

The first and second moment of the number of leading particles is computed in Section 6.1. In particular, a lengthy proof of a (relatively weak version of) a ballot estimate can be found in Section 6.1.1. Subsequently, Theorem 2.4 and Theorem 2.1 are then shown in Section 6.2. Section 7 proves the functional central limit theorem for the Fisher–KPP equation, Theorem 2.9. Finally, Section 8 discusses some open problems.

*Notational conventions.* For two functions  $f, g : [0, \infty) \rightarrow (0, \infty)$ , we write  $f \sim g$  when  $\lim_{t \rightarrow \infty} f(t)/g(t) = 1$ , and  $f \asymp g$  when  $0 < \inf_{t \in [0, \infty)} f(t)/g(t) \leq \sup_{t \in [0, \infty)} f(t)/g(t) < \infty$ . We use  $c$  and  $C$  to denote positive finite constants whose value may change during computations, and sometimes write  $c(\xi)$ , etc., in order to emphasize their dependence on realizations of the branching rates. Indexed constants such as  $c_1$  keep their value from their first time of occurrence. We use  $E[f; A]$  as an abbreviation for  $E[f \mathbf{1}_A]$ .

For  $x \in \mathbb{R} \setminus \mathbb{Z}$ , we define  $P_x$  by linear interpolation. More precisely, for  $x = \lfloor x \rfloor + \lambda$  we define  $P_x := (1 - \lambda)P_{\lfloor x \rfloor} + \lambda P_{\lfloor x \rfloor + 1}$ . Similarly, other quantities which are only defined for integers a priori are to be interpreted as the linear interpolation of the evaluations at their integer neighbors, which will usually be clear from the context.

While we have stated the precise assumptions needed in the main results given above,

we will from now on assume (POT), (INI) and (VEL) to be fulfilled

as standing assumptions without further mentioning. This helps in keeping notation lighter compared to mentioning a suitable subset of these assumptions at each of the numerous subsequent auxiliary results.

**4. Expected number of particles of given velocity.** In this section, we study the asymptotic behavior of  $\mathbb{E}_{u_0}^{\xi}[N(t, vt)]$  and related quantities, following the strategy described in Section 3.

4.1. *Tilted random walk measures.* We introduce the tilted distributions of random walk in random potential, and show that one can tilt the random walk in a suitable way to make the extremal behavior typical.

Recall that  $(X_t)_{t \geq 0}$  denotes continuous-time simple random walk on  $\mathbb{Z}$  with jump rate 1. For  $i \in \mathbb{Z}$ , we define the hitting time of  $i$  as

$$(4.1) \quad H_i := \inf\{s \in [0, \infty) : X_s = i\},$$

and set  $\tau_i := H_i - H_{i-1}$ . Recalling (POT) and writing

$$(4.2) \quad \zeta(x) := \xi(x) - \mathbf{es}, \quad x \in \mathbb{Z},$$

we infer

$$(4.3) \quad -\infty < \text{ess inf } \zeta < \text{ess sup } \zeta = 0.$$

For  $n \geq 1$ ,  $A \in \sigma(X_{s \wedge H_n}, s \in [0, \infty))$  and  $\eta \in \mathbb{R}$ , we define

$$(4.4) \quad P_{(n)}^{\zeta, \eta}(A) := (Z_{(n)}^{\zeta, \eta})^{-1} E_0 \left[ \exp \left\{ \int_0^{H_n} (\zeta(X_s) + \eta) ds \right\}; A \right],$$

where

$$Z_{(n)}^{\zeta,\eta} := E_0 \left[ \exp \left\{ \int_0^{H_n} (\zeta(X_s) + \eta) \, ds \right\} \right].$$

We will see below (cf. Lemma 4.1) that these quantities are finite if and only if  $\eta \leq 0$ .

It can be seen easily, using the strong Markov property, that  $P_{(n)}^{\zeta,\eta}(A) = P_{(m)}^{\zeta,\eta}(A)$  for every  $m > n$  and  $A \in \sigma(X_{s \wedge H_n}, s \in [0, \infty))$ . We may thus use Kolmogorov’s extension theorem to extend  $P_{(n)}^{\zeta,\eta}$  to a measure  $P^{\zeta,\eta}$  on  $\sigma(X_s, s \geq 0)$ . We write  $P^\zeta$  for  $P^{\zeta,0}$ .

It will be suitable to introduce the following logarithmic moment generating functions:

$$(4.5) \quad L_i^\zeta(\eta) := \ln E_{i-1} \left[ \exp \left\{ \int_0^{H_i} (\zeta(X_s) + \eta) \, ds \right\} \right],$$

$$(4.6) \quad \bar{L}_n^\zeta(\eta) := \frac{1}{n} \sum_{i=1}^n L_i^\zeta(\eta),$$

$$(4.7) \quad L(\eta) := \mathbb{E}[L_1^\zeta(\eta)].$$

By the strong Markov property again,

$$(4.8) \quad Z_{(n)}^{\zeta,\eta} = \exp \left\{ \sum_{i=1}^n L_i^\zeta(\eta) \right\} = \exp \{ n \bar{L}_n^\zeta(\eta) \}.$$

We now discuss the finiteness of the above objects.

LEMMA 4.1. *Under (POT), the quantities defined in (4.5)–(4.7) are finite if and only if  $\eta \leq 0$ .*

PROOF. Since  $\text{ess sup } \zeta(x) \leq 0$ , the “if” part of the lemma is trivial.

The “only if” part can be proved via the following strategy: For  $i \in \mathbb{Z}$  and  $\eta > 0$ , using the independence assumption of the potential in (POT), the random walk starting in  $i$  can find arbitrarily large islands to the left of  $i$ , where the potential  $\zeta + \eta$  takes values larger than  $\eta/2$ . Once such an island is large enough so that the cost of the random walk to stay inside this island is offset by the exponential gain of a potential value larger than  $\eta/2$  in the Feynman–Kac formula, one infers that  $L_i^\zeta(\eta)$  is infinite, and then the same applies to the remaining quantities in question.

Since in the case of random walk with a drift, the “only if” statement is a direct consequence of Proposition 3.1 in [19], we omit making the above proof rigorous. The lengthy proof of [19], however, can directly be transferred to the case of simple random walk without drift.  $\square$

Recalling that  $\tau_i = H_i - H_{i-1}$ , as an easy corollary of Lemma 4.1 we obtain  $\ln E^{\zeta,\eta}[e^{\lambda\tau_i}] = L_i^\zeta(\eta + \lambda) - L_i^\zeta(\eta)$  for every  $\eta \leq 0$  and  $\lambda \in \mathbb{R}$ , as well as

$$(4.9) \quad E^{\zeta,\eta}[e^{\lambda\tau_i}] < \infty \quad \text{for every } \eta \leq 0 \text{ and } \lambda \leq |\eta|.$$

Finally, Birkhoff’s ergodic theorem implies that

$$(4.10) \quad L(\eta) \stackrel{\mathbb{P}\text{-a.s.}}{=} \lim_{n \rightarrow \infty} \bar{L}_n^\zeta(\eta).$$

Other simple properties of functions  $L^\zeta$  and  $L$  are given in the [Appendix](#).

We will primarily be interested in those values  $\eta = \bar{\eta}_n^\zeta(v)$  which make certain large deviations events typical, more precisely for which

$$(4.11) \quad E^{\zeta, \bar{\eta}_n^\zeta(v)}[H_n] = \frac{n}{v}, \quad v > 0.$$

In order to discuss the existence of such  $\eta$ , which is random, we introduce, in the next lemma, its “typical value”  $\bar{\eta}(v)$ . We recall the *critical velocity*  $v_c$  introduced below (2.2). It will be shown in Proposition A.3, partially using the results of this paper, that the identity

$$(4.12) \quad (v_c)^{-1} = L'(0)$$

holds true, where the derivative is taken from the left only. Throughout the paper, we use (4.12) as the primary definition of  $v_c$ .

LEMMA 4.2. *For every  $v > v_c$ , there exists a unique  $\bar{\eta}(v) \in (-\infty, 0)$  such that*

$$(4.13) \quad L^*(1/v) = \sup_{\eta \in \mathbb{R}} (\eta/v - L(\eta)) = \bar{\eta}(v)/v - L(\bar{\eta}(v)).$$

Furthermore,  $\bar{\eta}(v)$  is characterized by

$$(4.14) \quad L'(\bar{\eta}(v)) = v^{-1}.$$

Moreover,  $(v_c, \infty) \ni v \mapsto \bar{\eta}(v)$  is a smooth strictly decreasing function.

PROOF. Due to Lemma A.1,  $L$  is smooth, strictly increasing and strictly convex on  $(-\infty, 0)$ , finite on  $(-\infty, 0]$ , and infinite on  $(0, \infty)$ ; cf. Lemma 4.1. In addition, it can be seen easily that  $\lim_{\eta \rightarrow -\infty} L'(\eta) = 0$  (see [19], Lemma 3.5, for the corresponding statement in the case of a random walk with drift; the proof for simple random walk proceeds in the same way and is omitted here). Therefore, recalling also (4.12), we see that the solution to (4.13) exists for every  $v > v_c$ . Furthermore, due to usual properties of the Legendre transform, it is characterized by (4.14). The last statement follows directly from the previously mentioned properties.  $\square$

We now show that  $\bar{\eta}_n^\zeta(v)$  fulfilling (4.11) exists  $\mathbb{P}$ -a.s. for  $v > v_c$  and  $n$  large enough and, in fact, concentrates around  $\bar{\eta}(v)$ .

PROPOSITION 4.3. *For each  $v > v_c$ , there exists a  $\mathbb{P}$ -a.s. finite random variable  $\mathcal{N} = \mathcal{N}(v)$  such that for all  $n \geq \mathcal{N}$  there exists  $\bar{\eta}_n^\zeta(v) \in (-\infty, 0)$  satisfying (4.11). Moreover, for every  $q \in \mathbb{N}$  and  $V \subset (v_c, \infty)$  compact, there exists a constant  $C = C(q, V) < \infty$  such that for all  $n \in \mathbb{N}$ ,*

$$(4.15) \quad \mathbb{P}\left(\sup_{v \in V} |\bar{\eta}(v) - \bar{\eta}_n^\zeta(v)| \geq C \sqrt{\frac{\ln n}{n}}\right) \leq Cn^{-q}$$

(defining, arbitrarily,  $\bar{\eta}_n^\zeta(v) = 0$  if the solution to (4.11) does not exist).

PROOF. By Lemma A.1, for  $\eta < 0$ ,  $E^{\zeta, \eta}[H_n] = n(\bar{L}_n^\zeta)'(\eta)$ . Hence, in combination with (4.11), we may define  $\bar{\eta}_n^\zeta(v)$  as the solution to

$$(4.16) \quad (\bar{L}_n^\zeta)'(\bar{\eta}_n^\zeta(v)) = 1/v,$$

if this solution exists, and by  $\bar{\eta}_n^\zeta(v) = 0$  otherwise. If we show that this  $\bar{\eta}_n^\zeta(v)$  satisfies (4.15), then the fact  $\bar{\eta}_n^\zeta(v) \in (-\infty, 0)$  for all  $n \geq \mathcal{N}$  follows by a Borel–Cantelli argument, using also that  $\sup_{v \in V} \bar{\eta}(v) < 0$  by Lemma 4.2, as well as the compactness of  $V$ .

Comparing (4.14) and (4.16), we see that we need to understand the concentration properties of  $(\bar{L}_n^\zeta)'$  first. We claim the following.

CLAIM 4.4. For every  $q \in \mathbb{N}$  and  $\Delta \subset (-\infty, 0)$  compact, there exists  $C = C(q, \Delta) < \infty$  such that for all  $n \in \mathbb{N}$ ,

$$(4.17) \quad \mathbb{P}\left(\sup_{\eta \in \Delta} |(\bar{L}_n^\zeta)'(\eta) - L'(\eta)| \geq C \sqrt{\frac{\ln n}{n}}\right) \leq Cn^{-q}.$$

PROOF. We apply a Hoeffding-type bound for mixing sequences which we recall in Lemma A.5. Define the  $\sigma$ -algebras  $\mathcal{F}_k := \sigma(\xi(i) : i \leq k)$ ,  $k \in \mathbb{Z}$ . By Lemma A.1,  $((L_i^\zeta)'(\eta) - L'(\eta))_{i \in \mathbb{Z}}$  is a stationary sequence of bounded random variables. By Lemma A.2, there is  $c < \infty$  such that  $|\mathbb{E}[(L_i^\zeta)'(\eta) | \mathcal{F}_k] - L'(\eta)| \leq ce^{-(i-k)/c}$  for all  $i \geq k$  and  $\eta \in \Delta$ . Hence, the assumptions of Lemma A.5 are satisfied with  $m_i = c$ , and thus uniformly over  $\eta \in \Delta$ , for  $C$  large enough,

$$\mathbb{P}\left(|(\bar{L}_n^\zeta)'(\eta) - L'(\eta)| \geq C \sqrt{\frac{\ln n}{n}}\right) \leq Ce^{-C \ln n} \leq Cn^{-q-1}.$$

Hence, by a union bound,

$$(4.18) \quad \mathbb{P}\left(\sup_{\eta \in \frac{1}{n}\mathbb{Z} \cap \Delta} |(\bar{L}_n^\zeta)'(\eta) - L'(\eta)| \geq C \sqrt{\frac{\ln n}{n}}\right) \leq Cn^{-q}.$$

Moreover, by Lemma A.1,  $L'$  and  $(\bar{L}_n^\zeta)'$  are both increasing on  $(-\infty, 0)$  with continuous and positive derivatives. Hence, for any  $\Delta \subset (-\infty, 0)$  compact, there is  $c < \infty$  such that

$$(4.19) \quad c^{-1} < \inf_{\Delta} L'' \leq \sup_{\Delta} L'' < c.$$

Combining this with (4.18) and the fact that  $L'$  and  $(\bar{L}_n^\zeta)'$  are increasing again, this implies the claim.  $\square$

To prove (4.15), fix a compact  $\Delta \subset (-\infty, 0)$  such that  $\bar{\eta}(V)$  is contained in the interior of  $\Delta$ , which is possible by Lemma 4.2, and set  $\delta = \text{dist}(\bar{\eta}(V), \Delta^c) > 0$ . By (4.14) and (4.16),  $\bar{\eta}(v)$  and  $\bar{\eta}_n^\zeta(v)$  are the respective solutions to  $L'(\bar{\eta}(v)) = v^{-1}$  and  $(\bar{L}_n^\zeta)'(\bar{\eta}_n^\zeta(v)) = v^{-1}$  (if the solution to the second equation exists). Moreover, by (4.19), the slope of  $L'$  on  $\Delta$  is at least  $c^{-1}$ . Therefore, on the complement of the event in the probability on the left-hand side of (4.17), for  $n$  large enough so that  $C\sqrt{\ln(n)/n} < c^{-1}\delta$ , we know that for all  $v \in V$  the equation (4.16) has a solution  $\bar{\eta}_n^\zeta(v)$  which satisfies  $|\bar{\eta}_n^\zeta(v) - \bar{\eta}(v)| \leq cC\sqrt{\ln(n)/n} < \delta$ . Hence, (4.15) follows from (4.17) by adjusting constants.  $\square$

For future reference, we recall that whenever  $\bar{\eta}_n^\zeta(v)$  exists, then it is characterized, due to the usual properties of the Legendre transform, by

$$(4.20) \quad (\bar{L}_n^\zeta)^*(1/v) := \sup_{\eta \in \mathbb{R}} \left(\frac{\eta}{v} - \bar{L}_n^\zeta(\eta)\right) = \frac{\bar{\eta}_n^\zeta(v)}{v} - \bar{L}_n^\zeta(\bar{\eta}_n^\zeta(v)).$$

*Technical assumption.* In order to keep the constants in the paper independent of the velocity  $v$ , for the rest of this paper we assume that

$$(4.21) \quad \begin{aligned} &\text{the velocities } v \text{ that we are considering are contained in a fixed compact interval} \\ &V \subset (v_c, \infty) \text{ which has } v_0 \text{ in its interior.} \end{aligned}$$

Such  $V$  exists due to (VEL). The constants appearing in the results below may depend on  $V$ . Using Proposition 4.3 and the monotonicity of  $\bar{\eta}$  and  $\bar{\eta}_n^\zeta$  in  $v$  and  $\zeta$ , it is then possible to fix a

compact interval  $\Delta \subset (-\infty, 0)$  such that there is a  $\mathbb{P}$ -a.s. finite random variable  $\mathcal{N}_1$  such that the event

$$(4.22) \quad \mathcal{H}_n := \mathcal{H}_n(V) := \{\bar{\eta}_n^\zeta(v) \in \Delta \text{ for all } v \in V\} \quad \text{occurs for all } n \geq \mathcal{N}_1.$$

We also recall that we arbitrarily set  $\bar{\eta}_n^\zeta(v) = 0$  in the case when (4.11) does not have any solution. This occurs on  $\mathcal{H}_n^c$  only.

For future use, we state the following easy estimate.

LEMMA 4.5. *For each  $\delta \in (0, 1)$ , there exists a constant  $C = C(\delta)$  such that  $\mathbb{P}$ -a.s. for all  $n$  large enough, uniformly for  $v \in V$  and  $h \leq n^{1-\delta}$ ,*

$$|\bar{\eta}_n^\zeta(v) - \bar{\eta}_{n+h}^\zeta(v)| \leq \frac{Ch}{n}.$$

PROOF. Let  $\Delta$  be as in (4.22). We claim that there exists a constant  $C < \infty$  such that for all  $n \geq 1$ ,  $h \leq n$  and  $\eta \in \Delta$ ,

$$(4.23) \quad |(\bar{L}_{n+h}^\zeta)'(\eta) - (\bar{L}_n^\zeta)'(\eta)| \leq \frac{Ch}{n}.$$

Indeed, plugging in the definitions we obtain

$$(\bar{L}_{n+h}^\zeta)'(\eta) - (\bar{L}_n^\zeta)'(\eta) = -\frac{h}{n(n+h)} \sum_{i=1}^n (L_i^\zeta)'(\eta) + \left(\frac{1}{n+h}\right) \sum_{i=n+1}^{n+h} (L_i^\zeta)'(\eta),$$

from which we can then deduce (4.23) by observing that  $(L_i^\zeta)'(\eta)$  can be bounded uniformly over  $\mathbb{P}$ -a.a. realizations of  $\zeta$  and  $\eta \in \Delta$ , by Lemma A.1. The claim of the lemma then follows from (4.14), (4.16), (4.19) and (4.23) by the same arguments as at the end of the proof of Proposition 4.3.  $\square$

4.2. *An invariance principle for the empirical Legendre transforms.* In this section, we show an invariance principle for the suitably centered and rescaled Legendre transforms of the functions  $\bar{L}_n^\zeta$  defined in (4.6). In order to state them, we introduce

$$(4.24) \quad V_i^{\zeta, v}(\eta) := \eta/v - L_i^\zeta(\eta),$$

$$(4.25) \quad \sigma_v^2 := \text{Var}_{\mathbb{P}}(V_1^{\zeta, v}(\bar{\eta}(v))) + 2 \sum_{j \geq 2} \text{Cov}_{\mathbb{P}}(V_1^{\zeta, v}(\bar{\eta}(v)), V_j^{\zeta, v}(\bar{\eta}(v))).$$

Using the nondegeneracy part of assumption (POT), and the exponential decay of correlations of the  $L_i^\zeta$  proved in Lemma A.2, we see that  $\sigma_v^2 \in (0, \infty)$ .

PROPOSITION 4.6. *For each  $v \in V$ , the sequence of processes*

$$(4.26) \quad t \mapsto W_n(t) := \frac{1}{\sigma_v} t \sqrt{n} ((\bar{L}_{nt}^\zeta)^*(1/v) - L^*(1/v)), \quad n \in \mathbb{N},$$

*converges as  $n \rightarrow \infty$ , in  $\mathbb{P}$ -distribution to standard Brownian motion.*

Heuristically, the proof of this proposition is based on the fact that the fluctuations of the Legendre transforms  $(\bar{L}_n^\zeta)^*$  are essentially given by the fluctuations of the functions  $\bar{L}_n^\zeta$ , whereas the influence of the fluctuations of the maximizing argument at which the supremum is attained in the definition (4.20) of the Legendre transform is negligible.



PROOF OF PROPOSITION 4.6. Recall that, due to (4.20), on  $\mathcal{H}_n$ ,

$$(4.27) \quad (\bar{L}_n^\zeta)^*(1/v) = \frac{\bar{\eta}_n^\zeta(v)}{v} - \bar{L}_n^\zeta(\bar{\eta}_n^\zeta(v)) = \frac{1}{n} \sum_{i=1}^n V_i^{\zeta,v}(\bar{\eta}_n^\zeta(v)) = \frac{1}{n} S_n^{\zeta,v}(\bar{\eta}_n^\zeta(v)),$$

where we set

$$(4.28) \quad S_n^{\zeta,v}(\eta) := \sum_{i=1}^n V_i^{\zeta,v}(\eta)$$

as a shorthand. Using this notation, we expand the quantity of interest as

$$(4.29) \quad \begin{aligned} tn((\bar{L}_{in}^\zeta)^*(1/v) - L^*(1/v)) &= (tn(\bar{L}_{in}^\zeta)^*(1/v) - S_{in}^{\zeta,v}(\bar{\eta}(v))) \\ &+ (S_{in}^{\zeta,v}(\bar{\eta}(v)) - \mathbb{E}[S_{in}^{\zeta,v}(\bar{\eta}(v))]) \\ &+ (\mathbb{E}[S_{in}^{\zeta,v}(\bar{\eta}(v))] - tnL^*(1/v)). \end{aligned}$$

We will show that the first and the third summand on the right-hand side are negligible in a suitable sense, and that the second summand converges in distribution after rescaling by  $\sigma_v \sqrt{n}$  to standard Brownian motion under  $\mathbb{P}$ .

The third summand in (4.29) is the easiest since it vanishes. Indeed, by (4.7) and (4.13),

$$tnL^*(1/v) = tn\left(\frac{\bar{\eta}(v)}{v} - L(\bar{\eta}(v))\right) = tn\mathbb{E}\left[\frac{\bar{\eta}(v)}{v} - \bar{L}_{in}^\zeta(\bar{\eta}(v))\right] = \mathbb{E}[S_{in}^{\zeta,v}(\bar{\eta}(v))].$$

The next lemma deals with the second summand in (4.29).

LEMMA 4.7. *The sequence of processes*

$$[0, \infty) \ni t \mapsto \widetilde{W}_n(t) := \frac{1}{\sigma_v \sqrt{n}} (S_{in}^{\zeta,v}(\bar{\eta}(v)) - \mathbb{E}[S_{in}^{\zeta,v}(\bar{\eta}(v))]), \quad n \in \mathbb{N},$$

converges as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to standard Brownian motion.

PROOF. By the definition of  $S_n^{\zeta,v}$ ,

$$\frac{1}{\sigma_v \sqrt{n}} (S_{in}^{\zeta,v}(\bar{\eta}(v)) - \mathbb{E}[S_{in}^{\zeta,v}(\bar{\eta}(v))]) = \frac{1}{\sigma_v \sqrt{n}} \sum_{i=1}^{tn} V_i^{\zeta,v}(\bar{\eta}(v)) - \mathbb{E}[V_i^{\zeta,v}(\bar{\eta}(v))].$$

The  $V_i^{\zeta,v}(\bar{\eta}(v))$  form a nondegenerate stationary sequence of random variables, which are coordinatewise decreasing in the  $\zeta$ 's. Therefore, by the FKG-inequality, they also form an associated sequence in the sense that any two coordinatewise decreasing functions of the  $V_i^{\zeta,v}(\bar{\eta}(v))$ 's of finite variance are nonnegatively correlated. Hence, the functional central limit theorem for associated random variables proved in [57], Theorem 3, supplies us with convergence in  $C([0, M])$  for each  $M \in (0, \infty)$ , and the result is then extended to  $C([0, \infty))$  in the standard fashion.  $\square$

Finally, for the first summand in (4.29), we have the following estimate.

LEMMA 4.8. *There is  $C < \infty$  such that  $\mathbb{P}$ -a.s. for every  $M \in (1, \infty)$  and  $v \in V$ ,*

$$\limsup_{n \rightarrow \infty} \frac{1}{\ln n} \sup_{t \in [0, M]} |tn(\bar{L}_{in}^\zeta)^*(1/v) - S_{in}^{\zeta,v}(\bar{\eta}(v))| \leq C.$$

PROOF. By Proposition 4.3 and (4.22), the representation (4.27) holds for all  $n \geq \mathcal{N}_1$ , with  $\mathcal{N}_1$  a  $\mathbb{P}$ -a.s. finite random variable. As a consequence, it is sufficient to show that  $\mathbb{P}$ -a.s.,

$$(4.30) \quad \limsup_{n \rightarrow \infty} \frac{1}{\ln n} \max_{\mathcal{N}_1 \leq k \leq Mn} |S_k^{\zeta, v}(\tilde{\eta}_k^\zeta(v)) - S_k^{\zeta, v}(\bar{\eta}(v))| \leq C.$$

Assuming  $k \geq \mathcal{N}_1$  in what follows, using a Taylor expansion of the smooth function  $S_k^{\zeta, v}$  around  $\tilde{\eta}_k^\zeta(v)$  we get

$$(4.31) \quad \begin{aligned} S_k^{\zeta, v}(\bar{\eta}(v)) - S_k^{\zeta, v}(\tilde{\eta}_k^\zeta(v)) &= (S_k^{\zeta, v})'(\tilde{\eta}_k^\zeta(v))(\bar{\eta}(v) - \tilde{\eta}_k^\zeta(v)) \\ &\quad + (S_k^{\zeta, v})''(\tilde{\eta}_k^\zeta(v)) \frac{(\bar{\eta}(v) - \tilde{\eta}_k^\zeta(v))^2}{2}, \end{aligned}$$

for some  $\tilde{\eta}_k^\zeta \in \Delta$  with  $|\tilde{\eta}_k^\zeta - \tilde{\eta}_k^\zeta(v)| \leq |\bar{\eta}(v) - \tilde{\eta}_k^\zeta(v)|$ .

By (4.20),  $S_k^{\zeta, v}(\eta)$  is maximized for  $\eta = \tilde{\eta}_k^\zeta(v)$ , so  $(S_k^{\zeta, v})'(\tilde{\eta}_k^\zeta(v)) = 0$  and the first term on the right-hand side of (4.31) vanishes.

To bound the second term, observe that  $(S_k^{\zeta, v})''(\tilde{\eta}_k^\zeta) = -k(\bar{L}_k^\zeta)''(\tilde{\eta}_k^\zeta)$ . By Lemma A.1,  $\mathbb{P}$ -a.s.,  $(L_1^\zeta)''(\eta)$  is bounded from above, uniformly over  $\eta \in \Delta$  (cf. (4.22)). Hence,  $\mathbb{P}$ -a.s.,

$$(4.32) \quad (S_k^{\zeta, v})''(\tilde{\eta}_k^\zeta) \in [-Ck, 0] \quad \text{for all } k \geq \mathcal{N}_1, v \in V.$$

Going back to (4.31),  $\mathbb{P}$ -a.s. for all  $k \geq \mathcal{N}_1$ ,

$$|S_k^{\zeta, v}(\bar{\eta}(v)) - S_k^{\zeta, v}(\tilde{\eta}_k^\zeta(v))| \leq ck|\bar{\eta}(v) - \tilde{\eta}_k^\zeta(v)|^2.$$

Using the concentration estimates for  $\tilde{\eta}_k^\zeta(v)$  from Proposition 4.3, it is possible to fix a constant  $C < \infty$  and a  $\mathbb{P}$ -a.s. finite random variable  $\mathcal{N}_2 \geq \mathcal{N}_1$  such that for all  $k \geq \mathcal{N}_2$ ,  $|\tilde{\eta}_k^\zeta(v) - \bar{\eta}(v)| \leq C\sqrt{\ln k/k}$ . Putting all together, this implies that  $\mathbb{P}$ -a.s. the left-hand side in (4.30) is bounded by

$$\limsup_{n \rightarrow \infty} \frac{1}{\ln n} \left\{ \max_{\mathcal{N}_1 \leq k \leq \mathcal{N}_2} |S_k^{\zeta, v}(\tilde{\eta}_k^\zeta(v)) - S_k^{\zeta, v}(\bar{\eta}(v))| + \max_{\mathcal{N}_2 \leq k \leq Mn} C \ln k \right\} \leq C.$$

This completes the proof.  $\square$

Proposition 4.6 now follows from (4.29) and Lemmas 4.7 and 4.8.  $\square$

The proof of Proposition 4.6 has the following corollary which provides a useful explicit approximation to  $W_n(t)$ .

COROLLARY 4.9. *There is a constant  $C < \infty$  such that  $\mathbb{P}$ -a.s. for every  $M \in (0, \infty)$  and  $v \in V$ ,*

$$\limsup_{n \rightarrow \infty} \frac{1}{\ln n} \sup_{t \in [0, M]} \left| \sigma_v \sqrt{n} W_n(t) - \sum_{i=1}^{nt} (L(\bar{\eta}(v)) - L_i^\zeta(\bar{\eta}(v))) \right| \leq C.$$

PROOF. It suffices to use the definition (4.26) of  $W_n(t)$  together with Lemma 4.8. The claim then follows after a straightforward computation by inserting the definition of  $S_{in}^{\zeta, v}(\bar{\eta}(v))$  and using that  $L^*(1/v) = \bar{\eta}(v)/v - L(\bar{\eta}(v))$ .  $\square$

4.3. *An auxiliary invariance principle.* We now prove an invariance principle for the logarithm of the auxiliary process

$$Y_v(n) := E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\}; H_n \leq \frac{n}{v} \right], \quad n \in \mathbb{N}, v \in V,$$

which we will relate to quantities considered in the Feynman–Kac representation (3.1) later on. Observe that this invariance principle can be seen as a first step to exact large deviation estimates, as explained in Section 3 above.

For convenience, we split the process  $Y_v$  into the two summands

$$(4.33) \quad \begin{aligned} Y_v^{\approx}(n) &:= E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\}; H_n \in \left[ \frac{n}{v} - K, \frac{n}{v} \right] \right] \quad \text{and} \\ Y_v^{<}(n) &:= E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\}; H_n < \frac{n}{v} - K \right], \end{aligned}$$

where  $K > 0$  is a large constant which will be fixed later on.

For  $n \in \mathbb{N}$  and  $v \in V$ , we define random variables  $\sigma_n^\zeta(v)$ :

$$(4.34) \quad \sigma_n^\zeta(v) := \begin{cases} |\bar{\eta}_n^\zeta(v)| \sqrt{\text{Var}_{P^{\zeta, \bar{\eta}_n^\zeta(v)}}[H_n]} & \text{on } \mathcal{H}_n, \\ \max \Delta \sqrt{\text{Var}_{P^{\zeta, \max \Delta}}[H_n]} & \text{on } \mathcal{H}_n^c. \end{cases}$$

Under every  $P^{\zeta, \eta}$ , we can write  $H_n = \sum_{i=1}^n \tau_i$  as a sum of independent random variables (see (4.1) and below). Moreover, by Lemma A.1, there is a constant  $c < \infty$  such that  $c^{-1} \leq \text{Var}_{P^{\zeta, \eta}}[\tau_i] \leq c$  for all  $n \in \mathbb{N}$ ,  $\eta \in \Delta$  and  $\mathbb{P}$ -a.e.  $\zeta$ , and thus

$$(4.35) \quad c^{-1} \sqrt{n} \leq \sigma_n^\zeta(v) \leq c \sqrt{n} \quad \text{for all } n \in \mathbb{N}, v \in V \text{ and } \mathbb{P}\text{-a.e. } \zeta.$$

PROPOSITION 4.10. *Let  $V$  be as in (4.21), and let  $K$  from (4.33) be a large enough fixed constant. Then there exists a constant  $C < \infty$  such that*

$$(4.36) \quad Y_v^{\approx}(n) \sigma_n^\zeta(v) \exp\{nL^*(1/v) + \sigma_v \sqrt{n} W_n(1)\} \in [C^{-1}, C]$$

for all  $v \in V$ ,  $n \in \mathbb{N}$  on  $\mathcal{H}_n$ , where  $W_n$  is given in (4.26) of Proposition 4.6 and  $\sigma_v \in (0, \infty)$  is as in (4.25). In addition, for some  $\tilde{C} < \infty$ ,

$$(4.37) \quad \frac{Y_v^{\approx}(n)}{Y_v^{<}(n)} \in [\tilde{C}^{-1}, \tilde{C}] \quad \text{for all } v \in V, n \in \mathbb{N}, \text{ on } \mathcal{H}_n.$$

In particular, each of the three sequences of processes

$$(4.38) \quad \begin{aligned} t &\mapsto \frac{1}{\sigma_v \sqrt{n}} (\ln Y_v^{\approx}(tn) + tnL^*(1/v)), \quad n \in \mathbb{N}, \\ t &\mapsto \frac{1}{\sigma_v \sqrt{n}} (\ln Y_v^{<}(tn) + tnL^*(1/v)), \quad n \in \mathbb{N}, \\ t &\mapsto \frac{1}{\sigma_v \sqrt{n}} (\ln Y_v(tn) + tnL^*(1/v)), \quad n \in \mathbb{N}, \end{aligned}$$

converges as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to standard Brownian motion.

PROOF. Throughout the proof, we assume that  $n$  is large enough so that  $\mathcal{H}_n$  occurs. To simplify the notation, we also omit the dependence of  $\bar{\eta}_n^\zeta$  and  $\sigma_n^\zeta$  on the parameter  $v$ .

Let  $\widehat{\tau}_i := \tau_i - E^{\zeta, \bar{\eta}_n^\zeta}[\tau_i]$ . Using the definition of the tilted measure  $P^{\zeta, \eta}$  (see (4.4) and below) with (4.8), and the fact that  $\sum_{i=1}^n E^{\zeta, \bar{\eta}_n^\zeta}[\tau_i] = E^{\zeta, \bar{\eta}_n^\zeta}[H_n] = n/v$ , we can rewrite  $Y_v^{\approx}(n)$  as

$$\begin{aligned}
 (4.39) \quad Y_v^{\approx}(n) &= E^{\zeta, \bar{\eta}_n^\zeta} \left[ \exp \left\{ -\bar{\eta}_n^\zeta \sum_{i=1}^n \widehat{\tau}_i \right\}; \sum_{i=1}^n \tau_i \in \left[ \frac{n}{v} - K, \frac{n}{v} \right] \right] \\
 &\quad \times e^{-n(v^{-1}\bar{\eta}_n^\zeta - \bar{L}_n^\zeta(\bar{\eta}_n^\zeta))} \\
 &= E^{\zeta, \bar{\eta}_n^\zeta} \left[ \exp \left\{ -\sigma_n^\zeta \frac{\bar{\eta}_n^\zeta}{\sigma_n^\zeta} \sum_{i=1}^n \widehat{\tau}_i \right\}; \frac{\bar{\eta}_n^\zeta}{\sigma_n^\zeta} \sum_{i=1}^n \widehat{\tau}_i \in \left[ 0, -\frac{K\bar{\eta}_n^\zeta}{\sigma_n^\zeta} \right] \right] \\
 &\quad \times e^{-n(\bar{L}_n^\zeta)^*(1/v)}.
 \end{aligned}$$

Writing  $\mu_n^\zeta$  for the distribution of  $\frac{\bar{\eta}_n^\zeta}{\sigma_n^\zeta} \sum_{i=1}^n \widehat{\tau}_i$  under  $P^{\zeta, \bar{\eta}_n^\zeta}$  (depending implicitly on  $v$ ), we obtain

$$(4.40) \quad Y_v^{\approx}(n) = e^{-n(\bar{L}_n^\zeta)^*(1/v)} \int_0^{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta} e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x)$$

and, in a similar vein,

$$(4.41) \quad Y_v^<(n) = e^{-n(\bar{L}_n^\zeta)^*(1/v)} \int_{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta}^\infty e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x).$$

The first factor in (4.40), (4.41) can be controlled by Proposition 4.6 and Corollary 4.9. The following lemma gives estimates for the second factors.

LEMMA 4.11. *Let  $V$  and  $K$  be as in Proposition 4.10. Then there exists  $C \in (1, \infty)$  such that on  $\mathcal{H}_n$ , for all  $v \in V$ ,*

$$(4.42) \quad \sigma_n^\zeta \int_0^{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta} e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x) \in [C^{-1}, C],$$

$$(4.43) \quad \sigma_n^\zeta \int_{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta}^\infty e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x) \in [C^{-1}, C].$$

In order not to hinder the flow of reading, we complete the proof of Proposition 4.10 first. Using Lemma 4.11, (4.40) and recalling the definition (4.26) of  $W_n$  directly yields (4.36). From (4.41), (4.40) and Lemma 4.11 we deduce (4.37). Finally, replacing  $n$  by  $nt$  in (4.36), observing that  $\sqrt{t}W_{nt}(1) = W_n(t)$ , and using (4.37), the fact that  $\mathcal{H}_n$  occurs  $\mathbb{P}$ -a.s. for  $n$  large, in combination with Proposition 4.6, yields the convergence of the three sequences in (4.38) to standard Brownian motion.  $\square$

We now show Lemma 4.11 which was used in the last proof.

PROOF OF LEMMA 4.11. We start with proving (4.42). Throughout the proof, we assume that  $\mathcal{H}_n$  occurs. Observe that the  $\widehat{\tau}_i$ ,  $1 \leq i \leq n$ , are independent under  $P^{\zeta, \bar{\eta}_n^\zeta}$  and have small exponential moments uniformly in  $n$  (cf. (4.9)). Moreover, recalling (4.34), the variance of the distribution  $\mu_n^\zeta$  is one by definition. A local central limit theorem for such independent normalized sequences, Theorem 13.3 (or formula (13.43)) of [6], thus yields

$$(4.44) \quad \sup_A |\mu_n^\zeta(A) - \Phi(A)| \leq Cn^{-1/2},$$

where the supremum runs over all intervals in  $\mathbb{R}$ , and  $\Phi$  denotes the standard Gaussian measure. Applying (4.44) to  $A = [0, -K\bar{\eta}_n^\zeta/\sigma_n^\zeta]$  and bearing in mind (4.35), this implies that for all  $K$  large enough, uniformly in  $v \in V$ ,

$$c^{-1}n^{-1/2} < \mu_n^\zeta([0, -K\bar{\eta}_n^\zeta/\sigma_n^\zeta]) < cn^{-1/2}.$$

Since the function  $e^{-\sigma_n^\zeta x}$  is uniformly bounded from above and below in this interval, (4.42) follows by another application of (4.35).

In order to show (4.43), we observe that uniformly in  $v \in V$ ,

$$\sigma_n^\zeta \int_{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta}^\infty e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x) \geq \sigma_n^\zeta \int_{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta}^{-2K\bar{\eta}_n^\zeta/\sigma_n^\zeta} e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x) \geq C^{-1}$$

by the same arguments as in the proof of (4.42). On the other hand, using (4.35) and (4.44) again, writing  $I_j = [-jK\bar{\eta}_n^\zeta/\sigma_n^\zeta, -(j+1)K\bar{\eta}_n^\zeta/\sigma_n^\zeta]$ ,

$$\begin{aligned} \sigma_n^\zeta \int_{-K\bar{\eta}_n^\zeta/\sigma_n^\zeta}^\infty e^{-\sigma_n^\zeta x} d\mu_n^\zeta(x) &\leq \sigma_n^\zeta \sum_{j=1}^\infty \mu_n^\zeta(I_j) e^{-jK|\bar{\eta}_n^\zeta|} \\ (4.45) \qquad \qquad \qquad &\leq c\sigma_n^\zeta \sum_{j=1}^\infty n^{-1/2} e^{-jK|\bar{\eta}_n^\zeta|} \leq C, \end{aligned}$$

uniformly in  $v \in V$ . This completes the proof of the lemma.  $\square$

REMARK 4.12. The arguments of the last proof can be used to show that for arbitrary  $a \in [0, n/v]$ ,  $n \in \mathbb{N}$ ,  $v \in V$  on  $\mathcal{H}_n$ ,

$$(4.46) \qquad \frac{1}{Y_v^\approx(n)} E_0 \left[ e^{\int_0^{H_n} \zeta(X_s) ds}; H_n \leq \frac{n}{v} - a \right] \leq C e^{-ca}.$$

Indeed, the expectation on the left-hand side of (4.46) can be written as in (4.41) with  $K$  replaced by  $a$ . Hence, with help of (4.42), the left-hand side of (4.46) is bounded by the left-hand side of (4.45) with  $K$  replaced by  $a$ . Recalling the last-but-one expression in (4.45), inequality (4.46) easily follows.

REMARK 4.13. The proof of Proposition 4.10 is the only occasion where the random tilting by  $\bar{\eta}_n^\zeta(v)$  is really necessary. The reason for this is the application of (4.44), the local central limit theorem in spirit, which is useful only for events of sufficiently large probability. Deterministic tilting by  $\bar{\eta}(v)$ , which would simplify the remaining parts of the paper, unfortunately requires dealing with events of much smaller probability.

4.4. *The walk lingers in the bulk.* We now show that the invariance principles of Proposition 4.10 are useful in order to analyze the Feynman–Kac representation (3.3) of  $\mathbb{E}_{u_0}^\xi [N^\geq(t, vt)]$ . We explore the fact that, under the considered distributions, conditioning on  $X_{n/v} = n$  (as in the Feynman–Kac representation) implies that with high probability  $H_n$  is close to  $n/v$ , that is the “walk lingers in the bulk.”

LEMMA 4.14. *Let  $K > 0$  and  $V$  be as in Proposition 4.10. Then there exists a constant  $c > 0$  such that for all  $n \in \mathbb{N}$  and  $v \in V$ , on  $\mathcal{H}_n$ ,*

$$\begin{aligned} (4.47) \qquad cY_v^\approx(n) &\leq E_0 \left[ \exp \left\{ \int_0^{n/v} \zeta(X_s) ds \right\}; X_{n/v} = n \right] \\ &\leq E_0 \left[ \exp \left\{ \int_0^{n/v} \zeta(X_s) ds \right\}; X_{n/v} \geq n \right] \leq c^{-1} Y_v^\approx(n). \end{aligned}$$

In particular,

$$(4.48) \quad ce^{esn/v} Y_v^\approx(n) \leq \mathbb{E}_0^\xi \left[ N\left(\frac{n}{v}, n\right) \right] \leq \mathbb{E}_0^\xi \left[ N^{\geq}\left(\frac{n}{v}, n\right) \right] \leq c^{-1} e^{esn/v} Y_v^\approx(n).$$

PROOF. The second claim of the lemma follows directly from the first one; it suffices to recall  $\xi(x) = \zeta(x) + es$  and (3.3).

To prove the first claim, we define

$$p_n^\zeta(s) := E_n \left[ \exp \left\{ \int_0^s \zeta(X_r) dr \right\}; X_s = n \right], \quad n \in \mathbb{Z}, s \geq 0,$$

and set  $t = n/v$ , to simplify notation. Using the strong Markov property,

$$\begin{aligned} & E_0 \left[ \exp \left\{ \int_0^t \zeta(X_s) ds \right\}; X_t = n \right] \\ &= E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\} p_n^\zeta(t - H_n); H_n \leq t \right] \\ &\geq E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\}; H_n \in [t - K, t] \right] \inf_{s \leq K} p_n^\zeta(s). \end{aligned}$$

Since the  $\zeta(x)$ 's are bounded from below by assumption (POT), the infimum on the right-hand side can be bounded from below by a deterministic constant  $c = c(K) > 0$ , implying the first inequality in (4.47).

The second inequality of (4.47) is obvious. For the third one, observe that  $\{X_t \geq n\} \subset \{H_n \leq t\}$ . Therefore, decomposing the integral according to the value of  $H_n$  and using the fact that  $\zeta \leq 0$ , we obtain

$$\begin{aligned} & E_0 \left[ \exp \left\{ \int_0^t \zeta(X_s) ds \right\}; X_t \geq n \right] \\ &= E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\} \exp \left\{ \int_{H_n}^t \zeta(X_s) ds \right\}; X_t \geq n \right] \\ &\leq E_0 \left[ \exp \left\{ \int_0^{H_n} \zeta(X_s) ds \right\}; H_n \leq t \right] = Y_v(n). \end{aligned}$$

By Proposition 4.10,  $Y_v(n)$  and  $Y_v^\approx(n)$  are comparable on  $\mathcal{H}_n$ , which proves the third inequality.  $\square$

4.5. *Initial condition stability.* The next lemma shows that initial conditions  $u_0$  satisfying assumption (INI) are comparable to the “one-particle” initial condition  $u_0 = \mathbf{1}_{\{0\}}$ .

LEMMA 4.15. *Let  $V$  be as in (4.21). There exists a finite constant  $C$  such that for all  $u_0$  as in (INI) and for all  $n \in \mathbb{N}$ ,  $t \geq 0$  such that  $n/t \in V$ , on  $\mathcal{H}_n$ ,*

$$(4.49) \quad 1 \leq \frac{\mathbb{E}_{u_0}^\xi [N^{\geq}(t, n)]}{\mathbb{E}_0^\xi [N(t, n)]} \leq C.$$

PROOF. The first inequality in (4.49) is obvious, so we proceed to the second one. Moreover, since  $\mathbb{E}_{u_0}^\xi [N^{\geq}(t, n)]$  is an increasing function of  $u_0(x)$  for every  $x \in -\mathbb{N}_0$ , we can assume that  $u_0 = c\mathbf{1}_{-\mathbb{N}_0}$ . Using the Feynman–Kac representation (3.3), and replacing  $\xi$  by  $\zeta$ , we see that

$$(4.50) \quad \frac{\mathbb{E}_{c\mathbf{1}_{-\mathbb{N}_0}}^\xi [N^{\geq}(t, n)]}{\mathbb{E}_0^\xi [N(t, n)]} = \frac{c \sum_{x \leq 0} E_x [\exp \{ \int_0^t \zeta(X_s) ds \}; X_t \geq n]}{E_0 [\exp \{ \int_0^t \zeta(X_s) ds \}; X_t = n]}.$$

Applying the strong Markov property on the numerator of the right-hand side, we obtain

$$\begin{aligned} & \sum_{x \leq 0} E_x \left[ e^{\int_0^t \zeta(X_s) ds}; X_t \geq n \right] \\ & \leq \sum_{x \leq 0} E_x \left[ e^{\int_0^{H_n} \zeta(X_s) ds}; H_n \leq t \right] \\ & = \sum_{x \leq 0} \int_0^t E_x \left[ e^{\int_0^{H_0} \zeta(X_s) ds}; H_0 \in da \right] E_0 \left[ e^{\int_0^{H_n} \zeta(X_s) ds}; H_n \leq t - a \right]. \end{aligned}$$

By (4.46), on  $\mathcal{H}_n$ , the second factor on the right-hand side can be bounded from above by  $Ce^{-ca}Y_{n/t}^{\approx}(n)$  for all  $n$  with  $n/t \in V$  and  $a \in [0, t]$ . This implies that the right-hand side of the last display is bounded from above by

$$CY_{n/t}^{\approx}(n) \sum_{x \leq 0} \int_0^t P_x(H_0 \in da) e^{-ca} \leq CY_{n/t}^{\approx}(n),$$

where for the last inequality we used that due to the stationarity of simple random walk we have  $\sum_{x \leq 0} P_x(H_0 \in da) = \sum_{x \geq 0} P_0(H_x \in da)$ , and the latter is the probability that an arbitrary point  $x \geq 0$  is visited for the first time at time  $da$ , so it is bounded by  $da$ . By Lemma 4.14, on  $\mathcal{H}_n$ ,  $Y_{n/t}^{\approx}(n)$  is comparable to the denominator of the right-hand side in (4.50), which completes the proof.  $\square$

4.6. *Proof of Theorem 2.8 (functional CLT for the PAM).* We have all ingredients to show our first main result, the invariance principle for the PAM, Theorem 2.8.

PROOF OF THEOREM 2.8. Recall that we have to show that the sequence of processes  $(\ln u(nt, \lfloor vnt \rfloor) - nt\lambda(v))/(\sigma_v\sqrt{vn})$ , with  $u(t, x) = \mathbb{E}_{u_0}^{\xi}[N(t, x)]$  as in (2.10), satisfies the functional central limit theorem under  $\mathbb{P}$  as  $n \rightarrow \infty$ .

By Lemma 4.15, we can assume without loss of generality that  $u_0 = \mathbf{1}_{\{0\}}$ . Moreover, by Lemma 4.14,  $\mathbb{P}$ -a.s. for all large  $t$ ,

$$(4.51) \quad cY_v^{\approx}(vt)e^{tes} \leq u(t, \lfloor vt \rfloor) \leq c^{-1}Y_v^{\approx}(vt)e^{tes}.$$

Replacing  $t$  by  $vt$  in Proposition 4.10, we see that

$$t \mapsto \frac{1}{\sigma_v\sqrt{nv}} (\ln Y_v^{\approx}(tvn) + tvnL^*(1/v))$$

converges as  $n \rightarrow \infty$  to standard Brownian motion. Combining this with (4.51) easily implies the theorem; incidentally, it also shows that the Lyapunov exponent  $\lambda(v)$  defined in (2.2) satisfies  $\lambda(v) = es - vL^*(1/v)$  for  $v > v_c$ , as claimed in (A.9) of Proposition A.3. This completes the proof of Theorem 2.8.  $\square$

REMARK 4.16. Theorem 2.8 remains valid when the function  $u(t, x)$  is replaced by  $\mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, x)]$  with  $u_0$  as in (INI). This is a consequence of Lemma 4.15 again.

REMARK 4.17. It will be useful to have a more explicit formula for  $\ln \mathbb{E}_0^{\xi}[N^{\geq}(n/v_0, n)]$ . Combining (4.48) with Corollary 4.9, Proposition 4.10 and (A.9) yields the existence of a constant  $C < \infty$  and a  $\mathbb{P}$ -a.s. finite random variable  $\mathcal{N}_3$  such that  $\mathbb{P}$ -a.s. for all  $n \geq \mathcal{N}_3$ ,

$$\left| \ln \mathbb{E}_0^{\xi}[N^{\geq}(n/v_0, n)] - \sum_{i=1}^n L_i^{\zeta}(\bar{\eta}(v_0)) + nL(\bar{\eta}(v_0)) \right| \leq C \ln n.$$



**5. Breakpoint behavior.** The goal of this section is to prove the functional central limit theorem for the breakpoint, Theorem 2.6. This is done in Section 5.2 after some additional preparations.

5.1. *Perturbation estimates.* The results of Section 4 provide a reasonably precise description of the behavior of expectations of  $N^{\geq}(t, vt)$ . We are now interested in how sensitive the expectation of  $N^{\geq}(\cdot, \cdot)$  is to perturbations in the space and time coordinate. The first lemma deals with space perturbations.

LEMMA 5.1. (a) *Let  $\tilde{\varepsilon}(t)$  be a positive function with  $\lim_{t \rightarrow \infty} \tilde{\varepsilon}(t)t^{\delta} = 0$  for some  $\delta > 0$ . Then for each  $\varepsilon > 0$  there exists a constant  $C_0 = C_0(\varepsilon) < \infty$  such that  $\mathbb{P}$ -a.s.,*

$$(5.1) \quad \limsup_{t \rightarrow \infty} \sup \left\{ \left| \frac{1}{h} \ln \frac{\mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, vt + h)]}{\mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, vt)]} - L(\bar{\eta}(v)) \right| : (h, v) \in \mathcal{E}_t \right\} \leq \varepsilon,$$

where  $\mathcal{E}_t = \{(h, v) : C_0 \ln t \leq |h| \leq t\tilde{\varepsilon}(t), v \in V, v + \frac{h}{t} \in V\}$ .

(b) *There exist constants  $C, c \in (0, \infty)$  and a  $\mathbb{P}$ -a.s. finite random variable  $\mathcal{T}_1$  such that  $\mathbb{P}$ -a.s. for all  $t \geq \mathcal{T}_1$ , uniformly for  $0 \leq h \leq t^{1/3}$  and  $v, v + h/t \in V$ ,*

$$ce^{-Ch} \mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, vt)] \leq \mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, vt + h)] \leq Ce^{-ch} \mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, vt)].$$

PROOF. (a) We set  $v' := v + \frac{h}{t}$ . Without loss of generality, we can assume  $t$  to be large enough so that the events  $\mathcal{H}_{vt}$  and  $\mathcal{H}_{v't}$  occur, and thus  $\bar{\eta}_{vt}^{\zeta}(v)$  and  $\bar{\eta}_{v't}^{\zeta}(v')$  exist and satisfy corresponding versions of (4.11). By Lemmas 4.14 and 4.15, the fraction in (5.1) can be approximated, up to a multiplicative constant that is irrelevant in the limit, by

$$\frac{Y_{v'}(v't)}{Y_v(vt)} = \frac{E_0[\exp\{\int_0^{H_{v't}} \zeta(X_s) ds\}; H_{v't} \leq t]}{E_0[\exp\{\int_0^{H_{vt}} \zeta(X_s) ds\}; H_{vt} \leq t]}.$$

Using the notation from (4.28), this can be rewritten in the same vein as in (4.39) as

$$(5.2) \quad \frac{E^{\zeta, \bar{\eta}_{v't}^{\zeta}(v')} [e^{-\bar{\eta}_{v't}^{\zeta}(v') \sum_{i=1}^{v't} \tilde{\tau}_i}; \sum_{i=1}^{v't} \tilde{\tau}_i \in (-\infty, 0]] \cdot e^{-S_{v't}^{\zeta, v'}(\bar{\eta}_{v't}^{\zeta}(v'))}}{E^{\zeta, \bar{\eta}_{vt}^{\zeta}(v)} [e^{-\bar{\eta}_{vt}^{\zeta}(v) \sum_{i=1}^{vt} \hat{\tau}_i}; \sum_{i=1}^{vt} \hat{\tau}_i \in (-\infty, 0]] \cdot e^{-S_{vt}^{\zeta, v}(\bar{\eta}_{vt}^{\zeta}(v))}},$$

where, similarly as before  $\hat{\tau}_i := \tau_i - E^{\zeta, \bar{\eta}_{vt}^{\zeta}(v)}[\tau_i]$  and  $\tilde{\tau}_i := \tau_i - E^{\zeta, \bar{\eta}_{v't}^{\zeta}(v')}[\tau_i]$ . By the same methods as in (4.40)–(4.43), the expectations in the numerator and denominator of (5.2) are both of order  $t^{-1/2}$ . Their ratio is thus bounded from above and below by positive finite constants and can be neglected in the limit taken in (5.1).

The remaining terms in (5.2) contribute to the minuend of (5.1) as

$$(5.3) \quad \frac{1}{h} (S_{vt}^{\zeta, v}(\bar{\eta}_{vt}^{\zeta}(v)) - S_{vt}^{\zeta, v}(\bar{\eta}_{vt}^{\zeta}(v'))) + \frac{1}{h} (S_{v't}^{\zeta, v}(\bar{\eta}_{v't}^{\zeta}(v')) - S_{v't}^{\zeta, v'}(\bar{\eta}_{v't}^{\zeta}(v'))).$$

In order to show that the first summand on the right-hand side of (5.3) is negligible uniformly as  $t \rightarrow \infty$ , we write

$$(5.4) \quad \begin{aligned} S_{vt}^{\zeta, v}(\bar{\eta}_{vt}^{\zeta}(v')) &= S_{vt}^{\zeta, v}(\bar{\eta}_{vt}^{\zeta}(v)) + (S_{vt}^{\zeta, v})'(\bar{\eta}_{vt}^{\zeta}(v))(\bar{\eta}_{vt}^{\zeta}(v') - \bar{\eta}_{vt}^{\zeta}(v)) \\ &\quad + (S_{vt}^{\zeta, v})''(\tilde{\eta})(\bar{\eta}_{vt}^{\zeta}(v') - \bar{\eta}_{vt}^{\zeta}(v))^2, \end{aligned}$$

for some  $\tilde{\eta} \in \Delta$  with  $|\tilde{\eta} - \bar{\eta}_{vt}^{\zeta}(v)| \leq |\bar{\eta}_{v't}^{\zeta}(v') - \bar{\eta}_{vt}^{\zeta}(v)|$ . As observed below (4.31), one has  $(S_{vt}^{\zeta, v})'(\bar{\eta}_{vt}^{\zeta}(v)) = 0$ , so the second term vanishes. For the third one, note that by Lemma 4.5,  $\mathbb{P}$ -a.s. for all  $t$  large enough,

$$(5.5) \quad |\bar{\eta}_{v't}^{\zeta}(v') - \bar{\eta}_{vt}^{\zeta}(v)| \leq \frac{Ch}{t}.$$

Moreover, by the characterizing property (4.16) of  $\bar{\eta}_{vt}^\xi(v)$ , Lemma A.1 and the implicit function theorem, we see that  $v \mapsto \bar{\eta}_{vt}^\xi(v)$  is differentiable on the interior of  $V$ , with uniformly bounded derivative. Therefore, on  $\mathcal{H}_{vt}$ , uniformly for  $v \in V$  and  $h$  as in (5.1),

$$(5.6) \quad \left| \bar{\eta}_{vt}^\xi(v') - \bar{\eta}_{vt}^\xi(v) \right| \leq \frac{Ch}{t}.$$

Recalling (4.32), we see that,  $\mathbb{P}$ -a.s.,  $(S_{vt}^{\xi,v})''(\tilde{\eta}) \leq Ct$  uniformly in  $v \in V$  and  $t$  large. Combined with (5.4) to (5.6), we thus deduce that  $\mathbb{P}$ -a.s., the first term on the right-hand side of (5.3) satisfies

$$(5.7) \quad \left| \frac{1}{h} (S_{vt}^{\xi,v}(\bar{\eta}_{vt}^\xi(v)) - S_{vt}^{\xi,v}(\bar{\eta}_{v't}^\xi(v'))) \right| \leq \frac{Ch}{t}$$

which is negligible in the limit considered in (5.1).

Plugging in the definitions, the second summand on the right-hand side of (5.3) satisfies

$$(5.8) \quad \frac{1}{h} (S_{vt}^{\xi,v}(\bar{\eta}_{v't}^\xi(v')) - S_{v't}^{\xi,v'}(\bar{\eta}_{v't}^\xi(v'))) = \frac{1}{h} \sum_{i=vt+1}^{v't} L_i^\xi(\bar{\eta}_{v't}^\xi(v'))$$

(where the sum should be interpreted as  $-\sum_{i=v't+1}^{vt}$  if  $v' < v$ ). The right-hand side of (5.8) can be approximated with the help of the following claim.

CLAIM 5.2. *For each  $\varepsilon > 0$  and each  $q \in \mathbb{N}$ , there exists a constant  $C = C(q, \varepsilon) < \infty$  such that for all  $t$  large enough,*

$$\mathbb{P} \left( \sup_{\substack{v \in V \\ C(q, \varepsilon) \ln t \leq |h| \leq t\tilde{\varepsilon}(t)}} \left| \frac{1}{h} \sum_{i=vt+1}^{v't} L_i^\xi(\bar{\eta}_{v't}^\xi(v')) - L(\bar{\eta}(v)) \right| > \varepsilon, \mathcal{H}_{v't}, \mathcal{H}_{vt} \right) \leq Ct^{-q}$$

with  $\tilde{\varepsilon}(t)$  and  $v'$  as in Lemma 5.1.

We postpone the proof of this claim after the proof of Lemma 5.1. Inequality (5.7) and Claim 5.2 together imply that the left-hand side of (5.3)  $\mathbb{P}$ -a.s. satisfies

$$\limsup_{t \rightarrow \infty} \sup_{(t,v)} \left\{ \left| \frac{1}{h} (S_{vt}^{\xi,v}(\bar{\eta}_{vt}^\xi(v)) - S_{v't}^{\xi,v'}(\bar{\eta}_{v't}^\xi(v'))) - L(\bar{\eta}(v)) \right| \right\} \leq \varepsilon,$$

where the supremum is taken over all  $(t, v)$  satisfying  $C(2, \varepsilon) \ln t \leq |h| \leq t\tilde{\varepsilon}(t)$  and  $v \in V$ . This is what is necessary to prove Lemma 5.1(a).

(b) Using the same arguments as in the proof of (a), it is sufficient show that the exponential factors in (5.2) are bounded from above and below by exponential functions, that is, the right-hand side of (5.3) is bounded away from 0 and  $\infty$ . However, for the second summand on the right-hand side this easily follows from (5.8), because  $c^{-1} < L_i^\xi(\bar{\eta}_{v't}^\xi(v')) < c < 0$  uniformly in  $i \geq 0$ ,  $v \in V$ , and  $\xi$  satisfying (POT). The first summand can be neglected for  $t$  sufficiently large due to (5.7).  $\square$

PROOF OF CLAIM 5.2. We rewrite

$$(5.9) \quad \begin{aligned} & \sum_{i=vt+1}^{v't} L_i^\xi(\bar{\eta}_{v't}^\xi(v')) - L(\bar{\eta}(v)) \\ &= \sum_{i=vt+1}^{v't} (L_i^\xi(\bar{\eta}_{v't}^\xi(v')) - L_i^\xi(\bar{\eta}(v))) + \sum_{i=vt+1}^{v't} (L_i^\xi(\bar{\eta}(v)) - L(\bar{\eta}(v))). \end{aligned}$$

Observing that the family of functions  $(\eta \mapsto L_i^\zeta(\eta))_{i \in \mathbb{Z}, -\text{es} \leq \zeta(j) \leq 0 \forall j \in \mathbb{Z}}$  is equicontinuous on  $\Delta$ , Proposition 4.3, (5.5) and (5.6) yield that

$$(5.10) \quad \mathbb{P} \left( \sup_{\substack{v \in V \\ \ln t \leq |h| \leq t\tilde{\varepsilon}(t)}} \left| \frac{1}{h} \sum_{i=vt+1}^{v't} (L_i^\zeta(\tilde{\eta}_{v't}^\zeta(v')) - L_i^\zeta(\tilde{\eta}(v))) \right| \geq \frac{\varepsilon}{2}, \mathcal{H}_{v't}, \mathcal{H}_{vt} \right) \leq Ct^{-q}.$$

Regarding the second summand on the right-hand side of (5.9), it suffices to observe that for  $C(q, \varepsilon)$  large enough,

$$(5.11) \quad \mathbb{P} \left( \sup_{C(q, \varepsilon) \ln t \leq |h| \leq t\tilde{\varepsilon}(t)} \left| \frac{1}{h} \sum_{i=vt+1}^{v't} L_i^\zeta(x) - L(x) \right| \geq \varepsilon/2 \right) \leq Ct^{-q},$$

which follows from the Hoeffding-type bound (Lemma A.5) using the same steps as in the proof of Claim 4.4. Combining (5.9)–(5.11) with (4.22) completes the proof of the claim.  $\square$

We now deal with time perturbations, where it is possible and useful to obtain more precise estimates.

LEMMA 5.3. (a) *Let  $\varepsilon(t)$  be a function such that  $\lim_{t \rightarrow \infty} \varepsilon(t) = 0$ . Then there exists a constant  $C \in (0, \infty)$  and a  $\mathbb{P}$ -a.s. finite random variable  $\mathcal{T}_2$  such that  $\mathbb{P}$ -a.s. for all  $t \geq \mathcal{T}_2$ ,*

$$(5.12) \quad \sup_{(h, v) \in \mathcal{E}_t} \left| \ln \frac{\mathbb{E}_{u_0}^\xi [N^\geq(t+h, vt)]}{\mathbb{E}_{u_0}^\xi [N^\geq(t, vt)]} - h(\text{es} - \bar{\eta}(v)) \right| \leq C + C|h| \left( \sqrt{\frac{\ln t}{t}} + \frac{|h|}{t} \right),$$

where  $\mathcal{E}_t = \{(h, v) : |h| \leq t\varepsilon(t), v \in V, vt/(t+h) \in V\}$ .

(b) *In particular, there exist constants  $C, c \in (0, \infty)$  such that for  $\mathcal{T}_2$  as in (a) the following holds true:  $\mathbb{P}$ -a.s. for all  $t \geq \mathcal{T}_2$ , uniformly in  $0 \leq h \leq t\varepsilon(t)$  and  $v, vt/(t+h) \in V$ ,*

$$ce^{ch} \mathbb{E}_{u_0}^\xi [N^\geq(t, vt)] \leq \mathbb{E}_{u_0}^\xi [N^\geq(t+h, vt)] \leq Ce^{Ch} \mathbb{E}_{u_0}^\xi [N^\geq(t, vt)].$$

REMARK 5.4. By interchanging the roles of  $vt$  and  $vt+h$  as well as of  $t$  and  $t+h$  in the claims (b) of Lemmas 5.1 and 5.3, respectively, it follows that they hold also for  $h \in [-t^{\frac{1}{3}}, 0]$  and  $h \in [-t\varepsilon(t), 0]$ , respectively, with minimal modifications: in Lemma 5.3, the prefactors  $ce^{ch}$  and  $Ce^{Ch}$  should be replaced by  $ce^{Ch}$  and  $Ce^{ch}$ , respectively; a similar replacement applies also in Lemma 5.1.

PROOF OF LEMMA 5.3. (a) Let  $v' := vt/(t+h)$ . Through the proof, we assume  $t$  to be large enough such that  $\mathcal{H}_{vt}$  and  $\mathcal{H}_{v't}$  hold true. Using Proposition 4.10 and the same arguments as in the proof of Lemma 5.1, the fraction in (5.12) satisfies, for some  $c \in (1, \infty)$ ,

$$c^{-1} e^{hes} \cdot \frac{Y_{v'}^\approx(vt)}{Y_v^\approx(vt)} \leq \frac{\mathbb{E}_{u_0}^\xi [N^\geq(t+h, vt)]}{\mathbb{E}_{u_0}^\xi [N^\geq(t, vt)]} \leq ce^{hes} \cdot \frac{Y_{v'}^\approx(vt)}{Y_v^\approx(vt)}.$$

In addition, similar to (4.39),

$$\frac{Y_{v'}^\approx(vt)}{Y_v^\approx(vt)} = e^{hes} \cdot \frac{E^{\zeta, \tilde{\eta}_{v't}^\zeta(v')} [e^{-\tilde{\eta}_{v't}^\zeta(v') \sum_{i=1}^{v't} \tilde{\tau}_i}; \sum_{i=1}^{v't} \tilde{\tau}_i \in [-K, 0]] \cdot e^{-S_{v't}^{\zeta, v'}(\tilde{\eta}_{v't}^\zeta(v'))}}{E^{\zeta, \tilde{\eta}_{vt}^\zeta(v)} [e^{-\tilde{\eta}_{vt}^\zeta(v) \sum_{i=1}^{vt} \hat{\tau}_i}; \sum_{i=1}^{vt} \hat{\tau}_i \in [-K, 0]] \cdot e^{-S_{vt}^{\zeta, v}(\tilde{\eta}_{vt}^\zeta(v))}},$$

where again  $\hat{\tau}_i := \tau_i - E^{\zeta, \tilde{\eta}_{vt}^\zeta(v)}[\tau_i]$  and  $\tilde{\tau}_i := \tau_i - E^{\zeta, \tilde{\eta}_{v't}^\zeta(v')}[\tau_i]$ . As in the proof of Lemma 5.1, the ratio of the expectations in the numerator and denominator is asymptotically bounded

from above and below. It follows that the expression in the supremum of (5.12) is bounded from above by

$$\begin{aligned}
 (5.13) \quad & C + |h\bar{\eta}(v) + S_{vt}^{\zeta,v}(\bar{\eta}_{vt}^{\zeta}(v)) - S_{vt}^{\zeta,v'}(\bar{\eta}_{vt}^{\zeta}(v'))| \\
 & \leq C + |h\bar{\eta}(v) + (S_{vt}^{\zeta,v}(\bar{\eta}_{vt}^{\zeta}(v')) - S_{vt}^{\zeta,v'}(\bar{\eta}_{vt}^{\zeta}(v')))| \\
 & \quad + |(S_{vt}^{\zeta,v}(\bar{\eta}_{vt}^{\zeta}(v)) - S_{vt}^{\zeta,v}(\bar{\eta}_{vt}^{\zeta}(v')))|.
 \end{aligned}$$

Plugging in the definitions (4.28) and (4.24), the second summand on the right-hand side satisfies

$$\begin{aligned}
 |h\bar{\eta}(v) + (S_{vt}^{\zeta,v}(\bar{\eta}_{vt}^{\zeta}(v')) - S_{vt}^{\zeta,v'}(\bar{\eta}_{vt}^{\zeta}(v')))| &= \left| h\bar{\eta}(v) + vt\bar{\eta}_{vt}^{\zeta}(v')\left(\frac{1}{v} - \frac{1}{v'}\right) \right| \\
 &= |h|\left|\bar{\eta}(v) - \bar{\eta}_{vt}^{\zeta}(v')\right|.
 \end{aligned}$$

Using (5.6) and (4.15) of Proposition 4.3 in combination with the Borel–Cantelli lemma, this can then be bounded by the right-hand side of (5.12). The last summand on the right-hand side of (5.13) can be shown to be smaller than  $Ch^2/t$  using the same steps as in (5.4)–(5.7) of the proof of Lemma 5.1, completing the proof of (a). Claim (b) directly follows from (a).  $\square$

5.2. *Proof of Theorem 2.6 (functional CLT for the breakpoint).* We now have all the ingredients to show our second main result, the invariance principle for the breakpoint, Theorem 2.6.

PROOF OF THEOREM 2.6. We must show that the sequence of processes  $\frac{1}{\sigma_v\sqrt{n}}(\bar{m}_v(nt) - vnt)$  converges to standard Brownian motion, where

$$\bar{m}_v(t) = \sup \left\{ n \in \mathbb{N} : \mathbb{E}_{u_0}^{\xi}[N^{\geq}(t, n)] \geq \frac{1}{2}e^{t\lambda(v)} \right\}$$

was defined in (2.7).

We assume that  $u_0 = \mathbf{1}_{\{0\}}$  first. Let  $u^{\geq}(t, x) := \mathbb{E}_0^{\xi}[N^{\geq}(t, x)]$ ,  $t \geq 0$ ,  $x \in \mathbb{Z}$ , and extend it to  $x \in \mathbb{R}$  by linear interpolation. Furthermore, set

$$(5.14) \quad U_v(t) := t\lambda(v) - \ln u^{\geq}(t, vt) - \ln 2.$$

Recalling the definition of  $\sigma_v^2$  from (4.25), by Remark 4.16,

$$(5.15) \quad \left( t \mapsto \frac{U_v(nt)}{\sqrt{\sigma_v^2 vn}} \right)_{n \in \mathbb{N}} \text{ converges as } n \rightarrow \infty \text{ to Brownian motion.}$$

Obviously,  $u^{\geq}(t, x)$  is decreasing in  $x$  with  $\lim_{t \rightarrow \infty} \frac{1}{t} \ln u^{\geq}(t, 0) = \lambda(0) > \lambda(v)$  and  $\lim_{x \rightarrow \infty} u^{\geq}(t, x) = 0$ ; see Proposition A.3. Let  $r = r(t)$  be the largest solution of the equation

$$(5.16) \quad u^{\geq}(t, vt + r) = \frac{1}{2}e^{t\lambda(v)},$$

which exists  $\mathbb{P}$ -a.s. for  $t$  large enough by the previous considerations. Moreover, by the definition of  $\bar{m}_v(t)$ ,

$$(5.17) \quad r(t) - 1 < \bar{m}_v(t) - vt \leq r(t).$$

Combining equations (5.14) and (5.16), we see that  $r(t)$  is the largest solution to

$$\ln \frac{u^{\geq}(t, vt + r(t))}{u^{\geq}(t, vt)} = U_v(t).$$

Let  $\tilde{\varepsilon}(t)$  be an arbitrary positive function with  $\tilde{\varepsilon}(t)t^{\frac{1}{4}} \rightarrow 0$  and  $\tilde{\varepsilon}(t)t^{\frac{1}{2}} \rightarrow \infty$  as  $t \rightarrow \infty$ . By the space perturbation Lemma 5.1, using also the monotonicity of  $u^{\geq}(t, \cdot)$  and the fact that  $L(\bar{\eta}(v)) < 0$ , we obtain that for every

$$(5.18) \quad \delta \in (0, |L(\bar{\eta}(v))|),$$

$\mathbb{P}$ -a.s. for all  $t$  large enough,

$$\bar{\varphi}_t(r(t))L(\bar{\eta}(v)) - \delta|r(t)| \leq \ln \frac{u^{\geq}(t, vt + r(t))}{u^{\geq}(t, vt)} \leq \underline{\varphi}_t(r(t))L(\bar{\eta}(v)) + \delta|r(t)|;$$

here, for  $C_0 = C_0(\delta)$ , the functions  $\bar{\varphi}_t$  and  $\underline{\varphi}_t$  are given by

$$\underline{\varphi}_t(r) = \sup\{s : s \leq r \text{ and } C_0 \ln t \leq |s| \leq t\tilde{\varepsilon}(t)\},$$

$$\bar{\varphi}_t(r) = \inf\{s : s \geq r \text{ and } C_0 \ln t \leq |s| \leq t\tilde{\varepsilon}(t)\},$$

and satisfy  $\underline{\varphi}_t(r) = \bar{\varphi}_t(r) = r$  for  $C_0 \ln t \leq |r| \leq t\varepsilon(t)$  and  $\underline{\varphi}_t \leq \bar{\varphi}_t$ . This implies that whenever

$$(5.19) \quad |U_v(t)| \in [C_0(|L(\bar{\eta}(v))| + \delta) \ln t, t\tilde{\varepsilon}(t)(|L(\bar{\eta}(v))| - \delta)],$$

then, due to (5.18),

$$r(t) \in \left[ \frac{U_v(t)}{L(\bar{\eta}(v)) \pm \delta}, \frac{U_v(t)}{L(\bar{\eta}(v)) \mp \delta} \right],$$

where the upper signs correspond to  $U_v(t) > 0$  and the lower signs to  $U_v(t) < 0$ . In particular, since  $U_v$  satisfies the invariance principle (5.15), property (5.19) is satisfied with probability tending to 1 as  $t \rightarrow \infty$ . Since  $\delta$  is arbitrary, it thus follows that in  $\mathbb{P}$ -distribution

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} r(n \cdot) = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \cdot \frac{U_v(n \cdot)}{L(\bar{\eta}(v))}$$

as processes defined on  $[0, \infty)$ , which together with (5.17) and (5.15) implies the claim of the theorem for

$$(5.20) \quad \bar{\sigma}_v = \frac{\sqrt{\sigma_v^2 v}}{|L(\bar{\eta}(v))|}.$$

The case of general  $u_0$  satisfying (INI) then follows from Lemmas 4.15 and 5.1. This completes the proof.  $\square$

**5.3. Invariance principle for the breakpoint inverse.** We will later on need the following invariance principle for a generalized inverse of the breakpoint defined by  $T_0 = 0$  and, for  $n \geq 1$ ,

$$(5.21) \quad T_n := \inf \left\{ t \geq 0 : \mathbb{E}_{u_0}^{\xi} [N^{\geq}(t, n)] \geq \frac{1}{2} \right\} = \inf \{ t \geq 0 : \bar{m}(t) \geq n \}.$$

Observe that by definition

$$(5.22) \quad T_{\bar{m}(t)} \leq t.$$

**THEOREM 5.5.** *There exists a  $\mathbb{P}$ -a.s. finite random variable  $C = C(\xi)$  and a constant  $C_1 < \infty$  such that  $\mathbb{P}$ -a.s. for all  $n \geq 1$ ,*

$$(5.23) \quad \left| T_n - \left( \frac{n}{v_0} + \frac{1}{v_0 L(\bar{\eta}(v_0))} \sum_{i=1}^n (L_i^{\xi}(\bar{\eta}(v_0)) - L(\bar{\eta}(v_0))) \right) \right| \leq C + C_1 \ln n.$$

In particular,

$$(5.24) \quad \lim_{n \rightarrow \infty} \frac{T_n}{n} = \frac{1}{v_0}, \quad \mathbb{P}\text{-a.s.},$$

and, a fortiori, the sequence

$$t \mapsto \frac{v_0 L(\bar{\eta}(v_0))}{\sqrt{\sigma_{v_0} n}} \left( T_{nt} - \frac{nt}{v_0} \right), \quad n \geq 0,$$

converges as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to standard Brownian motion.

PROOF. To show (5.23), we set

$$h_n = \frac{1}{v_0 L(\bar{\eta}(v_0))} \sum_{i=1}^n (L_i^\zeta(\bar{\eta}(v_0)) - L(\bar{\eta}(v_0))).$$

Observe that  $\mathbb{P}$ -a.s. for all  $n$  large enough

$$(5.25) \quad |h_n| \leq C \sqrt{n \ln \ln n}.$$

Indeed, the random variables  $L_i^\zeta(\bar{\eta}(v_0)) - L(\bar{\eta}(v_0))$  are centered and mixing as in (A.5). We can thus apply Azuma's inequality for mixing sequences, Lemma A.5, which can be turned into a maximal inequality using [41], Theorem 1, to deduce that for all  $a \geq 0$ ,

$$\mathbb{P}\left(\max_{k \leq n} |h_k| \geq a\right) \leq C e^{-ca^2/n}.$$

The usual steps of the proof of the upper bound in the classical law of the iterated logarithm then provide us with (5.25).

We now fix  $\alpha \in \mathbb{R}$  and estimate  $\ln \mathbb{E}_{u_0}^\xi [N^{\geq}(n/v_0 + h + \alpha \ln n, n)]$ . To this end, we use the time perturbation Lemma 5.3 which can be applied due to (5.25). Combining this with Remarks 4.16 and 4.17, in order to rewrite  $\ln \mathbb{E}_{u_0}^\xi [N^{\geq}(n/v_0, n)]$ , we obtain that

$$(5.26) \quad \begin{aligned} & \ln \mathbb{E}_{u_0}^\xi [N^{\geq}(n/v_0 + h_n + \alpha \ln n, n)] \\ &= \sum_{i=1}^n (L_i^\zeta(\bar{\eta}(v_0)) - L(\bar{\eta}(v_0))) + (h_n + \alpha \ln n)(\mathbf{es} - \bar{\eta}(v_0)) + \varepsilon(\alpha, n) \\ &= \alpha(\mathbf{es} - \bar{\eta}(v_0)) \ln n + \varepsilon(\alpha, n), \end{aligned}$$

where the last equality follows from  $\mathbf{es} - \bar{\eta}(v_0) + v_0 L(\bar{\eta}(v_0)) = 0$ ; cf. (A.9). Furthermore, the error term  $\varepsilon(\alpha, n)$  satisfies

$$|\varepsilon(\alpha, n)| \leq C + C(|h_n| + (|\alpha| \vee 1) \ln n) \left( \sqrt{\frac{\ln n}{n}} + \frac{|h_n| + (|\alpha| \vee 1) \ln n}{n} \right) + C \ln n,$$

with  $C$  depending on neither  $\alpha$  nor  $n$ . Choosing  $\alpha$  sufficiently large positive (resp., negative) the right-hand side of (5.26) converges to  $+\infty$  (resp.,  $-\infty$ ). Recalling the definition (5.21) of  $T_n$ , claim (5.23) follows for all  $n$  sufficiently large. Adjusting  $C$  then deals with the remaining  $n$ 's.

The law of large numbers (5.24) directly follows from (5.23) in combination with the ergodic theorem and the definitions from (4.6) and (4.7). The invariance principle is then again a consequence of this formula and Remark 4.16.  $\square$

Theorem 5.5 can be used to deduce a strong law of numbers for the breakpoint which does not follow easily from the previous argumentation.

COROLLARY 5.6. Under (POT), (INI) and (VEL),

$$\lim_{t \rightarrow \infty} \frac{\bar{m}(t)}{t} = v_0, \quad \mathbb{P}\text{-a.s.}$$

PROOF. Consider first the case  $u_0 = \delta_0$ . Then for  $\varepsilon > 0$ , by (2.2),  $\liminf \bar{m}(t)/t \geq (1 - \varepsilon)v_0$ ,  $\mathbb{P}$ -a.s. On the other hand, since  $\bar{m}(t)$  diverges  $\mathbb{P}$ -a.s. and  $t \geq T_{\bar{m}(t)}$  due to (5.22),

$$\liminf \frac{t}{\bar{m}(t)} \geq \liminf \frac{T_{\bar{m}(t)}}{\bar{m}(t)} = \frac{1}{v_0}, \quad \mathbb{P}\text{-a.s.},$$

by (5.24), completing the proof for  $u_0 = \delta_0$ . General  $u_0$  satisfying (INI) can then be handled using Lemmas 4.15 and 5.1.  $\square$

**6. The breakpoint approximates the maximum.** In this section, we prove the main results about the position of the rightmost particle  $M(t)$  and its median  $m(t)$ . We will see that those are well approximated by the breakpoint, and thus satisfy the same invariance principles.

It is elementary to obtain upper tail estimates for  $M(t)$  and an upper bound on  $m(t)$ : the definition of  $\bar{m}(t)$  and the Markov inequality imply directly that

$$(6.1) \quad \bar{m}(t) \geq m(t).$$

In addition, by Lemma 5.1(b),  $\mathbb{P}$ -a.s. for  $t$  large enough,

$$(6.2) \quad \mathbb{P}_{u_0}^\xi(M(t) \geq \bar{m}(t) + h) \leq \mathbb{E}_{u_0}^\xi[N(t, \bar{m}(t) + h)] \leq C e^{-ch}, \quad h \in (0, t^{1/3}).$$

Note that these estimates are rather coarse. One expects (6.2) to hold with  $m(t)$  instead of  $\bar{m}(t)$  and  $\bar{m}(t) - m(t) \asymp \ln t$ . These bounds, however, are more than sufficient to show the stated functional limit theorems.

As usual in the branching random walk literature, the lower bounds are more difficult, and are obtained via second moment estimates on the so-called leading particles. Since  $m(t)$  and  $M(t)$  are stochastically increasing in the initial condition, we will assume, without loss of generality, that  $u_0 = \mathbf{1}_{\{0\}}$  throughout this section.

6.1. *Leading particles.* We consider a special class of particles  $Y \in N(t)$  with trajectories satisfying

$$(6.3) \quad \begin{aligned} Y_t &\geq \bar{m}(t), & Y_{T_{\bar{m}(t)}} &\geq \bar{m}(t) \quad \text{and} \\ H_k^Y &\geq T_k - \alpha \psi^\xi(k) \quad \text{for all } 1 \leq k < \bar{m}(t), \end{aligned}$$

where  $H_k^Y = \inf\{s \geq 0 : Y_s = k\}$ ,  $\alpha > 2$  is a fixed constant,  $T_k$  is the breakpoint inverse introduced in (5.21) of Theorem 5.5, and  $\psi^\xi$  is defined by

$$(6.4) \quad \psi^\xi(k) = \mathcal{C}(\xi) + C_1(1 \vee \ln k),$$

where  $\mathcal{C}(\xi)$  and  $C_1$  are as in (5.23). Analogously to the literature on homogeneous branching random walk, we will call such particles *leading at time  $t$* . We further set

$$N_t^\mathcal{L} = |\{Y \in N(t) : Y \text{ is leading at time } t\}|.$$

The probability of finding a leading particle at time  $t$  is bounded from below in the following proposition.

PROPOSITION 6.1. *There exists a constant  $\gamma > 0$  such that  $\mathbb{P}$ -a.s. for all  $t$  large enough*

$$\mathbb{P}_0^\xi(N_t^\mathcal{L} \geq 1) \geq t^{-\gamma}.$$



The proof of this proposition will be based on the classical Paley–Zygmund inequality

$$(6.5) \quad \mathbb{P}_0^\xi(N_t^{\mathcal{L}} \geq 1) \geq \frac{\mathbb{E}_0^\xi[N_t^{\mathcal{L}}]^2}{\mathbb{E}_0^\xi[(N_t^{\mathcal{L}})^2]}.$$

Estimates for the expectations on the right-hand side are provided in the following two subsections. Since we do not strive to find the optimal constant  $\gamma$  in this paper, we use  $\gamma$  to denote a generic large constant whose value can change during the computations.

6.1.1. *First moment for the leading particles.*

LEMMA 6.2. *There exists a constant  $\gamma > 0$  such that  $\mathbb{P}$ -a.s. for all  $t$  large enough*

$$\mathbb{E}_0^\xi[N_t^{\mathcal{L}}] \geq t^{-\gamma}.$$

PROOF. Let  $\bar{t} = T_{\bar{m}(t)}$ . Then by (5.22),  $\bar{t} \leq t$ . Hence, every particle satisfying  $Y_{\bar{t}} \geq \bar{m}(t)$  has probability at least 1/2 to satisfy also  $Y_t \geq \bar{m}(t)$ . Further, by the definitions of  $\bar{t}$  and  $\bar{m}(t)$ , we have  $\mathbb{E}_0^\xi[N^{\geq}(\bar{t}, \bar{m}(t))] \geq 1/2$ . Therefore,

$$\mathbb{E}_0^\xi[N_t^{\mathcal{L}}] \geq \frac{\mathbb{E}_0^\xi[\{|Y \in N(\bar{t}) : Y_{\bar{t}} \geq \bar{m}(t), H_k \geq T_k - \alpha\psi^\xi(k) \forall k < \bar{m}(t)\}|]}{4\mathbb{E}_0^\xi[N^{\geq}(\bar{t}, \bar{m}(t))]}.$$

Using the Feynman–Kac representation (Proposition 3.1), this implies that

$$(6.6) \quad \mathbb{E}_0^\xi[N_t^{\mathcal{L}}] \geq \frac{E_0[e^{\int_0^{\bar{t}} \zeta(X_s) ds}; X_{\bar{t}} \geq \bar{m}(t), H_k \geq T_k - \alpha\psi^\xi(k) \forall k < \bar{m}(t)]}{4E_0[e^{\int_0^{\bar{t}} \zeta(X_s) ds}; X_{\bar{t}} \geq \bar{m}(t)]}.$$

Following the same steps as in the proof of the lower bound in Lemma 4.14, the numerator in (6.6) satisfies

$$\begin{aligned} & E_0[e^{\int_0^{\bar{t}} \zeta(X_s) ds}; X_{\bar{t}} \geq \bar{m}(t), H_k \geq T_k - \alpha\psi^\xi(k) \forall k < \bar{m}(t)] \\ & \geq E_0\left[e^{\int_0^{H_{\bar{m}(t)}} \zeta(X_s) ds} \times E_{\bar{m}(t)}[e^{\int_0^{\bar{t}} \zeta(X_s) ds}; X_r \geq \bar{m}(t)]\right]_{r=\bar{t}-H_{\bar{m}(t)}}; \\ & \quad H_{\bar{m}(t)} \in [\bar{t} - K, \bar{t}], H_k \geq T_k - \alpha\psi^\xi(k) \forall k < \bar{m}(t)] \\ & \geq cE_0[e^{\int_0^{H_{\bar{m}(t)}} \zeta(X_s) ds}; H_{\bar{m}(t)} \in [\bar{t} - K, \bar{t}], H_k \geq T_k - \alpha\psi^\xi(k) \forall k < \bar{m}(t)], \end{aligned}$$

where in the last step we used  $\text{ess inf}_{\zeta, r \leq K, x \in \mathbb{Z}} E_x[e^{\int_0^r \zeta(X_s) ds}; X_r \geq x] \geq c > 0$ , due to (POT). On the other hand, by Lemma 4.14, the denominator of (6.6) is bounded from above by  $CE_0[e^{\int_0^{H_{\bar{m}(t)}} \zeta(X_s) ds}; H_{\bar{m}(t)} \in [\bar{t} - K, \bar{t}]]$ . Replacing now  $\bar{m}(t)$  by  $n$ , and thus  $\bar{t}$  by  $T_n$ , and using the law of large numbers (5.24) for  $T_n$ , we observe from the previous reasoning that in order to show the lemma, it is sufficient to prove that  $\mathbb{P}$ -a.s., for all  $n$  large enough,

$$(6.7) \quad \frac{E_0[e^{\int_0^{H_n} \zeta(X_s) ds}; H_n \in [T_n - K, T_n], H_k \geq T_k - \alpha\psi^\xi(k) \forall k < n]}{E_0[e^{\int_0^{H_n} \zeta(X_s) ds}; H_n \in [T_n - K, T_n]]} \geq n^{-\gamma}.$$

To prove (6.7), we set  $\eta = \bar{\eta}(v_0)$  below and rewrite its left-hand side as

$$\begin{aligned} & \frac{E^{\zeta, \eta}[e^{-\eta H_n}; H_n \in [T_n - K, T_n], H_k \geq T_k - \alpha\psi^\xi(k) \forall k < n]}{E^{\zeta, \eta}[e^{-\eta H_n}; H_n \in [T_n - K, T_n]]} \\ & \geq c \cdot \frac{P^{\zeta, \eta}(H_n \in [T_n - K, T_n], H_k \geq T_k - \alpha\psi^\xi(k) \forall k < n)}{P^{\zeta, \eta}(H_n \in [T_n - K, T_n])} \\ & \geq c \cdot P^{\zeta, \eta}(H_n \in [T_n - K, T_n], H_k \geq T_k - \alpha\psi^\xi(k) \forall k < n). \end{aligned}$$

Setting  $\widehat{H}_n := H_n - E^{\zeta, \eta}[H_n]$  and  $R_n := T_n - E^{\zeta, \eta}[H_n]$  in the last formula, we thus see that (6.7) is equivalent to

$$(6.8) \quad P^{\zeta, \eta}(\widehat{H}_n \in [R_n - K, R_n], \widehat{H}_k \geq R_k - \alpha \psi^\xi(k) \ \forall k < n) \geq n^{-\gamma}$$

$\mathbb{P}$ -a.s. for all  $n$  large enough. Note that  $R_n$  depends on the random environment  $\xi$  only, so it is not random under  $P^{\zeta, \eta}$ .

The next two claims will show that, after rescaling, the processes  $R_n$  and  $\widehat{H}_n$  behave like Brownian motions. To approximate  $R_n$ , whose increments are not stationary, we introduce an auxiliary process with stationary increments

$$R'_n := \sum_{i=1}^n \rho_i, \quad n \geq 1,$$

where

$$(6.9) \quad \rho_i := \frac{1}{v_0 L(\eta)} (L_i^\zeta(\eta) - L(\eta)) - (E^{\zeta, \eta}[\tau_i] - \mathbb{E}[E^{\zeta, \eta}[\tau_i]]), \quad i \geq 1.$$

LEMMA 6.3. *The random variables  $R'_n$  are adapted to the filtration  $\mathcal{F}_n = \sigma(\xi(i) : i \leq n)$ , and  $R'_n$  approximates  $R_n$  in the sense that  $\mathbb{P}$ -a.s.,*

$$(6.10) \quad |R_n - R'_n| \leq \psi^\xi(n) \quad \text{for all } n \geq 0,$$

with  $\psi^\xi$  as in (6.4). Moreover, the sequence of increments  $(\rho_n)$  is bounded, stationary and there exist some constants  $c, C \in (0, \infty)$  such that

$$(6.11) \quad |\mathbb{E}[\rho_{n+m} \mid \mathcal{F}_n]| \leq C e^{-cm}.$$

Finally, there is  $\sigma_1^2 \in (0, \infty)$  such that both processes,  $[0, \infty) \ni t \mapsto n^{-1/2} R_{nt}$  and  $[0, \infty) \ni t \mapsto n^{-1/2} R'_{nt}$ , converge as  $n \rightarrow \infty$  in  $\mathbb{P}$ -distribution to a Brownian motion with variance  $\sigma_1^2$ .

PROOF. The adaptedness of  $(R'_n)$  to  $(\mathcal{F}_n)$ , as well as the stationarity and the boundedness of  $(\rho_n)$  follow directly from their definitions, recalling the assumption (POT). The estimate (6.10) follows from (5.23) of Theorem 5.5 after a straightforward computation. Furthermore, Lemma A.2 yields

$$|\mathbb{E}[L_{n+m}^\zeta(\eta) - L(\eta) \mid \mathcal{F}_n]| \leq C e^{-cm},$$

and analogically, bearing in mind that  $(L_i^\zeta)'(\eta) = E^{\zeta, \eta}[\tau_i]$ ,

$$|\mathbb{E}[E^{\zeta, \eta}[\tau_{n+m}] - \mathbb{E}[E^{\zeta, \eta}[\tau_{n+m}]] \mid \mathcal{F}_n]| \leq C e^{-cm},$$

proving (6.11).

Finally, observing that the increments of  $R'_n$  are centered, the functional central limit theorem for  $n^{-1/2} R'_n$  follows directly from a functional central limit theorem for stationary mixing sequences (see, e.g., Theorem 11 and Corollary 12 of [55]), the assumptions of which can be checked easily from (6.11). The functional central limit theorem for  $n^{-1/2} R_n$  then follows from (6.10).  $\square$

CLAIM 6.4. *There is  $\sigma_2^2 \in (0, \infty)$  such that  $\mathbb{P}$ -a.s., under  $P^{\zeta, \eta}$ ,  $n^{-1/2} \widehat{H}_n$  converges to a Brownian motion with variance  $\sigma_2^2$ .*

PROOF. Since the  $\tau_i$ 's are independent under  $P^{\zeta, \eta}$ ,  $\widehat{H}_n$  is a sum of independent and centered random variables, which have uniformly exponential tails. Moreover, the sequence of the variances of the increments is stationary under  $\mathbb{P}$ . The claim then follows easily by a functional version of the Lindeberg–Feller central limit theorem (see, e.g., [28], Theorem 9.3.1).  $\square$

REMARK 6.5. In view of the last claim and Lemma 6.3, the probability in (6.8) can approximatively be viewed as the probability that one Brownian motion stays above another, quenched, Brownian motion. This problem was recently studied in [51] for the case of two independent Brownian motions, where it was shown that this probability behaves like  $n^{-\gamma}$  with  $\gamma$  depending on the variances of the Brownian motions. More importantly, it was proved there that  $\gamma > 1/2$  whenever the variance of the quenched Brownian motion is positive. That implies that the price for a particle to be leading should be larger than in the homogeneous case, resulting thus in a larger backlog of  $m(t)$  behind  $\bar{m}(t)$ .

In this paper, the situation is more intricate due to the dependencies of the random variables involved. Hence, we do not strive for the optimal  $\gamma$ . Nevertheless, our proof partially builds on certain ideas appearing in [51].

We proceed by showing (6.8). In view of (6.10),

$$(6.12) \quad \begin{aligned} &P^{\zeta,\eta}(\widehat{H}_n \in [R_n - K, R_n], \widehat{H}_k \geq R_k - \alpha\psi^\xi(k) \forall 1 \leq k < n) \\ &\geq P^{\zeta,\eta}(\widehat{H}_n - R'_n \in I_n, \widehat{H}_k - R'_k \geq -(\alpha - 1)\psi^\xi(k) \forall 1 \leq k < n), \end{aligned}$$

where  $I_n = [R_n - R'_n - K, R_n - R'_n]$ . Note that since  $\alpha > 2$ , we have that  $R_n - R'_n - K \geq -(\alpha - 1)\psi^\xi(n)$  for  $n$  large enough.

On the right-hand side of (6.12), we require the process  $\widehat{H}_n - R'_n$  to stay above the barrier between times 0 and  $n$  and to be (almost) fixed at times 0 and  $n$ . It turns out useful to split the problem into two parts: distancing the barrier at 0, and distancing the barrier at  $n$ . Thus, we will consider two independent copies  $X^1$  and  $X^2$  of  $X$  under the same measure  $P^{\zeta,\eta}$ , and write  $\widehat{H}_k^i$ ,  $i = 1, 2$ , for the associated hitting times. We further consider a random variable  $\Sigma_n$  independent of  $X^1, X^2$ , which under  $P^{\zeta,\eta}$  is uniformly distributed on  $\{1, \dots, n - 1\}$ . We introduce

$$(6.13) \quad \beta_k^i = \widehat{H}_k^i - R'_k, \quad k \geq 0, i = 1, 2,$$

as a convenient abbreviation—mind, however, that  $\beta_k^i$  has a part,  $R'_k$ , that depends only on the random environment  $\xi$ , and another part,  $\widehat{H}_k^i$ , that depends on both,  $\xi$  and the random walk  $X^i$ . Furthermore, define

$$(6.14) \quad \beta_k = \begin{cases} \beta_k^1 & \text{for } 1 \leq k \leq \Sigma_n, \\ \beta_{\Sigma_n}^1 + (\beta_k^2 - \beta_{\Sigma_n}^2) & \text{for } \Sigma_n < k \leq n. \end{cases}$$

The process  $\beta$  has the increments of  $\beta^1$  before  $\Sigma_n$  and the increments of  $\beta^2$  after  $\Sigma_n$ . Since, under  $P^{\zeta,\eta}$ , the processes  $\widehat{H}^1$  and  $\widehat{H}^2$  are independent and have independent increments; it follows that the process  $\beta$  has, under  $P^{\zeta,\eta}$ , the same distribution as  $\widehat{H} - R'$ . Hence, (6.8) will follow if we show that,  $\mathbb{P}$ -a.s. for all  $n$  large enough,

$$(6.15) \quad P^{\zeta,\eta}(\beta_k \geq -(\alpha - 1)\psi^\xi(k) \forall 1 \leq k < n, \beta_n \in I_n) \geq n^{-\gamma}.$$

Finally, we write  $\bar{\beta}_k^1 = \beta_{k+1}^1 - \beta_1^1$ , and  $\bar{\beta}_k^2 = \beta_{n-k}^2 - \beta_n^2$ ,  $k = 0, \dots, n$ , for  $\beta^1$  shifted by one, and “ $\beta^2$  running backwards from  $n$ ,” respectively. Due to the independence of the increments of  $\beta^1$  under  $P^{\zeta,\eta}$ ,  $\beta_1^1$  is independent of  $\bar{\beta}^1$ . We then decompose  $\beta_n$  as

$$(6.16) \quad \beta_n = \beta_{\Sigma_n}^1 + (\beta_n^2 - \beta_{\Sigma_n}^2) = \beta_1^1 + \bar{\beta}_{\Sigma_n-1}^1 - \bar{\beta}_{n-\Sigma_n}^2.$$

The following lemma, the proof of which is postponed to the end of this subsection, provides a control on the processes  $\beta^1$  and  $\bar{\beta}$ .

LEMMA 6.6. (a) *There is  $\gamma' > 0$  such that  $\mathbb{P}$ -a.s. for all  $n$  large enough,*

$$P^{\zeta,\eta}(\bar{\beta}_k^1 \geq 0 \forall 1 \leq k \leq n, \bar{\beta}_n^1 \geq n^{1/4}) \geq n^{-\gamma'},$$

$$P^{\zeta,\eta}(\bar{\beta}_k^2 \geq 0 \forall 1 \leq k \leq n, \bar{\beta}_n^2 \geq n^{1/4}) \geq n^{-\gamma'}.$$

(b) *There is  $C_2 > 0$  such that  $\mathbb{P}$ -a.s. for  $n$  large enough,*

$$P^{\zeta,\eta}\left(\max_{1 \leq k \leq n} \max_{i=1,2} |\bar{\beta}_k^i - \bar{\beta}_{k-1}^i| \leq C_2 \ln n\right) \geq 1 - n^{-3\gamma'}.$$

(c) *Let  $\delta \in (0, 1)$ . There is  $c > 0$  such that  $\mathbb{P}$ -a.s., for all  $x > 0$ ,*

$$P^{\zeta,\eta}(\beta_1 \in [x, x + \delta]) \geq c\delta e^{-x/c}.$$

We now complete the proof of (6.15). With (6.14) and (6.16), we get

$$\begin{aligned} & \{\beta_k \geq -(\alpha - 1)\psi^\xi(k) \forall 1 \leq k < n, \beta_n \in I_n\} \\ & \supset \left(\{\bar{\beta}_k^1 \geq 0 \forall 1 \leq k \leq n, \bar{\beta}_n^1 \geq n^{1/4}\} \cap \left\{\max_{1 \leq k \leq n} |\bar{\beta}_k^1 - \bar{\beta}_{k-1}^1| \leq C_2 \ln n\right\}\right) \\ & \cap \left\{\bar{\beta}_k^2 \geq 0 \forall 0 \leq k \leq n, \bar{\beta}_n^2 \geq n^{1/4}\right\} \cap \left\{\max_{1 \leq k \leq n} |\bar{\beta}_k^2 - \bar{\beta}_{k-1}^2| \leq C_2 \ln n\right\} \\ & \cap \{\beta_1^1 \in (I_n - \bar{\beta}_{\Sigma_n-1}^1 + \bar{\beta}_{n-\Sigma_n}^2) \cap [0, \infty)\}. \end{aligned}$$

Indeed, the first and third event on the right-hand side ensure that the trajectories of  $\bar{\beta}^1$  and  $\bar{\beta}^2$  cross as on Figure 2, and stay above the barrier, which together with  $\beta_1^1 \geq 0$  of the fifth event ensures that  $\beta_k$  stays above the barrier as required. The second and the fourth event then ensure that at the time of crossing they are “sufficiently close” (which is not necessary for the inclusion to hold, but will be useful later). The fifth event in addition ensures that  $\beta_n \in I_n$ ; cf. (6.16).




FIG. 2. Construction of  $\beta$  (thick line) from  $\bar{\beta}^1, \bar{\beta}^2$  (thin lines) and  $\beta_1^1$ . For clarity of presentation,  $\bar{\beta}^1$  is drawn starting from 1, and  $\bar{\beta}^2$  is drawn running backwards from  $n$ .

By Lemma 6.6(a), (b) and the independence of  $\bar{\beta}^1, \bar{\beta}^2$  under  $P^{\zeta, \eta}$ , the probability of the intersection of the first four events on the right-hand side of the last display is at least  $n^{-2\gamma'}$ . In addition, if these four events occur, then there is  $J \in \{1, \dots, n\}$  such that  $I_n - \bar{\beta}_{J-1}^1 + \bar{\beta}_{n-J}^2 \subset [0, 2C_2 \ln n]$ . Moreover,  $\mathbb{P}(\Sigma_n = J) = 1/(n-1)$ . Hence, using Lemma 6.6(c), conditionally on the occurrence of the first four events and  $\Sigma_n = J$ , we can bound the probability of the fifth event on the right-hand side from below by  $c'n^{-1}e^{-C_2 \ln n/c} \geq n^{-\gamma''}$ . Combining these estimates proves that  $\mathbb{P}$ -a.s. for  $n$  large, (6.15), and thus also (6.8), is larger than  $n^{-\gamma}$  with  $\gamma > 1 + 2\gamma' + \gamma''$ . This completes the proof of (6.7), and thus of Lemma 6.2.  $\square$

We proceed by proving Lemma 6.6 which we used in the last proof.

**PROOF OF LEMMA 6.6.** Throughout the proof, we use the fact that the three processes  $\beta, \beta^1$  and  $\beta^2$  have, under  $P^{\zeta, \eta}$ , the same distribution as  $(\widehat{H}_k - R'_k)_{k \geq 0}$ . Since the statements of the lemma depend only on the respective distribution, we can and will therefore assume that these three processes are equal to  $(\widehat{H}_k - R'_k)_{k \geq 0}$ . In particular, their increments satisfy

$$(6.17) \quad \beta_k - \beta_{k-1} = \widehat{H}_k - R'_k - (\widehat{H}_{k-1} - R'_{k-1}) = \tau_k + \left( -\frac{L_k^\zeta(\eta)}{v_0 L(\eta)} \right),$$

where the last equality follows from definitions of  $\widehat{H}, R'$  and  $\rho$ . It is also useful to observe that, due to (POT), the second summand on the right-hand side of (6.17) satisfies

$$(6.18) \quad -\frac{1}{C} \geq -\frac{L_k^\zeta(\eta)}{v_0 L(\eta)} > -C \quad \text{for all } k \in \mathbb{N}, \mathbb{P}\text{-a.s.}$$

for some constant  $C \in (0, \infty)$ , and that the  $\tau_i$  are unbounded nonnegative random variables with uniform exponential tail (cf. (4.9)), that is, there exists  $c > 0$  such that for all  $k \geq 0$ ,  $\mathbb{P}$ -a.s.,

$$(6.19) \quad P^{\zeta, \eta}(\tau_k \geq u) \leq e^{-cu} \quad \text{for all } u \geq 0.$$

In particular, in combination with (6.18) we infer that there is a small constant  $c > 0$  such that  $\mathbb{P}$ -a.s.,

$$(6.20) \quad P^{\zeta, \eta}(\beta_k - \beta_{k-1} > c) > c \quad \text{and} \quad P^{\zeta, \eta}(\beta_k - \beta_{k-1} < -c) > c.$$

The claim (b) of the lemma then readily follows from (6.17)–(6.19), using a union bound and the fact that the increments of  $\bar{\beta}^i$  correspond directly to increments of  $\beta^i, i = 1, 2$ .

To prove (c), we write  $\beta_1^1 = \tau_1 - L_1^\zeta(\eta)/(v_0 L(\eta))$ , by (6.17). Recalling (6.18), it is sufficient to show that there exists  $c > 0$  such that,  $\mathbb{P}$ -a.s., we have  $P^{\zeta, \eta}(\tau_1 \in [x + y, x + y + \delta]) \geq c\delta e^{-x/c}$ , uniformly over  $y \in [0, C]$ . To see this, recall that under  $P^{\zeta, \eta}$ ,  $X$  is a Markov chain whose jump rate from 0 is bounded uniformly in  $\zeta$ , again by (POT). If the waiting time of  $X$  at 0 is in the required interval, and the first jump of  $X$  is to the right, then the required event is realized, proving (c).

Claim (a) is the most difficult. We first prove it for  $\bar{\beta}^1$ , and explain the modification required to show it for  $\bar{\beta}^2$  at the end of the proof. To simplify notation, we consider  $\beta$  instead of  $\bar{\beta}^1$ . This is possible since  $\bar{\beta}^1$  has the same distribution as  $\beta$  in the environment shifted by one.

In the proof, we often split the random environment  $\xi$  into two parts  $\underline{\xi}(j) = (\xi(k))_{k \leq j}$  and  $\bar{\xi}(j) = (\xi(k))_{k > j}$ . Set  $t_0 = t_{-1} = 0$  and  $t_i = 2^i$ , for  $i \geq 1$ . Fix  $a \in (0, \infty)$  and for  $i \geq 1$  define random variables  $Z_i$  by

$$\begin{aligned} Z_i &:= \operatorname{ess\,inf}_{\underline{\xi}(t_{i-2})} \inf_{x \geq at_{i-1}^{1/2}} P^{\zeta, \eta}(\beta_{t_i} \geq at_i^{1/2}, \beta_k \geq t_i^{1/4} \forall k \in \{t_{i-1}, \dots, t_i\} \mid \beta_{t_{i-1}} = x) \\ &= \operatorname{ess\,inf}_{\underline{\xi}(t_{i-2})} P^{\zeta, \eta}(\beta_{t_i} \geq at_i^{1/2}, \beta_k \geq t_i^{1/4} \forall k \in \{t_{i-1}, \dots, t_i\} \mid \beta_{t_{i-1}} = at_{i-1}^{1/2}). \end{aligned}$$

Here,  $\text{ess inf}_{\xi(t_{i-2})}$  means taking the essential infimum with respect to  $\xi(t_{i-2})$  and leaving the remaining  $\xi$  random. The second equality then follows from the obvious monotonicity of the considered event in the starting position. Observe that the random variable  $Z_i$  is  $\sigma(\xi(k), t_{i-2} < k \leq t_i)$  measurable, that is, the sequence  $(Z_i)$  is 1-dependent.

Setting  $i(n) = \lceil \log_2 n \rceil$ , using the Markov property, for  $n$  large enough,

$$P^{\zeta, \eta}(\beta_k \geq 0 \forall k \leq n, \beta_n \geq n^{1/4}) \geq \prod_{i=1}^{i(n)} Z_i = \exp\left\{\sum_{i=1}^{i(n)} \ln Z_i\right\}.$$

If we show that  $\mathbb{P}$ -a.s.,

$$(6.21) \quad \limsup_{k \rightarrow \infty} \frac{1}{k} \sum_{i=1}^k (-\ln Z_i) \leq c < \infty,$$

then the first half of claim (a) will follow with  $\gamma' > c/\ln 2$ .

First, we claim that,  $\mathbb{P}$ -a.s.,  $-\ln Z_i < \infty$ . Indeed, recalling (6.20), it is easy to see that the probability in the definition of  $Z_i$  is always positive. If we show that the  $(-\ln Z_i)$ 's have uniformly small exponential moments, then (6.21) will follow by standard arguments, using also the 1-dependence of the sequence  $Z_i$ .

To complete the proof, it is therefore sufficient to show that there exists some small  $\theta > 0$  such that for all  $i$  large enough

$$(6.22) \quad \mathbb{E}[\exp\{-\theta \ln Z_i\}] \leq c < \infty.$$

Throughout the proof of this inequality,  $i$  is considered fixed and we often omit it from the notation. To gain more independence, again we introduce  $\bar{\rho}_k^{(j)}$ ,  $j < k$ , by

$$(6.23) \quad \bar{\rho}_k^{(j)} := \text{ess sup}_{\xi^{(j)}} \rho_k.$$

Note that  $\rho_k$  is a  $\sigma(\xi(n), n \leq k)$ -measurable random variable, and thus  $\bar{\rho}_k^{(j)}$  is  $\sigma(\xi(n), j < n \leq k)$ -measurable. We further write  $\bar{R}_n^{(j)} = \sum_{k=1}^n \bar{\rho}_k^{(j)}$  and note that the increments of  $\bar{R}^{(j)}$  provide upper bounds for the increments of  $R'$ .

Let  $M_R$  be the essential supremum of the absolute value of the increments  $\rho_k$  of  $R'$ , which is finite by Lemma 6.3. Set

$$(6.24) \quad L := at_i^{1/2},$$

$r_0 := t_{i-1}$ , and define

$$s_0 := \inf\left\{k \geq r_0 : R'_k - R'_{r_0} \geq \frac{L}{8}\right\} \wedge t_i.$$

Further, recursively for  $j \geq 1$ , we define

$$(6.25) \quad \begin{aligned} r_{j+1} &:= s_j + \left\lceil \frac{L}{8M_R} \right\rceil, \\ s_{j+1} &:= \inf\left\{k \geq r_{j+1} : \bar{R}_k^{(s_j)} - \bar{R}_{r_{j+1}}^{(s_j)} \geq \frac{L}{8}\right\} \wedge (r_{j+1} + (t_i - t_{i-1})). \end{aligned}$$

Heuristically,  $s_j$  is the first time when  $\bar{R}$  (and thus possibly also  $R'$ ) “increases considerably after time  $r_j$ ”; due to the definition of  $\beta_k$  in (6.13), such a behavior of  $R'$  is potentially dangerous for the event in  $Z_i$ , in that it might lead to  $Z_i$  being very small. By definition,  $s_{j+1}$  depends only on  $\xi(l)$  with  $l > s_j$ , so the increments  $s_j - r_j$  are independent under  $\mathbb{P}$  and bounded by  $t_i - t_{i-1}$ .

For  $j \geq 0$ , consider the events

$$\mathcal{G}_j = \left\{ \beta_{s_j} \geq 2L, \inf_{r_j \leq l \leq s_j} \widehat{H}_l - \widehat{H}_{r_j} \geq -\frac{L}{8} \right\},$$

$$\mathcal{G}'_j = \left\{ \inf_{s_j \leq l \leq r_{j+1}} \widehat{H}_l - \widehat{H}_{s_j} \geq -\frac{L}{8} \right\},$$

and define

$$J = \inf\{j : s_j - r_j \geq t_i - t_{i-1}\}.$$

Finally, set  $\mathcal{G} = \bigcap_{j=0}^J \mathcal{G}_j \cap \bigcap_{j=0}^{J-1} \mathcal{G}'_j$ .

We claim that this construction ensures that

$$(6.26) \quad Z_i \geq P^{\zeta, \eta}(\mathcal{G} \mid \beta_{t_{i-1}} = at_{i-1}^{1/2}).$$

To see this, observe that in each of the time intervals  $[r_j, s_j]$  and  $[s_j, r_{j+1}]$ , the process  $\bar{R}$  (and thus also  $R'$ ) moves upward by at most  $L/8 + M_R$  by definition of these intervals. On the other hand, on  $\mathcal{G}$  the process  $\widehat{H}$  moves downward by at most  $L/8$  in any of these intervals. Since in the probability defining  $Z_i$  we condition on  $\beta_{r_0} = L/\sqrt{2} > L/2$  and, on  $\mathcal{G}$ ,  $\beta_{s_j} \geq 2L$ , this ensures that  $\beta_k \geq cL^{1/2} \geq t_i^{1/4}$  for  $k \in [r_0, s_0]$  and  $\beta_k \geq L$  for  $k \in [s_0, s_J]$ . Moreover, on  $\mathcal{G}$ ,  $s_J \geq t_i$ , proving (6.26).

Using the independence of the increments of  $\widehat{H}$  under the measure  $P^{\zeta, \eta}$ , the monotonicity of  $x \mapsto P^{\zeta, \eta}(\mathcal{G}_j \mid \beta_{r_j} = x)$ , and the fact that  $J$  is  $\sigma(\xi(x) : x \in \mathbb{Z})$ -measurable, we get

$$(6.27) \quad \begin{aligned} & P^{\zeta, \eta}(\mathcal{G} \mid \beta_{t_{i-1}} = at_{i-1}^{1/2}) \\ & \geq P^{\zeta, \eta}(\mathcal{G}_0 \mid \beta_{r_0} = L/\sqrt{2}) \prod_{j=1}^J P^{\zeta, \eta}(\mathcal{G}_j \mid \beta_{r_j} = 2L) \prod_{j=0}^{J-1} P^{\zeta, \eta}(\mathcal{G}'_j). \end{aligned}$$

It is not difficult to show, using the independence and the uniform exponential tail of the increments of  $\widehat{H}$  as well as the fact that they are centered, that if  $i$  is large enough,  $P^{\zeta, \eta}(\mathcal{G}'_j) \geq \frac{1}{2}$  for all  $j$ . On the other hand, for  $j \geq 1$ ,

$$(6.28) \quad \begin{aligned} & P^{\zeta, \eta}(\mathcal{G}_j \mid \beta_{r_j} = 2L) \\ & \geq P^{\zeta, \eta} \left( \widehat{H}_{s_j} - \widehat{H}_{r_j} \geq \frac{5L}{2}, \inf_{r_j \leq l \leq s_j} \widehat{H}_l - \widehat{H}_{r_j} \geq -\frac{L}{8} \right). \end{aligned}$$

Observing that the increments of  $\widehat{H}$  are independent under  $P^{\zeta, \eta}$  and the considered events are both increasing in those increments, we can use the Harris-FKG inequality to bound this from below by

$$(6.29) \quad \begin{aligned} & P^{\zeta, \eta} \left( \widehat{H}_{s_j} - \widehat{H}_{r_j} \geq \frac{5L}{2} \right) \times P^{\zeta, \eta} \left( \inf_{r_j \leq l \leq s_j} \widehat{H}_l - \widehat{H}_{r_j} \geq -\frac{L}{8} \right) \\ & \geq c \exp \left\{ -\frac{L^2}{c(s_j - r_j)} \right\}, \end{aligned}$$

with a sufficiently small constant  $c > 0$ . To obtain the last inequality, we used Azuma's inequality (together with the fact that  $\widehat{H}$  is a martingale under  $P^{\zeta, \eta}$  and the variances of its increments are uniformly bounded, by (POT)), and as well as Gaussian scaling to infer that the second factor is bounded from below by a constant (since  $s_j - r_j \leq cL^2$ ). By changing the constant  $c$ , the same lower bound holds for the first term on the right-hand side of (6.27) as well.



Coming back to (6.22), using (6.26)–(6.29), recalling that the increments of  $R'$  are exponentially mixing (cf. (6.11)) and that the intervals  $[r_j, s_j]$  are separated by spaces of length  $\frac{L}{8M_R}$ , we obtain

$$\begin{aligned}
 & \mathbb{E}[\exp\{-\theta \ln Z_i\}] \\
 & \leq C \mathbb{E} \left[ \exp \left\{ -\theta J \ln \frac{c}{2} + \theta \sum_{j=0}^J \frac{L^2}{c(s_j - r_j)} \right\} \right] \\
 (6.30) \quad & \leq C \sum_{k=1}^{\infty} \left( \frac{2}{c} \right)^{\theta k} \mathbb{E} \left[ e^{\frac{\theta L^2}{c(s_k - r_k)}} \mathbf{1}_{s_k - r_k = t_i - t_{i-1}} \prod_{j=0}^{k-1} e^{\frac{\theta L^2}{c(s_j - r_j)}} \mathbf{1}_{s_j - r_j < t_i - t_{i-1}} \right] \\
 & = C \sum_{k=1}^{\infty} \left( \frac{2}{c} \right)^{\theta k} e^{\frac{2\theta L^2}{ct_i} k-1} \prod_{j=0}^{k-1} \mathbb{E} \left[ e^{\frac{\theta L^2}{c(s_j - r_j)}} \mathbf{1}_{s_j - r_j < t_i/2} \right],
 \end{aligned}$$

where in the equality we used the independence of the  $s_j - r_j$ 's under  $\mathbb{P}$ . To upper bound the last expectation, we rewrite it as

$$\begin{aligned}
 (6.31) \quad & \int_0^{\infty} \mathbb{P}(e^{\frac{\theta L^2}{c(s_j - r_j)}} \mathbf{1}_{s_j - r_j < t_i/2} > a) \, da \\
 & \leq e^{\frac{2\theta L^2}{ct_i}} \mathbb{P}(s_j - r_j < t_i/2) + \int_e^{\infty} \frac{2\theta L^2}{ct_i} \mathbb{P}(e^{\frac{\theta L^2}{c(s_j - r_j)}} > a) \, da.
 \end{aligned}$$

Substituting  $a = \exp\{\frac{\theta L^2}{cy}\}$ , the second summand can be written as

$$(6.32) \quad \int_0^{t_i/2} \mathbb{P}(s_j - r_j < y) \frac{\theta L^2}{cy^2} e^{\frac{\theta L^2}{cy}} \, dy.$$

Recalling the definition (6.25) of  $s_j$ , for  $i$  sufficiently large we have for  $0 \leq y \leq t_i/2 = \frac{L^2}{2a^2}$  that

$$\begin{aligned}
 \mathbb{P}(s_j - r_j < y) &= \mathbb{P} \left( \max_{0 \leq m \leq y} \sum_{k=1}^m \bar{\rho}_{r_j+k}^{(s_{j-1})} \geq \frac{L}{8} \right) \\
 &\leq \mathbb{P} \left( \max_{1 \leq m \leq y} \sum_{k=1}^m \rho_{r_j+k} \geq \frac{L}{9} \right);
 \end{aligned}$$

here, to obtain the inequality one takes advantage of the estimates (A.6) and (A.8), which yield that uniformly in  $0 \leq j \leq k$ ,

$$0 \leq \bar{\rho}_k^{(j)} - \rho_k \leq C_{\Delta} e^{-(k-j)/C_{\Delta}},$$

and thus, using  $r_j - s_{j-1} \geq cL$ , that

$$0 \leq \bar{\rho}_{r_j+k}^{(s_{j-1})} - \rho_{r_j+k} \leq C e^{-cL} \quad \forall k \in \left\{ 0, \dots, \frac{L^2}{2a^2} \right\}.$$

Inequality (6.11) can then be used to verify the assumption of Azuma's inequality for mixing sequences of Lemma A.5 for the sequence  $\rho_k$ , and thus

$$\mathbb{P} \left( \sum_{k=1}^m \rho_{r_j+k} \geq \frac{L}{9} \right) \leq C e^{-cL^2/m},$$

for some constants  $C$  and  $c$  and all admissible  $m$ . This inequality extends to a maximal inequality, as follows from [41], Theorem 1,

$$\mathbb{P}\left(\max_{0 \leq m \leq y} \sum_{k=1}^m \rho_{r_j+k} \geq \frac{L}{9}\right) \leq C e^{-cL^2/y}.$$

Inserting these back into (6.32) implies that the second summand in (6.31) is smaller than

$$\int_0^{t_i/2} \frac{C\theta L^2}{y^2} e^{\frac{(\theta-c)L^2}{c'y}} dy = \int_0^{\frac{1}{2a^2}} \frac{C\theta}{z^2} e^{\frac{(\theta-c)}{c'z}} dz,$$

which can be made arbitrarily small by choosing  $\theta$  small. In addition, the first summand in (6.31) is strictly smaller than 1 by the functional central limit theorem from Lemma 6.3, hence the right-hand side of (6.31) is strictly smaller than one for all  $\theta > 0$  sufficiently small. Therefore, for  $\theta$  small enough the sum in (6.30) converges, which implies (6.22) and completes the proof of the first claim in Lemma 6.6(a).

The proof of the second claim is very similar, so we only explain the modifications which need to be introduced due to the fact that  $\bar{\beta}^2$  is “running backwards,” and thus its dependence on the environment  $\zeta$  is different. The first modification involves the definition of  $Z_i$  where the ess inf should be taken over  $\xi(n - t_{i+1})$ . This makes  $Z_i$  measurable with respect to  $\sigma(\xi(k) : n - t_{i+1} < k \leq n - t_{i-1})$ , and thus  $(Z_i)$  still is a 1-dependent sequence. Furthermore, the definitions of  $s_j, r_j$  should be replaced by  $r_0 = t_{i-1}$ , and

$$\begin{aligned} s_0 &:= \inf\left\{k \geq r_0 : \bar{R}_{n-k}^{(n-k-\ell)} - \bar{R}_{n-r_0}^{(n-k-\ell)} \geq \frac{L}{8}\right\} \wedge t_i, \\ r_{j+1} &:= s_j + \ell, \\ s_{j+1} &:= \inf\left\{k \geq r_{j+1} : \bar{R}_{n-k}^{(n-k-\ell)} - \bar{R}_{n-r_{j+1}}^{(n-k-\ell)} \geq \frac{L}{8}\right\} \wedge (r_{j+1} + (t_i - t_{i-1})), \end{aligned}$$

with  $\ell := \lceil L/8M_R \rceil$ , which again makes the increments  $(s_j - r_j)$  independent under  $\mathbb{P}$ .

With these modifications, the second claim in (a) can be shown almost exactly as the first one, which completes the proof of the lemma.  $\square$

6.1.2. *Second moment for the leading particles.* We now estimate the second moment of the number of leading particles needed for the application of (6.5). The proof is relatively short because we do not try to get the optimal power  $\gamma$  below.

LEMMA 6.7. *There exists a constant  $\gamma < \infty$  such that  $\mathbb{P}$ -a.s. for all  $t$  large enough,*

$$\mathbb{E}_0^\xi[(N_t^\mathcal{L})^2] \leq t^\gamma.$$

PROOF. Recall the definition (6.3) of leading particles. Since we look for an upper bound, we can ignore the condition  $Y_{\bar{m}(t)} \geq \bar{m}(t)$  there. We define a random function  $\varphi^\xi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  by

$$(6.33) \quad \varphi^\xi(s) := k \quad \text{for all } s \in [T_k - \alpha\psi^\xi(k), T_{k+1} - \alpha\psi^\xi(k+1)), k \in \mathbb{N}_0,$$

where  $\psi^\xi$  as in (6.4) and  $T_0 - \alpha\psi^\xi(0) := -\infty$ , by convention. By (3.2) of Proposition 3.1, we then have

$$\begin{aligned} (6.34) \quad & \mathbb{E}_0^\xi[(N_t^\mathcal{L})^2] \\ & \leq \mathbb{E}_0^\xi[N_t^\mathcal{L}] + 2 \int_0^t E_0 \left[ \exp\left\{ \int_0^s \xi(X_r) dr \right\} \xi(X_s) \mathbf{1}_{X_r \leq \varphi^\xi(r)} \forall r \in [0, s] \right. \\ & \quad \left. \times \left( E_{X_s} \left[ \exp\left\{ \int_0^{t-s} \xi(X_r) dr \right\} \mathbf{1}_{X_r \leq \varphi^\xi(s+r)} \forall r \in [0, t-s], X_{t-s} \geq \bar{m}(t) \right] \right)^2 \right] ds. \end{aligned}$$

In the upper bound of (6.34), we will repeatedly use the perturbation Lemmas 5.1 and 5.3 in a neighborhood  $(t, \bar{m}(t))$ . This is always justified  $\mathbb{P}$ -a.s. for  $t$  large enough, observing also that  $\bar{m}(t)/t \rightarrow v_0 \in V$ ,  $\mathbb{P}$ -a.s., by Corollary 5.6. In particular, by Lemma 5.1(b) and the definition of  $\bar{m}(t)$ ,  $\mathbb{P}$ -a.s. for  $t$  large enough,

$$(6.35) \quad \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t) + 1)] \leq C/2.$$

The first summand on the right-hand side of (6.34) then satisfies

$$\mathbb{E}_0^\xi[N_t^{\mathcal{L}}] \leq \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C.$$

Since  $\xi(X_s) \leq es$ , the second summand on the right-hand side of (6.34) is bounded from above by

$$(6.36) \quad \begin{aligned} & 2es \int_0^t \sum_{k=-\infty}^{\varphi^\xi(s)} \mathbb{E}_0[e^{\int_0^s \xi(X_r) dr} \mathbf{1}_{X_s=k} (E_k[e^{\int_0^{t-s} \xi(X_r) dr}; X_{t-s} \geq \bar{m}(t)])^2] ds \\ & = 2es \int_0^t \sum_{k=-\infty}^{\varphi^\xi(s)} \mathbb{E}_0^\xi[N(s, k)] \mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))]^2 ds. \end{aligned}$$

To find an upper bound for the integral on the right-hand side of (6.36), we remark that, by the first moment formula (3.1) of Proposition 3.1, the Markov property, and (6.35),  $\mathbb{P}$ -a.s. for  $t$  large enough,

$$\begin{aligned} \mathbb{E}_0^\xi[N(s, k)] \mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))] &= \mathbb{E}_0^\xi[|\{Y \in N(t), Y_t \geq \bar{m}(t), Y_s = k\}|] \\ &\leq \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C. \end{aligned}$$

Hence,  $\mathbb{P}$ -a.s. for  $t$  large enough, uniformly in  $s \in [0, t]$ ,  $k \leq \varphi^\xi(s)$ ,

$$(6.37) \quad \mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))] \leq C/\mathbb{E}_0^\xi[N(s, k)].$$

In order to take advantage of (6.37), we treat separately four ranges of parameters  $s \in [0, t]$  and  $k \leq \varphi^\xi(s)$  in (6.36).

(A) We start with considering the range

$$(6.38) \quad \mathcal{N}_1(\xi) \leq k \leq \varphi^\xi(s), \quad s \geq \mathcal{S}(\xi), \text{ such that } k/s \in V,$$

where  $\mathcal{N}_1(\xi)$  and  $\varphi^\xi$  are defined in (4.21), (4.22) and (6.33), respectively, and  $\mathcal{S}(\xi)$  is a  $\sigma(\xi)$ -measurable random variable which is a.s. finite and which will be specified below. In this case, by Lemma 4.15,

$$\mathbb{E}_0^\xi[N(s, k)] \geq c \mathbb{E}_0^\xi[N^\geq(s, k)] \geq c \mathbb{E}_0^\xi[N^\geq(s, \varphi^\xi(s))],$$

where the last inequality follows from  $k \leq \varphi^\xi(s)$ . Let  $l = l(s)$  be such that  $s \in [T_l - \alpha\psi^\xi(l), T_{l+1} - \alpha\psi^\xi(l+1))$ ; note that  $\varphi^\xi(s) = l$ . Let  $\mathcal{S}(\xi)$  be a  $\mathbb{P}$ -a.s. finite random variable such  $s \geq \mathcal{S}(\xi)$  implies that  $l/T_l \in V$ ,  $l/(T_l - \alpha\psi^\xi(l)) \in V$  and  $s \geq T_l - \alpha\psi^\xi(l) \geq \mathcal{T}_1 \vee \mathcal{T}_2$ , where  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are as in Lemmas 5.1(b) and 5.3(b). The existence of such  $\mathcal{S}$  is implied by the law of large numbers (5.24) for  $T_n$ . Using then repeatedly Lemma 5.3(b) and Remark 5.4, the right-hand side of the previous display can be bounded from below by

$$c \mathbb{E}_0^\xi[N^\geq(T_l - \alpha\psi^\xi(l), l)] \leq c \mathbb{E}_0^\xi[N^\geq(T_l, l)] e^{-c\psi^\xi(l)} \geq \mathcal{C}'(\xi) t^{-\gamma},$$

for some  $\gamma \in (0, \infty)$  and a positive random variable  $\mathcal{C}'(\xi)$ , where in the last inequality we used  $\mathbb{E}_0^\xi[N^\geq(T_l, l)] = 1/2$ , and  $\psi^\xi(l) = \mathcal{C}(\xi) + C_1(1 \vee \ln l)$ , the need for which emanates

from the randomness of  $\psi^\xi$ . Thus, combining the last two displays with (6.37) we infer that  $\mathbb{P}$ -a.s. for  $t$  large enough, uniformly for  $k, s$  as in (6.38),

$$(6.39) \quad \mathbb{E}_0^\xi[N(s, k)]\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]^2 \leq C\mathbb{E}_0^\xi[N(s, k)]^{-1} \leq C''(\xi)t^\gamma.$$

(B) We now consider the ranges

$$(6.40) \quad s \in [0, t] \quad \text{and} \quad k \quad \text{such that} \quad |k/s| \leq \bar{v}/2,$$

where  $\bar{v} > 0$  is the asymptotic speed of the maximal particle in the homogeneous branching random walk with branching rate  $\mathbf{ei}$  (cf. (POT)). We assume without loss of generality that  $V$  is fixed so that it contains  $\bar{v}/2$  in its interior. Since  $\xi(x) \geq \mathbf{ei}$ , by a straightforward comparison argument and properties of the homogeneous branching random walk, we infer the existence of some constant  $c > 0$  such that  $\mathbb{E}_0^\xi[N(s, k)] \geq c$  for all  $s$  and  $k$  as in (6.40). Therefore, by (6.37),  $\mathbb{P}$ -a.s. for  $t$  large enough, uniformly for  $s, k$  as in (6.40),

$$(6.41) \quad \mathbb{E}_0^\xi[N(s, k)]\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]^2 \leq c\mathbb{E}_0^\xi[N(s, k)]^{-1} \leq C.$$

(C) Now let

$$s \in [0, t] \quad \text{and} \quad k \leq 0.$$

By the Feynman–Kac formula, using also  $\text{ess inf } \xi \geq 0$ ,

$$\begin{aligned} 2\mathbb{E}_k^\xi[N^\geq(t, \bar{m}(t))] &\geq 2E_k \left[ \exp \left\{ \int_0^{t-s} \xi(X_r) dr \right\} \mathbf{1}_{X_{t-s} \geq \bar{m}(t)} \mathbf{1}_{X_t \geq \bar{m}(t)} \right] \\ &\geq 2\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]P_0(X_s \geq 0) \geq \mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]. \end{aligned}$$

Therefore,

$$\begin{aligned} &\mathbb{E}_0^\xi[N(s, k)]\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]^2 \\ &\leq 2\mathbb{E}_0^\xi[N(s, k)]\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]\mathbb{E}_k^\xi[N^\geq(t, \bar{m}(t))]. \end{aligned}$$

For  $k \leq 0$ , by the monotonicity in the initial condition, taking advantage of Lemma 4.15,  $\mathbb{P}$ -a.s. for  $t$  large enough,

$$\mathbb{E}_k^\xi[N^\geq(t, \bar{m}(t))] \leq \mathbb{E}_{\mathbf{1}_{-\mathbb{N}_0}}^\xi[N^\geq(t, \bar{m}(t))] \leq C\mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C.$$

Combining the last two inequalities, applying also Markov property and (6.35),  $\mathbb{P}$ -a.s. for  $t$  large enough, uniformly in  $s \in [0, t]$ ,

$$(6.42) \quad \begin{aligned} &\sum_{k=-\infty}^0 \mathbb{E}_0^\xi[N(s, k)]\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))]^2 \\ &\leq C \sum_{k=-\infty}^0 \mathbb{E}_0^\xi[N(s, k)]\mathbb{E}_k^\xi[N^\geq(t - s, \bar{m}(t))] \\ &\leq C\mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C. \end{aligned}$$

(D) The remaining part of the range of parameters relevant in (6.36), which is not controlled by (A)–(C), is a subset of

$$\mathcal{B}^\xi = \{(s, k) \in [0, \infty) \times \mathbb{N} : \bar{v}s/2 \leq k \leq \varphi^\xi(s), s + k \leq \tilde{C}(\xi)\}$$

for some finite random variable  $\tilde{C}(\xi)$  depending on  $\mathcal{N}_1$  and  $\mathcal{S}$ . Observe that  $\mathcal{B}^\xi$  is a bounded set for  $\mathbb{P}$ -a.e.  $\xi$ .

We start with observing that there is a constant  $L > 0$  such that

$$(6.43) \quad \text{for } t \text{ large enough, } \mathbb{P}\text{-a.s., } \bar{m}(t+1) - \bar{m}(t) \leq L.$$

Indeed, by the perturbation Lemmas 5.1(b) and 5.3(b), and (6.35),

$$\begin{aligned} \mathbb{E}_0^\xi[N^\geq(t+1, \bar{m}(t) + L)] &\leq C e^C \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t) + L)] \\ &\leq C' e^{-cL} \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C'' e^{-cL}. \end{aligned}$$

Choosing  $L$  to make the right-hand side smaller than 1 then yields (6.43).

The definition of  $\bar{m}(t)$ , inequality (6.43), and Lemma 5.1(b) imply together that,  $\mathbb{P}$ -a.s. for  $t$  large enough,

$$\begin{aligned} 1/2 &\geq \mathbb{E}_0^\xi[N^\geq(t+1, \bar{m}(t+1) + 1)] \\ &\geq c e^{-c(\bar{m}(t+1)+1-\bar{m}(t))} \mathbb{E}_0^\xi[N^\geq(t+1, \bar{m}(t))] \\ &\geq c e^{-cL} \mathbb{E}_0^\xi[N^\geq(t+1, \bar{m}(t))]. \end{aligned}$$

Hence with  $C = e^{cL}/(2c)$ ,  $\mathbb{P}$ -a.s. for  $t$  large enough, using also the Markov property,

$$\begin{aligned} C &\geq \mathbb{E}_0^\xi[N^\geq(t+1, \bar{m}(t))] \\ &\geq \mathbb{E}_0^\xi[N(s+1, k)] \mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))]. \end{aligned}$$

By the boundedness of  $\mathcal{B}^\xi$  and (POT), there is a random variable  $\mathcal{C}(\xi) \in (0, \infty)$  such that, for all  $(s, k) \in \mathcal{B}^\xi$ ,

$$\mathcal{C}(\xi) \geq \mathbb{E}_0^\xi[N(s+1, k)] \geq P_0[X_{s+1} = k] \geq \mathcal{C}(\xi)^{-1}.$$

Combining the last two inequalities then yields

$$\mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))] \leq \mathcal{C}'(\xi)$$

for all  $(s, k) \in \mathcal{B}^\xi$ ,  $\mathbb{P}$ -a.s. for  $t$  large enough. Hence, following the same arguments as before, using (6.35), we obtain

$$(6.44) \quad \begin{aligned} &\mathbb{E}_0^\xi[N(s, k)] \mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))]^2 \\ &\leq \mathcal{C}'(\xi) \mathbb{E}_0^\xi[N(s, k)] \mathbb{E}_k^\xi[N^\geq(t-s, \bar{m}(t))] \\ &\leq \mathcal{C}'(\xi) \mathbb{E}_0^\xi[N^\geq(t, \bar{m}(t))] \leq C \mathcal{C}'(\xi), \end{aligned}$$

uniformly for  $(s, k) \in \mathcal{B}^\xi$ ,  $\mathbb{P}$ -a.s. for  $t$  large enough.

Using inequalities (6.39), (6.41), (6.42) and (6.44) in their respective domains in the summation and integration in (6.36) (recalling that  $V$  contains  $\bar{v}/2$  in its interior), we can,  $\mathbb{P}$ -a.s. for  $t$  large enough, bound the second summand on the right-hand side of (6.34) from above by

$$C''(\xi) t^{\gamma+2} + C t^2 + C t + C \mathcal{C}'(\xi),$$

where the summands correspond to cases (A)–(D) above. This completes the proof of the lemma.  $\square$

Combining Lemmas 6.2 and 6.7 with (6.5) completes the proof of Proposition 6.1.

6.2. *Proof of Theorem 2.4 and Proposition 2.3.* By inserting the estimates from Lemmas 6.2 and 6.7 into the Paley–Zygmund inequality (6.5) we obtain  $\mathbb{P}_0^\xi(N_t^{\mathcal{L}} \geq 1) \geq t^{-\gamma}$  for all large  $t$ ,  $\mathbb{P}$ -a.s. To complete the proof of the lower bound in Theorem 2.4, we need to amplify this estimate, using a technique adapted from the homogeneous branching random walk literature (see, e.g., [53]). The first step is the following lemma guaranteeing that with very high probability the number of particles in the origin grows exponentially in time.

LEMMA 6.8. *There exists  $C_3 > 1$  and  $t_0 < \infty$  such that such that for all  $t \geq t_0$  and  $\mathbb{P}$ -a.e.  $\xi$ ,*

$$\mathbb{P}_0^\xi(N(t, 0) \leq C_3^t) \leq C_3^{-t}.$$

PROOF. Recall from (POT) that the essential infimum  $\text{ei}$  of the  $\xi$  is strictly positive. By the monotonicity of  $N(t, 0)$  in  $\xi$  which can be ensured by a straightforward coupling, it suffices to show the claim for the homogeneous branching random walk with branching rate  $\text{ei}$ . We write  $\mathbb{P}_0^{\text{ei}}$  for the law of this process starting in 0.

For  $t \geq 0$  and  $\varepsilon > 0$ , let  $D_\varepsilon(t/3)$  be the set of *direct* offsprings of the initial particle until time  $t/3$  which are at sites  $[-\varepsilon t, \varepsilon t]$  at time  $t/3$ . Then, for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$(6.45) \quad \mathbb{P}_0^{\text{ei}}(|D_\varepsilon(t/3)| \leq \delta t/3) \leq e^{-\delta t/3}.$$

Indeed, the probability that the initial particle<sup>2</sup> leaves  $[-\varepsilon t/2, \varepsilon t/2]$  before  $t/3$  is smaller than  $e^{-c(\varepsilon)t}$ . If it stays in this interval, it produces more than  $t\text{ei}/4$  direct offsprings with probability larger than  $1 - e^{-ct}$ , by large deviations for the Poisson distribution with parameter  $t\text{ei}/3$ , and every of these offsprings stays in  $[-\varepsilon t, \varepsilon t]$  with probability at least  $1 - e^{-c(\varepsilon)t}$  again.

For a particle  $Y \in D_\varepsilon(t/3) \subset N(t/3)$ , we denote by  $A_Y(2t/3)$  the set of *all* offsprings it produced between times  $t/3$  and  $2t/3$  and which are at the site  $Y_{t/3}$  at time  $2t/3$ . We claim that there exists  $c > 1$  such that

$$(6.46) \quad \mathbb{P}_0^{\text{ei}}(|A_Y(2t/3)| \geq c^t) > 0.$$

Indeed, under  $\mathbb{P}^{\text{ei}}$ , it is well known (e.g., it follows from the Feynman–Kac formula) that the expected number of particles in 0 grows exponentially. Hence, we can fix  $r > 0$  such that  $\mathbb{E}^{\text{ei}}[N(r, 0)] =: \mu > 1$ , and consider an auxiliary process evolving as follows:

- start with one particle at an arbitrary site  $x \in \mathbb{Z}$  at time 0,
- particles evolve independently as a continuous time simple random walk and split into two at rate  $\text{ei}$ ,
- and at each time  $rn, n \in \mathbb{N}$ , all the particles not at  $x$  are killed.

Let  $Z_n$  be the number of particles at  $x$  at time  $rn$  in this auxiliary process. It is easy to see that  $Z_n$  is a supercritical Galton–Watson process, and thus it survives with a positive probability,  $\mathbb{P}_0^{\text{ei}}(Z_n > 0 \forall n \geq 0) \geq p > 0$ , and on the event of survival it grows exponentially,  $\mathbb{P}_0^{\text{ei}}(Z_n \geq c^n \mid Z_n > 0 \forall n \geq 0) \geq 1/2$  for some  $c > 1$ . Hence, for every  $Y \in D_\varepsilon(t/3)$ ,  $\mathbb{P}_0^{\text{ei}}(|A_Y(2t/3)| \geq c^t) \geq p/2$  at all times such that  $2t/3 = rn$  for some  $n \in \mathbb{N}$ . A straightforward extension to all times then yields (6.46).

Combining (6.45) and (6.46) implies that

$$\mathbb{P}_0^{\text{ei}}(N(2t/3, [-\varepsilon t, \varepsilon t]) \geq c^t) \geq 1 - e^{-c^t}.$$

---

<sup>2</sup>For simplicity of redaction, in a slight abuse of notation we reformulate the original branching mechanism which replaces a particle by two new particles, by the equivalent branching mechanism where instead particles are not replaced and give birth to one more particle.

Moreover, the constant  $c$  does not depend on  $\varepsilon$ . As a consequence, choosing  $\varepsilon > 0$  small enough such that

$$P_{\varepsilon t}(X_{t/3} = 0) \geq \left(\frac{1+c}{2}\right)^{-t/3},$$

and using an easy large deviation argument we obtain that

$$P_0^{\text{ei}}\left(N(t, 0) < \frac{c^t}{2} \left(\frac{1+c}{2}\right)^{-t/3} \mid N(2t/3, [-\varepsilon t, \varepsilon t]) \geq c^t\right) \leq e^{-c''t}.$$

This completes the proof of the lemma.  $\square$

We now obtain a lower bound on  $M(t)$ .

**PROPOSITION 6.9.** *For any  $q \in \mathbb{N}$ , there exists a constant  $C^{(q)} < \infty$  such that for  $\mathbb{P}$ -a.a.  $\xi$ , for all  $t$  large enough*

$$P_0^\xi(M(t) \geq \bar{m}(t) - C^{(q)} \ln t) \geq 1 - 2t^{-q}.$$

**PROOF.** Without loss of generality, we assume that  $q > \gamma$  for  $\gamma$  as in Proposition 6.1. We fix  $r = c_1 \ln t$  where  $c_1$  is chosen so that for  $C_3$  of Lemma 6.8 we have  $C_3^{-r} = t^{-q}$ , and further choose  $C^{(q)}$  large enough so that  $\bar{m}(t - r) \geq \bar{m}(t) - C^{(q)} \ln t$ . To see that this is possible, observe that by the perturbation Lemmas 5.1 and 5.3 we have for some  $c, c' \in (0, \infty)$ ,

$$\begin{aligned} E_0^\xi[N^\geq(t - r, \bar{m}(t) - C^{(q)} \ln t)] &\geq e^{-cr} E_0^\xi[N^\geq(t, \bar{m}(t) - C^{(q)} \ln t)] \\ &\geq e^{-cr} e^{c' C^{(q)} \ln t} E_0^\xi[N^\geq(t, \bar{m}(t))] \\ &\geq \frac{1}{2} e^{-cr} e^{c' C^{(q)} \ln t}, \end{aligned}$$

and fix  $C^{(q)}$  so that the right-hand side is smaller than  $1/2$ .

Set  $x := \bar{m}(t) - C^{(q)} \ln t$  and observe that by considering separately the events  $\{N(r, 0) < C_3^r\}$ ,  $\{N(r, 0) \geq C_3^r\}$  and using the Markov property and the independence of the particles in the second case

$$\begin{aligned} P_0^\xi(M(t) \geq x) &= P_0^\xi(N^\geq(t, x) \geq 1) \\ &\geq 1 - P_0^\xi(N(r, 0) \leq C_3^r) - (P_0^\xi(N^\geq(t - r, x) < 1))^{C_3^r} \\ &\geq 1 - t^{-q} - (P_0^\xi(N_{t-r}^\mathcal{L} < 1))^{C_3^r}. \end{aligned}$$

Here, for the last inequality we used Lemma 6.8 as well as  $x \leq \bar{m}(t - r)$  and so  $N^\geq(t - r, x) \geq N_{t-r}^\mathcal{L}$ . Proposition 6.1 then implies

$$(P_0^\xi(N_{t-r}^\mathcal{L} < 1))^{C_3^r} \leq (1 - t^{-\gamma})^{t^q} \leq t^{-q}$$

for  $t$  large enough. This completes the proof.  $\square$

**PROOF OF THEOREM 2.4 AND PROPOSITION 2.3.** Proposition 6.9 and the Borel–Cantelli lemma (controlling noninteger  $t$  by standard estimates) imply that  $M(t) \geq \bar{m}(t) - C^{(2)} \ln t$ ,  $\mathbb{P} \times P_0^\xi$ -a.s. for all  $t$  large enough, and thus  $\mathbb{P}$ -a.s.,  $m(t) \geq \bar{m}(t) - C^{(2)} \ln t$ , for such  $t$  as well. By the monotonicity and the independence of the particles, these lower bounds hold for an arbitrary initial condition satisfying (INI). These facts combined with (6.2) and  $m(t) \leq \bar{m}(t)$  (cf. (6.1)), complete the proof of Theorem 2.4 and Proposition 2.3.  $\square$

**7. BRWRE and the randomized Fisher–KPP.** In this section, we prove the central limit theorem for the front of the randomized Fisher–KPP equation. We begin by establishing the connection between the BRWRE and this Fisher–KPP equation. Its proof is a straightforward adaptation of [54], who proved the corresponding result in the case of the homogeneous BBM (see also [38–40]).

PROPOSITION 7.1. *For a bounded  $(\xi(x))_{x \in \mathbb{Z}}$  and  $f : \mathbb{Z} \rightarrow [0, 1]$ ,*

$$w(t, x) := 1 - \mathbb{E}_x^\xi \left[ \prod_{Y \in N(t)} f(Y_t) \right]$$

solves

$$\frac{\partial w}{\partial t} = \Delta_d w + \xi(x)w(1 - w)$$

with initial condition  $w(0, \cdot) = 1 - f$ . In particular,  $w(t, x) = \mathbb{P}_x^\xi(M(t) \geq 0)$  solves this equation with  $f = \mathbf{1}_{-\mathbb{N}}$ , that is,  $w(0, \cdot) = \mathbf{1}_{\mathbb{N}_0}$ .

PROOF. Actually, we show that  $v := 1 - w$  solves

$$\frac{\partial v}{\partial t} = \Delta_d v - \xi(x)v(1 - v)$$

with initial condition  $v(0, \cdot) = f$ , which will establish the claim.

According to whether or not the original particle has split into two before time  $t$ , the Feynman–Kac formula in combination with the Markov property at time  $s$  supplies us with

$$v(t, x) = E_x \left[ e^{-\int_0^t \xi(X_r) dr} f(X_t) \right] + \int_0^t E_x \left[ \xi(X_s) e^{-\int_0^s \xi(X_r) dr} v^2(t - s, X_s) \right] ds.$$

Using the reversibility of the random walk, and substituting  $s$  by  $t - s$ , this can be written as

$$\begin{aligned} v(t, x) &= \sum_{y \in \mathbb{Z}} f(y) E_y \left[ e^{-\int_0^t \xi(X_r) dr} \mathbf{1}_x(X_t) \right] \\ (7.1) \quad &+ \int_0^t \sum_{y \in \mathbb{Z}} v^2(s, y) E_y \left[ \xi(y) e^{-\int_0^{t-s} \xi(X_r) dr} \mathbf{1}_x(X_{t-s}) \right] ds. \end{aligned}$$

Differentiation then yields

$$\begin{aligned} \frac{\partial v}{\partial t}(t, x) &= \sum_{y \in \mathbb{Z}} f(y) E_y \left[ -\xi(x) e^{-\int_0^t \xi(X_r) dr} \mathbf{1}_x(X_t) \right] \\ &+ \sum_{y \in \mathbb{Z}} f(y) E_y \left[ e^{-\int_0^t \xi(X_r) dr} (\Delta_d \mathbf{1}_x)(X_t) \right] \\ &+ \xi(x) v^2(t, x) \\ &+ \int_0^t \sum_{y \in \mathbb{Z}} v^2(s, y) E_y \left[ -\xi(y) \xi(x) e^{-\int_0^{t-s} \xi(X_r) dr} \mathbf{1}_x(X_{t-s}) \right] ds \\ &+ \int_0^t \sum_{y \in \mathbb{Z}} v^2(s, y) E_y \left[ \xi(y) e^{-\int_0^{t-s} \xi(X_r) dr} (\Delta_d \mathbf{1}_x)(X_{t-s}) \right] ds. \end{aligned}$$

Comparing this expression with the representation (7.1), the second and fifth summands together yield  $\Delta_d v$ , the third is  $\xi(x)v^2$ , and the first and fourth together supply us with  $-\xi(x)v$ ,



which completes the proof of the first claim. The second claim is a straightforward consequence of the first one.  $\square$

**PROOF OF THEOREM 2.9.** Observe that the initial conditions in Theorem 2.9 and the second claim of Proposition 7.1 are related by the reflection  $x \mapsto -x$ . Hence, setting  $\tilde{\xi}(x) = \xi(-x)$ , it is easy to see from the last proposition that the front  $\widehat{m}(t)$  of the Fisher-KPP equation defined in (2.15) can be represented as

$$\widehat{m}(t) = \sup \left\{ x \in \mathbb{Z} : \mathbb{P}_{-x}^{\tilde{\xi}}(M(t) \geq 0) \geq \frac{1}{2} \right\}.$$

Comparing this to the definition (2.4) of  $m(t)$ , we see that the role of  $x$  and the origin is reversed, and the environment is reflected. This complication is easy to resolve. By the translation and reflection invariance of the environment  $\xi$ , for every  $x \in \mathbb{Z}$ ,

$$\mathbb{P} \left( \mathbb{P}_{-x}^{\tilde{\xi}}(M(t) \geq 0) \geq \frac{1}{2} \right) = \mathbb{P} \left( \mathbb{P}_0^{\xi}(M(t) \geq x) \geq \frac{1}{2} \right).$$

The central limit theorem for  $\widehat{m}(t)$  then follows from the one for  $m(t)$ .  $\square$

**8. Open questions.** We collect here some open questions which naturally arise from the investigations of this article.

1. Can we say that  $m(t)$  lags at least  $\Omega(\ln t)$  behind  $\bar{m}(t)$ ?
2. For  $x \in \mathbb{Z}$  fixed, is the function  $[0, \infty) \ni t \mapsto u(t, x)$  increasing? It is not hard to see that generally this is not the case on  $[0, \infty)$ ; however, is it true for  $t$  large enough?
3. Is the family  $M(t) - m(t)$ ,  $t \geq 0$ , tight? In the case of homogeneous BBM, it already follows from the convergence to a traveling wave solution (see [43]) that this is the case. In the case of spatially random branching rates, this remains an open question.

We expect our results to transfer to the continuum setting where the space  $\mathbb{Z}$  is replaced by  $\mathbb{R}$  under suitable regularity and mixing assumptions on  $\xi$ .

### APPENDIX: AUXILIARY RESULTS

We prove here several auxiliary results that are used through the text. Most of them use rather standard techniques, but we did not find any suitable reference for them.

#### A.1. Properties of logarithmic moment generating functions.

**LEMMA A.1.** *The functions  $L$ ,  $L_i^\zeta$ , and  $\bar{L}_n^\zeta$  defined in (4.5)–(4.7) are infinitely differentiable on  $(-\infty, 0)$  and satisfy for  $\eta \in (-\infty, 0]$ ,*

$$(A.1) \quad L'(\eta) = \mathbb{E} \left[ \frac{E^\zeta[H_1 e^{\eta H_1}]}{E^\zeta[e^{\eta H_1}]} \right] = \mathbb{E}[E^{\zeta, \eta}[H_1]],$$

$$(A.2) \quad (L_i^\zeta)'(\eta) = \frac{E^\zeta[\tau_i e^{\eta \tau_i}]}{E^\zeta[e^{\eta \tau_i}]} = E^{\zeta, \eta}[\tau_i]$$

(where the derivative in 0 should be interpreted as the derivative from the left), and thus also  $(\bar{L}_n^\zeta)'(\eta) = \frac{1}{n} E^{\zeta, \eta}[H_n]$ . Further

$$(A.3) \quad L''(\eta) = \mathbb{E}[E^{\zeta, \eta}[H_1^2] - E^{\zeta, \eta}[H_1]^2] > 0,$$

$$(A.4) \quad (L_i^\zeta)''(\eta) = (E^{\zeta, \eta}[\tau_i^2] - E^{\zeta, \eta}[\tau_i]^2) > 0,$$

and thus also  $(\bar{L}_n^\zeta)''(\eta) = \frac{1}{n}(E^{\zeta,\eta}[H_n^2] - E^{\zeta,\eta}[H_n]^2)$ . Moreover, for every  $\Delta \subset (-\infty, 0)$  compact, there is  $c(\Delta) \in (0, \infty)$  such that

$$\sup_{\eta \in \Delta} \text{ess sup} |L_i^\zeta(\eta)| \leq c(\Delta),$$

and analogous statements hold for  $(L_i^\zeta)'$  and  $(L_i^\zeta)''$  as well.

PROOF. The fact that  $L$  and  $\bar{L}_n^\zeta$  are infinitely differentiable follows easily from the dominated convergence theorem which allows to interchange the differentiation with the expectations. The equalities (A.1) and (A.2) can then be obtained by a direct computation from the definitions of the corresponding functions. The equalities (A.3) and (A.4) follow from the definition (4.4) of  $P^{\zeta,\eta}$ . The strict inequalities in (A.3) and (A.4) follow from the fact that, as  $H_1, \tau_i$  are nondegenerate random variables, Jensen's inequality provides us with a strict inequality.

To prove the last claim, we start with observing that  $\zeta \mapsto L_i^\zeta$  and  $\zeta \mapsto (L_i^\zeta)'$  are increasing as mappings from  $[\text{ei} - \text{es}, 0]^\mathbb{Z}$  to  $(-\infty, 0]$ , and  $\zeta(x) \in [\text{ei} - \text{es}, 0]$   $\mathbb{P}$ -a.s. Therefore,

$$\begin{aligned} -\infty < \ln E_{i-1}[e^{H_i(\text{ei}-\text{es}+\min \Delta)}] &\leq \inf_{\eta \in \Delta} \text{ess inf} L_i^\zeta(\eta) \\ &\leq \sup_{\eta \in \Delta} \text{ess sup} L_i^\zeta(\eta) \leq \ln E_{i-1}[e^{H_i \max \Delta}] < \infty, \end{aligned}$$

where the finiteness of the expectations on the left- and right-hand side follows easily since both  $\text{ei} - \text{es} + \min \Delta$  and  $\max \Delta$  are negative. The proof for  $(L_i^\zeta)'$  is similar. For  $(L_i^\zeta)''$  we use (A.4) as well as the definition of  $E^{\zeta,\eta}$  to observe that for  $\eta \in \Delta$ ,

$$\begin{aligned} \sup_{\eta \in \Delta} \text{ess sup} |(L_i^\zeta)''(\eta)| &\leq \sup_{\eta \in \Delta} \text{ess sup} E^{\zeta,\eta}[\tau_i^2] \\ &\leq \frac{E_{i-1}[H_i^2 e^{H_i \max \Delta}]}{E_{i-1}[e^{H_i(\min \Delta + \text{ei} - \text{es})}]} < \infty, \end{aligned}$$

which finishes the proof.  $\square$

LEMMA A.2. Let  $\mathcal{F}_k = \sigma(\xi(i) : i \leq k)$  and  $\Delta$  be a compact interval in  $(-\infty, 0)$ . Then there exists a constant  $C_\Delta < \infty$  such that for all  $0 \leq i < j$  and  $\eta \in \Delta$ ,  $\mathbb{P}$ -a.s.,

$$(A.5) \quad |\mathbb{E}[L_j^\zeta(\eta) \mid \mathcal{F}_i] - L(\eta)| \leq C_\Delta e^{-(j-i)/C_\Delta},$$

$$(A.6) \quad 0 \leq (\text{ess sup}_{\zeta(k):k \leq i} L_j^\zeta(\eta)) - L_j^\zeta(\eta) \leq C_\Delta e^{-(j-i)/C_\Delta},$$

and similarly

$$(A.7) \quad |\mathbb{E}[(L_j^\zeta)'(\eta) \mid \mathcal{F}_i] - L'(\eta)| \leq C_\Delta e^{-(j-i)/C_\Delta},$$

$$(A.8) \quad 0 \leq (\text{ess sup}_{\zeta(k):k \leq i} (L_j^\zeta)'(\eta)) - (L_j^\zeta)'(\eta) \leq C_\Delta e^{-(j-i)/C_\Delta}.$$

PROOF. We only prove the inequalities (A.5) and (A.6), the remaining ones being derived in a similar manner.

By translation invariance, we may assume without loss of generality  $0 = i < j$ . Write  $L_j^\zeta(\eta) = \ln(A + B)$  where

$$A = E_{j-1} \left[ \exp \left\{ \int_0^{H_j} (\zeta(X_s) + \eta) ds \right\}, \inf_{0 \leq s \leq H_j} X_s > 0 \right],$$

$$B = E_{j-1} \left[ \exp \left\{ \int_0^{H_j} (\zeta(X_s) + \eta) ds \right\}, \inf_{0 \leq s \leq H_j} X_s \leq 0 \right].$$

Let  $K_t$  denote the number of jumps that the random walk  $(X_n)$  has made up to time  $t > 0$ , which has Poisson distribution with parameter  $t$ . Then, since  $\text{ess sup } \zeta = 0$ , for  $\delta > 0$  sufficiently small, uniformly over  $\eta \in \Delta$ ,

$$B \leq E_{j-1} [e^{\eta H_j}; H_j \geq \delta j] + P_{j-1}(K_{\delta j} \geq j) \leq ce^{-j/c},$$

where the last inequality follows from standard large deviations for the Poisson random variable. On the other hand, due to (4.3), there is  $c' \in (0, 1)$  such that  $\mathbb{P}$ -a.s. one has  $A \in (c', 1)$ . Therefore, since  $\ln(1+x) \leq x$ , we infer that  $\mathbb{P}$ -a.s.

$$\ln(A) \leq L_j^\zeta(\eta) \leq \ln(A) + \ln \left( 1 + \frac{B}{A} \right) \leq \ln(A) + ce^{-j/c}.$$

Since  $\ln(A)$  is independent of  $\mathcal{F}_0$  by definition, taking the essential supremum over  $\zeta(k)$ ,  $k \leq 0$ , this implies the second inequality of (A.6). The first inequality of (A.6) follows from the definitions. Using the independence of  $\ln(A)$  and  $\mathcal{F}_0$  once again, we also infer that

$$|\mathbb{E}[L_j^\zeta(\eta) - L(\eta) \mid \mathcal{F}_0]| \leq |\mathbb{E}[\ln(A) - \mathbb{E}[\ln(A)] \mid \mathcal{F}_0]| + 2ce^{-j/c} = 2ce^{-j/c},$$

which completes the proof of the lemma.  $\square$

**A.2. Basic properties of the Lyapunov exponent.** We prove here various properties of the Lyapunov exponent  $\lambda$  defined in (2.2) that are used throughout the paper. Some of these properties are standard, but for some of them we did not find any reference. In particular, the proof of  $v_c > 0$  is presumably new and of independent interest.

PROPOSITION A.3. *Assume (POT).*

(a) *The function  $\lambda : \mathbb{R} \rightarrow \mathbb{R}$  is well defined, nonrandom, even and concave. It satisfies  $\lambda(0) = \text{es}$ ,  $\lambda(v) < \text{es}$  for every  $v \neq 0$ , and  $\lim_{v \rightarrow \infty} \lambda(v)/v = -\infty$ . In particular, there exists a unique  $v_0 \in (0, \infty)$  such that  $\lambda(v_0) = 0$ .*

(b) *There is  $v_c \in (0, \infty)$  given by  $v_c = (L'(0))^{-1}$  (where the derivative is taken from the left only) such that  $\lambda$  is linear on  $[0, v_c]$ , and strictly concave on  $(v_c, \infty)$ . In addition, for every  $v \in [0, \infty)$ ,*

$$(A.9) \quad \lambda(v) = \text{es} - vL^*(1/v),$$

where for  $v = 0$  the right-hand side is defined as  $\text{es}$ .

PROOF. (a) For  $\alpha \in (0, 1)$  and  $v_1, v_2 \in \mathbb{R}$ , the Markov property yields

$$(A.10) \quad \begin{aligned} & \ln E_0 [e^{\int_0^t \xi(X_s) ds}; X_t = \lfloor (\alpha v_1 + (1-\alpha)v_2)t \rfloor] \\ & \geq \ln E_0 [e^{\int_0^{(1-\alpha)t} \xi(X_s) ds}; X_{(1-\alpha)t} = \lfloor (1-\alpha)v_2 t \rfloor] \\ & \quad + \ln E_{\lfloor (1-\alpha)v_2 t \rfloor} [e^{\int_0^{\alpha t} \xi(X_s) ds}; X_{\alpha t} = \lfloor (\alpha v_1 + (1-\alpha)v_2)t \rfloor]. \end{aligned}$$

Hence, choosing  $v_1 := v_2 := v$  and using Kingman’s subadditive ergodic theorem [47] as well as the Feynman–Kac formula (Proposition 3.1), we obtain that for each  $v \in \mathbb{R}$ , the limit  $\lambda(v)$  exists and is nonrandom. In addition,  $\lambda$  is an even function since  $X$  is symmetric simple random walk and the  $(\xi(x))_{x \in \mathbb{Z}}$  are i.i.d. by assumption.

Dividing both sides of inequality (A.10) by  $t$  and taking the limit  $t \rightarrow \infty$ , the left-hand side converges  $\mathbb{P}$ -a.s. to  $\lambda(\alpha v_1 + (1 - \alpha)v_2)$ , and the first summand on the right-hand side converges to  $(1 - \alpha)\lambda(v_2)$ . Further, the second summand converges to  $\alpha\lambda(v_1)$  in distribution, since it has the same distribution (up to possibly a small error introduced by the use of the floor function, and which is irrelevant in the limit) as

$$\frac{1}{t} \ln E_0 \left[ \exp \left\{ \int_0^{\alpha t} \xi(X_s) ds \right\} \mathbf{1}_{X_{\alpha t} = \lfloor \alpha v_1 t \rfloor} \right],$$

which converges  $\mathbb{P}$ -a.s. to  $\alpha\lambda(v_1)$ . The concavity of  $\lambda$  then follows.

The proof of  $\lambda(0) = \text{es}$  is standard but we include it for the sake of completeness. By the Feynman–Kac formula and (4.2),

$$(A.11) \quad \lambda(0) = \text{es} + \lim_{t \rightarrow \infty} \frac{1}{t} \ln E_0 \left[ \exp \left\{ \int_0^t \zeta(X_s) ds \right\}; X_t = 0 \right].$$

Since  $\zeta(x) \leq 0$ , the upper bound  $\lambda(0) \leq \text{es}$  follows trivially. To show the lower bound, fix  $\varepsilon > 0$  arbitrarily and note that by standard i.i.d. properties of  $\zeta$ ’s, there is  $c(\varepsilon) > 0$  such that  $\mathbb{P}$ -a.s. for  $t$  large enough, there is an interval  $I_t \subset [-t^{1/4}, t^{1/4}]$  of length at least  $c(\varepsilon) \ln t$  such  $\zeta(j) \geq -\varepsilon$  for all  $j \in I_t \cap \mathbb{Z}$ . Consider now the event  $\mathcal{A}_t = \{X_0 = X_t = 0, X_s \in I_t \ \forall s \in [t^{1/2}, t - t^{1/2}]\}$ . By a local central limit theorem,  $P_0(X_{t^{1/2}} \in I_t) \geq ct^{-1/4}$ . By standard spectral estimates for the simple random walk, for any  $m \in I_t$ ,

$$P_m(X_s \in I_t \ \forall s \leq t - 2t^{1/2}) \geq e^{-ct/\ln t},$$

and, by a local central limit theorem again,  $P_m(X_{t^{1/2}} = 0) \geq ct^{-1/4}$ . The Markov property thus yields  $P_0(\mathcal{A}_t) \geq e^{-ct/\ln t}$ . Going back to (A.11), restricting the expectation to  $\mathcal{A}_t$ ,

$$\lambda(0) \geq \text{es} + \frac{1}{t} \limsup_{t \rightarrow \infty} P_0(\mathcal{A}_t) e^{-2\text{es}t^{1/2}} e^{-\varepsilon(t-2t^{1/2})} \geq \text{es} - \varepsilon.$$

Since  $\varepsilon > 0$  was chosen arbitrarily,  $\lambda(0) = \text{es}$  follows.

The fact  $\lim_{v \rightarrow \infty} \lambda(v)/|v| = -\infty$  follows from (POT) and large deviation properties of the continuous time simple random walk  $X$ .

(b) The strict concavity of  $\lambda(v)$  on  $(v_c, \infty)$  is a consequence of the strict convexity of  $L^*(1/v)$  on this interval, which in turn follows from definition (4.12) of  $v_c$ , the strict convexity of  $L$  on  $(-\infty, 0)$  and standard properties of the Legendre transform. Also, for  $v \in (v_c, \infty)$ , claim (A.9) is shown in the proof of Theorem 2.8 in Section 4.6. By the continuity of  $\lambda$  (which follows from concavity and finiteness) and the monotonicity and lower-semicontinuity of  $L^*$  (which entails its left-continuity in  $1/v_c$ ), (A.9) also holds for  $v = v_c$ . We thus only need to show the linearity of  $\lambda$  and (A.9) on  $[0, v_c]$ , and then  $v_c \in (0, \infty)$ .

To show the linearity, observe that by the Feynman–Kac representation,

$$\begin{aligned} u(t, vt) &= e^{t\text{es}} E_0 \left[ \exp \left\{ \int_0^t \zeta(X_s) ds \right\}; X_t = vt \right] \\ &\leq e^{t\text{es}} E_0 \left[ \prod_{i=1}^{vt-1} \exp \left\{ \int_{H_{i-1}}^{H_i} \zeta(X_s) ds \right\} \right] = e^{t\text{es}} e^{\sum_{i=1}^{vt} L_i^\zeta(0)}. \end{aligned}$$

Taking logarithms and letting  $t \rightarrow \infty$ , it follows that

$$(A.12) \quad \lambda(v) \leq \text{es} + vL(0) = \text{es} - vL^*(1/v_c), \quad v \in [0, v_c],$$

where for the inequality we used (4.10), and for the equality we took advantage of (4.12) again. The concavity of  $\lambda$ , the linear upper bound (A.12), and the fact that  $\lambda$  coincides with this linear upper bound for  $v = 0$  (cf. part (a)) and  $v = v_c$  (by (A.9) for  $v = v_c$ ) then imply the matching lower bound, proving the claimed linearity on  $[0, v_c]$ . Claim (A.9) for  $v \in [0, v_c]$  then follows directly, since we know that the inequality in (A.12) is an equality, and  $L(0) = -L^*(1/v) = L^*(1/v_c)$  for  $v \in [0, v_c]$ , by standard properties of Legendre transform.

It remains to show that  $(L'(0))^{-1} \in (0, \infty)$ . We recall that by (A.1),  $L'(0) = \mathbb{E}[E^{\zeta, 0}[H_1]]$ . Taking advantage of the boundedness from below of  $\zeta$  and the definition of  $E^{\zeta, 0}$ , it is sufficient to show that

$$(A.13) \quad \mathbb{E}[E_0[H_1 e^{\int_0^{H_1} \zeta(X_s) ds}]] \in (0, \infty).$$

The lower bound follows easily, as  $H_1$  is a nontrivial, nonnegative random variable. For the upper bound, for arbitrary fixed  $h \in (0, \text{es} - \text{ei})$  we introduce an auxiliary random environment

$$(A.14) \quad \zeta^*(x) := \begin{cases} 0 & \text{if } \zeta(x) \in (-h, 0], \\ -h & \text{if } \zeta(x) \leq -h, \end{cases} \quad x \in \mathbb{Z}.$$

In particular, note that  $\zeta \leq \zeta^*$  and

$$(A.15) \quad p := \mathbb{P}(\zeta^*(0) = -h) = 1 - \mathbb{P}(\zeta^*(0) = 0) \in (0, 1).$$

Furthermore, defining for  $n \in \mathbb{N}$  the events

$$G_1(n) := \left\{ \inf_{s \in [0, H_1)} X_s \in (-n^{\frac{2}{3}}, -n^{\frac{1}{3}}) \right\},$$

we infer by standard large deviation estimates for simple random walk that there exist constants  $c, C \in (0, \infty)$  such that

$$(A.16) \quad P_0(H_1 \in [n, n + 1), G_1(n)^c) \leq C e^{-cn^{\frac{1}{3}}} \quad \forall n \in \mathbb{N}_0.$$

The next ingredient is [3], Theorem 1.3, which implies that thin points for the random walk are rare in the following sense: Write  $\ell$  for the local time process

$$(A.17) \quad \ell_t(x) := \int_0^t \delta_x(X_s) ds, \quad x \in \mathbb{Z}, t \in [0, \infty),$$

and, for  $M \in (0, \infty)$ , introduce the set of *thin points* by

$$\mathcal{T}_{t, M} := \{x \in \mathbb{Z} : \ell_t(x) \in (0, M)\}.$$

Then [3], Theorem 1.3, entails that setting  $G_2(t) := \{|\mathcal{T}_{t, M}| \leq t^{\frac{1}{6}}\}$ , there exist constants  $c, C \in (0, \infty)$  such that

$$(A.18) \quad P_0(G_2(H_1)^c \mid H_1 \in [n, n + 1)) \leq C e^{-cn^{\frac{1}{7}}} \quad \forall n \in \mathbb{N}_0.$$

The last ingredient is a simple large deviation bound for i.i.d. Bernoulli variables: recalling  $p$  from (A.15) and setting

$$G_3(n) := \left\{ \left| \{x \in \{-n, \dots, 0\} : \zeta^*(x) = -h\} \right| \geq \frac{pn}{2} \right\},$$

we have due to (POT) that for arbitrary  $\varepsilon > 0$ ,

$$(A.19) \quad \mathbb{P}(G_3(n)^c) \leq C e^{-n(I_p(p/2) - \varepsilon)} \leq C e^{-c_p n} \quad \forall n \in \mathbb{N},$$

where  $I_p$  is the usual rate function of the Bernoulli( $p$ ) distribution. Combining (A.16) to (A.19), we infer that

$$\begin{aligned}
 & \mathbb{E}[E_0[H_1 e^{\int_0^{H_1} \zeta(X_s) ds}]] \\
 & \leq \sum_{n=0}^{\infty} \mathbb{E}[E_0[H_1 e^{\int_0^{H_1} \zeta(X_s) ds}, H_1 \in [n, n+1)], G_3(n^{\frac{1}{3}})] \\
 & \quad + C(n+1)e^{-cn^{\frac{1}{3}}} \\
 (A.20) \quad & \leq \sum_{n=0}^{\infty} \mathbb{E}[E_0[H_1 e^{\int_0^{H_1} \zeta(X_s) ds}, H_1 \in [n, n+1), G_1(n), G_2(H_1)], G_3(n^{\frac{1}{3}})] \\
 & \quad + C(p)e^{-cn^{\frac{1}{7}}} \\
 & \leq C(p) + \sum_{n=0}^{\infty} e^{-phn^{\frac{1}{3}}/2} < \infty,
 \end{aligned}$$

and the upper bound for (A.13) follows.  $\square$

Finally, we shortly discuss the existence of random environments which satisfy the condition (VEL) requiring that the speed of the maximum particle,  $v_0$  is strictly larger than the critical speed  $v_c$ . The simple proof of the following result reveals that this is the case for a very rich family of environments, which heuristically can be interpreted as exhibiting sufficiently strong branching.

LEMMA A.4. *There exist environments  $\xi$  such that (POT) and (VEL) hold true.*

PROOF. Choose an arbitrary random environment  $\xi$  fulfilling (POT), and consider a family of environments  $\xi^h := (\xi(x) + h)_{x \in \mathbb{Z}}$ ,  $h \geq 0$ . Writing  $\lambda^h$  for the Lyapunov exponent associated to  $\xi^h$ , the Feynman–Kac representation (Proposition 3.1) and the definition (2.2) of  $\lambda$  yield that  $\lambda^h(v) = \lambda(v) + h$ . Hence, by Proposition A.3, the value of  $v_c$  does not change with  $h$ , and on the other hand,  $v_0 \rightarrow \infty$  as  $h \rightarrow \infty$ , which entails the desired statement.  $\square$

**A.3. Hoeffding-type inequality for mixing sequences.** We repeatedly make use of the following concentration inequality for mixing sequences. We state it here for reader’s convenience.

LEMMA A.5 ([62], Theorem 2.4). *Let  $(X_i)_{i \in \mathbb{Z}}$  be a sequence of real valued bounded random variables on some  $(\Omega, \mathcal{F}, \mathbb{P})$  and let  $\mathcal{F}_i = \sigma(X_j, j \leq i)$ . Suppose that there are real numbers  $m_i > 0$ ,  $i \in \{1, \dots, n\}$  such that*

$$\sup_{j \in \{i+1, \dots, n\}} \left( \|X_i^2\|_{\infty} + 2 \left\| X_i \sum_{k=i+1}^j \mathbb{E}[X_k | \mathcal{F}_i] \right\|_{\infty} \right) \leq m_i \quad \text{for all } i \leq n.$$

Then for every  $a > 0$ ,

$$\mathbb{P}(|X_1 + \dots + X_n| \geq a) \leq \sqrt{e} \exp \left\{ -\frac{a^2}{2 \sum_{i=1}^n m_i} \right\}.$$

**Acknowledgments.** The authors would like to thank the Columbia University Mathematics Department as well as the Mathematics Department of the University of Vienna for their hospitality and financial support. Part of this work has been completed during a stay at the *Random Geometry* semester programme of the Isaac Newton Institute for Mathematical Sciences in Cambridge, UK, in 2015. The authors would like to thank an anonymous referee as well as Lars Schmitz for carefully reading the manuscript and for valuable comments.

The second author was supported by the UoC Forum *Classical and quantum dynamics of interacting particle systems* of the Institutional Strategy of the University of Cologne within the German Excellence Initiative.

## REFERENCES

- [1] ADDARIO-BERRY, L. and REED, B. (2009). Minima in branching random walks. *Ann. Probab.* **37** 1044–1079. MR2537549 <https://doi.org/10.1214/08-AOP428>
- [2] AÏDÉKON, E. (2013). Convergence in law of the minimum of a branching random walk. *Ann. Probab.* **41** 1362–1426. MR3098680 <https://doi.org/10.1214/12-AOP750>
- [3] ATHREYA, S., DREWITZ, A. and SUN, R. (2017). Subdiffusivity of a random walk among a Poisson system of moving traps on  $\mathbb{Z}$ . *Math. Phys. Anal. Geom.* **20** Art. 1, 22. MR3590740 <https://doi.org/10.1007/s11040-016-9227-8>
- [4] BACHMANN, M. (2000). Limit theorems for the minimal position in a branching random walk with independent logconcave displacements. *Adv. in Appl. Probab.* **32** 159–176. MR1765165 <https://doi.org/10.1239/aap/1013540028>
- [5] BERESTYCKI, J., BRUNET, É., HARRIS, J. W., HARRIS, S. C. and ROBERTS, M. I. (2015). Growth rates of the population in a branching Brownian motion with an inhomogeneous breeding potential. *Stochastic Process. Appl.* **125** 2096–2145. MR3315624 <https://doi.org/10.1016/j.spa.2014.12.008>
- [6] BHATTACHARYA, R. N. and RANGA RAO, R. (1976). *Normal Approximation and Asymptotic Expansions*. Wiley Series in Probability and Mathematical Statistics. Wiley, New York. MR0436272
- [7] BIGGINS, J. D. (1976). The first- and last-birth problems for a multiplicity age-dependent branching process. *Adv. in Appl. Probab.* **8** 446–459. MR0420890 <https://doi.org/10.2307/1426138>
- [8] BOVIER, A. (2017). *Gaussian Processes on Trees: From Spin Glasses to Branching Brownian Motion*. Cambridge Studies in Advanced Mathematics **163**. Cambridge Univ. Press, Cambridge. MR3618123 <https://doi.org/10.1017/9781316675779>
- [9] BOVIER, A. and HARTUNG, L. (2014). The extremal process of two-speed branching Brownian motion. *Electron. J. Probab.* **19** no. 18, 28. MR3164771 <https://doi.org/10.1214/EJP.v19-2982>
- [10] BOVIER, A. and HARTUNG, L. (2015). Variable speed branching Brownian motion 1. Extremal processes in the weak correlation regime. *ALEA Lat. Am. J. Probab. Math. Stat.* **12** 261–291. MR3351476
- [11] BRAMSON, M. (1983). Convergence of solutions of the Kolmogorov equation to travelling waves. *Mem. Amer. Math. Soc.* **44** iv+190. MR0705746 <https://doi.org/10.1090/memo/0285>
- [12] BRAMSON, M. and ZEITOUNI, O. (2009). Tightness for a family of recursion equations. *Ann. Probab.* **37** 615–653. MR2510018 <https://doi.org/10.1214/08-AOP414>
- [13] BRAMSON, M. and ZEITOUNI, O. (2012). Tightness of the recentered maximum of the two-dimensional discrete Gaussian free field. *Comm. Pure Appl. Math.* **65** 1–20. MR2846636 <https://doi.org/10.1002/cpa.20390>
- [14] BRAMSON, M. D. (1978). Maximal displacement of branching Brownian motion. *Comm. Pure Appl. Math.* **31** 531–581. MR0494541 <https://doi.org/10.1002/cpa.3160310502>
- [15] CHAUVIN, B. and DRMOTA, M. (2006). The random multisection problem, travelling waves and the distribution of the height of  $m$ -ary search trees. *Algorithmica* **46** 299–327. MR2291958 <https://doi.org/10.1007/s00453-006-0107-7>
- [16] COMETS, F. and POPOV, S. (2007). On multidimensional branching random walks in random environment. *Ann. Probab.* **35** 68–114. MR2303944 <https://doi.org/10.1214/009117906000000926>
- [17] COMETS, F. and POPOV, S. (2007). Shape and local growth for multidimensional branching random walks in random environment. *ALEA Lat. Am. J. Probab. Math. Stat.* **3** 273–299. MR2365644
- [18] DEMBO, A. and ZEITOUNI, O. (1998). *Large Deviations Techniques and Applications*, 2nd ed. *Applications of Mathematics (New York)* **38**. Springer, New York. MR1619036 <https://doi.org/10.1007/978-1-4612-5320-4>
- [19] DREWITZ, A. (2008). Lyapunov exponents for the one-dimensional parabolic Anderson model with drift. *Electron. J. Probab.* **13** 2283–2336. MR2469612 <https://doi.org/10.1214/EJP.v13-586>



- [20] DRMOTA, M. (2003). An analytic approach to the height of binary search trees. II. *J. ACM* **50** 333–374. MR2146358 <https://doi.org/10.1145/765568.765572>
- [21] FANG, M. and ZEITOUNI, O. (2012). Branching random walks in time inhomogeneous environments. *Electron. J. Probab.* **17** no. 67, 18. MR2968674 <https://doi.org/10.1214/EJP.v17-2253>
- [22] FANG, M. and ZEITOUNI, O. (2012). Slowdown for time inhomogeneous branching Brownian motion. *J. Stat. Phys.* **149** 1–9. MR2981635 <https://doi.org/10.1007/s10955-012-0581-z>
- [23] FREIDLIN, M. (1985). *Functional Integration and Partial Differential Equations. Annals of Mathematics Studies* **109**. Princeton Univ. Press, Princeton, NJ. MR0833742 <https://doi.org/10.1515/9781400881598>
- [24] GANTERT, N., MÜLLER, S., POPOV, S. and VACHKOVSKAIA, M. (2010). Survival of branching random walks in random environment. *J. Theoret. Probab.* **23** 1002–1014. MR2735734 <https://doi.org/10.1007/s10959-009-0227-5>
- [25] GÄRTNER, J. and FREIDLIN, M. I. (1979). The propagation of concentration waves in periodic and random media. *Dokl. Akad. Nauk SSSR* **249** 521–525. MR0553200
- [26] GÄRTNER, J. and KÖNIG, W. (2005). The parabolic Anderson model. In *Interacting Stochastic Systems* 153–179. Springer, Berlin. MR2118574 [https://doi.org/10.1007/3-540-27110-4\\_8](https://doi.org/10.1007/3-540-27110-4_8)
- [27] GÄRTNER, J. and MOLCHANOV, S. A. (1990). Parabolic problems for the Anderson model. I. Intermittency and related topics. *Comm. Math. Phys.* **132** 613–655. MR1069840
- [28] GIKHMAN, I. I. and SKOROKHOD, A. V. (1969). *Introduction to the Theory of Random Processes. Translated from the Russian by Scripta Technica, Inc. W. B. Saunders Co., Philadelphia.* MR0247660
- [29] GREVEN, A. and DEN HOLLANDER, F. (1992). Branching random walk in random environment: Phase transitions for local and global growth rates. *Probab. Theory Related Fields* **91** 195–249. MR1147615 <https://doi.org/10.1007/BF01291424>
- [30] GÜN, O., KÖNIG, W. and SEKULOVIĆ, O. (2013). Moment asymptotics for branching random walks in random environment. *Electron. J. Probab.* **18** no. 63, 18. MR3078022 <https://doi.org/10.1214/ejp.v18-2212>
- [31] HAMEL, F., NOLEN, J., ROQUEJOFFRE, J.-M. and RYZHIK, L. (2016). The logarithmic delay of KPP fronts in a periodic medium. *J. Eur. Math. Soc. (JEMS)* **18** 465–505. MR3463416 <https://doi.org/10.4171/JEMS/595>
- [32] HAMMERSLEY, J. M. (1974). Postulates for subadditive processes. *Ann. Probab.* **2** 652–680. MR0370721 <https://doi.org/10.1214/aop/1176996611>
- [33] HARRIS, S. C. and ROBERTS, M. I. (2017). The many-to-few lemma and multiple spines. *Ann. Inst. Henri Poincaré Probab. Stat.* **53** 226–242. MR3606740 <https://doi.org/10.1214/15-AIH714>
- [34] HEIL, H., NAKASHIMA, M. and YOSHIDA, N. (2011). Branching random walks in random environment are diffusive in the regular growth phase. *Electron. J. Probab.* **16** 1316–1340. MR2827461 <https://doi.org/10.1214/EJP.v16-922>
- [35] HU, Y. and SHI, Z. (2009). Minimal position and critical martingale convergence in branching random walks, and directed polymers on disordered trees. *Ann. Probab.* **37** 742–789. MR2510023 <https://doi.org/10.1214/08-AOP419>
- [36] HU, Y. and YOSHIDA, N. (2009). Localization for branching random walks in random environment. *Stochastic Process. Appl.* **119** 1632–1651. MR2513122 <https://doi.org/10.1016/j.spa.2008.08.005>
- [37] HUANG, C. and LIU, Q. (2014). Branching random walk with a random environment in time. Preprint. Available at arXiv:1407.7623.
- [38] IKEDA, N., NAGASAWA, M. and WATANABE, S. (1968). Branching Markov processes. I. *J. Math. Kyoto Univ.* **8** 233–278. MR0232439 <https://doi.org/10.1215/kjm/1250524137>
- [39] IKEDA, N., NAGASAWA, M. and WATANABE, S. (1968). Branching Markov processes. II. *J. Math. Kyoto Univ.* **8** 365–410. MR0238401 <https://doi.org/10.1215/kjm/1250524059>
- [40] IKEDA, N., NAGASAWA, M. and WATANABE, S. (1969). Branching Markov processes. III. *J. Math. Kyoto Univ.* **9** 95–160. MR0246376 <https://doi.org/10.1215/kjm/1250524013>
- [41] KEVEI, P. and MASON, D. M. (2011). A note on a maximal Bernstein inequality. *Bernoulli* **17** 1054–1062. MR2817617 <https://doi.org/10.3150/10-BEJ304>
- [42] KINGMAN, J. F. C. (1975). The first birth problem for an age-dependent branching process. *Ann. Probab.* **3** 790–801. MR0400438 <https://doi.org/10.1214/aop/1176996266>
- [43] KOLMOGOROV, A., PRETROVSKII, I. and PISKUNOV, N. (1937). Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique. *Bull. Univ. État Moscou, Sér. Int., Sect. A: Math. et Mécan.* **1**, Fasc. **6** 1–25.
- [44] KÖNIG, W. (2016). *The Parabolic Anderson Model. Pathways in Mathematics.* Birkhäuser/Springer, Cham. MR3526112 <https://doi.org/10.1007/978-3-319-33596-4>
- [45] LALLEY, S. and SELLKE, T. (1988). Traveling waves in inhomogeneous branching Brownian motions. I. *Ann. Probab.* **16** 1051–1062. MR0942755



- [46] LALLEY, S. and SELLKE, T. (1989). Travelling waves in inhomogeneous branching Brownian motions. II. *Ann. Probab.* **17** 116–127. [MR0972775](#)
- [47] LIGGETT, T. M. (1985). An improved subadditive ergodic theorem. *Ann. Probab.* **13** 1279–1285. [MR0806224](#)
- [48] MACHADO, F. P. and POPOV, S. Y. (2000). One-dimensional branching random walks in a Markovian random environment. *J. Appl. Probab.* **37** 1157–1163. [MR1808881](#) <https://doi.org/10.1017/s0021900200018350>
- [49] MAILLARD, P. and ZEITOUNI, O. (2016). Slowdown in branching Brownian motion with inhomogeneous variance. *Ann. Inst. Henri Poincaré Probab. Stat.* **52** 1144–1160. [MR3531703](#) <https://doi.org/10.1214/15-AIHP675>
- [50] MALLEIN, B. (2015). Maximal displacement of a branching random walk in time-inhomogeneous environment. *Stochastic Process. Appl.* **125** 3958–4019. [MR3373310](#) <https://doi.org/10.1016/j.spa.2015.05.011>
- [51] MALLEIN, B. and MIŁOŚ, P. (2018). Brownian motion and random walk above quenched random wall. *Ann. Inst. Henri Poincaré Probab. Stat.* **54** 1877–1916. [MR3865661](#) <https://doi.org/10.1214/17-AIHP859>
- [52] MALLEIN, B. and MIŁOŚ, P. (2019). Maximal displacement of a supercritical branching random walk in a time-inhomogeneous random environment. *Stochastic Process. Appl.* **129** 3239–3260. [MR3985561](#) <https://doi.org/10.1016/j.spa.2018.09.008>
- [53] MCDIARMID, C. (1995). Minimal positions in a branching random walk. *Ann. Appl. Probab.* **5** 128–139. [MR1325045](#)
- [54] MCKEAN, H. P. (1975). Application of Brownian motion to the equation of Kolmogorov–Petrovskii–Piskunov. *Comm. Pure Appl. Math.* **28** 323–331. [MR0400428](#) <https://doi.org/10.1002/cpa.3160280302>
- [55] MERLEVÈDE, F., PELIGRAD, M. and UTEV, S. (2006). Recent advances in invariance principles for stationary sequences. *Probab. Surv.* **3** 1–36. [MR2206313](#) <https://doi.org/10.1214/154957806100000202>
- [56] NAKASHIMA, M. (2011). Almost sure central limit theorem for branching random walks in random environment. *Ann. Appl. Probab.* **21** 351–373. [MR2759206](#) <https://doi.org/10.1214/10-AAP699>
- [57] NEWMAN, C. M. and WRIGHT, A. L. (1981). An invariance principle for certain dependent sequences. *Ann. Probab.* **9** 671–675. [MR0624694](#)
- [58] NOLEN, J. (2011). A central limit theorem for pulled fronts in a random medium. *Netw. Heterog. Media* **6** 167–194. [MR2806072](#) <https://doi.org/10.3934/nhm.2011.6.167>
- [59] NOLEN, J., ROQUEJOFFRE, J.-M. and RYZHIK, L. (2015). Power-like delay in time inhomogeneous Fisher–KPP equations. *Comm. Partial Differential Equations* **40** 475–505. [MR3285242](#) <https://doi.org/10.1080/03605302.2014.972744>
- [60] ORTGIESE, M. and ROBERTS, M. I. (2016). Intermittency for branching random walk in Pareto environment. *Ann. Probab.* **44** 2198–2263. [MR3502604](#) <https://doi.org/10.1214/15-AOP1021>
- [61] REED, B. (2003). The height of a random binary search tree. *J. ACM* **50** 306–332. [MR2146357](#) <https://doi.org/10.1145/765568.765571>
- [62] RIO, E. (2013). Inequalities and limit theorems for weakly dependent sequences. Lecture.
- [63] SHI, Z. (2015). *Branching Random Walks. Lecture Notes in Math.* **2151**. Springer, Cham. [MR3444654](#) <https://doi.org/10.1007/978-3-319-25372-5>