

ERGODIC POISSON SPLITTINGS

BY ÉLISE JANVRESSE¹, EMMANUEL ROY² AND THIERRY DE LA RUE³

¹Laboratoire Amiénois de Mathématique Fondamentale et Appliquée, CNRS-UMR 7352, Université de Picardie Jules Verne, Elise.Janvresse@u-picardie.fr

²Laboratoire Analyse, Géométrie et Applications, Université Paris 13, roy@math.univ-paris13.fr

³Laboratoire de Mathématiques Raphaël Salem, CNRS, Université de Rouen Normandie, Thierry.de-la-Rue@univ-rouen.fr

In this paper, we study splittings of a Poisson point process which are equivariant under a conservative transformation. We show that, if the Cartesian powers of this transformation are all ergodic, the only ergodic splitting is the obvious one, that is, a collection of independent Poisson processes. We apply this result to the case of a marked Poisson process: under the same hypothesis, the marks are necessarily independent of the point process and i.i.d. Under additional assumptions on the transformation, a further application is derived, giving a full description of the structure of a random measure invariant under the action of the transformation.

1. Introduction. Thinning and splitting are classical operations when studying point processes. Thinning consists in removing points according to some rule, whereas the related notion of splitting means decomposing the point process as the sum of several other point processes. It is well known that thinning a Poisson point process by choosing to remove points according to independent coin tosses yields a new Poisson process of lower (but proportional) intensity. Moreover, this procedure gives rise to a splitting of the original Poisson process into a sum of two independent Poisson processes.

In recent years, some new results on thinnings of the Poisson process have emerged. In particular, it is shown in [2] that it is possible to deterministically choose points from a homogeneous Poisson point process on \mathbb{R} to get another homogeneous Poisson process of lower intensity. Moreover, it is possible to proceed in a translation equivariant way. This result has been further refined in [8] by extending it to \mathbb{R}^d and replacing translation equivariance by isometry equivariance. Moreover, the remaining points were also shown to form a homogeneous Poisson point process.

The isometry or translation equivariance plays a key role here as Meyerovitch showed in [14] that it is not possible to deterministically thin a Poisson point process in an equivariant way with respect to a transformation which is conservative ergodic on the base space (the translations of \mathbb{R}^d yield dissipativity).

In the present paper, we are also interested in thinnings and splittings of Poisson processes which are equivariant under some dynamics. The difference with [2, 8] is that we consider thinnings/splittings which are equivariant with respect to a conservative transformation. Moreover, contrary to the above-mentioned result of [14], we allow additional randomness (yet keeping ergodicity) in the thinning/splitting procedure. We get the following result, phrased in our terminology where equivariance and homogeneity are expressed through the concepts of *T-point processes* and *T-splittings* (see Section 1.1 for precise definitions): If T is an infinite measure preserving map with *infinite ergodic index* (i.e., all its Cartesian products are ergodic), then any ergodic T -splitting of a Poisson T -point process yields independent Poisson T -point processes (Theorem 2.6).

Received November 2018.

MSC2010 subject classifications. Primary 60G55, 60G57, 37A50; secondary 37A40.

Key words and phrases. Poisson point process, random measure, splitting, thinning, Poisson suspension, joinings.

In the second part of the paper, we derive some applications. In Section 3, we show that, for T with infinite ergodic index, the only way to get an ergodic marked T -point process out of a Poisson T -point process is to take i.i.d. marks, independent of the underlying process (Theorem 3.1). Then we come back to the problem we raised in [9]. In that paper, we gave conditions on T under which an ergodic T -point process with moments of all orders is necessarily a cluster-Poisson process (see [3], p. 175), best described as an independent superposition of shifted Poisson processes (a so-called *SuShi*). We extend this result in Section 4 to general random measures (Proposition 4.2 and Theorem 4.3). This more general framework allows to simplify and improve some disjointness results from [9] in the last section.

1.1. *T-Random measures and T-point processes.* Let X be a complete separable metric space and \mathcal{A} be its Borel σ -algebra. Define \tilde{X} to be the space of *boundedly finite measures* (also called *locally finite measures*) on (X, \mathcal{A}) , that is to say measures giving finite mass to any bounded Borel subset of X . We refer to [12] for the topological properties of \tilde{X} . In particular, \tilde{X} can be turned into a complete separable metric space, that we equip with its Borel σ -algebra $\tilde{\mathcal{A}}$. Convergence of a sequence (ξ_n) to ξ in \tilde{X} is characterized by the convergence

$$\int f d\xi_n \xrightarrow{n \rightarrow \infty} \int f d\xi$$

for all continuous function f on X with bounded support.

We denote by $X^* \subset \tilde{X}$ the subspace of simple counting measures, that is, whose elements are of the form

$$\xi = \sum_{i \in I} \delta_{x_i},$$

where I is at most countable, $x_i \neq x_j$ whenever $i \neq j$, and any bounded subset $A \subset X$ contains finitely many points of the family $\{x_i\}_i$. We define \mathcal{A}^* as the restriction of $\tilde{\mathcal{A}}$ to X^* .

Throughout the paper, we fix a boundedly finite and continuous measure μ on X with $\mu(X) = \infty$, and an invertible transformation T on X preserving μ . We set

$$\mathcal{A}_f := \{A \in \mathcal{A}, 0 < \mu(A) < +\infty\}.$$

For any measure ξ on X , we define $T_*(\xi)$ as the pushforward measure of ξ by T : for any $A \in \mathcal{A}$,

$$T_*(\xi)(A) := \xi(T^{-1}A).$$

In particular, if $\xi = \sum_{i \in I} \delta_{x_i}$, then $T_*(\xi) = \sum_{i \in I} \delta_{T(x_i)}$.

As we already noticed in [9], the property of bounded finiteness may be lost by the action of T . Nevertheless, if m is a σ -finite measure on \tilde{X} which is concentrated on $\bigcap_{n \in \mathbb{Z}} T_*^{-n} \tilde{X}$, it makes sense to consider the T_* -invariance of m . In this case, $T_*(\xi) \in \tilde{X}$ for m -almost all $\xi \in \tilde{X}$, and $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$ is an invertible measure preserving dynamical system.

The following definition generalizes the notion of T -point process introduced in [9] (which is formally the same, just replacing \tilde{X} by X^*).

DEFINITION 1.1. A *T-random measure* is a random variable N defined on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with values in $(\tilde{X}, \tilde{\mathcal{A}})$ such that:

- for any set $A \in \mathcal{A}$, $N(A) = 0$ \mathbb{P} -a.s. whenever $\mu(A) = 0$;
- there exists a measure preserving invertible transformation S on $(\Omega, \mathcal{F}, \mathbb{P})$, such that for any set $A \in \mathcal{A}$, $N(A) \circ S = N(T^{-1}A)$.

Thus, a T -random measure N implements a factor relationship between the dynamical systems $(\Omega, \mathcal{F}, \mathbb{P}, S)$ and $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$, where m is the pushforward measure of \mathbb{P} by N . We say that N is *ergodic* whenever $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$ is ergodic. In particular, N is ergodic as soon as $(\Omega, \mathcal{F}, \mathbb{P}, S)$ is itself ergodic.

The *intensity* of a T -random measure N is the T -invariant measure on X defined by the formula $A \in \mathcal{A} \mapsto \mathbb{E}[N(A)]$. It is absolutely continuous with respect to μ and if it is σ -finite, it is a multiple of μ , by ergodicity of (X, \mathcal{A}, μ, T) . In this case, we say that N is *integrable*. More generally, the higher order moment measures can be defined as follows.

DEFINITION 1.2. Let $n \geq 1$. A T -random measure N on $(\Omega, \mathcal{F}, \mathbb{P}, S)$ is said to have *moments of order n* if, for all bounded $A \in \mathcal{A}$, $\mathbb{E}[(N(A))^n] < +\infty$. In this case, the formula

$$M_n^N(A_1 \times \dots \times A_n) := \mathbb{E}[N(A_1) \times \dots \times N(A_n)]$$

$(A_1, \dots, A_n \in \mathcal{A})$ defines a boundedly finite $T^{\times n}$ -invariant measure M_n^N on $(X^n, \mathcal{A}^{\otimes n})$ called the *n -order moment measure*.

A T -random measure with moments of order 2 is said to be *square integrable*.

A *T -point process* is a T -random measure taking values in X^* . In this case, for $\omega \in \Omega$, we identify $N(\omega)$ with the corresponding set of points in X . The most important T -point processes are Poisson point processes; let us recall their definition.

DEFINITION 1.3. A random variable N with values in (X^*, \mathcal{A}^*) is a *Poisson point process of intensity μ* if for any $k \geq 1$, for any mutually disjoint sets $A_1, \dots, A_k \in \mathcal{A}_f$, the random variables $N(A_1), \dots, N(A_k)$ are independent and Poisson distributed with respective parameters $\mu(A_1), \dots, \mu(A_k)$.

Such a process always exists, and its distribution μ^* on X^* is uniquely determined by the preceding conditions.

Since T preserves μ , one easily checks that T_* preserves μ^* . And defining N on the probability space $(X^*, \mathcal{A}^*, \mu^*)$ as the identity map provides an example of a T -point process, the underlying measure-preserving transformation being $S = T_*$ in this case.

DEFINITION 1.4. The probability-preserving dynamical system $(X^*, \mathcal{A}^*, \mu^*, T_*)$ is called the *Poisson suspension* over the base (X, \mathcal{A}, μ, T) .

The basic result (see, e.g., [15]) about Poisson suspensions states that $(X^*, \mathcal{A}^*, \mu^*, T_*)$ is ergodic (and then weakly mixing) if and only if there is no T -invariant set in \mathcal{A}_f . In particular, this is the case if (X, \mathcal{A}, μ, T) is ergodic and μ infinite.

We also recall the classical isometry formula that will be useful several times in this paper: for $f, g \in L^1(\mu) \cap L^2(\mu)$,

$$\begin{aligned} & \mathbb{E}_{\mu^*} \left[\left(\int_X f(x) N(dx) - \int_X f(x) \mu(dx) \right) \right. \\ (1) \quad & \left. \times \left(\int_X g(x) N(dx) - \int_X g(x) \mu(dx) \right) \right] \\ & = \int_X f(x) g(x) \mu(dx). \end{aligned}$$

REMARK 1.5. The notion of T -random measure with intensity μ can be interpreted in terms of *quasifactors* as introduced by Glasner and Meyerovitch. Glasner defined in [7] a

quasifactor of a probability measure preserving system (X, \mathcal{A}, μ, T) as a probability measure preserving system $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$ where $\mathbb{E}_m[N(A)] = \mu(A)$. (Here, N is the random variable defined by the identity on \tilde{X} , and in the case where μ is a probability measure, m is in fact concentrated on the subset of probability measures on X .) Meyerovitch in [13] extended this definition to the case where μ is infinite (but m is still a probability measure). Thus:

- Poisson suspensions appear as natural example of ergodic quasifactors.
- any T -random measure N with intensity μ on $(\Omega, \mathcal{F}, \mathbb{P}, S)$ gives rise to the quasifactor defined by $m := N_*(\mathbb{P})$;
- any quasifactor $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$ is associated to the T -random measure $N := \text{Id}$ on the probability space $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$.

In Section 4.2.3, we will consider yet another case, namely when m is an infinite measure, and use the terminology ∞ -quasifactor in this case.

1.2. Splittings.

DEFINITION 1.6. Let N be a T -point process defined on the dynamical system $(\Omega, \mathcal{F}, \mathbb{P}, S)$. For $1 \leq k \leq \infty$, a T -splitting of order k of N is a finite or countable family of T -point processes $\{N_i\}_{0 \leq i < k}$ defined on $(\Omega, \mathcal{F}, \mathbb{P}, S)$ so that $N = \sum_{0 \leq i < k} N_i$.

We use the terminology “ T -splitting” in the above definition to insist on the equivariance of the splitting with respect to the underlying transformation T . In this paper, however, whenever we deal with T -point processes, the splittings are always assumed to be T -splittings, and we will omit the “ T ” in the sequel.

The splitting is said to be *ergodic* if the joining generated by $\{N_i\}_{0 \leq i < k}$ is ergodic, that is if the action of S on the factor sigma-algebra generated by $\{N_i\}_{0 \leq i < k}$ is ergodic. In the situation $N' \leq N$, the usual terminology considers N' as a *thinning* of N , and we get $(N', N - N')$ as a splitting of order 2.

A *Poisson splitting of order k* is a splitting such that $\{N_i\}_{0 \leq i < k}$ are independent Poisson processes. (In this case, N itself has to be a Poisson T -point process.)

As we mentioned in the [Introduction](#), under the assumption of conservativity and ergodicity of T , Meyerovitch [14] proved that, in the canonical space $(X^*, \mathcal{A}^*, \mu^*, T_*)$ of the Poisson suspension, there exists no splitting of the canonical Poisson T -point process. But of course a splitting can exist in a larger probability space (e.g., a product space, in which we can find two independent Poisson T -point processes).

1.3. *Properties of Cartesian powers of T* . Let us end the [Introduction](#) by saying a few words about some properties of the Cartesian products of the underlying infinite measure preserving transformation that we will be dealing with.

It is well known, in the finite measure setting, that a weak mixing transformation has infinite ergodic index (all its Cartesian powers are ergodic). Moreover, if it is not weak mixing then its Cartesian square is already not ergodic. The situation is therefore pretty clear. In the infinite measure case, however, the picture is definitely not as simple. It was first observed in [11] that all intermediate situations may occur: For any $k \geq 1$, there is a transformation with first k Cartesian products ergodic whereas the $k + 1$ Cartesian product is not. The same authors also give examples of infinite measure preserving transformations with infinite ergodic index. All these examples fall into the Markov chain category.

Since then, as the zoo of infinite measure preserving transformations developed, various examples of transformations having infinite ergodic index or not were built (see [1] where

the so-called *infinite Chacon transformation*—an infinite measure preserving version of the classical Chacon transformation—is shown to have infinite ergodic index).

In the last part of the paper, we will assume a much stronger property of T , which can be viewed as a strong version of the Radon minimal self-joinings property introduced by Danilenko in [4], and roughly saying that the Cartesian powers of the transformation admit as few invariant measures as possible (see Definition 4.1). An example of a transformation enjoying this property, the *nearly finite Chacon transformation*, is described in [10].

2. Splitting of Poisson T -point processes. For each $n \geq 1$, we denote by \mathcal{P}_n the set of all partitions of $\{1, \dots, n\}$. Given $\pi \in \mathcal{P}_n$, we define a measure on X^n by

$$m_\pi(A_1 \times \dots \times A_n) := \prod_{P \in \pi} \mu\left(\bigcap_{i \in P} A_i\right).$$

For a given n , these measures are $T^{\times n}$ -invariant and mutually singular. The measure corresponding to the trivial partition with a single atom is called the n -diagonal measure, and is concentrated on $D_n := \{(x_1, \dots, x_n) \in X^n : x_1 = \dots = x_n\}$.

It is well known that the n -order moment measure of the Poisson process of intensity μ takes the form

$$\sum_{\pi \in \mathcal{P}_n} c_\pi m_\pi$$

for positive coefficients c_π , $\pi \in \mathcal{P}_n$. Moreover, if N is a Poisson process of intensity $\alpha\mu$, then the n -order moment measure equals

$$\sum_{\pi \in \mathcal{P}_n} c_\pi \alpha^{\#\pi} m_\pi.$$

In the context of a T -point process, it turns out that the existence for each moment measure of a decomposition as a linear combination of the measures m_π characterizes Poisson processes. This is the object of the following theorem, whose proof is hidden in Theorem 3.2 in [9]. Although the argument is almost word-for-word the same, we repeat the proof here since the assumptions are far more general, and it would be cumbersome to explain the differences without giving all the details.

THEOREM 2.1. *Let (X, \mathcal{A}, μ, T) be an infinite measure preserving dynamical system with no invariant set of positive finite measure. Let N be a T -point process with moments of all orders defined on $(\Omega, \mathcal{F}, \mathbb{P}, S)$.*

Then N is a Poisson process if and only if N is ergodic and for all $n \geq 1$, there exist nonnegative numbers α_π such that

$$M_n^N = \sum_{\pi \in \mathcal{P}_n} \alpha_\pi m_\pi.$$

PROOF. Only one direction needs to be detailed.

We can assume that $(\Omega, \mathcal{F}, \mathbb{P}, S)$ is ergodic. For $n = 1$, we obtain the intensity of N as a multiple of μ , say $\alpha\mu$ for some $\alpha \geq 0$.

We first point out that for each $n \geq 1$, the weight of the n -diagonal measure is α (this is valid for any point process of intensity $\alpha\mu$). Indeed, using a set $A \in \mathcal{A}_f$, and $((A_i^\ell)_{1 \leq i \leq p_\ell})_{\ell \geq 1}$ a generating sequence of partitions of A , we get

$$\begin{aligned} \sum_{i=1}^{p_\ell} M_n^N(A_i^\ell \times \dots \times A_i^\ell) &= \mathbb{E}\left[\sum_{i=1}^{p_\ell} N(A_i^\ell) \cdots N(A_i^\ell)\right] \\ &\xrightarrow{\ell \rightarrow \infty} \mathbb{E}[N(A)] = \alpha\mu(A) = \alpha\mu(A \cap \dots \cap A). \end{aligned}$$

On the other hand,

$$\sum_{i=1}^{P\ell} M_n^N(A_i^\ell \times \cdots \times A_i^\ell) = M_n^N\left(\bigsqcup_{i=1}^{P\ell} A_i^\ell \times \cdots \times A_i^\ell\right) \xrightarrow{\ell \rightarrow \infty} M_n^N(D_n \cap A \times \cdots \times A).$$

Therefore, $M_n^N(D_n \cap A \times \cdots \times A) = \alpha\mu(A \cap \cdots \cap A)$. Since the n -diagonal measure is the only measure m_π charging D_n , this implies as claimed that the weight of the n -diagonal measure is α .

We now want to prove by induction that, for all $n \geq 1$, M_n^N is the n -order moment measure of a Poisson process of intensity $\alpha\mu$. The property is of course satisfied for $n = 1$. Let us assume it is satisfied up to some $n \geq 1$, and let A_1, \dots, A_{n+1} be sets in \mathcal{A}_f . Pick a nonempty subset $K \subsetneq \{1, \dots, n + 1\}$. By the ergodic theorem, we get

$$\begin{aligned} & \frac{1}{\ell} \sum_{k=1}^{\ell} \mathbb{E} \left[\prod_{i \in K} N(A_i) \left(\prod_{i \in K^c} N(A_i) \circ S^k \right) \right] \\ (2) \quad & \xrightarrow{\ell \rightarrow \infty} \mathbb{E} \left[\prod_{i \in K} N(A_i) \right] \mathbb{E} \left[\prod_{i \in K^c} N(A_i) \right] \\ & = M_{\#K}^N \left(\prod_{i \in K} A_i \right) M_{(n+1-\#K)}^N \left(\prod_{i \in K^c} A_i \right). \end{aligned}$$

On the other hand,

$$\begin{aligned} & \frac{1}{\ell} \sum_{k=1}^{\ell} \mathbb{E} \left[\prod_{i \in K} N(A_i) \left(\prod_{i \in K^c} N(A_i) \circ S^k \right) \right] \\ (3) \quad & = \frac{1}{\ell} \sum_{k=1}^{\ell} M_{n+1}^N(T^{-\epsilon_k(1)} A_1 \times \cdots \times T^{-\epsilon_k(n+1)} A_{n+1}) \\ & = \sum_{\pi \in \mathcal{P}_{n+1}} \alpha_\pi \frac{1}{\ell} \sum_{k=1}^{\ell} m_\pi(T^{-\epsilon_k(1)} A_1 \times \cdots \times T^{-\epsilon_k(n+1)} A_{n+1}) \end{aligned}$$

where $\epsilon_k(i) := 0$ if $i \in K$, and $\epsilon_k(i) := k$ otherwise. Coming back to the definition of m_π , we write

$$m_\pi(T^{-\epsilon_k(1)} A_1 \times \cdots \times T^{-\epsilon_k(n+1)} A_{n+1}) = \prod_{P \in \pi} \mu \left(\bigcap_{i \in P} T^{-\epsilon_k(i)} A_i \right).$$

Observe that, if K is a union of atoms of π , we have for any $1 \leq k \leq \ell$,

$$(4) \quad m_\pi(T^{-\epsilon_k(1)} A_1 \times \cdots \times T^{-\epsilon_k(n+1)} A_{n+1}) = m_\pi(A_1 \times \cdots \times A_{n+1}).$$

Otherwise, there exists an atom $P \in \pi$ containing indices $i \in K$ and $j \notin K$, hence with $\epsilon_k(i) = 0$ and $\epsilon_k(j) = k$. We get that for some constant C ,

$$m_\pi(T^{-\epsilon_k(1)} A_1 \times \cdots \times T^{-\epsilon_k(n+1)} A_{n+1}) \leq C\mu(T^{-k} A_j \cap A_i).$$

But, since there is no T -invariant set of positive finite measure,

$$\frac{1}{\ell} \sum_{k=1}^{\ell} \mu(T^{-k} A_j \cap A_i) \xrightarrow{\ell \rightarrow \infty} 0.$$

Let \mathcal{P}_{n+1}^K be the set of partitions $\pi \in \mathcal{P}_{n+1}$ where K is a union of atoms of π . The above proves that, in the limit as $\ell \rightarrow \infty$, the contribution in (3) of all partitions $\pi \in \mathcal{P}_{n+1} \setminus \mathcal{P}_{n+1}^K$ vanishes. Thus we get, using (2), (3) and (4),

$$M_{\#K}^N \left(\prod_{i \in K} A_i \right) M_{(n+1-\#K)}^N \left(\prod_{i \in K^c} A_i \right) = \sum_{\pi \in \mathcal{P}_{n+1}^K} \alpha_\pi m_\pi (A_1 \times \cdots \times A_{n+1}).$$

Since $\emptyset \neq K \subsetneq \{1, \dots, n + 1\}$, the decompositions of $M_{\#K}^N$ and $M_{(n+1-\#K)}^N$ only involve the coefficients α_π , $\pi \in \mathcal{P}_1 \cup \cdots \cup \mathcal{P}_n$. Using the mutual singularity of the measures on both sides of the above equality, we see that all of the coefficients α_π , $\pi \in \mathcal{P}_{n+1}^K$, are completely determined by the coefficients corresponding to partitions in $\mathcal{P}_1 \cup \cdots \cup \mathcal{P}_n$. Moreover, the above argument is valid in particular when N is the Poisson process of intensity $\alpha\mu$. By letting K run through all strict subsets of $\{1, \dots, n + 1\}$, and using the induction hypothesis, we identify all but one coefficients of the decomposition of M_{n+1}^N as those of the Poisson point process of intensity $\alpha\mu$. The only coefficient that cannot be determined by this method is the one associated to the trivial partition of $\{1, \dots, n + 1\}$ into a single atom. But this corresponds to the $(n + 1)$ -diagonal measure, and we already know that this coefficient is α .

We have proved that the moment measures of any order of N are those of a Poisson point process of intensity $\alpha\mu$. Lemma 3.1 in [9] ensures then that N is a Poisson point process of intensity $\alpha\mu$. \square

Observe that the conclusion of the proof fails if one does not assume the ergodicity of the point process N : think of a mixture of two Poisson point processes with different intensities.

The action of $T^{\times n}$ on $(X^n, \mathcal{A}^{\otimes n}, m_\pi)$ is isomorphic to $(X^{\#\pi}, \mathcal{A}^{\otimes \#\pi}, \mu^{\otimes \#\pi}, T^{\times \#\pi})$. It is therefore ergodic if we assume that T has infinite ergodic index. With this assumption, we get the following easy consequence for a thinning of Poisson T -point process.

PROPOSITION 2.2. *Assume T has infinite ergodic index. Let N and N' be T -point processes defined on the system $(\Omega, \mathcal{F}, \mathbb{P}, S)$ such that N is Poisson of intensity μ and $N' \leq N$. If N' is ergodic, then it is Poisson of intensity $\alpha\mu$ for some $0 \leq \alpha \leq 1$.*

PROOF. If T has infinite ergodic index, all the measures m_π , $\pi \in \mathcal{P}_n$, $n \geq 1$, are ergodic with respect to $T^{\times n}$, therefore, the formula $\sum_{\pi \in \mathcal{P}_n} c_\pi m_\pi$ is precisely the ergodic decomposition of the moment measure M_n^N of the Poisson process N , with respect to $T^{\times n}$.

Now, as $N' \leq N$, we get for each $n \geq 1$, $N' \otimes \cdots \otimes N' \leq N \otimes \cdots \otimes N$, and thus $M_n^{N'} = \mathbb{E}[N' \otimes \cdots \otimes N'(\cdot)] \leq \mathbb{E}[N \otimes \cdots \otimes N(\cdot)] = M_n^N$. We therefore obtain the ergodic decomposition of $M_n^{N'}$ in the form

$$M_n^{N'} = \sum_{\pi \in \mathcal{P}_n} \alpha_\pi m_\pi$$

for some nonnegative numbers α_π . We can now apply the preceding theorem to the ergodic T -point process N' to get the result. \square

EXAMPLE 2.3. Let us describe an example where the conclusion of the proposition fails. Consider the so-called ‘‘homogeneous Poisson process,’’ that is, the classical Poisson process on the real line with intensity equal to the Lebesgue measure. The base transformation is the translation $T : x \mapsto x + 1$ (in particular, the Poisson T -point process is ergodic). Now we can form a thinning N' by keeping the points of N that are separated by at least $\kappa > 0$ from every other point. It is easy to see that N' is an ergodic T -point process but obviously not a Poisson

process. Here, the proposition does not apply; not really because T is not ergodic (in fact, T can be embedded in an ergodic action of \mathbb{R}), but because the Cartesian powers fail to be ergodic (even if we consider the \mathbb{R} -action).

For the proof of the next theorem, we shall need some definitions and the following lemma which is a particular case of Lemma 2.6 in [9].

DEFINITION 2.4. We say that a T -point process N defined on $(\Omega, \mathcal{F}, \mathbb{P}, S)$ is *free* if for \mathbb{P} -almost all ω , for all $k \in \mathbb{Z}^*$, $N(\omega) \cap N(S^k\omega) = \emptyset$.

Two T -point processes N_1 and N_2 defined on $(\Omega, \mathcal{F}, \mathbb{P}, S)$ are said to be *dissociated* if, for \mathbb{P} -almost all ω , for all $k \in \mathbb{Z}$, $N_1(\omega) \cap N_2(S^k\omega) = \emptyset$.

Note that if N is a Poisson T -point process, then N is free (Proposition 2.7 in [9]).

LEMMA 2.5. Let N_1, \dots, N_n be n T -point processes defined on the ergodic system $(\Omega, \mathcal{F}, \mathbb{P}, S)$, having moments of all orders. Assume there exist a real number $c > 0$ and some $\pi \in \mathcal{P}_n$ such that for any sets A_1, \dots, A_n in \mathcal{A}_f ,

$$\mathbb{E}[N_1(A_1) \cdots N_n(A_n)] \geq cm_\pi(A_1 \times \cdots \times A_n).$$

Then, for any atom $P \in \pi$, any $A \in \mathcal{A}_f$,

$$\mathbb{P}\left(A \cap \bigcap_{i \in P} N_i \neq \emptyset\right) > 0.$$

In particular, for $i, j \in P$, the processes N_i and N_j are not dissociated.

THEOREM 2.6. If T has infinite ergodic index, then any ergodic splitting of a Poisson T -point process is Poisson.

PROOF. We start with a splitting of order $k < \infty$ of a Poisson T -point process N defined on the ergodic dynamical system $(\Omega, \mathcal{F}, \mathbb{P}, S)$, that is, we have k T -point process N_1, \dots, N_k so that $N = N_1 + \cdots + N_k$.

From Proposition 2.2, the N_j 's are Poisson processes with respective intensities $\alpha_1\mu, \dots, \alpha_k\mu$ such that $\alpha_1 + \cdots + \alpha_k = 1$.

It remains to show the mutual independence of N_1, \dots, N_k . Let n_1, \dots, n_k be positive numbers, $n := n_1 + \cdots + n_k$, and let $\{Q_1, \dots, Q_k\}$ be the partition of $\{1, \dots, n\}$ in subsets of consecutive integers of respective size n_1, \dots, n_k . For any $\{A_i\}_{1 \leq i \leq n}$ in \mathcal{A}_f , set

$$(5) \quad \sigma(A_1 \times \cdots \times A_n) := \mathbb{E}\left[\prod_{j=1}^k \prod_{i \in Q_j} N_j(A_i)\right].$$

This defines a $T^{\times n}$ -invariant measure σ on $(X^n, \mathcal{A}^{\otimes n})$, for which, as above, since for all $1 \leq j \leq k$ we have $N_j \leq N$, $\sigma \leq M_n^N$. Hence σ has at most countably many ergodic components, of the form m_π for some $\pi \in \mathcal{P}_n$. Observe that the processes N_1, \dots, N_k are mutually dissociated as, for all $1 \leq j \leq k$, $N_j \leq N$ and N is free. Therefore, by Lemma 2.5, if the measure m_π appears in the ergodic decomposition of σ , then π refines the partition $\{Q_1, \dots, Q_k\}$. Hence, any ergodic component m_π of σ has the form

$$m_\pi(A_1 \times \cdots \times A_n) = \prod_{j=1}^k \prod_{P \in \pi, P \subset Q_j} \mu\left(\bigcap_{i \in P} A_i\right) = \prod_{j=1}^k \nu_j\left(\times_{i \in Q_j} A_i\right),$$

where each ν_j is a $T^{\times n_j}$ -invariant measure. In particular, any ergodic component of σ is invariant by the transformation $(x_1, \dots, x_n) \mapsto (y_1, \dots, y_n)$, where $y_i := Tx_i$ if $i \in Q_k$, and $y_i := x_i$ otherwise. It follows that σ itself is invariant by this transformation, hence the expression defining $\sigma(A_1 \times \dots \times A_n)$ on the right-hand side of (5) is unchanged if we replace $N_k(A_i)$ by $N_k(T^{-1}A_i)$ for all $i \in Q_k$ simultaneously. Therefore, we can write for any $\{A_i\}_{1 \leq i \leq n}$ in \mathcal{A}_f and any $L \geq 1$,

$$\begin{aligned} \mathbb{E} \left[\prod_{j=1}^k \prod_{i \in Q_j} N_j(A_i) \right] &= \frac{1}{L} \sum_{1 \leq \ell \leq L} \mathbb{E} \left[\left(\prod_{j=1}^{k-1} \prod_{i \in Q_j} N_j(A_i) \right) \prod_{i \in Q_k} N_k(T^{-\ell}A_i) \right] \\ &= \mathbb{E} \left[\left(\prod_{j=1}^{k-1} \prod_{i \in Q_j} N_j(A_i) \right) \left(\frac{1}{L} \sum_{1 \leq \ell \leq L} \prod_{i \in Q_k} N_k \circ S^\ell(A_i) \right) \right]. \end{aligned}$$

By the ergodic theorem, this converges as $L \rightarrow \infty$ to

$$\mathbb{E} \left[\prod_{j=1}^{k-1} \prod_{i \in Q_j} N_j(A_i) \right] \mathbb{E} \left[\prod_{i \in Q_k} N_k(A_i) \right].$$

A straightforward induction on k then yields the equality

$$\mathbb{E} \left[\prod_{j=1}^k \prod_{i \in Q_j} N_j(A_i) \right] = \prod_{j=1}^k \mathbb{E} \left[\prod_{i \in Q_j} N_j(A_i) \right],$$

and this is sufficient to obtain the independence between the Poisson processes.

The case $k = \infty$ is easily deduced from the finite order case. \square

REMARK 2.7. For the conclusion of Theorem 2.6 to hold, it is in fact enough to assume only that (with the notation of the proof) all but one of the T -point processes N_j are ergodic. Indeed, we can then apply the proof in any ergodic component of the joining (N_1, \dots, N_k) , and see that in each of these ergodic components we have the same structure of independent Poisson processes. A posteriori we see that there is only one ergodic component. In particular, with the assumptions of Proposition 2.2, we get that N' and $(N - N')$ are independent Poisson processes, and thus form a Poisson splitting. Moreover, if we remove the assumption of ergodicity of N' in Proposition 2.2, we get that the ergodic components of $(N', N - N')$ are necessarily Poisson splittings.

3. Application to marked Poisson point processes. Here, we deal with so-called marked point processes. Roughly speaking, a marked point process on X with marks in some measurable space (K, \mathcal{K}) is a point process on X whose points carry some information, a mark, taking values in K . For a T -point process, we require the mark to have the *shadowing property*, meaning that it follows the point when the dynamics on the point process is applied. We thus consider a *marked T -point process* as a $(T \times \text{Id})$ -point process on the bigger space $(X \times K, \mathcal{A} \otimes \mathcal{K})$ with intensity measure $\tilde{\mu}$ that projects on μ .

THEOREM 3.1. *Let (X, \mathcal{A}, μ, T) be an infinite measure preserving dynamical system with infinite ergodic index. Let N be an ergodic $(T \times \text{Id})$ -point process on $X \times K$. Assume that the T -point process $N_0 : A \mapsto N(A \times K)$ is a Poisson point process with intensity μ . Then N is a Poisson point process with intensity $\mu \otimes \rho$ where ρ is some probability measure on K .*

REMARK 3.2. It is well known that when a Poisson point process is independently endowed with i.i.d. marks, then the resulting point process on the product space is a Poisson process whose intensity is the product measure of the original intensity and the distribution of the marks. The above result means that, for T with infinite ergodic index, the only way to get an ergodic marked T -point process out of a Poisson T -point process is precisely to take i.i.d. marks, independent of the underlying process.

PROOF OF THEOREM 3.1. Let us denote by $\tilde{\mu}$ the intensity of N on $X \times K$. Let $E = \bigsqcup_{i \in I} A_i \times B_i \subset X \times K$ be a finite union of pairwise disjoint product sets of finite $\tilde{\mu}$ -measure. Let $(\tilde{B}_j)_{j \in J}$ be the finite partition of K into nonempty subsets generated by the sets B_i . For each $j \in J$, denote by I_j the subset of $i \in I$ such that $\tilde{B}_j \subset B_i$, then observe that the sets A_i , $i \in I_j$ are pairwise disjoint. Therefore, E can be also written as the disjoint union

$$E = \bigsqcup_{j \in J} \bigsqcup_{i \in I_j} A_i \times \tilde{B}_j,$$

and this refines the original partition of E into product sets $A_i \times B_i$. Since the ergodic T -point processes $N_{\tilde{B}_j} := N(\cdot \times \tilde{B}_j)$, $j \in J$, form an ergodic splitting of N_0 , we get by Theorem 2.6 that they are independent Poisson T -point processes. Since for each $j \in J$ the sets A_i , $i \in I_j$, are pairwise disjoint, the random variables $N(A_i \times \tilde{B}_j)$, $j \in J$, $i \in I_j$ are independent Poisson random variables. Finally, the random variables $N(A_i \times B_i)$, $i \in I$ are also independent Poisson random variables. This is enough to prove that N is a Poisson point process.

To get the intensity, observe that, for each $B \in \mathcal{K}$, $N(\cdot \times B)$ is a thinning of N_0 whose intensity is μ . By ergodicity of (X, \mathcal{A}, μ, T) , there exists $0 \leq \rho(B) \leq 1$ such that $\rho(B)\mu$ is the intensity of $N(\cdot \times B)$. It is now clear that the map $B \mapsto \rho(B)$ defines a probability measure on (K, \mathcal{K}) . Finally, $\tilde{\mu} = \mu \otimes \rho$. \square

4. Extended SuShis: From simple point processes to general random measures.

In this section, we aim to extend the “rigidity” result obtained in [9] for simple point processes to general random measures. The hypothesis on (X, \mathcal{A}, μ, T) will be the same as in [9] but rephrased with the notation introduced earlier.

DEFINITION 4.1. We say that the infinite measure preserving system (X, \mathcal{A}, μ, T) has the (P) property if, for each $n \geq 1$ the following is true: whenever σ is a boundedly finite, $T^{\times n}$ -invariant measure on X^n , with marginals absolutely continuous with respect to μ , then σ is conservative, and its ergodic components are all of the form $(T^{k_1} \times \dots \times T^{k_n})_* m_\pi$ for some $\pi \in \mathcal{P}_n$ and integers k_1, \dots, k_n .

Note that the (P) property implies in particular that T has infinite ergodic index (otherwise the ergodic components of some product measure $\mu^{\otimes n}$ would not satisfy the required assumption).

Let $\tilde{X}_c \subset \tilde{X}$ and $\tilde{X}_d \subset \tilde{X}$ be respectively the spaces of continuous and discrete boundedly finite measures on (X, \mathcal{A}) .

If N is an integrable T -random measure, we can write N as $N_c + N_d$ where $N_c \in \tilde{X}_c$ and $N_d \in \tilde{X}_d$ (both N_c and N_d can be obtained deterministically from N). Of course N_c and N_d are still integrable T -random measures whose respective intensities are multiples of μ (by ergodicity of T). Thanks to this decomposition, we can study separately the case of a.s. continuous T -random measures, and the case of a.s. discrete T -random measures, which will be the objects of the two following sections.

4.1. *The continuous case.*

PROPOSITION 4.2. *Assume that T has the (P) property.*

If N is a square integrable T -random measure defined on some ergodic system $(\Omega, \mathcal{F}, \mathbb{P}, S)$, whose realizations are a.s. continuous, then there exists $\alpha \geq 0$ such that N is constant and a.s. equal to $\alpha\mu$.

If the realizations are a.s. absolutely continuous with respect to μ , no hypothesis on moments are required to get the same conclusion, although we might have $\alpha = +\infty$.

PROOF. First, assume N is square integrable. We know that its intensity is of the form $\alpha\mu$ for some $\alpha \geq 0$, and the moment measure of order two, defined on $\mathcal{A} \otimes \mathcal{A}$ by

$$M_2^N(A \times B) := \mathbb{E}[N(A)N(B)] = \mathbb{E}[N \otimes N(A \times B)]$$

is boundedly finite. This measure is $T \times T$ -invariant, and its marginals are absolutely continuous with respect to μ . Moreover, as N is a.s. continuous, $N \otimes N$ gives zero measure to the graphs of T^k , for all $k \in \mathbb{Z}$. Thanks to property (P), $M_2^N = \beta\mu \otimes \mu$ for some $\beta \geq 0$. Applying the ergodic theorem, we get for all $A \in \mathcal{A}_f$,

$$\frac{1}{n} \sum_{k=1}^n \mathbb{E}[N(A)N(A) \circ S^k] \xrightarrow{n \rightarrow \infty} \mathbb{E}[N(A)]^2 = \alpha^2 \mu(A)^2.$$

On the other hand,

$$\frac{1}{n} \sum_{k=1}^n \mathbb{E}[N(A)N(A) \circ S^k] = \frac{1}{n} \sum_{k=1}^n M_2^N(A \times T^{-k}A) = \frac{1}{n} \sum_{k=1}^n \beta\mu(A)\mu(T^{-k}A),$$

which by invariance of μ is equal to $\beta\mu(A)^2$. Therefore, $\beta = \alpha^2$ and

$$\mathbb{E}[(N(A) - \alpha\mu(A))^2] = M_2^N(A \times A) - \alpha^2 \mu(A)^2 = 0,$$

which implies the result.

If we assume that the realizations are a.s. absolutely continuous, then we can write for all set $A \in \mathcal{A}$,

$$N(\omega)(A) = \int_A f(\omega, x)\mu(dx).$$

We can therefore define N_n by

$$N_n(\omega)(A) = \int_A (f \wedge n)(\omega, x)\mu(dx).$$

N_n is still a continuous ergodic T -random measure but is now square integrable. From the first part of the proof, $N_n = \alpha_n\mu$ a.s. Since N_n increases to N , we get $N = \alpha\mu$ a.s. where α is the increasing limit of α_n . \square

4.2. *The discrete case.* We consider the set of sequences

$$\ell_1^+(\mathbb{Z}) := \left\{ (a_k)_{k \in \mathbb{Z}} : \forall k \in \mathbb{Z}, a_k \geq 0 \text{ and } \sum_{k \in \mathbb{Z}} a_k < \infty \right\}.$$

The ℓ_1 norm turns $\ell_1^+(\mathbb{Z})$ into a complete separable metric space. The goal of this section is to obtain the following result.

THEOREM 4.3. *Assume T has the (P) property. Let N be a nonzero T -random measure with moments of all orders defined on some ergodic system $(\Omega, \mathcal{F}, \mathbb{P}, S)$ and whose realizations are almost-surely discrete. Then there exists a probability distribution κ on $\ell_1^+(\mathbb{Z})$ and a positive number c such that N is distributed as*

$$A \mapsto \int_{X \times \ell_1^+(\mathbb{Z})} \sum_{k \in \mathbb{Z}} a_k \mathbb{1}_A(T^k x) \mathcal{N}(dx, d\{a_k\}_{k \in \mathbb{Z}})$$

where \mathcal{N} is a Poisson point process on $X \times \ell_1^+(\mathbb{Z})$ with intensity $c\mu \otimes \kappa$.

This result says that N has a cluster form which can be obtained in the following way: start from a Poisson point process of intensity $c\mu$, then replace independently each point x output by this Poisson point process with a random measure (the cluster) of the form

$$\sum_{k \in \mathbb{Z}} a_k \delta_{T^k x}$$

where $\{a_k\}_{k \in \mathbb{Z}}$ is chosen according to κ .

In the following, replacing if necessary μ by the intensity of N which is of the form $\alpha\mu$ for some $\alpha > 0$ by ergodicity of T , we assume that the intensity of N is μ .

4.2.1. Removing points with small weights. Consider N as in the statement of Theorem 4.3. For $\epsilon > 0$, we define N_ϵ from N by removing points with weights less than ϵ . We also define $N_{\epsilon,1}$ as the simple point process obtained from N_ϵ with all weights set to 1. We have for all $A \in \mathcal{A}_f$,

$$N_{\epsilon,1}(A) \leq \frac{1}{\epsilon} N_\epsilon(A) \leq \frac{1}{\epsilon} N(A),$$

therefore, N_ϵ and $N_{\epsilon,1}$ are both T -random measures with moments of all orders.

According to Proposition 2.1 in [9], any square-integrable T -point process almost surely gives finite measure to any T -orbit. In particular, $N_{\epsilon,1}$ almost surely belongs to the subset $\tilde{X}_{d,f} \subset \tilde{X}_d$ of measures ν satisfying the following property:

$$\forall x \in X \quad \#\{n \in \mathbb{Z} : \nu(T^n x) > 0\} < \infty.$$

Of course, $N_\epsilon \in \tilde{X}_{d,f}$ almost surely as well.

We construct an injective map Φ from $\tilde{X}_{d,f}$ to $(X \times \ell_1^+(\mathbb{Z}))^*$. For $\nu \in \tilde{X}_{d,f}$, $\Phi(\nu)$ is the simple counting measures supported on the following collection of points in $X \times \ell_1^+(\mathbb{Z})$: we select in each orbit seen by ν the first element x in the orbit with maximal weight, and consider the point $(x, (\nu(T^n x))_{n \in \mathbb{Z}})$. In other words, a point $(x, (\beta_n)_{n \in \mathbb{Z}})$ belongs to $\Phi(\nu)$ if and only if:

- $\beta_0 > \beta_n$ for each $n < 0$;
- $\beta_0 \geq \beta_n$ for each $n \geq 0$;
- $\nu(\{T^n x\}) = \beta_n$ for each $n \in \mathbb{Z}$.

Observe that we can recover ν from $\Phi(\nu)$ by the formula

$$(6) \quad \nu(A) = \int_{X \times \ell_1^+(\mathbb{Z})} \sum_{k \in \mathbb{N}} \beta_k \mathbb{1}_A(T^k x) \Phi(\nu)(dx, d\{\beta_k\}_{k \in \mathbb{N}}).$$

Now, $\Phi(N_\epsilon)$ is \mathbb{P} -a.s. well-defined as a $(T \times \text{Id})$ -point process on $X \times \ell_1^+(\mathbb{Z})$. Its projection on the first coordinate $(\Phi(N_\epsilon))_0 := \Phi(N_\epsilon)(\cdot \times \ell_1^+(\mathbb{Z}))$ is a T -point process on X with moments of all orders (since $\Phi(N_\epsilon)(\cdot \times \ell_1^+(\mathbb{Z})) \leq N_{\epsilon,1}(\cdot)$). By construction, this T -point process is free since we have selected one point in each orbit seen by N_ϵ .

Theorem 3.2 in [9] states that, under the (P) property, any free T -point process with moments of all orders defined on an ergodic system is Poisson. Thus we get that $(\Phi(N_\epsilon))_0$ is a Poisson process. Then by Theorem 3.1, we obtain that $\Phi(N_\epsilon)$ is a Poisson process on $(X \times \ell_1^+(\mathbb{Z}))$. Applying (6), we almost surely have

$$N_\epsilon(A) = \int_{X \times \mathbb{R}_+^{\mathbb{Z}}} \sum_{k \in \mathbb{N}} \alpha_k \mathbb{1}_A(T^k x) \Phi(N_\epsilon)(dx, d\{\alpha_k\}_{k \in \mathbb{N}}),$$

and thus we get the theorem for N_ϵ , for any $\epsilon > 0$.

To get the result for N , we need to take advantage of the infinite divisibility character of N_ϵ , that N will inherit in the limit. We therefore have to recall some general results about infinite divisibility.

4.2.2. *Infinitely divisible random measures.* The notion of infinite divisibility can be defined on any measurable semigroup although we only give it in the context we are interested in.

DEFINITION 4.4. let (Z, \mathcal{Z}) be complete separable metric space. A probability distribution σ on $(\tilde{Z}, \tilde{\mathcal{Z}})$ is *infinitely divisible (ID)* if, for any $k \in \mathbb{N}$, there exists a probability distribution σ_k such that

$$\sigma = \underbrace{\sigma_k \star \cdots \star \sigma_k}_{k \text{ times}},$$

where \star is the convolution of measures induced by the addition on \tilde{Z} . By extension, we say that a random measure on Z is ID if its distribution is.

The first examples of ID random measures are Poisson measures. Their ID character comes directly from the well-known fact that the sum of two independent Poisson processes on the same space with intensities μ_1 and μ_2 is again a Poisson process, but with intensity $\mu_1 + \mu_2$.

We recall the fundamental representation result that can be found in [12], Theorem 3.20.

THEOREM 4.5. A probability measure m on \tilde{X} is ID if and only if there exist a measure $\gamma \in \tilde{X}$ and a σ -finite measure ρ on $\tilde{X} \setminus \{0\}$ satisfying, for all bounded $B \subset \tilde{X}$

$$\int_{\tilde{X}} (\xi(B) \wedge 1) d\rho(\xi) < \infty,$$

such that m is the distribution of the following random measure:

$$\gamma + \int_{\tilde{X}} \xi(\cdot) d\omega(\xi),$$

where ω is a random element of $(\tilde{X})^*$ chosen according to the Poisson measure ρ^* on \tilde{X} with intensity ρ .

The measures γ and ρ are uniquely determined by m , and ρ is called the Lévy measure of m .

Here, we summarize useful properties about the case we are interested in.

PROPOSITION 4.6. Let (X, \mathcal{A}, μ, T) be a conservative ergodic infinite measure preserving system, and let m be the distribution of an ergodic square integrable ID random measure with intensity μ , and whose realizations are almost surely discrete. Assume that m corresponds to (γ, ρ) as in Theorem 4.5, then:

- $\gamma = 0$,
- ρ is T_* -invariant and supported by \tilde{X}_d ,
- $\int_{\tilde{X}} \xi(A)\rho(d\xi) = \mu(A)$ for all $A \in \mathcal{A}_f$,
- $\int_{\tilde{X}} \xi(A)^2\rho(d\xi) = \int_{\tilde{X}} (\xi(A) - \mu(A))^2 m(d\xi)$ for all $A \in \mathcal{A}_f$,
- $(\tilde{X}, \tilde{\mathcal{A}}, \rho, T_*)$ has no T_* -invariant sets of finite, nonzero measure.

PROOF. If m corresponds to (γ, ρ) , then $(T_*)_*(m)$ is ID, too, and corresponds to $(T_*(\gamma), (T_*)_*(\rho))$. Therefore, if m is T_* -invariant then γ is T -invariant and ρ is T_* -invariant. For all $A \in \mathcal{A}_f$,

$$\begin{aligned} \mu(A) &= \int_{\tilde{X}} \xi(A)m(d\xi) \\ &= \gamma(A) + \int_{\tilde{X}^*} \int_{\tilde{X}} \xi(A)\omega(d\xi)\rho^*(d\omega) \\ &= \gamma(A) + \int_{\tilde{X}} \xi(A)\rho(d\xi). \end{aligned}$$

In particular, $\gamma \ll \mu$, and by ergodicity of μ , γ is some multiple of μ . But then m could not be the distribution of a point process as it would possess a continuous part, unless γ vanishes. From the Poisson construction, we also get that ρ is supported on \tilde{X}_d .

Next, the second formula is an application of the isometry formula (1).

The last fact should come as no surprise for anyone familiar with ergodic properties of ID systems; however, there is no available proof for this particular case. We give one here:

Assume $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$ is ergodic and $(\tilde{X}, \tilde{\mathcal{A}}, \rho, T_*)$ possesses a T_* -invariant set K with $\rho(K) < \infty$. Let $((\tilde{X})^*, (\tilde{\mathcal{A}})^*, \rho^*, (T_*)_*)$ be the Poisson suspension over $(\tilde{X}, \tilde{\mathcal{A}}, \rho, T_*)$. The map

$$\omega \mapsto \int_{\tilde{X}} \xi(\cdot)\omega(d\xi)$$

is a factor map between the suspension and the ergodic system $(\tilde{X}, \tilde{\mathcal{A}}, m, T_*)$. Moreover, we have another factor in the suspension, generated by the stationary process

$$\omega \mapsto \{(T_*)_*^k(\omega)(K)\}_{k \in \mathbb{Z}}.$$

But as K is T_* -invariant, $(T_*)_*^k(\omega)(K) = \omega(K)$ for all $k \in \mathbb{Z}$, therefore, $(T_*)_*$ acts as the identity on this factor. By disjointness between the identity map and any ergodic map [5], we obtain that these two factors are independent inside the Poisson suspension. In particular, for any $A \in \mathcal{A}_f$, $\omega \mapsto \int_{\tilde{X}} \xi(A)\omega(d\xi)$ and $\omega \mapsto \omega(K)$ are independent. It follows that

$$\int_{(\tilde{X})^*} \left(\int_{\tilde{X}} \xi(A)\omega(d\xi) - \mu(A) \right) (\omega(K) - \rho(K))\rho^*(d\omega) = 0.$$

By the isometry formula (1), we get

$$\int_{\tilde{X}} \xi(A)\mathbb{1}_K(\xi)\rho(d\xi) = 0.$$

By monotone convergence, we can replace A by X in the preceding integral and, as $\xi(X) > 0$ ρ -a.e., we get that $\mathbb{1}_K = 0$ ρ -a.e., that is, $\rho(K) = 0$. \square

Now, let us explain how N inherits the ID character of N_ϵ in the limit. We have $N_\epsilon \ll N$, thus we can introduce the random density $g_\epsilon := \frac{dN_\epsilon}{dN}$. Almost surely, g_ϵ increases to 1 as

$\epsilon \rightarrow 0$. Let f be a bounded continuous function on X with bounded support. Then by the dominated convergence theorem,

$$\int_X f dN_\epsilon = \int_X f g_\epsilon dN \xrightarrow{\epsilon \rightarrow 0} \int_X f dN \quad \text{a.s.}$$

Let also h be a bounded continuous function on \mathbb{R} . Again by the dominated convergence theorem, we get

$$\mathbb{E} \left[h \left(\int_X f dN_\epsilon \right) \right] \xrightarrow{\epsilon \rightarrow 0} \mathbb{E} \left[h \left(\int_X f dN \right) \right].$$

By [12], Theorem 4.11, this characterizes the weak convergence of the distribution of N_ϵ to the distribution of N . Using [12], Lemma 4.24, we can conclude that N is ID as a limit of ID random measures.

N is thus square integrable and ID, hence admits a Lévy measure ρ which is a σ -finite measure on \tilde{X}_d , and is T_* -invariant. By ergodicity of N , the system $(\tilde{X}, \tilde{\mathcal{A}}, \rho, T_*)$ enjoys all the properties stated in Proposition 4.6.

When T has the (P) property, this system has a very simple structure that we fully describe below.

4.2.3. *Infinite quasifactors.* We start by extending the notion of quasifactor of Glasner and Meyerovitch to the case of an infinite measure on \tilde{X} .

DEFINITION 4.7. An ∞ -quasifactor of (X, \mathcal{A}, μ, T) is a dynamical system $(\tilde{X}, \tilde{\mathcal{A}}, \rho, T_*)$ where ρ is an infinite, σ -finite, T_* -invariant measure and

$$\forall A \in \mathcal{A}_f \quad \int_{\tilde{X}} \xi(A) \rho(d\xi) = \mu(A).$$

The ∞ -quasifactor $(\tilde{X}, \tilde{\mathcal{A}}, \rho, T_*)$ is said to be *square integrable* if

$$\forall A \in \mathcal{A}_f \quad \int_{\tilde{X}} (\xi(A))^2 \rho(d\xi) < \infty.$$

As a simple example of ∞ -quasifactor, we can consider the infinite measure preserving system (X^*, ρ, T_*) , where ρ is the pushforward of μ by $x \mapsto \delta_x$. It is actually the Lévy measure of the Poisson point process of intensity μ . More generally, from Proposition 4.6, any ergodic infinitely divisible and square integrable T -random measure with intensity μ gives rise to a square integrable ∞ -quasifactor through its Lévy measure.

Let us first consider the case of a *simple* ∞ -quasifactor, that is, a measure ρ concentrated on the set X^* of simple counting measures on X .

PROPOSITION 4.8. *If T has the (P) property and $(X^*, \mathcal{A}^*, \rho, T_*)$ is a simple square integrable ∞ -quasifactor without T_* -invariant set of nonzero finite ρ -measure, then ρ -a.e. $\xi \in X^*$ is concentrated on a finite subset of a single T -orbit.*

PROOF. As $(X^*, \mathcal{A}^*, \rho, T_*)$ is square integrable, the formula

$$m(A \times B) := \int_{X^*} \xi(A) \xi(B) \rho(d\xi)$$

defines a boundedly finite $T \times T$ -invariant measure on $X \times X$. Thanks to property (P), it can be written as

$$m = \alpha_\infty \mu \otimes \mu + \sum_{k \in \mathbb{Z}} \alpha_k \Delta_k,$$

where Δ_k is the measure supported on the graph of T^k defined by $\Delta_k(A \times B) := \mu(A \cap T^{-k}B)$. Observe that, as there is no T_* -invariant set of nonzero finite ρ -measure,

$$\frac{1}{n} \sum_{\ell=1}^n m(A \times T^{-\ell}B) = \frac{1}{n} \sum_{\ell=1}^n \int_{X^*} \xi(A) T_*^\ell \xi(B) \rho(d\xi) \rightarrow 0.$$

However,

$$\frac{1}{n} \sum_{\ell=1}^n \alpha_\infty \mu \otimes \mu(A \times T^{-\ell}B) = \alpha_\infty \mu \otimes \mu(A \times B),$$

therefore, $\alpha_\infty = 0$.

This means that m is concentrated on the graphs of the maps T^k , $k \in \mathbb{Z}$, therefore, for ρ -a.e. $\xi \in X^*$ the product $\xi \otimes \xi$ is concentrated on these graphs. It follows that ρ -a.e. $\xi \in X^*$ is concentrated on a single T -orbit. It remains to verify that ρ is almost surely concentrated on a finite number of points in this orbit. For this, we observe that Proposition 2.1 in [9] (whose statement is recalled in Section 4.2.1) can be generalized to the case of an infinite measure ρ on X^* . Indeed, the Palm measures of ρ are probability measures since $\int_{X^*} \xi(A) \rho(d\xi) = \mu(A) < +\infty$. Since ρ has moments of order 2, the proposition yields that ρ -a.e. $\xi \in X^*$ is concentrated on a finite subset of a single T -orbit. \square

PROPOSITION 4.9. *If T has the (P) property and if $(\tilde{X}_d, \tilde{\mathcal{A}}, \rho, T_*)$ is a square integrable ∞ -quasifactor without T_* -invariant set of nonzero finite ρ -measure, then there exists $c > 0$ and a factor map φ from $(\tilde{X}_d, \tilde{\mathcal{A}}, \rho, T_*)$ to $(X, \mathcal{A}, c\mu, T)$ and some T_* -invariant maps $\xi \mapsto a_k(\xi) \geq 0$, $k \in \mathbb{Z}$, such that*

$$\xi = \sum_{k \in \mathbb{Z}} a_k(\xi) \delta_{T^k \varphi(\xi)}.$$

PROOF. First, observe that $\{0\}$ is a T_* -invariant set. It cannot have infinite ρ -measure because ρ is σ -finite, hence $\rho(\{0\}) = 0$ from the hypotheses.

Let $\xi \in \tilde{X}_d$ and set $\xi_{|\varepsilon}$, $\varepsilon > 0$ to be ξ where we have forgotten points with weights less than ε and set the other weights to be 1. Then $\xi \mapsto \xi_{|\varepsilon}$ is a factor map and $\xi_{|\varepsilon}$ turns out to induce on $\tilde{X}_d \setminus \{\xi_{|\varepsilon} = \{0\}\}$ a simple square integrable ∞ -quasifactor without T_* -invariant set of nonzero finite measure. Therefore, by Proposition 4.8, $\xi_{|\varepsilon}$ has a finite number of points on its support, which all lie on a single T -orbit.

It follows that all the points of the support of ξ are ρ -a.s. on a single T -orbit, and only a finite number of them have a weight greater than any fixed positive constant. We can therefore see that the map $\varphi : \xi \mapsto \varphi(\xi)$ where $\varphi(\xi)$ is the point of the support with the highest weight and the lowest place in the orbit is well-defined. It satisfies

$$\varphi(T_* \xi) = T \varphi(\xi).$$

Now with this ‘‘origin’’ $\varphi(\xi)$, we can define maps $\xi \mapsto a_k(\xi) \geq 0$ so that

$$\xi := \sum_{k \in \mathbb{Z}} a_k(\xi) \delta_{T^k \varphi(\xi)}.$$

We have

$$T_* \xi = \sum_{k \in \mathbb{Z}} a_k(\xi) \delta_{T^{k+1} \varphi(\xi)} = \sum_{k \in \mathbb{Z}} a_k(\xi) \delta_{T^k \varphi(T_* \xi)}$$

in the one hand, and on the other hand,

$$T_* \xi := \sum_{k \in \mathbb{Z}} a_k(T_* \xi) \delta_{T^k \varphi(T_* \xi)}.$$

Therefore, the maps a_k are T_* -invariant.

We have for all A

$$\begin{aligned} \mu(A) &= \int_{\tilde{X}_d} \xi(A) \rho(d\xi) \\ &= \sum_{k \in \mathbb{Z}} \int_{\tilde{X}_d} a_k(\xi) \delta_{T^k \varphi(\xi)}(A) \rho(d\xi) \\ &= \sum_{k \in \mathbb{Z}} \int_{\tilde{X}_d} a_k(T_*^k \xi) \delta_{\varphi(T_*^k(\xi))}(A) \rho(d\xi) \quad \text{by } T_*\text{-invariance of } a_k \\ &= \sum_{k \in \mathbb{Z}} \int_{\tilde{X}_d} \delta_{\varphi(\xi)}(A) a_k(\xi) \rho(d\xi) \quad \text{by } T_*\text{-invariance of } \rho. \end{aligned}$$

Let us define for each $k \in \mathbb{Z}$ the measure ρ_k by $\frac{d\rho_k}{d\rho} := a_k$. Then we get

$$\mu(A) = \sum_{k \in \mathbb{Z}} \varphi_* \rho_k(A),$$

and in particular $\varphi_* \rho_0 \ll \mu$. But, as $a_0 > 0$ ρ -a.e., $\rho_0 \sim \rho$ and we also have $\varphi_* \rho \ll \mu$. By ergodicity, $\varphi_* \rho = c\mu$ for some $c > 0$.

Therefore, φ induces a factor map between $(\tilde{X}_d, \tilde{\mathcal{A}}, \rho, T_*)$ and $(X, \mathcal{A}, c\mu, T)$. \square

COROLLARY 4.10. *Assume that T has the (P) property. Let $(\tilde{X}_d, \tilde{\mathcal{A}}, \rho, T_*)$ be an infinite measure-preserving square integrable ∞ -quasifactor without T_* -invariant set of nonzero finite ρ -measure. Then there exists a probability measure κ on $\mathbb{R}_+^{\mathbb{Z}}$ such that $(\tilde{X}_d, \tilde{\mathcal{A}}, \rho, T_*)$ is isomorphic to $(X \times \mathbb{R}_+^{\mathbb{Z}}, \mathcal{A} \otimes \mathcal{B}^{\otimes \mathbb{Z}}, \mu \otimes (c\kappa), T \times \text{Id})$ (where c is given in Proposition 4.9).*

Moreover, we have

$$\frac{1}{c} = \int_{\mathbb{R}_+^{\mathbb{Z}}} \left(\sum_{k \in \mathbb{Z}} a_k \right) \kappa(d\{a_k\}_{k \in \mathbb{Z}}).$$

In particular, $\{a_k\}_{k \in \mathbb{Z}} \in \ell_1^+(\mathbb{Z})$ κ -a.s.

PROOF. Define Φ from \tilde{X}_d to $X \times \mathbb{R}_+^{\mathbb{Z}}$ by

$$\Phi(\xi) := (\varphi(\xi), (a_k(\xi))_{k \in \mathbb{Z}}).$$

Then $T \times \text{Id}$ preserves $m := \Phi_* \rho$ and Φ is an isomorphism between $(\tilde{X}_d, \tilde{\mathcal{A}}, \rho, T_*)$ and $(X \times \mathbb{R}_+^{\mathbb{Z}}, \mathcal{A} \otimes \mathcal{B}^{\otimes \mathbb{Z}}, m, T \times \text{Id})$. Since the σ -algebra generated by $\varphi(\xi)$ is σ -finite by the preceding proposition, we can disintegrate m with respect to the first coordinate: we get a family $(\kappa_x)_{x \in X}$ of probability measures on $\mathbb{R}_+^{\mathbb{Z}}$, such that

$$m(A \times B) = \int_A \kappa_x(B) c d\mu(x),$$

where c is given in the preceding proposition. By invariance of m under $T \times \text{Id}$, we get $\kappa_x = \kappa_{T_x}$ for μ -almost every x , and by ergodicity of T we conclude that there exists κ such that $\kappa_x = \kappa$ μ -almost everywhere. This yields $m = \mu \otimes (c\kappa)$.

Now, for each $A \in \mathcal{A}_f$, we have

$$\begin{aligned} \mu(A) &= \int_{\tilde{X}_d} \xi(A) \rho(d\xi) \\ &= c \int_{\mathbb{R}_+^{\mathbb{Z}}} \left(\int_X \sum_{k \in \mathbb{Z}} a_k \mathbb{1}_A(T^k x) \mu(dx) \right) \kappa(d\{a_k\}_{k \in \mathbb{Z}}) \\ &= c\mu(A) \int_{\mathbb{R}_+^{\mathbb{Z}}} \left(\sum_{k \in \mathbb{Z}} a_k \right) \kappa(d\{a_k\}_{k \in \mathbb{Z}}). \end{aligned}$$

\square

4.2.4. *End of the proof of Theorem 4.3.* We come back to the end of the proof of the main theorem of this section. Recall that under the assumptions of this theorem, we were left with the following situation: N is a T -random measure with moments of all orders, and we showed it is infinitely divisible. Hence the conclusion follows from the next proposition.

PROPOSITION 4.11. *Assume that T has the (P) property. Let N be a square integrable ID T -random measure defined on some ergodic system $(\Omega, \mathcal{F}, \mathbb{P}, S)$ whose realizations are almost surely discrete. Then there exists a probability distribution κ on $(\ell_1^+(\mathbb{Z}), \mathcal{B}^{\otimes \mathbb{Z}})$ and $c > 0$ such that N is distributed as*

$$A \mapsto \int_{X \times \ell_1^+(\mathbb{Z})} \sum_{k \in \mathbb{Z}} a_k \mathbb{1}_A(T^k x) \mathcal{N}(dx, d\{a_k\}_{k \in \mathbb{Z}})$$

where \mathcal{N} is a Poisson point process on $X \times \ell_1^+(\mathbb{Z})$ with intensity $c\mu \otimes \kappa$.

PROOF. By Theorem 4.5 and Proposition 4.6, the ID square integrable T -random measure N can be described by its Lévy measure ρ . The latter is nothing else than a square integrable ∞ -quasifactor, whose structure is completely given in Corollary 4.10: the measure space (\tilde{X}_d, ρ) is isomorphic to $(X \times \ell_1^+(\mathbb{Z}), c\mu \otimes \kappa)$. In this context, the representation of N as an integral with respect to a Poisson random measure takes the more concrete form explicated in the statement of the proposition. \square

REMARK 4.12. Note that the assumption that N has moments of all orders has only been used to obtain the ID character of the random measure. Once this is established, square integrability of N is sufficient to conclude. We do not know if square integrability alone implies the conclusion of Theorem 4.3.

5. Improved disjointness results. This last, short section deals with joinings and disjointness. The notion of joinings in ergodic theory is the dynamical counterpart of couplings in probability theory. It is particularly relevant for the classification of dynamical systems as we do below. For the reader unfamiliar with this notion, we refer to the seminal paper [5] and the book [6], that present modern ergodic theory through joinings and disjointness.

In [9], we obtained a series of disjointness results for Poisson suspensions over transformations satisfying the (P) property with the additional assumption that the base transformation should have a measurable law of large numbers, which is a very particular property. We were already convinced that this assumption was not necessary. The results proved in the present paper allow to get rid of it.

The following proposition is an example of how the simplification occurs.

PROPOSITION 5.1. *Assume T has the (P) property. If an ergodic probability preserving system $(\Omega, \mathcal{F}, \mathbb{P}, S)$ is not disjoint from $(X^*, \mathcal{A}^*, \mu^*, T_*)$, then it possesses $(X^*, \mathcal{A}^*, (\alpha\mu)^*, T_*)$ as a factor for some $\alpha > 0$.*

PROOF. Consider a nontrivial joining λ of $(\Omega, \mathcal{F}, \mathbb{P}, S)$ with the Poisson suspension $(X^*, \mathcal{A}^*, \mu^*, T_*)$, and denote by $\Psi : L^2(\mu^*) \rightarrow L^2(\mathbb{P})$ the associated Markov operator. Denote by N the canonical Poisson T -point process defined on $(X^*, \mathcal{A}^*, \mu^*, T_*)$. By positivity of Ψ , the map $A \in \mathcal{A}_f \mapsto \Psi(N(A))$ extends to a T -random measure on $(\Omega, \mathcal{F}, \mathbb{P}, S)$. Indeed, for $A \in \mathcal{A}_f$ we have

$$\begin{aligned} \Psi(N(T^{-1}(A))) &= \Psi(N(A) \circ T_*) \\ &= \Psi(U_{T_*} N(A)) \end{aligned}$$

$$\begin{aligned}
 &= U_S \Psi(N(A)) \\
 &= \Psi(N(A)) \circ S,
 \end{aligned}$$

and $\mathbb{E}_{\mathbb{P}}[\Psi(N(A))] = \mathbb{E}_{\mu^*}[N(A)] = \mu(A)$.

Moreover, this T -random measure has moments of all orders, as for any $A \in \mathcal{A}_f$ and any $n \geq 1$, since Ψ can be interpreted as a conditional expectation, we have

$$\mathbb{E}_{\mathbb{P}}[(\Psi(N(A)))^n] \leq \mathbb{E}_{\mathbb{P}}[\Psi(N(A)^n)] = \mathbb{E}_{\mu^*}[N(A)^n] < \infty.$$

From Proposition 4.2, there exists $0 \leq c \leq 1$ and a T -random measure M of intensity μ defined on $(\Omega, \mathcal{F}, \mathbb{P})$, supported on discrete measures, such that

$$\Psi(N(\cdot)) = c\mu + (1 - c)M$$

If $c = 1$, then for all $A \in \mathcal{A}_f$, $\Psi(N(A) - \mu(A)) = 0$, which means that Ψ vanishes on the first chaos. Let $\Psi^* : L^2(\mathbb{P}) \rightarrow L^2(\mu^*)$ be the adjoint Markov operator, we get that $\Psi^*\Psi$ is a Markov operator on $L^2(\mu^*)$ that vanishes on the first chaos. It can be written as an integral of indecomposable operators

$$\Psi^*\Psi = \int_W \Psi_w \rho(dw),$$

where (W, \mathcal{W}, ρ) is an auxiliary probability space. Now the proof follows the same lines as Proposition 4.11 in [9]. We get that $\Psi^*\Psi$ is the projection on constants and this implies in turn that the initial joining is trivial, hence a contradiction.

Therefore, $c < 1$. We deduce that M is a factor of $(\Omega, \mathcal{F}, \mathbb{P}, S)$, and much as in Section 4.2.1, we obtain a further factor which is a Poisson point process of intensity $\alpha\mu$ for some $\alpha > 0$. (In Section 4.2.1, we got such a factor by considering $(\Phi(N_\epsilon))_0$.) \square

In particular, following the same proofs as in Theorems 5.10 and 5.14 in [9], we obtain the same conclusions without needing the assumption of measurable law of large numbers.

THEOREM 5.2. *If T has the (P) property, then $(X^*, \mathcal{A}^*, \mu^*, T_*)$ is disjoint from any rank one transformation and any Gaussian dynamical system.*

REFERENCES

- [1] ADAMS, T., FRIEDMAN, N. and SILVA, C. E. (1997). Rank-one weak mixing for nonsingular transformations. *Israel J. Math.* **102** 269–281. [MR1489108](https://doi.org/10.1007/BF02773802) <https://doi.org/10.1007/BF02773802>
- [2] BALL, K. (2005). Poisson thinning by monotone factors. *Electron. Commun. Probab.* **10** 60–69. [MR2133893](https://doi.org/10.1214/ECP.v10-1134) <https://doi.org/10.1214/ECP.v10-1134>
- [3] DALEY, D. J. and VERE-JONES, D. (2003). *An Introduction to the Theory of Point Processes. Vol. I: Elementary Theory and Methods*, 2nd ed. *Probability and Its Applications (New York)*. Springer, New York. [MR1950431](https://doi.org/10.1007/978-1-4419-9760-8)
- [4] DANILENKO, A. I. (2018). Infinite measure preserving transformations with Radon MSJ. *Israel J. Math.* **228** 21–51. [MR3874835](https://doi.org/10.1007/s11856-018-1746-5) <https://doi.org/10.1007/s11856-018-1746-5>
- [5] FURSTENBERG, H. (1967). Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation. *Math. Syst. Theory* **1** 1–49. [MR0213508](https://doi.org/10.1007/BF01692494) <https://doi.org/10.1007/BF01692494>
- [6] GLASNER, E. (2003). *Ergodic Theory via Joinings. Mathematical Surveys and Monographs* **101**. Amer. Math. Soc., Providence, RI. [MR1958753](https://doi.org/10.1090/surv/101) <https://doi.org/10.1090/surv/101>
- [7] GLASNER, S. (1983). Quasifactors in ergodic theory. *Israel J. Math.* **45** 198–208. [MR0719119](https://doi.org/10.1007/BF02774016) <https://doi.org/10.1007/BF02774016>
- [8] HOLROYD, A. E., LYONS, R. and SOO, T. (2011). Poisson splitting by factors. *Ann. Probab.* **39** 1938–1982. [MR2884878](https://doi.org/10.1214/11-AOP651) <https://doi.org/10.1214/11-AOP651>
- [9] JANVRESSE, É., ROY, E. and DE LA RUE, T. (2017). Poisson suspensions and Sushis. *Ann. Sci. Éc. Norm. Supér. (4)* **50** 1301–1334. [MR3742194](https://doi.org/10.24033/asens.2646) <https://doi.org/10.24033/asens.2646>

- [10] JANVRESSE, É., ROY, E. and DE LA RUE, T. (2019). Nearly finite Chacon Transformation. *Ann. H. Lebesgue* **2** 369–414.
- [11] KAKUTANI, S. and PARRY, W. (1963). Infinite measure preserving transformations with “mixing”. *Bull. Amer. Math. Soc.* **69** 752–756. MR0153815 <https://doi.org/10.1090/S0002-9904-1963-11022-8>
- [12] KALLENBERG, O. (2017). *Random Measures, Theory and Applications. Probability Theory and Stochastic Modelling* **77**. Springer, Cham. MR3642325 <https://doi.org/10.1007/978-3-319-41598-7>
- [13] MEYEROVITCH, T. (2011). Quasi-factors and relative entropy for infinite-measure-preserving transformations. *Israel J. Math.* **185** 43–60. MR2837127 <https://doi.org/10.1007/s11856-011-0100-y>
- [14] MEYEROVITCH, T. (2013). Ergodicity of Poisson products and applications. *Ann. Probab.* **41** 3181–3200. MR3127879 <https://doi.org/10.1214/12-AOP824>
- [15] ROY, E. (2007). Ergodic properties of Poissonian ID processes. *Ann. Probab.* **35** 551–576. MR2308588 <https://doi.org/10.1214/009117906000000692>