

## ASYMPTOTIC NORMALITY AND THE BOOTSTRAP IN STRATIFIED SAMPLING

BY P. J. BICKEL<sup>1</sup> AND D. A. FREEDMAN<sup>2</sup>

*University of California, Berkeley*

This paper is about the asymptotic distribution of linear combinations of stratum means in stratified sampling, with and without replacement. Both the number of strata and their size is arbitrary. Lindeberg conditions are shown to guarantee asymptotic normality and consistency of variance estimators. The same conditions also guarantee the validity of the bootstrap approximation for the distribution of the  $t$ -statistic. Via a bound on the Mallows distance, situations will be identified in which the bootstrap approximation works even though the normal approximation fails. Without proper scaling, the naive bootstrap fails.

**1. Introduction.** Consider the problem of estimating a linear combination  $\gamma = \sum_{i=1}^p c_i \mu_i$  of the means  $\mu_1, \dots, \mu_p$  of  $p$  numerical populations  $X_1, \dots, X_p$  with corresponding distributions  $F_1, \dots, F_p$ . For each  $i = 1, \dots, p$  there is a sample  $X_{ij}$  from population  $\mathcal{X}_i$ ; the sample elements are indexed by  $j = 1, \dots, n_i$ . Thus,  $n_i$  is the size of the sample from the  $i$ th population. Two situations will be discussed:

(a) The populations  $\mathcal{X}_i$  are assumed arbitrary and the sampling is with replacement:  $X_{ij}$  for  $j = 1, \dots, n_i$  are identically distributed with common distribution  $F_i$ ; all the  $X_{ij}$  are independent.

(b) The populations are assumed finite;  $\mathcal{X}_i$  has known size  $N_i$ ; sampling is without replacement and independent in  $i$ ; in this case,  $F_i$  is uniform. Enumerate  $X_i$  as  $\{x_{i1}, \dots, x_{iN_i}\}$ .

For simplicity, the populations are supposed univariate.

The natural unbiased estimate of  $\gamma$  is

$$(1) \quad \hat{\gamma} = \sum_{i=1}^p c_i \bar{X}_i.$$

Here, the dot is the averaging operator.

Let  $\tau_a^2$  or  $\tau_b^2$  denote the variance of  $\hat{\gamma}$  under sampling schemes (a) and (b) respectively. Let  $\hat{\tau}_a^2$  or  $\hat{\tau}_b^2$  be the customary unbiased variance estimates. Inference about  $\gamma$  can be based either on the normal approximation to the distribution of  $(\hat{\gamma} - \gamma)/\hat{\tau}$  or on bootstrap approximations. This paper will discuss the validity of these approximations when the total sample size tends to  $\infty$  in any way

---

Received February 1983; revised December 1983.

<sup>1</sup> This work was performed with the partial support of Office of Naval Research Contract N00014-80-C-0163. The hospitality of the Hebrew University, Jerusalem is also gratefully acknowledged.

<sup>2</sup> Research partially supported by National Science Foundation Grant MCS80-02535.

AMS 1980 subject classification. Primary 60F05; secondary 62E20.

Key words and phrases. Bootstrap, asymptotic normality, stratified sampling, standard errors.

whatsoever, e.g., many small samples or a few large samples or some combination thereof. More precisely: suppose  $p$ , the  $c_i$ , the populations, the  $N_i$ , and  $n_i$  all depend on an index  $\nu$  such that  $n(\nu) = n_1(\nu) + \dots + n_p(\nu) \rightarrow \infty$  as  $\nu \rightarrow \infty$ . This index will be suppressed in the sequel.

Here are two examples.

(a) The  $X_{ij}$  are unbiased measurements of the same quantity  $\mu$ , taken with  $p$  different instruments. So the precision of  $X_{ij}$ , viz.,

$$\sigma_i^2 = \int (x - \mu)^2 dF_i(x)$$

depends on  $i$ . If  $\sigma_i^2$  is known to be proportional to  $r_i$ , then

$$\hat{\gamma} = \sum \frac{n_i}{r_i} X_{i\cdot} / \sum \frac{n_i}{r_i}$$

is the natural estimate of  $\mu$ .

(b) In the classical stratified sampling model a population  $\mathcal{X}$  of size  $N$  is broken up into disjoint strata  $\mathcal{X}_1, \dots, \mathcal{X}_p$  of sizes  $N_1, \dots, N_p$  respectively;  $\sum_{i=1}^p N_i = N$ . From stratum  $i$  the sample  $X_{ij}$  for  $j = 1, \dots, n_i$  is taken without replacement. Enumerate the  $i$ th stratum as  $\{x_{i1}, \dots, x_{iN_i}\}$ . The population mean is

$$\gamma = \frac{1}{N} \sum_{i=1}^p \sum_{j=1}^{N_i} x_{ij} = \sum_{i=1}^p N_i x_{i\cdot} / N$$

and  $\hat{\gamma} = \sum_{i=1}^p N_i X_{i\cdot} / N$  is the usual estimate of  $\gamma$ .

We first take up the normal approximation in case (a). Suppose

$$(2) \quad \int x^2 dF_i < \infty \quad \text{and} \quad n_i \geq 2 \quad \text{for} \quad i = 1, \dots, p.$$

Then

$$\tau_a^2 = \sum_{i=1}^p c_i^2 \sigma_i^2 / n_i \quad \text{where} \quad \sigma_i^2 = \text{var } X_{ij}$$

and

$$\hat{\tau}_a^2 = \sum_{i=1}^p c_i^2 s_i^2 / n_i$$

where

$$s_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - X_{i\cdot})^2.$$

Let

$$\begin{aligned} \phi(x, \varepsilon) &= x \quad \text{for} \quad |x| \geq \varepsilon \\ &= 0 \quad \text{otherwise} \\ \bar{\phi}(x, \varepsilon) &= x - \phi(x, \varepsilon). \end{aligned}$$

Suppose that for all  $\varepsilon > 0$ ,

$$(3) \quad \tau_i^{-2} \sum_{i=1}^p n_i^{-1} c_i^2 E\{\phi^2(X_{ij} - \mu_i, \varepsilon n_i \tau_a | c_i |^{-1})\} \rightarrow 0.$$

By the Lindeberg-Feller theorem,  $(\hat{\gamma} - \gamma)/\tau_a$  converges in law to  $\mathcal{N}(0, 1)$ , the standard normal distribution.

According to the first main theorem of this paper, conditions (2) and (3) are also sufficient to guarantee that  $\hat{\tau}_a^2$  has the right limiting behavior. However, before giving a precise statement, it may be helpful to reformulate condition (3). Let  $Y_{ij} = (X_{ij} - \mu_i)/\sigma_i$ . Define the ‘‘variance weight’’ of the  $i$ th stratum by

$$w_i^2 = c_i^2 \sigma_i^2 / n_i \tau_a^2 = \text{var} \{c_i X_{ij} / \tau_a\}.$$

Clearly,

$$\sum_{i=1}^p w_i^2 = 1.$$

Condition (3) can then be written

$$(4) \quad \sum_{i=1}^p E\{\phi^2(w_i Y_{ij}, \varepsilon \sqrt{n_i})\} \rightarrow 0 \quad \text{for all } \varepsilon > 0.$$

**THEOREM 1.** *If (2) and (4) hold in case (a), then  $\hat{\tau}_a^2/\tau_a^2 \rightarrow 1$  in probability.*

The proof is deferred.

**COROLLARY.**  *$(\hat{\gamma} - \gamma)/\hat{\tau}_a$  tends to  $\mathcal{N}(0, 1)$  in law.*

We consider next the bootstrap approximation in case (a); also see Babu and Singh (1983). For  $i = 1, \dots, p$ , let  $\hat{F}_i$  be the empirical distribution of  $X_{ij}$  for  $j = 1, \dots, n_i$ . Take samples of size  $n_i$  with replacement from  $\hat{F}_i$ . That is, let  $\{X_{ij}^*\}$  be conditionally independent given  $\mathcal{F}$ , the  $\sigma$ -field spanned by  $\{X_{ij}\}$ ; let  $X_{ij}^*$  have common distribution  $\hat{F}_i$  for  $j = 1, \dots, n_i$ . Let

$$\begin{aligned} \hat{\gamma}^* &= \sum_{i=1}^p c_i X_{i.}^*, & s_i^{*2} &= (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij}^* - X_{i.}^*)^2 \\ \hat{\tau}_a^{*2} &= \sum_{i=1}^p c_i^2 s_i^{*2} / n_i, & \hat{\tau}_a^2 &= \sum_{i=1}^p c_i^2 (n_i - 1) s_i^2 / n_i^2. \end{aligned}$$

**THEOREM 2.** *If (2) and (4) hold in case (a), then the conditional distribution of  $(\hat{\gamma}^* - \hat{\gamma})/\hat{\tau}_a$  converges weakly to  $\mathcal{N}(0, 1)$  in probability, and  $\hat{\tau}_a^*/\hat{\tau}_a$  converges to 1 in probability.*

The proof is deferred. The theorem points to a problem in using the bootstrap to make inferences: the scaling may go wrong. This is because  $X_{i.}^*$  has variance  $(n_i - 1)s_i^2/n_i^2$ , not  $s_i^2/n_i$ . To fix ideas, suppose there are many small strata: more particularly, that  $n_i \leq k$  for all  $i$ . Now

$$\hat{\tau}_a^2 \leq (k - 1)/k \cdot \hat{\tau}_a^2 \approx (k - 1)/k \cdot \tau_a^2.$$

The bootstrap distribution of  $\hat{\gamma}^* - \hat{\gamma}$  has asymptotic scale  $\hat{\tau}_a$ , while  $\hat{\gamma} - \gamma$  has the scale  $\tau_a$ .

We take up next the normal approximation in case (b). Suppose

$$(5) \quad 2 \leq n_i \leq N_i - 1.$$

Then

$$\tau_b^2 = \sum_{i=1}^p c_i^2 \frac{\sigma_i^2 (N_i - n_i)}{n_i (N_i - 1)}$$

and

$$\hat{\tau}_b^2 = \sum_{i=1}^p c_i^2 \frac{s_i^2 (N_i - n_i)}{n_i N_i}.$$

To state the regularity condition, let  $v_i^2$  be the “variance weight” in case (b):  $v_i^2 = c_i^2 \sigma_i^2 (N_i - n_i) / n_i \tau_b^2 (N_i - 1) = \text{var}\{c_i X_{i\cdot} / \tau_b\}$ . Let  $\rho_i$  be “the effective sample size:”  $\rho_i = n_i (N_i - 1) / (N_i - n_i)$ . Let  $\mathcal{Y}_i = \{y_{i1}, \dots, y_{iN_i}\}$  where  $y_{ij} = (x_{ij} - \mu_i) / \sigma_i$  and  $\sigma_i^2 = N_i^{-1} \sum_{j=1}^{N_i} (x_{ij} - \mu_i)^2$ . So  $Y_{ij} = (X_{ij} - \mu_i) / \sigma_i$  are sampled from  $\mathcal{Y}_i$ .

The condition is

$$(6) \quad \sum_{i=1}^p N_i^{-1} \sum_{j=1}^{N_i} \phi^2(v_i y_{ij}, \varepsilon \sqrt{\rho_i}) \rightarrow 0.$$

This may be compared with condition (4).

If  $\sup_{1 \leq i \leq p} E |Y_i|^3$  is uniformly bounded independent of the hidden index  $\nu$ , the Lindeberg conditions (4) and (6) are implied respectively by the natural conditions  $\max_i w_i / \sqrt{n_i} \rightarrow 0$  or  $\max_i v_i / \sqrt{\rho_i} \rightarrow 0$ . Thus if the standardized populations have reasonably light tails, asymptotic normality holds if for each stratum the variance weight contribution is small or the stratum is heavily sampled.

**THEOREM 3.** *If (5) and (6) hold in case (b), then*

$$i) \quad (\hat{\gamma} - \gamma) / \tau_b \rightarrow \mathcal{N}(0, 1) \text{ in law}$$

and

$$ii) \quad \hat{\tau}_b / \tau_b \rightarrow 1 \text{ in probability.}$$

The proof is deferred.

**COROLLARY.**  $(\hat{\gamma} - \gamma) / \hat{\tau}_b \rightarrow \mathcal{N}(0, 1)$  in law.

Finally, we consider the bootstrap in case (b). If  $N_i/n_i = k_i$  an integer for each  $i$ , the natural bootstrap procedure was suggested by Gross (1980): given  $\{X_{ij}\}$ , to create populations  $\mathcal{X}_i$  consisting of  $k_i$  copies of each  $X_{ij}$  for  $j = 1, \dots, n_i$ , then  $X_{ij}^*$  for  $j = 1, \dots, n_i$  are generated as a sample without replacement from  $\mathcal{X}_i$ , the samples being independent for different  $i = 1, \dots, p$ . In general, if  $N_i = k_i n_i + r_i$  with  $0 \leq r_i < n_i$ , form populations  $\mathcal{X}_{i0}$  and  $\mathcal{X}_{i1}$ , where  $\mathcal{X}_{i0}$  consists of  $k_i$

copies of each  $X_{ij}$ , for  $j = 1, \dots, n_i$ ; while  $\mathcal{X}_{i1}$  consists of  $k_i + 1$  copies. Let

$$\alpha_i = \left(1 - \frac{r_i}{n_i}\right) \left(1 - \frac{r_i}{N_i - 1}\right).$$

With probability  $\alpha_i$ , let  $(X_{i1}^*, \dots, X_{in_i}^*)$  be a sample without replacement of size  $n_i$  from  $\hat{\mathcal{X}}_{i0}$ ; with probability  $1 - \alpha_i$ , let  $(X_{i1}^*, \dots, X_{in_i}^*)$  be a sample without replacement of size  $n_i$  from  $\hat{\mathcal{X}}_{i1}$ . The virtue of this scheme is that both  $\hat{\mathcal{X}}_{i0}$  and  $\hat{\mathcal{X}}_{i1}$  have the same distribution  $\hat{F}_i$  and

$$\text{Var}(X_{i.}^* | \{X_{ij}\}) = \frac{n_i - 1}{n_i^2} s_i^2 \left(\frac{N_i - n_i}{N_i - 1}\right).$$

The proof of the following theorem is similar to that of Theorem 2 and is omitted. Define  $\hat{\gamma}^*$  as before, and  $\hat{\tau}_b^{*2}$  by substituting  $X_{ij}^*$  for  $X_{ij}$  in  $\hat{\tau}_b^2$ .

**THEOREM 4.** *Let  $\hat{\tau}_b^2$  be the variance of  $\hat{\gamma}^*$  given the data. Then, if (5) and (6) hold in case (b), the conditional distribution of  $(\hat{\gamma}^* - \hat{\gamma})/\hat{\tau}_b$  converges weakly to  $\mathcal{N}(0, 1)$  and  $\hat{\tau}_b^*/\hat{\tau}_b \rightarrow 1$  in probability.*

The same inference problem arises as in the case of Theorem 2. The variance of  $\hat{\gamma}^*$  given the data is an inconsistent estimate of the variance of  $\hat{\gamma}$ . We have side-stepped the issue by computing the scale externally to the bootstrap process. Other patches could be made: one is to rescale the elements of  $\mathcal{X}_i$ ; another is to adjust the constants  $c_i$ . These fixes are all a bit *ad hoc*.

If  $\gamma$  stays bounded as  $\nu \rightarrow \infty$ , our results extend easily to pivots

$$\frac{g(\hat{\gamma}) - g(\gamma)}{g'(\gamma)\hat{\tau}_b}$$

where  $g$  is nonlinear continuously differentiable. The same issue as before arises a fortiori for nonlinear functions. Neither the variance of  $g(\hat{\gamma}^*)$  given the data nor its natural approximation  $[g'(\hat{\gamma})]^2 \hat{\tau}_b^2$  are consistent estimates of the asymptotic variance of  $g(\hat{\gamma})$ . A fix which works if  $\sum_{i=1}^p |c_i \mu_i|$  stays bounded is as before to rescale the elements of  $\mathcal{X}_i$  or the  $c_i$  before applying the bootstrap. Alternatives (the jackknife, linearization, BRR) are discussed in Krewski and Rao (1981). For the case of one stratum, Theorem 4 was derived independently by Chao and Lo (1983).

The bootstrap can work even when Theorem 4 fails but the circumstances are artificial. Suppose we have only one stratum and  $N_1 - n_1 = k$  for all  $\nu$  i.e., all but  $k$  members are sampled. Since  $\sum_{j=1}^{N_1} (x_{1j} - \mu_1) = 0$ , the pivot  $(\hat{\gamma} - \gamma)/\tau_b$  is distributed as the standardized mean of a sample of size  $k$  taken without replacement from the population  $\mathcal{X}_1$ . No matter how large  $N_1$  is, if  $k$  is small and  $\mathcal{X}_1$  nonnormal, we would not expect the normal approximation to apply to  $\hat{\gamma}$ . To be specific let  $F_\nu$  be the uniform distribution on  $\mathcal{X}_1$  and suppose

(7)  $F_\nu$  converges to  $F$  in the Mallows  $d_2$ -metric,

i.e.,  $F_\nu \rightarrow F$  weakly and  $\int x^2 dF_\nu \rightarrow \int x^2 dF$ . Then  $(\hat{\gamma} - \gamma)/\tau_b$  is distributed in the

limit as the standardized mean of  $k$  independent variables identically distributed according to  $F$ . On the other hand, since we have sampled nearly the whole population we expect the bootstrap to work.

**THEOREM 5.** *If (7) holds, the conditional distribution of  $(\hat{\gamma}^* - \hat{\gamma})/\hat{\tau}_b$  converges weakly in probability to the same limit as that of the unconditional distribution of  $(\hat{\gamma} - \gamma)/\tau_b$ . Moreover,  $\hat{\tau}_b/\tau_b$  and  $\hat{\tau}_b^*/\hat{\tau}_b$  both tend to 1 in probability.*

We can extend this result somewhat by replacing (7) with a compactness-in- $d_2$  condition on  $\{F_\nu\}$

$$\limsup_{m \rightarrow \infty} \limsup_{\nu} N_1^{-1} \sum_{j=1}^{N_1} \phi^2(v_{1j} y_{1j}, m) = 0.$$

This condition is evidently weaker than (6) for  $p = 1$ . The conclusion now is that the  $d_2$ -distance between the conditional distribution of  $(\hat{\gamma}^* - \hat{\gamma})/\hat{\tau}_b^*$  and the unconditional distribution of  $(\hat{\gamma} - \gamma)/\hat{\tau}_b$  tends in probability to 0. A further extension to an arbitrary number of strata which includes both Theorems 4 and 5 is also possible but not worthwhile.

**2. Some lemmas.** Recall the truncation operator  $\phi$  from Section 1.

**LEMMA 1.** a) 
$$\left| \phi\left(\frac{1}{k} \sum_{i=1}^k y_i, \varepsilon\right) \right| \leq \sum_{i=1}^k |\phi(y_i, \varepsilon/k)|; \quad \text{equivalently,}$$

$$|\phi(\sum_{i=1}^k y_i, \varepsilon)| \leq k \sum_{i=1}^k |\phi(y_i, \varepsilon/k^2)|$$

b) Let  $Y_1, Y_2, \dots$  be independent and identically distributed. Then

$$E \left\{ \phi^2\left(\frac{1}{k} \sum_{i=1}^k Y_i, \varepsilon\right) \right\} \leq k^2 E\{\phi^2(Y_i, \varepsilon/k)\}.$$

**PROOF.** Claim a). As is easily verified,

$$\left| \phi\left(\frac{1}{k} \sum_{i=1}^k y_i, \varepsilon\right) \right| \leq \phi\left(\frac{1}{k} \sum_{i=1}^k |y_i|, \varepsilon\right).$$

Without loss of generality, suppose all  $y_i \geq 0$ . Let  $y_{(k)}$  be the largest  $y_i$ . If  $y_{(k)} < \varepsilon/k$ , both sides of the inequality vanish. If  $y_{(k)} \geq \varepsilon/k$ , the left side is the average of the  $y_i$ , or zero; the right side is at least the maximum  $y_{(k)}$ .

Claim b) follows by the Cauchy-Schwarz inequality.  $\square$

**LEMMA 2.** Let  $(X'_1, \dots, X'_n)$  and  $(X_1, \dots, X_n)$  be distributed respectively as samples with and without replacement from a finite population. Then

$$E\{\phi^2(\sum_{i=1}^n X_i, \varepsilon)\} \leq E\{\phi^2(\sum_{i=1}^n X'_i, \frac{1}{2}\varepsilon)\}.$$

**PROOF.** By a theorem of Hoeffding (1963), if  $g$  is convex, then

$$E\{g(\sum X_i)\} \leq E\{g(\sum X'_i)\}.$$

Let

$$\begin{aligned} g(x, \varepsilon) &= x^2 && \text{for } |x| \geq \varepsilon \\ &= 2\varepsilon|x| - \varepsilon^2 && \text{for } 1/2\varepsilon \leq |x| \leq \varepsilon \\ &= 0 && \text{otherwise.} \end{aligned}$$

Then  $g$  is convex and

$$\phi^2(x, \varepsilon) \leq g(x, \varepsilon) \leq \phi^2(x, 1/2\varepsilon).$$

So

$$E\{\phi^2(\sum X_i, \varepsilon)\} \leq E\{g(\sum X_i, \varepsilon)\} \leq E\{g(\sum X'_i, \varepsilon)\} \leq E\{\phi^2(\sum X'_i, 1/2\varepsilon)\}. \quad \square$$

The next result involves the Mallows metric  $d_2$ ; see Mallows (1972) or Bickel and Freedman (1981).

**LEMMA 3.** *Let  $\mathcal{X}$  and  $\mathcal{Y}$  be two finite populations of real numbers, of the same size  $N$ . Let  $F$  and  $G$  be the uniform distributions on  $\mathcal{X}$  and  $\mathcal{Y}$ . Suppose  $F$  and  $G$  have the same means. Let  $X_1, \dots, X_n$  be a sample of size  $n$ , drawn at random without replacement from  $\mathcal{X}$ ; let  $F_{(n)}$  be the law of  $X_1 + \dots + X_n$ . Likewise for  $Y_1, \dots, Y_n$  and  $G_{(n)}$ . Then*

$$d_2[F_{(n)}, G_{(n)}]^2 \leq \frac{n(N-n)}{N-1} d_2(F, G)^2.$$

**PROOF.** Enumerate  $\mathcal{X}$  as  $x_1 \leq x_2 \leq \dots \leq x_N$  and  $\mathcal{Y}$  as  $y_1 \leq y_2 \leq \dots \leq y_N$ . Then

$$(1/N) \sum_{i=1}^N (x_i - y_i)^2 = d_2(F, G)^2.$$

This follows from Bickel and Freedman (1981, Lemmas 8.2 and 8.3). Let  $Z = \{1, \dots, N\}$ . Let  $Z_1, \dots, Z_n$  be a sample of size  $n$ , drawn at random without replacement from  $Z$ . Set  $X_i = x_{Z_i}$  and  $Y_i = y_{Z_i}$ . Now

$$\begin{aligned} d_2[F_{(n)}, G_{(n)}]^2 &\leq E\{[\sum_{i=1}^n (X_i - Y_i)]^2\} = \frac{n(N-n)}{N-1} E[(X_i - Y_i)^2] \\ &= \frac{n(N-n)}{N-1} d_2(F, G)^2. \quad \square \end{aligned}$$

Here is an easy generalization of Lemma 3.

**LEMMA 4.** *For  $i = 0, 1$  let  $\mathcal{X}_i = \{x_{i1}, \dots, x_{iN_i}\}$  be finite populations and  $F_i$  the associated uniform distributions on  $\mathcal{X}_i$ . Let  $F_{ni}$  be the distribution of  $\sum_{j=1}^n X_j$  when  $X_1, \dots, X_n$  is a sample without replacement from  $\mathcal{X}_i$ . Let  $n \leq N_0 \leq N_1$ . If  $J$  is a subset of  $\{1, \dots, N_1\}$ , let  $F_{1J}$  be the uniform distribution on  $\{x_{1j}; j \in J\}$ .*

Then,

$$d_2(F_{n_0}, F_{n_1})^2 \leq \frac{n(N_0 - n)}{N_0 - 1} \frac{1}{\binom{N_1}{N_0}} \sum_J \{d_2(F_0, F_{1J})^2: |J| = N_0\}.$$

**LEMMA 5.** For  $\nu \geq 1$  let  $\mathcal{X}_\nu$  be a finite population of size  $N_\nu$ ,  $F_\nu$  the uniform distribution on  $\mathcal{X}_\nu$ ,  $X_1, \dots, X_{n_\nu}$ , a sample without replacement from  $\mathcal{X}_\nu$ ,  $\hat{F}_\nu$  the empirical df of the sample. If for some  $F$ ,  $d_2(F_\nu, F) \rightarrow 0$  as  $\nu \rightarrow \infty$  and  $n_\nu \rightarrow \infty$  then  $d_2^2(\hat{F}_\nu, F) \rightarrow 0$  in probability.

**PROOF.** If  $g$  is continuous and bounded

$$E \int g(x) d\hat{F}_\nu(x) = \int g(x) dF_\nu(x) \rightarrow \int g(x) dF(x),$$

$$\text{Var} \left( \int g(x) d\hat{F}_\nu(x) \right) \rightarrow 0.$$

So,

$$(8) \quad \int g(x) d\hat{F}_\nu(x) \rightarrow \int g(x) dF(x)$$

in probability. Moreover,

$$\limsup_\nu E \int \phi(x, M)^2 d\hat{F}_\nu(x) = \int \phi(x, M)^2 dF(x)$$

by Lemma 8.3c) of Bickel and Freedman (1981). Since we can make  $\int \phi(x, M)^2 dF(x)$  small for  $M$  large we conclude that (8) holds for  $g(x) = x^2$  also and the lemma follows.  $\square$

### 3. Proving the theorems in case (a).

**PROOF OF THEOREM 1.** Recall the variance weights  $w_i$  from Section 1. As is easily verified,  $\hat{\tau}_a^2/\tau_a^2 = 1 + \xi - \zeta$ , where

$$(9a) \quad \xi = \sum_{i=1}^p w_i^2 (n_i - 1)^{-1} \sum_{j=1}^{n_i} (Y_{ij}^2 - 1)$$

$$(9b) \quad \zeta = \sum_{i=1}^p w_i^2 (n_i - 1)^{-1} (n_i Y_i^2 - 1).$$

To prove the theorem, it is enough to show that  $\xi$  and  $\zeta$  are both small. But  $\xi = \xi_1 + \xi_2$ , where

$$(10a) \quad \xi_1 = \sum_{i=1}^p (n_i - 1)^{-1} \sum_{j=1}^{n_i} [\bar{\phi}^2(w_i Y_{ij}, \varepsilon \sqrt{n_i}) - E\{\bar{\phi}^2(w_i Y_{ij}, \varepsilon \sqrt{n_i})\}]$$

$$(10b) \quad \xi_2 = \sum_{i=1}^p (n_i - 1)^{-1} \sum_{j=1}^{n_i} [\phi^2(w_i Y_{ij}, \varepsilon \sqrt{n_i}) - E\{\phi^2(w_i Y_{ij}, \varepsilon \sqrt{n_i})\}].$$



Now

$$\begin{aligned}
 E(\xi_1^2) &= \text{var } \xi_1 = \sum_{i=1}^p (n_i - 1)^{-2} \sum_{j=1}^{n_i} \text{var}\{\bar{\phi}^2(w_i Y_{ij}, \varepsilon \sqrt{n_i})\} \\
 &\leq \sum_{i=1}^p (n_i - 1)^{-2} n_i E\{\bar{\phi}^4(w_i Y_{ij}, \varepsilon \sqrt{n_i})\} \\
 &\leq \varepsilon^2 \sum_{i=1}^p (n_i - 1)^{-2} n_i^2 E\{\bar{\phi}^2(w_i Y_{ij}, \varepsilon \sqrt{n_i})\} \\
 &\leq \varepsilon^2 \sum_{i=1}^p (n_i - 1)^{-2} n_i^2 w_i^2 E\{Y_{ij}^2\} \\
 &\leq 4\varepsilon^2 \sum_{i=1}^p w_i^2 = 4\varepsilon^2.
 \end{aligned}$$

On the other hand,  $E\{|\xi_2|\} \rightarrow 0$  for each  $\varepsilon > 0$ , by (4). This disposes of  $\xi$ .

The term  $\zeta$  in (9b) can be decomposed according to whether  $n_i > M$  or  $n_i \leq M$ . Since

$$\sum_i \{(n_i - 1)^{-1} w_i^2: n_i \geq M + 1\} \leq M^{-1}$$

and  $E\{n_i Y_{i\cdot}^2\} = 1$ , the strata  $i$  with  $n_i \geq M + 1$  are negligible. For the  $i$  with  $n_i \leq M$ ,  $\zeta = \zeta_1 + \zeta_2$  where

$$(11a) \quad \zeta_1 = \sum_i \frac{n_i}{n_i - 1} [\bar{\phi}^2(w_i Y_{i\cdot}, \varepsilon \sqrt{n_i}) - E\{\bar{\phi}^2(w_i Y_{i\cdot}, \varepsilon \sqrt{n_i})\}]$$

$$(11b) \quad \zeta_2 = \sum_i \frac{n_i}{n_i - 1} [\phi^2(w_i Y_{i\cdot}, \varepsilon \sqrt{n_i}) - E\{\phi^2(w_i Y_{i\cdot}, \varepsilon \sqrt{n_i})\}].$$

The sums need be extended only over  $i$  with  $2 \leq n_i \leq M$ . Now whatever  $n_i$  may be, as for  $\xi_1$ ,

$$(12) \quad E\{\zeta_1^2\} \leq 4\varepsilon^2$$

is small. Next,

$$\begin{aligned}
 (13) \quad E\{|\zeta_2|\} &\leq 2 \sum_i \frac{n_i}{n_i - 1} E\{\phi^2(w_i Y_{i\cdot}, \varepsilon \sqrt{n_i})\} \\
 &\leq 4M^2 \sum_i E\{\phi^2(w_i Y_{ij}, \varepsilon \sqrt{n_i}/M)\}
 \end{aligned}$$

because  $2 \leq n_i \leq M$ ; see Lemma 1. So  $\zeta_2$  is small too, by condition (4).  $\square$

**PROOF OF THEOREM 2.** The Lindeberg condition is applied, given  $\mathcal{F}$ . It is enough to check that for every  $\varepsilon > 0$ ,

$$(14) \quad \tilde{\tau}_a^{-2} \sum_{i=1}^p n_i^{-1} c_i^2 E\{\phi^2(X_{ij}^* - X_{i\cdot}, \varepsilon n_i \tilde{\tau}_a | c_i |^{-1}) | \mathcal{F}\} \rightarrow 0$$

in probability, where  $\tilde{\tau}_a^2 = \sum_{i=1}^p c_i^2 (n_i - 1) s_i^2 / n_i^2$  is the conditional variance of  $\hat{\gamma}^*$  given  $\mathcal{F}$ . For then, Theorem 1 can be applied to  $X_{ij}^*$ .

Since  $n_i \geq 2$ ,

$$(15) \quad \frac{1}{2} \hat{\tau}_a \leq \tilde{\tau}_a \leq \hat{\tau}_a.$$

Thus  $\hat{\tau}_a$  and hence  $\tau_a$  may be substituted in (14) for  $\tilde{\tau}_a$ . So (14) reduces to

$$\tau_a^{-2} \sum_{i=1}^p c_i^2 n_i^{-2} \sum_{j=1}^{n_i} \phi^2(X_{ij} - X_{i\cdot}, \varepsilon n_i \tau_a |c_i|^{-1}) \rightarrow 0$$

in probability. This in turn reduces to

$$(16) \quad \sum_{i=1}^p n_i^{-1} \sum_{j=1}^{n_i} \phi^2[w_i(X_{ij} - X_{i\cdot})/\sigma_i, \varepsilon \sqrt{n_i}] \rightarrow 0$$

in probability.

Now  $(X_{ij} - X_{i\cdot})/\sigma_i = Y_{ij} - Y_{i\cdot}$ . Use Lemma 1a) with  $k = 2$  to see that (16) follows from (17) and (18):

$$(17) \quad \sum_{i=1}^p n_i^{-1} \sum_{j=1}^{n_i} \phi^2(w_i Y_{ij}, \frac{1}{4}\varepsilon \sqrt{n_i}) \rightarrow 0 \quad \text{in probability}$$

and

$$(18) \quad \sum_{i=1}^p \phi^2(w_i Y_{i\cdot}, \frac{1}{4}\varepsilon \sqrt{n_i}) \rightarrow 0 \quad \text{in probability.}$$

Clearly, (17) follows from (4). We bound the expected value of the left side of (18). Take first those  $i$  with  $n_i \leq M$ . In view of Lemma 1b), the sum over such  $i$  is bounded above by

$$M^2 \sum_i E\{\phi^2(w_i Y_{ij}, \frac{1}{4}\varepsilon \sqrt{n_i}/M)\}$$

which tends to zero by condition (4). Take next those  $i$  with  $n_i > M$ . The sum over such  $i$  is bounded above by

$$\sum_i E\{(w_i Y_{i\cdot})^2\} = \sum_i w_i^2 n_i^{-1} < M^{-1} \sum_i w_i^2 \leq M^{-1}$$

which is small for  $M$  large.

That  $\tilde{\tau}_a^*/\tilde{\tau}_a \rightarrow 1$  follows from Theorem 1.  $\square$

**REMARKS.** (i) The Lindeberg-Feller theorem can be supplemented by direct bounds generalizing those of Berry-Esseen; see Petrov (1975, Theorem 3, page 111 or Theorem 8, page 118). These bounds may give estimates on the discrepancy between the bootstrap distribution and the true distribution.

(ii) The difference between the distribution of  $(\hat{\gamma} - \gamma)/\tau_a$  and the bootstrap distribution of  $(\hat{\gamma}^* - \hat{\gamma})/\tilde{\tau}_a$  can be estimated using the Mallows metric as in equation (2.2) of Bickel and Freedman (1981). The condition needed to push this through is stronger than (4).

(iii) The results can be extended in an obvious way to vector  $X_{ij}$ , and under further conditions to nonlinear statistics such as  $\sum_{i=1}^p [g_i(X_{i\cdot}) - g_i(\mu_i)]$ ; this covers ratio estimates.

#### 4. Proving the theorems in case (b)

**PROOF OF THEOREM 3.** The Lindeberg-Feller theorem does not apply to give us i) directly here, since the  $X_{ij}$  are dependent for fixed  $i$ ; however, essentially

the same ideas can be used. The proof we give is a bit complicated; an alternative but we believe no simpler approach is given by Dvoretzky (1971). Our argument is by cases, and the focus is on asymptotic normality. Without loss of generality, assume  $\mu_i \equiv 0$ ,  $c_i \equiv 1$ . In outline, the argument is as follows.

CASE 1. There is only one stratum, and  $n \leq \frac{1}{2}N$ ; we drop the unnecessary stratum subscript  $i$ . Then  $\rho^2$  is of order  $n$ , and asymptotic normality follows from Erdős-Rényi (1959). Also see Rosén (1967), Dvoretzky (1971).

CASE 2. There is only one stratum, and  $n > \frac{1}{2}N$ . Apply Case 1 to the "co-sample" consisting of the objects not in the sample.

CASE 3. The number of strata is bounded; no variance weight tends to zero. Case 1 or Case 2 applies to each stratum individually.

CASE 4. There are many strata, each of small variance weight; in each stratum,  $n_i \leq \frac{1}{2}N_i$ . Then  $\hat{\gamma}/\tau_b$  is the sum of  $p$  independent u.a.n. summands:  $\text{var} \{X_i/\tau_b\} = v_i^2$  being uniformly small by assumption. We must verify the Lindeberg condition on  $X_i/\tau_b$ , and do so by an indirect argument. Let  $X'_{ij}$  be sampled with replacement from  $\mathcal{X}_i$ . And let

$$\hat{\gamma}' = \sum_{i=1}^p \frac{1}{n_i} \sum_{j=1}^{n_i} X'_{ij}.$$

Since  $n_i \leq \frac{1}{2}N_i$ , the variance weights  $v_i^2$  and  $w_i^2$  are of the same order, as are the total variances  $\tau_a^2$  and  $\tau_b^2$ . In particular, condition (6) implies (4). Thus, the Lindeberg condition holds for the individual summands in  $\hat{\gamma}'/\tau_a$ , viz.,  $X'_{ij}/n_i\tau_a$ , and asymptotic normality of  $\hat{\gamma}'$  follows. By the converse to Lindeberg's theorem, his condition holds for the stratum averages  $(1/n_i) \sum_{j=1}^{n_i} X'_{ij}/\tau_a$ . Hence, by Lemma 2, the condition holds for the stratum averages taken without replacement, viz.,  $(1/n_i) \sum_{j=1}^{n_i} X_{ij}/\tau_b$ . Now a second application of the direct Lindeberg theorem gives asymptotic normality of  $\hat{\gamma}$ .

CASE 5. There are many strata, each of small variance weight; on each stratum,  $n_i > \frac{1}{2}N_i$ . Apply Case 4 to the co-samples.

CASE 6. There are many strata, each of small variance weight. Consider two groups of strata: in the first,  $n_i \leq \frac{1}{2}N_i$ ; in the second,  $n_i > \frac{1}{2}N_i$ . Case 4 applies to the first group, Case 5 to the second. (One of the two groups may be negligible.)

*The general case.* We combine cases 3 and 6. Let

$$J_k(\nu) = \left\{ i: v_i \geq \frac{1}{k} \right\}; \quad V_k(\nu) = \sum \{v_i^2: i \in J_k(\nu)\}$$

where dependence on the hidden index is made explicit. Given any subsequence of  $\{\nu\}$  we can extract a subsequence  $\{\nu_r\}$  such that for all  $k$ , as  $r \rightarrow \infty$ ,  $V_k(\nu_r)$  tends

to a finite limit  $V_k$ . If  $V_k = 0$  for all  $k$ , there must be  $k_r \rightarrow \infty$  such that  $V_{k_r}(\nu_r) \rightarrow 0$ . Hence, as  $r \rightarrow \infty$ ,

$$(19) \quad \sum \{X_{i\cdot}/\tau_b: i \in J_{k_r}(\nu_r)\} \rightarrow 0 \quad \text{in probability.}$$

But,  $\max\{v_i: i \notin J_{k_r}(\nu_r)\} \leq 1/k_r \rightarrow 0$ . So we can apply case 6 to get that

$$(20) \quad \sum \{X_{i\cdot}/\tau_b: i \notin J_{k_r}(\nu_r)\} \quad \text{is asymptotically } N(0, 1).$$

Combining (19) and (20), we get

$$(21) \quad \sum X_{i\cdot}/\tau_b \quad \text{is asymptotically } N(0, 1), \quad \text{as } r \rightarrow \infty.$$

On the other hand, suppose  $V_k > 0$  for some  $k$ . Since  $J_k(\nu_r)$  has at most  $k^2$  members, we can apply case 3 to see that for all  $k$ , as  $r \rightarrow \infty$ ,

$$\sum \{X_{i\cdot}/\tau_b: i \in J_k(\nu_r)\} \quad \text{is asymptotically } N(0, V_k).$$

By a standard argument, there are  $k_r \rightarrow \infty$  such that

$$(22) \quad \sum \{X_{i\cdot}/\tau_b: i \in J_{k_r}(\nu_r)\} \quad \text{is asymptotically } N(0, \sup_k V_k).$$

Applying case 6 as above,

$$(23) \quad \sum \{X_{i\cdot}/\tau_b: i \notin J_{k_r}(\nu_r)\} \quad \text{is asymptotically } N(0, 1 - \sup_k V_k).$$

Combining (22) and (23) we obtain (21) in this case also. Part (i) of the theorem follows by a standard compactness argument. The proof of (ii) follows the pattern of that of Theorem 1 and is omitted.  $\square$

**PROOF OF THEOREM 5.** We simplify the argument by supposing  $n_1$  divides  $N_1$  so we can use the naive bootstrap. (The general argument uses Lemma 4.) Moreover, without loss of generality let  $\mu_1 = 0, \sigma_1 = 1$ . Since  $p = 1$  we want to compare the distribution of the standardized mean of a sample of size  $n_1$  from the population  $\mathcal{Y}_1$  and the distribution of the standardized mean of a sample of size  $n_1$  from the population composed of  $N_1/n_1$  copies of the standardized sample:  $(X_{ij} - \hat{\mu}_1)/\hat{\sigma}_1, 1 \leq j \leq n_1$ , where  $\hat{\mu}_1$  is the sample mean and  $\hat{\sigma}_1$  is sample standard deviation. So by Lemma 3,

$$d_2^2 \left\{ \mathcal{L} \left( \frac{\hat{\gamma} - \gamma}{\tau_b} \right), \mathcal{L} \left( \frac{\hat{\gamma}^* - \hat{\gamma}}{\hat{\tau}_b} \right) \mid X_{1j}, 1 \leq j \leq n_1 \right\} \leq d_2^2 \{F_\nu, \hat{F}_\nu, \hat{F}_\nu(\hat{\sigma}_1 x + \hat{\mu}_1)\}.$$

By Lemma 5,  $d_2^2(F_\nu, \hat{F}_\nu), \hat{\mu}$ , and  $\hat{\sigma}_1 - 1$  all tend in probability to 0 as  $\nu \rightarrow \infty$ . A truncation argument of the type we have already used shows that  $\hat{\tau}_b/\tau_b$  and  $\hat{\tau}_b^*/\hat{\tau}_b$  both tend in probability to 1. The theorem follows.  $\square$

REFERENCES

BABU, G. J. and SINGH, K. (1983). Inference on means using the bootstrap. *Ann. Statist.* **11** 999-1003.  
 BICKEL, P. J. and FREEDMAN, D. A. (1981). Some asymptotic theory for the bootstrap. *Ann. Statist.* **9** 1196-1217.

- BICKEL, P. J. and KRIEGER, A. (1983). Using the bootstrap to set confidence bands for a distribution function: Monte Carlo and some theory. In preparation.
- CHAO, M. T. and LO, S. H. (1983). A bootstrap method for finite population. Preprint.
- DVORETZKY, A. (1971). Asymptotic normality for sums of dependent random variables. *Proc. Sixth Berkeley Symp. Math. Probab.* II 513. (Ed: Le Cam, Neyman, Scott). Univ. of Calif. Press, Berkeley.
- ERDŐS, P. and RENYI, A. (1959). On the central limit theorem for samples from a finite population. *Publ. Math. Inst. Hung. Acad. Sci.* 4 49–61.
- GROSS, S. (1980). Median estimation in sample surveys. Paper presented at 1980 Amer. Statist. Assoc. meeting.
- HOEFFDING, W. (1963). Probability inequalities for sums of bounded random variables. *J. Amer. Statist. Assoc.* 58 13–30.
- KREWSKI, D. and RAO, J. N. K. (1981). Inference from stratified samples: properties of linearization, jackknife, and balanced repeated replication. *Ann. Statist.* 9 1010–1019.
- MALLOWS, C. (1972). A note on asymptotic joint normality. *Ann. Math. Statist.* 43 508–515.
- PETROV, V. (1975). *Sums of Independent Random Variables*. Springer, Berlin.
- ROSÉN, B. (1967). On the central limit theorem for sums of dependent random variables. *Z. Wahrsch. verw. Gebiete* 7 48–82.

DEPARTMENT OF STATISTICS  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720