

Inverse exponential decay: Stochastic fixed point equation and ARMA models

KRZYSZTOF BURDZY¹, BARTOSZ KOŁODZIEJEK² and TVRTKO TADIĆ³

¹*Department of Mathematics, University of Washington, Box 354350, Seattle, WA 98195, USA.*

E-mail: burdzy@uw.edu

²*Faculty of Mathematics and Information Science, Warsaw University of Technology, Koszykowa 75, 00-662 Warsaw, Poland. E-mail: b.kolodziejek@mini.pw.edu.pl*

³*Microsoft Corporation, One Microsoft Way, Redmond, WA 98052, USA. E-mail: tvrtko@math.hr*

We study solutions to the stochastic fixed point equation $X \stackrel{d}{=} AX + B$ when the coefficients are nonnegative and B is an “inverse exponential decay” (IED) random variable. We provide theorems on the left tail of X which complement well-known tail results of Kesten and Goldie. We generalize our results to ARMA processes with nonnegative coefficients whose noise terms are from the IED class. We describe the lower envelope for these ARMA processes.

Keywords: ARMA models; inverse-gamma distribution; iterated random sequences; stochastic fixed point equation; tail estimates; time series

1. Introduction

The paper is devoted to studying properties, especially left tails, of positive random variables that arise in several closely related contexts – stochastic fixed point equations, ARMA models, and iterated random functions.

For a two-dimensional random vector (A, B) , an independent random variable X is said to satisfy the stochastic fixed point equation if

$$X \stackrel{d}{=} AX + B. \quad (1.1)$$

The behavior of the solution, especially the left and right tails, has been extensively studied. A classical result ([7,9]) says that under some assumptions on (A, B) , for some α , C_- , $C_+ > 0$,

$$\mathbb{P}(X > x) \sim C_+ x^{-\alpha} \quad \text{and} \quad \mathbb{P}(X < -x) \sim C_- x^{-\alpha}, \quad (1.2)$$

as $x \rightarrow \infty$. (See the precise statement in Theorem 7.14. Here \sim means that the ratio of the two quantities converges to 1.) An excellent review of the subject can be found in a recent book [4].

It can be shown that if A and B are nonnegative random variables then the solution X to (1.1) is also a nonnegative random variable. Under this extra assumption on (A, B) , the first estimate in (1.2) is still meaningful and informative. But the second one is not because for $x > 0$ we have $\mathbb{P}(X < -x) = 0$. It is natural to ask for a meaningful estimate for the left tail under these circumstances. We will examine the behavior of $\mathbb{P}(X < x)$ as $x \rightarrow 0^+$. This question does not

seem to be addressed anywhere in the literature; in particular, it does not seem to be examined in [4].

The motivation for the present paper comes from a project on a “Fleming–Viot” type process defined in [5]. We will explain in Section 8 how the problem arises in the setting of [1].

1.1. Review of the main results

This paper revolves around $\text{IED}_L^\rho(\lambda)$ random variables defined as follows.

Definition 1.1. We will say that a nonnegative random variable X has an inverse exponential decay of the left tail with index $\rho > 0$ if

$$\lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X < x) = -\lambda, \quad (1.3)$$

for a slowly varying function L at 0 and $\lambda \geq 0$. We will call such a random variable $\text{IED}_L^\rho(\lambda)$. In the case $\lim_{x \rightarrow 0^+} L(x) = 1$, we will write $\text{IED}_1^\rho(\lambda)$.

The best known $\text{IED}_L^\rho(\lambda)$ distributions are called “inverse-gamma;” in this case, $\rho = 1$ and $L(x) \equiv 1$ (see Definition 3.1).

In Section 4 (see especially Theorems 4.4 and 4.7), we will find conditions for a sequence (X_i) of independent $\text{IED}_L^\rho(\lambda_i)$ -random variables so that the series $\sum_{i=1}^\infty \alpha_i X_i$ is an $\text{IED}_L^\rho(\Lambda)$ -random variable with

$$\Lambda = \left(\sum_{i=1}^\infty \alpha_i^{\rho/(1+\rho)} \lambda_i^{1/(1+\rho)} \right)^{1+\rho}.$$

Consider an ARMA series of the form

$$X_n = \sum_{i=1}^p \phi_i X_{n-i} + B_n + \sum_{j=1}^q \theta_j B_{n-j}$$

with positive coefficients ϕ_i and θ_j and initial value 0. Assume that (B_i) are i.i.d. $\text{IED}_1^\rho(\lambda)$ -random variables. We will give conditions (see Theorem 6.3) so that X_n converge to an $\text{IED}_1^\rho(\Lambda)$ -random variable and

$$\liminf_{n \rightarrow \infty} (\log n)^{1/\rho} X_n = \Lambda^{1/\rho} > 0, \quad \text{a.s.,}$$

where Λ is an explicit function of λ , ρ and the coefficients of the recursion.

We will also study the stochastic fixed point equation $X \stackrel{d}{=} AX + B$ where the vector (A, B) is independent of X , B is an $\text{IED}_L^\rho(\Lambda)$ -random variable, and A and B are nonnegative and positively quadrant dependent (see Theorem 7.6). If A and B are not positively quadrant dependent, we will prove by example that Theorem 7.6) (ii) need not be satisfied (see Section 8).

1.2. Organization of the paper

In Section 3, we introduce IED random variables and we prove that this class is closed under addition of finitely many independent summands. Section 4 is devoted to infinite series of independent IED random variables – we show that the sum may or may not be IED. In Section 5, we discuss the autoregressive equation, that is, the fixed point equation with the multiplicative coefficient that is a constant. In Section 6, we expand our results to ARMA models with positive coefficients and the noise from the IED class. In Section 7, we give estimates for left tails of solutions to the fixed point equation when the coefficients are positively quadrant dependent random variables. In Section 8, we show that if the coefficients are not positively quadrant dependent then these results no longer hold and the analysis is more demanding.

2. Preliminaries

We will write $a^+ = \max(0, a)$ for any real a .

We will use the convention that for any sequence (d_n) and $i > j$, $\sum_{n=i}^j d_n = 0$ and $\prod_{n=i}^j d_n = 1$.

Recall that the essential infimum of a random variable A is defined as follows,

$$\text{ess inf}(A) = \sup\{x \in \mathbb{R} : \mathbb{P}(A < x) = 0\}. \quad (2.1)$$

If $\lim_{x \rightarrow 0^+} f(x)/g(x) = 1$, then we will write $f(x) \sim g(x)$. The same notation will be used if the limit holds when $x \rightarrow \infty$.

Definition 2.1 (See [2], (1.2.1), Sections 1.4.1–1.4.2). A function $f : (0, \infty) \rightarrow (0, \infty)$ is called slowly varying at 0 if for all $a > 0$ we have $\lim_{x \rightarrow 0^+} f(ax)/f(x) = 1$. A function $f : (0, \infty) \rightarrow (0, \infty)$ is called regularly varying of index ρ at 0 if for all $a > 0$ we have $\lim_{x \rightarrow 0^+} f(ax)/f(x) = a^\rho$. A function f is called regularly varying of index ρ at infinity if $x \mapsto 1/f(1/x)$ is a regularly varying function of index ρ at 0.

Lemma 2.2 (See [2], (1.2.1), Thm. 1.4.1). A positive function f is regularly varying of index ρ at 0 if and only if $f(x) = x^\rho L(x)$ for some slowly varying function L at 0.

Definition 2.3 (See [2], (1.5.10)). If $f(x)$ is defined and locally bounded on some interval $(0, a]$ and $\lim_{x \rightarrow 0^+} f(x) = \infty$, then the generalized inverse of f is defined by

$$f^{\leftarrow}(y) = \sup\{x > 0 : f(x) > y\}.$$

If $f(x)$ is strictly positive on some interval $(0, a]$ and $\lim_{x \rightarrow 0^+} f(x) = 0$, then the generalized inverse of f is defined by

$$f^{\leftarrow}(y) = \inf\{x > 0 : f(x) > y\}.$$

Lemma 2.4. *If $\alpha > 0$ and f is α -regularly varying at 0 then there exists a function g which is $1/\alpha$ -regularly varying 0 and such that*

$$f(g(x)) \sim g(f(x)) \sim x \quad (2.2)$$

as $x \rightarrow 0^+$. The function g , called an asymptotic inverse of f , is determined up to asymptotic equivalence and one version of g is f^{\leftarrow} .

Proof. The proof is routine so it is left to the reader. See [2], Section 1.5.7, in particular Theorem 1.5.12. \square

3. Inverse exponential decay

The definition of random variables with inverse exponential decay of the left tail is inspired, in part, by inverse gamma distributions. These are used in Bayesian statistics (see [8]). One way to define inverse gamma distributions is by saying that the reciprocal of a random variable with a gamma distribution has the inverse gamma distribution. A more direct definition follows.

Definition 3.1. For a positive random variable X we say it has the inverse gamma distribution with parameters $\alpha, \beta > 0$ if its density function has the form

$$f(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{-\alpha-1} e^{-\beta/x},$$

for $x \in (0, \infty)$.

Recall Definition 1.1. The notation L, L_1 , etc., will be used exclusively for slowly varying functions at 0 unless stated otherwise.

Lemma 3.2. *Suppose that $\rho_1 < \rho_2$. If X is an $\text{IED}_{L_1}^{\rho_1}(\lambda)$ -random variable then X is $\text{IED}_{L_2}^{\rho_2}(0)$ -random variable for every slowly varying function L_2 at 0.*

Proof. The proof is routine and left to the reader. \square

Lemma 3.3. *For any non-negative random variable X ,*

$$\lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X < x) = \lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X \leq x), \quad (3.1)$$

in the sense that if one of the limits exists then the other one exists as well and they are equal.

Proof. For all $x > 0$ and $\varepsilon \in (0, 1)$,

$$\mathbb{P}(X \leq x(1 - \varepsilon)) \leq \mathbb{P}(X < x) \leq \mathbb{P}(X \leq x) \leq \mathbb{P}(X < x(1 + \varepsilon)). \quad (3.2)$$

Therefore

$$\limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X < x) \leq \liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X \leq x). \quad (3.3)$$

Assume that the limit $\lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X < x) = -\lambda$ exists. In view of the last inequality in (3.2), for any $\varepsilon > 0$,

$$\begin{aligned} & \limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X \leq x) \\ & \leq \limsup_{x \rightarrow 0^+} \frac{L(x)}{L(x(1+\varepsilon))} \cdot \frac{1}{(1+\varepsilon)^\rho} \cdot ([x(1+\varepsilon)]^\rho L(x(1+\varepsilon)) \log \mathbb{P}(X < x(1+\varepsilon))) \\ & = \limsup_{x \rightarrow 0^+} \frac{L(x)}{L(x(1+\varepsilon))} \cdot \frac{1}{(1+\varepsilon)^\rho} \cdot (-\lambda) = -\frac{\lambda}{(1+\varepsilon)^\rho}. \end{aligned}$$

Letting $\varepsilon \rightarrow 0^+$ and combining the resulting inequality with (3.3) yields (3.1).

In the case when $\lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X \leq x)$ exists, a similar argument, based on (3.3) and the first inequality in (3.2) proves (3.1). \square

Example 3.4. We will show that the positive limit in (1.3) might not exist for any fixed ρ and L . It is easy to see that there exists a c.d.f. F with the property that

$$F(2^{-(3k+1)}) = e^{-2^{3k+1}} \quad \text{and} \quad F(2^{-(3k+2)}) = e^{-2^{3k+3}} \quad (3.4)$$

for $k = 1, 2, \dots$, because F restricted to the arguments listed in (3.4) is increasing. If X is a random variable with c.d.f. F , then

$$\lim_{k \rightarrow \infty} 2^{-(3k+1)} \log \mathbb{P}(X \leq 2^{-(3k+1)}) = -1, \quad (3.5)$$

$$\lim_{k \rightarrow \infty} 2^{-(3k+2)} \log \mathbb{P}(X \leq 2^{-(3k+2)}) = -2. \quad (3.6)$$

Assume that there exist ρ , a function L slowly varying at 0, and $\lambda > 0$ such that $\log P(X < x) \sim -\lambda(x^\rho L(x))^{-1}$ as $x \rightarrow 0^+$. Then (3.5) shows that $\rho = 1$. Lemma 3.3 and (3.5)–(3.6) imply that

$$\begin{aligned} \frac{1}{2} &= \lim_{k \rightarrow \infty} \frac{2^{-(3k+1)} \log \mathbb{P}(X \leq 2^{-(3k+1)})}{2^{-(3k+2)} \log \mathbb{P}(X \leq 2^{-(3k+2)})} = \frac{1}{2} \lim_{k \rightarrow \infty} \frac{\log \mathbb{P}(X \leq 2 \cdot 2^{-(3k+2)})}{\log \mathbb{P}(X \leq 2^{-(3k+2)})} \\ &= \frac{1}{2} \lim_{k \rightarrow \infty} \frac{2^{-(3k+2)} L(2^{-(3k+2)})}{2 \cdot 2^{-(3k+2)} L(2 \cdot 2^{-(3k+2)})} = \frac{1}{4}. \end{aligned}$$

This contradiction proves our claim.

Lemma 3.5. If X has inverse gamma distribution with parameters $\alpha, \beta > 0$ (see Definition 3.1), then X is an $\text{IED}_1^1(\beta)$ -random variable.

Proof. Consider any $\varepsilon \in (0, 1)$. If X is a random variable with the inverse gamma distribution with parameters $\alpha, \beta > 0$ then for any $\varepsilon \in (0, 1)$ and sufficiently small $x > 0$,

$$\begin{aligned}\mathbb{P}(X < x) &\geq \int_{(1-\varepsilon)x}^x \frac{\beta^\alpha}{\Gamma(\alpha)} e^{-\beta/t} t^{-\alpha-1} dt \\ &\geq \frac{\beta^\alpha}{\Gamma(\alpha)} \exp\left(-\frac{\beta}{(1-\varepsilon)x}\right) ((1-\varepsilon)x)^{-\alpha-1} \cdot \varepsilon x.\end{aligned}$$

This implies that $\liminf_{x \rightarrow 0^+} x \log \mathbb{P}(X < x) \geq -\beta/(1-\varepsilon)$, for all $\varepsilon \in (0, 1)$. Hence, $\liminf_{x \rightarrow 0^+} x \log \mathbb{P}(X < x) \geq -\beta$.

On the other hand,

$$\begin{aligned}\mathbb{P}(X < x) &= \int_0^x \frac{\beta^\alpha}{\Gamma(\alpha)} e^{-\beta/t} t^{-\alpha-1} dt = \int_0^x \frac{\beta^\alpha}{\Gamma(\alpha)} e^{-\beta(1-\varepsilon)/t} e^{-\beta\varepsilon/t} t^{-\alpha-1} dt \\ &\leq e^{-\beta(1-\varepsilon)/x} \int_0^x \frac{\beta^\alpha}{\Gamma(\alpha)} e^{-\beta\varepsilon/t} t^{-\alpha-1} dt \leq e^{-\beta(1-\varepsilon)/x} \int_0^\infty \frac{\beta^\alpha}{\Gamma(\alpha)} e^{-\beta\varepsilon/t} t^{-\alpha-1} dt.\end{aligned}$$

Since the last integral is finite and independent of x , we have

$$\limsup_{x \rightarrow 0^+} x \log \mathbb{P}(X < x) \leq -\beta(1-\varepsilon)$$

for all $\varepsilon \in (0, 1)$. Hence, $\limsup_{x \rightarrow 0^+} x \log \mathbb{P}(X < x) \leq -\beta$. \square

The following two propositions are elementary so their proofs are left to the reader.

Proposition 3.6. Suppose that X is an $\text{IED}_L^\rho(\lambda)$ random variable, $\gamma > 0$ and $L_1(x) \equiv L(x^{1/\gamma})$. Then X^γ is an $\text{IED}_{L_1}^{\rho/\gamma}(\lambda)$ random variable.

In particular, if X is an $\text{IED}_1^\rho(\lambda)$ random variable and $\gamma > 0$, then X^γ is an $\text{IED}_1^{\rho/\gamma}(\lambda)$ random variable.

Proposition 3.7. If X is an $\text{IED}_L^\rho(\lambda)$ random variable and $\alpha > 0$, then αX is an $\text{IED}_L^\rho(\alpha^\rho \lambda)$ random variable.

Example 3.8. Note that if X is a nonnegative random variable with the property

$$\lim_{x \rightarrow \infty} \frac{\log \mathbb{P}(X > x)}{x^\rho} = -\lambda \leq 0,$$

then X^{-1} is an $\text{IED}_1^\rho(\lambda)$ -random variable. We give two natural examples.

- (a) If X has the exponential distribution with parameter λ (i.e., its mean is $1/\lambda$), then X^{-1} is an $\text{IED}_1^1(\lambda)$ -random variable.
- (b) If X has the normal distribution with mean μ and variance σ^2 then X^{-2} is an $\text{IED}_1^1((2\sigma^2)^{-1})$ -random variable. It follows from Proposition 3.6 that $|X|^{-1} = (X^{-2})^{\frac{1}{2}}$ is an $\text{IED}_1^2((2\sigma^2)^{-1})$ -random variable.

The following is an alternative characterization of IED-random variables that we will use often.

Lemma 3.9. *A positive random variable X is $\text{IED}_L^\rho(\lambda)$ if and only if for every $\delta > 0$ there exist $x_0 > 0$, $c_\delta > 0$ and $C_\delta > 0$ such that for all $x \in (0, x_0)$,*

$$c_\delta \exp\left(-\frac{\lambda + \delta}{x^\rho L(x)}\right) \leq \mathbb{P}(X < x) \leq \mathbb{P}(X \leq x) \leq C_\delta \exp\left(-\frac{\lambda(1 - \delta)}{x^\rho L(x)}\right). \quad (3.7)$$

Proof. The claim follows easily from the definition (1.3). \square

The following theorem, the springboard for the rest of the paper, says that IED random variables form a closed class under natural operations.

Theorem 3.10. *Suppose that X_1 and X_2 are independent random variables and X_k is $\text{IED}_L^\rho(\lambda_k)$, for $k = 1, 2$. Then $X_1 + X_2$ is an $\text{IED}_L^\rho([\lambda_1^{1/(1+\rho)} + \lambda_2^{1/(1+\rho)}]^{1+\rho})$ random variable.*

Before proving the theorem we recall without proof de Bruijn's Tauberian theorem (see [2], Thm. 4.12.9). See Definition 2.3 for the generalized inverse f^\leftarrow .

Theorem 3.11. *Suppose that μ is a measure on $(0, \infty)$ with the finite Laplace transform*

$$M(z) := \int_0^\infty e^{-zx} \mu(dx) < \infty, \quad \text{for all } z > 0.$$

Suppose that $\alpha < 0$ and $\phi(x)$ is a regularly varying function with index α at 0. Then,

$$\lim_{x \rightarrow 0^+} \phi^\leftarrow(1/x) \log \mu((0, x]) = -\lambda \quad (3.8)$$

if and only if

$$-\log M(z) \sim (1 - \alpha)(-\lambda/\alpha)^{\alpha/(\alpha-1)} / \psi^\leftarrow(z), \quad \text{as } z \rightarrow \infty, \quad (3.9)$$

where $\psi(z) = \phi(z)/z$.

Proof of Theorem 3.10. Let \mathbb{P}_{X_j} denote the distribution of X_j and

$$M_j(z) = \int_0^\infty e^{-zx} \mathbb{P}_{X_j}(dx), \quad j = 1, 2.$$

It is clear that for $M_j(z) \leq 1 < \infty$ for all $z > 0$ and $j = 1, 2$.

If L_2 is slowly varying at 0 then $\phi(x) := (1/x)^{1/\rho} L_2(1/x)$ is regularly varying with index $-1/\rho$ at infinity. Arguments analogous to those in [2], Section 1.5.7 (see also Lemma 2.4) show

that we can choose L_2 so that $\phi^{\leftarrow}(1/x) \sim x^\rho L(x)$ when $x \rightarrow 0^+$. With this choice of ϕ , the assumption that X_j is $\text{IED}_L^\rho(\lambda_j)$ matches (3.8), so (3.9) holds, that is, for $j = 1, 2$,

$$-\log M_j(z) \sim (1 + 1/\rho)(\lambda_j \rho)^{1/(1+\rho)} / \psi^{\leftarrow}(z), \quad \text{as } z \rightarrow \infty,$$

where $\psi(z) = \phi(z)/z$. Since X_1 and X_2 are independent,

$$\begin{aligned} -\log M_{X_1+X_2}(z) &= -\log M_{X_1}(z) - \log M_{X_2}(z) \\ &\sim (1 + 1/\rho)\rho^{1/(1+\rho)} [(\lambda_1^{1/(1+\rho)} + \lambda_2^{1/(1+\rho)})^{1+\rho}]^{1/(1+\rho)} / \psi^{\leftarrow}(z), \end{aligned}$$

as $z \rightarrow \infty$. The proof is completed by reversing our argument, using Theorem 3.11 and applying it to $X_1 + X_2$. \square

Proposition 3.12. *Suppose that $\rho_1 < \rho_2 < \dots < \rho_n$ and for $i = 1, \dots, n$ and $j = 1, \dots, m_i$, X_{ij} is $\text{IED}_{L_i}^{\rho_i}(\lambda_{ij})$, and all these random variables are independent. Then*

$$S := \sum_{i=1}^n \sum_{j=1}^{m_i} \alpha_{ij} X_{ij}$$

is an $\text{IED}_{L_n}^{\rho_n}(\Lambda)$ -random variable, where

$$\Lambda = \left(\sum_{j=1}^{m_n} \alpha_{nj}^{\rho_n/(1+\rho_n)} \lambda_{nj}^{1/(1+\rho_n)} \right)^{1+\rho_n}.$$

Proof. Lemma 3.2 and Proposition 3.7 imply that

$$\lim_{x \rightarrow 0^+} x^{\rho_n} L_n(x) \log \mathbb{P}(\alpha_{ij} X_{ij} < x) = \begin{cases} 0, & i \neq n; \\ -\alpha_{ij}^{\rho_n} \lambda_{ij}, & i = n. \end{cases}$$

The proposition follows from Theorem 3.10 and induction. \square

4. Convergence of infinite IED-series

It is a natural question whether Proposition 3.12 holds for infinite series with independent IED summands. The short answer is “no” but “yes” under extra assumptions.

Proposition 4.1. *Suppose that $(X_i)_{i \geq 1}$ are independent and X_i is $\text{IED}_L^\rho(\lambda_i)$, for $i \geq 1$. Let $S = \sum_{i \geq 1} X_i$ and $\Lambda = (\sum_{i \geq 1} \lambda_i^{1/(1+\rho)})^{1+\rho}$. Then*

$$\limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(S < x) \leq -\Lambda. \quad (4.1)$$

Proof. Since X_i 's are nonnegative random variables, for every n we have $\mathbb{P}(S < x) \leq \mathbb{P}(\sum_{i=1}^n X_i < x)$. This and Proposition 3.12 imply

$$\begin{aligned} \limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(S < x) &\leq \limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}\left(\sum_{i=1}^n X_i < x\right) \\ &= -\left(\sum_{i=1}^n \lambda_i^{1/(1+\rho)}\right)^{1+\rho}. \end{aligned}$$

If we let $n \rightarrow \infty$, the claim follows. \square

Example 4.2. Recall the notation and assumptions from Proposition 4.1. Two examples given below show that, in general, (4.1) cannot be strengthened to equality. The first example is a little bit more elegant than the second one. But $S \equiv \infty$ in the first example, suggesting that divergence of the sum $\sum_{i \geq 1} X_i$ is the only possible obstacle to having equality in (4.1). For this reason we present another example with $S \leq 1$, a.s.

(a) Suppose that X_i 's have inverse gamma distributions with parameters $\alpha_i = 1/4$ and $\beta_i = 1/i^4$, for $i \geq 1$ (see Definition 3.1). According to Lemma 3.5, X_i is an $\text{IED}_1^1(\beta_i)$ -random variable, for every $i \geq 1$. Let $\rho = 1$ and $\lambda_i = \beta_i = 1/i^4$ for all i . Then, for $i \geq 1$,

$$\mathbb{P}(X_i > 1) = \int_1^\infty \frac{1}{i\Gamma(1/4)} x^{-1/4-1} e^{-1/(i^4 x)} dx \geq \frac{1}{i} \int_1^\infty \frac{1}{\Gamma(1/4)} x^{-1/4-1} e^{-1/x} dx.$$

Hence,

$$\sum_{i \geq 1} \mathbb{P}(X_i > 1) = \infty.$$

Therefore, by the Borel–Cantelli lemma, $X_i > 1$ for infinitely many i 's. It follows that $S = \infty$, a.s., regardless of the fact that

$$\Lambda := \left(\sum_{i \geq 1} \lambda_i^{1/(1+1)}\right)^{1+1} = \left(\sum_{i=1}^\infty \frac{1}{i^2}\right)^2 < \infty.$$

(b) For some $\tau_i > 0$, $i \geq 1$, to be specified later, we give X_i 's the following cumulative distribution functions,

$$F_i(x) = \begin{cases} 0 & \text{for } x \leq 0; \\ e^{1-\tau_i} e^{-1/(2^i x)} & \text{for } 0 < x < 2^{-i}; \\ 1 & \text{for } 2^{-i} \leq x. \end{cases}$$

Note that each X_i is an $\text{IED}_1^1(2^{-i})$ -random variable. Let $\lambda_i = 2^{-i}$ and note that $S := \sum_{i=1}^\infty X_i \leq 1$, a.s., because $X_i \leq 2^{-i}$, a.s., for all i . Although $\sqrt{\Lambda} = \sum_{i=1}^\infty \sqrt{\lambda_i} < \infty$, we will show that

S is not an $\text{IED}_1^1(\Lambda)$ -random variable. Consider an integer $n \geq 1$ and let $x = 2^{-(n+1/2)}$. Hence $2^{-(n+1)} < x < 2^{-n}$. Since each X_i is bounded by 2^{-i} , we have

$$\{S < x\} \subset \bigcap_{i=1}^n \{X_i < 2^{-i}\}.$$

Hence, using the fact that $\mathbb{P}(X_i < 2^{-i}) = e^{-\tau_i}$, we have

$$x \log \mathbb{P}(S < x) \leq x \sum_{i=1}^n \log \mathbb{P}(X_i < 2^{-i}) = -x \sum_{i=1}^n \tau_i. \quad (4.2)$$

If we choose $\tau_i = c2^{i-1/2}$, then

$$x \sum_{i=1}^n \tau_i = x \sum_{i=1}^n c2^{i-1/2} = 2^{-(n+1/2)} c2^{-1/2} (2^{n+1} - 1) = c(1 - 2^{-n-1}).$$

This and (4.2) imply that $\liminf_{x \rightarrow 0^+} x \log \mathbb{P}(S < x) \leq -c$. If we set $-c < -\Lambda$, then $\liminf_{x \rightarrow 0^+} x \log \mathbb{P}(S < x) < -\Lambda$.

We will give sufficient conditions for the equality in (4.1) in Theorems 4.4 and 4.7. The main technical part of the proof is contained in the following lemma.

Lemma 4.3. *Suppose that $(X_i)_{i \geq 1}$ is a sequence of independent random variables such that*

$$\lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X_i < x) = -\lambda_i,$$

for a sequence $(\lambda_i)_{i \geq 1}$ of non-negative real numbers and a slowly varying function L at 0. Assume that

$$\Lambda^{1/(1+\rho)} := \sum_{i \geq 1} \lambda_i^{1/(1+\rho)} < \infty.$$

Suppose that a random variable B satisfies the following conditions.

(i) *B has the property*

$$\lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(B < x) = -1.$$

(ii) *B is stochastically greater than $X_i/\lambda_i^{1/\rho}$ for each $i \geq 1$, that is, for all $x \in \mathbb{R}$ and $i \geq 1$,*

$$\mathbb{P}(B \leq x) \leq \mathbb{P}\left(\frac{X_i}{\lambda_i^{1/\rho}} \leq x\right).$$

(iii) There exist positive real numbers $(\gamma_i)_{i \geq 1}$ such that $\sum_{i=1}^{\infty} \gamma_i = 1$, $\sum_{i=1}^{\infty} \lambda_i / \gamma_i^{\rho} < \infty$, and

$$\lim_{x \rightarrow 0^+} x^{\rho} L(x) \sum_{i=1}^{\infty} \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) = 0. \quad (4.3)$$

Then $\sum_{i=1}^{\infty} X_i$ converges to an a.s. finite random variable S satisfying

$$\lim_{x \rightarrow 0^+} x^{\rho} L(x) \log \mathbb{P}(S < x) = -\Lambda.$$

Proof. Since B is stochastically greater than $X_i / \lambda_i^{1/\rho}$, we have

$$\mathbb{P}(X_i \geq \gamma_i x) = \mathbb{P}(X_i / \lambda_i^{1/\rho} \geq \gamma_i x / \lambda_i^{1/\rho}) \leq \mathbb{P}(B \geq \gamma_i x / \lambda_i^{1/\rho}).$$

It follows from (4.3) that for small $x > 0$, we have

$$\sum_{i=1}^{\infty} \mathbb{P}(B \geq \gamma_i x / \lambda_i^{1/\rho}) < \infty. \quad (4.4)$$

Hence, for small $x > 0$, $\sum_{i=1}^{\infty} \mathbb{P}(X_i \geq \gamma_i x) < \infty$. By the Borel–Cantelli lemma, the sequence (X_i) is eventually dominated by $(\gamma_i x)$ a.s. Recall that $\sum_{i=1}^{\infty} \gamma_i = 1$ to see that $\sum_{n=1}^{\infty} X_n$ converges to an a.s. finite random variable S .

Let $S_n = \sum_{k=1}^n X_k$. We have

$$\{S < x\} \supset \left\{ S_n < x \sum_{k=1}^n \gamma_k \right\} \cap \bigcap_{k=n+1}^{\infty} \{X_k < \gamma_k x\}$$

so, by assumption (ii),

$$\begin{aligned} \mathbb{P}(S < x) &\geq \mathbb{P}\left(S_n < x \sum_{k=1}^n \gamma_k\right) \prod_{k=n+1}^{\infty} \mathbb{P}(X_k < \gamma_k x) \\ &\geq \mathbb{P}\left(S_n < x \sum_{k=1}^n \gamma_k\right) \prod_{k=n+1}^{\infty} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right). \end{aligned} \quad (4.5)$$

By assumption (i), for $\varepsilon > 0$ there exists $x_0 > 0$ such that

$$\mathbb{P}(B < x) \geq \exp(-2(x^{\rho} L(x))^{-1}),$$

for $0 < x < x_0$. Let

$$F_n(x_0) = \left\{ k \geq n+1 : x_0 > \frac{\gamma_k x}{\lambda_k^{1/\rho}} \right\}, \quad G_n(x_0) = \left\{ k \geq n+1 : x_0 \leq \frac{\gamma_k x}{\lambda_k^{1/\rho}} \right\}.$$

Using this notation, we can break the last product in (4.5) as follows:

$$\begin{aligned} \prod_{k=n+1}^{\infty} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) &= \prod_{k \in F_n(x_0)} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) \prod_{k \in G_n(x_0)} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) \\ &\geq \exp\left[-\frac{2}{x^\rho} \sum_{k \in F_n(x_0)} \frac{\lambda_k}{\gamma_k^\rho} L\left(\frac{\gamma_k x}{\lambda_k^{1/\rho}}\right)^{-1}\right] \\ &\quad \times \prod_{k \in G_n(x_0)} \left[1 - \mathbb{P}\left(B \geq \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right)\right]. \end{aligned} \quad (4.6)$$

By the definition of $G_n(x_0)$, for all $k \in G_n(x_0)$ we have

$$\mathbb{P}\left(B \geq \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) \leq \mathbb{P}(B \geq x_0) < 1.$$

Standard calculus arguments show that there exists $m > 1$ such that $-ma \leq \log(1 - a)$ for $0 < a \leq \mathbb{P}(B < x_0)$. Hence, by (4.6),

$$\begin{aligned} \prod_{k=n+1}^{\infty} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) &\geq \exp\left(-\frac{2}{x^\rho} \sum_{k \in F_n(x_0)} \frac{\lambda_k}{\gamma_k^\rho} L\left(\frac{\gamma_k x}{\lambda_k^{1/\rho}}\right)^{-1} - m \sum_{k \in G_n(x_0)} \mathbb{P}\left(B \geq \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right)\right) \\ &\geq \exp\left(-\frac{2}{x^\rho} \sum_{k=n+1}^{\infty} \frac{\lambda_k}{\gamma_k^\rho} L\left(\frac{\gamma_k x}{\lambda_k^{1/\rho}}\right)^{-1} - m \sum_{k=1}^{\infty} \mathbb{P}\left(B \geq \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right)\right). \end{aligned} \quad (4.7)$$

Applying Proposition 3.12 to the first factor on the last line of (4.5),

$$\begin{aligned} \liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}\left(S_n < x \sum_{k=1}^n \gamma_k\right) &= \liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}\left(\left(\sum_{k=1}^n \gamma_k\right)^{-1} S_n < x\right) \\ &\geq -\left(\sum_{i \geq 1} \lambda_i^{1/(1+\rho)}\right)^{1+\rho} \left(\sum_{k=1}^n \gamma_k\right)^{-\rho}. \end{aligned} \quad (4.8)$$

The estimate (4.7) and the assumption (4.3) yield for the second factor in (4.5),

$$\begin{aligned} \liminf_{x \rightarrow 0^+} x^\rho L(x) \log \left(\prod_{k=n+1}^{\infty} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) \right) \\ \geq \liminf_{x \rightarrow 0^+} x^\rho L(x) \left(-\frac{2}{x^\rho} \sum_{k=n+1}^{\infty} \frac{\lambda_k}{\gamma_k^\rho} L\left(\frac{\gamma_k x}{\lambda_k^{1/\rho}}\right)^{-1} - m \sum_{k=1}^{\infty} \mathbb{P}\left(B \geq \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) \right). \end{aligned} \quad (4.9)$$

Set $f(z) = z^\rho L(z)$. By [2], Thm. 1.5.2,

$$\lim_{z \rightarrow 0} f(z)/f(z/h) = h^\rho, \quad (4.10)$$

uniformly in h , on each fixed interval of the form $[b, \infty)$. Since $\gamma_i/\lambda_i^{1/\rho} \rightarrow \infty$ as $i \rightarrow \infty$, there is $b > 0$ such that all the values of this sequence are in $[b, \infty)$. We apply (4.10) to see that for some $C_1 < \infty$ and $x_1 > 0$, for all $x \in (0, x_1)$ and $i \geq 1$,

$$\frac{\lambda_i}{\gamma_i^\rho} L(x) L\left(\frac{\gamma_i x}{\lambda_i^{1/\rho}}\right)^{-1} = \frac{f(x)}{f(\gamma_i x/\lambda_i^{1/\rho})} \leq C_1 \frac{\lambda_i}{\gamma_i^\rho}. \quad (4.11)$$

This, (4.4) and (4.9) imply that

$$\liminf_{x \rightarrow 0^+} x^\rho L(x) \log \left(\prod_{k=n+1}^{\infty} \mathbb{P}\left(B < \frac{\gamma_k}{\lambda_k^{1/\rho}} x\right) \right) \geq -2C_1 \sum_{i=n+1}^{\infty} \frac{\lambda_i}{\gamma_i^\rho}. \quad (4.12)$$

Combining (4.5), (4.8) and (4.12) gives

$$\liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(S < x) \geq -\left(\sum_{i \geq 1} \lambda_i^{1/(1+\rho)}\right)^{1+\rho} \left(\sum_{k=1}^n \gamma_k\right)^{-\rho} - 2C_1 \sum_{i=n+1}^{\infty} \frac{\lambda_i}{\gamma_i^\rho}. \quad (4.13)$$

Recall that we have assumed that $\Lambda^{1/(1+\rho)} = \sum_{i \geq 1} \lambda_i^{1/(1+\rho)} < \infty$, $\sum_{i=1}^{\infty} \gamma_i = 1$, and $\sum_{i=1}^{\infty} \lambda_i/\gamma_i^\rho < \infty$. Thus, when we let $n \rightarrow \infty$ in (4.13), we obtain

$$\liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(S < x) \geq -\Lambda.$$

The opposite inequality follows from Proposition 4.1. □

Theorem 4.4. Suppose that $(B_i)_{i \geq 1}$ is a sequence of i.i.d. $\text{IED}_L^\rho(1)$ -random variables satisfying

$$\limsup_{x \rightarrow \infty} x^\rho \mathbb{P}(B_1 \geq x) < \infty, \quad (4.14)$$

and $(\alpha_i)_{i \geq 1}$ is a sequence of strictly positive real numbers satisfying $\sum_{i=1}^{\infty} \alpha_i^{\rho/(1+\rho)} < \infty$. Then the series $\sum_{i=1}^{\infty} \alpha_i B_i$ converges a.s. to an $\text{IED}_L^\rho(\Lambda)$ random variable, where

$$\Lambda = \left(\sum_{i=1}^{\infty} \alpha_i^{\rho/(1+\rho)} \right)^{1+\rho}.$$

Remark 4.5. Condition (4.14) implies that $\mathbb{E}[B_1^\tau] < \infty$ for $\tau \in [0, \rho)$. If $\mathbb{E}[B_1^\rho] < \infty$ then (4.14) is satisfied.

Proof of Theorem 4.4. We will apply Lemma 4.3. Let $X_i = \alpha_i B_i$ and $\lambda_i = \alpha_i^\rho$, for $i \geq 1$. By Proposition 3.7, B_i is $\text{IED}_L^\rho(\alpha_i^\rho)$ -random variable, for $i \geq 1$. Let B be distributed as B_1 . It is easy to see that assumptions (i) and (ii) of Lemma 4.3 are satisfied by X_i 's and B .

Let $c = (\sum_{i=1}^\infty \alpha_i^{\rho/(1+\rho)})^{-1}$ and $\gamma_i = c\alpha_i^{\rho/(1+\rho)}$ for $i \geq 1$. Then $\sum_{i=1}^\infty \gamma_i = 1$ and

$$\sum_{i=1}^\infty \frac{\lambda_i}{\gamma_i^\rho} = \sum_{i=1}^\infty \frac{\alpha_i^\rho}{c^\rho \alpha_i^{\rho^2/(1+\rho)}} = \frac{1}{c^\rho} \sum_{i=1}^\infty \alpha_i^{\rho/(1+\rho)} = c^{-1-\rho} < \infty, \quad (4.15)$$

so two conditions listed in assumption (iii) of Lemma 4.3 are satisfied. It remains to verify (4.3).

Without loss of generality we can assume that $\lim_{x \rightarrow \infty} L(x) = 1$. Then (4.14) is equivalent to $\limsup_{x \rightarrow \infty} x^\rho L(x) \mathbb{P}(B \geq x) < \infty$. We have $\lim_{x \rightarrow 0^+} x^\rho L(x) \mathbb{P}(B \geq x) = 0$ because $\rho > 0$ and L is slowly varying at 0. The two conditions imply that there exists $C > 0$ such that $\mathbb{P}(B \geq x) \leq Cx^{-\rho} L(x)^{-1}$ for all $x > 0$. In particular, we have for all $x > 0$,

$$\mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) \leq Cx^{-\rho} \frac{\lambda_i}{\gamma_i^\rho} L\left(\frac{\gamma_i x}{\lambda_i^{1/\rho}}\right)^{-1}. \quad (4.16)$$

For every fixed i , $\lim_{x \rightarrow 0^+} x^\rho L(x) \mathbb{P}(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x) = 0$, so for every fixed n ,

$$\limsup_{x \rightarrow 0^+} x^\rho L(x) \sum_{i=1}^\infty \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) = \limsup_{x \rightarrow 0^+} x^\rho L(x) \sum_{i=n}^\infty \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right). \quad (4.17)$$

By (4.16),

$$x^\rho L(x) \sum_{i=n}^\infty \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) \leq C \sum_{i=n}^\infty \frac{\lambda_i}{\gamma_i^\rho} L(x) L\left(\frac{\gamma_i x}{\lambda_i^{1/\rho}}\right)^{-1}. \quad (4.18)$$

This, (4.11), (4.17), (4.18) and (4.15) imply that

$$\limsup_{x \rightarrow 0^+} x^\rho L(x) \sum_{i=1}^\infty \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) \leq CC_1 \sum_{i=n}^\infty \frac{\lambda_i}{\gamma_i^\rho} < \infty.$$

Letting $n \rightarrow \infty$, we obtain

$$\lim_{x \rightarrow 0^+} x^\rho L(x) \sum_{i=n}^\infty \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) = 0.$$

We see that (4.3) holds and, therefore, the theorem follows from Lemma 4.3. \square

Example 4.6. Suppose that $(B_i)_{i \geq 1}$ are i.i.d. $\text{IED}_1^\rho(1)$ -random variables with finite ρ -th moment. Theorem 4.4 and Remark 4.5 imply that for all $c, \varepsilon > 0$, the series $\sum_{i=1}^\infty ci^{-(1+\varepsilon)(1+\rho)/\rho} B_i$ converges a.s. The limit is an $\text{IED}_1^\rho(\Lambda)$ -random variable with parameter $\Lambda = c^\rho (\sum_{i=1}^\infty i^{-1-\varepsilon})^{1+\rho}$.

The following theorem shows that if the parameters α_i decrease at a geometric rate then we can weaken the condition on the moments of B_i and obtain the same conclusion as in Theorem 4.4.

Theorem 4.7. *Suppose that $(B_i)_{i \geq 1}$ is a sequence of i.i.d. $\text{IED}_L^\rho(1)$ -random variables satisfying $\mathbb{E}[\log^+ B_i] < \infty$. For any sequence of strictly positive real numbers $(\alpha_i)_{i \geq 1}$ with the property $\limsup_{i \rightarrow \infty} \sqrt[i]{\alpha_i} = \kappa \in (0, 1)$, the series $\sum_{i=1}^{\infty} \alpha_i B_i$ converges a.s., and the limit is an $\text{IED}_L^\rho(\Lambda)$ -random variable with*

$$\Lambda = \left(\sum_{i=1}^{\infty} \alpha_i^{\rho/(1+\rho)} \right)^{1+\rho}.$$

Proof. We will apply Lemma 4.3. Let $X_i = \alpha_i B_i$ and $\lambda_i = \alpha_i^\rho$ for $i \geq 1$, and let B be a random variable with the same distribution as B_1 . It is easy to see that assumptions (i) and (ii) of Lemma 4.3 are satisfied. It only remains to show that condition (iii) is satisfied.

Pick $\zeta \in (\kappa, 1)$, and set $\gamma_i = \zeta^{i-1}(1 - \zeta)$. Note that $\sum_{i=1}^{\infty} \gamma_i = 1$. It is clear that there exist $c > 0$ and κ_1 such that $0 < \kappa < \kappa_1 < \zeta$ and $0 \leq \lambda_i^{1/\rho} = \alpha_i \leq c\kappa_1^i$ for all $i \geq 1$. Hence,

$$\sum_{i=1}^{\infty} \frac{\lambda_i}{\gamma_i^\rho} \leq c_1 \frac{\zeta^\rho}{(1 - \zeta)^\rho} \sum_{i=1}^{\infty} (\kappa_1/\zeta)^{i\rho} < \infty,$$

so two conditions listed in assumption (iii) of Lemma 4.3 are satisfied. It remains to verify (4.3).

We will use the following well known inequality, saying that for any positive random variable X we have

$$\sum_{k=1}^{\infty} \mathbb{P}(X \geq k) \leq \mathbb{E}X + 1. \quad (4.19)$$

The above inequality is used to justify the second inequality below,

$$\begin{aligned} \sum_{i=1}^{\infty} \mathbb{P}\left(B \geq \frac{\gamma_i}{\lambda_i^{1/\rho}} x\right) &\leq \sum_{i=1}^{\infty} \mathbb{P}\left(B \geq \frac{1 - \zeta}{c\zeta} \left(\frac{\zeta}{\kappa_1}\right)^i x\right) \\ &= \sum_{i=1}^{\infty} \mathbb{P}(\log B - \log x + \log(c\zeta(1 - \zeta)^{-1}) \geq i \log(\zeta/\kappa_1)) \\ &\leq \frac{\mathbb{E}(\log B - \log x + \log(c\zeta(1 - \zeta)^{-1}))^+}{\log(\zeta/\kappa_1)} + 1 \\ &\leq \frac{\mathbb{E}[\log^+ B] + |\log x| + |\log(c\zeta(1 - \zeta)^{-1})|}{\log(\zeta/\kappa_1)} + 1. \end{aligned}$$

The last estimate implies that $\lim_{x \rightarrow 0^+} x^\rho L(x) \sum_{i=1}^{\infty} \mathbb{P}(B \geq \gamma_i x / \lambda_i) = 0$ because we have assumed that $\mathbb{E}[\log^+ B_i] < \infty$. We conclude that assumption (iii) of Lemma 4.3 holds. \square

5. Autoregressive equation

We will consider solutions to the autoregressive equation, a simple ARMA model, in this section. More general ARMA models will be considered in subsequent sections.

We start by recalling a known result. We would like to point out that random variables A and B need not be independent for the following to hold.

Theorem 5.1. *If $\mathbb{E}[\log |A|] < 0$ and $\mathbb{E}[\log^+ |B|] < \infty$ then (1.1) has a unique solution. Suppose that $(A_i, B_i)_{i \geq 1}$ are i.i.d. two-dimensional vectors distributed as (A, B) .*

- (a) *The distribution of the solution to (1.1) is the stationary distribution for the Markov chain given by*

$$X_n = A_n X_{n-1} + B_n. \quad (5.1)$$

- (b) *The series*

$$S = \sum_{i=1}^{\infty} \left(\prod_{j=1}^{i-1} A_j \right) B_i \quad (5.2)$$

converges a.s. and the distribution of the limit is the same as that of the solution to (1.1).

Proof. By [4], Theorem 2.1.3, and [6], Theorem 2.1, the sequence (5.1) has a unique ergodic invariant stationary distribution. Moreover, (5.2) is a representation of that distribution. By [4], Lemma 2.2.7, this distribution is the unique solution to the fixed point equation (1.1). \square

In the rest of this section we will take a look at the nonnegative solution to the autoregressive equation

$$X \stackrel{d}{=} rX + B, \quad (5.3)$$

where $0 < r < 1$.

Corollary 5.2. *If $\mathbb{E}[\log^+ B] < \infty$ then (5.3) has a unique solution. Suppose that (B_i) are i.i.d. random variables distributed as B .*

- (a) *The distribution of the solution to (5.3) is the stationary distribution of the Markov chain given by*

$$X_n = rX_{n-1} + B_n. \quad (5.4)$$

- (b) *The series*

$$\sum_{i=1}^{\infty} r^{i-1} B_i \quad (5.5)$$

converges a.s. and the distribution of the limit is the same as that of the solution to (5.3).




Figure 1. The figure shows the the graph of $(\log(n)X_n)_{n \geq 1}$, where $(X_n)_{n \geq 1}$ is the ARMA process $X_{n+1} = \frac{1}{4}X_n + B_{n+1}$, where $B_n = \min(E_n^{-1}, 1)$, E_n 's are i.i.d. exponential with parameter $1/2$ and $X_0 = 0$. It follows from Proposition 5.4 that $\Lambda = 2$. This is visible in the graph as the black line segments occasionally reach to the horizontal line at level 2.

Proof. The corollary follows from Theorem 5.1. □

Corollary 5.3. *If B is an $\text{IED}_L^\rho(\lambda)$ -random variable such that $\mathbb{E}[\log^+ B] < \infty$, and X is the solution to (5.3) then X is an $\text{IED}_L^\rho(\Lambda)$ -random variable with*

$$\Lambda = \frac{\lambda}{(1 - r^{\rho/(1+\rho)})^{1+\rho}}. \quad (5.6)$$

Proof. The corollary follows from Theorem 4.7 and Corollary 5.2(b). □

The following result is a special case of Theorem 6.3 so we leave it without proof.

Proposition 5.4. *If B is $\text{IED}_1^\rho(\lambda)$, $\mathbb{E}[(\log^+ B)^s] < \infty$ for all $s > 0$, Λ is defined in (5.6), and X_n 's are defined in (5.4) then*

$$\liminf_{n \rightarrow \infty} (\log(n))^{1/\rho} X_n = \Lambda^{1/\rho}, \quad a.s.$$

This proposition (for case $\rho = 1$) is illustrated in Figure 1. Note that the result holds under mild assumptions on the right tail of B .

6. ARMA models with IED noise

This section is devoted to autoregressive moving average (ARMA) models more general than those in the previous section.

Definition 6.1. An ARMA(p, q) sequence has the form

$$X_n = \sum_{i=1}^p \phi_i X_{n-i} + B_n + \sum_{j=1}^q \theta_j B_{n-j}, \quad (6.1)$$

where $(\phi_i)_{i=1,\dots,p}$ and $(\theta_j)_{j=1,\dots,q}$ are positive constants and $(B_i)_{i \geq 1}$ are i.i.d.

Our analysis of ARMA models will be based on [3], Chapter 3. Using the notation from Definition 6.1, we define complex polynomials Φ and Θ by

$$\Phi(z) = 1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p \quad \text{and} \quad \Theta(z) = 1 + \theta_1 z + \dots + \theta_q z^q.$$

Theorem 6.2. Suppose that $(B_n)_{n \geq 1}$ are i.i.d. $\text{IED}_L^\rho(\lambda)$ random variables and we have $\mathbb{E}[\log^+ B_1] < \infty$. Assume that $\Phi(z) \neq 0$ for $|z| \leq 1$ and Φ and Θ have no common roots. Then $\Psi(z) := \Theta(z)/\Phi(z)$ is analytic on a neighborhood of the unit disc $\{|z| \leq 1\}$, and its Taylor series, i.e.,

$$\Psi(z) = \sum_{k=0}^{\infty} \psi_k z^k, \quad (6.2)$$

has positive coefficients.

(i) Random variables X_n in (6.1) can be represented as

$$X_n = \sum_{k=0}^n \psi_k B_{n-k}. \quad (6.3)$$

(ii) Each X_n is an $\text{IED}_L^\rho(\Lambda_n)$ -random variable with

$$\Lambda_n = \lambda \left(\sum_{k=0}^n \psi_k^{\rho/(1+\rho)} \right)^{1+\rho}.$$

(iii) When $n \rightarrow \infty$, $X_n \xrightarrow{d} X := \sum_{k=0}^{\infty} \psi_k B_k$, and the limit is a finite $\text{IED}_L^\rho(\Lambda)$ -random variable with

$$\Lambda = \lambda \left(\sum_{k=0}^{\infty} \psi_k^{\rho/(1+\rho)} \right)^{1+\rho}.$$

Proof. There exists $\varepsilon > 0$ such that $|\phi_1 z + \phi_2 z^2 + \dots + \phi_p z^p| < 1$ for $|z| < \varepsilon$. For such z ,

$$\Psi(z) = \sum_{j=0}^{\infty} (\phi_1 z + \phi_2 z^2 + \dots + \phi_p z^p)^j (1 + \theta_1 z + \dots + \theta_q z^q).$$

It is evident from this formula that for $|z| < \varepsilon$, $\Psi(z)$ can be represented as a series with positive coefficients. By the uniqueness of Taylor series, all ψ_k 's are positive.

The function Ψ is analytic on a disc around 0 whose radius is greater than 1. Hence, the Taylor series of Ψ around 0 has a convergence radius $R > 1$. By the Cauchy–Hadamard formula, $\limsup_{k \rightarrow \infty} |\psi_k|^{1/k} = R^{-1} < 1$. Therefore, there exist $C > 0$ and $0 < \beta < 1$ such that $\psi_k < C\beta^k$ for $k \geq 0$. This implies that both series $\sum_{k \geq 1} \psi_k$ and $\sum_{k \geq 1} \sqrt{\psi_k}$ converge.

Part (i) follows from [3], Thm. 3.1.1. Part (ii) follows from Proposition 3.12. Part (iii) follows from Proposition 3.12 and Theorem 4.7. \square

We will now prove a generalization of Proposition 5.4.

Theorem 6.3. Consider an ARMA sequence (6.1) satisfying the assumptions of Theorem 6.2 and recall the notation from (6.2). If $\mathbb{E}[(\log^+ B)^r] < \infty$ for all $r > 0$ then

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} = \Lambda^{1/\rho} := \lambda^{1/\rho} \left(\sum_{k=0}^{\infty} \psi_k^{\rho/(1+\rho)} \right)^{(1+\rho)/\rho}, \quad a.s.,$$

where g is the generalized inverse of the function $x \mapsto x^\rho L(x)$ at 0.

The proof will be preceded by a few lemmas.

Lemma 6.4. (i) For every $\varepsilon > 0$, the events

$$\left\{ X_n \leq g \left(\frac{\Lambda}{(1+\varepsilon)\log n} \right) \right\}$$

happen finitely often, a.s.

(ii) We have

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} \geq \Lambda^{1/\rho}, \quad a.s.$$

Proof. (i) For any $\varepsilon > 0$ there exists $\delta \in (0, 1)$ such that $\gamma := (1 - \delta)(1 + \varepsilon/2) > 1$. By Theorem 6.2, there exist C_δ, x_0 and n_0 such that

$$\mathbb{P}(X_{n_0} \leq x) \leq C_\delta \exp \left(-\frac{\Lambda(1-\delta)}{x^\rho L(x)} \right)$$

for all $x \in (0, x_0)$.

Random variables B_k are i.i.d., so (6.3) implies that

$$X_n = \sum_{k=0}^n \psi_k B_{n-k} \stackrel{d}{=} \sum_{k=0}^n \psi_k B_k.$$

Since ψ_k 's and B_k 's are nonnegative, it follows that X_{n+1} stochastically majorizes X_n for all n . Hence, $n \mapsto \mathbb{P}(X_n \leq x)$ is a non-increasing sequence and, therefore, for $n \geq n_0$ and $x \in (0, x_0)$,

$$\mathbb{P}(X_n \leq x) \leq C_\delta \exp\left(-\frac{\Lambda(1-\delta)}{x^\rho L(x)}\right).$$

It follows that for large n ,

$$\mathbb{P}\left(X_n \leq g\left(\frac{\Lambda}{(1+\varepsilon)\log n}\right)\right) \leq C_\delta e^{-(1+\varepsilon/2)(1-\delta)\log n} = C_\delta n^{-\gamma}.$$

Hence,

$$\sum_{n=1}^{\infty} \mathbb{P}\left(X_n \leq g\left(\frac{\Lambda}{(1+\varepsilon)\log n}\right)\right) < \infty,$$

and the claim follows by the Borel–Cantelli lemma.

(ii) It follows from part (i) that for every $\varepsilon > 0$,

$$\liminf_{n \rightarrow \infty} g\left(\frac{\Lambda}{(1+\varepsilon)\log n}\right)^{-1} X_n \geq 1, \quad \text{a.s.}$$

By Lemma 2.4, g is $1/\rho$ -regularly varying at 0. Hence,

$$g\left(\frac{\Lambda}{(1+\varepsilon)\log n}\right) \sim \left(\frac{\Lambda}{1+\varepsilon}\right)^{1/\rho} g(1/\log n).$$

Therefore,

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} \geq \Lambda^{1/\rho} / (1+\varepsilon)^{1/\rho}.$$

Part (ii) follows by letting $\varepsilon \rightarrow 0$. □

It will be convenient to use the following notation, reminiscent of (6.3),

$$X_n^m = \sum_{k=0}^{n-m-1} \psi_k B_{n-k}. \quad (6.4)$$

Recall that $\lfloor a \rfloor$ denotes the largest integer less than or equal to a .

Lemma 6.5. Fix $\varepsilon > 0$ and suppose that $1 < 1 + \delta < \sqrt{1 + \varepsilon}$ and $\alpha \in (1, 1 + \delta)$. Events

$$\left\{ X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right) \right\}$$

happen infinitely often a.s.

Proof. Note that the random variables $X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}$, $n \geq 2$, are jointly independent.

The random variable $S := \sum_{k=0}^{\infty} \psi_k B_k$ stochastically majorizes every $X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}$. By Theorem 6.2(iii) and Lemma 3.9, for large n ,

$$\begin{aligned} \mathbb{P}\left(X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) &\geq \mathbb{P}\left(S \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) \\ &\geq c_\delta \exp\left(-\frac{\Lambda(1+\delta)\log n^\alpha}{\Lambda(1+\varepsilon)}\right) = c_\delta n^{-\alpha(1+\delta)/(1+\varepsilon)} \geq n^{-\alpha/(1+\delta)}. \end{aligned}$$

It follows that

$$\sum_{n=2}^{\infty} \mathbb{P}\left(X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) = \infty,$$

and, therefore, the claim follows by the Borel–Cantelli lemma. \square

Lemma 6.6. *If $\alpha > 1$ then,*

$$\lim_{n \rightarrow \infty} \frac{X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}}{g(1/\log \lfloor n^\alpha \rfloor)} = 0, \quad a.s.$$

Proof. We have

$$X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} = \sum_{k=\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor}^{\lfloor n^\alpha \rfloor} \psi_k B_{\lfloor n^\alpha \rfloor - k} = \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} \psi_{\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor + j} B_{\lfloor (n-1)^\alpha \rfloor - j}.$$

This and the estimate $\psi_k \leq C\beta^k$ from the proof of Theorem 6.2 yield

$$\begin{aligned} X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} &\leq \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} C\beta^{\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor + j} B_{\lfloor (n-1)^\alpha \rfloor - j} \\ &= C\beta^{\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor} \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} \beta^j B_{\lfloor (n-1)^\alpha \rfloor - j}. \end{aligned} \quad (6.5)$$

Recall that $\beta \in (0, 1)$ and $\alpha > 1$. It is not hard to show that there exists $c > 0$ such that $\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor \geq cn^{\alpha-1}$ for large n , so

$$\begin{aligned} X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} &\leq C\beta^{cn^{\alpha-1}} \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} \beta^j B_{\lfloor (n-1)^\alpha \rfloor - j} \\ &\leq C\beta^{(c/2)n^{\alpha-1}} \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} \beta^j (\beta^{(c/2)n^{\alpha-1}} B_{\lfloor (n-1)^\alpha \rfloor - j}) \end{aligned}$$

$$\begin{aligned}
&\leq C\beta^{(c/2)n^{\alpha-1}} \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} \beta^{\lfloor (n-1)^\alpha \rfloor - j} \left(\beta^{(c/2)(n^\alpha)^{(\alpha-1)/\alpha}} B_j \right) \\
&\leq C\beta^{(c/2)n^{\alpha-1}} \sum_{j=0}^{\lfloor (n-1)^\alpha \rfloor} \beta^{\lfloor (n-1)^\alpha \rfloor - j} \left(\beta^{(c/2)j^{(\alpha-1)/\alpha}} B_j \right). \tag{6.6}
\end{aligned}$$

We use the assumption that $\mathbb{E}[(\log^+ B)^{\alpha/(\alpha-1)}] < \infty$ and inequality (4.19) to see that, for any $c_1 > 0$,

$$\begin{aligned}
\sum_{n=1}^{\infty} \mathbb{P}(B_n \geq \beta^{-c_1 n^{(\alpha-1)/\alpha}}) &= \sum_{n=1}^{\infty} \mathbb{P}(\log B_n \geq c_1 n^{(\alpha-1)/\alpha} \log \beta^{-1}) \\
&\leq \sum_{n=1}^{\infty} \mathbb{P}(\log^+ B_n \geq c_1 n^{(\alpha-1)/\alpha} \log \beta^{-1}) \\
&= \sum_{n=1}^{\infty} \mathbb{P}\left(\frac{1}{c_1 \log \beta^{-1}} \log^+ B_n \geq n^{(\alpha-1)/\alpha}\right) \\
&= \sum_{n=1}^{\infty} \mathbb{P}\left(\left(\frac{1}{c_1 \log \beta^{-1}}\right)^{\alpha/(\alpha-1)} (\log^+ B_n)^{\alpha/(\alpha-1)} \geq n\right) \\
&\leq \mathbb{E}\left(\left(\frac{1}{c_1 \log \beta^{-1}}\right)^{\alpha/(\alpha-1)} (\log^+ B_n)^{\alpha/(\alpha-1)}\right) + 1 < \infty.
\end{aligned}$$

If we take $c_1 = c/2$ then, by the Borel–Cantelli lemma, with probability 1,

$$K := \sup_{n \geq 1} \beta^{(c/2)n^{(\alpha-1)/\alpha}} B_n < \infty.$$

This and (6.6) imply that

$$X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{(n-1)^\alpha} \leq C\beta^{(c/2)n^{\alpha-1}} \sum_{j=0}^{\infty} \beta^j K = C\beta^{(c/2)n^{\alpha-1}} \frac{1}{1-\beta} K.$$

Thus, a.s.,

$$\limsup_{n \rightarrow \infty} \frac{X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{(n-1)^\alpha}}{g(1/\log \lfloor n^\alpha \rfloor)} \leq \limsup_{n \rightarrow \infty} \frac{C\beta^{\frac{\varepsilon}{2}n^{\alpha-1}}}{g(1/\log \lfloor n^\alpha \rfloor)} \cdot \frac{1}{1-\beta} K = 0. \quad \square$$

Proof of Theorem 6.3. By Lemmas 2.4, 6.5 and 6.6, for every $\varepsilon > 0$,

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} \leq \liminf_{n \rightarrow \infty} \frac{X_{\lfloor n^\alpha \rfloor}}{g(1/\log \lfloor n^\alpha \rfloor)}$$

$$\begin{aligned}
&= \liminf_{n \rightarrow \infty} \left(\frac{X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{[(n-1)^\alpha]}}{g(1/\log \lfloor n^\alpha \rfloor)} + \frac{X_{\lfloor n^\alpha \rfloor}^{[(n-1)^\alpha]}}{g(1/\log \lfloor n^\alpha \rfloor)} \right) \\
&= \liminf_{n \rightarrow \infty} \left(\frac{X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{[(n-1)^\alpha]}}{g(1/\log \lfloor n^\alpha \rfloor)} + \frac{X_{\lfloor n^\alpha \rfloor}^{[(n-1)^\alpha]}}{g(\Lambda(1+\varepsilon)/\log \lfloor n^\alpha \rfloor)} \Lambda^{1/\rho} (1+\varepsilon)^{1/\rho} \right) \\
&\leq \Lambda^{1/\rho} (1+\varepsilon)^{1/\rho}.
\end{aligned}$$

Hence, $\liminf_{n \rightarrow \infty} X_n/g(1/\log n) \leq \Lambda^{1/\rho}$, a.s. The theorem follows from this and Lemma 6.4(ii). \square

7. Random multiplicative coefficient

So far, we only considered products of IED random variables with constants. In (5.3), the multiplicative coefficient in the stochastic fixed point equation was a constant. In this section, we will look into the case when these constants are replaced with nonnegative random variables independent of other random elements of the model.

In order to solve the stochastic fixed point equation $X \stackrel{d}{=} AX + B$ we will need an assumption on the form of dependence between random variables A and B . In this paper we will assume that A and B are positively quadrant dependent. This is a well-known dependence condition, used in various models in insurance and actuarial sciences. We start with the standard definition of positive quadrant dependence.

Definition 7.1. We will call random variables X and Y positively quadrant dependent if

$$\mathbb{P}(X > x, Y > y) \geq \mathbb{P}(X > x)\mathbb{P}(Y > y), \quad (7.1)$$

for all $x, y \in \mathbb{R}$.

Remark 7.2. Note that if two random variables are independent then they are also positively quadrant dependent.

For the purposes of this paper the following characterization of positive quadrant dependence will be more useful than the original definition.

Lemma 7.3. The random variables X and Y are positively quadrant dependent if and only if

$$\mathbb{P}(X \leq x, Y \leq y) \geq \mathbb{P}(X \leq x)\mathbb{P}(Y \leq y) \quad (7.2)$$

for all $x, y \in \mathbb{R}$.

Proof. We add $-\mathbb{P}(X > x)$ to both sides of (7.1) to obtain

$$-\mathbb{P}(X > x, Y \leq y) \geq -\mathbb{P}(X > x)\mathbb{P}(Y \leq y).$$

We add $\mathbb{P}(Y \leq y)$ to both sides of the last inequality to obtain (7.2). This process can be reversed so (7.1) can be derived from (7.2). \square

Recall Definition 2.1 of essential infimum of a random variable.

Theorem 7.4. *Suppose that A is a nonnegative random variable and its essential infimum is equal to a . If X is an $\text{IED}_L^\rho(\lambda)$ -random variable and X and A are positively quadrant dependent then AX is an $\text{IED}_L^\rho(a^\rho\lambda)$ -random variable.*

Proof. Since a is the essential infimum of A , we have $a \leq A$, a.s., so

$$\mathbb{P}(AX < x) = \mathbb{P}(AX < x, a \leq A) \leq \mathbb{P}(aX < x).$$

Using Proposition 3.7,

$$\limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(AX < x) \leq \lim_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(aX < x) = -a^\rho \lambda.$$

Let $\varepsilon > 0$. The assumption that A and X are positively quadrant dependent implies that

$$\begin{aligned} \mathbb{P}(AX < x) &\geq \mathbb{P}(AX < x, a \leq A \leq a + \varepsilon) \geq \mathbb{P}((a + \varepsilon)X < x, A \leq a + \varepsilon) \\ &\geq \mathbb{P}((a + \varepsilon)X < x) \mathbb{P}(A \leq a + \varepsilon). \end{aligned}$$

By Proposition 3.7,

$$\begin{aligned} &\liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(AX < x) \\ &\geq \lim_{x \rightarrow 0^+} x^\rho L(x) (\log \mathbb{P}((a + \varepsilon)X < x) + \log \mathbb{P}(a \leq A \leq a + \varepsilon)) \\ &= -(a + \varepsilon)^\rho \lambda. \end{aligned}$$

The proof is completed by letting $\varepsilon \rightarrow 0^+$. \square

Corollary 7.5. *Suppose that independent random vectors (A_i, X_i) are such that for all $i = 1, \dots, n$,*

- (a) A_i and X_i are nonnegative and positively quadrant dependent;
- (b) X_i is an $\text{IED}_L^\rho(\lambda_i)$ -random variable.

Then $A_1 X_1 + \dots + A_n X_n$ is an $\text{IED}_L^\rho(\Lambda)$ -random variable with the parameter

$$\Lambda = ((\text{ess inf}(A_1)^\rho \lambda_1)^{1/(1+\rho)} + \dots + (\text{ess inf}(A_n)^\rho \lambda_n)^{1/(1+\rho)})^{1+\rho}.$$

Proof. The corollary follows from Theorem 7.4 and Proposition 3.12. \square

Theorem 7.6. *Let $(A_i, B_i)_{i \geq 1}$ be an i.i.d. sequence of two-dimensional vectors with the following properties.*

- (i) A_1 and B_1 are nonnegative and positively quadrant dependent.
- (ii) $\mathbb{E}[\log A_1] < 0$ and $\mathbb{E}[\log^+ B_1] < \infty$.
- (iii) B_1 is an $\text{IED}_L^\rho(\lambda)$ -random variable.

(a) The series

$$\sum_{i=1}^{\infty} \left(\prod_{j=1}^{i-1} A_j \right) B_i$$

converges a.s. to a finite $\text{IED}_L^\rho(\Lambda)$ -random variable S , where

$$\Lambda = (1 - \text{ess inf}(A_1)^{\rho/(1+\rho)})^{-1-\rho} \lambda. \quad (7.3)$$

(b) The stochastic fixed point equation $X \stackrel{d}{=} A_1 X + B_1$, where X and (A_1, B_1) are independent, has a unique solution with the same distribution as that of S .

We will need the following lemma.

Lemma 7.7. Assume that for all $a, b \in \mathbb{R}$,

$$\mathbb{P}(A \leq a, B \leq b) \geq \mathbb{P}(A' \leq a, B' \leq b),$$

with $A \stackrel{d}{=} A'$ and $B \stackrel{d}{=} B'$. If $h(x, y)$ is bounded and $\frac{\partial^2}{\partial x \partial y} h \geq 0$, then

$$\mathbb{E}[h(A, B)] \geq \mathbb{E}[h(A', B')].$$

Proof. The lemma is a special case of [11], Thm. 2. □

Proof of Theorem 7.6. (a) To simplify notation, let (A, B) have the same distribution as (A_1, B_1) . It follows from Theorem 5.1 that S is the solution to the stochastic fixed point equation $X \stackrel{d}{=} AX + B$. We set $a = \text{ess inf}(A)$, $f_B(z) = -\log \mathbb{E}[e^{-zB}]$ and $f_S(z) = -\log \mathbb{E}[e^{-zS}]$. We have assumed that $\mathbb{E}[\log A] < 0$ so $a \in [0, 1)$. By Theorem 3.11, f_B is regularly varying at infinity with index $\rho/(1+\rho)$. By the same theorem, it will suffice to show

$$\lim_{z \rightarrow \infty} \frac{f_S(z)}{f_B(z)} = (1 - a^{\rho/(1+\rho)})^{-1}. \quad (7.4)$$

If S is independent of (A, B) then $S \stackrel{d}{=} AS + B$ and, therefore,

$$e^{-f_S(z)} = \mathbb{E}e^{-zS} = \mathbb{E}e^{-z(AS+B)} \leq \mathbb{E}e^{-z(aS+B)} = \mathbb{E}e^{-zaS} \mathbb{E}e^{-zB} = e^{-f_S(az)-f_B(z)}, \quad (7.5)$$

and

$$\begin{aligned} e^{-f_S(z)} &= \mathbb{E}e^{-zS} = \mathbb{E}e^{-z(AS+B)} = \mathbb{E}(\mathbb{E}(e^{-z(AS+B)} \mid A, B)) \\ &= \mathbb{E}(e^{-zB} \mathbb{E}(e^{-zaS} \mid A, B)) = \mathbb{E}(e^{-zB} e^{-f_S(za)}) = \mathbb{E}(e^{-f_S(za)-zB}). \end{aligned} \quad (7.6)$$

It follows from (7.5) that $f_S(z) \geq f_S(az) + f_B(z)$ and

$$\begin{aligned} \liminf_{z \rightarrow \infty} \frac{f_S(z)}{f_B(z)} &\geq \liminf_{z \rightarrow \infty} \frac{f_S(az)}{f_B(z)} + 1 = \liminf_{z \rightarrow \infty} \frac{f_B(az)}{f_B(z)} \frac{f_S(az)}{f_B(az)} + 1 \\ &= a^{\rho/(1+\rho)} \liminf_{z \rightarrow \infty} \frac{f_S(z)}{f_B(z)} + 1, \end{aligned}$$

hence

$$\liminf_{z \rightarrow \infty} \frac{f_S(z)}{f_B(z)} \geq (1 - a^{\rho/(1+\rho)})^{-1}. \quad (7.7)$$

We will apply Lemma 7.7 to the function

$$h(x, y) = \exp(-f_S(xz) - yz) = \mathbb{E} \exp(-xzS - yz)$$

and independent random variables A' and B' such that $A' \stackrel{d}{=} A$ and $B' \stackrel{d}{=} B$. Since A and B are positively quadrant dependent, Lemma 7.3 implies that

$$\mathbb{P}(A \leq a, B \leq b) \geq \mathbb{P}(A \leq a)\mathbb{P}(B \leq b) = \mathbb{P}(A' \leq a)\mathbb{P}(B' \leq b) = \mathbb{P}(A' \leq a, B' \leq b).$$

It is easy to check that $\frac{\partial^2}{\partial x \partial y} h \geq 0$. Hence, by Lemma 7.7, for a fixed $\varepsilon \in (0, 1 - a)$,

$$\begin{aligned} \mathbb{E}[e^{-f_S(Az) - zB}] &\geq \mathbb{E}[e^{-f_S(A'z) - zB'}] = \mathbb{E}[e^{-f_S(A'z)}] \mathbb{E}[e^{-zB'}] = \mathbb{E}[e^{-f_S(Az)}] \mathbb{E}[e^{-zB}] \\ &\geq \mathbb{E}[e^{-f_S(Az)} \mathbf{1}_{\{A < a + \varepsilon\}}] e^{-f_B(z)} \geq e^{-f_S((a+\varepsilon)z) - f_B(z)} \mathbb{P}(A < a + \varepsilon). \end{aligned}$$

This and (7.6) imply that

$$f_S((a + \varepsilon)z) + f_B(z) - \log \mathbb{P}(A < a + \varepsilon) \geq f_S(z).$$

Substituting $(a + \varepsilon)^k z$ for z in the last formula yields

$$f_S((a + \varepsilon)^{k+1} z) + f_B((a + \varepsilon)^k z) - \log \mathbb{P}(A < a + \varepsilon) \geq f_S((a + \varepsilon)^k z).$$

The telescoping sum argument gives

$$\begin{aligned} f_S(z) - f_S((a + \varepsilon)^{n+1} z) &\leq \sum_{k=0}^n f_B((a + \varepsilon)^k z) - (n + 1) \log \mathbb{P}(A < a + \varepsilon) \\ &\leq \sum_{k=0}^{\infty} f_B((a + \varepsilon)^k z) - (n + 1) \log \mathbb{P}(A < a + \varepsilon). \end{aligned} \quad (7.8)$$

Note that

$$\sum_{k=0}^{\infty} f_B((a + \varepsilon)^k z) = -\log \mathbb{E} \left[\exp \left(-z \sum_{k=0}^{\infty} (a + \varepsilon)^k B_k \right) \right].$$

It follows from Proposition 3.7 and Theorem 4.7 that $\sum_{k=0}^{\infty} (a + \varepsilon)^k B_k$ is a finite $\text{IED}_L^\rho(\Lambda_1)$ -random variable, where

$$\Lambda_1 = \lambda \left(\sum_{k=0}^{\infty} (a + \varepsilon)^{k\rho/(1+\rho)} \right)^{1+\rho} = \frac{\lambda}{(1 - (a + \varepsilon)^{\rho/(1+\rho)})^{1+\rho}}.$$

Hence, by Theorem 3.11 we have

$$\sum_{k=0}^{\infty} f_B((a + \varepsilon)^k z) \sim (1 - (a + \varepsilon)^{\rho/(1+\rho)})^{-1} f_B(z). \quad (7.9)$$

If we take $n = \lceil -\log z / \log(a + \varepsilon) \rceil$ then $(a + \varepsilon)^2 \leq (a + \varepsilon)^{n+1} z \leq 1$ and, therefore $|f_S((a + \varepsilon)^{n+1} z)| < c_1$. Also,

$$|(n+1) \log \mathbb{P}(A < a + \varepsilon)| \leq \frac{|2 \log \mathbb{P}(A < a + \varepsilon)|}{\log(a + \varepsilon)} \log z.$$

These observations, the fact that f_B is regularly varying at infinity with index $\rho/(1 + \rho) > 0$, (7.8) and (7.9), imply that

$$\limsup_{z \rightarrow \infty} \frac{f_S(z)}{f_B(z)} \leq \lim_{z \rightarrow \infty} \sum_{k=0}^{\infty} \frac{f_B((a + \varepsilon)^k z)}{f_B(z)} = (1 - (a + \varepsilon)^{\rho/(1+\rho)})^{-1}. \quad (7.10)$$

This completes the proof of (7.4) because the above estimate holds for all sufficiently small $\varepsilon > 0$ and we already have (7.7).

(b) Part (b) follows from Theorem 5.1 and part (a). \square

Methods similar to those in the proof of Theorem 7.6 were used in [10] to analyze light-tailed solutions to $X \stackrel{d}{=} AX + B$.

We will now interpret the parameter Λ in (7.3) in a way similar to that in Proposition 5.4.

Theorem 7.8. *Let $(A_i, B_i)_{i=1}^{\infty}$ be an i.i.d. sequence of two-dimensional vectors with the following properties.*

- (i) A_1 and B_1 are nonnegative and positively quadrant dependent random variables.
- (ii) There exists $\beta \in (0, 1)$ such that $A_1 \leq \beta$, a.s.
- (iii) $\mathbb{E}[(\log^+ B_1)^s] < \infty$ for all $s > 0$.
- (iv) B_1 is an $\text{IED}_L^\rho(\lambda)$ -random variable.

Let $X_0 = 0$ and

$$X_n = A_n X_{n-1} + B_n, \quad n \geq 1. \quad (7.11)$$

Recall Λ defined in (7.3) and let g be the asymptotic inverse of $x \mapsto x^\rho L(x)$ at 0. Then

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} = \Lambda^{1/\rho}, \quad \text{a.s.}$$

The structure of the proof will be similar to that of Theorem 6.3, however some new technical steps will be needed. In all lemmas preceding the proof of Theorem 7.8, we will use the same notation and make the same assumptions as in the theorem.

Lemma 7.9.

(a) Let S be defined as in (5.2). We have $X_n \xrightarrow{d} S$ as $n \rightarrow \infty$ and

$$\mathbb{P}(S \leq x) \leq \mathbb{P}(X_n \leq x) \leq \mathbb{P}(X_{n-1} \leq x), \quad (7.12)$$

for all $n \geq 1$ and $x \geq 0$.

(b) Let $a = \text{ess inf } A_1$. For every $n \geq 1$, X_n is an $\text{IED}_L^\rho(\Lambda_n)$ -random variable with

$$\Lambda_n = \lambda \left(\frac{1 - a^{\frac{n\rho}{1+\rho}}}{1 - a^{\rho/(1+\rho)}} \right)^{1+\rho}. \quad (7.13)$$

Proof. Set $S_n = \sum_{i=1}^n (\prod_{j=1}^{i-1} A_j) B_i$. Definition (7.11) implies the following representation, $X_n = \sum_{i=1}^n (\prod_{j=n-i+2}^n A_j) B_{n-i+1}$. This and the change of index $i \mapsto n - i + 1$ easily show that $X_n \stackrel{d}{=} S_n$. Therefore, $\mathbb{P}(X_n \leq x) = \mathbb{P}(S_n \leq x)$. Since A_j 's and B_j 's are non-negative, $S_n \uparrow S$ a.s. All claims made in part (a) follow easily from these observations.

The definition of the essential infimum a and the assumption that A_i 's and B_i 's are non-negative imply that

$$\mathbb{P}(S_n \leq x) \leq \mathbb{P}(a^{n-1} B_n + a^{n-2} B_{n-2} + \cdots + a B_2 + B_1 \leq x). \quad (7.14)$$

We use the assumption that (A_k, B_k) , $k \geq 1$ are i.i.d., (A_1, B_1) are positively quadrant dependent, and Lemma 7.3 to see that for $\varepsilon > 0$,

$$\begin{aligned} \mathbb{P}(S_n \leq x) &\geq \mathbb{P}(S_n \leq x, A_1 \leq a + \varepsilon, A_2 \leq a + \varepsilon, \dots, A_{n-1} \leq a + \varepsilon) \\ &\geq \mathbb{P}((a + \varepsilon)^{n-1} B_n + (a + \varepsilon)^{n-2} B_{n-2} + \cdots + (a + \varepsilon) B_2 + B_1 \leq x, \\ &\quad A_1 \leq a + \varepsilon, A_2 \leq a + \varepsilon, \dots, A_{n-1} \leq a + \varepsilon) \\ &\geq \mathbb{P}((a + \varepsilon)^{n-1} B_n + (a + \varepsilon)^{n-2} B_{n-2} + \cdots + (a + \varepsilon) B_2 + B_1 \leq x, \\ &\quad A_2 \leq a + \varepsilon, \dots, A_{n-1} \leq a + \varepsilon) \mathbb{P}(A_1 \leq a + \varepsilon) \\ &\geq \mathbb{P}((a + \varepsilon)^{n-1} B_n + (a + \varepsilon)^{n-2} B_{n-2} + \cdots + (a + \varepsilon) B_2 + B_1 \leq x, \\ &\quad A_3 \leq a + \varepsilon, \dots, A_{n-1} \leq a + \varepsilon) \mathbb{P}(A_1 \leq a + \varepsilon) \mathbb{P}(A_2 \leq a + \varepsilon) \\ &\quad \dots \\ &\geq \mathbb{P}((a + \varepsilon)^{n-1} B_n + (a + \varepsilon)^{n-2} B_{n-2} + \cdots + (a + \varepsilon) B_2 + B_1 \leq x) \\ &\quad \times \mathbb{P}(A_1 \leq a + \varepsilon)^{n-1}. \end{aligned} \quad (7.15)$$

We have

$$\sum_{k=0}^{n-1} (a^k)^{\rho/(1+\rho)} = \frac{1 - a^{n\rho/(1+\rho)}}{1 - a^{\rho/(1+\rho)}}, \quad \sum_{k=0}^{n-1} ((a + \varepsilon)^k)^{\rho/(1+\rho)} = \frac{1 - (a + \varepsilon)^{n\rho/(1+\rho)}}{1 - (a + \varepsilon)^{\rho/(1+\rho)}}. \quad (7.16)$$

Recall that $\mathbb{P}(X_n \leq x) = \mathbb{P}(S_n \leq x)$ and use (7.14), (7.15), (7.16) and Proposition 3.12 to see that

$$\begin{aligned} \limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X_n \leq x) &= \limsup_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(S_n \leq x) \leq \lambda \left(\frac{1 - a^{\frac{n\rho}{1+\rho}}}{1 - a^{\frac{\rho}{1+\rho}}} \right)^{1+\rho}, \\ \liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(X_n \leq x) &= \liminf_{x \rightarrow 0^+} x^\rho L(x) \log \mathbb{P}(S_n \leq x) \geq \lambda \left(\frac{1 - (a + \varepsilon)^{\frac{n\rho}{1+\rho}}}{1 - (a + \varepsilon)^{\frac{\rho}{1+\rho}}} \right)^{1+\rho}. \end{aligned}$$

We complete the proof of (b) by letting $\varepsilon \rightarrow 0$. □

Lemma 7.10. (i) For every $\varepsilon > 0$,

$$\left\{ X_n \leq g \left(\frac{\Lambda}{(1 + \varepsilon) \log n} \right) \right\}$$

happens finitely often almost surely.

(ii) We have

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} \geq \Lambda^{1/\rho} \quad a.s.$$

Proof. (i) For any $\varepsilon > 0$ there exists $\delta \in (0, 1)$ such that $\gamma := (1 - \delta)(1 + \varepsilon/2) > 1$. Assumption (ii) of Theorem 7.8 implies that $a < 1$ so Λ_n defined in (7.13) converge to Λ as $n \rightarrow \infty$. By Lemma 3.9 and part (b) of Lemma 7.9, there exist C_δ , n_0 and $x_0 > 0$ such that $\mathbb{P}(X_{n_0} \leq x) \leq C_\delta e^{-\Lambda(1-\delta)/(x^\rho L(x))}$ for all $x \in (0, x_0)$. By part (a) of Lemma 7.9, for $n \geq n_0$ and $x \in (0, x_0)$,

$$\mathbb{P}(X_n \leq x) \leq C_\delta e^{-\Lambda(1-\delta)/(x^\rho L(x))}.$$

It follows that, for large n ,

$$\mathbb{P} \left(X_n \leq g \left(\frac{\Lambda}{(1 + \varepsilon) \log n} \right) \right) \leq C_\delta e^{-(1+\varepsilon/2)(1-\delta) \log n} = C_\delta n^{-\gamma}.$$

Hence,

$$\sum_{n=1}^{\infty} \mathbb{P} \left(X_n \leq g \left(\frac{\Lambda}{(1 + \varepsilon) \log n} \right) \right) < \infty,$$

and the claim follows by the Borel–Cantelli lemma.

(ii) Part (i) implies that for every $\varepsilon > 0$,

$$\liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} \geq \frac{\Lambda^{1/\rho}}{(1+\varepsilon)^{1/\rho}}.$$

Part (ii) follows by letting $\varepsilon \rightarrow 0$. □

From definition (7.11), we obtain for $n \geq m$,

$$\begin{aligned} X_n &= B_n + A_n X_{n-1} = B_n + A_n B_{n-1} + A_n A_{n-1} X_{n-2} = \cdots \\ &= \sum_{j=m+1}^n \left(\prod_{k=j+1}^n A_k \right) B_j + \left(\prod_{k=m+1}^n A_k \right) X_m. \end{aligned} \quad (7.17)$$

We rearrange terms and define new random variables,

$$X_n^m := X_n - \left(\prod_{k=m+1}^n A_k \right) X_m = \sum_{j=m+1}^n \left(\prod_{k=j+1}^n A_k \right) B_j. \quad (7.18)$$

The notation X_n^m is the same as in (6.4) but the meaning is different. We have chosen the same notation for a different object because this will allow us to reuse a part of the proof of Lemma 6.6

Lemma 7.11.

(a) Let $Y_0 = 0$ and $Y_k := X_{m+k}^m$ for $k \geq 1$. The sequence $(Y_k, k \geq 0)$ satisfies

$$Y_{k+1} = A_{m+k+1} Y_k + B_{m+k+1}.$$

(b) For every $m \geq 0$, the sequence $(X_{m+k}^m, k \geq 0)$ has the same distribution as $(X_k, k \geq 0)$.

Proof. The claim in (a) follows from (7.18). Part (b) follows from (a). □

Lemma 7.12. Fix $\varepsilon > 0$ and suppose that $1 < 1 + \delta < \sqrt{1 + \varepsilon}$ and $\alpha \in (1, 1 + \delta)$. Events

$$\left\{ X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right) \right\}$$

happen infinitely often a.s.

Proof. We use Lemma 7.11(b) and (7.12) to see that

$$\begin{aligned} \sum_{n=1}^{\infty} \mathbb{P}\left(X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) &= \sum_{n=1}^{\infty} \mathbb{P}\left(X_{\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) \\ &\geq \sum_{n=1}^{\infty} \mathbb{P}\left(S \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right). \end{aligned}$$

The random variable S is $\text{IED}_L^\rho(\Lambda)$ by Theorem 7.6, so we can use Lemma 3.9 to write

$$\mathbb{P}\left(S \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) \geq \mathbb{P}\left(S \leq g\left(\frac{\Lambda(1+\delta)}{\log n^{\alpha/(1+\delta)}}\right)\right) \geq c_\delta n^{-\alpha/(1+\delta)}.$$

It follows that

$$\sum_{n=2}^{\infty} \mathbb{P}\left(X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} \leq g\left(\frac{\Lambda(1+\varepsilon)}{\log n^\alpha}\right)\right) = \infty.$$

This, the fact that the random variables $X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}$, $n \geq 2$, are jointly independent and the Borel–Cantelli lemma imply the claim made in the lemma. \square

Lemma 7.13. *When $n \rightarrow \infty$,*

$$\frac{X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}}{g(\log \lfloor n^\alpha \rfloor)} \rightarrow 0, \quad a.s.$$

Proof. We first use (7.18) and then (7.17) with $m = 0$ to obtain

$$\begin{aligned} X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} &= \left(\prod_{k=\lfloor (n-1)^\alpha \rfloor + 1}^{\lfloor n^\alpha \rfloor} A_k \right) X_{\lfloor (n-1)^\alpha \rfloor} \\ &= \left(\prod_{k=\lfloor (n-1)^\alpha \rfloor + 1}^{\lfloor n^\alpha \rfloor} A_k \right) \left(\sum_{j=1}^{\lfloor (n-1)^\alpha \rfloor} \left(\prod_{k=j+1}^{\lfloor (n-1)^\alpha \rfloor} A_k \right) B_j \right) \\ &= \sum_{j=1}^{\lfloor (n-1)^\alpha \rfloor} \left(\prod_{k=j+1}^{\lfloor n^\alpha \rfloor} A_k \right) B_j. \end{aligned}$$

We apply assumption (ii) of Theorem 7.8, a part of assumption (i) (namely, that A_k 's and B_k 's are nonnegative), and the change of index $i = \lfloor (n-1)^\alpha \rfloor - j$, to see that

$$\begin{aligned} X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor} &= \sum_{j=1}^{\lfloor (n-1)^\alpha \rfloor} \left(\prod_{k=j+1}^{\lfloor n^\alpha \rfloor} A_k \right) B_j \leq \sum_{j=1}^{\lfloor (n-1)^\alpha \rfloor} \beta^{\lfloor n^\alpha \rfloor - j} B_j \\ &= \sum_{i=0}^{\lfloor (n-1)^\alpha \rfloor - 1} \beta^{\lfloor n^\alpha \rfloor - \lfloor (n-1)^\alpha \rfloor + i} B_{\lfloor (n-1)^\alpha \rfloor - i}. \end{aligned}$$

The rest of the proof is the same as the proof of Lemma 6.6 starting at (6.5), with $C = 1$. \square

Proof of Theorem 7.8. Note that g is regularly varying with index $1/\rho$ at 0. By Lemmas 7.12 and 7.13, for every $\varepsilon > 0$,

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{X_n}{g(1/\log n)} &\leq \liminf_{n \rightarrow \infty} \frac{X_{\lfloor n^\alpha \rfloor}}{g(1/\log(\lfloor n^\alpha \rfloor))} \\ &= \liminf_{n \rightarrow \infty} \left(\frac{X_{\lfloor n^\alpha \rfloor} - X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}}{g(1/\log(\lfloor n^\alpha \rfloor))} + \frac{X_{\lfloor n^\alpha \rfloor}^{\lfloor (n-1)^\alpha \rfloor}}{g(1/\log(\lfloor n^\alpha \rfloor))} \right) \\ &\leq \Lambda^{1/\rho} (1 + \varepsilon)^{1/\rho}. \end{aligned}$$

Hence, $\liminf_{n \rightarrow \infty} X_n/g(1/\log(n)) \leq \Lambda^{1/\rho}$ a.s. The theorem follows from this and Lemma 7.10(ii). \square

The following theorem is a version of the well-known results by Kesten [9] and Goldie [7], formulated in [4], Theorem 2.4.4.

Theorem 7.14. Assume that (A, B) satisfy the following conditions.

- (i) $A \geq 0$, a.s., and the law of $\log A$ conditioned on $\{A > 0\}$ is non-arithmetic, i.e., it is not supported on $a\mathbb{Z}$ for any $a > 0$.
- (ii) There exists $\alpha > 0$ such that $\mathbb{E}[A^\alpha] = 1$, $\mathbb{E}[|B|^\alpha] < \infty$ and $\mathbb{E}[A^\alpha \log^+ A] < \infty$.
- (iii) $\mathbb{P}(Ax + B = x) < 1$ for every $x \in \mathbb{R}$.

Then the equation $X \stackrel{d}{=} AX + B$ has a solution. There exist constants c_+, c_- such that $c_+ + c_- > 0$ and

$$\mathbb{P}(X > x) \sim c_+ x^{-\alpha} \quad \text{and} \quad \mathbb{P}(X < -x) \sim c_- x^{-\alpha}, \quad \text{when } x \rightarrow \infty. \quad (7.19)$$

The constants c_+, c_- are given by

$$c_+ = \frac{1}{\alpha m_\alpha} \mathbb{E}[(AX + B)_+^\alpha - (AX)_+^\alpha], \quad c_- = \frac{1}{\alpha m_\alpha} \mathbb{E}[(AX + B)_-^\alpha - (AX)_-^\alpha],$$

where $m_\alpha = \mathbb{E}[A^\alpha \log A]$.

Next we will combine the results of Kesten and Goldie with our own.

Theorem 7.15. Assume that (A, B) satisfy the following conditions.

- (i) A and B are nonnegative, non-constant and positively quadrant dependent random variables.
- (ii) There exists $\alpha > 0$ such that $\mathbb{E}[A^\alpha] = 1$, $\mathbb{E}[A^\alpha \log A] < \infty$ and $\log A$ conditioned on $\{A > 0\}$ is a non-arithmetic random variable.
- (iii) B is an $\text{IED}_L^\rho(\lambda)$ -random variable and $\mathbb{E}[B^\alpha] < \infty$.

Then the stochastic fixed point equation $X \stackrel{d}{=} AX + B$ has a unique solution which is an $\text{IED}_L^\rho(\Lambda)$ random variable, where Λ is defined in (7.3).

Moreover,

$$\lim_{x \rightarrow \infty} x^\alpha \mathbb{P}(X > x) \quad (7.20)$$

exists and is a positive number.

Proof. We will show that assumptions of Theorem 7.6 are satisfied. Since $x \mapsto \log x$ is concave and A is non-constant, we have $\alpha \mathbb{E}[\log A] < \log \mathbb{E}[A^\alpha] = 0$, so assumption (ii) of Theorem 7.6 holds. The other assumptions also hold so the first claim follows from Theorem 7.6.

We note that assumptions (i) and (ii) of Theorem 7.14 hold. We will verify assumption (iii). The function $\log A$ is non-arithmetic when conditioned on $\{A > 0\}$, hence $\mathbb{P}(A \neq 1) > 0$. Since $\mathbb{E}[A^\alpha] = 1$, there exists $\alpha \in (0, 1)$ such that $\mathbb{P}(A \leq \alpha) > 0$. Random variables A and B are positive quadrant dependent and B is an $\text{IED}_L^\rho(\lambda)$ -random variable so

$$\mathbb{P}(Ax + B \neq x) \geq \mathbb{P}(Ax \leq \alpha x, B \leq (1 - \alpha/2)x) \geq \mathbb{P}(A \leq \alpha) \mathbb{P}(B \leq (1 - \alpha/2)x) > 0.$$

All assumptions of Theorem 7.14 have been verified, so the claim (7.20) is a consequence of (7.19). \square

8. Dependent coefficients in the fixed point equation

This section has a double purpose. First, we will explain how the questions studied in this paper arose in a different project. That project is devoted to a rather different topic so we will only sketch some of its ideas. Needless to say, we hope that our present results will be used to study other models.

Second, the fixed point equation (1.1) coming from the other project has coefficients A and B dependent in a different way than in the previous sections of this paper. We plan to develop a theory for such equations in a future article. Here we will limit ourselves to showing that the lack of positive quadrant dependence can make a substantial difference to the main results on IED random variables.

8.1. Motivation

In the rest of this section, we will assume that the vector (A, B) has the following density.

$$\mathbb{P}(A \in da, B \in db) = \frac{e^{-(a^{1/2} - \frac{1}{2})^2/b} - e^{-(a^{1/2} + \frac{1}{2})^2/b}}{2\sqrt{\pi ab}} \cdot \frac{e^{-1/(4b)}}{\sqrt{\pi b^{3/2}}}, \quad a, b > 0. \quad (8.1)$$

We will now explain how this density arose in a project on the Fleming–Viot type process (see [5]). Our new results are in preparation but one can find the following basic scheme in [1]. Let:

- $W_1 = (W_1(t) : t \geq 0)$ and $W_2 = (W_2(t) : t \geq 0)$ be two independent Brownian motions;

- τ_1 and τ_2 , respectively, be the first times W_1 and W_2 hit 0;
- $T = \min\{\tau_1, \tau_2\}$ and $Y = \max\{W_1(T), W_2(T)\}$.

One can show that

$$\mathbb{P}(Y \in dy, T \in dt) = \frac{\exp(-\frac{(1-y)^2+1}{2t}) - \exp(-\frac{(1+y)^2+1}{2t})}{\pi t^2} dt dy, \quad y, t > 0.$$

The distribution given in (8.1) is obtained by the substitution $A = Y^{-2}$, $B = TY^{-2}$.

Let Y_n denote the position and let T_n denote the time of the n -th renewal of the Fleming–Viot type process. Self-similarity of the process implies that $(T_n/Y_n^2)_{n \geq 1}$ is an iterated random sequence, whose limiting behavior is described by the stochastic fixed point equation $X \stackrel{d}{=} AX + B$. We are interested in the right tail behavior of $Y_n/\sqrt{T_n}$ for large n , so we could show that $\liminf_{n \rightarrow \infty} Y_n/\sqrt{T_n \log \log T_n}$ is a constant. It turns out that this is equivalent to estimating $\mathbb{P}(X < x)$ as $x \rightarrow 0^+$.

8.2. Dependent coefficients

We start with some basic facts about the distribution defined in (8.1).

Recall that \sim means that the ratio of two quantities converges to 1.

Proposition 8.1. *Assume that the vector (A, B) has the distribution given by (8.1).*

(a) *The density of A is*

$$\mathbb{P}(A \in da) = \frac{4}{\pi(4a^2 + 1)} da. \quad (8.2)$$

(b) *The density of B is*

$$\mathbb{P}(B \in db) = \frac{1}{\sqrt{\pi}} \left(\int_{-\frac{1}{\sqrt{2b}}}^{\frac{1}{\sqrt{2b}}} \frac{e^{-v^2/2}}{\sqrt{2\pi}} dv \right) \frac{e^{-1/(4b)}}{b^{3/2}} db. \quad (8.3)$$

Moreover,

$$\mathbb{P}(B < x) \sim \int_0^x \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db \quad \text{as } x \rightarrow 0^+, \quad (8.4)$$

$$\mathbb{P}(B > x) \sim \frac{1}{\pi x} \quad \text{as } x \rightarrow \infty. \quad (8.5)$$

Random variable B is $\text{IED}_1^1(1/4)$.

Proof. (a) We integrate the density of (A, B) with respect to b over $(0, \infty)$ to compute the density of A .

$$\mathbb{P}(A \in da) = \int_0^\infty \frac{\exp(-\frac{(a^{1/2}-\frac{1}{2})^2+\frac{1}{4}}{b}) - \exp(-\frac{(a^{1/2}+\frac{1}{2})^2+\frac{1}{4}}{b})}{2\pi\sqrt{ab^2}} db da$$

$$\begin{aligned}
&= \frac{1}{2\pi\sqrt{a}} \left[\int_0^\infty \frac{\exp(-\frac{(a^{1/2}-\frac{1}{2})^2+\frac{1}{4}}{b})}{b^2} db - \int_0^\infty \frac{\exp(-\frac{(a^{1/2}+\frac{1}{2})^2+\frac{1}{4}}{b})}{b^2} db \right] da \\
&= \frac{1}{2\pi\sqrt{a}} \left[\frac{1}{(a^{1/2}-\frac{1}{2})^2+\frac{1}{4}} - \frac{1}{(a^{1/2}+\frac{1}{2})^2+\frac{1}{4}} \right] da \\
&= \frac{da}{\pi(a^2+\frac{1}{4})}.
\end{aligned}$$

(b) In the following calculation, we use the substitution $u = \sqrt{a}$.

$$\begin{aligned}
&\mathbb{P}(B \in db) \\
&= \int_0^\infty \frac{\exp(-\frac{(a^{1/2}-\frac{1}{2})^2}{b}) - \exp(-\frac{(a^{1/2}+\frac{1}{2})^2}{b})}{2\sqrt{\pi ab}} da \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db \\
&= \int_0^\infty \frac{\exp(-\frac{(u-\frac{1}{2})^2}{b}) - \exp(-\frac{(u+\frac{1}{2})^2}{b})}{\sqrt{\pi b}} du \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db \\
&= \left(\int_0^\infty \frac{\exp(-\frac{(u-\frac{1}{2})^2}{b})}{\sqrt{\pi b}} du - \int_0^\infty \frac{\exp(-\frac{(u+\frac{1}{2})^2}{b})}{\sqrt{\pi b}} du \right) \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db \\
&= \left(\int_{-\frac{1}{2}}^\infty \frac{e^{-u^2/b}}{\sqrt{\pi b}} du - \int_{\frac{1}{2}}^\infty \frac{e^{-u^2/b}}{\sqrt{\pi b}} du \right) \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db \\
&= \left(\int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{e^{-u^2/b}}{\sqrt{\pi b}} du \right) \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db \\
&= \left(\int_{-\frac{1}{\sqrt{2b}}}^{\frac{1}{\sqrt{2b}}} \frac{e^{-v^2/2}}{\sqrt{2\pi}} dv \right) \frac{e^{-1/(4b)}}{\sqrt{\pi} b^{3/2}} db.
\end{aligned}$$

This proves (8.3). We use (8.3) and the following facts,

$$\lim_{b \rightarrow \infty} \sqrt{b/2} \int_{-\frac{1}{\sqrt{2b}}}^{\frac{1}{\sqrt{2b}}} \frac{e^{-v^2/2}}{\sqrt{2\pi}} dv = \frac{1}{\sqrt{2\pi}} \quad \text{and} \quad \lim_{b \rightarrow \infty} e^{-1/(4b)} = 1,$$

to conclude that, when $x \rightarrow \infty$,

$$\mathbb{P}(B > x) \sim \frac{1}{\pi} \int_x^\infty b^{-2} db = \frac{1}{\pi x}.$$

This proves (8.5). Since

$$\lim_{b \rightarrow 0^+} \int_{-\frac{1}{\sqrt{2b}}}^{\frac{1}{\sqrt{2b}}} \frac{e^{-v^2/2}}{\sqrt{2\pi}} dv = 1,$$

we obtain for $x \rightarrow 0^+$,

$$\mathbb{P}(B < x) \sim \int_0^x \frac{e^{-1/(4b)}}{\sqrt{\pi}b^{3/2}} db.$$

This proves (8.4). The claim that B is an $\text{IED}_1^1(\frac{1}{4})$ -random variable follows from (8.4) by the same arguments as in Lemma 3.5. \square

Lemma 8.2. *If (A_1, B_1) and (A_2, B_2) are independent random vectors with the density (8.1) then*

$$\limsup_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(A_2 B_1 + B_2 < \varepsilon) \leq -\frac{3}{10} < -\frac{1}{4}. \quad (8.6)$$

Remark 8.3. Suppose that (A_1, B_1) and (A_2, B_2) are i.i.d., and A_1 and B_1 are independent with distributions (8.2) and (8.3). Then $\text{ess inf}(A_1) = 0$ and B_1 is $\text{IED}_1^1(\frac{1}{4})$. By Corollary 7.5, $\lim_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(A_2 B_1 + B_2 < \varepsilon) = -\frac{1}{4}$. However, (8.6) shows that we do not have the same conclusion when A_1 and B_1 are not independent.

Proof of Lemma 8.2. First, recall that by Lemma 3.9, for every $\delta > 0$ there exists $C > 0$ such that for all $x > 0$,

$$\mathbb{P}(B_1 < x) \leq C \exp\left(-\left(\frac{1}{4} - \delta\right)x^{-1}\right).$$

The second inequality in the following calculation is justified by the above formula. Later in the calculation, we will use the substitution $y = a^{1/2}$.

$$\begin{aligned} & \mathbb{P}(A_2 B_1 + B_2 < \varepsilon) \\ &= \int_0^\varepsilon \int_0^\infty \mathbb{P}(a B_1 + b < \varepsilon) \frac{\exp(-\frac{(a^{1/2} - \frac{1}{2})^2 + \frac{1}{4}}{b}) - \exp(-\frac{(a^{1/2} + \frac{1}{2})^2 + \frac{1}{4}}{b})}{2\pi b^2 \sqrt{a}} da db \\ &\leq \int_0^\varepsilon \int_0^\infty \mathbb{P}\left(B_1 < \frac{\varepsilon - b}{a}\right) \frac{\exp(-\frac{(a^{1/2} - \frac{1}{2})^2 + \frac{1}{4}}{b})}{2\pi b^2 \sqrt{a}} da db \\ &\leq \int_0^\varepsilon \int_0^\infty C \exp\left(-\left(\frac{1}{4} - \delta\right) \frac{a}{\varepsilon - b}\right) \frac{\exp(-\frac{(a^{1/2} - \frac{1}{2})^2 + \frac{1}{4}}{b})}{2\pi b^2 \sqrt{a}} da db \\ &\leq \int_0^\varepsilon \int_0^\infty C \exp\left(-\left(\frac{1}{4} - \delta\right) \frac{a}{\varepsilon}\right) \frac{\exp(-\frac{(a^{1/2} - \frac{1}{2})^2}{\varepsilon} - \frac{1}{4b})}{2\pi b^2 \sqrt{a}} da db \\ &= C \int_0^\infty \frac{1}{2\sqrt{\pi}a} \exp\left(-\varepsilon^{-1} \left[\left(a^{1/2} - \frac{1}{2}\right)^2 + \left(\frac{1}{4} - \delta\right)a\right]\right) da \int_0^\varepsilon \frac{e^{-1/(4b)}}{\sqrt{\pi}b^2} db \\ &= C \int_0^\infty \frac{1}{2\sqrt{\pi}a} \exp\left(-\varepsilon^{-1} \left[\left(\frac{5}{4} - \delta\right) \left(a^{1/2} - \frac{2}{5-4\delta}\right)^2 + \frac{1}{4} - \frac{1}{5-4\delta}\right]\right) da \end{aligned}$$

$$\begin{aligned}
& \times \int_0^\varepsilon \frac{e^{-1/(4b)}}{\sqrt{\pi}b^2} db \\
& \leq C\sqrt{2} \left[\frac{\varepsilon}{2} \left(\frac{5}{4} - \delta \right)^{-1} \right]^{1/2} \int_{-\infty}^\infty \frac{1}{\sqrt{2\pi}[\frac{\varepsilon}{2}(\frac{5}{4} - \delta)^{-1}]} \exp\left(-\frac{(y - \frac{2}{5-4\delta})^2}{2[\frac{\varepsilon}{2}(\frac{5}{4} - \delta)^{-1}]}\right) dy \\
& \quad \times \exp\left(-\varepsilon^{-1}\left(\frac{1}{4} - \frac{1}{5-4\delta}\right)\right) \int_0^\varepsilon \frac{e^{-1/(4b)}}{\sqrt{\pi}b^2} db \\
& = C\sqrt{2} \left[\frac{\varepsilon}{2} \left(\frac{5}{4} - \delta \right)^{-1} \right]^{1/2} \exp\left(-\varepsilon^{-1}\left(\frac{1}{4} - \frac{1}{5-4\delta}\right)\right) \int_0^\varepsilon \frac{e^{-1/(4b)}}{\sqrt{\pi}b^2} db.
\end{aligned}$$

By the same argument as in the proof of Lemma 3.5,

$$\limsup_{\varepsilon \rightarrow 0^+} \varepsilon \log \int_0^\varepsilon \frac{e^{-1/(4b)}}{\sqrt{\pi}b^2} db = -1/4.$$

This and the previous estimate yield

$$\limsup_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(A_2 B_1 + B_2 < \varepsilon) \leq -\frac{1}{4} - \frac{1}{4} + \frac{1}{5-4\delta}.$$

The proof of the lemma is completed by letting $\delta \rightarrow 0^+$. □

Proposition 8.4. Assume that the vector (A, B) has the distribution given by (8.1). The stochastic fixed point equation

$$X \stackrel{d}{=} AX + B, \quad (8.7)$$

has a unique solution and we have

$$\limsup_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(X < \varepsilon) \leq -\frac{3}{10} < -\frac{1}{4}. \quad (8.8)$$

Proof. We use the substitution $a = x/2$ in the following calculation.

$$\begin{aligned}
\mathbb{E}[\log A] &= \int_0^\infty \frac{4 \log a}{\pi(4a^2 + 1)} da = 2 \int_0^\infty \frac{\log(x/2)}{\pi(x^2 + 1)} dx = 2 \int_0^\infty \frac{\log x - \log 2}{\pi(x^2 + 1)} dx \\
&= -2 \log 2 \int_0^\infty \frac{1}{\pi(x^2 + 1)} dx + \int_0^\infty \frac{\log x}{\pi(x^2 + 1)} dx.
\end{aligned}$$

It is easy to see that $\int_0^\infty \frac{1}{\pi(x^2 + 1)} dx = 1/2$. The substitution $y = x^{-1}$ yields $\int_0^\infty \frac{\log x}{\pi(x^2 + 1)} dx = -\int_0^\infty \frac{\log y}{\pi(y^2 + 1)} dy$, so these integrals must be equal to 0. Hence, $\mathbb{E}[\log A] = -\log 2 < 0$.

It follows from (8.5) that $\mathbb{E}[\log^+ B] < \infty$. The assumptions of Theorem 5.1 are satisfied so (8.7) has a unique solution.

Suppose that X is the solution to (8.7). It is non-negative because it has the representation (5.2), where all random variables are non-negative. Suppose that $(A_1, B_1) \stackrel{d}{=} (A, B)$, $(A_2, B_2) \stackrel{d}{=} (A, B)$, and (A_1, B_1) , (A_2, B_2) and X are jointly independent. Then

$$\mathbb{P}(X < \varepsilon) = \mathbb{P}(A_1 X + B_1 < \varepsilon) = \mathbb{P}(A_2(A_1 X + B_1) + B_2 < \varepsilon) \leq \mathbb{P}(A_2 B_1 + B_2 < \varepsilon).$$

Now (8.8) follows from Lemma 8.2. \square

Remark 8.5. One can actually show that

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(X < \varepsilon) = -\frac{1}{2}.$$

However, the proof would take several additional pages so we will only sketch it. An appropriate modification of the argument in the proof of Lemma 8.2 shows that for the sequence $X_n = A_n X_{n-1} + B_n$ with $X_0 = 0$, we have $\limsup_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(X_n < \varepsilon) \leq \tau_n$, where $\tau_n \downarrow -\frac{1}{2}$. Using the fact that $\mathbb{P}(X < \varepsilon) \leq \mathbb{P}(X_n < \varepsilon)$ it follows that

$$\limsup_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(X < \varepsilon) \leq -\frac{1}{2}.$$

On the other hand for every $\delta > 0$, one can find a bounded positive function g_δ on $(0, \infty)$ such that

$$\mathbb{P}(Ax + B < \varepsilon) \geq g_\delta(x) e^{-(\frac{1}{2} + \delta)\varepsilon^{-1}},$$

for all $x > 0$. Hence, $\mathbb{P}(AX + B < \varepsilon) \geq \mathbb{E}[g_\delta(X)] e^{-(\frac{1}{2} + \delta)\varepsilon^{-1}}$, and, therefore,

$$\liminf_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbb{P}(X < \varepsilon) \geq -\left(\frac{1}{2} + \delta\right).$$

The claim follows by letting $\delta \rightarrow 0^+$.

Acknowledgements

The authors would like to thank Bojan Basrak, Hrvoje Planinić and anonymous referees for the most useful advice. The third author would like to thank the Department of Mathematics at the University of Washington in Seattle, where the project took place, for the hospitality. The third author is also grateful to the Microsoft Corporation for allowance on Azure cloud service where he ran many simulations.

Research of the first author was supported in part by Simons Foundation Grant 506732. Research of the third author was supported in part by Croatian Science Foundation grant 3526.

References

- [1] Bieniek, M., Burdzy, K. and Pal, S. (2012). Extinction of Fleming–Viot-type particle systems with strong drift. *Electron. J. Probab.* **17** no. 11, 15 pp. [MR2878790](#)
- [2] Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1987). *Regular Variation. Encyclopedia of Mathematics and Its Applications* **27**. Cambridge: Cambridge Univ. Press. [MR0898871](#)
- [3] Brockwell, P.J. and Davis, R.A. (1991). *Time Series: Theory and Methods*, 2nd ed. *Springer Series in Statistics*. New York: Springer. [MR1093459](#)
- [4] Buraczewski, D., Damek, E. and Mikosch, T. (2016). *Stochastic Models with Power-Law Tails: The Equation $X = AX + B$* . *Springer Series in Operations Research and Financial Engineering*. Cham: Springer. [MR3497380](#)
- [5] Burdzy, K., Holyst, R. and March, P. (2000). A Fleming–Viot particle representation of the Dirichlet Laplacian. *Comm. Math. Phys.* **214** 679–703. [MR1800866](#)
- [6] Diaconis, P. and Freedman, D. (1999). Iterated random functions. *SIAM Rev.* **41** 45–76. [MR1669737](#)
- [7] Goldie, C.M. (1991). Implicit renewal theory and tails of solutions of random equations. *Ann. Appl. Probab.* **1** 126–166. [MR1097468](#)
- [8] Hoff, P.D. (2009). *A First Course in Bayesian Statistical Methods. Springer Texts in Statistics*. New York: Springer. [MR2648134](#)
- [9] Kesten, H. (1973). Random difference equations and renewal theory for products of random matrices. *Acta Math.* **131** 207–248. [MR0440724](#)
- [10] Kołodziejek, B. (2018). On perpetuities with light tails. *Adv. in Appl. Probab.* **50** 1119–1154. [MR3881112](#)
- [11] Tchen, A.H. (1980). Inequalities for distributions with given marginals. *Ann. Probab.* **8** 814–827. [MR0577318](#)

Received May 2018 and revised January 2019